

If you are this age, talk to your health care professional about these vaccines

If you are this age,	Flu Influenza	Tdap or Td Tetanus, diphtheria, pertussis	Shingles Zoster		Pneumococcal		Meningococcal		MMR Measles, mumps, rubella	HPV Human papillomavirus		Chickenpox Varicella	Hepatitis A	Hepatitis B	Hib <i>Haemophilus influenzae type b</i>
			RZV	ZVL	PCV13	PPSV23	MenACWY	MenB		for women	for men				
19 - 21 years	Dark	Dark			Dark	Dark	Dark	Dark	Dark			Dark	Dark	Dark	Dark
22 - 26 years	Dark	Dark			Dark	Dark	Dark	Dark	Dark			Dark	Dark	Dark	Dark
27 - 49 years	Dark	Dark			Dark	Dark	Dark	Dark	Dark			Dark	Dark	Dark	Dark
50 - 64 years	Dark	Dark	Dark		Dark	Dark	Dark	Dark	Dark	Dark		Dark	Dark	Dark	Dark
65+ year	Dark	Dark		Dark	Dark	Dark	Dark	Dark	Dark	Dark		Dark	Dark	Dark	Dark

More Information:

You should get flu vaccine every year.

You should get 1 dose of Tdap if you did not get it as a child or adult. You should also get a Td booster every 10 years. Women should get 1 dose of Tdap during every pregnancy.

There are 2 types of zoster vaccine. You should get 2 doses of RZV at age 50 years or older (preferred) or 1 dose of ZVL at age 60 years or older, even if you had shingles before.

There are 2 types of pneumococcal vaccine. You should get 1 dose of PCV13 and at least 1 dose of PPSV23 depending on your age and health condition.

There are 2 types of meningococcal vaccine. You may need one or both types depending on your health condition.

You should get this vaccine if you did not get it when you were a child.

You should get HPV vaccine if you are a woman through age 26 years or a man through age 21 years and did not already complete the series.



Recommended For You: This vaccine is recommended for you *unless* your health care professional tells you that you do not need it or should not get it.



May Be Recommended For You: This vaccine is recommended for you if you have certain risk factors due to your health condition. Talk to your health care professional to see if you need this vaccine.

If you are traveling outside the United States, you may need additional vaccines.
Ask your health care professional about which vaccines you may need at least 6 weeks before you travel.

For more information, call 1-800-CDC-INFO (1-800-232-4636) or visit www.cdc.gov/vaccines



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

If you have this health condition,

talk to your health care professional about these vaccines



	Flu Influenza	Tdap or Td Tetanus, diphtheria, pertussis	Shingles Zoster		Pneumococcal		Meningococcal		MMR Measles, mumps, rubella	HPV Human papillomavirus		Chickenpox Varicella	Hepatitis A	Hepatitis B	Hib <i>Haemophilus influenzae type b</i>
			RZV	ZVL	PCV13	PPSV23	MenACWY	MenB		for women	for men				
Pregnancy	Dark	Dark		Light	Dark	Dark	Dark	Dark	Light	Dark	Dark	Light	Dark	Dark	Dark
Weakened Immune System	Dark	Dark		Light	Dark	Dark	Dark	Dark	Light	Dark	Dark	Light	Dark	Dark	Dark
HIV: CD4 count less than 200	Dark	Dark		Light	Dark	Dark	Dark	Dark	Light	Dark	Dark	Light	Dark	Dark	Dark
HIV: CD4 count 200 or greater	Dark	Dark		Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark
Kidney disease or poor kidney function	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark
Spleen removed or does not work well	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark
Heart disease Chronic lung disease Chronic alcoholism	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark
Diabetes (Type 1 or Type 2)	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark
Chronic Liver Disease	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark	Dark

More Information:

You should get flu vaccine every year.

You should get 1 dose of Tdap if you did not get it as a child or adult. You should also get a Td booster every 10 years. Women should get 1 dose of Tdap vaccine during every pregnancy.

There are 2 types of zoster vaccine. You should get 2 doses of RZV at age 50 years or older (preferred) or 1 dose of ZVL at age 60 years or older, even if you had shingles before.

There are 2 types of pneumococcal vaccine. You should get 1 dose of PCV13 and at least 1 dose of PPSV23 depending on your age and health condition.

There are 2 types of meningococcal vaccine. You may need one or both types depending on your health condition.

You should get this vaccine if you did not get it when you were a child.

You should get HPV vaccine if you are a woman through age 26 years or a man through age 21 years and did not already complete the series.

You should get Hib vaccine if you do not have a spleen, have sickle cell disease, or received a bone marrow transplant.

Recommended For You: This vaccine is recommended for you *unless* your health care professional tells you that you do not need it or should not get it.

May Be Recommended For You: This vaccine is recommended for you if you have certain other risk factors due to your health condition. Talk to your health care professional to see if you need this vaccine.

YOU SHOULD NOT GET THIS VACCINE

For more information, call 1-800-CDC-INFO (1-800-232-4636) or visit www.cdc.gov/vaccines



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention