

Influenza Vaccine: Who Should Get It, and Who Should Not

FIGHT FLU



Everyone 6 months and older is recommended for annual influenza vaccination, with rare exceptions. **For the 2018-19 flu season, ACIP recommends annual influenza vaccination for everyone 6 months and older with any licensed, appropriate influenza vaccine (IIV, RIV4, or LAIV4) with no preference expressed for any one vaccine over another. Some vaccines are not recommended in some situations and health conditions, and some people should not receive influenza vaccines at all (though this is uncommon).**

Different flu shots are approved for people of different ages. Everyone should get a vaccine that is appropriate for their age.

The Influenza (Flu) Shot

People who can get the flu shot

- There are inactivated influenza vaccines that are approved for people as young as 6 months of age.
- Some vaccines are only approved for adults. For example, the recombinant influenza vaccine is for people aged 18 years and older, and the adjuvanted and high-dose inactivated vaccines are for people aged 65 years and older.
- Flu shots are recommended for pregnant women and people with chronic health conditions.

People who SHOULD NOT get the flu shot

- Children younger than 6 months of age are too young to get a flu shot.
- People with severe, life-threatening allergies to flu vaccine or any ingredient in the vaccine. This might include gelatin, antibiotics, or other ingredients. See special considerations regarding egg allergy for more information about egg allergies and flu vaccine.

People who should talk to their doctor before getting the flu shot:

If you have one of the following conditions, talk with your healthcare provider. He or she can help decide whether vaccination is right for you, and select the best vaccine for your situation:

- If you have an allergy to eggs or any of the ingredients in the vaccine. Talk to your doctor about your allergy. See special considerations regarding egg allergy at the bottom of this document for more information about egg allergies and flu vaccine.
- If you ever had Guillain-Barré Syndrome (a severe paralyzing illness, also called GBS). Some people with a history of GBS should not get this vaccine. Talk to your doctor about your GBS history.
- If you are not feeling well, talk to your doctor about your symptoms.

For more information, visit:
www.cdc.gov/flu or www.flu.gov
or call **1-800-CDC-INFO**



**U.S. Department of
Health and Human Services**
Centers for Disease
Control and Prevention

Nasal Spray Flu Vaccine:

People who can get the nasal spray flu vaccine:

The nasal spray vaccine is approved for use in people 2 years through 49 years of age. It is an option for healthy, non-pregnant people in this age group.

People who SHOULD NOT get the nasal spray vaccine:

- Children younger than 2 years
- Adults 50 years and older
- Pregnant women
- People with a history of severe allergic reaction to any component of the vaccine or to a previous dose of any influenza vaccine
- Children 2 years through 17 years of age who are receiving aspirin- or salicylate-containing medications.
- People who are immunocompromised (those who have weakened immune systems)
- Children 2 years through 4 years who have asthma or who have had a history of wheezing in the past 12 months.
- People who have taken influenza antiviral drugs within the previous 48 hours.
- People who care for severely immunocompromised persons who require a protected environment (or otherwise avoid contact with those persons for 7 days after getting the nasal spray vaccine).

People who should talk to their healthcare provider before getting nasal spray vaccine:

If you have one of the following condition, talk with your healthcare provider. He or she can help decide whether vaccination is right for you, and select the best vaccine for your situation:

- People with Asthma aged 5 years and older.
- People with other underlying medical conditions that can put them at higher risk of serious flu complications. These include conditions such as chronic lung diseases, heart disease (except isolated hypertension), kidney disease, liver disorders, neurologic and neuromuscular disorders, blood disorders, or metabolic disorders (such as diabetes). See "People at High Risk of Developing Flu-Related Complications."
- People with moderate or severe acute illness with or without fever.
- People with Guillain-Barré Syndrome within 6 weeks following a previous dose of influenza vaccine.

Special Consideration Regarding Egg Allergy

People with egg allergies can receive any licensed, recommended age-appropriate influenza vaccine (IIV, RIV4, or LAIV4) that is otherwise appropriate. People who have a history of severe egg allergy (those who have had any symptom other than hives after exposure to egg) should be vaccinated in a medical setting, supervised by a health care provider who is able to recognize and manage severe allergic