
100 Years of Advances Against Cancer

The National Institutes of Health and the National Cancer Institute in partnership and cooperation with the scientific research community.



Created by the National Cancer Institute's Grants Administration Branch.

Gratwick Laboratories (Roswell Park Memorial Institute). First cancer research laboratories established in 1898, moved in 1901.

1900s-1920's



1903 Radium found effective in treatment of tumors.

1911 P. Rous discovered a virus that causes cancer in chickens.

Black and white drawing of N.Y. Cancer Hospital, Memorial Sloan Kettering

(Est. 1884)



1912 Cancer cells are grown in the laboratory, the first long-term "tissue culture."

1913 Publication of first known article on cancer's warning signs:

Ladies' Home Journal

1915 Coal tar gives rabbits cancer in experimental proof of carcinogenesis.

1904: View of American Oncologic Hospital, the first precursor of Fox Chase Cancer Center.

1927: View of Institute for Cancer Research founded in 1927 at Lankenau Research Hospital. The two organizations joined in 1966 to become Fox Chase Cancer Center.



'Conquer Cancer' Adopted as Battle Cry of the Public Health Service

FatalitiesFromDisease **AreIncreasingRapidly**

Surgeon General Parran Believes 25,000 Lives Can Be Saved Yearly by Application of Information Already Available.


By Christine Sadler,

By Christine Sadler.

In the natural order of advancement cancer should have diagnostic and the belief in three-baseds and the belief in three-based and grant That is today the second and the belief in three-based and the based of the same and the s

Cancer, the second greatest killer in the United States, has become the country's No. 1 medical problem. And the Public Health Service is now engaged in a Nation-wide campaign to wipe out this ancient scourge. Last week President Roosevelt signed a bill authorizing the erection of a \$750,000 National Cancer Institute to be built on a site near Bethesda, Md. The land was donated by the late Luke Wilson, a trustee of Woodward & Lothrop, who was a victim of cancer.

A famous cancer expert, Dr. Carl Voegtlin, of the National Institute of Health, has been suggested as head of the new foundation. He is shown below with Dr. R. H. Fitch (left) and Dr. Herbert Kaher (right) at work on a phase of their studies of malignant growths.



The super voltage X-ray tube above was designed to test high-voltage X-ray machines and compare the



U.S. Creates New Division For Research


Bills Rushed by Congress Are Signed by President.

Victim of Dreaded Maladv Donated Land for Laboratories.

sion of the Public Health Service, is expected to get one of the applications. Other experts, from whom applications of the experts, and the experts of the experts of the Memoral Hospital for the Treatment of Casore and Alliest Mays and his stroker, Dr. William Mays, of the Mayo Clinic: Dr. Personte Carter Wood, head of the Mayo Charter of the Mayo Clinic Dr. Mayo and his herber, Dr. William Mays, of the Mayo Clinic; Dr. Mayo and his herber, Dr. William Mays, of the Mayo Clinic; Dr. Marc Culfer, Mayor and Carter Wood, Bed of the Mayor Chicago, Chicago, Chicago, Chicago, Ling of Chicago, Dr. Temple Payor, and the Chicago, and the Chicago

desideratum, the one on which all campaigns against canner are hased."

A major task undertaken by the elimitatus will be the obecking of announced "canner cures." Thousands of Americans are dupded annually by decites whose purported in the field of annex. The cures are not recognized by the "Verytilla Mentiseed" Verytilla Mentiseed.





1930 The National Institute of Health is established by the Ransdell Act.

1938 NCI Budget: \$400,000

June 6, 1938

Members of the first National Advisory Cancer Council at the groundbreaking ceremonies at the NCI's building 6.



June 24, 1939 Cornerstone laying for NCI's building



Shown: Mrs. Luke Wilson, whose husband, a cancer victim, donated the land Dr. Thomas Parton, Surgeon General of the U.S. Public Health Service & Henry Morgenthau, Secretary of the Treasury (Holding trowel)

Welcome to the National Cancer Institute





1939 The National Cancer Institute's first home, "Building 6".

1937 The first NCI staff was comprised of Harvard personnel.



November 27,1937: First NCI grant was awarded for \$27,550 to Louis F. Fieser to investigate chemical structure and carcinogenic activity.

1940's

1940 First issue of the Journal of the National Cancer Institute is published.

1943 Pap smear is introduced into medical practice.

1944 DNA found by O. Avery, C. MacLeod, and M. McCarty to be the basic cell material.



NCI's First Scientific Staff Source: *Life Magazine*, June 17, 1940


1940-1949 NCI appropriations \$42 million!

October 31, 1940 Pres. Franklin Roosevelt dedicates the first 6 buildings of NIH.



National Institutes of Health Campus - 1940

1947 S. Farber finds that a folic acid derivative inhibits acute leukemia.



1948 G. Hitchings synthesizes 6-mercaptopurine (6-MP) to combat childhood leukemia.

1949 FDA approves nitrogen mustard (methclorethanine), a drug that interacts with DNA chemically to kill cancer cells.

S. Farber also founds Children's Hospital Cancer Research Foundation in the next decade.

1950 E. Wynder, E. Graham, and Sir R. Doll confirm cigarette smoking-cancer link.

19503


1952 DNA found to be genetic material in some viruses.

1955 R. Hertz and M. Chiu Li achieve total cure of a human solid tumor, choriocarcinoma.

1953 FDA approves methotrexate as anti-cancer drug.

1953 J. Watson and F. Crick discover the structure of DNA.

1955 National Chemotherapy Program begins.



1950-1959 NCI appropriations \$330 million!

1960 Chromosome abnormality associated with leukemias.

1961 M. Nirenberg and others prove triplet code governs DNA action.

1962 Royal College of Physicians issues report on smoking and health.

1964

- •U.S. Surgeon General issues Report on Smoking and Health
- •A virus (Epstein-Barr virus) is linked to human cancer for the first time.
- •American Society of Clinical Oncology established.



Dr. Emil Freireich working with a blood cell separator centrifuge at M.D. Anderson Hospital.



1960-1969 NCI appropriations \$1.8 billion!

FDA Approvals:

- 1961 Vinblastine: a drug that binds to tubulin. Derived from the ornamental shrub, vinca rosea.
- **1962** 5-FU
- 1963 Vincristine: a sister drug to Vinblastine.
- 1964 Melphalan (L-PAM) approved for marketing.

1966 NCI standardizes testing of cancer-causing chemicals.

1969 R. Heubner & G. Todaro propose the oncogene hypothesis.





1970-1979 NCI appropriations \$6.1 billion.

1970 H. Temin & D. Baltimore discover reverse transcriptase enzyme: a key to gene engineering.

1971 President Richard M.
Nixon converted Army's
former biological warfare
facilities at Ft. Detrick, MD, to
house research activities on the
causes, treatment, and
prevention of cancer.

1973

- •Computed tomography (CT) introduced in the United States
- •Recombinant DNA techniques developed for cloning genes
- •Surveillance, Epidemiology, and End Results (SEER) Program established.

December 23, 1971 President Nixon signs the National Cancer Act of 1971.



1974

- •CANCERLINE, a national database of published cancer research established.
- •FDA approves doxorubicin, an anti-tumor antibiotic from Streptomyces bacteria.

1976

- •Cancer Information Service (1-800-4-CANCER) opened.
- •Interleukin-2 discovered.
- •First human proto-oncogenes discovered.

1975 Methods developed to identify and sequence DNA fragments.

1977 First national cancer patient education program founded (I Can Cope).

1978

- First human testing of a biological therapy (alpha-interferon).
- Tamoxifen approved by FDA for marketing as a treatment drug.
- FDA approves cisplatin, a powerful anti-cancer drug.
- Metastic cells shown to arise from pre-existing subpopulations in primary tumors.



1979

- •P53 discovered, most frequently mutated gene in human cancer.
- •Modified radical mastectomy replaces radical mastectomy for breast cancer.

1970's

- •Studies in human populations link cancer risk to infectious agents, such as human papillomavirus (cervical cancer) and hepatitis B (liver cancer).
- •Statistical methods developed to control simultaneously for several factors in the analysis of studies and to quantify cancer risks.
 - •Studies clarify the patterns of cancer risk following exposure to ionizing radiation.

•Studies link cancer risks to hormonal drugs, such as diethylstilbestrol (DES) taken during pregnancy and hormonal replacement therapy.

1980-1989 NCI appropriations \$12 billion!

1980's

•Magnetic resonance imaging (MRI) introduced.



•Flexible sigmoidoscopy and colonoscopy developed to help find and remove precancerous growths.

•Continuous pain medication infusion pumps developed.



•First highly effective anti-nausea drugs developed to alleviate side effects of chemotherapy.

•Biochemical and genetic assays integrated into epidemiologic studies (molecular epidemiology).

1981 Introduction of first human viral vaccine that can prevent cancer (hepatitis B virus vaccine for liver cancer).

1985 Lumpectomy plus radiation found equivalent to mastectomy for breast cancer.

1986 First tumor suppressor gene cloned (Rb).

Adjuvant chemotherapy:

- •1988 Proven to increase disease-free survival in early breast cancer
- •1989 Proven to increase survival in colon cancer




19905

Breast cancer death rates began to decline!

•Multi-step nature of carcinogenesis proven.

•Transition from film-based radiology to digital computer-assisted medical imaging.

•Several common genetic variants linked to the risk of lung and other cancers.




•Fluorescence in situ hybridization technique developed (FISH).

1990-1999 NCI appropriations \$21.8 billion!

1990 First chemoprevention trial to show efficacy – vitamin A analogue against mouth and throat tumors.

1991 Adjuvant radiation and chemotherapy found to improve survival in rectal cancer.



Cloning Advances:

- •1993 First of the hereditary nonpolyposis colon cancer genes
- •1994 BRCA1, the first inherited breast cancer gene
- •1995 BRCA2

FDA Approvals:

- •1995 Tretinoin, the first successful differentiating agent
- •1995 Porfimer sodium, a drug that sensitizes tumors to light, permitting photodynamic therapy in the U.S.
- •1996 Topecan, first of a class of drugs that interferes with the enzyme topoisomerase
- •1997 Rituximab, first biotechnology product approved by FDA to treat patients with cancer
- •1998 Trastuzumab (Herceptin), targets cancer cells that produce a protein found in high number of women with metastic breast cancer

1990's Studies and Trials

1992 Breast Cancer Prevention Trial begins, testing tamoxifen as a preventive agent in women at increased risk of the disease (1998: Results found that tamoxifen reduced the chances of women at risk of developing breast cancer by half.)

1993

- •NCI-sponsored studies in China show importance of nutrition in preventing cancer.
- •Prostate Cancer Prevention Trial begins, testing finasteride, a drug used to reduce symptoms of prostate enlargement.
- •The Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial begins recruiting 148,000 volunteers (the largest early detection study).
- 1997 Cancer Genome Anatomy Project is launched, a multi-year project to assemble the first index of genes involved in cancer.

A New Century...

NIH-Funded Centers to Seek Early Environmental Exposures that May Lead to Breast Cancer

10/14/2003 -- National Institutes of Health announced the funding of four new Breast Cancer and the Environment Research Centers to study the prenatal-to-adult environmental exposures that may predispose a woman to breast cancer.

NIH Funds Eight New Grants Focused on Aging and Cancer

10/10/2003 – Seventy-seven percent of all cancers are diagnosed at 55 and older. With cancer death rates highest among people 65 years and older, the National Cancer Institute (NCI) and the National Institute on Aging (NIA) are launching a new initiative to accelerate research into the relationship between aging and cancer.

New Advances

New Treatment Significantly Improves Long-Term Outlook for Breast Cancer Survivors

10/9/2003 International clinical trial concludes women should consider taking letrozole after five years of tamoxifen treatment to continue to reduce risk of recurrence.

Decline Shown in Death Rates from Four Leading Cancers:

09/02/2003 - Death rates from the four most common cancers - lung, breast, prostate, and colorectal - continued to decline in the late 1990s according to new data from the "Annual Report to the Nation on the Status of Cancer, 1975-2000."

1938 – 2003 NCI appropriations total \$57.5 billion!

Making a Difference...

Clinicians Scientists

Researchers

Administrators

Volunteers

Patients

American Public

In 2001, there were 9,600,000 cancer survivors in the U.S.

We...are Changing the World.

Timeline information and Pictures extracted from the following website: http://cancer.gov

Our Challenge Goal to the Nation

Eliminate the suffering and death due to cancer by 2015.

Andrew C. von Eschenbach, M.D.

Director National Cancer Institute

October 2003