Table 6.31A Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Numbers in Thousands, 2002

	Total	AGE GROUP (Years)		
Location of Treatment/Counseling ¹		18-25	26-49	50 or Older
Outpatient Mental Health Clinic or Center	3,136	479	1,956	702
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	8,461	1,091	4,914	2,456
Doctor's Office - Not Part of a Clinic	3,796	383	2,173	1,239
Outpatient Medical Clinic	950	167	400	383
Partial Day Hospital or Day Treatment Program	289	72	199	19
School or University Setting/Clinic/Center ²	107	70	37	*
Church//Religious Setting ²	82	25	57	*
Some Other Place	389	56	229	104

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 6.31B Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Percentages, 2002

Location of Treatment/Counseling ¹	Total	AGE GROUP (Years)		
		18-25	26-49	50 or Older
Outpatient Mental Health Clinic or Center	20.1	22.7	21.7	15.6
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	54.1	51.7	54.5	54.5
Doctor's Office - Not Part of a Clinic	24.3	18.2	24.1	27.5
Outpatient Medical Clinic	6.1	7.9	4.4	8.5
Partial Day Hospital or Day Treatment Program	1.9	3.4	2.2	0.4
School or University Setting/Clinic/Center ²	0.7	3.3	0.4	*
Church//Religious Setting ²	0.5	1.2	0.6	*
Some Other Place	2.5	2.7	2.5	2.3

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 6.32A Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Numbers in Thousands, 2002

	Total	AGE GROUP (Years)		
Location of Treatment/Counseling ¹		18-25	26-49	50 or Older
Outpatient Mental Health Clinic or Center	1,415	263	928	*
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	3,034	554	1,861	*
Doctor's Office - Not Part of a Clinic	1,528	194	1,044	*
Outpatient Medical Clinic	370	65	166	*
Partial Day Hospital or Day Treatment Program	100	21	64	*
School or University Setting/Clinic/Center ²	36	27	9	*
Church//Religious Setting ²	25	8	17	*
Some Other Place	138	21	89	*

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic* and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings.*

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

Table 6.32BLocation of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness
Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Percentages, 2002

	Total	AGE GROUP (Years)		
Location of Treatment/Counseling ¹		18-25	26-49	50 or Older
Outpatient Mental Health Clinic or Center	24.7	26.7	25.9	*
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	52.9	56.2	51.9	*
Doctor's Office - Not Part of a Clinic	26.6	19.7	29.1	*
Outpatient Medical Clinic	6.5	6.7	4.6	*
Partial Day Hospital or Day Treatment Program	1.7	2.2	1.8	*
School or University Setting/Clinic/Center ²	0.6	2.7	0.3	*
Church//Religious Setting ²	0.4	0.8	0.5	*
Some Other Place	2.4	2.2	2.5	*

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for alcohol or drug use. Respondents with missing treatment/counseling information are excluded.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4th edition of the *Diagnostic* and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.6 of Appendix B of the *Results from the 2002 National Survey on Drug Use and Health: National Findings.*

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.