

May 3, 2001

Dear Colleague:

Information related to the 2000 BRFSS summary prevalence report of selected risk factors and preventive health measures from the core and modules is attached.

This report consists of four parts:

An introduction that contains:

- An overview of the 2000 summary prevalence report.
- An explanation of the tables for 2000 Behavioral Risk Factor Surveillance System (BRFSS) prevalence data.
- Definitions of BRFSS risk factors and preventive health measures. When appropriate, Year 2000 Objectives are identified.

Analysis of selected questions from the 2000 BRFSS questionnaire. The set of analyses is divided into three sections:

Section I Selected risk factors and preventive health measures in the BRFSS core.

Section II Selected risk factors and preventive health measures from modules used by at least 13 states

Section III Preventive health measures that concern women only.

As soon as your schedule permits, please examine the enclosed materials closely.

Sincerely yours,

Eve Powell-Griner, Ph. D.
Acting Chief, Behavioral Surveillance Branch
Division of Adult and Community Health
National Center for Chronic Disease Prevention and Health

Promotion

2000 BRFSS SUMMARY PREVALENCE REPORT

BEHAVIORAL SURVEILLANCE BRANCH

DIVISION OF ADULT AND COMMUNITY HEALTH

NATIONAL CENTER FOR CHRONIC DISEASE PREVENTION AND HEALTH PROMOTION

CENTERS FOR DISEASE CONTROL AND PREVENTION

UNITED STATES DEPARTMENT OF HEALTH AND HUMAN SERVICES

TABLE OF CONTENTS

OVERVIEW OF THE 2000 SUMMARY REPORT	VII
EXPLANATION OF TABLES.....	VII
DEFINITIONS OF TERMS AND RISK FACTORS	X
LIMITATIONS OF THE DATA.....	XV

SECTION I PREVALENCE REPORT FOR CORE QUESTIONS

GENERAL HEALTH STATUS

TABLE 1.1 SELF REPORTED HEALTH STATUS: FAIR OR POOR	1
1.2 SELF REPORTED HEALTH STATUS: FAIR OR POOR-BY SEX	2
1.3 SELF REPORTED HEALTH STATUS: FAIR OR POOR, CHANGE 1999-2000	3

QUALITY OF LIFE: PHYSICAL AND MENTAL HEALTH

TABLE 2.1 PHYSICAL HEALTH NOT GOOD.....	4
2.2 PHYSICAL HEALTH NOT GOOD-BY SEX.....	5
2.3 PHYSICAL HEALTH NOT GOOD, CHANGE 1999-2000.....	6
TABLE 3.1 MENTAL HEALTH NOT GOOD.....	7
3.2 MENTAL HEALTH NOT GOOD-BY SEX.....	8
3.3 MENTAL HEALTH NOT GOOD, CHANGE 1999-2000.....	9
TABLE 4.1 ACTIVITIES LIMITED.....	10
4.2 ACTIVITIES LIMITED-BY SEX.....	11
4.3 ACTIVITIES LIMITED, CHANGE 1999-2000.....	12

HEALTH INSURANCE

TABLE 5.1 NO HEALTH INSURANCE, AGE 18 AND OVER.....	13
5.2 NO HEALTH INSURANCE, AGE 18 AND OVER-BY SEX.....	14
5.3 NO HEALTH INSURANCE, AGE 18 AND OVER, CHANGE 1999-2000.....	15
TABLE 6.1 NO HEALTH INSURANCE, AGE 18-64 YEARS	16
6.2 NO HEALTH INSURANCE, AGE 18-64 YEAR-BY SEX	17
6.3 NO HEALTH INSURANCE, AGE 18-64 YEARS, CHANGE 1999-2000	18

DIABETES AWARENESS

TABLE 7.1 EVER TOLD YOU HAVE DIABETES	19
7.2 EVER TOLD YOU HAVE DIABETES-BY SEX	20
7.3 EVER TOLD YOU HAVE DIABETES, CHANGE 1999-2000	21

SMOKING STATUS

TABLE 8.1 EVER SMOKED	22
8.2 EVER SMOKED-BY SEX	23
8.3 EVER SMOKED, CHANGE 1999-2000	24
TABLE 9.1 CURRENT SMOKER	25
9.2 CURRENT SMOKER-BY SEX	26
9.3 CURRENT SMOKER, CHANGE 1999-2000.....	27

PHYSICAL ACTIVITY

TABLE 10.1	NO LEISURE TIME PHYSICAL ACTIVITY.....	28
10.2	NO LEISURE TIME PHYSICAL ACTIVITY BY SEX.....	29
10.3	NO LEISURE TIME PHYSICAL ACTIVITY, CHANGE 1998-2000.....	30
TABLE 11.1	NO REGULAR OR SUSTAINED PHYSICAL ACTIVITY	31
11.2	NO REGULAR OR SUSTAINED PHYSICAL ACTIVITY-BY SEX.....	32
11.3	NO REGULAR OR SUSTAINED PHYSICAL ACTIVITY, CHANGE 1998-2000	33
TABLE 12.1	NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY.....	34
12.2	NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY-BY SEX.....	35
12.3	NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY, CHANGE 1998-2000.....	36

FRUITS AND VEGETABLES

TABLE 13.1	DOES NOT EAT 5 SERVINGS A DAY.....	37
13.2	DOES NOT EAT 5 SERVINGS A DAY-BY SEX.....	38
13.3	DOES NOT EAT 5 SERVINGS A DAY, CHANGE 1998-2000.....	39

WEIGHT CONTROL

TABLE 14.1	DOES NOT EAT FEWER CALORIES AND/LESS FAT OR EXERCISE TO LOSE WEIGHT.....	40
14.2	DOES NOT EAT FEWER CALORIES AND/LESS FAT OR EXERCISE TO LOSE WEIGHT-BY SEX.....	41
14.3	DOES NOT EAT FEWER CALORIES AND/LESS FAT OR EXERCISE TO LOSE WEIGHT CHANGE 1998-2000	42

OVERWEIGHT

TABLE 15.1	OVERWEIGHT BASED ON BODY MASS INDEX (BMI > 25.0).....	43
15.2	OVERWEIGHT BASED ON BODY MASS INDEX (BMI > 25.0)-BY SEX	44
15.3	OVERWEIGHT BASED ON BODY MASS INDEX (BMI > 25.0), CHANGE 1999-2000.....	45

HIV/AIDS

TABLE 16.1	WOULD NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64	46
16.2	WOULD NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64-BY SEX	47
16.3	WOULD NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64, CHANGE 1999-2000	48
TABLE 17.1	CHANCES OF GETTING HIV MEDIUM OR HIGH, AGE 18-64	49
17.2	CHANCES OF GETTING HIV MEDIUM OR HIGH, AGE 18-64-BY SEX	50
17.3	CHANCES OF GETTING HIV MEDIUM OR HIGH, AGE 18-64, CHANGE 1999-2000	51
TABLE 18.1	NOT TESTED FOR HIV WITHIN PAST 12 MONTHS, AGE 18-64.....	52
18.2	NOT TESTED FOR HIV WITHIN PAST 12 MONTHS, AGE 18-64	53
18.3	NOT TESTED FOR HIV WITHIN PAST 12 MONTHS, AGE 18-64, CHANGE 1999-2000.....	54

SECTION II

PREVALENCE REPORT FOR THE MODULES

DIABETES

TABLE 19.1	AGE AT ONSET OF DIABETES.....	55
19.2	AGE AT ONSET OF DIABETES-BY SEX.....	56
TABLE 20.1	DIABETICS USING INSULIN.....	57
20.2	DIABETICS USING INSULIN-BY SEX.....	58
TABLE 21.1	HEALTH PROFESSIONAL VISITS IN PAST YEAR (by DIABETICS	59
21.2	HEALTH PROFESSIONAL VISITS IN PAST YEAR (by DIABETICS)-BY SEX.....	60

SMOKELESS TOBACCO

TABLE 22.1	EVER USE SMOKELESS TOBACCO.....	61
22.2	EVER USE SMOKELESS TOBACCO-BY SEX.....	62
TABLE 23.1	CURRENTLY USES SMOKELESS TOBACCO	63
23.2	CURRENTLY USES SMOKELESS TOBACCO-BY SEX	64

IMMUNIZATION

TABLE 24.1	DID NOT HAVE FLU SHOT WITHIN 12 MONTHS, AGE 65 AND OVER	65
24.2	DID NOT HAVE FLU SHOT WITHIN 12 MONTHS, AGE 65 AND OVER-BY SEX	66
TABLE 25.1	NEVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER	67
25.2	NEVER HAD PNEUMONIA VACCINATION, AGE 65 AND OVER-BY SEX	68

CARDIOVASCULAR DISEASE

TABLE 26.1	EVER TOLD YOU HAD A HEART ATTACK OR MYOCARDIAL INFARCTION.....	69
26.2	EVER TOLD YOU HAD A HEART ATTACK OR MYOCARDIAL INFARCTION-BY SEX.....	70
TABLE 27.1	EVER TOLD YOU HAD A STROKE.....	71
27.2	EVER TOLD YOU HAD A STROKE-BY SEX.....	72

SECTION III

PREVALENCE REPORT FOR WOMEN'S HEALTH ISSUES

MAMMOGRAPHY

TABLE 28.1	NEVER HAD MAMMOGRAM, AGE 18 AND OVER	73
28.2	NEVER HAD MAMMOGRAM, AGE 18 AND OVER, CHANGE 1999-2000	74
TABLE 29.1	NEVER HAD MAMMOGRAM, AGE 40 AND OVER	75
29.2	NEVER HAD MAMMOGRAM, AGE 40 AND OVER, CHANGE 1999-2000	76
TABLE 30.1	DID NOT HAVE A MAMMOGRAM WITHIN PAST TWO YEARS, AGE 50 AND OVER.....	77
30.2	DID NOT HAVE A MAMMOGRAM WITHIN PAST TWO YEARS, AGE 50 AND OVER, CHANGE 1999-2000.....	78

BREAST EXAM

TABLE 31.1	NEVER HAD BREAST EXAM, AGE 18 AND OVER.....	79
31.2	NEVER HAD BREAST EXAM, AGE 18 AND OVER, CHANGE 1999-2000.....	80
TABLE 32.1	NEVER HAD BREAST EXAM, AGE 40 AND OVER.....	81
32.2	NEVER HAD BREAST EXAM, AGE 40 AND OVER, CHANGE 1999-2000.....	82
TABLE 33.1	DID NOT HAVE A BREAST EXAM WITHIN PAST TWO YEARS, AGE 50 AND OVER.....	83
33.2	DID NOT HAVE A BREAST EXAM WITHIN PAST TWO YEARS, AGE 50 AND OVER, CHANGE 1999-2000.....	84

MAMMMOGRAM AND BREAST EXAM

TABLE 34.1	NEVER HAD MAMMOGRAM & BREAST EXAM, AGE 40 AND OVER.....	85
34.2	NEVER HAD MAMMOGRAM & BREAST EXAM, AGE 40 AND OVER, CHANGE 1999-2000.....	86
TABLE 35.1	DID NOT HAVE MAMMOGRAM & BREAST EXAM W/IN PAST 2 YRS, AGE 50 AND OVER.....	87
35.2	DID NOT HAVE MAMMOGRAM & BREAST EXAM W/IN PAST 2 YRS, AGE 50 AND OVER, CHANGE 1999-2000.....	88

CERVICAL CANCER

TABLE 36.1	NEVER HAD PAP SMEAR, WOMEN AGE 18 AND OVER	89
36.2	NEVER HAD PAP SMEAR, WOMEN AGE 18 AND OVER, CHANGE 1999-2000	90
TABLE 37.1	NEVER HAD PAP SMEAR, WOMEN WITH INTACT CERVIX.....	91
37.2	NEVER HAD PAP SMEAR, WOMEN WITH INTACT CERVIX CHANGE 1999-2000.....	92
TABLE 38.1	DID NOT HAVE PAP SMEAR IN PAST 3 YEARS, WOMEN WITH INTACT CERVIX	93
38.2	DID NOT HAVE PAP SMEAR IN PAST 3 YEARS, WOMEN WITH INTACT CERVIX, CHANGE 1999-2000.....	94

OVERVIEW OF THE 2000 SUMMARY REPORT

The report is divided into three sections. *Section I* displays tables for selected variables in the core. In *Section I*, there are three tables for each risk factor or preventive measure unless otherwise noted. Estimated prevalence and relevant statistics are provided for the total population sampled and by sex. Prevalence estimates for both 1999 and 2000, or where appropriate 1998 and 2000, and the percentage point difference between the estimates are given for variables in the core.

Section II shows tables for selected variables in modules used by at least 13 states. *Section II* contains two tables for each risk factor or preventive measure of interest in the modules. These tables show the estimated prevalence and relevant statistics for the total sampled population and both sex groups who responded to selected questions in the optional modules.

Section III contains tables for preventive health measures that are related to women's health issues. Estimated prevalence and relevant statistics are provided for the total population sampled. Prevalence estimates for both 1999 and 2000 and the percentage point difference between them are given.

Three tables (Tables 1.1-1.3) from Section I will be used as examples to explain the information contained in this report. Each table describes a behavioral risk factor or preventive health measure from the Behavioral Risk Factor Surveillance System (BRFSS) instrument.

Table 1.1 (page 1) shows the estimated prevalence and relevant statistics for each participant. (Participants refer to all states, and the District of Columbia and territories of the U.S.). The table shows the estimated prevalence and relevant statistics for "fair or poor health status" in the population surveyed by each BRFSS participant. This prevalence estimate excludes respondents with unknown information on self reported health status or respondents choosing not to answer the question.

The columns in Table 1.1 provide the following information:

COLUMN HEADING

CONTENTS

<u>Participant</u>	Name of the participant. Participants refers to all states including the District of Columbia and territories of the United States.
<u>Sample size</u>	The number of individuals who responded to the specific BRFSS question. These are unweighted numbers. These numbers <u>exclude</u> individuals who reported "not sure, don't know", those who refused to answer the question, or had no response coded (missing or skipped).
<u>Percent/Mean</u>	An estimate of the percent (or prevalence or mean number) of residents who report the behavioral risk factor of interest. This is weighted to the participant's population to estimate the true prevalence or mean in the population.
<u>Standard error</u>	A measure of the amount of variability that exists in the calculation of the prevalence estimate.
<u>95% confidence interval</u>	A range which contains the true population prevalence estimate 95% of the time, if repeated sampling of the population is performed.

SUMMARY STATISTICS are given at the bottom of the table:

<u>Number of participants</u>	The total number of participants who asked question(s) regarding the respective "risk factor".
<u>Median</u>	The middle value of all estimated prevalences.
<u>Range</u>	The lowest prevalence and the highest prevalence among all participants.

For example, in Table 1.1, there were 2237 individuals in the state of Alabama who responded to the "self reported health status" question. Of those responding, the estimated prevalence of Alabama's population who reported that they had fair or poor health status is 19.3%. This prevalence estimate has a standard error of 0.9. The 95% confidence interval for this estimate is (17.6,21.0). Ninety-five percent (95%) of confidence intervals generated under repeated sampling will contain the true population prevalence of Alabama's population who "had fair or poor health status".

Table 1.2 (page 2) shows the estimated prevalence and relevant statistics by sex for each state participant. The same statistics are displayed in Table 1.2 as in Table 1.1, except that Table 1.2 shows the estimated prevalence of self reported fair or poor health status^A for males and females separately. This side by side display of the tables permits each participant to compare estimates for males and females in their respective state.

Table 1.3 (page 3) shows the percent difference between the prevalence estimates of self reported fair or poor health status in 1999 and 2000 survey years. A negative change indicates a decrease in prevalence from 1999 to 2000, while a positive change indicates an increase in prevalence between the two years. This difference gives state health officials a quick method to examine changes that may have occurred in their respective state. Some of these changes may have resulted from legislative actions (or inactions) in the state, programs funded, or programs that came into existence because of results that states may have shown from the BRFSS. Since changes between two years may be attributable to many factors, conclusions drawn from this table should be interpreted with extreme caution.

Sample Size. Sample sizes vary by question. Some estimates may be based on responses from less than 50 respondents. Interpreting estimates that are based on a small number of respondents can mislead the reader into believing that a given finding is much more precise than it actually is. The BRFSS recommends not interpreting percentages where the denominator is based upon fewer than 50 unweighted respondents (sample size).

DEFINITIONS OF TERMS AND RISK FACTORS IN THIS REPORT

AIDS KNOWLEDGE/ATTITUDES

HIV TEST - Respondents, age 18-64, who did not have an HIV test in the past 12 months.

CHANCES OF GETTING HIV - Respondents, age 18-64, who believe that their chances of getting infected with HIV, the virus that causes AIDS are medium or high.

TEEN CONDOM USE - Respondents, age 18-64, who would not encourage sexually active teenagers to use a condom.

BREAST EXAM

BREAST EXAM (1)- Female respondents, age 18 and older, who report they never had a breast exam by a doctor.

BREAST EXAM (2)- Female respondents, age 40 and older, who report they never had a breast exam by a doctor.

BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they did not have a breast exam within the past two years.

CARDIOVASCULAR DISEASE

HEART ATTACK OR MYOCARDIAL INFARCTION - Respondents who report they were told by a doctor they had a heart attack or myocardial infarction.

STROKE - Respondents who report they were told by a doctor they had a stroke.

CERVICAL CANCER

PAP SMEAR (1)- Female respondents, age 18 and older who report they never had a pap smear. Year 2000 Objective 16.12. - Target $\geq 95\%$ who have ever had a pap smear.

PAP SMEAR, WITH UTERINE CERVIX (2)- Female respondents, with uterine cervix, age 18 and older, who report they never had a pap smear.

PAP SMEAR WITHIN 3 YEARS - Female respondents, with uterine cervix, age 18 and older, who report they did not have a pap smear within the last three years.

CLINICAL BREAST EXAM See BREAST EXAM

DIABETES

AGE AT ONSET - Mean age when respondent was told that they have diabetes, among those who report they were told by a doctor that they have diabetes.

DIABETES AWARENESS - Respondents who report they were told by a doctor that they have diabetes.

NOW TAKING INSULIN - Respondents who are now taking insulin, among those who report they were told by a doctor that they have diabetes.

NUMBER OF TIMES SEEN HEALTH PROFESSIONAL - Mean number of times respondents report they have seen a doctor, nurse, or other health professional for their diabetes in the past year, among those who report they were told by a doctor that they have diabetes.

FRUITS AND VEGETABLES

FRUITS AND VEGETABLES - Respondents who report they are not consuming five or more servings of fruits and vegetables a day. Year 2000 Objective 2.6. Target $\geq 50\%$ who are consuming five or more servings of fruits or vegetables a day.

HEALTH INSURANCE

HEALTH INSURANCE (1)- Respondents age 18 and older who report they have no health care coverage.

HEALTH INSURANCE (2)- Respondents age 18-64 years who report they have no health care coverage.

HEALTH STATUS

ACTIVITIES LIMITED - Mean number of days in past 30 days activities were limited among respondents reporting days when physical health or mental health was not good.

MENTAL HEALTH NOT GOOD - Mean number of days in past 30 days mental health was not good as reported by respondents.

PHYSICAL HEALTH NOT GOOD - Mean number of days in past 30 days physical health was not good as reported by respondents.

SELF REPORTED HEALTH STATUS - Respondents who report having a general health status of fair or poor.

IMMUNIZATION

FLU SHOT - Respondents, age 65 and older, who report they did not have a flu shot within the past 12 months. Year 2000 Objective 20.11 - Target $\geq 80\%$ who had a flu shot in the past 12 months.

PNEUMONIA VACCINATION - Respondents, age 65 and older, who report they never had a pneumonia vaccination. Year 2000 Objective 20.11 - Target $\geq 80\%$ who have had a pneumonia vaccination.

MAMMOGRAPHY

MAMMOGRAM (1) - Female respondents age 18 and older, who report that they have never had a mammogram.

MAMMOGRAM (2) - Female respondents, age 40 and older, who report that they have never had a mammogram.

MAMMOGRAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they did not have a mammogram within the past two years.

MAMMOGRAPHY AND BREAST EXAM

MAMMOGRAM & BREAST EXAM - Female respondents, age 40 and older, who report that they have never had a mammogram and a breast exam.

MAMMOGRAM & BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they did not have a mammogram and a breast exam within the last two years. Year 2000 Objective 16.11 - Target $\geq 60\%$ women, age 50 and over who had a mammogram and a breast exam in the preceding 1-2 years.

NUTRITION See FRUITS and VEGETABLES

OVERWEIGHT

OVERWEIGHT: BY BODY MASS INDEX - Respondents with body mass index (BMI) equal to or greater than 25.0. BMI is defined as weight in kilograms divided by height in meters squared (w/h^2). Year 2000 Objective 2.3 - Target $\leq 20\%$ at or above the BMI level. *Note: BRFSS may underestimate the prevalence of overweight because people tend to under report their weight.*

PAP SMEAR See CERVICAL CANCER

PHYSICAL ACTIVITY

NO LEISURE TIME PHYSICAL ACTIVITY - Respondents who report no leisure-time physical activity during the past month. Year 2000 Objective 1.5 - Target $\leq 15\%$ with no-leisure time activity.

REGULAR AND SUSTAINED PHYSICAL ACTIVITY - Respondents who report no physical activity defined as: 5 or more sessions per week, 30 minutes or more per session, regardless of intensity. Year 2000 Objective 1.3 - Target $\geq 30\%$ who engage in physical activity for at least 30 minutes per day.

REGULAR AND VIGOROUS PHYSICAL ACTIVITY - Respondents who report no physical activity or a pair of activities defined as: 3 or more sessions per week, 20 minutes or more per session, at 50% or more capacity. Year 2000 Objective 1.4 - Target $\geq 20\%$ who engage in physical activity 3 or more days per week for 20 or more minutes.

SMOKELESS TOBACCO

EVER USED - Respondents who have ever used or tried smokeless tobacco.

CURRENT USE - Respondents who report that they currently use smokeless tobacco.

SMOKING STATUS

EVER SMOKED - Respondents who have ever smoked 100 cigarettes in their lifetime.

CURRENT SMOKER - Respondents who have smoked 100 cigarettes in their lifetime and reported smoking every day or some days. Year 2000 Objective 3.4 - Target $\leq 15\%$ who smoke cigarettes..

WEIGHT CONTROL

CALORIES AND PHYSICAL ACTIVITY - Respondents with BMI calculated as 25.0

or more (at- risk for weight) who report that they are not either eating fewer calories or less fat and not using physical activity or exercise to lose or maintain weight.

LIMITATIONS OF THE DATA

This report consolidates the most current data on selected risk factors and preventive health measures for states, the District of Columbia and Puerto Rico. The information was obtained from the data files of individual states. Therefore, while all participants use telephone surveys to collect the data, the data in this report vary somewhat with respect to data collection, sampling design and sample size.

A. Data Collection, Sample Design and Sample Size

All states conduct interviews using a computer-assisted telephone interviewing (CATI) software program. All states use probability-based methods to select their sample. The BRFSS standard for sample designs is that sample records must be justifiable as a probability sample of all households with telephones in the state. All participating areas met this criterion in 2000. For data year 2000, one state, Minnesota used the Mitofsky-Waksberg method of sampling. Forty-nine states and the District of Columbia used a disproportionate stratified sample (DSS) design. Puerto Rico used a simple random sample design.

In a few states, a portion of sample records intended for use during one month may have been completed in another month. This deviation should only affect analyses based on monthly, rather than annual data. Illinois used a dual questionnaire or split sample survey design.

Several states did not collect data for all twelve months of the year. Ohio did not collect data January through March, or July through September. Their data include interviews completed in January and February of 2001. Arizona is missing data in July, August, September and December. The District of Columbia did not start the survey until April.

Hawaii did not release telephone numbers in accordance with protocol and may have introduced bias into the survey. More information about the quality of the survey data can be found in the *2000 BRFSS Summary Quality Control Report*.

B. Sample Size, Estimates and Other Limitations of the Data

Although almost all states collect data monthly, the annual number of interviews range from 1717 in the District of Columbia to 8149 in Massachusetts. Estimates for some states are therefore based upon smaller sample sizes, and for rare events (e.g. diabetes), may yield unstable estimates. Some items such as immunizations are affected by seasonality, and estimates for these items may be biased for subpopulations.

As in previous years, California modified the wording of mammography, breast exam and Papanicolaou (PAP) smear questions, and are therefore excluded from those tables in the report.

The physical activity, fruits and vegetables, and weight control questions were asked of only half of Illinois sample due to their dual survey design. These estimates are presented in Tables 10.1-10.3 through 14.1-14.3. More than 7%, (n=309) of Puerto Rico's respondents did not answer the fruit and vegetable questions and are counted as missing values in the calculation of the five or more servings a day estimate in Tables 13.1-13.3.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. Respondents may not remember essential information, a question may

not mean the same thing to different respondents, and some individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their impact on the data.

The data user must make his or her own evaluation of the data. Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. When the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the estimates and/or differences between groups and areas.

TABLE 1.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2237	19.3	0.9	(17.6, 21.0)
ALASKA	2079	10.3	1.1	(8.2, 12.3)
ARIZONA	2671	14.8	1.4	(12.1, 17.6)
ARKANSAS	3000	19.0	0.8	(17.5, 20.5)
CALIFORNIA	3905	16.7	0.8	(15.1, 18.2)
COLORADO	3053	12.7	0.9	(10.9, 14.5)
CONNECTICUT	3909	13.9	0.7	(12.6, 15.2)
DELAWARE	2757	12.4	0.8	(10.8, 14.0)
DISTRICT OF CO	1713	12.3	0.9	(10.5, 14.0)
FLORIDA	5184	15.3	0.6	(14.2, 16.4)
GEORGIA	4103	15.2	0.7	(13.8, 16.6)
HAWAII	6011	12.4	0.6	(11.3, 13.5)
IDAHO	4973	13.1	0.6	(12.0, 14.2)
ILLINOIS	3792	13.0	0.7	(11.7, 14.3)
INDIANA	2916	14.1	0.7	(12.7, 15.4)
IOWA	3607	10.9	0.6	(9.7, 12.0)
KANSAS	4224	12.4	0.6	(11.3, 13.5)
KENTUCKY	6410	21.6	0.7	(20.3, 22.9)
LOUISIANA	5002	16.3	0.6	(15.1, 17.5)
MAINE	4593	14.7	0.9	(12.9, 16.5)
MARYLAND	4583	12.8	0.7	(11.5, 14.1)
MASSACHUSETTS	8127	13.5	0.5	(12.6, 14.4)
MICHIGAN	2598	13.7	0.7	(12.2, 15.1)
MINNESOTA	2851	9.7	0.6	(8.5, 10.9)
MISSISSIPPI	2174	20.2	1.0	(18.3, 22.1)
MISSOURI	4382	15.3	0.7	(13.9, 16.7)
MONTANA	3016	11.3	0.7	(9.9, 12.8)
NEBRASKA	3092	11.3	0.6	(10.1, 12.5)
NEVADA	2100	15.8	1.3	(13.2, 18.3)
NEW HAMPSHIRE	1952	10.1	0.7	(8.6, 11.5)
NEW JERSEY	3773	15.7	0.7	(14.4, 17.1)
NEW MEXICO	3244	17.1	0.7	(15.7, 18.5)
NEW YORK	3349	14.8	0.8	(13.2, 16.3)
NORTH CAROLINA	3010	16.7	0.7	(15.2, 18.1)
NORTH DAKOTA	1913	11.5	0.8	(9.9, 13.0)
OHIO	3246	13.3	0.9	(11.6, 14.9)
OKLAHOMA	3682	15.4	0.7	(14.1, 16.6)
OREGON	3742	16.9	0.7	(15.4, 18.3)
PENNSYLVANIA	3532	14.4	0.7	(13.0, 15.7)
RHODE ISLAND	3538	14.8	0.7	(13.5, 16.2)
SOUTH CAROLINA	3312	15.0	0.7	(13.6, 16.4)
SOUTH DAKOTA	4985	12.1	0.5	(11.2, 13.1)
TENNESSEE	3032	18.3	0.8	(16.8, 19.9)
TEXAS	5004	20.2	0.7	(18.9, 21.5)
UTAH	2890	11.6	0.8	(10.0, 13.1)
VERMONT	3624	11.2	0.6	(10.0, 12.3)
VIRGINIA	1987	13.1	0.9	(11.4, 14.8)
WASHINGTON	3572	11.9	0.6	(10.7, 13.1)
WEST VIRGINIA	2351	25.4	1.0	(23.5, 27.3)
WISCONSIN	2710	12.8	0.7	(11.3, 14.2)
WYOMING	2387	13.1	0.8	(11.6, 14.7)
PUERTO RICO	4201	32.8	0.9	(31.0, 34.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 14.0 RANGE = 9.7-32.8

* HEALTH STATUS IS FAIR OR POOR

April 17, 2001

TABLE 1.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----					-----FEMALES-----				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	835	17.0	1.3	(14.3,	19.6)	1402	21.4	1.2	(19.0,	23.7)
ALASKA	988	10.2	1.6	(7.1,	13.2)	1091	10.4	1.4	(7.7,	13.0)
ARIZONA	1083	13.7	2.3	(9.2,	18.1)	1588	16.0	1.7	(12.6,	19.3)
ARKANSAS	1153	16.4	1.1	(14.1,	18.6)	1847	21.4	1.1	(19.4,	23.5)
CALIFORNIA	1647	15.9	1.3	(13.4,	18.4)	2258	17.5	1.0	(15.5,	19.4)
COLORADO	1290	11.0	1.2	(8.6,	13.3)	1763	14.4	1.3	(11.8,	16.9)
CONNECTICUT	1523	13.5	1.0	(11.4,	15.5)	2386	14.2	0.8	(12.6,	15.8)
DELAWARE	1133	11.4	1.2	(9.0,	13.8)	1624	13.3	1.1	(11.2,	15.4)
DISTRICT OF CO	674	11.7	1.5	(8.9,	14.6)	1039	12.7	1.1	(10.5,	14.9)
FLORIDA	2092	13.5	0.8	(11.9,	15.2)	3092	16.9	0.8	(15.4,	18.5)
GEORGIA	1570	12.9	1.1	(10.7,	15.0)	2533	17.3	0.9	(15.6,	19.1)
HAWAII	2679	12.0	0.9	(10.3,	13.7)	3332	12.8	0.8	(11.3,	14.3)
IDAHO	2117	12.0	0.8	(10.4,	13.6)	2856	14.2	0.8	(12.7,	15.7)
ILLINOIS	1566	10.8	0.9	(8.9,	12.6)	2226	15.1	0.9	(13.3,	16.8)
INDIANA	1206	12.5	1.0	(10.5,	14.5)	1710	15.6	0.9	(13.7,	17.4)
IOWA	1424	9.7	0.9	(8.1,	11.4)	2183	11.9	0.8	(10.4,	13.4)
KANSAS	1641	11.9	0.9	(10.2,	13.6)	2583	12.8	0.7	(11.5,	14.2)
KENTUCKY	2553	20.7	1.0	(18.7,	22.6)	3857	22.5	0.9	(20.7,	24.3)
LOUISIANA	1877	14.3	0.9	(12.5,	16.2)	3125	18.1	0.8	(16.6,	19.5)
MAINE	1883	14.4	1.5	(11.5,	17.4)	2710	14.9	1.1	(12.7,	17.1)
MARYLAND	1776	13.4	1.1	(11.3,	15.5)	2807	12.2	0.8	(10.6,	13.7)
MASSACHUSETTS	3219	13.5	0.7	(12.1,	15.0)	4908	13.4	0.6	(12.3,	14.5)
MICHIGAN	1120	12.7	1.1	(10.6,	14.8)	1478	14.5	1.0	(12.6,	16.5)
MINNESOTA	1257	9.0	0.9	(7.3,	10.8)	1594	10.4	0.8	(8.7,	12.0)
MISSISSIPPI	819	19.0	1.5	(16.0,	21.9)	1355	21.3	1.2	(18.9,	23.7)
MISSOURI	1695	15.3	1.1	(13.1,	17.5)	2687	15.3	0.9	(13.6,	17.0)
MONTANA	1289	10.1	1.1	(8.0,	12.2)	1727	12.5	1.0	(10.6,	14.4)
NEBRASKA	1244	9.7	0.9	(7.9,	11.4)	1848	12.8	0.9	(11.1,	14.5)
NEVADA	1044	14.8	1.8	(11.2,	18.4)	1056	16.7	1.8	(13.2,	20.2)
NEW HAMPSHIRE	788	8.3	1.0	(6.3,	10.4)	1164	11.7	1.1	(9.6,	13.7)
NEW JERSEY	1473	14.0	1.0	(12.0,	16.0)	2300	17.3	0.9	(15.5,	19.1)
NEW MEXICO	1467	14.3	1.0	(12.4,	16.2)	1777	19.8	1.1	(17.7,	21.8)
NEW YORK	1334	14.0	1.2	(11.7,	16.3)	2015	15.4	1.0	(13.4,	17.4)
NORTH CAROLINA	1142	16.1	1.2	(13.8,	18.4)	1868	17.2	0.9	(15.3,	19.0)
NORTH DAKOTA	814	11.2	1.2	(8.9,	13.5)	1099	11.7	1.1	(9.6,	13.9)
OHIO	1272	12.3	1.4	(9.6,	14.9)	1974	14.2	1.1	(12.1,	16.3)
OKLAHOMA	1569	13.7	1.0	(11.8,	15.5)	2113	16.9	0.9	(15.1,	18.7)
OREGON	1494	17.0	1.2	(14.7,	19.4)	2248	16.7	0.9	(14.9,	18.4)
PENNSYLVANIA	1422	12.7	1.1	(10.6,	14.8)	2110	15.9	0.9	(14.1,	17.6)
RHODE ISLAND	1371	13.8	1.1	(11.7,	15.9)	2167	15.8	0.9	(13.9,	17.6)
SOUTH CAROLINA	1321	13.1	1.1	(11.0,	15.1)	1991	16.7	1.0	(14.8,	18.6)
SOUTH DAKOTA	1928	12.5	0.8	(10.9,	14.0)	3057	11.8	0.6	(10.6,	13.0)
TENNESSEE	997	15.6	1.3	(13.2,	18.1)	2035	20.8	1.0	(18.8,	22.7)
TEXAS	2004	19.2	1.0	(17.2,	21.2)	3000	21.1	0.9	(19.4,	22.7)
UTAH	1263	10.8	1.2	(8.5,	13.1)	1627	12.3	1.0	(10.3,	14.3)
VERMONT	1517	10.5	0.8	(8.9,	12.1)	2107	11.8	0.8	(10.3,	13.3)
VIRGINIA	853	11.6	1.2	(9.2,	14.0)	1134	14.5	1.2	(12.1,	16.8)
WASHINGTON	1486	12.0	1.0	(10.2,	13.9)	2086	11.7	0.8	(10.2,	13.3)
WEST VIRGINIA	953	27.2	1.5	(24.2,	30.2)	1398	23.7	1.2	(21.3,	26.1)
WISCONSIN	1219	12.7	1.0	(10.6,	14.7)	1491	12.9	1.0	(10.9,	14.9)
WYOMING	1008	11.8	1.1	(9.6,	14.0)	1379	14.5	1.1	(12.3,	16.6)
PUERTO RICO	1543	26.3	1.4	(23.5,	29.0)	2658	38.7	1.3	(36.3,	41.2)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 13.0
 RANGE = 8.3-27.2

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 15.0
 RANGE = 10.4-38.7

* HEALTH STATUS IS FAIR OR POOR

April 17, 2001

TABLE 1.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	18.4	19.3	0.9
ALASKA	10.7	10.3	-0.4
ARIZONA	8.4	14.8	6.4
ARKANSAS	19.7	19.0	-0.6
CALIFORNIA	15.8	16.7	0.9
COLORADO	11.5	12.7	1.2
CONNECTICUT	11.6	13.9	2.2
DELAWARE	12.4	12.4	-0.0
DISTRICT OF CO	13.0	12.3	-0.8
FLORIDA	15.3	15.3	0.0
GEORGIA	15.0	15.2	0.2
HAWAII	14.3	12.4	-1.9
IDAHO	12.9	13.1	0.3
ILLINOIS	14.7	13.0	-1.7
INDIANA	12.8	14.1	1.3
IOWA	12.1	10.9	-1.3
KANSAS	12.9	12.4	-0.5
KENTUCKY	21.6	21.6	-0.0
LOUISIANA	16.9	16.3	-0.6
MAINE	12.9	14.7	1.8
MARYLAND	14.2	12.8	-1.5
MASSACHUSETTS	11.6	13.5	1.9
MICHIGAN	11.5	13.7	2.2
MINNESOTA	10.0	9.7	-0.3
MISSISSIPPI	20.9	20.2	-0.6
MISSOURI	15.7	15.3	-0.4
MONTANA	10.9	11.3	0.4
NEBRASKA	12.4	11.3	-1.1
NEVADA	13.8	15.8	2.0
NEW HAMPSHIRE	10.6	10.1	-0.6
NEW JERSEY	12.7	15.7	3.1
NEW MEXICO	16.9	17.1	0.3
NEW YORK	13.8	14.8	1.0
NORTH CAROLINA	17.9	16.7	-1.3
NORTH DAKOTA	12.2	11.5	-0.7
OHIO	13.8	13.3	-0.5
OKLAHOMA	17.4	15.4	-2.1
OREGON	13.7	16.9	3.1
PENNSYLVANIA	13.7	14.4	0.6
RHODE ISLAND	12.6	14.8	2.3
SOUTH CAROLINA	13.9	15.0	1.1
SOUTH DAKOTA	13.1	12.1	-0.9
TENNESSEE	19.9	18.3	-1.6
TEXAS	19.2	20.2	1.0
UTAH	10.0	11.6	1.5
VERMONT	9.9	11.2	1.3
VIRGINIA	11.7	13.1	1.4
WASHINGTON	12.0	11.9	-0.1
WEST VIRGINIA	24.0	25.4	1.4
WISCONSIN	11.9	12.8	0.9
WYOMING	11.3	13.1	1.8
PUERTO RICO	33.0	32.8	-0.2

* HEALTH STATUS IS FAIR OR POOR

April 17, 2001

TABLE 2.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2204	3.9	0.2	(3.5, 4.3)
ALASKA	2055	2.8	0.2	(2.4, 3.2)
ARIZONA	2640	3.3	0.3	(2.6, 3.9)
ARKANSAS	2950	4.1	0.2	(3.8, 4.5)
CALIFORNIA	3891	3.5	0.2	(3.1, 3.8)
COLORADO	3035	3.3	0.2	(2.9, 3.7)
CONNECTICUT	3851	3.2	0.1	(3.0, 3.5)
DELAWARE	2736	3.3	0.2	(2.9, 3.6)
DISTRICT OF CO	1684	3.2	0.2	(2.8, 3.6)
FLORIDA	5114	3.4	0.1	(3.2, 3.7)
GEORGIA	3976	3.2	0.2	(2.9, 3.5)
HAWAII	5983	2.6	0.1	(2.3, 2.8)
IDAHO	4910	3.2	0.1	(3.0, 3.4)
ILLINOIS	3769	3.0	0.1	(2.8, 3.3)
INDIANA	2888	3.5	0.2	(3.2, 3.8)
IOWA	3551	2.9	0.1	(2.6, 3.1)
KANSAS	4174	2.9	0.1	(2.7, 3.1)
KENTUCKY	6351	4.0	0.2	(3.7, 4.3)
LOUISIANA	4875	3.2	0.1	(3.0, 3.5)
MAINE	4512	3.5	0.2	(3.1, 3.9)
MARYLAND	4532	3.3	0.2	(3.0, 3.6)
MASSACHUSETTS	8019	3.4	0.1	(3.2, 3.6)
MICHIGAN	2549	3.3	0.2	(3.0, 3.6)
MINNESOTA	2825	2.6	0.1	(2.4, 2.9)
MISSISSIPPI	2111	3.5	0.2	(3.1, 3.9)
MISSOURI	4307	3.2	0.1	(2.9, 3.5)
MONTANA	2979	2.7	0.2	(2.4, 3.0)
NEBRASKA	3070	2.7	0.2	(2.4, 3.0)
NEVADA	2077	3.6	0.3	(3.1, 4.2)
NEW HAMPSHIRE	1923	3.1	0.2	(2.7, 3.4)
NEW JERSEY	3721	3.1	0.1	(2.8, 3.4)
NEW MEXICO	3210	3.5	0.2	(3.2, 3.8)
NEW YORK	3294	3.6	0.2	(3.2, 3.9)
NORTH CAROLINA	2979	3.7	0.2	(3.4, 4.1)
NORTH DAKOTA	1887	2.7	0.2	(2.3, 3.0)
OHIO	3210	3.3	0.2	(2.9, 3.7)
OKLAHOMA	3654	3.1	0.1	(2.8, 3.4)
OREGON	3690	3.7	0.2	(3.4, 4.0)
PENNSYLVANIA	3495	3.2	0.1	(2.9, 3.5)
RHODE ISLAND	3493	3.6	0.2	(3.3, 3.9)
SOUTH CAROLINA	3259	3.3	0.2	(3.0, 3.6)
SOUTH DAKOTA	4918	2.8	0.1	(2.6, 3.1)
TENNESSEE	2936	3.3	0.2	(3.0, 3.6)
TEXAS	4905	3.4	0.1	(3.1, 3.6)
UTAH	2872	3.5	0.2	(3.1, 3.8)
VERMONT	3587	3.1	0.1	(2.8, 3.3)
VIRGINIA	1937	2.7	0.2	(2.4, 3.0)
WASHINGTON	3529	3.2	0.1	(3.0, 3.5)
WEST VIRGINIA	2323	5.2	0.2	(4.7, 5.6)
WISCONSIN	2686	3.6	0.2	(3.2, 3.9)
WYOMING	2367	3.0	0.2	(2.7, 3.3)
PUERTO RICO	4120	3.3	0.2	(3.0, 3.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.3 RANGE = 2.6- 5.2

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

April 17, 2001

TABLE 2.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----			
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	826	3.4	0.3	(2.8, 4.0)	*	1378	4.3	0.3	(3.9, 4.8)
ALASKA	977	2.1	0.2	(1.6, 2.6)	*	1078	3.6	0.3	(2.9, 4.2)
ARIZONA	1080	3.1	0.6	(2.0, 4.2)	*	1560	3.5	0.3	(2.8, 4.1)
ARKANSAS	1144	3.6	0.3	(3.0, 4.1)	*	1806	4.6	0.2	(4.2, 5.1)
CALIFORNIA	1640	3.0	0.3	(2.4, 3.5)	*	2251	4.0	0.2	(3.6, 4.4)
COLORADO	1285	2.9	0.3	(2.4, 3.4)	*	1750	3.7	0.3	(3.2, 4.3)
CONNECTICUT	1501	2.8	0.2	(2.4, 3.2)	*	2350	3.6	0.2	(3.3, 4.0)
DELAWARE	1130	2.8	0.3	(2.3, 3.3)	*	1606	3.7	0.3	(3.1, 4.2)
DISTRICT OF CO	662	2.7	0.3	(2.1, 3.3)	*	1022	3.6	0.3	(3.1, 4.2)
FLORIDA	2073	3.0	0.2	(2.7, 3.4)	*	3041	3.8	0.2	(3.4, 4.2)
GEORGIA	1532	2.7	0.2	(2.2, 3.1)	*	2444	3.7	0.2	(3.3, 4.1)
HAWAII	2670	2.4	0.2	(2.0, 2.7)	*	3313	2.8	0.2	(2.5, 3.0)
IDAHO	2092	2.7	0.2	(2.3, 3.0)	*	2818	3.7	0.2	(3.3, 4.0)
ILLINOIS	1559	2.4	0.2	(2.0, 2.8)	*	2210	3.6	0.2	(3.2, 4.0)
INDIANA	1197	2.9	0.2	(2.4, 3.3)	*	1691	4.0	0.2	(3.6, 4.4)
IOWA	1406	2.4	0.2	(2.1, 2.8)	*	2145	3.3	0.2	(2.9, 3.7)
KANSAS	1631	2.6	0.2	(2.2, 2.9)	*	2543	3.2	0.2	(2.9, 3.5)
KENTUCKY	2536	3.8	0.2	(3.4, 4.3)	*	3815	4.1	0.2	(3.7, 4.5)
LOUISIANA	1838	3.0	0.2	(2.6, 3.4)	*	3037	3.5	0.2	(3.2, 3.8)
MAINE	1857	3.2	0.3	(2.6, 3.7)	*	2655	3.8	0.3	(3.3, 4.3)
MARYLAND	1766	3.4	0.3	(2.9, 4.0)	*	2766	3.2	0.2	(2.8, 3.5)
MASSACHUSETTS	3183	3.1	0.2	(2.8, 3.4)	*	4836	3.7	0.1	(3.4, 4.0)
MICHIGAN	1106	3.0	0.2	(2.6, 3.5)	*	1443	3.6	0.2	(3.2, 4.0)
MINNESOTA	1253	2.3	0.2	(1.9, 2.6)	*	1572	3.0	0.2	(2.6, 3.4)
MISSISSIPPI	799	2.9	0.3	(2.3, 3.5)	*	1312	4.0	0.3	(3.5, 4.5)
MISSOURI	1675	3.3	0.2	(2.8, 3.8)	*	2632	3.2	0.2	(2.8, 3.5)
MONTANA	1276	2.2	0.2	(1.8, 2.5)	*	1703	3.3	0.2	(2.9, 3.7)
NEBRASKA	1240	2.3	0.2	(1.8, 2.7)	*	1830	3.1	0.2	(2.7, 3.5)
NEVADA	1039	3.5	0.4	(2.7, 4.4)	*	1038	3.7	0.4	(3.0, 4.4)
NEW HAMPSHIRE	775	2.3	0.2	(1.8, 2.8)	*	1148	3.8	0.3	(3.3, 4.3)
NEW JERSEY	1458	2.8	0.2	(2.4, 3.2)	*	2263	3.4	0.2	(3.0, 3.7)
NEW MEXICO	1459	2.8	0.2	(2.4, 3.2)	*	1751	4.2	0.2	(3.7, 4.6)
NEW YORK	1313	3.2	0.3	(2.7, 3.7)	*	1981	3.9	0.2	(3.4, 4.3)
NORTH CAROLINA	1141	3.3	0.3	(2.8, 3.8)	*	1838	4.1	0.2	(3.7, 4.5)
NORTH DAKOTA	808	2.3	0.2	(1.8, 2.7)	*	1079	3.1	0.2	(2.6, 3.5)
OHIO	1267	2.7	0.3	(2.1, 3.2)	*	1943	3.9	0.3	(3.4, 4.4)
OKLAHOMA	1562	2.7	0.2	(2.3, 3.1)	*	2092	3.5	0.2	(3.2, 3.9)
OREGON	1479	3.1	0.2	(2.7, 3.6)	*	2211	4.2	0.2	(3.8, 4.6)
PENNSYLVANIA	1408	2.5	0.2	(2.1, 2.9)	*	2087	3.8	0.2	(3.4, 4.2)
RHODE ISLAND	1358	3.2	0.2	(2.8, 3.6)	*	2135	4.0	0.2	(3.6, 4.4)
SOUTH CAROLINA	1310	2.6	0.2	(2.2, 3.0)	*	1949	3.9	0.2	(3.5, 4.3)
SOUTH DAKOTA	1906	2.6	0.2	(2.2, 2.9)	*	3012	3.1	0.1	(2.8, 3.4)
TENNESSEE	974	2.6	0.3	(2.1, 3.1)	*	1962	3.9	0.2	(3.5, 4.3)
TEXAS	1964	3.0	0.2	(2.6, 3.3)	*	2941	3.7	0.2	(3.4, 4.0)
UTAH	1262	2.8	0.3	(2.3, 3.3)	*	1610	4.1	0.3	(3.6, 4.6)
VERMONT	1505	2.7	0.2	(2.4, 3.1)	*	2082	3.4	0.2	(3.0, 3.8)
VIRGINIA	833	2.3	0.2	(1.8, 2.7)	*	1104	3.2	0.2	(2.7, 3.6)
WASHINGTON	1477	2.9	0.2	(2.5, 3.2)	*	2052	3.6	0.2	(3.2, 3.9)
WEST VIRGINIA	944	5.1	0.3	(4.5, 5.8)	*	1379	5.2	0.3	(4.7, 5.7)
WISCONSIN	1211	3.2	0.2	(2.8, 3.7)	*	1475	3.9	0.2	(3.4, 4.4)
WYOMING	1001	2.2	0.2	(1.8, 2.6)	*	1366	3.8	0.2	(3.3, 4.2)
PUERTO RICO	1508	2.7	0.2	(2.3, 3.2)	*	2612	3.8	0.2	(3.5, 4.2)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 2.8
 RANGE = 2.1- 5.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.7
 RANGE = 2.8- 5.2

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

April 17, 2001

TABLE 2.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 MEAN DIFFERENCES FROM 1999 TO 2000 PHYSICAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 MEAN	2000 MEAN	DIFFERENCE
ALABAMA	4.0	3.9	-0.1
ALASKA	2.8	2.8	0.0
ARIZONA	0.9	3.3	2.4
ARKANSAS	4.0	4.1	0.1
CALIFORNIA	3.3	3.5	0.2
COLORADO	3.3	3.3	0.0
CONNECTICUT	3.2	3.2	0.0
DELAWARE	3.3	3.3	-0.1
DISTRICT OF CO	2.7	3.2	0.5
FLORIDA	3.4	3.4	0.1
GEORGIA	2.9	3.2	0.3
HAWAII	2.4	2.6	0.1
IDAHO	3.2	3.2	-0.1
ILLINOIS	3.2	3.0	-0.2
INDIANA	3.1	3.5	0.4
IOWA	3.4	2.9	-0.5
KANSAS	2.6	2.9	0.3
KENTUCKY	4.1	4.0	-0.1
LOUISIANA	3.3	3.2	-0.0
MAINE	3.5	3.5	0.0
MARYLAND	3.1	3.3	0.2
MASSACHUSETTS	3.2	3.4	0.2
MICHIGAN	3.4	3.3	-0.1
MINNESOTA	2.6	2.6	0.0
MISSISSIPPI	3.5	3.5	-0.0
MISSOURI	3.5	3.2	-0.3
MONTANA	3.1	2.7	-0.4
NEBRASKA	3.1	2.7	-0.4
NEVADA	3.5	3.6	0.1
NEW HAMPSHIRE	3.1	3.1	-0.0
NEW JERSEY	3.2	3.1	-0.1
NEW MEXICO	3.7	3.5	-0.2
NEW YORK	3.3	3.6	0.2
NORTH CAROLINA	3.7	3.7	0.0
NORTH DAKOTA	2.9	2.7	-0.2
OHIO	3.1	3.3	0.2
OKLAHOMA	3.0	3.1	0.2
OREGON	3.6	3.7	0.0
PENNSYLVANIA	3.2	3.2	0.0
RHODE ISLAND	3.0	3.6	0.7
SOUTH CAROLINA	3.2	3.3	0.1
SOUTH DAKOTA	3.0	2.8	-0.1
TENNESSEE	3.4	3.3	-0.1
TEXAS	3.3	3.4	0.0
UTAH	3.2	3.5	0.3
VERMONT	2.9	3.1	0.2
VIRGINIA	2.7	2.7	0.0
WASHINGTON	3.5	3.2	-0.3
WEST VIRGINIA	4.0	5.2	1.1
WISCONSIN	3.6	3.6	0.0
WYOMING	2.9	3.0	0.1
PUERTO RICO	3.8	3.3	-0.5

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

April 17, 2001

TABLE 3.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2199	3.4	0.2	(3.1, 3.8)
ALASKA	2045	2.8	0.2	(2.4, 3.2)
ARIZONA	2636	2.7	0.2	(2.3, 3.2)
ARKANSAS	2947	3.3	0.2	(3.0, 3.6)
CALIFORNIA	3895	3.2	0.1	(3.0, 3.5)
COLORADO	3024	3.1	0.2	(2.8, 3.5)
CONNECTICUT	3854	2.9	0.1	(2.7, 3.2)
DELAWARE	2734	3.1	0.2	(2.7, 3.5)
DISTRICT OF CO	1689	3.5	0.2	(3.1, 3.9)
FLORIDA	5122	3.2	0.1	(2.9, 3.4)
GEORGIA	3983	3.4	0.2	(3.1, 3.7)
HAWAII	5973	2.3	0.1	(2.1, 2.5)
IDAHO	4910	3.4	0.1	(3.2, 3.7)
ILLINOIS	3772	2.8	0.1	(2.5, 3.0)
INDIANA	2888	3.4	0.2	(3.1, 3.7)
IOWA	3557	2.5	0.1	(2.2, 2.7)
KANSAS	4180	2.8	0.1	(2.6, 3.0)
KENTUCKY	6329	4.4	0.2	(4.1, 4.7)
LOUISIANA	4917	3.0	0.1	(2.7, 3.2)
MAINE	4512	3.2	0.2	(2.9, 3.6)
MARYLAND	4539	3.1	0.1	(2.8, 3.4)
MASSACHUSETTS	8036	3.3	0.1	(3.1, 3.5)
MICHIGAN	2540	3.6	0.2	(3.3, 4.0)
MINNESOTA	2821	2.7	0.1	(2.4, 2.9)
MISSISSIPPI	2117	3.6	0.2	(3.2, 4.0)
MISSOURI	4329	3.2	0.2	(2.9, 3.5)
MONTANA	2968	2.3	0.2	(2.0, 2.6)
NEBRASKA	3058	2.4	0.1	(2.1, 2.7)
NEVADA	2075	3.4	0.2	(3.0, 3.9)
NEW HAMPSHIRE	1918	2.5	0.2	(2.2, 2.8)
NEW JERSEY	3721	3.2	0.1	(2.9, 3.5)
NEW MEXICO	3221	3.5	0.2	(3.2, 3.8)
NEW YORK	3297	3.2	0.2	(2.9, 3.5)
NORTH CAROLINA	2956	2.9	0.1	(2.6, 3.1)
NORTH DAKOTA	1889	2.6	0.2	(2.3, 2.9)
OHIO	3202	3.4	0.2	(3.0, 3.7)
OKLAHOMA	3654	2.2	0.1	(2.0, 2.5)
OREGON	3691	3.7	0.1	(3.4, 3.9)
PENNSYLVANIA	3497	3.2	0.1	(2.9, 3.4)
RHODE ISLAND	3488	3.4	0.2	(3.1, 3.7)
SOUTH CAROLINA	3266	3.1	0.2	(2.8, 3.4)
SOUTH DAKOTA	4918	2.6	0.1	(2.4, 2.8)
TENNESSEE	2925	2.7	0.1	(2.5, 3.0)
TEXAS	4912	3.6	0.1	(3.3, 3.8)
UTAH	2871	3.4	0.2	(3.1, 3.7)
VERMONT	3582	2.9	0.1	(2.6, 3.1)
VIRGINIA	1946	3.3	0.2	(2.9, 3.7)
WASHINGTON	3512	3.2	0.1	(2.9, 3.4)
WEST VIRGINIA	2337	4.1	0.2	(3.7, 4.5)
WISCONSIN	2681	3.7	0.2	(3.3, 4.0)
WYOMING	2365	3.3	0.2	(2.9, 3.6)
PUERTO RICO	4094	2.1	0.1	(1.9, 2.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.2 RANGE = 2.1- 4.4

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

April 17, 2001

TABLE 3.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				*	-----FEMALES-----				*
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	824	2.8	0.3	(2.3, 3.2)	*	1375	4.1	0.2	(3.6, 4.5)	*
ALASKA	977	1.9	0.2	(1.5, 2.3)	*	1068	3.7	0.3	(3.1, 4.4)	*
ARIZONA	1067	2.3	0.3	(1.7, 2.8)	*	1569	3.2	0.4	(2.5, 3.9)	*
ARKANSAS	1140	2.3	0.2	(1.9, 2.7)	*	1807	4.2	0.2	(3.7, 4.6)	*
CALIFORNIA	1639	2.6	0.2	(2.2, 2.9)	*	2256	3.9	0.2	(3.5, 4.2)	*
COLORADO	1280	2.4	0.2	(2.0, 2.9)	*	1744	3.8	0.3	(3.2, 4.3)	*
CONNECTICUT	1509	2.4	0.2	(2.1, 2.8)	*	2345	3.4	0.2	(3.1, 3.7)	*
DELAWARE	1124	2.4	0.3	(1.8, 2.9)	*	1610	3.8	0.3	(3.3, 4.3)	*
DISTRICT OF CO	662	2.6	0.3	(2.1, 3.2)	*	1027	4.2	0.3	(3.6, 4.7)	*
FLORIDA	2066	2.6	0.2	(2.2, 2.9)	*	3056	3.7	0.2	(3.4, 4.0)	*
GEORGIA	1528	2.7	0.2	(2.3, 3.1)	*	2455	4.0	0.2	(3.6, 4.4)	*
HAWAII	2659	2.0	0.2	(1.6, 2.3)	*	3314	2.7	0.2	(2.4, 3.0)	*
IDAHO	2091	2.7	0.2	(2.4, 3.0)	*	2819	4.2	0.2	(3.8, 4.5)	*
ILLINOIS	1561	2.0	0.2	(1.7, 2.2)	*	2211	3.5	0.2	(3.1, 3.9)	*
INDIANA	1198	2.7	0.2	(2.3, 3.1)	*	1690	4.1	0.2	(3.7, 4.6)	*
IOWA	1403	2.0	0.2	(1.6, 2.4)	*	2154	2.9	0.2	(2.6, 3.2)	*
KANSAS	1634	2.1	0.2	(1.8, 2.5)	*	2546	3.4	0.2	(3.1, 3.7)	*
KENTUCKY	2528	3.4	0.2	(3.0, 3.9)	*	3801	5.3	0.2	(4.8, 5.7)	*
LOUISIANA	1849	2.3	0.2	(1.9, 2.6)	*	3068	3.6	0.2	(3.3, 3.9)	*
MAINE	1858	2.7	0.3	(2.2, 3.3)	*	2654	3.7	0.3	(3.2, 4.2)	*
MARYLAND	1763	2.7	0.2	(2.2, 3.1)	*	2776	3.5	0.2	(3.1, 3.8)	*
MASSACHUSETTS	3180	2.9	0.2	(2.6, 3.2)	*	4856	3.8	0.1	(3.5, 4.0)	*
MICHIGAN	1102	2.8	0.2	(2.3, 3.2)	*	1438	4.5	0.3	(4.0, 4.9)	*
MINNESOTA	1242	2.1	0.2	(1.7, 2.4)	*	1579	3.2	0.2	(2.8, 3.5)	*
MISSISSIPPI	808	2.9	0.3	(2.3, 3.4)	*	1309	4.2	0.3	(3.7, 4.7)	*
MISSOURI	1679	2.7	0.2	(2.3, 3.1)	*	2650	3.8	0.2	(3.4, 4.1)	*
MONTANA	1271	1.6	0.2	(1.2, 2.0)	*	1697	3.0	0.3	(2.5, 3.4)	*
NEBRASKA	1234	1.7	0.2	(1.3, 2.1)	*	1824	3.0	0.2	(2.7, 3.4)	*
NEVADA	1037	2.7	0.3	(2.2, 3.2)	*	1038	4.2	0.4	(3.5, 4.9)	*
NEW HAMPSHIRE	774	1.9	0.2	(1.5, 2.3)	*	1144	3.1	0.2	(2.6, 3.5)	*
NEW JERSEY	1451	2.8	0.2	(2.4, 3.2)	*	2270	3.5	0.2	(3.2, 3.9)	*
NEW MEXICO	1459	2.6	0.2	(2.2, 2.9)	*	1762	4.4	0.2	(3.9, 4.8)	*
NEW YORK	1317	2.5	0.2	(2.1, 2.9)	*	1980	3.8	0.2	(3.4, 4.2)	*
NORTH CAROLINA	1127	2.2	0.2	(1.8, 2.6)	*	1829	3.5	0.2	(3.1, 3.9)	*
NORTH DAKOTA	806	2.3	0.3	(1.8, 2.8)	*	1083	2.9	0.2	(2.5, 3.3)	*
OHIO	1256	2.7	0.3	(2.2, 3.2)	*	1946	4.0	0.3	(3.4, 4.5)	*
OKLAHOMA	1558	1.7	0.2	(1.4, 2.1)	*	2096	2.7	0.2	(2.4, 3.0)	*
OREGON	1475	3.0	0.2	(2.5, 3.4)	*	2216	4.3	0.2	(3.9, 4.7)	*
PENNSYLVANIA	1415	2.3	0.2	(2.0, 2.7)	*	2082	3.9	0.2	(3.5, 4.3)	*
RHODE ISLAND	1361	3.0	0.2	(2.6, 3.5)	*	2127	3.8	0.2	(3.4, 4.2)	*
SOUTH CAROLINA	1311	2.1	0.2	(1.8, 2.5)	*	1955	4.0	0.2	(3.6, 4.5)	*
SOUTH DAKOTA	1908	2.1	0.1	(1.8, 2.4)	*	3010	3.0	0.2	(2.7, 3.3)	*
TENNESSEE	975	1.9	0.2	(1.5, 2.3)	*	1950	3.5	0.2	(3.2, 3.9)	*
TEXAS	1968	3.0	0.2	(2.7, 3.3)	*	2944	4.1	0.2	(3.8, 4.5)	*
UTAH	1258	2.8	0.2	(2.3, 3.2)	*	1613	4.0	0.2	(3.6, 4.5)	*
VERMONT	1500	2.5	0.2	(2.2, 2.9)	*	2082	3.2	0.2	(2.9, 3.5)	*
VIRGINIA	835	2.5	0.2	(2.0, 3.0)	*	1111	4.0	0.3	(3.5, 4.5)	*
WASHINGTON	1464	2.7	0.2	(2.3, 3.1)	*	2048	3.6	0.2	(3.3, 4.0)	*
WEST VIRGINIA	948	3.8	0.3	(3.2, 4.4)	*	1389	4.3	0.3	(3.8, 4.8)	*
WISCONSIN	1205	2.9	0.2	(2.4, 3.3)	*	1476	4.4	0.3	(4.0, 4.9)	*
WYOMING	1004	2.6	0.2	(2.1, 3.0)	*	1361	4.0	0.2	(3.5, 4.4)	*
PUERTO RICO	1509	1.6	0.2	(1.3, 2.0)	*	2585	2.6	0.2	(2.2, 3.0)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 2.5
 RANGE = 1.6- 3.8

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.8
 RANGE = 2.6- 5.3

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

April 17, 2001

TABLE 3.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 MEAN DIFFERENCES FROM 1999 TO 2000 MENTAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 MEAN	2000 MEAN	DIFFERENCE
ALABAMA	3.6	3.4	-0.2
ALASKA	3.2	2.8	-0.4
ARIZONA	0.7	2.7	2.1
ARKANSAS	3.2	3.3	0.1
CALIFORNIA	3.3	3.2	-0.0
COLORADO	3.3	3.1	-0.2
CONNECTICUT	2.8	2.9	0.1
DELAWARE	3.3	3.1	-0.2
DISTRICT OF CO	2.8	3.5	0.7
FLORIDA	2.9	3.2	0.2
GEORGIA	3.1	3.4	0.3
HAWAII	2.2	2.3	0.1
IDAHO	3.1	3.4	0.4
ILLINOIS	2.8	2.8	0.0
INDIANA	3.1	3.4	0.3
IOWA	2.8	2.5	-0.3
KANSAS	2.4	2.8	0.4
KENTUCKY	4.7	4.4	-0.3
LOUISIANA	3.2	3.0	-0.2
MAINE	2.7	3.2	0.6
MARYLAND	3.7	3.1	-0.6
MASSACHUSETTS	2.8	3.3	0.6
MICHIGAN	3.4	3.6	0.3
MINNESOTA	2.8	2.7	-0.2
MISSISSIPPI	3.5	3.6	0.1
MISSOURI	2.9	3.2	0.3
MONTANA	2.7	2.3	-0.4
NEBRASKA	2.8	2.4	-0.4
NEVADA	4.1	3.4	-0.7
NEW HAMPSHIRE	3.3	2.5	-0.8
NEW JERSEY	2.9	3.2	0.3
NEW MEXICO	3.8	3.5	-0.3
NEW YORK	2.9	3.2	0.3
NORTH CAROLINA	3.0	2.9	-0.1
NORTH DAKOTA	2.5	2.6	0.1
OHIO	2.3	3.4	1.1
OKLAHOMA	2.4	2.2	-0.2
OREGON	3.9	3.7	-0.3
PENNSYLVANIA	2.7	3.2	0.5
RHODE ISLAND	2.7	3.4	0.7
SOUTH CAROLINA	2.7	3.1	0.4
SOUTH DAKOTA	2.8	2.6	-0.2
TENNESSEE	3.0	2.7	-0.2
TEXAS	3.4	3.6	0.2
UTAH	3.3	3.4	0.1
VERMONT	2.9	2.9	0.0
VIRGINIA	3.1	3.3	0.2
WASHINGTON	3.1	3.2	0.0
WEST VIRGINIA	3.0	4.1	1.1
WISCONSIN	3.2	3.7	0.5
WYOMING	2.8	3.3	0.5
PUERTO RICO	2.4	2.1	-0.3

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

April 17, 2001

TABLE 4.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1207	4.6	0.3	(4.1, 5.1)
ALASKA	1078	2.8	0.3	(2.2, 3.4)
ARIZONA	1176	3.8	0.4	(3.1, 4.5)
ARKANSAS	1593	4.2	0.2	(3.8, 4.7)
CALIFORNIA	2141	3.9	0.2	(3.4, 4.3)
COLORADO	1654	3.2	0.3	(2.7, 3.6)
CONNECTICUT	2091	3.1	0.2	(2.7, 3.5)
DELAWARE	1255	4.3	0.3	(3.7, 5.0)
DISTRICT OF CO	940	3.6	0.3	(3.0, 4.1)
FLORIDA	2402	4.7	0.2	(4.2, 5.1)
GEORGIA	2112	3.9	0.2	(3.5, 4.4)
HAWAII	2712	3.4	0.2	(3.0, 3.8)
IDAHO	2842	3.3	0.2	(3.0, 3.7)
ILLINOIS	1921	3.4	0.2	(3.0, 3.8)
INDIANA	1609	3.5	0.2	(3.1, 3.9)
IOWA	1741	3.0	0.2	(2.6, 3.4)
KANSAS	2073	3.0	0.2	(2.6, 3.3)
KENTUCKY	2878	6.5	0.3	(6.0, 7.0)
LOUISIANA	2269	4.7	0.2	(4.3, 5.1)
MAINE	2249	4.1	0.3	(3.5, 4.6)
MARYLAND	2392	3.5	0.2	(3.1, 4.0)
MASSACHUSETTS	4529	3.3	0.1	(3.0, 3.5)
MICHIGAN	1467	3.3	0.2	(2.9, 3.7)
MINNESOTA	1410	3.4	0.2	(3.0, 3.8)
MISSISSIPPI	1075	4.1	0.3	(3.5, 4.7)
MISSOURI	2307	3.0	0.2	(2.6, 3.4)
MONTANA	1281	3.5	0.3	(3.0, 4.0)
NEBRASKA	1313	3.9	0.3	(3.3, 4.4)
NEVADA	1176	3.8	0.4	(3.0, 4.5)
NEW HAMPSHIRE	1024	3.1	0.3	(2.6, 3.6)
NEW JERSEY	1955	2.9	0.2	(2.6, 3.3)
NEW MEXICO	1761	3.6	0.2	(3.2, 4.0)
NEW YORK	1836	3.4	0.2	(3.0, 3.8)
NORTH CAROLINA	1456	4.1	0.2	(3.7, 4.6)
NORTH DAKOTA	938	3.0	0.3	(2.5, 3.5)
OHIO	1721	3.4	0.3	(2.9, 3.9)
OKLAHOMA	1432	4.6	0.3	(4.1, 5.1)
OREGON	2255	3.5	0.2	(3.1, 3.8)
PENNSYLVANIA	1906	3.2	0.2	(2.8, 3.5)
RHODE ISLAND	1886	3.4	0.2	(3.0, 3.8)
SOUTH CAROLINA	1614	4.3	0.2	(3.8, 4.8)
SOUTH DAKOTA	2433	2.9	0.2	(2.6, 3.1)
TENNESSEE	1486	3.7	0.2	(3.2, 4.1)
TEXAS	2795	3.7	0.2	(3.4, 4.0)
UTAH	1768	2.8	0.2	(2.4, 3.2)
VERMONT	1907	3.2	0.2	(2.8, 3.6)
VIRGINIA	1091	2.8	0.2	(2.4, 3.3)
WASHINGTON	2135	3.1	0.2	(2.8, 3.5)
WEST VIRGINIA	1209	6.1	0.3	(5.5, 6.8)
WISCONSIN	1620	3.2	0.2	(2.8, 3.7)
WYOMING	1285	3.1	0.2	(2.7, 3.6)
PUERTO RICO	1544	7.0	0.3	(6.3, 7.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.5 RANGE = 2.8- 7.0

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

April 17, 2001

TABLE 4.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED * -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----					-----FEMALES-----				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	381	4.9	0.5	(4.0, 5.9)	*	826	4.4	0.3	(3.7, 5.0)	*
ALASKA	449	2.7	0.4	(1.8, 3.6)	*	629	2.9	0.4	(2.1, 3.7)	*
ARIZONA	413	4.0	0.6	(2.8, 5.2)	*	763	3.7	0.5	(2.8, 4.6)	*
ARKANSAS	524	4.0	0.4	(3.3, 4.8)	*	1069	4.4	0.3	(3.8, 5.0)	*
CALIFORNIA	793	3.5	0.3	(2.9, 4.1)	*	1348	4.2	0.3	(3.6, 4.7)	*
COLORADO	634	2.8	0.3	(2.1, 3.5)	*	1020	3.5	0.4	(2.8, 4.1)	*
CONNECTICUT	752	3.0	0.3	(2.4, 3.6)	*	1339	3.2	0.2	(2.7, 3.6)	*
DELAWARE	437	4.5	0.5	(3.4, 5.5)	*	818	4.3	0.4	(3.5, 5.0)	*
DISTRICT OF CO	328	3.6	0.5	(2.6, 4.5)	*	612	3.6	0.3	(3.0, 4.3)	*
FLORIDA	867	4.9	0.4	(4.2, 5.7)	*	1535	4.5	0.3	(3.9, 5.0)	*
GEORGIA	680	4.1	0.4	(3.3, 4.9)	*	1432	3.8	0.3	(3.3, 4.3)	*
HAWAII	1072	3.6	0.3	(2.9, 4.2)	*	1640	3.3	0.2	(2.8, 3.8)	*
IDAHO	1031	3.4	0.3	(2.9, 4.0)	*	1811	3.2	0.2	(2.8, 3.6)	*
ILLINOIS	673	3.3	0.4	(2.6, 4.0)	*	1248	3.5	0.3	(3.0, 4.0)	*
INDIANA	581	3.6	0.4	(2.9, 4.2)	*	1028	3.4	0.2	(3.0, 3.9)	*
IOWA	604	3.2	0.4	(2.5, 3.9)	*	1137	2.9	0.2	(2.4, 3.3)	*
KANSAS	687	3.2	0.3	(2.6, 3.8)	*	1386	2.8	0.2	(2.4, 3.2)	*
KENTUCKY	1038	7.0	0.4	(6.2, 7.8)	*	1840	6.1	0.3	(5.4, 6.7)	*
LOUISIANA	715	5.0	0.4	(4.2, 5.8)	*	1554	4.5	0.3	(4.0, 5.0)	*
MAINE	820	4.0	0.5	(3.1, 4.8)	*	1429	4.2	0.3	(3.5, 4.8)	*
MARYLAND	847	4.0	0.4	(3.2, 4.8)	*	1545	3.2	0.2	(2.7, 3.7)	*
MASSACHUSETTS	1622	3.2	0.2	(2.7, 3.6)	*	2907	3.3	0.2	(3.0, 3.7)	*
MICHIGAN	562	3.2	0.3	(2.5, 3.8)	*	905	3.4	0.3	(2.9, 3.9)	*
MINNESOTA	538	3.1	0.3	(2.5, 3.7)	*	872	3.6	0.3	(3.1, 4.1)	*
MISSISSIPPI	334	4.5	0.5	(3.4, 5.5)	*	741	3.9	0.3	(3.2, 4.6)	*
MISSOURI	791	3.4	0.4	(2.6, 4.1)	*	1516	2.8	0.2	(2.4, 3.1)	*
MONTANA	464	4.1	0.5	(3.2, 4.9)	*	817	3.1	0.3	(2.6, 3.7)	*
NEBRASKA	432	3.7	0.4	(2.9, 4.5)	*	881	4.0	0.3	(3.3, 4.6)	*
NEVADA	522	4.1	0.7	(2.7, 5.5)	*	654	3.5	0.4	(2.7, 4.3)	*
NEW HAMPSHIRE	362	3.0	0.4	(2.1, 3.8)	*	662	3.3	0.3	(2.6, 3.9)	*
NEW JERSEY	700	2.9	0.3	(2.3, 3.6)	*	1255	2.9	0.2	(2.5, 3.3)	*
NEW MEXICO	675	3.5	0.3	(2.9, 4.1)	*	1086	3.6	0.3	(3.2, 4.1)	*
NEW YORK	671	3.4	0.4	(2.7, 4.1)	*	1165	3.4	0.3	(2.9, 3.9)	*
NORTH CAROLINA	467	4.4	0.4	(3.5, 5.2)	*	989	4.0	0.3	(3.4, 4.5)	*
NORTH DAKOTA	342	3.0	0.4	(2.2, 3.7)	*	596	3.0	0.3	(2.4, 3.6)	*
OHIO	575	3.0	0.4	(2.2, 3.9)	*	1146	3.7	0.3	(3.1, 4.4)	*
OKLAHOMA	483	5.0	0.5	(4.0, 5.9)	*	949	4.3	0.3	(3.8, 4.9)	*
OREGON	782	3.2	0.3	(2.5, 3.8)	*	1473	3.7	0.2	(3.2, 4.1)	*
PENNSYLVANIA	655	3.3	0.3	(2.6, 3.9)	*	1251	3.1	0.2	(2.7, 3.5)	*
RHODE ISLAND	650	3.7	0.4	(3.0, 4.4)	*	1236	3.2	0.2	(2.8, 3.6)	*
SOUTH CAROLINA	536	4.2	0.4	(3.3, 5.0)	*	1078	4.4	0.3	(3.8, 5.0)	*
SOUTH DAKOTA	819	3.2	0.3	(2.7, 3.8)	*	1614	2.6	0.2	(2.2, 2.9)	*
TENNESSEE	397	3.2	0.4	(2.4, 4.0)	*	1089	4.0	0.3	(3.5, 4.5)	*
TEXAS	990	3.5	0.3	(3.0, 4.0)	*	1805	3.8	0.2	(3.4, 4.3)	*
UTAH	668	2.8	0.3	(2.1, 3.4)	*	1100	2.9	0.3	(2.4, 3.4)	*
VERMONT	704	3.3	0.3	(2.8, 3.9)	*	1203	3.1	0.2	(2.6, 3.5)	*
VIRGINIA	421	3.0	0.4	(2.3, 3.8)	*	670	2.7	0.3	(2.1, 3.2)	*
WASHINGTON	786	3.2	0.3	(2.6, 3.7)	*	1349	3.1	0.2	(2.7, 3.5)	*
WEST VIRGINIA	447	7.6	0.6	(6.4, 8.7)	*	762	5.0	0.4	(4.3, 5.8)	*
WISCONSIN	631	3.2	0.3	(2.6, 3.7)	*	989	3.3	0.3	(2.8, 3.9)	*
WYOMING	454	2.9	0.4	(2.2, 3.6)	*	831	3.3	0.3	(2.7, 3.9)	*
PUERTO RICO	458	7.6	0.6	(6.4, 8.8)	*	1086	6.5	0.4	(5.8, 7.3)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 3.5	MEDIAN	= 3.5
	RANGE	= 2.7- 7.6	RANGE	= 2.6- 6.5

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

April 17, 2001

TABLE 4.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 MEAN DIFFERENCES FROM 1999 TO 2000 ACTIVITIES LIMITED *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 MEAN	2000 MEAN	DIFFERENCE
ALABAMA	4.3	4.6	0.3
ALASKA	3.8	2.8	-1.0
ARIZONA	2.9	3.8	0.9
ARKANSAS	4.2	4.2	0.1
CALIFORNIA	4.0	3.9	-0.2
COLORADO	3.1	3.2	0.0
CONNECTICUT	3.5	3.1	-0.4
DELAWARE	3.3	4.3	1.1
DISTRICT OF CO	2.7	3.6	0.9
FLORIDA	4.6	4.7	0.1
GEORGIA	3.4	3.9	0.5
HAWAII	2.8	3.4	0.6
IDAHO	3.2	3.3	0.1
ILLINOIS	3.5	3.4	-0.1
INDIANA	3.5	3.5	-0.0
IOWA	3.1	3.0	-0.1
KANSAS	2.8	3.0	0.1
KENTUCKY	6.2	6.5	0.3
LOUISIANA	4.5	4.7	0.2
MAINE	4.3	4.1	-0.3
MARYLAND	3.2	3.5	0.4
MASSACHUSETTS	3.8	3.3	-0.5
MICHIGAN	3.1	3.3	0.2
MINNESOTA	4.3	3.4	-1.0
MISSISSIPPI	4.2	4.1	-0.0
MISSOURI	3.7	3.0	-0.7
MONTANA	3.4	3.5	0.1
NEBRASKA	3.8	3.9	0.0
NEVADA	3.2	3.8	0.6
NEW HAMPSHIRE	2.8	3.1	0.3
NEW JERSEY	3.6	2.9	-0.7
NEW MEXICO	3.8	3.6	-0.3
NEW YORK	3.3	3.4	0.1
NORTH CAROLINA	4.6	4.1	-0.5
NORTH DAKOTA	2.9	3.0	0.1
OHIO	3.5	3.4	-0.1
OKLAHOMA	4.1	4.6	0.5
OREGON	3.6	3.5	-0.1
PENNSYLVANIA	3.9	3.2	-0.7
RHODE ISLAND	3.7	3.4	-0.2
SOUTH CAROLINA	4.4	4.3	-0.1
SOUTH DAKOTA	3.0	2.9	-0.1
TENNESSEE	3.9	3.7	-0.2
TEXAS	3.7	3.7	-0.0
UTAH	2.7	2.8	0.2
VERMONT	3.0	3.2	0.2
VIRGINIA	3.0	2.8	-0.2
WASHINGTON	3.4	3.1	-0.2
WEST VIRGINIA	7.2	6.1	-1.0
WISCONSIN	2.9	3.2	0.3
WYOMING	3.1	3.1	0.0
PUERTO RICO	6.4	7.0	0.5

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

April 17, 2001

TABLE 5.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

13

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2238	16.1	1.0	(14.2, 18.1)
ALASKA	2076	18.1	1.3	(15.6, 20.6)
ARIZONA	2671	17.1	1.5	(14.2, 20.0)
ARKANSAS	2999	17.2	0.8	(15.6, 18.8)
CALIFORNIA	3905	18.4	0.8	(16.8, 20.0)
COLORADO	3052	13.5	1.0	(11.6, 15.4)
CONNECTICUT	3908	9.3	0.6	(8.1, 10.4)
DELAWARE	2752	8.3	0.7	(6.8, 9.7)
DISTRICT OF CO	1709	11.5	0.9	(9.8, 13.3)
FLORIDA	5193	17.1	0.6	(15.8, 18.3)
GEORGIA	4098	14.9	0.8	(13.4, 16.4)
HAWAII	6010	7.0	0.5	(6.1, 7.9)
IDAHO	4960	17.1	0.6	(15.8, 18.3)
ILLINOIS	3780	10.9	0.7	(9.6, 12.2)
INDIANA	2913	10.4	0.7	(9.1, 11.7)
IOWA	3607	8.9	0.6	(7.7, 10.1)
KANSAS	4227	10.7	0.6	(9.6, 11.8)
KENTUCKY	6405	14.1	0.6	(12.9, 15.3)
LOUISIANA	4987	21.9	0.7	(20.5, 23.3)
MAINE	4599	13.6	0.9	(11.8, 15.4)
MARYLAND	4582	9.9	0.6	(8.8, 11.1)
MASSACHUSETTS	8129	8.4	0.4	(7.6, 9.2)
MICHIGAN	2597	8.5	0.7	(7.1, 9.8)
MINNESOTA	2851	7.1	0.6	(5.9, 8.2)
MISSISSIPPI	2176	19.3	1.0	(17.3, 21.3)
MISSOURI	4380	11.2	0.7	(9.8, 12.5)
MONTANA	3013	14.9	0.9	(13.1, 16.7)
NEBRASKA	3092	9.1	0.6	(7.9, 10.3)
NEVADA	2094	13.6	1.0	(11.6, 15.6)
NEW HAMPSHIRE	1956	9.0	0.8	(7.4, 10.5)
NEW JERSEY	3770	12.9	0.7	(11.5, 14.3)
NEW MEXICO	3241	23.5	0.9	(21.7, 25.2)
NEW YORK	3352	13.1	0.8	(11.7, 14.6)
NORTH CAROLINA	3012	12.7	0.8	(11.3, 14.2)
NORTH DAKOTA	1912	11.8	0.9	(10.1, 13.5)
OHIO	3245	10.2	0.8	(8.6, 11.7)
OKLAHOMA	3678	16.9	0.8	(15.4, 18.3)
OREGON	3740	15.1	0.7	(13.7, 16.5)
PENNSYLVANIA	3526	9.0	0.6	(7.9, 10.2)
RHODE ISLAND	3538	11.1	0.7	(9.7, 12.5)
SOUTH CAROLINA	3304	14.5	0.7	(13.0, 15.9)
SOUTH DAKOTA	4985	10.5	0.5	(9.5, 11.5)
TENNESSEE	3032	11.9	0.7	(10.5, 13.3)
TEXAS	5001	23.0	0.7	(21.5, 24.4)
UTAH	2883	11.1	0.7	(9.6, 12.5)
VERMONT	3624	10.0	0.6	(8.8, 11.2)
VIRGINIA	1992	11.4	0.9	(9.6, 13.1)
WASHINGTON	3578	9.8	0.6	(8.7, 10.9)
WEST VIRGINIA	2348	19.1	1.0	(17.2, 21.0)
WISCONSIN	2715	7.5	0.6	(6.4, 8.6)
WYOMING	2387	17.2	0.9	(15.5, 19.0)
PUERTO RICO	4205	9.0	0.7	(7.7, 10.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 11.8 RANGE = 7.0-23.5
 RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID
 PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

April 17, 2001

TABLE 5.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18 AND OVER* -BY SEX

14

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	835	17.5	1.6	(14.4, 20.6)	*	1403	14.9	1.2	(12.5, 17.3)	
ALASKA	983	19.2	1.9	(15.4, 22.9)	*	1093	16.9	1.7	(13.6, 20.1)	
ARIZONA	1083	16.7	1.7	(13.4, 20.0)	*	1588	17.5	2.3	(12.9, 22.0)	
ARKANSAS	1151	17.1	1.3	(14.6, 19.6)	*	1848	17.3	1.1	(15.2, 19.4)	
CALIFORNIA	1647	19.3	1.2	(16.9, 21.7)	*	2258	17.5	1.0	(15.5, 19.5)	
COLORADO	1291	14.2	1.6	(11.2, 17.3)	*	1761	12.8	1.2	(10.4, 15.2)	
CONNECTICUT	1523	11.2	1.0	(9.2, 13.2)	*	2385	7.5	0.7	(6.1, 8.8)	
DELAWARE	1130	9.5	1.3	(7.1, 12.0)	*	1622	7.1	0.8	(5.6, 8.6)	
DISTRICT OF CO	672	13.5	1.5	(10.5, 16.4)	*	1037	9.8	1.0	(7.8, 11.9)	
FLORIDA	2095	17.6	1.0	(15.6, 19.6)	*	3098	16.6	0.8	(15.0, 18.2)	
GEORGIA	1562	15.6	1.3	(13.1, 18.0)	*	2536	14.3	0.9	(12.6, 16.0)	
HAWAII	2676	8.1	0.7	(6.7, 9.5)	*	3334	5.8	0.5	(4.7, 6.8)	
IDAHO	2108	17.8	1.0	(15.8, 19.7)	*	2852	16.4	0.8	(14.9, 18.0)	
ILLINOIS	1560	11.8	1.0	(9.8, 13.7)	*	2220	10.1	0.8	(8.6, 11.7)	
INDIANA	1202	10.8	1.0	(8.8, 12.8)	*	1711	10.0	0.9	(8.3, 11.7)	
IOWA	1424	10.1	1.0	(8.3, 12.0)	*	2183	7.8	0.8	(6.3, 9.2)	
KANSAS	1642	11.4	0.9	(9.6, 13.2)	*	2585	10.0	0.7	(8.7, 11.4)	
KENTUCKY	2548	15.5	1.0	(13.6, 17.5)	*	3857	12.8	0.7	(11.4, 14.2)	
LOUISIANA	1867	21.7	1.2	(19.4, 23.9)	*	3120	22.2	0.9	(20.5, 23.8)	
MAINE	1885	15.7	1.5	(12.8, 18.6)	*	2714	11.7	1.1	(9.6, 13.9)	
MARYLAND	1775	10.6	1.0	(8.7, 12.5)	*	2807	9.3	0.7	(7.9, 10.7)	
MASSACHUSETTS	3216	10.8	0.7	(9.4, 12.1)	*	4913	6.3	0.5	(5.4, 7.2)	
MICHIGAN	1117	8.9	1.1	(6.8, 11.0)	*	1480	8.1	0.8	(6.4, 9.7)	
MINNESOTA	1255	9.3	1.0	(7.4, 11.2)	*	1596	5.0	0.6	(3.7, 6.2)	
MISSISSIPPI	820	21.2	1.7	(17.8, 24.6)	*	1356	17.6	1.2	(15.3, 19.8)	
MISSOURI	1692	12.9	1.2	(10.6, 15.2)	*	2688	9.6	0.7	(8.1, 11.0)	
MONTANA	1285	15.7	1.5	(12.8, 18.6)	*	1728	14.2	1.1	(12.0, 16.3)	
NEBRASKA	1244	9.6	1.0	(7.6, 11.5)	*	1848	8.7	0.8	(7.3, 10.2)	
NEVADA	1043	14.2	1.4	(11.3, 17.0)	*	1051	13.0	1.4	(10.3, 15.7)	
NEW HAMPSHIRE	791	8.9	1.3	(6.4, 11.3)	*	1165	9.0	1.0	(7.1, 10.9)	
NEW JERSEY	1470	13.6	1.1	(11.4, 15.8)	*	2300	12.3	0.8	(10.6, 13.9)	
NEW MEXICO	1464	24.9	1.4	(22.2, 27.6)	*	1777	22.1	1.2	(19.9, 24.4)	
NEW YORK	1334	16.0	1.3	(13.5, 18.5)	*	2018	10.5	0.8	(8.9, 12.2)	
NORTH CAROLINA	1143	13.5	1.3	(10.9, 16.0)	*	1869	12.1	0.9	(10.4, 13.8)	
NORTH DAKOTA	815	12.8	1.3	(10.2, 15.4)	*	1097	10.8	1.0	(8.8, 12.8)	
OHIO	1267	10.6	1.3	(8.1, 13.0)	*	1978	9.8	1.0	(7.8, 11.8)	
OKLAHOMA	1566	17.7	1.2	(15.3, 20.0)	*	2112	16.1	1.0	(14.2, 18.0)	
OREGON	1495	17.4	1.2	(15.1, 19.6)	*	2245	13.0	0.9	(11.3, 14.6)	
PENNSYLVANIA	1415	10.7	1.0	(8.7, 12.6)	*	2111	7.6	0.7	(6.2, 8.9)	
RHODE ISLAND	1373	15.1	1.2	(12.7, 17.5)	*	2165	7.5	0.8	(6.0, 9.0)	
SOUTH CAROLINA	1317	14.5	1.1	(12.2, 16.7)	*	1987	14.5	0.9	(12.6, 16.3)	
SOUTH DAKOTA	1924	10.8	0.8	(9.3, 12.4)	*	3061	10.1	0.7	(8.8, 11.4)	
TENNESSEE	997	13.9	1.3	(11.5, 16.4)	*	2035	10.0	0.8	(8.5, 11.5)	
TEXAS	2008	24.0	1.2	(21.7, 26.3)	*	2993	22.0	0.9	(20.2, 23.7)	
UTAH	1258	11.3	1.1	(9.2, 13.5)	*	1625	10.9	1.0	(8.9, 12.8)	
VERMONT	1515	12.3	1.0	(10.3, 14.2)	*	2109	7.9	0.8	(6.4, 9.3)	
VIRGINIA	854	10.1	1.3	(7.7, 12.6)	*	1138	12.5	1.3	(10.1, 15.0)	
WASHINGTON	1488	10.0	0.8	(8.3, 11.6)	*	2090	9.6	0.7	(8.1, 11.1)	
WEST VIRGINIA	951	20.9	1.6	(17.8, 23.9)	*	1397	17.5	1.2	(15.2, 19.8)	
WISCONSIN	1218	8.8	0.9	(7.1, 10.5)	*	1497	6.3	0.7	(5.0, 7.6)	
WYOMING	1009	18.4	1.4	(15.6, 21.1)	*	1378	16.1	1.1	(13.9, 18.3)	
PUERTO RICO	1545	10.9	1.2	(8.6, 13.2)	*	2660	7.4	0.8	(5.8, 9.0)	

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 13.5
 RANGE = 8.1-24.9

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 10.8
 RANGE = 5.0-22.2

RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

April 17, 2001

TABLE 5.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO HEALTH INSURANCE, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	15.1	16.1	1.1
ALASKA	23.3	18.1	-5.2
ARIZONA	14.2	17.1	2.9
ARKANSAS	16.1	17.2	1.1
CALIFORNIA	19.3	18.4	-0.9
COLORADO	13.9	13.5	-0.4
CONNECTICUT	10.1	9.3	-0.9
DELAWARE	9.6	8.3	-1.4
DISTRICT OF CO	13.2	11.5	-1.6
FLORIDA	16.2	17.1	0.9
GEORGIA	13.6	14.9	1.3
HAWAII	9.1	7.0	-2.2
IDAHO	16.7	17.1	0.4
ILLINOIS	11.8	10.9	-0.9
INDIANA	12.4	10.4	-2.0
IOWA	8.8	8.9	0.1
KANSAS	10.4	10.7	0.3
KENTUCKY	14.5	14.1	-0.4
LOUISIANA	22.4	21.9	-0.5
MAINE	13.2	13.6	0.4
MARYLAND	9.9	9.9	0.1
MASSACHUSETTS	7.2	8.4	1.2
MICHIGAN	9.8	8.5	-1.3
MINNESOTA	5.8	7.1	1.2
MISSISSIPPI	17.1	19.3	2.2
MISSOURI	10.9	11.2	0.3
MONTANA	17.0	14.9	-2.1
NEBRASKA	8.1	9.1	1.1
NEVADA	18.1	13.6	-4.5
NEW HAMPSHIRE	11.1	9.0	-2.1
NEW JERSEY	11.9	12.9	1.0
NEW MEXICO	22.0	23.5	1.5
NEW YORK	14.6	13.1	-1.4
NORTH CAROLINA	11.5	12.7	1.3
NORTH DAKOTA	11.1	11.8	0.7
OHIO	10.0	10.2	0.1
OKLAHOMA	17.3	16.9	-0.4
OREGON	15.4	15.1	-0.3
PENNSYLVANIA	10.2	9.0	-1.1
RHODE ISLAND	10.6	11.1	0.5
SOUTH CAROLINA	15.6	14.5	-1.2
SOUTH DAKOTA	10.5	10.5	0.0
TENNESSEE	12.4	11.9	-0.5
TEXAS	22.6	23.0	0.4
UTAH	12.4	11.1	-1.3
VERMONT	12.8	10.0	-2.8
VIRGINIA	10.0	11.4	1.4
WASHINGTON	11.9	9.8	-2.2
WEST VIRGINIA	18.1	19.1	1.0
WISCONSIN	10.2	7.5	-2.7
WYOMING	16.9	17.2	0.4
PUERTO RICO	11.7	9.0	-2.6

RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

April 17, 2001

TABLE 6.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1794	19.4	1.2	(17.1, 21.8)
ALASKA	1873	19.1	1.4	(16.4, 21.8)
ARIZONA	2029	20.7	1.8	(17.2, 24.2)
ARKANSAS	2336	20.9	1.0	(19.0, 22.9)
CALIFORNIA	3318	21.3	0.9	(19.5, 23.0)
COLORADO	2551	15.8	1.1	(13.5, 18.0)
CONNECTICUT	3190	10.6	0.7	(9.2, 12.0)
DELAWARE	2235	9.7	0.9	(8.0, 11.4)
DISTRICT OF CO	1452	12.8	1.0	(10.8, 14.8)
FLORIDA	3948	21.6	0.8	(20.0, 23.1)
GEORGIA	3421	16.5	0.9	(14.9, 18.2)
HAWAII	4841	8.3	0.5	(7.2, 9.3)
IDAHO	4162	20.4	0.7	(18.9, 21.8)
ILLINOIS	3168	12.8	0.8	(11.3, 14.3)
INDIANA	2328	12.3	0.8	(10.7, 13.8)
IOWA	2734	10.9	0.8	(9.4, 12.3)
KANSAS	3345	12.9	0.7	(11.6, 14.3)
KENTUCKY	5007	16.6	0.7	(15.2, 18.1)
LOUISIANA	4085	25.6	0.8	(24.0, 27.2)
MAINE	3603	16.3	1.1	(14.1, 18.4)
MARYLAND	3825	11.2	0.7	(9.8, 12.5)
MASSACHUSETTS	6686	9.9	0.5	(8.9, 10.8)
MICHIGAN	2122	10.1	0.8	(8.5, 11.7)
MINNESOTA	2331	8.3	0.7	(7.0, 9.7)
MISSISSIPPI	1772	22.7	1.2	(20.3, 25.0)
MISSOURI	3399	13.4	0.8	(11.8, 15.0)
MONTANA	2382	18.0	1.1	(15.9, 20.2)
NEBRASKA	2326	11.1	0.8	(9.7, 12.6)
NEVADA	1762	16.0	1.2	(13.7, 18.3)
NEW HAMPSHIRE	1626	10.3	0.9	(8.5, 12.1)
NEW JERSEY	3052	15.4	0.8	(13.7, 17.0)
NEW MEXICO	2634	27.7	1.0	(25.6, 29.7)
NEW YORK	2772	15.3	0.9	(13.6, 17.1)
NORTH CAROLINA	2372	15.1	0.9	(13.4, 16.9)
NORTH DAKOTA	1513	14.3	1.0	(12.2, 16.3)
OHIO	2627	12.3	1.0	(10.4, 14.2)
OKLAHOMA	2829	20.1	0.9	(18.4, 21.9)
OREGON	3057	18.1	0.9	(16.4, 19.8)
PENNSYLVANIA	2831	11.1	0.7	(9.7, 12.5)
RHODE ISLAND	2831	13.6	0.9	(11.8, 15.3)
SOUTH CAROLINA	2774	16.9	0.9	(15.2, 18.6)
SOUTH DAKOTA	3895	12.7	0.6	(11.5, 14.0)
TENNESSEE	2483	13.2	0.8	(11.5, 14.8)
TEXAS	4245	26.9	0.8	(25.2, 28.5)
UTAH	2436	12.8	0.8	(11.1, 14.4)
VERMONT	2927	11.7	0.7	(10.2, 13.1)
VIRGINIA	1685	12.9	1.0	(10.9, 15.0)
WASHINGTON	2991	11.5	0.7	(10.2, 12.8)
WEST VIRGINIA	1862	23.5	1.2	(21.2, 25.8)
WISCONSIN	2233	8.9	0.7	(7.6, 10.2)
WYOMING	1983	20.2	1.0	(18.1, 22.2)
PUERTO RICO	3236	10.1	0.8	(8.6, 11.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 13.9 RANGE = 8.3-27.7 YEAR 2000 OBJECTIVE 21.4
 RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID
 PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

April 17, 2001

TABLE 6.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO HEALTH INSURANCE, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	18.0	19.4	1.5
ALASKA	24.8	19.1	-5.7
ARIZONA	16.7	20.7	4.0
ARKANSAS	19.8	20.9	1.1
CALIFORNIA	22.2	21.3	-1.0
COLORADO	16.3	15.8	-0.5
CONNECTICUT	12.3	10.6	-1.7
DELAWARE	11.3	9.7	-1.6
DISTRICT OF CO	15.2	12.8	-2.4
FLORIDA	20.4	21.6	1.1
GEORGIA	15.6	16.5	0.9
HAWAII	10.3	8.3	-2.0
IDAHO	20.1	20.4	0.3
ILLINOIS	13.6	12.8	-0.8
INDIANA	15.1	12.3	-2.8
IOWA	10.9	10.9	-0.1
KANSAS	12.5	12.9	0.4
KENTUCKY	17.3	16.6	-0.7
LOUISIANA	25.8	25.6	-0.2
MAINE	16.1	16.3	0.2
MARYLAND	11.1	11.2	0.1
MASSACHUSETTS	8.3	9.9	1.5
MICHIGAN	11.4	10.1	-1.3
MINNESOTA	6.8	8.3	1.5
MISSISSIPPI	20.3	22.7	2.4
MISSOURI	13.1	13.4	0.3
MONTANA	20.9	18.0	-2.9
NEBRASKA	9.8	11.1	1.4
NEVADA	21.3	16.0	-5.2
NEW HAMPSHIRE	13.2	10.3	-2.8
NEW JERSEY	14.1	15.4	1.3
NEW MEXICO	25.8	27.7	1.8
NEW YORK	17.4	15.3	-2.0
NORTH CAROLINA	13.6	15.1	1.5
NORTH DAKOTA	13.8	14.3	0.4
OHIO	12.2	12.3	0.1
OKLAHOMA	20.5	20.1	-0.3
OREGON	18.6	18.1	-0.5
PENNSYLVANIA	12.5	11.1	-1.4
RHODE ISLAND	12.6	13.6	0.9
SOUTH CAROLINA	18.2	16.9	-1.2
SOUTH DAKOTA	13.0	12.7	-0.3
TENNESSEE	14.2	13.2	-1.1
TEXAS	26.3	26.9	0.6
UTAH	14.2	12.8	-1.4
VERMONT	14.8	11.7	-3.1
VIRGINIA	11.2	12.9	1.8
WASHINGTON	14.1	11.5	-2.6
WEST VIRGINIA	22.3	23.5	1.2
WISCONSIN	12.1	8.9	-3.2
WYOMING	19.9	20.2	0.3
PUERTO RICO	13.2	10.1	-3.0

RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

April 17, 2001

TABLE 7.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2239	7.4	0.6	(6.2, 8.6)
ALASKA	2078	3.8	0.7	(2.5, 5.1)
ARIZONA	2676	5.9	1.1	(3.8, 8.0)
ARKANSAS	2998	6.2	0.5	(5.3, 7.2)
CALIFORNIA	3903	6.8	0.6	(5.6, 7.9)
COLORADO	3057	5.1	0.6	(3.9, 6.3)
CONNECTICUT	3908	5.5	0.4	(4.7, 6.2)
DELAWARE	2754	6.4	0.6	(5.2, 7.7)
DISTRICT OF CO	1714	7.2	0.7	(5.8, 8.6)
FLORIDA	5195	6.9	0.4	(6.1, 7.7)
GEORGIA	4104	6.8	0.5	(5.9, 7.7)
HAWAII	6011	5.2	0.4	(4.5, 5.9)
IDAHO	4973	4.9	0.4	(4.2, 5.6)
ILLINOIS	3791	6.2	0.5	(5.3, 7.1)
INDIANA	2921	6.0	0.5	(5.1, 7.0)
IOWA	3608	6.1	0.5	(5.2, 7.0)
KANSAS	4225	5.9	0.4	(5.1, 6.7)
KENTUCKY	6408	6.5	0.4	(5.7, 7.3)
LOUISIANA	5006	6.7	0.4	(5.9, 7.4)
MAINE	4599	6.0	0.6	(4.8, 7.1)
MARYLAND	4591	6.5	0.5	(5.5, 7.4)
MASSACHUSETTS	8137	5.8	0.3	(5.2, 6.4)
MICHIGAN	2602	7.0	0.5	(6.0, 8.0)
MINNESOTA	2853	4.9	0.5	(4.0, 5.8)
MISSISSIPPI	2181	7.6	0.6	(6.4, 8.8)
MISSOURI	4384	6.7	0.5	(5.6, 7.7)
MONTANA	3019	4.9	0.5	(3.9, 5.8)
NEBRASKA	3092	4.9	0.5	(4.1, 5.8)
NEVADA	2098	6.8	1.0	(4.7, 8.8)
NEW HAMPSHIRE	1956	4.4	0.5	(3.4, 5.4)
NEW JERSEY	3775	5.8	0.4	(5.0, 6.6)
NEW MEXICO	3244	6.5	0.5	(5.5, 7.5)
NEW YORK	3357	6.3	0.5	(5.3, 7.3)
NORTH CAROLINA	3014	6.4	0.5	(5.5, 7.3)
NORTH DAKOTA	1915	5.2	0.6	(4.1, 6.4)
OHIO	3250	6.4	0.6	(5.2, 7.5)
OKLAHOMA	3685	5.5	0.4	(4.7, 6.3)
OREGON	3742	6.0	0.5	(5.1, 6.9)
PENNSYLVANIA	3531	7.1	0.5	(6.1, 8.1)
RHODE ISLAND	3537	6.0	0.5	(5.1, 6.9)
SOUTH CAROLINA	3311	7.1	0.5	(6.1, 8.1)
SOUTH DAKOTA	4995	5.7	0.4	(5.0, 6.4)
TENNESSEE	3037	7.2	0.6	(6.1, 8.3)
TEXAS	5018	6.2	0.4	(5.4, 6.9)
UTAH	2889	5.4	0.6	(4.3, 6.5)
VERMONT	3628	4.4	0.4	(3.7, 5.1)
VIRGINIA	1993	6.2	0.7	(4.7, 7.6)
WASHINGTON	3580	5.5	0.4	(4.7, 6.3)
WEST VIRGINIA	2353	7.6	0.6	(6.5, 8.7)
WISCONSIN	2721	6.1	0.5	(5.1, 7.1)
WYOMING	2390	5.0	0.5	(4.0, 6.0)
PUERTO RICO	4203	8.6	0.5	(7.5, 9.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 6.1 RANGE = 3.8- 8.6

* EVER TOLD YOU HAVE DIABETES

April 17, 2001

TABLE 7.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----				-----FEMALES-----			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	836	7.3	1.0	(5.4, 9.3)	1403	7.4	0.7	(6.0, 8.9)
ALASKA	985	3.6	0.8	(2.1, 5.2)	1093	4.0	1.1	(1.9, 6.1)
ARIZONA	1086	6.0	1.8	(2.4, 9.6)	1590	5.8	1.2	(3.4, 8.3)
ARKANSAS	1151	6.1	0.8	(4.6, 7.7)	1847	6.3	0.6	(5.1, 7.5)
CALIFORNIA	1646	7.4	1.0	(5.5, 9.4)	2257	6.1	0.7	(4.8, 7.5)
COLORADO	1293	6.1	1.0	(4.2, 8.0)	1764	4.1	0.7	(2.7, 5.6)
CONNECTICUT	1524	5.9	0.6	(4.6, 7.1)	2384	5.1	0.5	(4.1, 6.0)
DELAWARE	1130	7.4	1.1	(5.1, 9.6)	1624	5.6	0.7	(4.3, 6.8)
DISTRICT OF CO	674	6.1	1.1	(4.0, 8.2)	1040	8.2	1.0	(6.2, 10.1)
FLORIDA	2095	7.0	0.6	(5.8, 8.3)	3100	6.9	0.5	(5.8, 7.9)
GEORGIA	1567	6.0	0.6	(4.7, 7.3)	2537	7.5	0.7	(6.3, 8.8)
HAWAII	2677	5.8	0.6	(4.7, 6.9)	3334	4.7	0.5	(3.8, 5.6)
IDAHO	2116	4.6	0.5	(3.6, 5.7)	2857	5.1	0.5	(4.2, 6.0)
ILLINOIS	1567	5.6	0.7	(4.2, 7.0)	2224	6.8	0.6	(5.6, 7.9)
INDIANA	1209	5.5	0.7	(4.1, 6.9)	1712	6.5	0.6	(5.3, 7.8)
IOWA	1424	6.1	0.7	(4.7, 7.5)	2184	6.1	0.6	(5.0, 7.3)
KANSAS	1641	6.3	0.7	(5.0, 7.6)	2584	5.5	0.5	(4.6, 6.4)
KENTUCKY	2550	7.0	0.7	(5.7, 8.2)	3858	6.1	0.5	(5.1, 7.1)
LOUISIANA	1877	5.7	0.6	(4.6, 6.9)	3129	7.5	0.5	(6.5, 8.5)
MAINE	1886	6.5	1.0	(4.6, 8.4)	2713	5.5	0.7	(4.1, 6.8)
MARYLAND	1777	7.1	0.8	(5.6, 8.7)	2814	5.8	0.5	(4.8, 6.9)
MASSACHUSETTS	3219	6.0	0.5	(5.0, 6.9)	4918	5.6	0.4	(4.8, 6.4)
MICHIGAN	1119	7.4	0.8	(5.8, 8.9)	1483	6.7	0.7	(5.3, 8.1)
MINNESOTA	1257	4.7	0.7	(3.4, 6.0)	1596	5.1	0.6	(3.9, 6.3)
MISSISSIPPI	823	6.9	0.9	(5.1, 8.6)	1358	8.2	0.8	(6.6, 9.8)
MISSOURI	1695	7.6	0.9	(5.7, 9.4)	2689	5.9	0.6	(4.7, 7.0)
MONTANA	1290	4.4	0.7	(3.0, 5.7)	1729	5.3	0.7	(4.1, 6.6)
NEBRASKA	1244	5.4	0.7	(4.0, 6.7)	1848	4.5	0.6	(3.4, 5.7)
NEVADA	1043	8.7	1.7	(5.4, 12.0)	1055	4.8	1.2	(2.6, 7.1)
NEW HAMPSHIRE	789	4.8	0.8	(3.2, 6.4)	1167	4.0	0.6	(2.9, 5.2)
NEW JERSEY	1473	6.3	0.7	(5.0, 7.7)	2302	5.4	0.5	(4.3, 6.4)
NEW MEXICO	1469	6.3	0.7	(4.9, 7.6)	1775	6.7	0.7	(5.3, 8.2)
NEW YORK	1337	6.6	0.8	(5.0, 8.2)	2020	6.0	0.6	(4.9, 7.2)
NORTH CAROLINA	1143	6.1	0.7	(4.7, 7.5)	1871	6.7	0.6	(5.5, 7.9)
NORTH DAKOTA	814	5.3	0.9	(3.6, 7.0)	1101	5.2	0.8	(3.7, 6.7)
OHIO	1273	6.4	1.0	(4.5, 8.3)	1977	6.3	0.8	(4.8, 7.8)
OKLAHOMA	1568	4.9	0.6	(3.8, 5.9)	2117	6.0	0.6	(4.9, 7.2)
OREGON	1494	6.3	0.8	(4.8, 7.8)	2248	5.8	0.5	(4.7, 6.8)
PENNSYLVANIA	1421	6.8	0.8	(5.3, 8.3)	2110	7.4	0.7	(6.0, 8.9)
RHODE ISLAND	1373	6.9	0.7	(5.4, 8.3)	2164	5.2	0.5	(4.2, 6.3)
SOUTH CAROLINA	1321	7.3	0.8	(5.7, 8.9)	1990	7.0	0.7	(5.7, 8.3)
SOUTH DAKOTA	1933	6.0	0.6	(4.9, 7.1)	3062	5.4	0.5	(4.5, 6.3)
TENNESSEE	999	6.8	0.9	(5.1, 8.5)	2038	7.6	0.7	(6.2, 9.0)
TEXAS	2014	6.3	0.6	(5.1, 7.4)	3004	6.1	0.5	(5.2, 7.1)
UTAH	1263	5.5	0.8	(3.9, 7.1)	1626	5.2	0.8	(3.7, 6.7)
VERMONT	1515	4.7	0.6	(3.6, 5.8)	2113	4.1	0.5	(3.3, 5.0)
VIRGINIA	853	5.4	0.8	(3.7, 7.0)	1140	6.9	1.2	(4.6, 9.2)
WASHINGTON	1488	5.4	0.6	(4.1, 6.6)	2092	5.7	0.5	(4.6, 6.7)
WEST VIRGINIA	955	7.6	0.9	(6.0, 9.3)	1398	7.6	0.7	(6.1, 9.0)
WISCONSIN	1222	5.7	0.7	(4.3, 7.2)	1499	6.5	0.7	(5.0, 7.9)
WYOMING	1010	5.4	0.8	(3.8, 7.0)	1380	4.6	0.6	(3.4, 5.8)
PUERTO RICO	1544	8.7	0.8	(7.1, 10.4)	2659	8.4	0.7	(7.1, 9.7)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 6.1
 RANGE = 3.6- 8.7

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 5.9
 RANGE = 4.0- 8.4

* EVER TOLD YOU HAVE DIABETES

TABLE 7.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000
 DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	7.4	7.4	0.0
ALASKA	3.5	3.8	0.3
ARIZONA	4.3	5.9	1.7
ARKANSAS	6.6	6.2	-0.4
CALIFORNIA	6.1	6.8	0.7
COLORADO	3.8	5.1	1.3
CONNECTICUT	4.3	5.5	1.1
DELAWARE	6.0	6.4	0.4
DISTRICT OF CO	6.5	7.2	0.7
FLORIDA	6.9	6.9	0.1
GEORGIA	5.6	6.8	1.3
HAWAII	5.2	5.2	0.1
IDAHO	4.8	4.9	0.1
ILLINOIS	6.4	6.2	-0.2
INDIANA	6.6	6.0	-0.6
IOWA	5.2	6.1	0.9
KANSAS	5.4	5.9	0.5
KENTUCKY	6.4	6.5	0.1
LOUISIANA	6.1	6.7	0.6
MAINE	5.4	6.0	0.6
MARYLAND	6.8	6.5	-0.3
MASSACHUSETTS	5.0	5.8	0.7
MICHIGAN	5.5	7.0	1.6
MINNESOTA	4.8	4.9	0.1
MISSISSIPPI	7.9	7.6	-0.3
MISSOURI	6.1	6.7	0.6
MONTANA	5.9	4.9	-1.1
NEBRASKA	4.3	4.9	0.6
NEVADA	5.8	6.8	1.0
NEW HAMPSHIRE	4.3	4.4	0.1
NEW JERSEY	5.4	5.8	0.4
NEW MEXICO	5.5	6.5	1.1
NEW YORK	5.7	6.3	0.6
NORTH CAROLINA	6.2	6.4	0.3
NORTH DAKOTA	5.0	5.2	0.3
OHIO	6.1	6.4	0.3
OKLAHOMA	5.8	5.5	-0.3
OREGON	4.6	6.0	1.4
PENNSYLVANIA	6.4	7.1	0.7
RHODE ISLAND	5.3	6.0	0.7
SOUTH CAROLINA	6.4	7.1	0.7
SOUTH DAKOTA	4.9	5.7	0.8
TENNESSEE	6.0	7.2	1.2
TEXAS	6.2	6.2	-0.0
UTAH	4.2	5.4	1.2
VERMONT	4.3	4.4	0.2
VIRGINIA	6.1	6.2	0.0
WASHINGTON	5.2	5.5	0.3
WEST VIRGINIA	7.3	7.6	0.3
WISCONSIN	5.3	6.1	0.8
WYOMING	4.6	5.0	0.4
PUERTO RICO	9.6	8.6	-1.0

* EVER TOLD YOU HAVE DIABETES

April 17, 2001

TABLE 8.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2234	49.2	1.3	(46.8, 51.7)
ALASKA	2072	53.2	1.6	(50.0, 56.3)
ARIZONA	2672	40.3	1.7	(36.9, 43.8)
ARKANSAS	3001	49.6	1.1	(47.5, 51.6)
CALIFORNIA	3903	43.3	0.9	(41.4, 45.1)
COLORADO	3055	46.2	1.3	(43.6, 48.8)
CONNECTICUT	3904	47.8	0.9	(46.0, 49.6)
DELAWARE	2751	46.8	1.3	(44.2, 49.3)
DISTRICT OF CO	1712	42.9	1.4	(40.1, 45.6)
FLORIDA	5183	50.6	0.8	(49.0, 52.2)
GEORGIA	4095	46.3	1.0	(44.3, 48.2)
HAWAII	6002	43.8	0.9	(42.1, 45.6)
IDAHO	4969	46.9	0.8	(45.2, 48.5)
ILLINOIS	3791	44.9	0.9	(43.1, 46.8)
INDIANA	2918	49.6	1.0	(47.5, 51.6)
IOWA	3608	46.3	1.0	(44.4, 48.2)
KANSAS	4209	44.6	0.9	(42.9, 46.4)
KENTUCKY	6404	53.5	0.9	(51.8, 55.2)
LOUISIANA	4991	43.6	0.8	(42.0, 45.1)
MAINE	4587	53.8	1.3	(51.3, 56.2)
MARYLAND	4572	44.1	0.9	(42.2, 45.9)
MASSACHUSETTS	8122	48.1	0.7	(46.8, 49.4)
MICHIGAN	2599	49.6	1.1	(47.4, 51.8)
MINNESOTA	2848	42.9	1.1	(40.8, 45.0)
MISSISSIPPI	2176	44.0	1.2	(41.6, 46.4)
MISSOURI	4383	51.9	1.1	(49.8, 53.9)
MONTANA	3013	48.6	1.2	(46.3, 51.0)
NEBRASKA	3095	45.6	1.0	(43.6, 47.6)
NEVADA	2098	55.8	1.6	(52.7, 58.8)
NEW HAMPSHIRE	1953	55.1	1.3	(52.5, 57.6)
NEW JERSEY	3768	43.4	0.9	(41.6, 45.3)
NEW MEXICO	3243	49.5	1.0	(47.5, 51.4)
NEW YORK	3348	47.2	1.0	(45.2, 49.1)
NORTH CAROLINA	3005	50.4	1.1	(48.3, 52.5)
NORTH DAKOTA	1910	46.7	1.3	(44.2, 49.1)
OHIO	3245	49.4	1.3	(47.0, 51.9)
OKLAHOMA	3680	42.6	0.9	(40.8, 44.5)
OREGON	3737	49.1	0.9	(47.2, 50.9)
PENNSYLVANIA	3530	48.9	1.0	(47.0, 50.8)
RHODE ISLAND	3537	50.1	1.0	(48.2, 52.0)
SOUTH CAROLINA	3303	46.8	1.0	(44.8, 48.8)
SOUTH DAKOTA	4996	46.5	0.8	(44.9, 48.1)
TENNESSEE	3029	47.9	1.1	(45.8, 50.0)
TEXAS	5005	43.8	0.8	(42.2, 45.4)
UTAH	2891	31.2	1.1	(29.0, 33.5)
VERMONT	3618	51.5	1.0	(49.6, 53.4)
VIRGINIA	1986	44.7	1.3	(42.2, 47.3)
WASHINGTON	3572	49.5	0.9	(47.7, 51.4)
WEST VIRGINIA	2350	51.6	1.1	(49.4, 53.8)
WISCONSIN	2715	51.2	1.1	(49.0, 53.3)
WYOMING	2386	48.9	1.1	(46.7, 51.1)
PUERTO RICO	4205	28.4	1.0	(26.5, 30.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 47.5 RANGE = 28.4-55.8

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

April 17, 2001

TABLE 8.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----					-----FEMALES-----				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	836	60.0	2.0	(56.1, 63.8)	*	1398	39.6	1.5	(36.7, 42.6)	*
ALASKA	984	56.2	2.3	(51.7, 60.6)	*	1088	49.9	2.2	(45.5, 54.3)	*
ARIZONA	1084	41.8	2.6	(36.7, 46.9)	*	1588	38.9	2.4	(34.3, 43.6)	*
ARKANSAS	1153	58.2	1.7	(54.9, 61.4)	*	1848	41.8	1.3	(39.3, 44.4)	*
CALIFORNIA	1645	51.4	1.5	(48.5, 54.3)	*	2258	35.3	1.2	(33.0, 37.6)	*
COLORADO	1292	47.2	2.0	(43.3, 51.0)	*	1763	45.3	1.7	(42.0, 48.5)	*
CONNECTICUT	1521	50.1	1.5	(47.2, 52.9)	*	2383	45.7	1.2	(43.4, 47.9)	*
DELAWARE	1132	51.6	2.0	(47.7, 55.5)	*	1619	42.4	1.6	(39.2, 45.5)	*
DISTRICT OF CO	671	45.8	2.2	(41.5, 50.1)	*	1041	40.4	1.8	(37.0, 43.8)	*
FLORIDA	2088	56.6	1.3	(54.1, 59.0)	*	3095	45.2	1.0	(43.2, 47.2)	*
GEORGIA	1564	53.0	1.6	(49.8, 56.1)	*	2531	40.1	1.2	(37.7, 42.5)	*
HAWAII	2678	50.7	1.3	(48.1, 53.3)	*	3324	36.8	1.1	(34.6, 39.0)	*
IDAHO	2111	51.2	1.3	(48.7, 53.7)	*	2858	42.8	1.1	(40.6, 44.9)	*
ILLINOIS	1567	49.9	1.5	(47.0, 52.7)	*	2224	40.4	1.2	(38.1, 42.7)	*
INDIANA	1208	54.2	1.6	(51.1, 57.3)	*	1710	45.3	1.4	(42.6, 48.0)	*
IOWA	1425	54.5	1.5	(51.5, 57.5)	*	2183	38.8	1.2	(36.4, 41.1)	*
KANSAS	1634	52.9	1.4	(50.2, 55.7)	*	2575	36.8	1.1	(34.7, 38.9)	*
KENTUCKY	2548	60.8	1.3	(58.2, 63.4)	*	3856	46.9	1.1	(44.7, 49.1)	*
LOUISIANA	1873	50.0	1.3	(47.4, 52.5)	*	3118	37.8	1.0	(35.9, 39.7)	*
MAINE	1879	59.0	1.9	(55.3, 62.7)	*	2708	48.9	1.6	(45.7, 52.0)	*
MARYLAND	1773	48.4	1.5	(45.4, 51.3)	*	2799	40.1	1.2	(37.8, 42.3)	*
MASSACHUSETTS	3217	49.8	1.1	(47.7, 51.9)	*	4905	46.6	0.9	(44.9, 48.3)	*
MICHIGAN	1119	53.3	1.7	(50.0, 56.7)	*	1480	46.2	1.5	(43.3, 49.1)	*
MINNESOTA	1254	48.2	1.6	(45.1, 51.3)	*	1594	37.9	1.4	(35.1, 40.7)	*
MISSISSIPPI	821	52.4	2.0	(48.5, 56.2)	*	1355	36.5	1.6	(33.5, 39.6)	*
MISSOURI	1695	58.6	1.6	(55.4, 61.8)	*	2688	45.8	1.3	(43.2, 48.4)	*
MONTANA	1288	51.0	1.9	(47.3, 54.6)	*	1725	46.4	1.6	(43.3, 49.4)	*
NEBRASKA	1247	51.7	1.6	(48.6, 54.9)	*	1848	39.9	1.3	(37.3, 42.5)	*
NEVADA	1043	59.7	2.2	(55.4, 64.0)	*	1055	51.7	2.3	(47.3, 56.2)	*
NEW HAMPSHIRE	789	58.5	2.1	(54.4, 62.5)	*	1164	51.8	1.7	(48.5, 55.1)	*
NEW JERSEY	1468	47.6	1.5	(44.7, 50.6)	*	2300	39.6	1.2	(37.3, 41.9)	*
NEW MEXICO	1466	56.7	1.5	(53.8, 59.6)	*	1777	42.6	1.3	(40.1, 45.2)	*
NEW YORK	1330	50.5	1.6	(47.3, 53.6)	*	2018	44.2	1.3	(41.7, 46.7)	*
NORTH CAROLINA	1142	58.1	1.7	(54.8, 61.5)	*	1863	43.4	1.3	(40.9, 45.9)	*
NORTH DAKOTA	814	54.4	2.0	(50.5, 58.2)	*	1096	39.1	1.6	(36.0, 42.3)	*
OHIO	1268	53.8	2.0	(49.9, 57.7)	*	1977	45.5	1.6	(42.4, 48.6)	*
OKLAHOMA	1567	46.8	1.5	(43.9, 49.6)	*	2113	38.8	1.2	(36.4, 41.1)	*
OREGON	1491	54.9	1.4	(52.1, 57.7)	*	2246	43.6	1.2	(41.3, 45.8)	*
PENNSYLVANIA	1420	53.3	1.5	(50.4, 56.3)	*	2110	44.9	1.2	(42.5, 47.2)	*
RHODE ISLAND	1372	53.1	1.6	(50.0, 56.1)	*	2165	47.5	1.2	(45.0, 49.9)	*
SOUTH CAROLINA	1320	54.0	1.6	(50.9, 57.1)	*	1983	40.2	1.3	(37.8, 42.7)	*
SOUTH DAKOTA	1931	52.5	1.3	(50.0, 55.0)	*	3065	40.8	1.0	(38.8, 42.8)	*
TENNESSEE	999	57.2	1.8	(53.7, 60.6)	*	2030	39.5	1.2	(37.2, 41.9)	*
TEXAS	2007	51.6	1.3	(49.1, 54.1)	*	2998	36.5	1.0	(34.5, 38.4)	*
UTAH	1264	38.4	1.8	(34.9, 41.9)	*	1627	24.4	1.4	(21.7, 27.1)	*
VERMONT	1513	56.4	1.5	(53.5, 59.2)	*	2105	46.9	1.2	(44.5, 49.3)	*
VIRGINIA	849	49.8	2.0	(45.9, 53.6)	*	1137	40.0	1.6	(36.8, 43.2)	*
WASHINGTON	1484	55.2	1.4	(52.3, 58.0)	*	2088	44.1	1.2	(41.8, 46.5)	*
WEST VIRGINIA	954	58.6	1.8	(55.1, 62.1)	*	1396	45.3	1.5	(42.4, 48.1)	*
WISCONSIN	1218	54.2	1.6	(51.0, 57.4)	*	1497	48.4	1.5	(45.4, 51.3)	*
WYOMING	1007	52.5	1.7	(49.2, 55.9)	*	1379	45.4	1.5	(42.4, 48.3)	*
PUERTO RICO	1545	38.5	1.7	(35.2, 41.7)	*	2660	19.4	1.0	(17.4, 21.5)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 52.9
 RANGE = 38.4-60.8

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 42.5
 RANGE = 19.4-51.8

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

April 17, 2001

TABLE 8.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	47.7	49.2	1.5
ALASKA	53.0	53.2	0.2
ARIZONA	37.9	40.3	2.4
ARKANSAS	50.3	49.6	-0.7
CALIFORNIA	45.1	43.3	-1.8
COLORADO	48.3	46.2	-2.1
CONNECTICUT	48.2	47.8	-0.4
DELAWARE	49.4	46.8	-2.7
DISTRICT OF CO	36.7	42.9	6.2
FLORIDA	47.4	50.6	3.2
GEORGIA	46.3	46.3	-0.0
HAWAII	41.5	43.8	2.4
IDAHO	45.8	46.9	1.1
ILLINOIS	48.1	44.9	-3.1
INDIANA	49.0	49.6	0.6
IOWA	48.1	46.3	-1.8
KANSAS	43.6	44.6	1.1
KENTUCKY	52.9	53.5	0.6
LOUISIANA	44.2	43.6	-0.6
MAINE	52.8	53.8	1.0
MARYLAND	44.5	44.1	-0.4
MASSACHUSETTS	45.1	48.1	3.0
MICHIGAN	48.7	49.6	0.9
MINNESOTA	39.5	42.9	3.4
MISSISSIPPI	43.2	44.0	0.8
MISSOURI	51.8	51.9	0.1
MONTANA	47.9	48.6	0.7
NEBRASKA	45.3	45.6	0.3
NEVADA	60.6	55.8	-4.8
NEW HAMPSHIRE	51.3	55.1	3.8
NEW JERSEY	41.9	43.4	1.5
NEW MEXICO	47.1	49.5	2.4
NEW YORK	44.9	47.2	2.3
NORTH CAROLINA	48.4	50.4	2.1
NORTH DAKOTA	46.9	46.7	-0.3
OHIO	51.6	49.4	-2.2
OKLAHOMA	43.9	42.6	-1.3
OREGON	49.6	49.1	-0.5
PENNSYLVANIA	46.4	48.9	2.4
RHODE ISLAND	47.8	50.1	2.3
SOUTH CAROLINA	44.0	46.8	2.8
SOUTH DAKOTA	47.6	46.5	-1.1
TENNESSEE	45.6	47.9	2.4
TEXAS	45.7	43.8	-1.9
UTAH	31.2	31.2	0.0
VERMONT	48.4	51.5	3.0
VIRGINIA	46.7	44.7	-2.0
WASHINGTON	49.2	49.5	0.3
WEST VIRGINIA	51.0	51.6	0.6
WISCONSIN	52.3	51.2	-1.1
WYOMING	48.8	48.9	0.1
PUERTO RICO	29.1	28.4	-0.7

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

April 17, 2001

TABLE 9.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2234	25.2	1.1	(23.0, 27.5)
ALASKA	2072	25.0	1.4	(22.3, 27.8)
ARIZONA	2672	18.6	1.6	(15.5, 21.7)
ARKANSAS	3000	25.1	0.9	(23.3, 26.8)
CALIFORNIA	3901	17.2	0.7	(15.8, 18.7)
COLORADO	3055	20.0	1.0	(18.0, 22.0)
CONNECTICUT	3902	19.9	0.8	(18.4, 21.4)
DELAWARE	2751	22.9	1.1	(20.8, 25.0)
DISTRICT OF CO	1712	20.9	1.1	(18.7, 23.1)
FLORIDA	5180	23.2	0.7	(21.8, 24.6)
GEORGIA	4094	23.5	0.9	(21.8, 25.2)
HAWAII	6002	19.7	0.7	(18.2, 21.1)
IDAHO	4967	22.3	0.7	(20.9, 23.7)
ILLINOIS	3790	22.3	0.8	(20.8, 23.9)
INDIANA	2916	26.9	0.9	(25.1, 28.7)
IOWA	3607	23.2	0.8	(21.6, 24.8)
KANSAS	4208	21.1	0.7	(19.6, 22.5)
KENTUCKY	6403	30.5	0.8	(28.9, 32.1)
LOUISIANA	4990	24.1	0.7	(22.7, 25.4)
MAINE	4586	23.8	1.1	(21.6, 25.9)
MARYLAND	4570	20.5	0.8	(19.0, 22.0)
MASSACHUSETTS	8122	20.0	0.5	(18.9, 21.0)
MICHIGAN	2598	24.1	1.0	(22.2, 26.1)
MINNESOTA	2848	19.8	0.9	(18.1, 21.5)
MISSISSIPPI	2175	23.5	1.1	(21.3, 25.7)
MISSOURI	4381	27.2	1.0	(25.3, 29.1)
MONTANA	3011	18.8	0.9	(17.0, 20.6)
NEBRASKA	3093	21.2	0.9	(19.6, 22.9)
NEVADA	2096	29.0	1.4	(26.2, 31.8)
NEW HAMPSHIRE	1952	25.3	1.2	(23.0, 27.7)
NEW JERSEY	3766	21.0	0.8	(19.4, 22.5)
NEW MEXICO	3242	23.6	0.9	(21.9, 25.3)
NEW YORK	3347	21.6	0.8	(20.0, 23.2)
NORTH CAROLINA	3004	26.1	1.0	(24.2, 28.0)
NORTH DAKOTA	1910	23.2	1.1	(21.1, 25.3)
OHIO	3245	26.2	1.1	(24.0, 28.4)
OKLAHOMA	3680	23.3	0.8	(21.7, 24.9)
OREGON	3737	20.8	0.8	(19.3, 22.2)
PENNSYLVANIA	3529	24.3	0.8	(22.7, 26.0)
RHODE ISLAND	3536	23.4	0.8	(21.8, 25.0)
SOUTH CAROLINA	3303	24.9	0.9	(23.1, 26.7)
SOUTH DAKOTA	4996	21.9	0.7	(20.5, 23.2)
TENNESSEE	3026	25.7	0.9	(23.8, 27.5)
TEXAS	5005	21.9	0.7	(20.6, 23.2)
UTAH	2891	12.9	0.8	(11.2, 14.5)
VERMONT	3618	21.5	0.8	(19.9, 23.0)
VIRGINIA	1986	21.4	1.1	(19.3, 23.5)
WASHINGTON	3571	20.7	0.8	(19.2, 22.2)
WEST VIRGINIA	2345	26.1	1.0	(24.2, 28.1)
WISCONSIN	2715	24.1	0.9	(22.3, 26.0)
WYOMING	2385	23.8	1.0	(21.9, 25.6)
PUERTO RICO	4205	13.1	0.8	(11.7, 14.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 23.2 RANGE = 12.9-30.5 YEAR 2000 OBJ. 3.4

* SMOKES EVERYDAY OR SOME DAYS

April 17, 2001

TABLE 9.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	836	28.9	1.9	(25.1, 32.6)	*	1398	22.0	1.3	(19.5, 24.5)	
ALASKA	984	26.8	2.1	(22.7, 30.9)	*	1088	23.1	1.8	(19.5, 26.7)	
ARIZONA	1084	18.3	2.3	(13.9, 22.7)	*	1588	19.0	2.4	(14.4, 23.6)	
ARKANSAS	1153	26.1	1.5	(23.3, 29.0)	*	1847	24.1	1.1	(21.9, 26.3)	
CALIFORNIA	1645	20.1	1.2	(17.7, 22.5)	*	2256	14.4	0.8	(12.8, 16.0)	
COLORADO	1292	19.5	1.5	(16.5, 22.4)	*	1763	20.5	1.4	(17.8, 23.2)	
CONNECTICUT	1520	20.4	1.2	(18.0, 22.8)	*	2382	19.4	0.9	(17.6, 21.2)	
DELAWARE	1132	25.7	1.7	(22.3, 29.1)	*	1619	20.3	1.3	(17.7, 22.8)	
DISTRICT OF CO	671	22.1	1.8	(18.5, 25.6)	*	1041	19.9	1.4	(17.2, 22.6)	
FLORIDA	2086	24.5	1.1	(22.3, 26.6)	*	3094	22.1	0.9	(20.3, 23.8)	
GEORGIA	1563	26.3	1.4	(23.6, 29.0)	*	2531	20.9	1.0	(18.9, 23.0)	
HAWAII	2678	22.9	1.1	(20.7, 25.1)	*	3324	16.4	0.9	(14.6, 18.1)	
IDAHO	2110	22.8	1.1	(20.7, 24.8)	*	2857	21.8	0.9	(20.0, 23.7)	
ILLINOIS	1566	24.9	1.3	(22.4, 27.4)	*	2224	20.0	1.0	(18.1, 21.9)	
INDIANA	1208	28.4	1.4	(25.6, 31.2)	*	1708	25.5	1.2	(23.2, 27.9)	
IOWA	1425	25.7	1.3	(23.1, 28.3)	*	2182	20.9	1.1	(18.8, 22.9)	
KANSAS	1634	24.2	1.2	(21.9, 26.5)	*	2574	18.1	0.8	(16.5, 19.8)	
KENTUCKY	2547	33.4	1.3	(30.8, 35.9)	*	3856	27.9	1.0	(25.9, 29.9)	
LOUISIANA	1872	26.8	1.2	(24.5, 29.0)	*	3118	21.7	0.8	(20.1, 23.3)	
MAINE	1878	24.6	1.7	(21.2, 28.0)	*	2708	23.1	1.4	(20.4, 25.7)	
MARYLAND	1771	22.0	1.2	(19.6, 24.4)	*	2799	19.1	0.9	(17.3, 20.9)	
MASSACHUSETTS	3217	20.2	0.9	(18.5, 21.8)	*	4905	19.7	0.7	(18.4, 21.1)	
MICHIGAN	1119	26.0	1.5	(23.0, 28.9)	*	1479	22.4	1.3	(19.9, 25.0)	
MINNESOTA	1254	20.7	1.3	(18.2, 23.2)	*	1594	18.9	1.1	(16.7, 21.1)	
MISSISSIPPI	821	25.2	1.7	(21.8, 28.6)	*	1354	21.9	1.4	(19.1, 24.7)	
MISSOURI	1694	30.1	1.6	(27.1, 33.2)	*	2687	24.5	1.1	(22.3, 26.7)	
MONTANA	1287	18.0	1.4	(15.2, 20.7)	*	1724	19.7	1.2	(17.3, 22.1)	
NEBRASKA	1246	21.9	1.3	(19.3, 24.4)	*	1847	20.6	1.1	(18.4, 22.8)	
NEVADA	1041	28.6	1.8	(25.1, 32.1)	*	1055	29.4	2.2	(25.2, 33.7)	
NEW HAMPSHIRE	788	27.0	1.9	(23.3, 30.8)	*	1164	23.7	1.5	(20.9, 26.6)	
NEW JERSEY	1468	23.6	1.3	(21.1, 26.2)	*	2298	18.6	0.9	(16.8, 20.3)	
NEW MEXICO	1465	26.2	1.4	(23.5, 28.8)	*	1777	21.2	1.1	(19.0, 23.3)	
NEW YORK	1330	22.6	1.3	(20.0, 25.2)	*	2017	20.7	1.0	(18.7, 22.6)	
NORTH CAROLINA	1142	28.4	1.6	(25.3, 31.6)	*	1862	23.9	1.2	(21.7, 26.2)	
NORTH DAKOTA	814	25.7	1.7	(22.4, 29.0)	*	1096	20.8	1.3	(18.2, 23.4)	
OHIO	1268	26.6	1.8	(23.2, 30.0)	*	1977	25.9	1.4	(23.1, 28.7)	
OKLAHOMA	1567	23.7	1.2	(21.3, 26.1)	*	2113	22.9	1.1	(20.9, 25.0)	
OREGON	1491	22.2	1.2	(19.9, 24.6)	*	2246	19.4	1.0	(17.5, 21.2)	
PENNSYLVANIA	1420	25.5	1.4	(22.8, 28.1)	*	2109	23.3	1.0	(21.3, 25.3)	
RHODE ISLAND	1372	23.8	1.3	(21.2, 26.4)	*	2164	23.0	1.1	(21.0, 25.1)	
SOUTH CAROLINA	1320	28.7	1.5	(25.7, 31.6)	*	1983	21.5	1.1	(19.4, 23.6)	
SOUTH DAKOTA	1931	22.5	1.1	(20.4, 24.6)	*	3065	21.3	0.9	(19.5, 23.0)	
TENNESSEE	998	27.6	1.6	(24.6, 30.7)	*	2028	23.9	1.1	(21.8, 26.0)	
TEXAS	2007	25.2	1.1	(23.1, 27.4)	*	2998	18.7	0.8	(17.2, 20.3)	
UTAH	1264	14.5	1.3	(12.0, 17.0)	*	1627	11.3	1.0	(9.3, 13.3)	
VERMONT	1513	21.7	1.3	(19.2, 24.2)	*	2105	21.2	1.0	(19.2, 23.2)	
VIRGINIA	849	24.2	1.7	(20.8, 27.6)	*	1137	18.8	1.3	(16.3, 21.3)	
WASHINGTON	1484	21.8	1.2	(19.4, 24.2)	*	2087	19.7	1.0	(17.8, 21.6)	
WEST VIRGINIA	952	27.7	1.6	(24.6, 30.8)	*	1393	24.7	1.2	(22.3, 27.1)	
WISCONSIN	1218	24.4	1.4	(21.6, 27.2)	*	1497	23.9	1.2	(21.4, 26.3)	
WYOMING	1007	23.2	1.4	(20.5, 26.0)	*	1378	24.3	1.3	(21.8, 26.8)	
PUERTO RICO	1545	16.8	1.3	(14.2, 19.3)	*	2660	9.9	0.8	(8.3, 11.6)	

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 24.4
 RANGE = 14.5-33.4

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 21.2
 RANGE = 9.9-29.4

* SMOKES EVERYDAY OR SOME DAYS

April 17, 2001

TABLE 9.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	23.5	25.2	1.8
ALASKA	27.3	25.0	-2.3
ARIZONA	20.1	18.6	-1.4
ARKANSAS	27.2	25.1	-2.2
CALIFORNIA	18.7	17.2	-1.5
COLORADO	22.5	20.0	-2.5
CONNECTICUT	22.8	19.9	-2.9
DELAWARE	25.5	22.9	-2.6
DISTRICT OF CO	20.6	20.9	0.3
FLORIDA	20.6	23.2	2.6
GEORGIA	23.8	23.5	-0.2
HAWAII	18.5	19.7	1.2
IDAHO	21.5	22.3	0.8
ILLINOIS	24.2	22.3	-1.9
INDIANA	27.0	26.9	-0.1
IOWA	23.5	23.2	-0.3
KANSAS	21.0	21.1	0.0
KENTUCKY	29.7	30.5	0.8
LOUISIANA	23.5	24.1	0.6
MAINE	23.3	23.8	0.5
MARYLAND	20.3	20.5	0.2
MASSACHUSETTS	19.3	20.0	0.7
MICHIGAN	25.1	24.1	-1.0
MINNESOTA	19.5	19.8	0.3
MISSISSIPPI	22.9	23.5	0.5
MISSOURI	27.1	27.2	0.1
MONTANA	20.2	18.8	-1.4
NEBRASKA	23.2	21.2	-2.0
NEVADA	31.5	29.0	-2.5
NEW HAMPSHIRE	22.3	25.3	3.0
NEW JERSEY	20.6	21.0	0.3
NEW MEXICO	22.5	23.6	1.1
NEW YORK	21.8	21.6	-0.2
NORTH CAROLINA	25.1	26.1	0.9
NORTH DAKOTA	22.1	23.2	1.1
OHIO	27.6	26.2	-1.4
OKLAHOMA	25.2	23.3	-1.8
OREGON	21.4	20.8	-0.7
PENNSYLVANIA	23.1	24.3	1.2
RHODE ISLAND	22.3	23.4	1.1
SOUTH CAROLINA	23.6	24.9	1.3
SOUTH DAKOTA	22.5	21.9	-0.6
TENNESSEE	24.8	25.7	0.8
TEXAS	22.4	21.9	-0.5
UTAH	14.0	12.9	-1.1
VERMONT	21.7	21.5	-0.2
VIRGINIA	21.4	21.4	0.0
WASHINGTON	22.4	20.7	-1.7
WEST VIRGINIA	27.1	26.1	-1.0
WISCONSIN	23.7	24.1	0.4
WYOMING	23.9	23.8	-0.1
PUERTO RICO	13.7	13.1	-0.6

* SMOKES EVERYDAY OR SOME DAYS

April 17, 2001

TABLE 10.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO LEISURE TIME PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2241	31.6	1.2	(29.3, 33.9)
ALASKA	2081	20.0	1.2	(17.6, 22.4)
ARIZONA	2678	34.2	1.7	(30.8, 37.6)
ARKANSAS	3005	28.1	0.9	(26.3, 29.8)
CALIFORNIA	3907	26.5	0.9	(24.7, 28.2)
COLORADO	3057	19.8	1.1	(17.7, 21.8)
CONNECTICUT	3913	25.2	0.8	(23.6, 26.7)
DELAWARE	2757	28.0	1.1	(25.8, 30.3)
DISTRICT OF CO	1715	20.8	1.1	(18.5, 23.0)
FLORIDA	5200	28.8	0.7	(27.4, 30.3)
GEORGIA	4108	29.0	0.9	(27.2, 30.7)
HAWAII	6016	23.2	0.8	(21.7, 24.7)
IDAHO	4974	19.8	0.6	(18.5, 21.0)
ILLINOIS	1853	30.9	1.3	(28.4, 33.4)
INDIANA	2921	25.4	0.9	(23.6, 27.1)
IOWA	3610	27.3	0.9	(25.6, 28.9)
KANSAS	4232	30.4	0.8	(28.8, 32.0)
KENTUCKY	6415	41.1	0.9	(39.4, 42.7)
LOUISIANA	5009	36.2	0.8	(34.7, 37.8)
MAINE	4600	27.2	1.1	(25.0, 29.4)
MARYLAND	4589	24.2	0.8	(22.7, 25.8)
MASSACHUSETTS	8144	24.6	0.6	(23.5, 25.7)
MICHIGAN	2604	22.9	0.9	(21.1, 24.7)
MINNESOTA	2850	24.8	1.0	(22.9, 26.7)
MISSISSIPPI	2181	33.3	1.2	(31.0, 35.7)
MISSOURI	4385	28.8	0.9	(27.0, 30.6)
MONTANA	3018	23.3	1.0	(21.3, 25.3)
NEBRASKA	3091	29.6	1.0	(27.7, 31.5)
NEVADA	2101	25.0	1.4	(22.2, 27.7)
NEW HAMPSHIRE	1956	26.7	1.2	(24.4, 28.9)
NEW JERSEY	3776	28.6	0.9	(26.9, 30.3)
NEW MEXICO	3247	24.4	0.8	(22.7, 26.0)
NEW YORK	3357	29.4	1.0	(27.5, 31.2)
NORTH CAROLINA	3016	30.4	1.0	(28.6, 32.3)
NORTH DAKOTA	1916	24.3	1.1	(22.2, 26.4)
OHIO	3250	31.3	1.2	(29.0, 33.6)
OKLAHOMA	3686	34.4	0.9	(32.6, 36.2)
OREGON	3743	20.1	0.8	(18.6, 21.5)
PENNSYLVANIA	3535	23.0	0.8	(21.4, 24.6)
RHODE ISLAND	3541	27.5	0.9	(25.8, 29.3)
SOUTH CAROLINA	3312	28.1	0.9	(26.3, 29.9)
SOUTH DAKOTA	4994	26.7	0.7	(25.3, 28.0)
TENNESSEE	3036	32.7	1.0	(30.8, 34.6)
TEXAS	5016	28.6	0.7	(27.1, 30.0)
UTAH	2890	15.5	0.9	(13.8, 17.2)
VERMONT	3629	23.2	0.8	(21.6, 24.8)
VIRGINIA	1993	25.0	1.1	(22.9, 27.2)
WASHINGTON	3578	16.9	0.7	(15.5, 18.2)
WEST VIRGINIA	2353	33.6	1.1	(31.5, 35.7)
WISCONSIN	2717	22.1	0.9	(20.4, 23.9)
WYOMING	2389	22.6	1.0	(20.7, 24.5)
PUERTO RICO	4206	54.1	1.1	(51.9, 56.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 26.9 RANGE = 15.5-54.1

April 17, 2001

TABLE 10.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO LEISURE TIME PHYSICAL ACTIVITY* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----					-----FEMALES-----				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	836	26.8	1.9	(23.1, 30.5)	*	1405	35.9	1.5	(33.0, 38.7)	*
ALASKA	987	20.0	1.8	(16.6, 23.5)	*	1094	20.0	1.7	(16.6, 23.4)	*
ARIZONA	1085	31.5	2.4	(26.7, 36.2)	*	1593	36.8	2.4	(32.1, 41.5)	*
ARKANSAS	1155	23.7	1.3	(21.1, 26.3)	*	1850	32.0	1.2	(29.6, 34.4)	*
CALIFORNIA	1647	24.0	1.4	(21.2, 26.7)	*	2260	29.0	1.2	(26.7, 31.3)	*
COLORADO	1294	16.3	1.5	(13.4, 19.2)	*	1763	23.1	1.5	(20.1, 26.0)	*
CONNECTICUT	1526	22.5	1.2	(20.1, 24.8)	*	2387	27.7	1.1	(25.6, 29.7)	*
DELAWARE	1134	23.2	1.6	(20.0, 26.4)	*	1623	32.5	1.5	(29.5, 35.5)	*
DISTRICT OF CO	674	17.4	1.8	(14.0, 20.9)	*	1041	23.6	1.5	(20.7, 26.6)	*
FLORIDA	2098	25.5	1.1	(23.3, 27.7)	*	3102	31.8	1.0	(29.9, 33.7)	*
GEORGIA	1569	23.2	1.3	(20.6, 25.9)	*	2539	34.3	1.2	(32.0, 36.5)	*
HAWAII	2681	18.3	1.0	(16.4, 20.3)	*	3335	28.2	1.1	(26.1, 30.3)	*
IDAHO	2117	18.2	0.9	(16.4, 20.0)	*	2857	21.3	0.9	(19.6, 23.1)	*
ILLINOIS	785	27.4	1.9	(23.7, 31.2)	*	1068	34.2	1.7	(30.9, 37.4)	*
INDIANA	1209	21.2	1.3	(18.7, 23.7)	*	1712	29.2	1.3	(26.7, 31.6)	*
IOWA	1424	26.8	1.3	(24.2, 29.4)	*	2186	27.7	1.1	(25.6, 29.8)	*
KANSAS	1643	28.5	1.3	(26.0, 31.0)	*	2589	32.2	1.0	(30.2, 34.3)	*
KENTUCKY	2553	40.0	1.3	(37.4, 42.5)	*	3862	42.1	1.1	(39.9, 44.3)	*
LOUISIANA	1878	34.1	1.2	(31.7, 36.5)	*	3131	38.1	1.0	(36.2, 40.0)	*
MAINE	1886	26.4	1.7	(23.0, 29.8)	*	2714	27.9	1.5	(25.1, 30.8)	*
MARYLAND	1778	20.4	1.2	(18.0, 22.8)	*	2811	27.8	1.1	(25.7, 29.8)	*
MASSACHUSETTS	3225	23.0	0.9	(21.3, 24.8)	*	4919	26.0	0.8	(24.5, 27.5)	*
MICHIGAN	1120	19.5	1.3	(16.9, 22.1)	*	1484	25.9	1.3	(23.5, 28.4)	*
MINNESOTA	1255	23.2	1.4	(20.5, 25.9)	*	1595	26.3	1.3	(23.7, 28.9)	*
MISSISSIPPI	822	29.0	1.8	(25.5, 32.4)	*	1359	37.2	1.6	(34.1, 40.4)	*
MISSOURI	1695	28.4	1.5	(25.6, 31.3)	*	2690	29.2	1.2	(26.9, 31.5)	*
MONTANA	1290	21.7	1.4	(18.9, 24.6)	*	1728	24.8	1.5	(21.9, 27.7)	*
NEBRASKA	1247	28.1	1.5	(25.3, 31.0)	*	1844	31.0	1.2	(28.5, 33.4)	*
NEVADA	1046	24.5	2.0	(20.5, 28.5)	*	1055	25.4	2.0	(21.6, 29.2)	*
NEW HAMPSHIRE	789	25.6	1.9	(21.9, 29.2)	*	1167	27.7	1.4	(24.9, 30.5)	*
NEW JERSEY	1475	26.5	1.4	(23.9, 29.2)	*	2301	30.6	1.1	(28.4, 32.7)	*
NEW MEXICO	1468	22.9	1.2	(20.5, 25.3)	*	1779	25.8	1.2	(23.5, 28.0)	*
NEW YORK	1338	26.9	1.5	(23.9, 29.8)	*	2019	31.6	1.2	(29.2, 34.0)	*
NORTH CAROLINA	1145	27.3	1.5	(24.4, 30.2)	*	1871	33.3	1.2	(30.9, 35.7)	*
NORTH DAKOTA	816	23.8	1.6	(20.7, 26.9)	*	1100	24.8	1.4	(22.0, 27.6)	*
OHIO	1273	30.1	1.8	(26.5, 33.7)	*	1977	32.3	1.5	(29.3, 35.2)	*
OKLAHOMA	1569	34.0	1.4	(31.2, 36.8)	*	2117	34.8	1.2	(32.5, 37.1)	*
OREGON	1495	19.0	1.1	(16.8, 21.2)	*	2248	21.1	1.0	(19.1, 23.0)	*
PENNSYLVANIA	1423	20.8	1.2	(18.4, 23.2)	*	2112	24.9	1.1	(22.8, 27.0)	*
RHODE ISLAND	1373	24.1	1.4	(21.4, 26.8)	*	2168	30.6	1.2	(28.3, 32.9)	*
SOUTH CAROLINA	1322	25.6	1.4	(22.9, 28.3)	*	1990	30.4	1.2	(28.1, 32.7)	*
SOUTH DAKOTA	1933	26.5	1.1	(24.4, 28.6)	*	3061	26.8	0.9	(25.0, 28.5)	*
TENNESSEE	999	30.0	1.6	(26.9, 33.2)	*	2037	35.1	1.2	(32.8, 37.4)	*
TEXAS	2012	26.5	1.1	(24.3, 28.7)	*	3004	30.5	1.0	(28.6, 32.4)	*
UTAH	1264	13.7	1.2	(11.4, 16.0)	*	1626	17.3	1.2	(14.9, 19.7)	*
VERMONT	1516	23.2	1.3	(20.7, 25.6)	*	2113	23.3	1.0	(21.3, 25.3)	*
VIRGINIA	854	21.9	1.6	(18.7, 25.1)	*	1139	27.9	1.5	(25.0, 30.8)	*
WASHINGTON	1488	15.8	1.0	(13.8, 17.8)	*	2090	17.9	0.9	(16.1, 19.7)	*
WEST VIRGINIA	954	32.0	1.6	(28.8, 35.2)	*	1399	35.0	1.4	(32.3, 37.8)	*
WISCONSIN	1219	19.1	1.2	(16.8, 21.5)	*	1498	24.9	1.3	(22.4, 27.5)	*
WYOMING	1010	22.0	1.4	(19.2, 24.8)	*	1379	23.2	1.3	(20.7, 25.7)	*
PUERTO RICO	1545	49.1	1.7	(45.7, 52.5)	*	2661	58.5	1.3	(55.9, 61.0)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 24.0
 RANGE = 13.7-49.1

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 28.6
 RANGE = 17.3-58.5

* NO PHYSICAL ACTIVITY

April 17, 2001

TABLE 10.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 2000 NO LEISURE TIME PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	29.7	31.6	1.9
ALASKA	23.5	20.0	-3.5
ARIZONA	51.3	34.2	-17.1
ARKANSAS	35.9	28.1	-7.8
CALIFORNIA	25.5	26.5	0.9
COLORADO	21.3	19.8	-1.5
CONNECTICUT	27.2	25.2	-2.0
DELAWARE	35.4	28.0	-7.3
DISTRICT OF CO	38.5	20.8	-17.7
FLORIDA	31.1	28.8	-2.3
GEORGIA	29.6	29.0	-0.7
HAWAII	18.0	23.2	5.3
IDAHO	20.4	19.8	-0.6
ILLINOIS	27.1	30.9	3.8
INDIANA	27.1	25.4	-1.7
IOWA	26.7	27.3	0.6
KANSAS	38.3	30.4	-7.9
KENTUCKY	42.7	41.1	-1.6
LOUISIANA	32.2	36.2	4.0
MAINE	27.7	27.2	-0.5
MARYLAND	20.3	24.2	3.9
MASSACHUSETTS	25.4	24.6	-0.8
MICHIGAN	21.4	22.9	1.5
MINNESOTA	25.5	24.8	-0.6
MISSISSIPPI	33.8	33.3	-0.5
MISSOURI	27.9	28.8	0.9
MONTANA	25.2	23.3	-1.9
NEBRASKA	26.1	29.6	3.5
NEVADA	24.1	25.0	0.9
NEW HAMPSHIRE	24.9	26.7	1.8
NEW JERSEY	32.6	28.6	-4.0
NEW MEXICO	23.0	24.4	1.4
NEW YORK	31.0	29.4	-1.6
NORTH CAROLINA	27.7	30.4	2.7
NORTH DAKOTA	33.1	24.3	-8.8
OHIO	29.8	31.3	1.4
OKLAHOMA	43.0	34.4	-8.5
OREGON	18.9	20.1	1.2
PENNSYLVANIA	32.7	23.0	-9.8
RHODE ISLAND	29.9	27.5	-2.4
SOUTH CAROLINA	33.7	28.1	-5.6
SOUTH DAKOTA	33.3	26.7	-6.6
TENNESSEE	35.8	32.7	-3.1
TEXAS	27.9	28.6	0.7
UTAH	17.1	15.5	-1.6
VERMONT	26.0	23.2	-2.8
VIRGINIA	24.8	25.0	0.2
WASHINGTON	17.6	16.9	-0.7
WEST VIRGINIA	43.7	33.6	-10.1
WISCONSIN	23.4	22.1	-1.3
WYOMING	21.1	22.6	1.5
PUERTO RICO	57.4	54.1	-3.3

* NO PHYSICAL ACTIVITY

April 17, 2001

TABLE 11.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND SUSTAINED PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2241	78.1	1.0	(76.0, 80.1)
ALASKA	2081	73.7	1.4	(70.9, 76.5)
ARIZONA	2678	78.7	1.3	(76.1, 81.3)
ARKANSAS	3005	78.7	0.9	(76.9, 80.4)
CALIFORNIA	3907	76.0	0.8	(74.4, 77.6)
COLORADO	3057	75.3	1.1	(73.1, 77.5)
CONNECTICUT	3913	76.3	0.8	(74.8, 77.9)
DELAWARE	2757	79.1	1.1	(77.0, 81.3)
DISTRICT OF CO	1715	70.8	1.3	(68.2, 73.3)
FLORIDA	5200	77.8	0.7	(76.5, 79.1)
GEORGIA	4108	79.5	0.8	(77.8, 81.1)
HAWAII	6016	71.1	0.8	(69.5, 72.7)
IDAHO	4974	74.8	0.7	(73.4, 76.2)
ILLINOIS	1853	79.4	1.1	(77.3, 81.5)
INDIANA	2921	78.3	0.9	(76.5, 80.0)
IOWA	3610	80.1	0.8	(78.5, 81.6)
KANSAS	4232	82.1	0.7	(80.8, 83.4)
KENTUCKY	6415	85.7	0.6	(84.5, 86.9)
LOUISIANA	5009	85.3	0.6	(84.2, 86.4)
MAINE	4600	79.2	1.0	(77.3, 81.1)
MARYLAND	4589	77.7	0.8	(76.1, 79.2)
MASSACHUSETTS	8144	76.4	0.6	(75.2, 77.5)
MICHIGAN	2604	75.2	1.0	(73.2, 77.1)
MINNESOTA	2850	79.5	0.9	(77.8, 81.1)
MISSISSIPPI	2181	81.4	1.0	(79.5, 83.4)
MISSOURI	4385	79.8	0.9	(78.1, 81.5)
MONTANA	3018	76.1	1.1	(73.9, 78.2)
NEBRASKA	3091	80.4	0.8	(78.8, 82.0)
NEVADA	2101	75.5	1.3	(73.0, 78.0)
NEW HAMPSHIRE	1956	78.2	1.1	(76.0, 80.3)
NEW JERSEY	3776	78.6	0.8	(77.0, 80.1)
NEW MEXICO	3247	76.3	0.9	(74.6, 77.9)
NEW YORK	3357	79.5	0.8	(77.9, 81.0)
NORTH CAROLINA	3016	81.7	0.8	(80.1, 83.3)
NORTH DAKOTA	1916	77.9	1.1	(75.7, 80.0)
OHIO	3250	81.7	1.0	(79.8, 83.6)
OKLAHOMA	3686	81.8	0.7	(80.3, 83.2)
OREGON	3743	73.0	0.8	(71.4, 74.6)
PENNSYLVANIA	3535	77.3	0.8	(75.7, 78.9)
RHODE ISLAND	3541	75.6	0.8	(74.0, 77.3)
SOUTH CAROLINA	3312	79.3	0.8	(77.7, 80.9)
SOUTH DAKOTA	4994	78.6	0.7	(77.3, 79.9)
TENNESSEE	3036	85.5	0.8	(84.0, 87.0)
TEXAS	5016	79.3	0.6	(78.0, 80.6)
UTAH	2890	74.0	1.1	(71.9, 76.1)
VERMONT	3629	75.3	0.8	(73.6, 76.9)
VIRGINIA	1993	77.3	1.2	(75.0, 79.5)
WASHINGTON	3578	73.0	0.8	(71.3, 74.6)
WEST VIRGINIA	2353	81.9	0.9	(80.2, 83.7)
WISCONSIN	2717	76.9	0.9	(75.1, 78.7)
WYOMING	2389	73.5	1.0	(71.5, 75.5)
PUERTO RICO	4206	83.8	0.8	(82.1, 85.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 78.2 RANGE = 70.8-85.7
 * 5+ TIMES PER WEEK, 30+ MINUTES PER SESSION, REGARDLESS OF INTENSITY

TABLE 11.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND SUSTAINED PHYSICAL ACTIVITY* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	836	73.2	1.8	(69.7, 76.7)	*	1405	82.3	1.1	(80.1, 84.5)	
ALASKA	987	73.0	2.1	(69.0, 77.0)	*	1094	74.4	2.0	(70.6, 78.3)	
ARIZONA	1085	77.6	2.0	(73.6, 81.6)	*	1593	79.8	1.8	(76.3, 83.2)	
ARKANSAS	1155	75.8	1.5	(72.9, 78.8)	*	1850	81.2	1.1	(79.1, 83.2)	
CALIFORNIA	1647	75.4	1.2	(73.0, 77.8)	*	2260	76.6	1.1	(74.5, 78.7)	
COLORADO	1294	76.3	1.7	(72.9, 79.6)	*	1763	74.5	1.5	(71.6, 77.3)	
CONNECTICUT	1526	75.0	1.3	(72.5, 77.5)	*	2387	77.5	1.0	(75.7, 79.4)	
DELAWARE	1134	75.3	1.8	(71.8, 78.8)	*	1623	82.6	1.3	(80.2, 85.1)	
DISTRICT OF CO	674	68.4	2.1	(64.3, 72.4)	*	1041	72.8	1.6	(69.6, 75.9)	
FLORIDA	2098	77.7	1.1	(75.6, 79.8)	*	3102	77.9	0.9	(76.2, 79.6)	
GEORGIA	1569	77.1	1.4	(74.4, 79.9)	*	2539	81.6	1.0	(79.7, 83.6)	
HAWAII	2681	67.4	1.3	(64.9, 69.9)	*	3335	74.9	1.0	(73.0, 76.9)	
IDAHO	2117	75.8	1.1	(73.7, 77.9)	*	2857	73.9	1.0	(72.0, 75.7)	
ILLINOIS	785	78.6	1.7	(75.3, 81.8)	*	1068	80.1	1.4	(77.4, 82.9)	
INDIANA	1209	73.3	1.5	(70.4, 76.2)	*	1712	82.8	1.0	(80.8, 84.7)	
IOWA	1424	79.5	1.3	(77.0, 81.9)	*	2186	80.7	1.0	(78.8, 82.6)	
KANSAS	1643	81.9	1.1	(79.8, 84.0)	*	2589	82.4	0.8	(80.8, 84.0)	
KENTUCKY	2553	84.6	1.0	(82.7, 86.6)	*	3862	86.7	0.8	(85.1, 88.2)	
LOUISIANA	1878	83.9	1.0	(82.0, 85.9)	*	3131	86.6	0.7	(85.2, 87.9)	
MAINE	1886	80.8	1.4	(78.1, 83.5)	*	2714	77.7	1.3	(75.0, 80.3)	
MARYLAND	1778	75.8	1.3	(73.3, 78.4)	*	2811	79.4	1.0	(77.5, 81.2)	
MASSACHUSETTS	3225	75.6	0.9	(73.8, 77.4)	*	4919	77.1	0.7	(75.7, 78.5)	
MICHIGAN	1120	74.5	1.5	(71.6, 77.5)	*	1484	75.8	1.3	(73.1, 78.4)	
MINNESOTA	1255	80.0	1.3	(77.5, 82.4)	*	1595	79.0	1.1	(76.8, 81.2)	
MISSISSIPPI	822	79.9	1.6	(76.8, 83.1)	*	1359	82.8	1.1	(80.5, 85.0)	
MISSOURI	1695	80.2	1.4	(77.5, 82.8)	*	2690	79.4	1.1	(77.2, 81.6)	
MONTANA	1290	77.3	1.6	(74.2, 80.4)	*	1728	74.9	1.5	(72.0, 77.8)	
NEBRASKA	1247	81.1	1.2	(78.7, 83.6)	*	1844	79.7	1.1	(77.5, 81.8)	
NEVADA	1046	74.2	1.9	(70.4, 78.0)	*	1055	76.8	1.7	(73.4, 80.2)	
NEW HAMPSHIRE	789	78.9	1.6	(75.7, 82.2)	*	1167	77.4	1.4	(74.7, 80.2)	
NEW JERSEY	1475	77.2	1.3	(74.7, 79.6)	*	2301	79.9	1.0	(77.9, 81.8)	
NEW MEXICO	1468	76.1	1.3	(73.5, 78.6)	*	1779	76.4	1.1	(74.3, 78.6)	
NEW YORK	1338	78.8	1.3	(76.3, 81.2)	*	2019	80.1	1.0	(78.1, 82.1)	
NORTH CAROLINA	1145	80.7	1.3	(78.1, 83.3)	*	1871	82.6	1.0	(80.6, 84.5)	
NORTH DAKOTA	816	79.4	1.7	(76.1, 82.7)	*	1100	76.4	1.4	(73.6, 79.1)	
OHIO	1273	79.3	1.7	(76.0, 82.5)	*	1977	83.9	1.1	(81.7, 86.1)	
OKLAHOMA	1569	80.4	1.2	(78.1, 82.7)	*	2117	83.0	0.9	(81.1, 84.8)	
OREGON	1495	73.4	1.3	(70.9, 75.9)	*	2248	72.7	1.1	(70.6, 74.7)	
PENNSYLVANIA	1423	77.8	1.2	(75.3, 80.2)	*	2112	76.9	1.0	(74.9, 78.9)	
RHODE ISLAND	1373	75.0	1.3	(72.4, 77.7)	*	2168	76.1	1.1	(74.1, 78.2)	
SOUTH CAROLINA	1322	78.9	1.3	(76.3, 81.5)	*	1990	79.7	1.0	(77.7, 81.7)	
SOUTH DAKOTA	1933	78.7	1.0	(76.7, 80.8)	*	3061	78.5	0.9	(76.9, 80.2)	
TENNESSEE	999	85.1	1.4	(82.4, 87.7)	*	2037	85.9	0.9	(84.2, 87.6)	
TEXAS	2012	79.0	1.0	(77.1, 81.0)	*	3004	79.6	0.8	(77.9, 81.2)	
UTAH	1264	74.0	1.6	(70.8, 77.2)	*	1626	74.1	1.4	(71.3, 76.9)	
VERMONT	1516	76.0	1.2	(73.6, 78.4)	*	2113	74.6	1.1	(72.4, 76.7)	
VIRGINIA	854	78.8	1.6	(75.8, 81.9)	*	1139	75.8	1.7	(72.6, 79.1)	
WASHINGTON	1488	72.8	1.3	(70.2, 75.3)	*	2090	73.2	1.1	(71.1, 75.2)	
WEST VIRGINIA	954	79.7	1.5	(76.9, 82.6)	*	1399	83.9	1.1	(81.8, 86.0)	
WISCONSIN	1219	75.5	1.4	(72.7, 78.2)	*	1498	78.3	1.2	(76.0, 80.5)	
WYOMING	1010	73.4	1.5	(70.4, 76.4)	*	1379	73.6	1.3	(71.1, 76.2)	
PUERTO RICO	1545	81.6	1.4	(78.8, 84.4)	*	2661	85.7	1.0	(83.8, 87.6)	

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 77.4	MEDIAN	= 78.8
	RANGE	= 67.4-85.1	RANGE	= 72.7-86.7

* 5+ TIMES PER WEEK, 30+ MINUTES PER SESSION, REGARDLESS OF INTENSITY

TABLE 11.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 2000 NO REGULAR AND SUSTAINED PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	78.6	78.1	-0.5
ALASKA	75.2	73.7	-1.5
ARIZONA	86.5	78.7	-7.8
ARKANSAS	84.6	78.7	-5.9
CALIFORNIA	75.3	76.0	0.7
COLORADO	75.9	75.3	-0.5
CONNECTICUT	79.5	76.3	-3.2
DELAWARE	82.8	79.1	-3.7
DISTRICT OF CO	79.7	70.8	-8.9
FLORIDA	79.5	77.8	-1.7
GEORGIA	82.0	79.5	-2.5
HAWAII	69.7	71.1	1.4
IDAHO	72.9	74.8	2.0
ILLINOIS	78.1	79.4	1.3
INDIANA	80.5	78.3	-2.2
IOWA	80.2	80.1	-0.1
KANSAS	85.7	82.1	-3.6
KENTUCKY	87.0	85.7	-1.3
LOUISIANA	83.9	85.3	1.5
MAINE	75.9	79.2	3.3
MARYLAND	75.0	77.7	2.7
MASSACHUSETTS	78.0	76.4	-1.6
MICHIGAN	76.4	75.2	-1.2
MINNESOTA	78.3	79.5	1.2
MISSISSIPPI	80.8	81.4	0.7
MISSOURI	80.9	79.8	-1.1
MONTANA	77.5	76.1	-1.5
NEBRASKA	80.9	80.4	-0.5
NEVADA	74.4	75.5	1.1
NEW HAMPSHIRE	76.3	78.2	1.8
NEW JERSEY	80.9	78.6	-2.3
NEW MEXICO	74.2	76.3	2.0
NEW YORK	80.3	79.5	-0.8
NORTH CAROLINA	81.5	81.7	0.2
NORTH DAKOTA	81.3	77.9	-3.4
OHIO	81.0	81.7	0.7
OKLAHOMA	84.3	81.8	-2.6
OREGON	73.3	73.0	-0.3
PENNSYLVANIA	80.6	77.3	-3.3
RHODE ISLAND	79.6	75.6	-4.0
SOUTH CAROLINA	82.2	79.3	-2.9
SOUTH DAKOTA	81.8	78.6	-3.2
TENNESSEE	83.5	85.5	2.0
TEXAS	79.2	79.3	0.1
UTAH	72.9	74.0	1.2
VERMONT	74.5	75.3	0.8
VIRGINIA	78.7	77.3	-1.5
WASHINGTON	74.5	73.0	-1.6
WEST VIRGINIA	86.5	81.9	-4.6
WISCONSIN	79.8	76.9	-2.9
WYOMING	71.8	73.5	1.7
PUERTO RICO	84.5	83.8	-0.7

* 5+ TIMES PER WEEK, 30+ MINUTES PER SESSION, REGARDLESS OF INTENSITY

TABLE 12.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 34
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2241	89.5	0.8	(87.8, 91.1)
ALASKA	2081	81.7	1.3	(79.1, 84.2)
ARIZONA	2678	84.6	1.2	(82.3, 86.9)
ARKANSAS	3005	88.4	0.7	(87.0, 89.7)
CALIFORNIA	3907	83.6	0.7	(82.3, 84.9)
COLORADO	3057	83.8	0.9	(82.0, 85.7)
CONNECTICUT	3913	83.0	0.7	(81.7, 84.4)
DELAWARE	2757	87.1	0.9	(85.4, 88.7)
DISTRICT OF CO	1715	84.0	1.0	(82.0, 85.9)
FLORIDA	5200	84.4	0.6	(83.3, 85.5)
GEORGIA	4108	88.0	0.7	(86.7, 89.2)
HAWAII	6016	82.1	0.7	(80.7, 83.4)
IDAHO	4974	85.7	0.6	(84.6, 86.9)
ILLINOIS	1853	87.5	0.8	(85.9, 89.1)
INDIANA	2921	87.2	0.7	(85.8, 88.6)
IOWA	3610	87.7	0.6	(86.5, 88.9)
KANSAS	4232	88.1	0.6	(87.0, 89.2)
KENTUCKY	6415	90.5	0.5	(89.5, 91.4)
LOUISIANA	5009	91.1	0.4	(90.2, 92.0)
MAINE	4600	85.9	0.8	(84.3, 87.5)
MARYLAND	4589	83.2	0.7	(81.7, 84.6)
MASSACHUSETTS	8144	82.8	0.5	(81.8, 83.8)
MICHIGAN	2604	85.1	0.8	(83.6, 86.6)
MINNESOTA	2850	84.4	0.8	(82.9, 85.9)
MISSISSIPPI	2181	90.5	0.7	(89.2, 91.8)
MISSOURI	4385	87.4	0.7	(86.1, 88.8)
MONTANA	3018	81.7	1.0	(79.7, 83.6)
NEBRASKA	3091	86.1	0.7	(84.7, 87.4)
NEVADA	2101	85.6	1.1	(83.5, 87.7)
NEW HAMPSHIRE	1956	83.6	0.9	(81.8, 85.5)
NEW JERSEY	3776	85.6	0.7	(84.4, 86.9)
NEW MEXICO	3247	85.3	0.7	(84.0, 86.6)
NEW YORK	3357	86.0	0.7	(84.7, 87.4)
NORTH CAROLINA	3016	87.6	0.7	(86.1, 89.0)
NORTH DAKOTA	1916	86.6	0.8	(84.9, 88.2)
OHIO	3250	86.9	0.8	(85.3, 88.5)
OKLAHOMA	3686	87.5	0.6	(86.3, 88.6)
OREGON	3743	82.9	0.7	(81.5, 84.2)
PENNSYLVANIA	3535	86.0	0.6	(84.8, 87.3)
RHODE ISLAND	3541	82.4	0.7	(81.0, 83.8)
SOUTH CAROLINA	3312	87.2	0.6	(86.0, 88.4)
SOUTH DAKOTA	4994	87.3	0.6	(86.2, 88.4)
TENNESSEE	3036	89.4	0.6	(88.1, 90.6)
TEXAS	5016	87.4	0.5	(86.4, 88.4)
UTAH	2890	84.8	0.9	(83.0, 86.6)
VERMONT	3629	82.4	0.7	(81.0, 83.8)
VIRGINIA	1993	84.2	0.9	(82.4, 86.0)
WASHINGTON	3578	83.4	0.7	(82.1, 84.8)
WEST VIRGINIA	2353	88.9	0.7	(87.5, 90.2)
WISCONSIN	2717	85.4	0.7	(84.0, 86.8)
WYOMING	2389	86.2	0.8	(84.7, 87.7)
PUERTO RICO	4206	94.7	0.4	(93.9, 95.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 86.0 RANGE = 81.7-94.7

* 3+ TIMES/WEEK 20+ MINUTE/SESSIONS, 50% OF CAPACITY

April 17, 2001

PREVALENCE DIFFERENCES FROM 1998 TO 2000 NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	90.0	89.5	-0.5
ALASKA	82.5	81.7	-0.8
ARIZONA	90.2	84.6	-5.6
ARKANSAS	89.7	88.4	-1.3
CALIFORNIA	83.6	83.6	0.0
COLORADO	85.2	83.8	-1.4
CONNECTICUT	85.9	83.0	-2.8
DELAWARE	87.6	87.1	-0.6
DISTRICT OF CO	83.4	84.0	0.5
FLORIDA	84.4	84.4	0.0
GEORGIA	87.4	88.0	0.6
HAWAII	82.3	82.1	-0.2
IDAHO	84.1	85.7	1.6
ILLINOIS	85.8	87.5	1.7
INDIANA	87.1	87.2	0.1
IOWA	89.4	87.7	-1.7
KANSAS	91.2	88.1	-3.1
KENTUCKY	92.9	90.5	-2.4
LOUISIANA	89.3	91.1	1.8
MAINE	85.7	85.9	0.2
MARYLAND	83.6	83.2	-0.5
MASSACHUSETTS	84.4	82.8	-1.6
MICHIGAN	85.9	85.1	-0.8
MINNESOTA	84.5	84.4	-0.1
MISSISSIPPI	89.6	90.5	0.9
MISSOURI	88.6	87.4	-1.2
MONTANA	86.7	81.7	-5.0
NEBRASKA	89.5	86.1	-3.4
NEVADA	85.7	85.6	-0.1
NEW HAMPSHIRE	87.7	83.6	-4.0
NEW JERSEY	87.6	85.6	-2.0
NEW MEXICO	85.1	85.3	0.2
NEW YORK	85.2	86.0	0.9
NORTH CAROLINA	88.4	87.6	-0.8
NORTH DAKOTA	89.0	86.6	-2.5
OHIO	88.4	86.9	-1.5
OKLAHOMA	88.8	87.5	-1.3
OREGON	84.9	82.9	-2.0
PENNSYLVANIA	86.8	86.0	-0.7
RHODE ISLAND	84.5	82.4	-2.1
SOUTH CAROLINA	88.9	87.2	-1.6
SOUTH DAKOTA	90.3	87.3	-2.9
TENNESSEE	90.4	89.4	-1.0
TEXAS	86.2	87.4	1.3
UTAH	85.0	84.8	-0.2
VERMONT	83.0	82.4	-0.6
VIRGINIA	84.6	84.2	-0.4
WASHINGTON	85.5	83.4	-2.1
WEST VIRGINIA	92.7	88.9	-3.8
WISCONSIN	86.8	85.4	-1.4
WYOMING	84.4	86.2	1.8
PUERTO RICO	91.4	94.7	3.3

* 3+ TIMES/WEEK 20+ MINUTE/SESSIONS, 50% OF CAPACITY

April 17, 2001

TABLE 13.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FRUITS AND VEGETABLES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2241	77.3	1.1	(75.1, 79.4)
ALASKA	2083	76.3	1.4	(73.5, 79.0)
ARIZONA	2679	63.1	1.6	(60.0, 66.2)
ARKANSAS	3005	77.5	0.9	(75.8, 79.2)
CALIFORNIA	3906	73.3	0.8	(71.7, 75.0)
COLORADO	3057	76.7	1.1	(74.6, 78.7)
CONNECTICUT	3914	70.7	0.8	(69.0, 72.3)
DELAWARE	2757	77.5	1.1	(75.4, 79.7)
DISTRICT OF CO	1716	68.1	1.3	(65.6, 70.6)
FLORIDA	5201	76.8	0.7	(75.4, 78.1)
GEORGIA	4111	77.7	0.8	(76.1, 79.3)
HAWAII	6015	77.6	0.7	(76.2, 79.0)
IDAHO	4978	78.9	0.7	(77.5, 80.2)
ILLINOIS	1855	76.7	1.2	(74.4, 78.9)
INDIANA	2922	80.0	0.8	(78.3, 81.6)
IOWA	3608	81.9	0.7	(80.5, 83.3)
KANSAS	4235	76.6	0.7	(75.2, 78.0)
KENTUCKY	6412	77.3	0.7	(76.0, 78.7)
LOUISIANA	5009	84.2	0.6	(83.1, 85.3)
MAINE	4600	75.5	1.0	(73.5, 77.5)
MARYLAND	4592	72.6	0.9	(70.9, 74.2)
MASSACHUSETTS	8147	70.0	0.6	(68.8, 71.2)
MICHIGAN	2605	76.9	0.9	(75.2, 78.7)
MINNESOTA	2853	75.7	0.9	(73.9, 77.4)
MISSISSIPPI	2181	81.4	1.0	(79.4, 83.4)
MISSOURI	4384	79.3	0.9	(77.6, 81.0)
MONTANA	3010	77.2	1.0	(75.2, 79.3)
NEBRASKA	3089	79.4	0.8	(77.8, 81.0)
NEVADA	2102	78.7	1.3	(76.1, 81.3)
NEW HAMPSHIRE	1958	73.8	1.2	(71.6, 76.1)
NEW JERSEY	3779	72.6	0.9	(70.9, 74.3)
NEW MEXICO	3245	79.5	0.8	(78.0, 81.1)
NEW YORK	3358	72.5	0.9	(70.7, 74.3)
NORTH CAROLINA	3011	77.9	0.9	(76.2, 79.5)
NORTH DAKOTA	1917	76.8	1.1	(74.7, 78.9)
OHIO	3255	78.6	1.0	(76.6, 80.6)
OKLAHOMA	3685	81.8	0.7	(80.5, 83.2)
OREGON	3746	73.2	0.8	(71.6, 74.8)
PENNSYLVANIA	3536	76.7	0.8	(75.2, 78.3)
RHODE ISLAND	3543	70.8	0.9	(69.1, 72.6)
SOUTH CAROLINA	3313	75.4	0.9	(73.7, 77.1)
SOUTH DAKOTA	4994	80.1	0.6	(78.9, 81.3)
TENNESSEE	3033	65.9	1.0	(63.9, 67.9)
TEXAS	5018	76.6	0.7	(75.3, 78.0)
UTAH	2891	79.4	1.0	(77.5, 81.4)
VERMONT	3628	71.3	0.8	(69.6, 72.9)
VIRGINIA	1996	74.4	1.2	(72.2, 76.7)
WASHINGTON	3583	75.3	0.8	(73.8, 76.8)
WEST VIRGINIA	2353	78.6	1.0	(76.7, 80.5)
WISCONSIN	2720	78.3	0.9	(76.5, 80.1)
WYOMING	2390	79.6	0.9	(77.8, 81.5)
PUERTO RICO	3897	92.8	0.5	(91.7, 93.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 76.9 RANGE = 63.1-92.8

* NOT EATING 5 SERVINGS A DAY

April 17, 2001

TABLE 13.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FRUITS AND VEGETABLES* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*				*-----FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	837	77.4	1.9	(73.7, 81.0)	1404	77.2	1.2	(74.8, 79.7)
ALASKA	988	78.1	2.1	(74.1, 82.1)	1095	74.3	1.9	(70.5, 78.1)
ARIZONA	1086	65.2	2.4	(60.5, 69.9)	1593	61.1	2.2	(56.8, 65.4)
ARKANSAS	1155	79.7	1.4	(77.0, 82.4)	1850	75.5	1.1	(73.4, 77.7)
CALIFORNIA	1646	80.1	1.2	(77.9, 82.4)	2260	66.7	1.2	(64.4, 69.0)
COLORADO	1294	80.3	1.5	(77.4, 83.3)	1763	73.1	1.5	(70.2, 76.0)
CONNECTICUT	1526	74.1	1.3	(71.6, 76.6)	2388	67.5	1.1	(65.4, 69.6)
DELAWARE	1133	81.4	1.7	(78.1, 84.6)	1624	74.0	1.4	(71.2, 76.8)
DISTRICT OF CO	674	71.7	2.0	(67.8, 75.5)	1042	65.1	1.7	(61.8, 68.4)
FLORIDA	2098	80.9	1.0	(78.9, 82.8)	3103	73.0	0.9	(71.2, 74.8)
GEORGIA	1571	80.3	1.2	(77.9, 82.7)	2540	75.3	1.1	(73.1, 77.4)
HAWAII	2680	81.1	1.0	(79.2, 83.1)	3335	74.0	1.0	(72.0, 75.9)
IDAHO	2119	84.0	0.9	(82.2, 85.8)	2859	73.9	1.0	(72.0, 75.8)
ILLINOIS	786	79.5	1.7	(76.1, 82.8)	1069	74.0	1.6	(70.9, 77.0)
INDIANA	1209	83.6	1.2	(81.3, 86.0)	1713	76.6	1.1	(74.3, 78.8)
IOWA	1422	86.8	1.1	(84.7, 88.8)	2186	77.5	1.0	(75.5, 79.4)
KANSAS	1645	82.3	1.0	(80.4, 84.3)	2590	71.2	1.0	(69.2, 73.2)
KENTUCKY	2552	81.0	1.0	(79.0, 82.9)	3860	74.1	1.0	(72.2, 75.9)
LOUISIANA	1879	85.9	0.9	(84.1, 87.6)	3130	82.7	0.7	(81.2, 84.1)
MAINE	1886	82.7	1.3	(80.1, 85.3)	2714	68.8	1.5	(65.9, 71.7)
MARYLAND	1779	78.1	1.3	(75.5, 80.6)	2813	67.5	1.1	(65.4, 69.7)
MASSACHUSETTS	3226	76.1	0.9	(74.3, 77.9)	4921	64.6	0.8	(62.9, 66.2)
MICHIGAN	1122	82.4	1.3	(79.9, 84.8)	1483	72.0	1.3	(69.4, 74.5)
MINNESOTA	1257	80.6	1.2	(78.2, 83.0)	1596	71.0	1.3	(68.5, 73.6)
MISSISSIPPI	822	83.2	1.5	(80.3, 86.0)	1359	79.9	1.5	(77.0, 82.7)
MISSOURI	1695	83.2	1.3	(80.7, 85.8)	2689	75.8	1.1	(73.5, 78.0)
MONTANA	1284	82.1	1.4	(79.4, 84.8)	1726	72.6	1.5	(69.6, 75.5)
NEBRASKA	1244	83.8	1.2	(81.5, 86.1)	1845	75.3	1.2	(73.1, 77.6)
NEVADA	1046	81.7	1.8	(78.1, 85.3)	1056	75.7	1.9	(71.9, 79.5)
NEW HAMPSHIRE	791	79.1	1.7	(75.8, 82.5)	1167	68.9	1.5	(65.9, 71.9)
NEW JERSEY	1475	76.0	1.3	(73.4, 78.6)	2304	69.4	1.1	(67.2, 71.6)
NEW MEXICO	1467	83.8	1.1	(81.8, 85.9)	1778	75.5	1.2	(73.2, 77.8)
NEW YORK	1336	78.3	1.3	(75.7, 80.9)	2022	67.3	1.2	(64.9, 69.7)
NORTH CAROLINA	1142	81.1	1.3	(78.6, 83.7)	1869	74.9	1.1	(72.6, 77.1)
NORTH DAKOTA	815	82.3	1.5	(79.4, 85.2)	1102	71.5	1.5	(68.6, 74.4)
OHIO	1275	82.3	1.6	(79.2, 85.3)	1980	75.3	1.3	(72.7, 77.9)
OKLAHOMA	1569	85.3	1.0	(83.4, 87.1)	2116	78.7	1.0	(76.7, 80.7)
OREGON	1496	78.0	1.2	(75.6, 80.5)	2250	68.6	1.1	(66.4, 70.8)
PENNSYLVANIA	1423	82.5	1.1	(80.4, 84.7)	2113	71.5	1.1	(69.3, 73.7)
RHODE ISLAND	1375	77.2	1.3	(74.7, 79.8)	2168	65.1	1.2	(62.7, 67.5)
SOUTH CAROLINA	1322	78.0	1.4	(75.4, 80.7)	1991	73.0	1.1	(70.8, 75.2)
SOUTH DAKOTA	1932	84.5	0.9	(82.7, 86.2)	3062	76.0	0.9	(74.3, 77.7)
TENNESSEE	999	68.6	1.7	(65.3, 71.8)	2034	63.5	1.2	(61.2, 65.8)
TEXAS	2014	81.4	1.0	(79.5, 83.3)	3004	72.1	0.9	(70.3, 74.0)
UTAH	1264	84.0	1.4	(81.3, 86.7)	1627	75.0	1.4	(72.3, 77.8)
VERMONT	1517	78.1	1.2	(75.8, 80.4)	2111	64.9	1.2	(62.6, 67.1)
VIRGINIA	855	79.5	1.6	(76.4, 82.6)	1141	69.7	1.7	(66.4, 73.0)
WASHINGTON	1490	80.5	1.1	(78.3, 82.6)	2093	70.4	1.1	(68.3, 72.5)
WEST VIRGINIA	954	83.4	1.4	(80.7, 86.1)	1399	74.2	1.3	(71.6, 76.8)
WISCONSIN	1222	84.4	1.2	(82.0, 86.8)	1498	72.6	1.3	(70.1, 75.2)
WYOMING	1010	84.2	1.3	(81.6, 86.8)	1380	75.2	1.3	(72.6, 77.7)
PUERTO RICO	1404	93.4	0.8	(91.9, 94.9)	2493	92.2	0.7	(90.8, 93.6)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 81.1
 RANGE = 65.2-93.4

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 73.1
 RANGE = 61.1-92.2

* NOT EATING 5 SERVINGS A DAY

April 17, 2001

TABLE 13.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 2000 FRUITS AND VEGETABLES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	76.1	77.3	1.2
ALASKA	76.9	76.3	-0.6
ARIZONA	90.9	63.1	-27.8
ARKANSAS	72.1	77.5	5.4
CALIFORNIA	72.5	73.3	0.8
COLORADO	74.0	76.7	2.7
CONNECTICUT	72.0	70.7	-1.4
DELAWARE	73.3	77.5	4.2
DISTRICT OF CO	82.3	68.1	-14.2
FLORIDA	75.1	76.8	1.7
GEORGIA	79.3	77.7	-1.6
HAWAII	72.5	77.6	5.1
IDAHO	76.2	78.9	2.7
ILLINOIS	77.3	76.7	-0.6
INDIANA	76.5	80.0	3.4
IOWA	81.4	81.9	0.5
KANSAS	76.5	76.6	0.1
KENTUCKY	84.4	77.3	-7.0
LOUISIANA	82.7	84.2	1.5
MAINE	73.6	75.5	1.9
MARYLAND	69.9	72.6	2.6
MASSACHUSETTS	69.0	70.0	1.1
MICHIGAN	72.6	76.9	4.4
MINNESOTA	68.1	75.7	7.6
MISSISSIPPI	84.4	81.4	-2.9
MISSOURI	80.0	79.3	-0.7
MONTANA	76.2	77.2	1.0
NEBRASKA	82.3	79.4	-2.9
NEVADA	77.9	78.7	0.8
NEW HAMPSHIRE	72.1	73.8	1.7
NEW JERSEY	73.9	72.6	-1.3
NEW MEXICO	79.3	79.5	0.2
NEW YORK	74.5	72.5	-2.0
NORTH CAROLINA	78.6	77.9	-0.7
NORTH DAKOTA	77.2	76.8	-0.4
OHIO	84.0	78.6	-5.4
OKLAHOMA	79.7	81.8	2.1
OREGON	75.1	73.2	-1.9
PENNSYLVANIA	75.2	76.7	1.6
RHODE ISLAND	75.4	70.8	-4.6
SOUTH CAROLINA	78.2	75.4	-2.8
SOUTH DAKOTA	80.0	80.1	0.1
TENNESSEE	70.3	65.9	-4.4
TEXAS	77.5	76.6	-0.9
UTAH	73.8	79.4	5.7
VERMONT	70.8	71.3	0.5
VIRGINIA	73.1	74.4	1.4
WASHINGTON	74.2	75.3	1.1
WEST VIRGINIA	81.4	78.6	-2.8
WISCONSIN	74.3	78.3	4.0
WYOMING	78.8	79.6	0.9
PUERTO RICO	91.9	92.8	0.9

* NOT EATING 5 SERVINGS A DAY

April 17, 2001

TABLE 14.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NOT EATING FEWER CALORIES AND/OR LESS FAT*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED AND BMI LT 25.0

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1092	53.8	1.7	(50.4, 57.3)
ALASKA	953	50.1	2.3	(45.6, 54.6)
ARIZONA	982	47.5	2.8	(42.0, 53.0)
ARKANSAS	1356	55.1	1.6	(52.0, 58.2)
CALIFORNIA	1780	45.5	1.4	(42.8, 48.2)
COLORADO	1204	49.2	2.2	(44.9, 53.5)
CONNECTICUT	1653	50.5	1.4	(47.8, 53.2)
DELAWARE	1218	52.7	1.9	(49.0, 56.5)
DISTRICT OF CO	749	50.0	2.1	(45.9, 54.0)
FLORIDA	2114	55.3	1.3	(52.8, 57.7)
GEORGIA	1840	51.8	1.5	(48.9, 54.7)
HAWAII	2284	50.6	1.4	(47.7, 53.4)
IDAHO	2203	50.7	1.3	(48.2, 53.2)
ILLINOIS	834	54.3	2.0	(50.4, 58.2)
INDIANA	1382	53.9	1.5	(50.9, 56.9)
IOWA	1662	54.7	1.4	(51.9, 57.5)
KANSAS	1819	53.1	1.3	(50.4, 55.7)
KENTUCKY	2448	59.5	1.4	(56.6, 62.3)
LOUISIANA	2088	55.8	1.3	(53.3, 58.2)
MAINE	2054	53.0	1.9	(49.3, 56.7)
MARYLAND	2011	49.3	1.4	(46.5, 52.0)
MASSACHUSETTS	3402	50.6	1.1	(48.5, 52.6)
MICHIGAN	1282	48.5	1.6	(45.4, 51.7)
MINNESOTA	1070	51.4	1.7	(48.0, 54.8)
MISSISSIPPI	1016	52.8	1.8	(49.3, 56.3)
MISSOURI	1871	52.3	1.6	(49.1, 55.5)
MONTANA	1208	58.0	2.0	(54.2, 61.9)
NEBRASKA	1158	51.3	1.7	(48.1, 54.6)
NEVADA	856	52.9	2.6	(47.7, 58.1)
NEW HAMPSHIRE	818	51.2	2.0	(47.3, 55.1)
NEW JERSEY	1688	54.5	1.4	(51.7, 57.3)
NEW MEXICO	1362	51.7	1.5	(48.7, 54.7)
NEW YORK	1423	53.3	1.6	(50.3, 56.4)
NORTH CAROLINA	1363	59.1	1.6	(56.0, 62.1)
NORTH DAKOTA	847	54.5	1.9	(50.7, 58.2)
OHIO	1415	52.3	1.9	(48.6, 56.0)
OKLAHOMA	1497	57.2	1.4	(54.4, 60.0)
OREGON	1655	49.8	1.4	(47.1, 52.5)
PENNSYLVANIA	1612	53.4	1.4	(50.6, 56.2)
RHODE ISLAND	1503	49.9	1.5	(47.0, 52.9)
SOUTH CAROLINA	1507	53.5	1.5	(50.5, 56.4)
SOUTH DAKOTA	2173	52.3	1.2	(49.9, 54.8)
TENNESSEE	1188	59.6	1.7	(56.3, 62.9)
TEXAS	2296	53.8	1.2	(51.5, 56.2)
UTAH	1203	43.3	1.9	(39.7, 46.9)
VERMONT	1533	50.6	1.4	(47.8, 53.3)
VIRGINIA	892	49.0	1.9	(45.3, 52.7)
WASHINGTON	1542	47.5	1.4	(44.7, 50.2)
WEST VIRGINIA	1008	58.6	1.7	(55.2, 62.0)
WISCONSIN	1288	45.2	1.6	(42.1, 48.3)
WYOMING	1007	51.2	1.7	(47.8, 54.6)
PUERTO RICO	1982	65.1	1.5	(62.2, 67.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 52.3 RANGE = 43.3-65.1
 * AND NOT EXERCISING TO LOSE WEIGHT

April 30, 2001

TABLE 14.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NOT EATING FEWER CALORIES AND/OR LESS FAT* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED AND BMI LT 25.0

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	463	51.9	2.6	(46.7, 57.1)	*	629	55.9	2.3	(51.5, 60.3)	*
ALASKA	501	49.9	3.1	(43.9, 55.9)	*	452	50.5	3.5	(43.7, 57.2)	*
ARIZONA	468	45.6	4.0	(37.7, 53.4)	*	514	50.2	3.8	(42.9, 57.6)	*
ARKANSAS	586	56.6	2.3	(52.1, 61.2)	*	770	53.4	2.0	(49.4, 57.4)	*
CALIFORNIA	895	45.6	2.0	(41.8, 49.4)	*	885	45.4	2.0	(41.5, 49.2)	*
COLORADO	593	47.5	3.1	(41.3, 53.6)	*	611	51.5	2.9	(45.7, 57.2)	*
CONNECTICUT	766	53.1	2.0	(49.2, 57.0)	*	887	47.1	1.9	(43.4, 50.8)	*
DELAWARE	578	52.2	2.8	(46.8, 57.6)	*	640	53.4	2.6	(48.3, 58.4)	*
DISTRICT OF CO	294	52.1	3.3	(45.6, 58.5)	*	455	48.1	2.6	(43.0, 53.2)	*
FLORIDA	980	55.2	1.9	(51.5, 58.8)	*	1134	55.4	1.7	(52.0, 58.7)	*
GEORGIA	773	50.1	2.3	(45.7, 54.6)	*	1067	53.6	1.9	(49.9, 57.4)	*
HAWAII	1234	51.0	1.9	(47.2, 54.8)	*	1050	49.9	2.2	(45.6, 54.2)	*
IDAHO	1044	52.2	1.8	(48.6, 55.8)	*	1159	49.0	1.7	(45.6, 52.4)	*
ILLINOIS	399	56.5	2.8	(51.0, 62.0)	*	435	51.9	2.7	(46.5, 57.2)	*
INDIANA	663	55.2	2.2	(51.0, 59.4)	*	719	52.4	2.1	(48.2, 56.5)	*
IOWA	751	57.8	2.1	(53.8, 61.8)	*	911	51.2	1.9	(47.4, 55.0)	*
KANSAS	818	55.6	2.0	(51.6, 59.5)	*	1001	50.0	1.7	(46.6, 53.4)	*
KENTUCKY	1071	60.8	2.1	(56.6, 64.9)	*	1377	58.1	1.9	(54.3, 61.8)	*
LOUISIANA	835	54.1	2.0	(50.2, 57.9)	*	1253	57.5	1.6	(54.4, 60.5)	*
MAINE	942	52.9	2.7	(47.6, 58.2)	*	1112	53.1	2.5	(48.1, 58.0)	*
MARYLAND	919	50.5	2.1	(46.4, 54.5)	*	1092	47.9	1.9	(44.2, 51.5)	*
MASSACHUSETTS	1631	51.4	1.5	(48.5, 54.3)	*	1771	49.5	1.4	(46.7, 52.3)	*
MICHIGAN	622	53.4	2.3	(48.9, 57.9)	*	660	42.7	2.1	(38.5, 46.9)	*
MINNESOTA	532	54.0	2.5	(49.1, 58.8)	*	538	48.4	2.4	(43.6, 53.1)	*
MISSISSIPPI	429	54.1	2.7	(48.9, 59.4)	*	587	51.3	2.3	(46.9, 55.8)	*
MISSOURI	797	55.5	2.5	(50.7, 60.4)	*	1074	48.6	2.1	(44.5, 52.7)	*
MONTANA	575	61.3	2.7	(55.9, 66.6)	*	633	53.9	2.8	(48.5, 59.3)	*
NEBRASKA	528	54.0	2.4	(49.3, 58.7)	*	630	48.2	2.2	(43.8, 52.6)	*
NEVADA	506	54.0	3.3	(47.6, 60.3)	*	350	51.4	4.2	(43.2, 59.7)	*
NEW HAMPSHIRE	405	51.4	2.8	(45.8, 56.9)	*	413	51.0	2.7	(45.6, 56.3)	*
NEW JERSEY	809	55.8	2.0	(51.9, 59.8)	*	879	52.8	1.9	(49.0, 56.6)	*
NEW MEXICO	687	56.0	2.2	(51.8, 60.2)	*	675	46.8	2.1	(42.6, 51.0)	*
NEW YORK	668	52.8	2.3	(48.4, 57.3)	*	755	53.9	2.1	(49.8, 58.0)	*
NORTH CAROLINA	611	62.0	2.3	(57.5, 66.5)	*	752	55.6	2.1	(51.6, 59.6)	*
NORTH DAKOTA	425	59.1	2.6	(53.9, 64.2)	*	422	48.6	2.7	(43.3, 53.9)	*
OHIO	645	53.8	2.8	(48.3, 59.3)	*	770	50.5	2.6	(45.4, 55.6)	*
OKLAHOMA	723	59.7	2.0	(55.7, 63.7)	*	774	54.4	2.0	(50.4, 58.3)	*
OREGON	733	51.7	2.1	(47.6, 55.7)	*	922	47.7	1.9	(44.0, 51.3)	*
PENNSYLVANIA	776	53.6	2.1	(49.5, 57.7)	*	836	53.2	1.9	(49.4, 57.0)	*
RHODE ISLAND	706	48.9	2.1	(44.7, 53.1)	*	797	51.1	2.1	(46.9, 55.2)	*
SOUTH CAROLINA	681	57.2	2.2	(53.0, 61.4)	*	826	49.3	2.0	(45.4, 53.1)	*
SOUTH DAKOTA	955	53.9	1.8	(50.4, 57.4)	*	1218	50.5	1.7	(47.3, 53.8)	*
TENNESSEE	440	61.6	2.6	(56.5, 66.7)	*	748	57.3	2.1	(53.3, 61.3)	*
TEXAS	1077	52.7	1.7	(49.3, 56.0)	*	1219	55.3	1.6	(52.1, 58.5)	*
UTAH	596	45.5	2.6	(40.4, 50.6)	*	607	40.6	2.6	(35.5, 45.6)	*
VERMONT	738	54.6	2.0	(50.7, 58.5)	*	795	45.7	1.9	(41.9, 49.5)	*
VIRGINIA	457	50.1	2.6	(44.9, 55.2)	*	435	47.6	2.7	(42.4, 52.8)	*
WASHINGTON	756	51.4	2.0	(47.5, 55.4)	*	786	42.2	1.9	(38.5, 46.0)	*
WEST VIRGINIA	474	60.3	2.5	(55.5, 65.2)	*	534	56.6	2.3	(52.0, 61.2)	*
WISCONSIN	660	45.1	2.2	(40.7, 49.5)	*	628	45.4	2.2	(41.0, 49.8)	*
WYOMING	479	56.4	2.5	(51.5, 61.3)	*	528	44.9	2.4	(40.2, 49.6)	*
PUERTO RICO	755	64.5	2.3	(60.0, 68.9)	*	1227	65.7	1.9	(62.0, 69.3)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 53.9
 RANGE = 45.1-64.5

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 50.7
 RANGE = 40.6-65.7

* NOT EXERCISING TO LOSE WEIGHT

April 30, 2001

TABLE 14.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1998 TO 2000 NOT EATING FEWER CALORIES AND/OR LESS FAT*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, REFUSED AND BMI LT 25.0

PARTICIPANT	1998 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	52.1	53.8	1.8
ALASKA	43.4	50.1	6.7
ARIZONA	58.7	47.5	-11.2
ARKANSAS	61.5	55.1	-6.4
CALIFORNIA	48.1	45.5	-2.6
COLORADO	50.3	49.2	-1.1
CONNECTICUT	46.1	50.5	4.4
DELAWARE	55.3	52.7	-2.5
DISTRICT OF CO	62.9	50.0	-12.9
FLORIDA	59.6	55.3	-4.3
GEORGIA	51.8	51.8	0.0
HAWAII	44.6	50.6	6.0
IDAHO	46.2	50.7	4.5
ILLINOIS	54.0	54.3	0.3
INDIANA	48.9	53.9	5.0
IOWA	51.8	54.7	2.8
KANSAS	53.1	53.1	-0.1
KENTUCKY	61.0	59.5	-1.6
LOUISIANA	55.5	55.8	0.3
MAINE	53.9	53.0	-0.9
MARYLAND	48.0	49.3	1.2
MASSACHUSETTS	46.3	50.6	4.3
MICHIGAN	52.1	48.5	-3.5
MINNESOTA	48.8	51.4	2.6
MISSISSIPPI	55.3	52.8	-2.6
MISSOURI	52.5	52.3	-0.2
MONTANA	52.6	58.0	5.4
NEBRASKA	49.6	51.3	1.7
NEVADA	52.3	52.9	0.6
NEW HAMPSHIRE	54.9	51.2	-3.7
NEW JERSEY	53.4	54.5	1.1
NEW MEXICO	47.4	51.7	4.4
NEW YORK	51.6	53.3	1.8
NORTH CAROLINA	60.4	59.1	-1.4
NORTH DAKOTA	55.3	54.5	-0.9
OHIO	52.5	52.3	-0.2
OKLAHOMA	63.1	57.2	-5.9
OREGON	45.1	49.8	4.7
PENNSYLVANIA	54.6	53.4	-1.2
RHODE ISLAND	53.4	49.9	-3.4
SOUTH CAROLINA	56.4	53.5	-2.9
SOUTH DAKOTA	50.6	52.3	1.8
TENNESSEE	51.0	59.6	8.6
TEXAS	53.3	53.8	0.6
UTAH	47.8	43.3	-4.5
VERMONT	52.6	50.6	-2.0
VIRGINIA	47.2	49.0	1.9
WASHINGTON	47.2	47.5	0.3
WEST VIRGINIA	60.2	58.6	-1.6
WISCONSIN	45.5	45.2	-0.2
WYOMING	47.3	51.2	3.9
PUERTO RICO	70.0	65.1	-4.9

* AND NOT EXERCISING TO LOSE WEIGHT

April 30, 2001

TABLE 15.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2156	60.8	1.2	(58.3, 63.2)
ALASKA	2026	59.2	1.6	(56.1, 62.4)
ARIZONA	2528	55.9	1.8	(52.4, 59.5)
ARKANSAS	2891	59.9	1.0	(57.8, 61.9)
CALIFORNIA	3791	57.3	1.0	(55.4, 59.2)
COLORADO	2939	48.0	1.3	(45.4, 50.6)
CONNECTICUT	3710	53.7	0.9	(51.9, 55.6)
DELAWARE	2607	55.8	1.3	(53.2, 58.4)
DISTRICT OF CO	1627	53.3	1.4	(50.4, 56.1)
FLORIDA	4998	53.9	0.8	(52.3, 55.5)
GEORGIA	3919	59.4	1.0	(57.4, 61.4)
HAWAII	5880	50.2	0.9	(48.5, 51.9)
IDAHO	4781	55.7	0.9	(54.0, 57.3)
ILLINOIS	3633	58.9	1.0	(57.1, 60.8)
INDIANA	2829	58.4	1.1	(56.3, 60.4)
IOWA	3425	60.0	1.0	(58.1, 61.9)
KANSAS	4022	58.7	0.9	(57.0, 60.4)
KENTUCKY	6098	61.0	0.9	(59.3, 62.8)
LOUISIANA	4787	60.0	0.8	(58.4, 61.6)
MAINE	4405	56.3	1.3	(53.8, 58.7)
MARYLAND	4367	56.7	1.0	(54.8, 58.6)
MASSACHUSETTS	7749	52.8	0.7	(51.4, 54.2)
MICHIGAN	2508	61.0	1.1	(58.8, 63.2)
MINNESOTA	2744	55.0	1.1	(52.9, 57.1)
MISSISSIPPI	2067	61.7	1.3	(59.2, 64.2)
MISSOURI	4225	56.4	1.1	(54.4, 58.5)
MONTANA	2884	53.1	1.3	(50.6, 55.5)
NEBRASKA	2883	58.5	1.1	(56.4, 60.6)
NEVADA	2025	53.2	1.6	(50.1, 56.4)
NEW HAMPSHIRE	1863	54.6	1.4	(51.9, 57.3)
NEW JERSEY	3595	56.7	1.0	(54.8, 58.6)
NEW MEXICO	3103	55.5	1.0	(53.5, 57.5)
NEW YORK	3186	56.9	1.0	(54.9, 58.9)
NORTH CAROLINA	2875	59.2	1.1	(57.1, 61.3)
NORTH DAKOTA	1795	60.4	1.3	(57.9, 62.9)
OHIO	3082	57.3	1.3	(54.8, 59.8)
OKLAHOMA	3506	56.3	1.0	(54.4, 58.1)
OREGON	3568	57.6	0.9	(55.8, 59.4)
PENNSYLVANIA	3389	57.7	1.0	(55.8, 59.6)
RHODE ISLAND	3371	53.8	1.0	(51.8, 55.7)
SOUTH CAROLINA	3177	59.0	1.0	(57.0, 61.0)
SOUTH DAKOTA	4779	58.8	0.8	(57.2, 60.4)
TENNESSEE	2872	59.4	1.1	(57.3, 61.5)
TEXAS	4758	59.8	0.8	(58.2, 61.4)
UTAH	2795	54.1	1.3	(51.7, 56.6)
VERMONT	3477	52.8	1.0	(50.9, 54.7)
VIRGINIA	1906	56.1	1.3	(53.5, 58.7)
WASHINGTON	3410	55.1	1.0	(53.3, 57.0)
WEST VIRGINIA	2242	59.8	1.2	(57.5, 62.0)
WISCONSIN	2615	57.8	1.1	(55.6, 59.9)
WYOMING	2311	55.1	1.2	(52.9, 57.4)
PUERTO RICO	4003	61.0	1.1	(58.9, 63.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 57.1 RANGE = 48.0-61.7 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 25.0 AND LE 99.8 FOR MALES AND FEMALES April 30, 2001

TABLE 15.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY NEW BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*				-----FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	824	68.2	1.8	(64.7, 71.8)	1332	53.9	1.6	(50.7, 57.0)
ALASKA	981	67.6	2.2	(63.2, 72.0)	1045	49.7	2.3	(45.2, 54.2)
ARIZONA	1056	66.6	2.4	(61.8, 71.4)	1472	45.4	2.4	(40.6, 50.1)
ARKANSAS	1144	66.8	1.6	(63.7, 69.9)	1747	53.3	1.3	(50.7, 55.9)
CALIFORNIA	1624	66.6	1.4	(64.0, 69.3)	* 2167	47.8	1.3	(45.3, 50.3)
COLORADO	1275	56.6	2.0	(52.7, 60.5)	* 1664	39.4	1.7	(36.0, 42.7)
CONNECTICUT	1503	62.6	1.4	(59.8, 65.4)	* 2207	45.1	1.2	(42.7, 47.4)
DELAWARE	1119	62.8	2.0	(58.9, 66.7)	* 1488	49.0	1.7	(45.6, 52.4)
DISTRICT OF CO	651	54.6	2.3	(50.2, 59.0)	* 976	52.1	1.8	(48.5, 55.7)
FLORIDA	2064	63.0	1.2	(60.6, 65.4)	* 2934	45.2	1.1	(43.1, 47.2)
GEORGIA	1545	67.0	1.5	(64.1, 69.9)	* 2374	52.1	1.3	(49.5, 54.6)
HAWAII	2658	60.5	1.3	(58.0, 63.0)	* 3222	39.5	1.2	(37.2, 41.8)
IDAHO	2102	63.1	1.3	(60.7, 65.6)	* 2679	48.1	1.1	(45.9, 50.3)
ILLINOIS	1544	65.6	1.4	(62.8, 68.4)	* 2089	52.4	1.3	(50.0, 54.9)
INDIANA	1203	67.2	1.5	(64.2, 70.2)	* 1626	49.9	1.4	(47.1, 52.7)
IOWA	1403	68.2	1.5	(65.4, 71.1)	* 2022	52.0	1.3	(49.5, 54.4)
KANSAS	1625	68.0	1.3	(65.5, 70.6)	* 2397	49.3	1.2	(47.1, 51.6)
KENTUCKY	2522	68.9	1.3	(66.5, 71.4)	* 3576	53.4	1.2	(51.0, 55.7)
LOUISIANA	1857	66.3	1.3	(63.8, 68.7)	* 2930	54.1	1.0	(52.1, 56.1)
MAINE	1872	64.1	1.9	(60.4, 67.8)	* 2533	48.6	1.7	(45.3, 51.9)
MARYLAND	1754	64.0	1.5	(61.1, 66.9)	* 2613	49.6	1.2	(47.2, 52.0)
MASSACHUSETTS	3168	64.3	1.0	(62.3, 66.3)	* 4581	41.9	0.9	(40.1, 43.6)
MICHIGAN	1110	71.3	1.5	(68.4, 74.2)	* 1398	51.2	1.5	(48.2, 54.2)
MINNESOTA	1243	64.8	1.5	(61.8, 67.7)	* 1501	45.3	1.5	(42.4, 48.3)
MISSISSIPPI	811	70.3	1.8	(66.7, 73.9)	* 1256	53.6	1.6	(50.4, 56.8)
MISSOURI	1676	66.2	1.6	(63.1, 69.4)	* 2549	47.2	1.3	(44.6, 49.8)
MONTANA	1277	62.6	1.8	(59.0, 66.2)	* 1607	43.4	1.7	(40.1, 46.7)
NEBRASKA	1201	69.8	1.5	(66.8, 72.8)	* 1682	47.3	1.4	(44.6, 50.0)
NEVADA	1039	62.9	2.1	(58.8, 67.0)	* 986	42.8	2.4	(38.1, 47.5)
NEW HAMPSHIRE	782	66.4	2.0	(62.5, 70.3)	* 1081	42.8	1.7	(39.4, 46.2)
NEW JERSEY	1459	66.8	1.4	(64.0, 69.5)	* 2136	47.0	1.2	(44.5, 49.4)
NEW MEXICO	1440	63.4	1.5	(60.5, 66.3)	* 1663	47.7	1.4	(45.1, 50.4)
NEW YORK	1318	65.0	1.5	(62.0, 68.0)	* 1868	49.2	1.4	(46.5, 51.8)
NORTH CAROLINA	1136	67.2	1.7	(64.0, 70.5)	* 1739	51.3	1.4	(48.6, 54.0)
NORTH DAKOTA	806	70.5	1.8	(67.0, 74.0)	* 989	49.6	1.7	(46.2, 53.0)
OHIO	1257	65.1	1.9	(61.3, 68.9)	* 1825	49.7	1.7	(46.4, 52.9)
OKLAHOMA	1543	66.4	1.4	(63.7, 69.1)	* 1963	46.4	1.3	(43.8, 48.9)
OREGON	1475	65.4	1.4	(62.7, 68.1)	* 2093	49.7	1.2	(47.3, 52.1)
PENNSYLVANIA	1413	67.5	1.4	(64.7, 70.3)	* 1976	48.4	1.3	(45.9, 50.9)
RHODE ISLAND	1360	63.4	1.5	(60.4, 66.4)	* 2011	44.6	1.3	(42.1, 47.1)
SOUTH CAROLINA	1309	66.1	1.6	(63.0, 69.1)	* 1868	52.2	1.3	(49.6, 54.7)
SOUTH DAKOTA	1913	66.6	1.2	(64.2, 69.0)	* 2866	51.0	1.1	(48.9, 53.1)
TENNESSEE	977	68.0	1.7	(64.8, 71.3)	* 1895	51.2	1.3	(48.7, 53.7)
TEXAS	1970	69.4	1.2	(67.1, 71.7)	* 2788	50.2	1.1	(48.1, 52.4)
UTAH	1260	62.2	1.8	(58.7, 65.7)	* 1535	46.0	1.7	(42.7, 49.2)
VERMONT	1508	61.4	1.5	(58.4, 64.3)	* 1969	44.3	1.3	(41.8, 46.7)
VIRGINIA	849	65.3	1.9	(61.5, 69.0)	* 1057	46.9	1.8	(43.4, 50.3)
WASHINGTON	1473	64.2	1.4	(61.5, 67.0)	* 1937	45.8	1.2	(43.3, 48.2)
WEST VIRGINIA	942	69.1	1.7	(65.7, 72.4)	* 1300	50.9	1.5	(47.9, 53.9)
WISCONSIN	1213	67.9	1.6	(64.8, 71.0)	* 1402	47.7	1.5	(44.8, 50.7)
WYOMING	1000	64.8	1.7	(61.5, 68.0)	* 1311	45.3	1.5	(42.3, 48.2)
PUERTO RICO	1504	66.4	1.7	(62.9, 69.8)	* 2499	56.2	1.4	(53.5, 58.9)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 66.2
 RANGE = 54.6-71.3

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 48.8
 RANGE = 39.4-56.2

* BODY MASS INDEX GE 25.0 AND LE 99.8 FOR MALES AND FEMALES

April 30, 2001

TABLE 15.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 OVERWEIGHT: BY NEW BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	60.9	60.8	-0.1
ALASKA	61.1	59.2	-1.9
ARIZONA	51.6	55.9	4.3
ARKANSAS	58.8	59.9	1.0
CALIFORNIA	54.8	57.3	2.5
COLORADO	47.9	48.0	0.1
CONNECTICUT	52.6	53.7	1.2
DELAWARE	55.5	55.8	0.3
DISTRICT OF CO	50.9	53.3	2.4
FLORIDA	56.8	53.9	-3.0
GEORGIA	57.6	59.4	1.8
HAWAII	50.2	50.2	0.1
IDAHO	55.2	55.7	0.4
ILLINOIS	56.9	58.9	2.0
INDIANA	56.7	58.4	1.6
IOWA	58.4	60.0	1.6
KANSAS	56.1	58.7	2.6
KENTUCKY	58.5	61.0	2.5
LOUISIANA	58.3	60.0	1.7
MAINE	54.8	56.3	1.4
MARYLAND	55.4	56.7	1.3
MASSACHUSETTS	50.3	52.8	2.5
MICHIGAN	59.6	61.0	1.5
MINNESOTA	55.9	55.0	-0.8
MISSISSIPPI	62.3	61.7	-0.6
MISSOURI	57.7	56.4	-1.2
MONTANA	52.7	53.1	0.4
NEBRASKA	57.8	58.5	0.7
NEVADA	53.7	53.2	-0.4
NEW HAMPSHIRE	50.1	54.6	4.5
NEW JERSEY	54.5	56.7	2.2
NEW MEXICO	55.3	55.5	0.3
NEW YORK	54.0	56.9	2.9
NORTH CAROLINA	57.9	59.2	1.3
NORTH DAKOTA	60.1	60.4	0.3
OHIO	56.9	57.3	0.3
OKLAHOMA	57.8	56.3	-1.6
OREGON	54.5	57.6	3.1
PENNSYLVANIA	56.2	57.7	1.4
RHODE ISLAND	54.2	53.8	-0.5
SOUTH CAROLINA	58.3	59.0	0.7
SOUTH DAKOTA	59.3	58.8	-0.5
TENNESSEE	56.6	59.4	2.8
TEXAS	58.3	59.8	1.5
UTAH	50.9	54.1	3.2
VERMONT	52.3	52.8	0.5
VIRGINIA	58.7	56.1	-2.5
WASHINGTON	54.4	55.1	0.8
WEST VIRGINIA	62.0	59.8	-2.2
WISCONSIN	55.7	57.8	2.1
WYOMING	55.4	55.1	-0.3
PUERTO RICO	58.5	61.0	2.6

* BODY MASS INDEX GE 25.0 AND LE 99.8 FOR MALES AND FEMALES

April 30, 2001

TABLE 16.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TEEN CONDOM USE, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1764	12.1	1.0	(10.2, 14.0)
ALASKA	1827	9.8	1.1	(7.6, 12.0)
ARIZONA	1965	10.8	1.2	(8.5, 13.1)
ARKANSAS	2291	14.2	0.9	(12.5, 15.9)
CALIFORNIA	3239	7.2	0.6	(6.1, 8.3)
COLORADO	2504	7.5	0.7	(6.1, 8.9)
CONNECTICUT	3128	9.1	0.6	(7.9, 10.3)
DELAWARE	2196	9.2	0.8	(7.6, 10.7)
DISTRICT OF CO	1423	5.8	0.7	(4.4, 7.2)
FLORIDA	3829	8.1	0.5	(7.1, 9.1)
GEORGIA	3322	12.8	0.8	(11.3, 14.3)
HAWAII	4719	10.0	0.6	(8.9, 11.1)
IDAHO	4082	12.0	0.6	(10.8, 13.2)
ILLINOIS	3115	11.2	0.7	(9.8, 12.5)
INDIANA	2277	14.1	0.8	(12.5, 15.7)
IOWA	2668	9.2	0.6	(8.0, 10.5)
KANSAS	3261	13.0	0.7	(11.7, 14.3)
KENTUCKY	4839	11.9	0.6	(10.6, 13.1)
LOUISIANA	4023	12.6	0.7	(11.3, 13.9)
MAINE	3540	8.2	0.7	(6.7, 9.6)
MARYLAND	3749	10.5	0.7	(9.2, 11.7)
MASSACHUSETTS	6530	6.9	0.4	(6.1, 7.6)
MICHIGAN	2086	9.7	0.7	(8.3, 11.1)
MINNESOTA	2272	26.8	1.1	(24.6, 29.0)
MISSISSIPPI	1736	13.7	1.0	(11.8, 15.7)
MISSOURI	3315	13.8	0.8	(12.2, 15.4)
MONTANA	2335	17.2	1.1	(15.1, 19.3)
NEBRASKA	2228	12.9	0.8	(11.3, 14.5)
NEVADA	1748	11.1	1.1	(8.9, 13.2)
NEW HAMPSHIRE	1536	7.4	0.8	(5.9, 8.9)
NEW JERSEY	2985	9.2	0.6	(8.0, 10.4)
NEW MEXICO	2573	8.0	0.6	(6.9, 9.1)
NEW YORK	2707	8.2	0.7	(6.9, 9.6)
NORTH CAROLINA	2342	14.5	0.9	(12.7, 16.3)
NORTH DAKOTA	1477	13.1	0.9	(11.2, 14.9)
OHIO	2568	13.4	1.0	(11.4, 15.3)
OKLAHOMA	2800	15.0	0.8	(13.5, 16.4)
OREGON	2989	9.5	0.6	(8.2, 10.7)
PENNSYLVANIA	2798	10.5	0.7	(9.2, 11.7)
RHODE ISLAND	2769	7.3	0.6	(6.1, 8.4)
SOUTH CAROLINA	2730	11.3	0.7	(9.9, 12.6)
SOUTH DAKOTA	3793	11.9	0.6	(10.8, 13.1)
TENNESSEE	2355	21.0	1.0	(19.0, 22.9)
TEXAS	4155	11.4	0.6	(10.3, 12.4)
UTAH	2395	14.8	1.0	(12.9, 16.7)
VERMONT	2876	6.1	0.5	(5.1, 7.0)
VIRGINIA	1646	9.5	0.8	(7.9, 11.0)
WASHINGTON	2929	8.6	0.6	(7.4, 9.7)
WEST VIRGINIA	1838	12.0	0.8	(10.3, 13.6)
WISCONSIN	2178	7.2	0.6	(6.1, 8.4)
WYOMING	1959	11.9	0.8	(10.3, 13.5)
PUERTO RICO	3121	13.6	0.8	(12.1, 15.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 11.1 RANGE = 5.8-26.8

* RESPONDENT WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM April 30, 2001

TABLE 16.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR TEEN CONDOM USE, AGE 18-64 YEARS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*				-----FEMALES-----*			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	686	13.2	1.6	(10.0, 16.3)	1078	11.1	1.0	(9.0, 13.1)
ALASKA	874	11.8	1.8	(8.3, 15.4)	953	7.5	1.2	(5.1, 9.8)
ARIZONA	798	9.7	1.7	(6.3, 13.1)	1167	11.9	1.6	(8.8, 15.0)
ARKANSAS	930	16.6	1.4	(13.9, 19.3)	1361	11.9	1.0	(9.9, 13.8)
CALIFORNIA	1385	6.2	0.7	(4.8, 7.6)	1854	8.2	0.9	(6.5, 9.9)
COLORADO	1077	7.9	1.1	(5.8, 9.9)	1427	7.1	0.9	(5.3, 8.9)
CONNECTICUT	1251	10.6	1.0	(8.6, 12.5)	1877	7.7	0.7	(6.3, 9.0)
DELAWARE	916	10.1	1.3	(7.6, 12.5)	1280	8.3	1.0	(6.3, 10.2)
DISTRICT OF CO	577	5.7	1.1	(3.6, 7.9)	846	5.9	0.9	(4.0, 7.7)
FLORIDA	1605	7.8	0.8	(6.3, 9.3)	2224	8.3	0.7	(7.0, 9.7)
GEORGIA	1303	14.2	1.2	(11.8, 16.5)	2019	11.4	0.9	(9.6, 13.2)
HAWAII	2180	8.5	0.8	(7.0, 10.0)	2539	11.6	0.8	(9.9, 13.2)
IDAHO	1776	12.5	0.9	(10.7, 14.2)	2306	11.5	0.8	(10.0, 13.1)
ILLINOIS	1340	12.7	1.1	(10.6, 14.8)	1775	9.6	0.8	(8.0, 11.2)
INDIANA	1001	15.4	1.2	(13.0, 17.8)	1276	12.8	1.1	(10.8, 14.9)
IOWA	1129	9.5	1.0	(7.5, 11.4)	1539	9.0	0.8	(7.4, 10.6)
KANSAS	1348	13.8	1.0	(11.8, 15.9)	1913	12.1	0.8	(10.5, 13.7)
KENTUCKY	2015	12.4	1.0	(10.5, 14.3)	2824	11.3	0.8	(9.8, 12.9)
LOUISIANA	1564	14.8	1.1	(12.6, 17.0)	2459	10.4	0.7	(9.1, 11.8)
MAINE	1509	9.2	1.2	(6.8, 11.6)	2031	7.1	0.8	(5.5, 8.7)
MARYLAND	1493	10.0	1.1	(7.9, 12.0)	2256	10.9	0.8	(9.4, 12.5)
MASSACHUSETTS	2668	7.0	0.6	(5.9, 8.2)	3862	6.7	0.5	(5.7, 7.7)
MICHIGAN	922	9.1	1.0	(7.1, 11.0)	1164	10.4	1.0	(8.4, 12.4)
MINNESOTA	1042	30.3	1.6	(27.1, 33.5)	1230	23.3	1.4	(20.6, 26.0)
MISSISSIPPI	661	17.8	1.8	(14.4, 21.3)	1075	9.9	1.0	(8.0, 11.8)
MISSOURI	1337	16.7	1.4	(13.9, 19.4)	1978	11.0	0.9	(9.2, 12.8)
MONTANA	1032	19.3	1.7	(16.0, 22.7)	1303	15.1	1.3	(12.6, 17.5)
NEBRASKA	930	13.9	1.3	(11.4, 16.4)	1298	11.9	1.0	(10.0, 13.7)
NEVADA	902	10.6	1.3	(8.1, 13.1)	846	11.6	1.8	(8.0, 15.2)
NEW HAMPSHIRE	629	8.5	1.3	(6.0, 11.0)	907	6.4	0.9	(4.7, 8.1)
NEW JERSEY	1202	9.6	1.0	(7.7, 11.5)	1783	8.9	0.8	(7.4, 10.3)
NEW MEXICO	1190	8.7	0.9	(6.9, 10.4)	1383	7.4	0.8	(5.9, 8.9)
NEW YORK	1108	8.9	1.1	(6.8, 11.0)	1599	7.6	0.8	(6.0, 9.2)
NORTH CAROLINA	927	15.7	1.5	(12.8, 18.6)	1415	13.4	1.1	(11.3, 15.4)
NORTH DAKOTA	652	12.8	1.4	(10.1, 15.6)	825	13.3	1.3	(10.9, 15.8)
OHIO	1034	15.8	1.7	(12.5, 19.1)	1534	11.0	1.2	(8.7, 13.3)
OKLAHOMA	1235	14.5	1.1	(12.3, 16.6)	1565	15.4	1.0	(13.4, 17.4)
OREGON	1226	9.0	0.9	(7.1, 10.8)	1763	10.0	0.8	(8.4, 11.5)
PENNSYLVANIA	1156	11.3	1.0	(9.3, 13.3)	1642	9.6	0.8	(8.0, 11.2)
RHODE ISLAND	1095	7.1	1.0	(5.2, 9.1)	1674	7.4	0.7	(6.1, 8.7)
SOUTH CAROLINA	1122	13.1	1.1	(10.9, 15.3)	1608	9.5	0.8	(7.9, 11.0)
SOUTH DAKOTA	1523	12.8	1.0	(10.9, 14.7)	2270	11.0	0.7	(9.6, 12.4)
TENNESSEE	802	21.8	1.6	(18.7, 24.9)	1553	20.2	1.2	(17.9, 22.4)
TEXAS	1671	12.6	0.9	(10.8, 14.3)	2484	10.1	0.7	(8.8, 11.4)
UTAH	1089	15.3	1.4	(12.5, 18.1)	1306	14.2	1.3	(11.7, 16.8)
VERMONT	1226	5.7	0.7	(4.3, 7.1)	1650	6.4	0.7	(5.1, 7.7)
VIRGINIA	738	10.2	1.3	(7.7, 12.7)	908	8.7	1.0	(6.8, 10.6)
WASHINGTON	1261	8.5	0.9	(6.8, 10.3)	1668	8.6	0.7	(7.1, 10.0)
WEST VIRGINIA	774	12.8	1.3	(10.3, 15.4)	1064	11.1	1.1	(9.0, 13.2)
WISCONSIN	1004	7.4	0.9	(5.7, 9.0)	1174	7.1	0.8	(5.6, 8.6)
WYOMING	854	14.7	1.3	(12.1, 17.2)	1105	9.2	1.0	(7.2, 11.1)
PUERTO RICO	1131	11.4	1.1	(9.2, 13.6)	1990	15.6	1.0	(13.6, 17.6)

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 11.6	MEDIAN	= 10.3
	RANGE	= 5.7-30.3	RANGE	= 5.9-23.3

* RESPONDENT WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

April 30, 2001

TABLE 16.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 TEEN CONDOM USE, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	9.5	12.1	2.6
ALASKA	10.3	9.8	-0.5
ARIZONA	6.3	10.8	4.5
ARKANSAS	12.6	14.2	1.6
CALIFORNIA	N/A	7.2	N/A
COLORADO	8.0	7.5	-0.5
CONNECTICUT	8.9	9.1	0.2
DELAWARE	7.6	9.2	1.5
DISTRICT OF CO	7.4	5.8	-1.6
FLORIDA	10.2	8.1	-2.1
GEORGIA	14.1	12.8	-1.3
HAWAII	9.6	10.0	0.4
IDAHO	11.8	12.0	0.2
ILLINOIS	12.2	11.2	-1.0
INDIANA	10.4	14.1	3.7
IOWA	7.7	9.2	1.5
KANSAS	12.4	13.0	0.6
KENTUCKY	11.4	11.9	0.4
LOUISIANA	13.1	12.6	-0.5
MAINE	8.0	8.2	0.2
MARYLAND	7.9	10.5	2.6
MASSACHUSETTS	8.6	6.9	-1.7
MICHIGAN	10.9	9.7	-1.2
MINNESOTA	28.9	26.8	-2.1
MISSISSIPPI	12.4	13.7	1.4
MISSOURI	10.3	13.8	3.5
MONTANA	15.5	17.2	1.6
NEBRASKA	11.2	12.9	1.7
NEVADA	10.2	11.1	0.9
NEW HAMPSHIRE	7.2	7.4	0.2
NEW JERSEY	9.6	9.2	-0.4
NEW MEXICO	10.6	8.0	-2.6
NEW YORK	8.7	8.2	-0.4
NORTH CAROLINA	12.4	14.5	2.1
NORTH DAKOTA	10.3	13.1	2.8
OHIO	8.6	13.4	4.7
OKLAHOMA	12.3	15.0	2.6
OREGON	10.8	9.5	-1.3
PENNSYLVANIA	11.7	10.5	-1.3
RHODE ISLAND	9.9	7.3	-2.7
SOUTH CAROLINA	12.6	11.3	-1.3
SOUTH DAKOTA	9.8	11.9	2.2
TENNESSEE	17.9	21.0	3.1
TEXAS	10.5	11.4	0.8
UTAH	11.6	14.8	3.2
VERMONT	6.6	6.1	-0.6
VIRGINIA	10.5	9.5	-1.0
WASHINGTON	8.4	8.6	0.2
WEST VIRGINIA	12.2	12.0	-0.2
WISCONSIN	8.5	7.2	-1.2
WYOMING	8.5	11.9	3.4
PUERTO RICO	15.1	13.6	-1.5

* RESPONDENT WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

April 30, 2001

49

TABLE 17.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCES OF GETTING HIV, AGE 18-64 YEARS*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1771	6.0	0.7	(4.6, 7.3)
ALASKA	1845	5.7	0.7	(4.2, 7.1)
ARIZONA	2003	5.1	0.9	(3.3, 6.9)
ARKANSAS	2301	6.1	0.6	(4.9, 7.2)
CALIFORNIA	3247	11.8	0.8	(10.3, 13.2)
COLORADO	2522	5.5	0.7	(4.2, 6.8)
CONNECTICUT	3136	6.2	0.5	(5.1, 7.2)
DELAWARE	2200	6.1	0.7	(4.8, 7.4)
DISTRICT OF CO	1399	11.7	1.0	(9.8, 13.7)
FLORIDA	3816	7.6	0.5	(6.5, 8.6)
GEORGIA	3302	6.6	0.6	(5.4, 7.8)
HAWAII	4723	7.9	0.6	(6.8, 9.0)
IDAHO	4144	4.3	0.4	(3.6, 5.0)
ILLINOIS	3128	6.9	0.6	(5.8, 8.0)
INDIANA	2312	5.1	0.6	(4.1, 6.2)
IOWA	2692	5.2	0.5	(4.2, 6.3)
KANSAS	3292	5.1	0.5	(4.1, 6.0)
KENTUCKY	4914	5.7	0.5	(4.7, 6.6)
LOUISIANA	4006	6.2	0.4	(5.4, 7.1)
MAINE	3560	4.9	0.6	(3.8, 6.0)
MARYLAND	3760	6.1	0.5	(5.1, 7.1)
MASSACHUSETTS	6540	6.6	0.4	(5.8, 7.4)
MICHIGAN	2081	6.5	0.6	(5.3, 7.7)
MINNESOTA	2297	6.8	0.6	(5.6, 7.9)
MISSISSIPPI	1711	6.3	0.7	(4.9, 7.6)
MISSOURI	3356	6.1	0.6	(4.9, 7.2)
MONTANA	2365	4.2	0.5	(3.2, 5.2)
NEBRASKA	2254	4.9	0.5	(3.9, 5.9)
NEVADA	1738	8.7	1.0	(6.7, 10.7)
NEW HAMPSHIRE	1547	5.3	0.8	(3.8, 6.8)
NEW JERSEY	2989	5.7	0.5	(4.7, 6.7)
NEW MEXICO	2592	7.0	0.6	(5.9, 8.1)
NEW YORK	2709	7.7	0.6	(6.5, 8.9)
NORTH CAROLINA	2341	5.2	0.6	(4.0, 6.4)
NORTH DAKOTA	1499	5.7	0.7	(4.4, 7.0)
OHIO	2603	6.4	0.7	(5.0, 7.8)
OKLAHOMA	2815	4.3	0.4	(3.5, 5.1)
OREGON	3004	6.1	0.5	(5.1, 7.2)
PENNSYLVANIA	2795	6.0	0.5	(4.9, 7.0)
RHODE ISLAND	2788	4.8	0.5	(3.9, 5.7)
SOUTH CAROLINA	2741	6.3	0.6	(5.1, 7.4)
SOUTH DAKOTA	3858	5.1	0.4	(4.3, 6.0)
TENNESSEE	2425	6.4	0.7	(5.1, 7.7)
TEXAS	4167	9.2	0.5	(8.2, 10.2)
UTAH	2422	4.6	0.5	(3.5, 5.6)
VERMONT	2896	6.0	0.5	(5.0, 7.0)
VIRGINIA	1641	5.2	0.6	(4.0, 6.4)
WASHINGTON	2966	5.2	0.5	(4.3, 6.1)
WEST VIRGINIA	1840	4.4	0.5	(3.4, 5.4)
WISCONSIN	2213	7.0	0.7	(5.7, 8.3)
WYOMING	1970	6.0	0.6	(4.9, 7.1)
PUERTO RICO	3087	9.3	0.8	(7.8, 10.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 6.0 RANGE = 4.2-11.8

* CHANCES OF GETTING INFECTED WITH HIV MEDIUM/HIGH

April 30, 2001

TABLE 17.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCES OF GETTING HIV, AGE 18-64 YEARS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	688	6.7	1.1	(4.5, 8.8)	*	1083	5.3	0.8	(3.7, 6.9)	*
ALASKA	885	6.4	1.1	(4.2, 8.6)	*	960	4.8	0.9	(3.1, 6.6)	*
ARIZONA	820	4.9	1.3	(2.3, 7.5)	*	1183	5.3	1.3	(2.7, 7.8)	*
ARKANSAS	930	6.3	1.0	(4.4, 8.2)	*	1371	5.9	0.7	(4.6, 7.3)	*
CALIFORNIA	1383	13.9	1.2	(11.7, 16.2)	*	1864	9.5	0.9	(7.6, 11.3)	*
COLORADO	1089	4.7	0.9	(3.0, 6.4)	*	1433	6.3	1.0	(4.5, 8.2)	*
CONNECTICUT	1259	7.2	0.9	(5.5, 8.9)	*	1877	5.1	0.6	(3.9, 6.3)	*
DELAWARE	913	6.4	1.1	(4.3, 8.4)	*	1287	5.9	0.9	(4.2, 7.6)	*
DISTRICT OF CO	561	15.5	1.7	(12.1, 18.8)	*	838	8.4	1.1	(6.2, 10.6)	*
FLORIDA	1603	8.0	0.8	(6.4, 9.6)	*	2213	7.1	0.7	(5.8, 8.5)	*
GEORGIA	1293	7.7	1.0	(5.8, 9.6)	*	2009	5.6	0.8	(4.1, 7.1)	*
HAWAII	2190	9.0	0.9	(7.2, 10.8)	*	2533	6.7	0.6	(5.5, 8.0)	*
IDAHO	1797	4.9	0.6	(3.8, 6.1)	*	2347	3.8	0.4	(2.9, 4.6)	*
ILLINOIS	1344	7.8	0.9	(6.1, 9.6)	*	1784	5.9	0.7	(4.5, 7.3)	*
INDIANA	1016	5.0	0.8	(3.5, 6.5)	*	1296	5.3	0.8	(3.7, 6.9)	*
IOWA	1141	5.6	0.8	(3.9, 7.2)	*	1551	4.9	0.7	(3.5, 6.3)	*
KANSAS	1369	5.3	0.7	(3.9, 6.6)	*	1923	4.9	0.7	(3.5, 6.3)	*
KENTUCKY	2068	6.2	0.8	(4.7, 7.7)	*	2846	5.2	0.6	(4.1, 6.3)	*
LOUISIANA	1555	7.0	0.7	(5.6, 8.4)	*	2451	5.5	0.5	(4.5, 6.5)	*
MAINE	1519	5.9	0.9	(4.1, 7.7)	*	2041	4.0	0.7	(2.6, 5.3)	*
MARYLAND	1496	6.9	0.8	(5.3, 8.4)	*	2264	5.3	0.6	(4.2, 6.5)	*
MASSACHUSETTS	2669	7.9	0.7	(6.6, 9.3)	*	3871	5.3	0.5	(4.4, 6.2)	*
MICHIGAN	918	6.2	0.9	(4.5, 7.9)	*	1163	6.8	0.8	(5.2, 8.4)	*
MINNESOTA	1058	7.9	0.9	(6.1, 9.6)	*	1239	5.7	0.7	(4.3, 7.1)	*
MISSISSIPPI	653	7.7	1.2	(5.3, 10.2)	*	1058	4.9	0.7	(3.5, 6.3)	*
MISSOURI	1356	7.3	1.0	(5.4, 9.3)	*	2000	4.9	0.7	(3.6, 6.2)	*
MONTANA	1044	4.5	0.8	(2.9, 6.0)	*	1321	3.9	0.7	(2.6, 5.2)	*
NEBRASKA	936	5.0	0.7	(3.5, 6.4)	*	1318	4.9	0.7	(3.5, 6.3)	*
NEVADA	894	10.2	1.6	(7.1, 13.2)	*	844	7.1	1.2	(4.7, 9.5)	*
NEW HAMPSHIRE	639	6.1	1.3	(3.5, 8.6)	*	908	4.5	0.9	(2.8, 6.3)	*
NEW JERSEY	1201	7.2	0.9	(5.6, 8.9)	*	1788	4.2	0.5	(3.1, 5.2)	*
NEW MEXICO	1199	7.4	0.9	(5.7, 9.1)	*	1393	6.6	0.7	(5.2, 8.1)	*
NEW YORK	1101	9.2	1.0	(7.2, 11.2)	*	1608	6.3	0.7	(5.0, 7.6)	*
NORTH CAROLINA	926	5.6	1.1	(3.6, 7.7)	*	1415	4.7	0.6	(3.5, 5.9)	*
NORTH DAKOTA	669	5.1	0.9	(3.2, 6.9)	*	830	6.4	0.9	(4.5, 8.2)	*
OHIO	1054	7.1	1.0	(5.1, 9.1)	*	1549	5.8	1.0	(3.9, 7.7)	*
OKLAHOMA	1240	4.4	0.6	(3.2, 5.6)	*	1575	4.1	0.5	(3.1, 5.2)	*
OREGON	1230	7.6	0.9	(5.7, 9.4)	*	1774	4.7	0.6	(3.6, 5.8)	*
PENNSYLVANIA	1156	6.3	0.9	(4.6, 8.1)	*	1639	5.6	0.6	(4.4, 6.8)	*
RHODE ISLAND	1109	5.4	0.7	(3.9, 6.8)	*	1679	4.3	0.6	(3.2, 5.5)	*
SOUTH CAROLINA	1125	7.2	1.0	(5.3, 9.1)	*	1616	5.4	0.6	(4.2, 6.6)	*
SOUTH DAKOTA	1552	5.6	0.7	(4.3, 7.0)	*	2306	4.6	0.5	(3.6, 5.7)	*
TENNESSEE	827	7.3	1.2	(5.0, 9.6)	*	1598	5.5	0.7	(4.2, 6.8)	*
TEXAS	1689	10.0	0.8	(8.4, 11.6)	*	2478	8.5	0.6	(7.2, 9.7)	*
UTAH	1098	5.8	0.9	(4.0, 7.5)	*	1324	3.4	0.6	(2.2, 4.5)	*
VERMONT	1232	6.3	0.8	(4.7, 7.9)	*	1664	5.7	0.7	(4.3, 7.1)	*
VIRGINIA	740	5.8	1.0	(3.8, 7.7)	*	901	4.6	0.8	(3.1, 6.0)	*
WASHINGTON	1277	6.2	0.7	(4.7, 7.6)	*	1689	4.2	0.6	(3.1, 5.3)	*
WEST VIRGINIA	776	3.9	0.8	(2.3, 5.4)	*	1064	5.0	0.7	(3.6, 6.3)	*
WISCONSIN	1028	7.1	1.0	(5.1, 9.0)	*	1185	6.9	0.9	(5.1, 8.6)	*
WYOMING	862	5.9	0.9	(4.3, 7.6)	*	1108	6.0	0.8	(4.5, 7.5)	*
PUERTO RICO	1125	9.9	1.3	(7.5, 12.4)	*	1962	8.7	0.9	(6.9, 10.4)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 6.4
 RANGE = 3.9-15.5

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 5.3
 RANGE = 3.4- 9.5

* CHANCES OF GETTING INFECTED WITH HIV MEDIUM/HIGH

April 30, 2001

TABLE 17.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 CHANCES OF GETTING HIV, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	5.0	6.0	1.0
ALASKA	6.1	5.7	-0.4
ARIZONA	2.7	5.1	2.4
ARKANSAS	6.4	6.1	-0.3
CALIFORNIA	N/A	11.8	N/A
COLORADO	5.9	5.5	-0.4
CONNECTICUT	8.0	6.2	-1.8
DELAWARE	7.4	6.1	-1.3
DISTRICT OF CO	10.6	11.7	1.2
FLORIDA	7.5	7.6	0.0
GEORGIA	6.8	6.6	-0.2
HAWAII	8.9	7.9	-0.9
IDAHO	5.0	4.3	-0.7
ILLINOIS	6.7	6.9	0.2
INDIANA	7.0	5.1	-1.9
IOWA	7.6	5.2	-2.4
KANSAS	7.0	5.1	-1.9
KENTUCKY	6.6	5.7	-0.9
LOUISIANA	6.8	6.2	-0.6
MAINE	4.5	4.9	0.4
MARYLAND	7.1	6.1	-1.0
MASSACHUSETTS	8.4	6.6	-1.8
MICHIGAN	6.9	6.5	-0.4
MINNESOTA	8.8	6.8	-2.0
MISSISSIPPI	7.5	6.3	-1.2
MISSOURI	6.1	6.1	-0.0
MONTANA	3.9	4.2	0.3
NEBRASKA	6.8	4.9	-1.9
NEVADA	7.9	8.7	0.8
NEW HAMPSHIRE	5.8	5.3	-0.5
NEW JERSEY	8.0	5.7	-2.3
NEW MEXICO	7.6	7.0	-0.6
NEW YORK	7.3	7.7	0.4
NORTH CAROLINA	6.0	5.2	-0.8
NORTH DAKOTA	5.6	5.7	0.1
OHIO	7.4	6.4	-1.0
OKLAHOMA	5.0	4.3	-0.8
OREGON	5.3	6.1	0.9
PENNSYLVANIA	5.6	6.0	0.4
RHODE ISLAND	5.1	4.8	-0.3
SOUTH CAROLINA	6.7	6.3	-0.4
SOUTH DAKOTA	4.9	5.1	0.3
TENNESSEE	5.9	6.4	0.5
TEXAS	8.0	9.2	1.2
UTAH	4.5	4.6	0.0
VERMONT	5.0	6.0	1.0
VIRGINIA	6.4	5.2	-1.2
WASHINGTON	3.8	5.2	1.4
WEST VIRGINIA	6.9	4.4	-2.5
WISCONSIN	7.0	7.0	0.0
WYOMING	6.4	6.0	-0.4
PUERTO RICO	11.8	9.3	-2.5

* CHANCES OF GETTING INFECTED WITH HIV MEDIUM/HIGH

April 30, 2001

TABLE 18.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HIV TEST IN PAST 12 MONTHS, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	311	63.3	3.5	(56.5, 70.0)
ALASKA	263	64.8	4.7	(55.7, 74.0)
ARIZONA	254	64.9	5.3	(54.6, 75.3)
ARKANSAS	371	65.0	3.1	(59.0, 71.0)
CALIFORNIA	509	68.9	2.6	(63.7, 74.0)
COLORADO	387	69.3	3.1	(63.2, 75.3)
CONNECTICUT	454	63.4	2.6	(58.4, 68.5)
DELAWARE	411	57.9	3.4	(51.3, 64.5)
DISTRICT OF CO	264	53.7	3.6	(46.6, 60.7)
FLORIDA	690	59.0	2.2	(54.6, 63.4)
GEORGIA	572	56.4	2.9	(50.6, 62.1)
HAWAII	549	49.8	2.8	(44.3, 55.3)
IDAHO	557	68.7	2.4	(64.1, 73.3)
ILLINOIS	455	66.4	2.6	(61.4, 71.5)
INDIANA	324	68.4	2.9	(62.6, 74.1)
IOWA	284	67.3	3.3	(60.8, 73.8)
KANSAS	391	59.2	3.0	(53.3, 65.0)
KENTUCKY	566	62.5	2.8	(57.1, 67.9)
LOUISIANA	699	62.2	2.1	(58.2, 66.2)
MAINE	454	69.3	3.5	(62.4, 76.1)
MARYLAND	720	61.0	2.4	(56.2, 65.7)
MASSACHUSETTS	1049	64.1	1.9	(60.4, 67.7)
MICHIGAN	313	65.5	3.2	(59.2, 71.8)
MINNESOTA	202	68.7	4.0	(60.8, 76.5)
MISSISSIPPI	320	52.9	3.2	(46.6, 59.2)
MISSOURI	473	59.0	3.3	(52.6, 65.4)
MONTANA	262	67.4	3.9	(59.8, 74.9)
NEBRASKA	202	59.9	3.9	(52.3, 67.6)
NEVADA	334	71.8	3.1	(65.6, 77.9)
NEW HAMPSHIRE	257	65.0	3.6	(58.0, 71.9)
NEW JERSEY	432	62.9	2.7	(57.5, 68.3)
NEW MEXICO	351	67.3	2.9	(61.7, 72.9)
NEW YORK	398	60.0	2.9	(54.4, 65.6)
NORTH CAROLINA	352	61.7	3.3	(55.3, 68.1)
NORTH DAKOTA	168	58.0	4.1	(49.9, 66.1)
OHIO	352	60.2	3.8	(52.8, 67.6)
OKLAHOMA	378	65.9	2.8	(60.4, 71.3)
OREGON	441	66.2	2.7	(60.9, 71.4)
PENNSYLVANIA	412	65.7	2.7	(60.4, 71.1)
RHODE ISLAND	447	63.0	2.8	(57.5, 68.5)
SOUTH CAROLINA	502	54.4	2.8	(48.9, 59.8)
SOUTH DAKOTA	427	65.2	2.8	(59.6, 70.7)
TENNESSEE	365	62.7	2.9	(57.0, 68.4)
TEXAS	735	60.9	2.1	(56.7, 65.0)
UTAH	316	63.0	3.7	(55.7, 70.3)
VERMONT	404	73.6	2.4	(68.8, 78.3)
VIRGINIA	347	54.2	3.1	(48.2, 60.3)
WASHINGTON	488	64.6	2.5	(59.6, 69.5)
WEST VIRGINIA	202	66.8	3.6	(59.7, 73.9)
WISCONSIN	252	64.7	3.6	(57.6, 71.8)
WYOMING	278	67.8	3.3	(61.3, 74.2)
PUERTO RICO	359	65.4	3.4	(58.7, 72.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 64.3 RANGE = 49.8-73.6

* NO HIV TEST OUTSIDE OF BLOOD DONATIONS

April 30, 2001

TABLE 18.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HIV TEST IN PAST 12 MONTHS, AGE 18-64 YEARS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	143	64.8	4.5	(55.9, 73.7)	*	168	61.5	5.2	(51.2, 71.7)	*
ALASKA	161	58.6	6.2	(46.5, 70.7)	*	102	80.4	4.9	(70.7, 90.1)	*
ARIZONA	123	63.8	7.5	(49.2, 78.4)	*	131	66.8	6.6	(53.9, 79.8)	*
ARKANSAS	170	69.2	4.4	(60.6, 77.9)	*	201	60.2	4.1	(52.1, 68.3)	*
CALIFORNIA	256	64.2	3.8	(56.9, 71.6)	*	253	75.4	3.0	(69.5, 81.3)	*
COLORADO	202	69.6	4.0	(61.7, 77.5)	*	185	68.7	4.8	(59.4, 78.1)	*
CONNECTICUT	205	62.6	3.8	(55.1, 70.1)	*	249	64.4	3.4	(57.8, 71.0)	*
DELAWARE	209	54.0	4.9	(44.4, 63.6)	*	202	63.0	4.4	(54.5, 71.6)	*
DISTRICT OF CO	131	54.7	5.0	(44.9, 64.5)	*	133	52.3	5.1	(42.4, 62.2)	*
FLORIDA	355	56.3	3.1	(50.2, 62.5)	*	335	62.9	3.1	(56.9, 68.9)	*
GEORGIA	289	54.3	4.0	(46.5, 62.2)	*	283	59.6	4.0	(51.9, 67.4)	*
HAWAII	305	45.6	3.8	(38.1, 53.0)	*	244	56.6	4.2	(48.3, 64.9)	*
IDAHO	288	68.4	3.2	(62.1, 74.7)	*	269	69.1	3.4	(62.5, 75.7)	*
ILLINOIS	239	65.2	3.6	(58.1, 72.2)	*	216	68.3	3.5	(61.4, 75.1)	*
INDIANA	168	66.3	4.1	(58.2, 74.3)	*	156	71.1	4.1	(63.0, 79.2)	*
IOWA	142	66.4	4.5	(57.5, 75.2)	*	142	68.6	4.8	(59.1, 78.1)	*
KANSAS	188	56.6	4.3	(48.2, 65.0)	*	203	62.7	3.9	(55.1, 70.3)	*
KENTUCKY	277	64.2	3.8	(56.9, 71.6)	*	289	60.2	4.0	(52.3, 68.2)	*
LOUISIANA	303	61.7	3.0	(55.7, 67.6)	*	396	62.9	2.7	(57.6, 68.1)	*
MAINE	225	68.6	4.9	(58.9, 78.2)	*	229	70.4	4.6	(61.5, 79.3)	*
MARYLAND	321	58.4	3.6	(51.4, 65.4)	*	399	64.1	3.1	(58.0, 70.2)	*
MASSACHUSETTS	527	62.2	2.6	(57.2, 67.2)	*	522	66.8	2.6	(61.7, 71.9)	*
MICHIGAN	170	59.6	4.6	(50.6, 68.6)	*	143	74.1	3.9	(66.5, 81.8)	*
MINNESOTA	101	70.1	4.9	(60.6, 79.7)	*	101	67.0	6.3	(54.8, 79.3)	*
MISSISSIPPI	159	55.0	4.5	(46.1, 63.9)	*	161	50.1	4.4	(41.5, 58.7)	*
MISSOURI	214	60.6	4.5	(51.7, 69.4)	*	259	57.0	4.6	(47.9, 66.0)	*
MONTANA	135	72.7	5.0	(62.9, 82.4)	*	127	60.3	5.9	(48.8, 71.8)	*
NEBRASKA	104	55.0	5.4	(44.4, 65.5)	*	98	67.5	5.2	(57.2, 77.7)	*
NEVADA	193	65.1	4.4	(56.6, 73.6)	*	141	80.7	4.1	(72.7, 88.6)	*
NEW HAMPSHIRE	118	59.5	5.4	(49.0, 70.0)	*	139	71.1	4.4	(62.4, 79.8)	*
NEW JERSEY	226	62.2	3.7	(54.9, 69.5)	*	206	64.1	3.9	(56.5, 71.6)	*
NEW MEXICO	195	64.4	3.9	(56.9, 72.0)	*	156	71.5	4.2	(63.2, 79.7)	*
NEW YORK	191	59.1	4.1	(51.0, 67.1)	*	207	61.3	3.8	(53.8, 68.8)	*
NORTH CAROLINA	172	62.5	4.7	(53.3, 71.7)	*	180	60.6	4.3	(52.2, 69.0)	*
NORTH DAKOTA	98	57.6	5.4	(47.0, 68.3)	*	70	58.6	6.2	(46.6, 70.7)	*
OHIO	149	58.5	5.7	(47.4, 69.7)	*	203	62.0	4.9	(52.4, 71.7)	*
OKLAHOMA	194	66.8	3.7	(59.6, 74.1)	*	184	64.6	4.2	(56.4, 72.8)	*
OREGON	205	62.4	4.0	(54.5, 70.3)	*	236	70.6	3.3	(64.0, 77.1)	*
PENNSYLVANIA	190	62.7	4.1	(54.8, 70.6)	*	222	69.2	3.5	(62.3, 76.2)	*
RHODE ISLAND	203	58.5	4.1	(50.5, 66.5)	*	244	68.6	3.7	(61.2, 75.9)	*
SOUTH CAROLINA	258	53.0	4.0	(45.1, 60.8)	*	244	56.4	3.7	(49.1, 63.7)	*
SOUTH DAKOTA	220	59.0	3.9	(51.4, 66.7)	*	207	75.0	3.4	(68.4, 81.6)	*
TENNESSEE	170	60.1	4.1	(52.1, 68.2)	*	195	66.6	3.9	(58.8, 74.3)	*
TEXAS	357	59.4	3.0	(53.5, 65.3)	*	378	63.0	3.0	(57.2, 68.8)	*
UTAH	156	53.8	5.4	(43.2, 64.4)	*	160	74.8	4.4	(66.1, 83.4)	*
VERMONT	199	72.5	3.4	(65.8, 79.2)	*	205	74.9	3.5	(68.0, 81.7)	*
VIRGINIA	172	51.9	4.3	(43.4, 60.4)	*	175	57.4	4.4	(48.9, 65.9)	*
WASHINGTON	245	61.8	3.6	(54.6, 68.9)	*	243	68.3	3.3	(61.8, 74.8)	*
WEST VIRGINIA	106	70.2	4.8	(60.7, 79.6)	*	96	61.5	5.4	(50.9, 72.0)	*
WISCONSIN	132	64.4	5.1	(54.4, 74.4)	*	120	65.1	5.1	(55.1, 75.1)	*
WYOMING	140	69.8	4.4	(61.2, 78.3)	*	138	65.0	4.9	(55.4, 74.6)	*
PUERTO RICO	217	64.4	4.4	(55.7, 73.0)	*	142	67.4	4.9	(57.8, 77.0)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 62.2
 RANGE = 45.6-72.7

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 65.1
 RANGE = 50.1-80.7

* NO HIV TEST OUTSIDE OF BLOOD DONATIONS

April 30, 2001

TABLE 18.3 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000
 NO HIV TEST IN PAST 12 MONTHS, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	57.2	63.3	6.1
ALASKA	66.0	64.8	-1.2
ARIZONA	67.6	64.9	-2.7
ARKANSAS	64.5	65.0	0.5
CALIFORNIA	N/A	68.9	N/A
COLORADO	61.6	69.3	7.7
CONNECTICUT	64.4	63.4	-1.0
DELAWARE	61.4	57.9	-3.5
DISTRICT OF CO	49.7	53.7	4.0
FLORIDA	60.8	59.0	-1.8
GEORGIA	57.6	56.4	-1.2
HAWAII	50.1	49.8	-0.3
IDAHO	71.4	68.7	-2.7
ILLINOIS	N/A	66.4	N/A
INDIANA	57.4	68.4	10.9
IOWA	65.7	67.3	1.6
KANSAS	62.0	59.2	-2.8
KENTUCKY	56.7	62.5	5.8
LOUISIANA	53.3	62.2	8.9
MAINE	59.7	69.3	9.6
MARYLAND	63.1	61.0	-2.1
MASSACHUSETTS	72.1	64.1	-8.0
MICHIGAN	64.3	65.5	1.3
MINNESOTA	67.5	68.7	1.1
MISSISSIPPI	60.6	52.9	-7.7
MISSOURI	56.7	59.0	2.3
MONTANA	68.7	67.4	-1.3
NEBRASKA	66.8	59.9	-6.8
NEVADA	65.8	71.8	6.0
NEW HAMPSHIRE	59.9	65.0	5.1
NEW JERSEY	64.3	62.9	-1.4
NEW MEXICO	65.2	67.3	2.0
NEW YORK	52.5	60.0	7.5
NORTH CAROLINA	53.7	61.7	8.0
NORTH DAKOTA	60.6	58.0	-2.6
OHIO	73.3	60.2	-13.0
OKLAHOMA	60.0	65.9	5.9
OREGON	65.6	66.2	0.6
PENNSYLVANIA	64.1	65.7	1.6
RHODE ISLAND	59.0	63.0	4.0
SOUTH CAROLINA	56.7	54.4	-2.4
SOUTH DAKOTA	57.8	65.2	7.4
TENNESSEE	57.1	62.7	5.6
TEXAS	61.9	60.9	-1.1
UTAH	69.9	63.0	-6.9
VERMONT	68.9	73.6	4.6
VIRGINIA	54.7	54.2	-0.5
WASHINGTON	70.2	64.6	-5.7
WEST VIRGINIA	66.0	66.8	0.8
WISCONSIN	70.5	64.7	-5.8
WYOMING	64.5	67.8	3.2
PUERTO RICO	62.6	65.4	2.7

* NO HIV TEST OUTSIDE OF BLOOD DONATIONS

April 30, 2001

TABLE 19.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET OF DIABETES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	185	47.2	1.3	(44.7, 49.7)
ALASKA	68	41.2	2.7	(35.8, 46.6)
ARIZONA	133	51.2	2.0	(47.2, 55.2)
ARKANSAS	194	48.9	1.6	(45.7, 52.1)
CALIFORNIA	211	47.2	1.3	(44.6, 49.7)
COLORADO	143	51.1	1.8	(47.5, 54.6)
CONNECTICUT	212	49.2	1.1	(47.1, 51.3)
DISTRICT OF CO	110	48.7	1.5	(45.8, 51.5)
FLORIDA	342	51.8	0.9	(50.1, 53.6)
GEORGIA	285	46.3	1.1	(44.1, 48.6)
HAWAII	322	48.0	1.0	(46.0, 50.0)
IDAHO	245	49.1	1.3	(46.6, 51.5)
INDIANA	180	50.1	1.2	(47.7, 52.6)
IOWA	216	52.1	1.6	(49.0, 55.3)
KANSAS	241	49.1	1.1	(47.0, 51.3)
KENTUCKY	474	48.6	1.0	(46.6, 50.6)
LOUISIANA	326	47.1	1.1	(45.0, 49.3)
MAINE	276	48.7	1.6	(45.5, 51.9)
MASSACHUSETTS	458	49.4	0.9	(47.6, 51.3)
MICHIGAN	184	47.9	1.2	(45.6, 50.2)
MINNESOTA	136	52.1	1.5	(49.2, 54.9)
MISSISSIPPI	171	46.3	1.3	(43.7, 48.9)
MISSOURI	292	50.8	1.3	(48.2, 53.3)
MONTANA	187	52.5	1.7	(49.2, 55.8)
NEBRASKA	156	51.9	1.6	(48.8, 55.0)
NEVADA	113	53.6	2.1	(49.5, 57.7)
NEW HAMPSHIRE	89	50.2	1.6	(47.0, 53.3)
NEW JERSEY	222	48.0	1.1	(45.8, 50.2)
NEW MEXICO	205	48.2	1.0	(46.2, 50.2)
NORTH CAROLINA	222	50.7	1.2	(48.4, 53.0)
NORTH DAKOTA	99	49.4	2.5	(44.6, 54.2)
OHIO	217	50.2	1.7	(46.9, 53.6)
OKLAHOMA	211	46.9	1.3	(44.4, 49.5)
PENNSYLVANIA	222	50.8	1.3	(48.3, 53.2)
RHODE ISLAND	205	47.8	1.5	(44.9, 50.6)
SOUTH CAROLINA	213	47.2	1.1	(45.1, 49.3)
SOUTH DAKOTA	282	48.8	1.3	(46.2, 51.3)
TENNESSEE	198	47.2	1.1	(45.0, 49.3)
TEXAS	296	47.0	1.0	(45.1, 48.9)
UTAH	144	48.2	1.5	(45.2, 51.1)
VERMONT	168	50.3	1.4	(47.6, 53.0)
VIRGINIA	111	52.3	1.7	(49.0, 55.7)
WASHINGTON	195	47.5	1.5	(44.5, 50.4)
WEST VIRGINIA	192	51.2	1.0	(49.2, 53.1)
WISCONSIN	173	48.6	1.5	(45.6, 51.6)
WYOMING	112	49.3	1.9	(45.6, 53.0)
PUERTO RICO	450	47.5	1.0	(45.5, 49.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 47 MEDIAN = 48.9 RANGE = 41.2-53.6

* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

April 30, 2001

TABLE 19.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET OF DIABETES* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	70	49.3	2.0	(45.4, 53.2)	*	115	45.4	1.6	(42.3, 48.5)	*
ALASKA	35	37.6	4.2	(29.3, 45.9)	*	33	44.8	3.0	(38.9, 50.7)	*
ARIZONA	60	49.6	2.9	(43.9, 55.3)	*	73	52.8	2.6	(47.7, 57.9)	*
ARKANSAS	72	46.6	2.8	(41.1, 52.0)	*	122	50.9	1.7	(47.6, 54.3)	*
CALIFORNIA	95	46.5	2.1	(42.5, 50.6)	*	116	48.0	1.4	(45.1, 50.8)	*
COLORADO	80	50.7	2.4	(46.0, 55.5)	*	63	51.6	2.7	(46.4, 56.7)	*
CONNECTICUT	90	51.2	1.4	(48.4, 53.9)	*	122	47.1	1.7	(43.9, 50.4)	*
DISTRICT OF CO	34	48.4	2.7	(43.1, 53.7)	*	76	48.8	1.6	(45.6, 52.0)	*
FLORIDA	142	52.5	1.2	(50.2, 54.8)	*	200	51.2	1.4	(48.6, 53.9)	*
GEORGIA	106	46.5	1.9	(42.7, 50.3)	*	179	46.2	1.4	(43.5, 48.9)	*
HAWAII	168	47.9	1.3	(45.4, 50.5)	*	154	48.0	1.6	(44.8, 51.2)	*
IDAHO	102	51.6	2.0	(47.8, 55.5)	*	143	46.8	1.5	(43.8, 49.7)	*
INDIANA	67	49.3	1.9	(45.5, 53.1)	*	113	50.8	1.6	(47.7, 54.0)	*
IOWA	84	51.9	2.7	(46.5, 57.3)	*	132	52.3	1.8	(48.8, 55.9)	*
KANSAS	94	50.2	1.6	(47.1, 53.2)	*	147	48.1	1.5	(45.1, 51.1)	*
KENTUCKY	201	47.6	1.5	(44.7, 50.5)	*	273	49.6	1.4	(46.8, 52.3)	*
LOUISIANA	108	48.0	1.8	(44.4, 51.5)	*	218	46.6	1.4	(43.9, 49.2)	*
MAINE	114	49.8	2.6	(44.8, 54.8)	*	162	47.5	2.0	(43.6, 51.3)	*
MASSACHUSETTS	191	50.6	1.3	(48.1, 53.2)	*	267	48.3	1.3	(45.7, 50.8)	*
MICHIGAN	91	49.8	1.5	(46.8, 52.7)	*	93	45.9	1.7	(42.5, 49.2)	*
MINNESOTA	60	52.5	2.0	(48.6, 56.5)	*	76	51.7	2.1	(47.6, 55.9)	*
MISSISSIPPI	62	46.3	2.0	(42.4, 50.3)	*	109	46.3	1.8	(42.7, 49.8)	*
MISSOURI	120	51.3	2.0	(47.4, 55.2)	*	172	50.1	1.6	(47.0, 53.3)	*
MONTANA	75	50.5	3.0	(44.6, 56.3)	*	112	54.1	1.9	(50.5, 57.7)	*
NEBRASKA	72	50.0	2.0	(46.0, 54.0)	*	84	54.1	2.4	(49.3, 58.8)	*
NEVADA	66	53.5	2.8	(48.1, 58.9)	*	47	53.9	3.2	(47.6, 60.1)	*
NEW HAMPSHIRE	41	47.9	2.0	(43.9, 51.9)	*	48	52.8	2.4	(48.1, 57.6)	*
NEW JERSEY	101	47.3	1.6	(44.2, 50.3)	*	121	48.8	1.6	(45.7, 51.9)	*
NEW MEXICO	93	49.4	1.4	(46.6, 52.2)	*	112	47.1	1.5	(44.2, 50.0)	*
NORTH CAROLINA	81	49.5	1.9	(45.8, 53.2)	*	141	51.8	1.5	(48.9, 54.7)	*
NORTH DAKOTA	46	47.6	4.1	(39.6, 55.5)	*	53	51.3	2.4	(46.5, 56.0)	*
OHIO	90	54.4	2.5	(49.4, 59.3)	*	127	46.4	2.0	(42.5, 50.3)	*
OKLAHOMA	85	48.8	1.9	(45.0, 52.6)	*	126	45.5	1.7	(42.2, 48.9)	*
PENNSYLVANIA	94	50.5	2.0	(46.6, 54.4)	*	128	51.0	1.6	(48.0, 54.1)	*
RHODE ISLAND	98	47.1	2.1	(43.0, 51.2)	*	107	48.6	2.0	(44.6, 52.5)	*
SOUTH CAROLINA	89	47.9	1.5	(45.0, 50.8)	*	124	46.5	1.5	(43.5, 49.5)	*
SOUTH DAKOTA	120	46.4	1.9	(42.6, 50.1)	*	162	51.4	1.8	(47.9, 54.9)	*
TENNESSEE	67	45.0	1.8	(41.4, 48.5)	*	131	49.0	1.3	(46.5, 51.5)	*
TEXAS	133	48.4	1.3	(45.8, 51.0)	*	163	45.6	1.4	(42.8, 48.4)	*
UTAH	61	49.4	1.9	(45.8, 53.0)	*	83	46.9	2.3	(42.4, 51.3)	*
VERMONT	73	48.9	2.1	(44.9, 53.0)	*	95	51.8	1.8	(48.3, 55.2)	*
VIRGINIA	46	53.2	2.8	(47.8, 58.5)	*	65	51.6	2.0	(47.6, 55.6)	*
WASHINGTON	79	47.6	2.6	(42.5, 52.6)	*	116	47.3	1.7	(43.9, 50.7)	*
WEST VIRGINIA	84	51.6	1.5	(48.8, 54.4)	*	108	50.7	1.3	(48.2, 53.3)	*
WISCONSIN	74	50.1	2.1	(46.1, 54.1)	*	99	47.3	2.2	(43.1, 51.5)	*
WYOMING	49	48.8	3.0	(43.0, 54.6)	*	63	49.8	2.2	(45.5, 54.2)	*
PUERTO RICO	178	44.5	1.4	(41.7, 47.3)	*	272	50.3	1.4	(47.6, 53.0)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 47
 MEDIAN = 49.3
 RANGE = 37.6-54.4

NUMBER OF PARTICIPANTS = 47
 MEDIAN = 48.8
 RANGE = 44.8-54.1

* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

April 30, 2001

TABLE 20.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETICS TAKING INSULIN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	189	35.1	4.1	(27.2, 43.0)
ALASKA	73	37.8	8.2	(21.7, 54.0)
ARIZONA	135	43.7	10.4	(23.4, 64.0)
ARKANSAS	198	21.2	3.7	(14.0, 28.4)
CALIFORNIA	214	20.3	3.3	(13.9, 26.7)
COLORADO	149	29.8	5.5	(19.0, 40.6)
CONNECTICUT	219	23.4	2.9	(17.6, 29.1)
DISTRICT OF CO	117	35.4	5.0	(25.7, 45.2)
FLORIDA	357	25.0	2.5	(20.2, 29.8)
GEORGIA	313	34.8	3.3	(28.3, 41.3)
HAWAII	335	18.7	2.6	(13.5, 23.8)
IDAHO	252	27.7	3.3	(21.2, 34.3)
INDIANA	186	29.3	3.6	(22.2, 36.4)
IOWA	231	35.8	3.8	(28.4, 43.2)
KANSAS	253	35.4	3.4	(28.7, 42.1)
KENTUCKY	492	30.8	3.0	(24.9, 36.7)
LOUISIANA	343	31.3	2.9	(25.7, 36.9)
MAINE	284	30.7	4.9	(21.0, 40.3)
MASSACHUSETTS	474	28.0	2.5	(23.0, 32.9)
MICHIGAN	193	28.7	3.5	(21.9, 35.6)
MINNESOTA	136	32.3	4.4	(23.7, 40.8)
MISSISSIPPI	183	34.3	4.0	(26.6, 42.1)
MISSOURI	306	28.9	3.9	(21.4, 36.5)
MONTANA	190	29.1	4.3	(20.7, 37.5)
NEBRASKA	159	24.5	4.3	(16.1, 32.8)
NEVADA	115	39.0	7.7	(23.9, 54.1)
NEW HAMPSHIRE	92	28.4	5.3	(18.0, 38.7)
NEW JERSEY	228	26.0	3.3	(19.6, 32.4)
NEW MEXICO	213	29.8	3.7	(22.4, 37.1)
NORTH CAROLINA	230	25.7	3.1	(19.6, 31.8)
NORTH DAKOTA	102	41.0	5.7	(29.8, 52.1)
OHIO	228	28.6	4.6	(19.5, 37.6)
OKLAHOMA	220	31.4	3.7	(24.0, 38.7)
PENNSYLVANIA	229	26.5	3.4	(19.8, 33.2)
RHODE ISLAND	213	36.7	3.8	(29.3, 44.1)
SOUTH CAROLINA	221	36.4	3.7	(29.0, 43.7)
SOUTH DAKOTA	288	37.1	3.3	(30.7, 43.5)
TENNESSEE	211	28.3	3.7	(21.1, 35.5)
TEXAS	307	24.5	2.7	(19.3, 29.8)
UTAH	147	33.4	4.9	(23.9, 43.0)
VERMONT	174	24.6	3.6	(17.6, 31.7)
VIRGINIA	118	36.4	6.8	(23.1, 49.7)
WASHINGTON	201	25.5	3.3	(19.0, 31.9)
WEST VIRGINIA	198	24.9	3.3	(18.5, 31.3)
WISCONSIN	178	35.4	4.4	(26.8, 43.9)
WYOMING	113	26.4	4.4	(17.9, 35.0)
PUERTO RICO	481	32.7	3.0	(26.8, 38.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 47 MEDIAN = 29.8 RANGE = 18.7-43.7

* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

April 17, 2001

TABLE 20.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETICS TAKING INSULIN* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	70	25.6	5.8	(14.1, 37.0)	*	119	43.5	5.1	(33.4, 53.6)	
ALASKA	36	50.6	11.0	(29.1, 72.1)	*	37	25.1	9.6	(6.3, 43.9)	
ARIZONA	62	53.1	15.2	(23.4, 82.8)	*	73	34.4	9.6	(15.6, 53.2)	
ARKANSAS	72	24.2	6.6	(11.3, 37.2)	*	126	18.6	3.6	(11.5, 25.7)	
CALIFORNIA	96	15.2	3.8	(7.7, 22.7)	*	118	26.5	5.1	(16.4, 36.5)	
COLORADO	81	35.4	7.9	(20.0, 50.9)	*	68	21.7	6.6	(8.8, 34.6)	
CONNECTICUT	94	18.9	3.9	(11.3, 26.6)	*	125	28.1	4.3	(19.7, 36.5)	
DISTRICT OF CO	35	32.8	8.5	(16.1, 49.6)	*	82	37.0	6.0	(25.1, 48.8)	
FLORIDA	146	18.9	3.4	(12.2, 25.6)	*	211	30.7	3.5	(23.9, 37.4)	
GEORGIA	115	28.3	4.9	(18.7, 38.0)	*	198	39.6	4.4	(30.9, 48.3)	
HAWAII	172	18.0	3.6	(10.9, 25.1)	*	163	19.5	3.9	(12.0, 27.1)	
IDAHO	103	28.5	5.4	(17.9, 39.2)	*	149	27.1	4.1	(19.1, 35.0)	
INDIANA	68	25.3	5.7	(14.2, 36.4)	*	118	32.3	4.7	(23.2, 41.4)	
IOWA	87	35.9	6.0	(24.1, 47.7)	*	144	35.7	4.7	(26.6, 44.9)	
KANSAS	102	28.7	5.2	(18.6, 38.9)	*	151	42.6	4.4	(34.0, 51.2)	
KENTUCKY	206	34.1	4.7	(24.9, 43.2)	*	286	27.4	3.7	(20.2, 34.5)	
LOUISIANA	112	22.5	4.4	(14.0, 31.1)	*	231	37.3	3.6	(30.2, 44.4)	
MAINE	117	35.6	7.8	(20.3, 50.9)	*	167	25.3	5.4	(14.7, 35.9)	
MASSACHUSETTS	195	28.3	3.8	(20.8, 35.8)	*	279	27.6	3.3	(21.1, 34.1)	
MICHIGAN	92	30.7	5.0	(20.9, 40.5)	*	101	26.7	4.9	(17.2, 36.2)	
MINNESOTA	60	25.0	6.1	(13.0, 37.0)	*	76	38.3	6.3	(26.0, 50.6)	
MISSISSIPPI	64	44.2	6.8	(30.8, 57.5)	*	119	27.0	4.4	(18.5, 35.6)	
MISSOURI	124	28.7	5.9	(17.2, 40.2)	*	182	29.2	4.8	(19.7, 38.6)	
MONTANA	76	31.5	7.2	(17.4, 45.6)	*	114	27.2	5.2	(17.1, 37.3)	
NEBRASKA	72	24.2	5.7	(13.1, 35.2)	*	87	24.8	6.4	(12.2, 37.4)	
NEVADA	67	42.2	9.7	(23.3, 61.1)	*	48	33.2	13.7	(6.4, 60.0)	
NEW HAMPSHIRE	42	31.5	8.2	(15.4, 47.5)	*	50	24.9	6.3	(12.5, 37.3)	
NEW JERSEY	102	31.5	5.1	(21.5, 41.5)	*	126	20.0	3.8	(12.5, 27.5)	
NEW MEXICO	96	28.8	5.4	(18.2, 39.5)	*	117	30.6	5.3	(20.3, 40.9)	
NORTH CAROLINA	82	25.4	5.1	(15.5, 35.4)	*	148	26.0	3.8	(18.5, 33.4)	
NORTH DAKOTA	46	55.1	8.1	(39.2, 71.0)	*	56	27.1	6.3	(14.7, 39.5)	
OHIO	93	25.6	7.2	(11.6, 39.7)	*	135	31.3	5.9	(19.7, 42.8)	
OKLAHOMA	88	25.6	5.2	(15.4, 35.9)	*	132	35.5	5.1	(25.5, 45.5)	
PENNSYLVANIA	96	23.4	4.8	(14.0, 32.7)	*	133	29.0	4.8	(19.7, 38.4)	
RHODE ISLAND	102	37.9	5.4	(27.2, 48.5)	*	111	35.4	5.1	(25.3, 45.4)	
SOUTH CAROLINA	90	33.9	5.6	(23.0, 44.8)	*	131	38.7	5.0	(28.9, 48.4)	
SOUTH DAKOTA	122	39.7	4.7	(30.4, 49.0)	*	166	34.3	4.5	(25.5, 43.2)	
TENNESSEE	70	28.5	6.4	(16.0, 41.0)	*	141	28.1	4.1	(20.0, 36.2)	
TEXAS	135	21.0	3.8	(13.6, 28.5)	*	172	27.9	3.7	(20.6, 35.2)	
UTAH	63	28.8	6.5	(16.1, 41.6)	*	84	38.1	7.1	(24.1, 52.1)	
VERMONT	76	21.6	5.1	(11.6, 31.6)	*	98	27.8	5.0	(18.0, 37.6)	
VIRGINIA	47	29.3	7.6	(14.4, 44.2)	*	71	41.6	9.8	(22.4, 60.8)	
WASHINGTON	81	28.6	5.3	(18.2, 39.0)	*	120	22.6	4.0	(14.8, 30.4)	
WEST VIRGINIA	85	19.8	4.6	(10.8, 28.7)	*	113	29.6	4.6	(20.5, 38.6)	
WISCONSIN	74	27.1	5.9	(15.5, 38.7)	*	104	42.2	6.0	(30.3, 54.0)	
WYOMING	49	19.8	5.6	(8.7, 30.8)	*	64	34.0	6.4	(21.4, 46.6)	
PUERTO RICO	186	31.0	4.6	(21.9, 40.0)	*	295	34.2	3.8	(26.7, 41.7)	

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 47
 MEDIAN = 28.6
 RANGE = 15.2-55.1

NUMBER OF PARTICIPANTS = 47
 MEDIAN = 29.2
 RANGE = 18.6-43.5

* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

April 17, 2001

TABLE 21.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR VISITS TO HEALTH PROFESSIONALS IN PAST YEAR*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	185	4.5	0.4	(3.8, 5.2)
ALASKA	71	3.9	0.5	(2.9, 4.9)
ARIZONA	128	4.2	0.8	(2.7, 5.8)
ARKANSAS	189	4.9	0.4	(4.1, 5.7)
CALIFORNIA	214	4.7	0.5	(3.8, 5.7)
COLORADO	147	3.7	0.5	(2.6, 4.7)
CONNECTICUT	212	5.3	0.5	(4.4, 6.2)
DISTRICT OF CO	112	5.7	0.8	(4.2, 7.1)
FLORIDA	330	5.1	0.3	(4.5, 5.7)
GEORGIA	284	4.9	0.4	(4.2, 5.6)
HAWAII	330	5.7	0.4	(4.9, 6.4)
IDAHO	243	5.1	0.7	(3.7, 6.5)
INDIANA	179	4.2	0.4	(3.6, 4.9)
IOWA	220	3.6	0.2	(3.1, 4.0)
KANSAS	246	4.1	0.3	(3.5, 4.7)
KENTUCKY	472	4.6	0.2	(4.1, 5.0)
LOUISIANA	322	4.7	0.3	(4.1, 5.2)
MAINE	279	3.8	0.3	(3.2, 4.4)
MASSACHUSETTS	461	4.6	0.3	(4.0, 5.2)
MICHIGAN	181	4.3	0.4	(3.5, 5.2)
MINNESOTA	133	4.2	0.4	(3.4, 5.0)
MISSISSIPPI	171	4.3	0.3	(3.7, 5.0)
MISSOURI	291	4.6	0.4	(3.8, 5.4)
MONTANA	182	4.6	0.4	(3.7, 5.4)
NEBRASKA	158	5.5	0.6	(4.4, 6.5)
NEVADA	110	3.9	0.5	(2.9, 5.0)
NEW HAMPSHIRE	88	4.4	0.5	(3.4, 5.3)
NEW JERSEY	223	6.3	0.7	(5.0, 7.6)
NEW MEXICO	204	5.6	0.7	(4.3, 7.0)
NORTH CAROLINA	223	4.6	0.3	(4.0, 5.3)
NORTH DAKOTA	100	3.9	0.5	(2.9, 4.8)
OHIO	219	4.7	0.7	(3.5, 6.0)
OKLAHOMA	212	4.4	0.3	(3.8, 5.0)
PENNSYLVANIA	223	5.3	0.5	(4.5, 6.2)
RHODE ISLAND	208	4.3	0.3	(3.7, 5.0)
SOUTH CAROLINA	210	4.0	0.4	(3.3, 4.7)
SOUTH DAKOTA	271	4.6	0.4	(3.9, 5.3)
TENNESSEE	196	4.2	0.3	(3.6, 4.7)
TEXAS	294	4.6	0.3	(3.9, 5.2)
UTAH	143	4.4	0.6	(3.3, 5.5)
VERMONT	168	4.3	0.4	(3.5, 5.1)
VIRGINIA	111	4.7	0.7	(3.2, 6.2)
WASHINGTON	198	4.5	0.6	(3.4, 5.5)
WEST VIRGINIA	195	5.3	0.5	(4.4, 6.2)
WISCONSIN	173	3.7	0.3	(3.1, 4.2)
WYOMING	109	3.9	0.7	(2.6, 5.2)
PUERTO RICO	462	6.8	0.3	(6.2, 7.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 47 MEDIAN = 4.6 RANGE = 3.6- 6.8

* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

April 17, 2001

TABLE 21.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR VISITS TO HEALTH PROFESSIONALS IN PAST YEAR* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	68	4.0	0.5	(3.1, 4.9)	*	117	5.0	0.5	(4.0, 6.0)	*
ALASKA	36	2.6	0.4	(1.8, 3.4)	*	35	5.3	0.8	(3.7, 6.9)	*
ARIZONA	60	5.1	1.1	(3.0, 7.1)	*	68	3.4	1.1	(1.2, 5.6)	*
ARKANSAS	70	4.9	0.7	(3.4, 6.3)	*	119	4.9	0.5	(4.1, 5.8)	*
CALIFORNIA	96	4.3	0.5	(3.3, 5.4)	*	118	5.2	0.8	(3.6, 6.8)	*
COLORADO	79	3.1	0.6	(1.9, 4.3)	*	68	4.5	1.0	(2.6, 6.3)	*
CONNECTICUT	90	5.6	0.8	(4.1, 7.1)	*	122	4.9	0.6	(3.9, 6.0)	*
DISTRICT OF CO	33	5.9	1.7	(2.6, 9.2)	*	79	5.5	0.7	(4.2, 6.9)	*
FLORIDA	140	4.9	0.5	(4.0, 5.8)	*	190	5.3	0.5	(4.4, 6.2)	*
GEORGIA	107	3.7	0.4	(3.0, 4.5)	*	177	5.7	0.5	(4.7, 6.7)	*
HAWAII	170	5.5	0.5	(4.6, 6.4)	*	160	5.9	0.7	(4.6, 7.1)	*
IDAHO	100	5.0	1.1	(2.9, 7.1)	*	143	5.2	1.0	(3.2, 7.1)	*
INDIANA	67	3.9	0.5	(2.9, 5.0)	*	112	4.5	0.5	(3.5, 5.4)	*
IOWA	84	3.6	0.3	(3.0, 4.2)	*	136	3.5	0.3	(3.0, 4.1)	*
KANSAS	101	3.9	0.5	(3.0, 4.8)	*	145	4.3	0.4	(3.5, 5.0)	*
KENTUCKY	198	4.5	0.3	(3.9, 5.2)	*	274	4.6	0.3	(4.0, 5.2)	*
LOUISIANA	108	4.3	0.5	(3.3, 5.3)	*	214	4.9	0.3	(4.3, 5.5)	*
MAINE	115	4.0	0.5	(3.0, 5.0)	*	164	3.6	0.4	(2.7, 4.4)	*
MASSACHUSETTS	190	4.8	0.6	(3.7, 5.9)	*	271	4.4	0.3	(3.8, 5.0)	*
MICHIGAN	90	4.0	0.4	(3.2, 4.9)	*	91	4.7	0.8	(3.2, 6.2)	*
MINNESOTA	59	4.1	0.6	(2.9, 5.4)	*	74	4.3	0.5	(3.3, 5.2)	*
MISSISSIPPI	62	4.5	0.5	(3.4, 5.6)	*	109	4.2	0.4	(3.4, 5.0)	*
MISSOURI	119	4.4	0.5	(3.4, 5.5)	*	172	4.8	0.6	(3.7, 5.9)	*
MONTANA	73	4.1	0.5	(3.0, 5.1)	*	109	4.9	0.7	(3.6, 6.3)	*
NEBRASKA	72	5.7	0.8	(4.2, 7.3)	*	86	5.1	0.8	(3.7, 6.6)	*
NEVADA	65	4.2	0.7	(2.9, 5.6)	*	45	3.4	0.6	(2.1, 4.6)	*
NEW HAMPSHIRE	42	4.5	0.6	(3.2, 5.7)	*	46	4.2	0.7	(2.9, 5.6)	*
NEW JERSEY	99	6.3	1.0	(4.4, 8.2)	*	124	6.3	0.9	(4.6, 7.9)	*
NEW MEXICO	94	5.9	1.1	(3.8, 8.0)	*	110	5.4	0.9	(3.6, 7.1)	*
NORTH CAROLINA	78	3.9	0.4	(3.0, 4.7)	*	145	5.3	0.5	(4.3, 6.2)	*
NORTH DAKOTA	46	3.5	0.6	(2.3, 4.7)	*	54	4.3	0.7	(2.9, 5.7)	*
OHIO	91	4.9	1.2	(2.6, 7.2)	*	128	4.6	0.6	(3.4, 5.7)	*
OKLAHOMA	87	3.7	0.4	(2.9, 4.4)	*	125	4.9	0.5	(4.0, 5.8)	*
PENNSYLVANIA	95	6.0	0.9	(4.3, 7.7)	*	128	4.8	0.4	(4.1, 5.5)	*
RHODE ISLAND	101	3.7	0.5	(2.8, 4.6)	*	107	5.1	0.5	(4.1, 6.0)	*
SOUTH CAROLINA	90	3.4	0.3	(2.8, 4.0)	*	120	4.6	0.7	(3.3, 6.0)	*
SOUTH DAKOTA	118	4.2	0.5	(3.3, 5.1)	*	153	5.0	0.5	(4.0, 6.0)	*
TENNESSEE	67	3.8	0.5	(3.0, 4.7)	*	129	4.4	0.4	(3.7, 5.1)	*
TEXAS	133	3.7	0.4	(2.9, 4.5)	*	161	5.5	0.5	(4.4, 6.5)	*
UTAH	62	3.7	0.7	(2.3, 5.1)	*	81	5.1	0.9	(3.3, 6.8)	*
VERMONT	75	4.6	0.7	(3.2, 6.0)	*	93	4.0	0.4	(3.3, 4.7)	*
VIRGINIA	44	4.3	0.8	(2.8, 5.8)	*	67	5.0	1.1	(2.8, 7.2)	*
WASHINGTON	80	4.6	0.9	(2.8, 6.3)	*	118	4.3	0.7	(3.0, 5.7)	*
WEST VIRGINIA	84	5.3	0.6	(4.1, 6.5)	*	111	5.4	0.7	(4.1, 6.6)	*
WISCONSIN	71	3.5	0.5	(2.6, 4.4)	*	102	3.8	0.4	(3.1, 4.5)	*
WYOMING	48	3.2	0.4	(2.4, 4.1)	*	61	4.6	1.3	(2.1, 7.2)	*
PUERTO RICO	177	6.0	0.4	(5.3, 6.8)	*	285	7.5	0.4	(6.7, 8.2)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 47
 MEDIAN = 4.3
 RANGE = 2.6- 6.3

NUMBER OF PARTICIPANTS = 47
 MEDIAN = 4.9
 RANGE = 3.4- 7.5

* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

April 17, 2001

TABLE 22.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	2080	26.2	1.4	(23.4, 29.0)
COLORADO	3019	21.0	1.1	(18.8, 23.1)
IDAHO	4935	18.3	0.7	(17.0, 19.6)
IOWA	3520	20.2	0.9	(18.5, 21.9)
LOUISIANA	4956	15.7	0.6	(14.5, 16.9)
MARYLAND	4538	10.6	0.6	(9.4, 11.7)
MISSISSIPPI	2125	24.0	1.1	(21.8, 26.3)
MONTANA	3015	24.5	1.1	(22.4, 26.7)
NEBRASKA	3005	17.5	0.8	(16.0, 19.1)
NEVADA	2100	16.4	1.1	(14.2, 18.5)
NORTH CAROLINA	3006	18.6	0.9	(16.9, 20.3)
OHIO	3204	16.9	1.0	(14.9, 18.9)
OKLAHOMA	3680	15.6	0.8	(14.2, 17.1)
TEXAS	4898	18.6	0.7	(17.3, 19.8)
VIRGINIA	1993	12.6	0.9	(10.8, 14.4)
WASHINGTON	3565	19.3	0.8	(17.8, 20.9)
WEST VIRGINIA	2344	28.4	1.1	(26.3, 30.4)
WYOMING	2385	30.7	1.1	(28.6, 32.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 18.6 RANGE = 10.6-30.7

* EVER USED OR TRIED PRODUCTS SUCH AS CHEWING TOBACCO OR SNUFF

April 17, 2001

TABLE 22.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALASKA	986	43.8	2.3	(39.3, 48.3)	*	1094	7.1	1.0	(5.2, 9.0)	*
COLORADO	1275	38.2	2.0	(34.4, 42.1)	*	1744	4.5	0.7	(3.2, 5.9)	*
IDAHO	2101	32.6	1.2	(30.3, 34.9)	*	2834	4.5	0.4	(3.7, 5.4)	*
IOWA	1389	39.0	1.5	(36.0, 41.9)	*	2131	3.0	0.5	(2.1, 3.9)	*
LOUISIANA	1863	29.6	1.2	(27.3, 31.9)	*	3093	3.2	0.3	(2.5, 3.9)	*
MARYLAND	1767	20.0	1.2	(17.7, 22.2)	*	2771	1.9	0.3	(1.3, 2.5)	*
MISSISSIPPI	802	44.3	2.0	(40.4, 48.1)	*	1323	6.1	0.7	(4.7, 7.6)	*
MONTANA	1287	46.1	1.9	(42.4, 49.7)	*	1728	4.0	0.6	(2.9, 5.2)	*
NEBRASKA	1210	32.4	1.5	(29.5, 35.4)	*	1795	3.7	0.5	(2.8, 4.7)	*
NEVADA	1045	30.0	2.0	(26.1, 33.9)	*	1055	2.6	0.5	(1.6, 3.5)	*
NORTH CAROLINA	1141	32.8	1.6	(29.5, 36.0)	*	1865	5.6	0.6	(4.6, 6.7)	*
OHIO	1251	32.4	1.9	(28.8, 36.1)	*	1953	2.9	0.6	(1.8, 4.0)	*
OKLAHOMA	1567	29.5	1.4	(26.8, 32.2)	*	2113	2.9	0.4	(2.1, 3.6)	*
TEXAS	1967	34.7	1.2	(32.3, 37.0)	*	2931	3.2	0.4	(2.5, 4.0)	*
VIRGINIA	853	24.4	1.7	(21.0, 27.8)	*	1140	1.5	0.4	(0.8, 2.3)	*
WASHINGTON	1480	35.0	1.4	(32.3, 37.8)	*	2085	4.3	0.5	(3.3, 5.3)	*
WEST VIRGINIA	950	54.5	1.8	(51.0, 57.9)	*	1394	5.1	0.6	(3.8, 6.3)	*
WYOMING	1009	53.3	1.7	(49.9, 56.7)	*	1376	8.5	0.9	(6.7, 10.3)	*

SUMMARY STATISTICS:
 NUMBER OF PARTICIPANTS = 18
 MEDIAN = 33.7
 RANGE = 20.0-54.5

NUMBER OF PARTICIPANTS = 18
 MEDIAN = 3.9
 RANGE = 1.5- 8.5

* EVER USED OR TRIED PRODUCTS SUCH AS CHEWING TOBACCO OR SNUFF

April 17, 2001

TABLE 23.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	2079	5.7	0.8	(4.1, 7.3)
COLORADO	3019	4.1	0.6	(2.9, 5.4)
IDAHO	4935	3.3	0.3	(2.7, 3.9)
IOWA	3517	3.0	0.4	(2.3, 3.8)
LOUISIANA	4956	3.5	0.3	(2.8, 4.1)
MARYLAND	4538	1.4	0.2	(1.0, 1.9)
MISSISSIPPI	2123	7.3	0.7	(6.0, 8.7)
MONTANA	3015	6.3	0.6	(5.1, 7.5)
NEBRASKA	3005	3.9	0.5	(3.0, 4.8)
NEVADA	2099	2.6	0.4	(1.9, 3.4)
NORTH CAROLINA	3005	5.2	0.5	(4.3, 6.1)
OHIO	3203	3.4	0.5	(2.4, 4.4)
OKLAHOMA	3680	4.5	0.5	(3.4, 5.5)
TEXAS	4897	4.1	0.4	(3.4, 4.8)
VIRGINIA	1993	3.1	0.5	(2.1, 4.0)
WASHINGTON	3565	3.1	0.4	(2.4, 3.9)
WEST VIRGINIA	2344	8.8	0.7	(7.5, 10.2)
WYOMING	2385	7.5	0.6	(6.2, 8.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 4.0 RANGE = 1.4- 8.8

April 17, 2001

TABLE 23.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALASKA	985	9.5	1.5	(6.5, 12.4)	*	1094	1.7	0.3	(1.0, 2.3)	*
COLORADO	1275	8.1	1.2	(5.7, 10.6)	*	1744	0.3	0.2	(-0.1, 0.8)	*
IDAHO	2101	6.5	0.6	(5.3, 7.6)	*	2834	0.3	0.1	(0.0, 0.5)	*
IOWA	1388	6.3	0.8	(4.8, 7.8)	*	2129	0.0	0.0	(0.0, 0.0)	*
LOUISIANA	1863	6.7	0.6	(5.4, 7.9)	*	3093	0.6	0.2	(0.3, 0.9)	*
MARYLAND	1767	2.8	0.5	(1.8, 3.8)	*	2771	0.2	0.1	(0.0, 0.3)	*
MISSISSIPPI	801	14.0	1.4	(11.3, 16.7)	*	1322	1.4	0.3	(0.7, 2.0)	*
MONTANA	1287	12.6	1.2	(10.2, 15.0)	*	1728	0.3	0.2	(0.0, 0.6)	*
NEBRASKA	1210	7.9	0.9	(6.1, 9.6)	*	1795	0.2	0.1	(-0.0, 0.4)	*
NEVADA	1044	5.2	0.8	(3.7, 6.7)	*	1055	0.0	0.0	(-0.0, 0.1)	*
NORTH CAROLINA	1140	8.9	0.9	(7.1, 10.7)	*	1865	1.8	0.3	(1.2, 2.4)	*
OHIO	1250	6.9	1.1	(4.7, 9.0)	*	1953	0.2	0.1	(-0.0, 0.5)	*
OKLAHOMA	1567	9.2	1.1	(7.1, 11.3)	*	2113	0.1	0.1	(-0.0, 0.3)	*
TEXAS	1966	8.3	0.7	(6.9, 9.6)	*	2931	0.2	0.1	(0.0, 0.4)	*
VIRGINIA	853	6.0	0.9	(4.1, 7.8)	*	1140	0.3	0.1	(0.0, 0.6)	*
WASHINGTON	1480	6.3	0.8	(4.9, 7.8)	*	2085	0.1	0.1	(-0.1, 0.2)	*
WEST VIRGINIA	950	17.9	1.3	(15.3, 20.5)	*	1394	0.7	0.2	(0.3, 1.2)	*
WYOMING	1009	14.3	1.2	(11.9, 16.7)	*	1376	0.8	0.3	(0.3, 1.3)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 18
 MEDIAN = 8.0
 RANGE = 2.8-17.9

NUMBER OF PARTICIPANTS = 18
 MEDIAN = 0.3
 RANGE = 0.0- 1.8

April 17, 2001

TABLE 24.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO FLU SHOT, AGE 65 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	623	28.8	2.8	(23.3, 34.4)
CALIFORNIA	576	30.4	2.4	(25.8, 35.0)
GEORGIA	648	36.4	2.4	(31.7, 41.0)
ILLINOIS	305	36.5	3.2	(30.3, 42.7)
IOWA	817	26.3	1.7	(22.9, 29.7)
KANSAS	850	32.0	1.8	(28.5, 35.4)
MINNESOTA	518	25.1	2.2	(20.8, 29.4)
MONTANA	626	27.7	2.3	(23.2, 32.2)
NEBRASKA	712	30.5	2.0	(26.7, 34.3)
NEW JERSEY	666	34.3	2.2	(30.1, 38.6)
NORTH CAROLINA	619	34.7	2.1	(30.6, 38.8)
NORTH DAKOTA	376	30.7	2.6	(25.6, 35.9)
OHIO	590	34.0	2.7	(28.7, 39.4)
SOUTH CAROLINA	504	30.3	2.3	(25.8, 34.8)
SOUTH DAKOTA	1066	29.1	1.6	(25.9, 32.3)
TENNESSEE	532	32.7	2.4	(28.1, 37.3)
VERMONT	668	30.7	1.9	(26.9, 34.5)
VIRGINIA	282	36.6	3.3	(30.2, 42.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 30.7 RANGE = 25.1-36.6 YEAR 2000 OBJ. 20.11
 * DID NOT HAVE FLU SHOT WITHIN PAST YEAR
 April 17, 2001

67

TABLE 25.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PNEUMONIA VACCINATION, AGE 65 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	612	26.9	2.7	(21.7, 32.2)
CALIFORNIA	567	39.5	2.6	(34.4, 44.6)
GEORGIA	633	39.8	2.4	(35.2, 44.4)
ILLINOIS	305	48.6	3.2	(42.3, 54.9)
IOWA	805	31.6	1.8	(28.0, 35.1)
KANSAS	835	41.4	1.9	(37.7, 45.1)
MINNESOTA	510	34.4	2.4	(29.7, 39.0)
MONTANA	617	35.1	2.5	(30.1, 40.0)
NEBRASKA	706	37.2	2.1	(33.1, 41.2)
NEW JERSEY	651	43.2	2.3	(38.7, 47.6)
NORTH CAROLINA	615	42.2	2.2	(37.9, 46.5)
NORTH DAKOTA	364	37.1	2.7	(31.7, 42.4)
OHIO	576	37.6	2.8	(32.1, 43.1)
SOUTH CAROLINA	493	40.3	2.5	(35.5, 45.2)
SOUTH DAKOTA	1035	46.7	1.8	(43.2, 50.1)
TENNESSEE	353	23.3	2.5	(18.5, 28.1)
VERMONT	650	34.4	2.0	(30.4, 38.4)
VIRGINIA	271	35.9	3.3	(29.4, 42.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 37.4 RANGE = 23.3-48.6 YEAR 2000 OBJ. 20.11
 * NEVER HAD PNEUMONIA VACCINATION
 April 17, 2001

TABLE 25.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PNEUMONIA VACCINATION, AGE 65 AND OVER* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ARIZONA	235	31.4	4.9	(21.8,	41.0)	377	23.7	3.0	(17.9,	29.5)
CALIFORNIA	219	43.7	4.4	(34.9,	52.4)	348	36.6	3.0	(30.6,	42.6)
GEORGIA	210	44.1	4.1	(36.0,	52.1)	423	37.0	2.8	(31.5,	42.4)
ILLINOIS	100	53.4	5.5	(42.6,	64.2)	205	45.2	3.8	(37.8,	52.5)
IOWA	236	30.5	3.2	(24.2,	36.7)	569	32.3	2.2	(28.0,	36.5)
KANSAS	245	41.9	3.4	(35.2,	48.6)	590	41.0	2.2	(36.7,	45.3)
MINNESOTA	178	37.8	4.0	(29.9,	45.6)	332	32.0	2.8	(26.4,	37.6)
MONTANA	221	36.4	4.1	(28.4,	44.4)	396	34.1	3.2	(27.8,	40.4)
NEBRASKA	249	35.4	3.4	(28.8,	42.1)	457	38.4	2.6	(33.2,	43.5)
NEW JERSEY	224	44.5	3.7	(37.2,	51.9)	427	42.3	2.8	(36.8,	47.7)
NORTH CAROLINA	195	43.2	3.8	(35.8,	50.6)	420	41.6	2.6	(36.4,	46.8)
NORTH DAKOTA	128	37.8	4.6	(28.8,	46.7)	236	36.5	3.4	(29.9,	43.1)
OHIO	194	40.3	5.0	(30.6,	50.1)	382	35.9	3.3	(29.4,	42.3)
SOUTH CAROLINA	165	43.4	4.2	(35.3,	51.5)	328	38.3	3.1	(32.3,	44.3)
SOUTH DAKOTA	334	41.7	2.8	(36.1,	47.3)	701	50.1	2.2	(45.8,	54.4)
TENNESSEE	82	18.5	4.3	(10.1,	26.9)	271	26.2	3.0	(20.4,	32.0)
VERMONT	247	37.2	3.3	(30.7,	43.6)	403	32.5	2.6	(27.5,	37.6)
VIRGINIA	87	42.1	6.1	(30.3,	54.0)	184	31.8	3.6	(24.8,	38.9)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 18
 MEDIAN = 41.0
 RANGE = 18.5-53.4

NUMBER OF PARTICIPANTS = 18
 MEDIAN = 36.6
 RANGE = 23.7-50.1

* NEVER HAD PNEUMONIA VACCINATION

April 17, 2001

69

TABLE 26.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD A HEART ATTACK OR MYOCARDIAL INFARCTION*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
DELAWARE	2720	4.2	0.5	(3.2, 5.1)
DISTRICT OF CO	1683	3.0	0.5	(2.0, 4.0)
GEORGIA	3999	3.7	0.3	(3.1, 4.4)
INDIANA	2903	5.2	0.4	(4.3, 6.0)
IOWA	3521	4.1	0.4	(3.4, 4.8)
KENTUCKY	6338	5.4	0.4	(4.7, 6.1)
MISSISSIPPI	2136	5.3	0.5	(4.3, 6.3)
MONTANA	3015	3.4	0.4	(2.7, 4.2)
OHIO	3201	5.4	0.6	(4.3, 6.6)
OKLAHOMA	3680	4.0	0.4	(3.3, 4.7)
PENNSYLVANIA	3508	4.6	0.4	(3.8, 5.3)
SOUTH CAROLINA	3273	4.5	0.5	(3.5, 5.4)
VIRGINIA	1958	4.2	0.5	(3.2, 5.1)
WEST VIRGINIA	2346	7.6	0.6	(6.5, 8.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 14 MEDIAN = 4.3 RANGE = 3.0- 7.6

* TOLD BY A DOCTOR THEY HAD A HEART ATTACK OR MYOCARDIAL INFARCTION

April 17, 2001

TABLE 26.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 70
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD A HEART ATTACK OR MYOCARDIAL INFARCTION* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
DELAWARE	1123	4.9	0.8	(3.4, 6.5)	*	1597	3.5	0.6	(2.3, 4.6)	*
DISTRICT OF CO	666	2.8	0.8	(1.2, 4.3)	*	1017	3.2	0.6	(2.0, 4.5)	*
GEORGIA	1535	4.6	0.6	(3.5, 5.7)	*	2464	2.9	0.4	(2.2, 3.6)	*
INDIANA	1202	6.4	0.7	(4.9, 7.9)	*	1701	4.0	0.5	(3.1, 5.0)	*
IOWA	1389	5.7	0.6	(4.4, 7.0)	*	2132	2.6	0.4	(1.9, 3.3)	*
KENTUCKY	2526	7.0	0.6	(5.8, 8.1)	*	3812	3.9	0.4	(3.2, 4.7)	*
MISSISSIPPI	806	6.2	0.9	(4.5, 7.8)	*	1330	4.6	0.6	(3.3, 5.8)	*
MONTANA	1289	4.3	0.7	(3.0, 5.7)	*	1726	2.6	0.4	(1.8, 3.4)	*
OHIO	1253	7.4	1.1	(5.4, 9.5)	*	1948	3.6	0.6	(2.5, 4.8)	*
OKLAHOMA	1568	4.1	0.5	(3.2, 5.1)	*	2112	3.9	0.5	(2.9, 4.9)	*
PENNSYLVANIA	1412	5.5	0.6	(4.2, 6.7)	*	2096	3.8	0.5	(2.8, 4.7)	*
SOUTH CAROLINA	1310	5.6	0.9	(3.9, 7.3)	*	1963	3.5	0.5	(2.6, 4.3)	*
VIRGINIA	845	5.3	0.8	(3.6, 6.9)	*	1113	3.1	0.6	(2.0, 4.2)	*
WEST VIRGINIA	950	9.5	1.0	(7.5, 11.4)	*	1396	6.0	0.7	(4.7, 7.3)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 14
 MEDIAN = 5.5
 RANGE = 2.8- 9.5

NUMBER OF PARTICIPANTS = 14
 MEDIAN = 3.5
 RANGE = 2.6- 6.0

* TOLD BY A DOCTOR THEY HAD A HEART ATTACK OR MYOCARDIAL INFARCTION

April 17, 2001

TABLE 27.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD YOU HAD A STROKE*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
DELAWARE	2723	2.3	0.4	(1.6, 3.0)
DISTRICT OF CO	1687	2.7	0.5	(1.8, 3.6)
GEORGIA	4009	2.2	0.3	(1.6, 2.7)
INDIANA	2913	2.5	0.3	(1.9, 3.1)
IOWA	3523	1.9	0.2	(1.5, 2.4)
KENTUCKY	6330	2.8	0.3	(2.3, 3.3)
MISSISSIPPI	2138	2.6	0.4	(1.9, 3.3)
MONTANA	3016	2.3	0.3	(1.6, 2.9)
OHIO	3208	2.5	0.4	(1.7, 3.2)
OKLAHOMA	3676	1.7	0.2	(1.3, 2.2)
PENNSYLVANIA	3515	2.4	0.3	(1.9, 3.0)
SOUTH CAROLINA	3273	1.7	0.2	(1.2, 2.2)
VIRGINIA	1962	2.1	0.4	(1.4, 2.8)
WEST VIRGINIA	2346	3.1	0.4	(2.3, 3.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 14 MEDIAN = 2.4 RANGE = 1.7- 3.1

* RESPONDENTS TOLD BY A DOCTOR THEY HAD A STROKE

April 17, 2001

TABLE 27.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD YOU HAD A STROKE* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
DELAWARE	1123	2.3	0.6	(1.2, 3.4)	*	1600	2.3	0.5	(1.4, 3.1)	*
DISTRICT OF CO	668	2.2	0.7	(0.9, 3.4)	*	1019	3.1	0.7	(1.8, 4.4)	*
GEORGIA	1544	2.5	0.5	(1.6, 3.4)	*	2465	1.9	0.3	(1.3, 2.4)	*
INDIANA	1207	2.4	0.5	(1.5, 3.3)	*	1706	2.6	0.4	(1.9, 3.3)	*
IOWA	1389	1.7	0.3	(1.0, 2.3)	*	2134	2.2	0.3	(1.5, 2.8)	*
KENTUCKY	2524	2.6	0.4	(1.9, 3.4)	*	3806	3.0	0.4	(2.3, 3.8)	*
MISSISSIPPI	807	1.8	0.5	(0.9, 2.7)	*	1331	3.4	0.6	(2.3, 4.5)	*
MONTANA	1289	2.1	0.5	(1.1, 3.0)	*	1727	2.5	0.4	(1.6, 3.3)	*
OHIO	1255	2.2	0.6	(1.1, 3.3)	*	1953	2.7	0.5	(1.6, 3.7)	*
OKLAHOMA	1564	1.4	0.3	(0.9, 2.0)	*	2112	2.0	0.3	(1.4, 2.6)	*
PENNSYLVANIA	1415	2.5	0.5	(1.6, 3.4)	*	2100	2.4	0.4	(1.6, 3.2)	*
SOUTH CAROLINA	1311	1.7	0.4	(0.9, 2.5)	*	1962	1.7	0.3	(1.1, 2.2)	*
VIRGINIA	846	1.8	0.5	(0.9, 2.7)	*	1116	2.3	0.5	(1.4, 3.3)	*
WEST VIRGINIA	951	2.9	0.6	(1.8, 4.0)	*	1395	3.3	0.5	(2.3, 4.2)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 14
 MEDIAN = 2.2
 RANGE = 1.4- 2.9

NUMBER OF PARTICIPANTS = 14
 MEDIAN = 2.4
 RANGE = 1.7- 3.4

* RESPONDENTS TOLD BY A DOCTOR THEY HAD A STROKE

April 17, 2001

TABLE 28.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1393	35.2	1.5	(32.2, 38.2)
ALASKA	1083	43.6	2.3	(39.2, 48.0)
ARIZONA	1574	37.6	2.4	(32.9, 42.3)
ARKANSAS	1834	38.0	1.3	(35.4, 40.5)
COLORADO	1745	40.9	1.7	(37.6, 44.2)
CONNECTICUT	2333	35.1	1.1	(32.9, 37.3)
DELAWARE	1606	33.3	1.6	(30.2, 36.4)
DISTRICT OF CO	1028	34.4	1.7	(31.1, 37.7)
FLORIDA	3049	32.1	1.0	(30.2, 34.0)
GEORGIA	2497	38.0	1.3	(35.6, 40.5)
HAWAII	3302	36.8	1.1	(34.6, 39.0)
IDAHO	2839	42.6	1.1	(40.4, 44.7)
ILLINOIS	2211	38.8	1.2	(36.5, 41.1)
INDIANA	1691	40.2	1.4	(37.5, 42.9)
IOWA	2143	38.0	1.2	(35.6, 40.4)
KANSAS	2556	37.4	1.1	(35.3, 39.5)
KENTUCKY	3820	37.0	1.1	(34.8, 39.1)
LOUISIANA	3078	38.2	1.0	(36.3, 40.1)
MAINE	2689	37.4	1.6	(34.3, 40.6)
MARYLAND	2758	35.9	1.1	(33.6, 38.1)
MASSACHUSETTS	4834	36.5	0.9	(34.9, 38.2)
MICHIGAN	1467	35.8	1.5	(32.9, 38.6)
MINNESOTA	1594	39.0	1.4	(36.2, 41.7)
MISSISSIPPI	1330	40.7	1.6	(37.4, 43.9)
MISSOURI	2669	39.0	1.3	(36.4, 41.6)
MONTANA	1726	38.9	1.7	(35.6, 42.2)
NEBRASKA	1773	38.1	1.4	(35.4, 40.7)
NEVADA	1048	35.7	2.1	(31.6, 39.9)
NEW HAMPSHIRE	1123	40.5	1.7	(37.2, 43.7)
NEW JERSEY	2261	38.4	1.2	(36.0, 40.7)
NEW MEXICO	1766	40.6	1.4	(38.0, 43.3)
NEW YORK	1970	36.2	1.3	(33.8, 38.7)
NORTH CAROLINA	1854	35.6	1.3	(33.1, 38.2)
NORTH DAKOTA	1082	39.6	1.6	(36.4, 42.8)
OHIO	1952	37.0	1.5	(34.0, 40.0)
OKLAHOMA	2099	41.7	1.2	(39.3, 44.1)
OREGON	2231	36.0	1.1	(33.8, 38.3)
PENNSYLVANIA	2092	34.6	1.2	(32.3, 36.9)
RHODE ISLAND	2138	33.5	1.2	(31.2, 35.9)
SOUTH CAROLINA	1969	35.4	1.3	(32.9, 37.8)
SOUTH DAKOTA	3047	39.9	1.0	(37.8, 41.9)
TENNESSEE	2019	37.3	1.2	(34.9, 39.6)
TEXAS	2974	40.6	1.0	(38.6, 42.6)
UTAH	1616	45.2	1.6	(42.0, 48.4)
VERMONT	2090	42.4	1.3	(39.9, 44.8)
VIRGINIA	1116	37.1	1.7	(33.7, 40.5)
WASHINGTON	2085	38.5	1.2	(36.2, 40.8)
WEST VIRGINIA	1388	36.3	1.4	(33.5, 39.0)
WISCONSIN	1484	37.6	1.5	(34.7, 40.5)
WYOMING	1369	39.7	1.5	(36.8, 42.7)
PUERTO RICO	2644	44.7	1.4	(42.0, 47.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 38.0 RANGE = 32.1-45.2

* NEVER HAD A MAMMOGRAM

April 17, 2001

TABLE 28.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO MAMMOGRAM, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	37.6	35.2	-2.5
ALASKA	44.3	43.6	-0.7
ARIZONA	34.2	37.6	3.4
ARKANSAS	41.8	38.0	-3.8
COLORADO	42.2	40.9	-1.2
CONNECTICUT	33.5	35.1	1.6
DELAWARE	33.8	33.3	-0.5
DISTRICT OF CO	40.7	34.4	-6.3
FLORIDA	31.0	32.1	1.1
GEORGIA	40.0	38.0	-1.9
HAWAII	38.2	36.8	-1.4
IDAHO	43.6	42.6	-1.0
ILLINOIS	39.1	38.8	-0.3
INDIANA	39.6	40.2	0.7
IOWA	38.6	38.0	-0.6
KANSAS	38.1	37.4	-0.7
KENTUCKY	38.6	37.0	-1.6
LOUISIANA	41.7	38.2	-3.5
MAINE	37.6	37.4	-0.2
MARYLAND	36.5	35.9	-0.7
MASSACHUSETTS	37.5	36.5	-1.0
MICHIGAN	36.5	35.8	-0.7
MINNESOTA	42.4	39.0	-3.4
MISSISSIPPI	41.2	40.7	-0.6
MISSOURI	40.0	39.0	-1.0
MONTANA	39.5	38.9	-0.6
NEBRASKA	40.0	38.1	-1.9
NEVADA	33.7	35.7	2.1
NEW HAMPSHIRE	41.0	40.5	-0.6
NEW JERSEY	39.5	38.4	-1.2
NEW MEXICO	42.7	40.6	-2.1
NEW YORK	39.4	36.2	-3.1
NORTH CAROLINA	38.8	35.6	-3.2
NORTH DAKOTA	39.5	39.6	0.2
OHIO	37.5	37.0	-0.5
OKLAHOMA	42.7	41.7	-1.0
OREGON	35.6	36.0	0.5
PENNSYLVANIA	36.8	34.6	-2.2
RHODE ISLAND	36.0	33.5	-2.4
SOUTH CAROLINA	38.1	35.4	-2.7
SOUTH DAKOTA	41.0	39.9	-1.1
TENNESSEE	38.2	37.3	-0.9
TEXAS	41.2	40.6	-0.6
UTAH	44.9	45.2	0.3
VERMONT	41.3	42.4	1.1
VIRGINIA	40.4	37.1	-3.3
WASHINGTON	39.1	38.5	-0.6
WEST VIRGINIA	39.6	36.3	-3.4
WISCONSIN	39.8	37.6	-2.2
WYOMING	41.6	39.7	-1.9
PUERTO RICO	45.8	44.7	-1.1

* NEVER HAD A MAMMOGRAM

April 17, 2001

TABLE 29.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM, WOMEN AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	908	11.3	1.2	(8.9, 13.7)
ALASKA	623	13.8	2.1	(9.8, 17.9)
ARIZONA	1006	10.5	1.6	(7.5, 13.6)
ARKANSAS	1219	14.7	1.2	(12.4, 16.9)
COLORADO	1104	12.4	1.5	(9.6, 15.3)
CONNECTICUT	1427	7.3	0.8	(5.7, 8.8)
DELAWARE	994	7.1	1.0	(5.1, 9.1)
DISTRICT OF CO	570	7.0	1.2	(4.6, 9.3)
FLORIDA	2017	11.1	0.8	(9.5, 12.7)
GEORGIA	1483	12.6	1.1	(10.6, 14.7)
HAWAII	2197	10.5	0.8	(8.9, 12.2)
IDAHO	1657	15.9	1.1	(13.8, 18.0)
ILLINOIS	1335	12.6	1.0	(10.5, 14.6)
INDIANA	1052	13.5	1.2	(11.1, 15.8)
IOWA	1455	13.1	1.0	(11.2, 15.0)
KANSAS	1617	11.7	0.9	(10.0, 13.5)
KENTUCKY	2592	12.9	0.9	(11.2, 14.6)
LOUISIANA	1844	14.2	0.9	(12.5, 16.0)
MAINE	1829	9.3	1.1	(7.2, 11.4)
MARYLAND	1654	9.2	0.9	(7.4, 10.9)
MASSACHUSETTS	2819	7.9	0.6	(6.8, 9.0)
MICHIGAN	922	8.8	1.1	(6.6, 10.9)
MINNESOTA	970	12.7	1.1	(10.5, 14.9)
MISSISSIPPI	841	16.4	1.4	(13.7, 19.2)
MISSOURI	1741	14.7	1.1	(12.5, 16.9)
MONTANA	1189	12.6	1.3	(10.2, 15.1)
NEBRASKA	1156	13.6	1.1	(11.4, 15.8)
NEVADA	631	10.9	1.7	(7.6, 14.1)
NEW HAMPSHIRE	712	9.0	1.1	(6.9, 11.2)
NEW JERSEY	1404	14.0	1.1	(11.8, 16.2)
NEW MEXICO	1143	12.7	1.1	(10.5, 14.9)
NEW YORK	1147	9.8	1.1	(7.7, 11.9)
NORTH CAROLINA	1176	10.9	1.0	(8.9, 12.9)
NORTH DAKOTA	675	12.0	1.4	(9.3, 14.7)
OHIO	1194	10.4	1.3	(7.9, 12.8)
OKLAHOMA	1357	17.7	1.2	(15.5, 20.0)
OREGON	1428	9.4	0.9	(7.8, 11.1)
PENNSYLVANIA	1356	10.0	0.9	(8.2, 11.7)
RHODE ISLAND	1336	8.1	0.8	(6.4, 9.7)
SOUTH CAROLINA	1210	10.0	1.0	(8.0, 12.1)
SOUTH DAKOTA	1989	13.0	0.8	(11.5, 14.6)
TENNESSEE	1244	11.9	1.0	(9.9, 13.9)
TEXAS	1695	15.1	1.0	(13.1, 17.0)
UTAH	930	13.6	1.5	(10.7, 16.5)
VERMONT	1346	12.6	1.0	(10.7, 14.6)
VIRGINIA	655	11.4	1.3	(8.8, 14.0)
WASHINGTON	1271	10.7	1.0	(8.8, 12.7)
WEST VIRGINIA	905	13.6	1.2	(11.2, 16.0)
WISCONSIN	951	11.8	1.1	(9.7, 14.0)
WYOMING	892	13.6	1.2	(11.1, 16.0)
PUERTO RICO	1737	17.2	1.2	(14.9, 19.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 12.0 RANGE = 7.0-17.7

* NEVER HAD A MAMMOGRAM

April 17, 2001

TABLE 29.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO MAMMOGRAM, WOMEN AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.0	11.3	-2.7
ALASKA	17.4	13.8	-3.5
ARIZONA	12.0	10.5	-1.5
ARKANSAS	18.9	14.7	-4.2
COLORADO	15.6	12.4	-3.2
CONNECTICUT	8.1	7.3	-0.9
DELAWARE	8.8	7.1	-1.7
DISTRICT OF CO	10.5	7.0	-3.5
FLORIDA	9.5	11.1	1.6
GEORGIA	14.5	12.6	-1.9
HAWAII	11.5	10.5	-0.9
IDAHO	16.0	15.9	-0.1
ILLINOIS	13.4	12.6	-0.8
INDIANA	14.6	13.5	-1.1
IOWA	13.9	13.1	-0.8
KANSAS	14.1	11.7	-2.3
KENTUCKY	14.4	12.9	-1.4
LOUISIANA	17.9	14.2	-3.6
MAINE	9.1	9.3	0.2
MARYLAND	10.7	9.2	-1.5
MASSACHUSETTS	9.8	7.9	-1.9
MICHIGAN	10.5	8.8	-1.8
MINNESOTA	16.0	12.7	-3.3
MISSISSIPPI	19.4	16.4	-3.0
MISSOURI	16.9	14.7	-2.2
MONTANA	14.2	12.6	-1.6
NEBRASKA	15.7	13.6	-2.1
NEVADA	11.1	10.9	-0.2
NEW HAMPSHIRE	11.8	9.0	-2.7
NEW JERSEY	15.4	14.0	-1.4
NEW MEXICO	16.4	12.7	-3.7
NEW YORK	14.2	9.8	-4.4
NORTH CAROLINA	13.3	10.9	-2.4
NORTH DAKOTA	13.4	12.0	-1.4
OHIO	11.6	10.4	-1.2
OKLAHOMA	16.9	17.7	0.8
OREGON	8.5	9.4	0.9
PENNSYLVANIA	11.8	10.0	-1.9
RHODE ISLAND	9.0	8.1	-0.9
SOUTH CAROLINA	12.6	10.0	-2.6
SOUTH DAKOTA	13.7	13.0	-0.7
TENNESSEE	15.7	11.9	-3.8
TEXAS	13.8	15.1	1.3
UTAH	13.9	13.6	-0.3
VERMONT	12.2	12.6	0.5
VIRGINIA	13.7	11.4	-2.3
WASHINGTON	10.0	10.7	0.7
WEST VIRGINIA	17.4	13.6	-3.8
WISCONSIN	11.2	11.8	0.6
WYOMING	16.7	13.6	-3.1
PUERTO RICO	19.2	17.2	-2.0

* NEVER HAD A MAMMOGRAM

April 17, 2001

TABLE 30.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM, WOMEN AGE 50 AND OVER*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	606	23.3	1.9	(19.6, 26.9)
ALASKA	309	14.4	2.6	(9.2, 19.6)
ARIZONA	707	15.2	1.8	(11.6, 18.7)
ARKANSAS	879	25.2	1.6	(22.1, 28.3)
COLORADO	700	19.6	2.2	(15.4, 23.8)
CONNECTICUT	926	15.4	1.3	(12.8, 18.0)
DELAWARE	685	9.7	1.3	(7.2, 12.3)
DISTRICT OF CO	377	14.3	2.1	(10.3, 18.3)
FLORIDA	1407	16.8	1.1	(14.6, 19.0)
GEORGIA	951	19.5	1.5	(16.6, 22.4)
HAWAII	1436	17.9	1.3	(15.3, 20.5)
IDAHO	1060	27.9	1.6	(24.8, 31.1)
ILLINOIS	895	21.5	1.5	(18.6, 24.4)
INDIANA	711	27.5	1.8	(24.0, 31.1)
IOWA	1025	21.4	1.3	(18.7, 24.0)
KANSAS	1092	21.4	1.3	(18.8, 24.0)
KENTUCKY	1814	23.1	1.3	(20.6, 25.5)
LOUISIANA	1214	23.0	1.4	(20.3, 25.6)
MAINE	1287	17.1	1.8	(13.7, 20.5)
MARYLAND	1043	14.6	1.4	(11.9, 17.3)
MASSACHUSETTS	1860	14.0	0.9	(12.2, 15.8)
MICHIGAN	620	15.7	1.6	(12.6, 18.8)
MINNESOTA	646	23.7	1.9	(20.0, 27.4)
MISSISSIPPI	554	29.0	2.1	(24.8, 33.2)
MISSOURI	1179	24.2	1.6	(21.1, 27.2)
MONTANA	790	20.7	1.7	(17.3, 24.0)
NEBRASKA	799	22.0	1.6	(18.8, 25.2)
NEVADA	396	22.3	2.9	(16.6, 27.9)
NEW HAMPSHIRE	428	15.9	1.8	(12.3, 19.5)
NEW JERSEY	898	22.7	1.6	(19.5, 25.8)
NEW MEXICO	765	21.2	1.7	(17.9, 24.6)
NEW YORK	730	18.4	1.8	(14.9, 21.8)
NORTH CAROLINA	817	19.7	1.5	(16.8, 22.7)
NORTH DAKOTA	439	20.5	2.2	(16.3, 24.7)
OHIO	791	18.9	1.9	(15.2, 22.7)
OKLAHOMA	970	25.7	1.5	(22.8, 28.6)
OREGON	964	17.3	1.3	(14.7, 19.8)
PENNSYLVANIA	899	21.3	1.5	(18.3, 24.2)
RHODE ISLAND	875	13.3	1.2	(11.0, 15.7)
SOUTH CAROLINA	789	16.3	1.5	(13.4, 19.3)
SOUTH DAKOTA	1359	21.4	1.2	(19.0, 23.7)
TENNESSEE	847	23.6	1.7	(20.3, 26.8)
TEXAS	1025	26.0	1.5	(22.9, 29.0)
UTAH	595	22.8	2.2	(18.6, 27.1)
VERMONT	869	19.9	1.5	(17.1, 22.8)
VIRGINIA	405	18.8	2.1	(14.7, 22.8)
WASHINGTON	817	20.4	1.5	(17.4, 23.4)
WEST VIRGINIA	644	23.2	1.8	(19.7, 26.7)
WISCONSIN	600	23.1	1.8	(19.5, 26.7)
WYOMING	576	29.1	2.2	(24.9, 33.3)
PUERTO RICO	1281	28.3	1.7	(25.0, 31.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 21.2 RANGE = 9.7-29.1

* DID NOT HAVE A MAMMOGRAM IN PAST TWO YEARS

April 17, 2001

TABLE 30.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO MAMMOGRAM, WOMEN AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	23.8	23.3	-0.5
ALASKA	22.9	14.4	-8.5
ARIZONA	15.5	15.2	-0.3
ARKANSAS	29.8	25.2	-4.6
COLORADO	24.8	19.6	-5.2
CONNECTICUT	13.9	15.4	1.6
DELAWARE	14.0	9.7	-4.3
DISTRICT OF CO	16.7	14.3	-2.4
FLORIDA	16.6	16.8	0.1
GEORGIA	24.8	19.5	-5.3
HAWAII	20.7	17.9	-2.8
IDAHO	31.4	27.9	-3.5
ILLINOIS	23.7	21.5	-2.2
INDIANA	31.3	27.5	-3.7
IOWA	25.5	21.4	-4.2
KANSAS	20.1	21.4	1.4
KENTUCKY	26.5	23.1	-3.5
LOUISIANA	30.4	23.0	-7.4
MAINE	18.2	17.1	-1.1
MARYLAND	16.9	14.6	-2.3
MASSACHUSETTS	16.9	14.0	-2.9
MICHIGAN	16.7	15.7	-1.0
MINNESOTA	26.6	23.7	-3.0
MISSISSIPPI	32.9	29.0	-4.0
MISSOURI	25.7	24.2	-1.5
MONTANA	28.8	20.7	-8.1
NEBRASKA	25.8	22.0	-3.8
NEVADA	25.0	22.3	-2.7
NEW HAMPSHIRE	15.6	15.9	0.3
NEW JERSEY	22.2	22.7	0.4
NEW MEXICO	27.2	21.2	-6.0
NEW YORK	21.5	18.4	-3.1
NORTH CAROLINA	19.8	19.7	-0.1
NORTH DAKOTA	23.3	20.5	-2.8
OHIO	23.2	18.9	-4.3
OKLAHOMA	28.2	25.7	-2.5
OREGON	22.6	17.3	-5.4
PENNSYLVANIA	22.1	21.3	-0.8
RHODE ISLAND	14.9	13.3	-1.5
SOUTH CAROLINA	18.7	16.3	-2.3
SOUTH DAKOTA	25.1	21.4	-3.7
TENNESSEE	24.9	23.6	-1.3
TEXAS	25.1	26.0	0.9
UTAH	27.7	22.8	-4.9
VERMONT	22.3	19.9	-2.4
VIRGINIA	26.0	18.8	-7.2
WASHINGTON	24.5	20.4	-4.0
WEST VIRGINIA	28.0	23.2	-4.8
WISCONSIN	25.2	23.1	-2.1
WYOMING	29.2	29.1	-0.1
PUERTO RICO	29.5	28.3	-1.3

* DID NOT HAVE A MAMMOGRAM IN PAST TWO YEARS

April 17, 2001

TABLE 31.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CLINICAL BREAST EXAM, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1399	10.1	1.1	(7.9, 12.4)
ALASKA	1090	7.0	1.1	(4.8, 9.1)
ARIZONA	1586	19.8	2.4	(15.1, 24.5)
ARKANSAS	1831	9.8	0.8	(8.2, 11.4)
COLORADO	1754	11.3	1.2	(8.9, 13.8)
CONNECTICUT	2371	9.8	0.8	(8.3, 11.3)
DELAWARE	1611	10.2	1.0	(8.2, 12.2)
DISTRICT OF CO	1020	11.7	1.2	(9.4, 13.9)
FLORIDA	3042	14.8	0.8	(13.4, 16.3)
GEORGIA	2501	10.5	0.8	(9.0, 12.0)
HAWAII	3329	15.3	0.9	(13.6, 17.1)
IDAHO	2846	9.5	0.7	(8.2, 10.8)
ILLINOIS	2212	14.5	0.9	(12.7, 16.2)
INDIANA	1707	11.3	1.0	(9.4, 13.1)
IOWA	2150	8.9	0.7	(7.4, 10.3)
KANSAS	2565	12.3	0.7	(10.9, 13.8)
KENTUCKY	3817	12.2	0.7	(10.7, 13.6)
LOUISIANA	3096	14.7	0.7	(13.3, 16.2)
MAINE	2696	7.5	0.8	(6.0, 9.0)
MARYLAND	2795	9.5	0.7	(8.1, 11.0)
MASSACHUSETTS	4880	8.7	0.5	(7.8, 9.7)
MICHIGAN	1472	10.2	1.0	(8.3, 12.2)
MINNESOTA	1592	7.6	0.8	(6.1, 9.1)
MISSISSIPPI	1346	12.1	1.0	(10.1, 14.1)
MISSOURI	2675	12.5	0.9	(10.9, 14.2)
MONTANA	1725	6.8	1.1	(4.7, 8.9)
NEBRASKA	1781	11.3	1.0	(9.4, 13.2)
NEVADA	1049	11.8	1.5	(8.8, 14.8)
NEW HAMPSHIRE	1127	5.6	0.8	(3.9, 7.2)
NEW JERSEY	2288	15.0	0.9	(13.2, 16.8)
NEW MEXICO	1769	11.9	0.9	(10.1, 13.8)
NEW YORK	2004	13.1	1.0	(11.2, 15.0)
NORTH CAROLINA	1867	8.5	0.8	(7.0, 10.0)
NORTH DAKOTA	1093	9.8	1.0	(7.8, 11.8)
OHIO	1971	11.6	1.1	(9.5, 13.7)
OKLAHOMA	2110	10.1	0.8	(8.6, 11.7)
OREGON	2236	8.8	0.8	(7.3, 10.3)
PENNSYLVANIA	2105	9.2	0.7	(7.7, 10.6)
RHODE ISLAND	2155	9.2	0.7	(7.8, 10.7)
SOUTH CAROLINA	1984	11.2	0.8	(9.6, 12.8)
SOUTH DAKOTA	3051	8.6	0.6	(7.4, 9.7)
TENNESSEE	2025	13.6	0.9	(11.9, 15.3)
TEXAS	2991	16.3	0.8	(14.6, 17.9)
UTAH	1618	9.4	1.0	(7.6, 11.3)
VERMONT	2106	10.3	0.9	(8.6, 12.1)
VIRGINIA	1121	12.2	1.3	(9.6, 14.7)
WASHINGTON	2077	6.7	0.6	(5.4, 7.9)
WEST VIRGINIA	1395	10.1	1.0	(8.2, 12.1)
WISCONSIN	1493	9.4	0.9	(7.7, 11.1)
WYOMING	1377	8.3	0.8	(6.6, 9.9)
PUERTO RICO	2637	21.6	1.2	(19.2, 23.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 10.2 RANGE = 5.6-21.6

* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

April 17, 2001

TABLE 31.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO CLINICAL BREAST EXAM, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	12.3	10.1	-2.2
ALASKA	9.6	7.0	-2.6
ARIZONA	18.7	19.8	1.1
ARKANSAS	10.5	9.8	-0.7
COLORADO	9.0	11.3	2.4
CONNECTICUT	13.0	9.8	-3.2
DELAWARE	9.1	10.2	1.1
DISTRICT OF CO	8.8	11.7	2.9
FLORIDA	12.4	14.8	2.5
GEORGIA	10.4	10.5	0.1
HAWAII	13.4	15.3	2.0
IDAHO	9.2	9.5	0.3
ILLINOIS	13.5	14.5	1.0
INDIANA	10.6	11.3	0.6
IOWA	8.6	8.9	0.3
KANSAS	14.8	12.3	-2.4
KENTUCKY	11.4	12.2	0.8
LOUISIANA	15.1	14.7	-0.4
MAINE	5.0	7.5	2.5
MARYLAND	10.5	9.5	-1.0
MASSACHUSETTS	12.3	8.7	-3.6
MICHIGAN	9.0	10.2	1.2
MINNESOTA	6.3	7.6	1.2
MISSISSIPPI	12.9	12.1	-0.8
MISSOURI	13.6	12.5	-1.1
MONTANA	6.2	6.8	0.6
NEBRASKA	10.0	11.3	1.3
NEVADA	11.1	11.8	0.7
NEW HAMPSHIRE	6.2	5.6	-0.7
NEW JERSEY	18.2	15.0	-3.2
NEW MEXICO	11.4	11.9	0.6
NEW YORK	16.3	13.1	-3.2
NORTH CAROLINA	7.4	8.5	1.1
NORTH DAKOTA	7.8	9.8	1.9
OHIO	8.9	11.6	2.7
OKLAHOMA	14.3	10.1	-4.2
OREGON	9.3	8.8	-0.4
PENNSYLVANIA	12.8	9.2	-3.6
RHODE ISLAND	14.8	9.2	-5.6
SOUTH CAROLINA	13.9	11.2	-2.7
SOUTH DAKOTA	7.5	8.6	1.1
TENNESSEE	14.0	13.6	-0.4
TEXAS	15.8	16.3	0.4
UTAH	8.5	9.4	1.0
VERMONT	10.3	10.3	0.0
VIRGINIA	11.1	12.2	1.1
WASHINGTON	8.7	6.7	-2.0
WEST VIRGINIA	11.5	10.1	-1.4
WISCONSIN	8.3	9.4	1.1
WYOMING	10.8	8.3	-2.5
PUERTO RICO	23.7	21.6	-2.2

* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

April 17, 2001

TABLE 32.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CLINICAL BREAST EXAM, WOMEN AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	903	9.8	1.1	(7.6, 12.0)
ALASKA	621	4.5	1.1	(2.4, 6.7)
ARIZONA	1004	17.6	2.1	(13.6, 21.7)
ARKANSAS	1211	9.7	0.9	(7.9, 11.5)
COLORADO	1100	8.6	1.3	(6.0, 11.2)
CONNECTICUT	1420	8.4	0.9	(6.6, 10.2)
DELAWARE	987	8.2	1.1	(6.0, 10.3)
DISTRICT OF CO	563	13.6	1.7	(10.3, 16.9)
FLORIDA	2003	13.2	0.9	(11.5, 14.9)
GEORGIA	1478	10.6	1.0	(8.7, 12.5)
HAWAII	2193	12.4	1.0	(10.4, 14.3)
IDAHO	1653	8.7	0.8	(7.1, 10.2)
ILLINOIS	1332	11.7	1.0	(9.7, 13.6)
INDIANA	1048	11.1	1.1	(8.9, 13.3)
IOWA	1453	8.2	0.8	(6.7, 9.7)
KANSAS	1610	11.8	0.9	(10.1, 13.5)
KENTUCKY	2581	11.4	0.8	(9.8, 13.1)
LOUISIANA	1829	13.4	0.9	(11.7, 15.1)
MAINE	1824	6.8	0.9	(5.1, 8.6)
MARYLAND	1650	8.7	0.9	(6.9, 10.4)
MASSACHUSETTS	2805	7.1	0.6	(6.0, 8.2)
MICHIGAN	919	9.5	1.1	(7.3, 11.7)
MINNESOTA	968	6.9	0.9	(5.1, 8.7)
MISSISSIPPI	840	11.2	1.2	(8.8, 13.5)
MISSOURI	1732	12.0	1.1	(9.9, 14.1)
MONTANA	1187	4.8	0.8	(3.3, 6.3)
NEBRASKA	1150	9.0	1.0	(7.1, 10.9)
NEVADA	630	11.7	2.0	(7.7, 15.7)
NEW HAMPSHIRE	708	4.3	0.8	(2.8, 5.9)
NEW JERSEY	1398	13.1	1.1	(10.9, 15.2)
NEW MEXICO	1142	9.3	1.0	(7.4, 11.3)
NEW YORK	1143	11.3	1.2	(8.9, 13.6)
NORTH CAROLINA	1173	7.9	0.9	(6.2, 9.6)
NORTH DAKOTA	669	7.6	1.1	(5.4, 9.8)
OHIO	1193	9.4	1.2	(7.0, 11.8)
OKLAHOMA	1354	8.5	0.9	(6.8, 10.2)
OREGON	1421	7.2	0.8	(5.6, 8.7)
PENNSYLVANIA	1355	9.2	0.9	(7.5, 11.0)
RHODE ISLAND	1329	8.5	0.9	(6.8, 10.1)
SOUTH CAROLINA	1210	10.8	1.0	(8.8, 12.8)
SOUTH DAKOTA	1981	7.6	0.6	(6.3, 8.8)
TENNESSEE	1238	13.5	1.1	(11.3, 15.7)
TEXAS	1688	13.9	1.0	(11.9, 15.9)
UTAH	929	5.8	1.0	(3.9, 7.8)
VERMONT	1344	8.8	0.9	(7.2, 10.5)
VIRGINIA	651	10.8	1.4	(8.1, 13.4)
WASHINGTON	1262	5.0	0.7	(3.6, 6.4)
WEST VIRGINIA	904	9.4	1.1	(7.4, 11.5)
WISCONSIN	950	8.7	1.0	(6.8, 10.6)
WYOMING	889	7.5	1.0	(5.5, 9.6)
PUERTO RICO	1728	13.2	1.0	(11.2, 15.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 9.3 RANGE = 4.3-17.6

* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

April 17, 2001

TABLE 32.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO CLINICAL BREAST EXAM, WOMEN AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	13.1	9.8	-3.3
ALASKA	6.9	4.5	-2.3
ARIZONA	11.9	17.6	5.7
ARKANSAS	11.0	9.7	-1.2
COLORADO	9.3	8.6	-0.7
CONNECTICUT	12.0	8.4	-3.6
DELAWARE	8.2	8.2	0.0
DISTRICT OF CO	7.5	13.6	6.1
FLORIDA	10.8	13.2	2.5
GEORGIA	11.0	10.6	-0.4
HAWAII	11.1	12.4	1.2
IDAHO	7.2	8.7	1.5
ILLINOIS	10.6	11.7	1.1
INDIANA	9.7	11.1	1.4
IOWA	8.3	8.2	-0.1
KANSAS	11.8	11.8	0.0
KENTUCKY	10.8	11.4	0.7
LOUISIANA	13.1	13.4	0.3
MAINE	4.4	6.8	2.4
MARYLAND	8.9	8.7	-0.3
MASSACHUSETTS	9.2	7.1	-2.1
MICHIGAN	5.3	9.5	4.2
MINNESOTA	5.5	6.9	1.4
MISSISSIPPI	13.6	11.2	-2.4
MISSOURI	13.0	12.0	-1.0
MONTANA	4.9	4.8	-0.0
NEBRASKA	8.8	9.0	0.2
NEVADA	9.3	11.7	2.4
NEW HAMPSHIRE	5.8	4.3	-1.5
NEW JERSEY	15.5	13.1	-2.5
NEW MEXICO	10.4	9.3	-1.0
NEW YORK	12.7	11.3	-1.4
NORTH CAROLINA	5.8	7.9	2.1
NORTH DAKOTA	6.4	7.6	1.2
OHIO	7.8	9.4	1.6
OKLAHOMA	10.9	8.5	-2.3
OREGON	8.6	7.2	-1.4
PENNSYLVANIA	12.5	9.2	-3.3
RHODE ISLAND	13.0	8.5	-4.6
SOUTH CAROLINA	12.6	10.8	-1.8
SOUTH DAKOTA	7.3	7.6	0.2
TENNESSEE	12.2	13.5	1.3
TEXAS	12.6	13.9	1.3
UTAH	4.6	5.8	1.3
VERMONT	8.5	8.8	0.3
VIRGINIA	9.5	10.8	1.3
WASHINGTON	7.7	5.0	-2.7
WEST VIRGINIA	11.0	9.4	-1.6
WISCONSIN	6.6	8.7	2.1
WYOMING	9.1	7.5	-1.5
PUERTO RICO	13.8	13.2	-0.6

* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

April 17, 2001

TABLE 33.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CLINICAL BREAST EXAM, WOMEN AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	600	25.7	1.9	(21.9, 29.5)
ALASKA	302	14.1	2.7	(8.8, 19.3)
ARIZONA	707	26.2	2.5	(21.3, 31.0)
ARKANSAS	857	26.9	1.7	(23.7, 30.2)
COLORADO	692	23.4	2.3	(18.9, 28.0)
CONNECTICUT	923	15.9	1.4	(13.0, 18.7)
DELAWARE	679	16.6	1.8	(13.1, 20.2)
DISTRICT OF CO	368	22.2	2.5	(17.4, 27.1)
FLORIDA	1386	23.8	1.3	(21.3, 26.4)
GEORGIA	933	20.6	1.6	(17.5, 23.7)
HAWAII	1430	21.2	1.5	(18.3, 24.1)
IDAHO	1056	27.6	1.6	(24.5, 30.7)
ILLINOIS	887	23.4	1.6	(20.3, 26.5)
INDIANA	708	28.0	1.9	(24.3, 31.7)
IOWA	1015	21.0	1.4	(18.4, 23.7)
KANSAS	1071	23.5	1.4	(20.8, 26.2)
KENTUCKY	1812	24.2	1.3	(21.6, 26.7)
LOUISIANA	1203	24.3	1.4	(21.6, 27.1)
MAINE	1279	16.1	1.7	(12.9, 19.4)
MARYLAND	1032	17.6	1.5	(14.6, 20.6)
MASSACHUSETTS	1848	15.5	1.0	(13.6, 17.4)
MICHIGAN	607	22.2	1.9	(18.4, 26.0)
MINNESOTA	644	23.4	1.9	(19.7, 27.1)
MISSISSIPPI	547	28.9	2.1	(24.7, 33.1)
MISSOURI	1167	25.8	1.6	(22.6, 29.1)
MONTANA	788	16.6	1.6	(13.4, 19.7)
NEBRASKA	793	22.3	1.7	(19.1, 25.6)
NEVADA	392	32.2	3.5	(25.3, 39.1)
NEW HAMPSHIRE	423	11.8	1.6	(8.6, 15.0)
NEW JERSEY	886	25.8	1.7	(22.4, 29.1)
NEW MEXICO	761	22.3	1.7	(18.9, 25.6)
NEW YORK	723	19.6	1.8	(16.1, 23.1)
NORTH CAROLINA	812	19.1	1.5	(16.1, 22.1)
NORTH DAKOTA	428	20.8	2.2	(16.5, 25.1)
OHIO	790	20.8	2.0	(16.9, 24.7)
OKLAHOMA	965	21.9	1.4	(19.0, 24.7)
OREGON	956	21.1	1.5	(18.2, 23.9)
PENNSYLVANIA	898	21.5	1.5	(18.6, 24.5)
RHODE ISLAND	866	15.1	1.3	(12.5, 17.6)
SOUTH CAROLINA	787	19.1	1.6	(15.9, 22.2)
SOUTH DAKOTA	1337	20.7	1.2	(18.3, 23.0)
TENNESSEE	841	25.1	1.7	(21.8, 28.4)
TEXAS	1017	27.6	1.6	(24.6, 30.7)
UTAH	589	23.5	2.2	(19.1, 27.9)
VERMONT	859	22.0	1.6	(19.0, 25.1)
VIRGINIA	402	22.7	2.3	(18.3, 27.1)
WASHINGTON	810	21.1	1.6	(18.0, 24.3)
WEST VIRGINIA	635	23.4	1.8	(19.8, 27.0)
WISCONSIN	600	22.6	2.1	(18.5, 26.7)
WYOMING	570	28.4	2.1	(24.3, 32.4)
PUERTO RICO	1270	24.2	1.5	(21.2, 27.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 22.3 RANGE = 11.8-32.2

* DID NOT HAVE EXAM IN PAST TWO YEARS

April 17, 2001

TABLE 33.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO CLINICAL BREAST EXAM, WOMEN AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	26.8	25.7	-1.1
ALASKA	20.1	14.1	-6.0
ARIZONA	18.5	26.2	7.6
ARKANSAS	29.9	26.9	-3.0
COLORADO	25.2	23.4	-1.7
CONNECTICUT	16.8	15.9	-0.9
DELAWARE	17.8	16.6	-1.1
DISTRICT OF CO	15.9	22.2	6.3
FLORIDA	19.7	23.8	4.1
GEORGIA	26.1	20.6	-5.5
HAWAII	23.7	21.2	-2.5
IDAHO	26.5	27.6	1.1
ILLINOIS	22.5	23.4	0.8
INDIANA	31.1	28.0	-3.0
IOWA	23.8	21.0	-2.7
KANSAS	22.9	23.5	0.6
KENTUCKY	24.4	24.2	-0.2
LOUISIANA	27.3	24.3	-3.0
MAINE	11.5	16.1	4.6
MARYLAND	19.9	17.6	-2.2
MASSACHUSETTS	16.9	15.5	-1.4
MICHIGAN	17.5	22.2	4.7
MINNESOTA	21.9	23.4	1.5
MISSISSIPPI	29.3	28.9	-0.4
MISSOURI	26.5	25.8	-0.7
MONTANA	23.2	16.6	-6.7
NEBRASKA	21.2	22.3	1.1
NEVADA	28.9	32.2	3.3
NEW HAMPSHIRE	14.4	11.8	-2.6
NEW JERSEY	23.2	25.8	2.6
NEW MEXICO	26.2	22.3	-4.0
NEW YORK	21.2	19.6	-1.5
NORTH CAROLINA	16.7	19.1	2.4
NORTH DAKOTA	20.8	20.8	-0.0
OHIO	21.2	20.8	-0.4
OKLAHOMA	25.3	21.9	-3.4
OREGON	23.8	21.1	-2.8
PENNSYLVANIA	23.9	21.5	-2.3
RHODE ISLAND	20.2	15.1	-5.1
SOUTH CAROLINA	21.7	19.1	-2.7
SOUTH DAKOTA	22.5	20.7	-1.9
TENNESSEE	23.7	25.1	1.4
TEXAS	26.6	27.6	1.0
UTAH	21.7	23.5	1.8
VERMONT	22.5	22.0	-0.4
VIRGINIA	23.7	22.7	-1.0
WASHINGTON	24.1	21.1	-2.9
WEST VIRGINIA	27.9	23.4	-4.5
WISCONSIN	22.0	22.6	0.6
WYOMING	26.5	28.4	1.9
PUERTO RICO	29.6	24.2	-5.5

* DID NOT HAVE EXAM IN PAST TWO YEARS

April 17, 2001

TABLE 34.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	903	18.6	1.5	(15.6, 21.5)
ALASKA	620	15.5	2.2	(11.3, 19.7)
ARIZONA	1002	24.4	2.3	(19.9, 29.0)
ARKANSAS	1208	20.5	1.3	(17.9, 23.0)
COLORADO	1100	18.8	1.8	(15.4, 22.2)
CONNECTICUT	1419	13.4	1.1	(11.2, 15.5)
DELAWARE	986	13.3	1.4	(10.7, 16.0)
DISTRICT OF CO	562	18.2	1.9	(14.5, 21.9)
FLORIDA	2002	20.1	1.0	(18.1, 22.1)
GEORGIA	1475	19.7	1.3	(17.3, 22.2)
HAWAII	2193	18.6	1.1	(16.4, 20.8)
IDAHO	1653	20.9	1.2	(18.6, 23.2)
ILLINOIS	1331	19.8	1.2	(17.3, 22.2)
INDIANA	1047	20.4	1.4	(17.6, 23.2)
IOWA	1450	17.8	1.1	(15.7, 20.0)
KANSAS	1608	19.0	1.1	(16.9, 21.2)
KENTUCKY	2576	18.7	1.0	(16.6, 20.7)
LOUISIANA	1828	21.8	1.1	(19.7, 23.9)
MAINE	1824	13.4	1.3	(10.9, 15.8)
MARYLAND	1650	15.0	1.1	(12.8, 17.1)
MASSACHUSETTS	2803	12.5	0.7	(11.1, 13.9)
MICHIGAN	918	15.4	1.4	(12.7, 18.2)
MINNESOTA	966	16.6	1.3	(14.1, 19.1)
MISSISSIPPI	837	22.3	1.6	(19.2, 25.4)
MISSOURI	1729	21.8	1.3	(19.2, 24.5)
MONTANA	1186	15.7	1.4	(13.0, 18.4)
NEBRASKA	1149	17.8	1.3	(15.4, 20.3)
NEVADA	629	19.7	2.4	(15.0, 24.3)
NEW HAMPSHIRE	708	11.7	1.3	(9.2, 14.1)
NEW JERSEY	1398	22.1	1.3	(19.5, 24.7)
NEW MEXICO	1142	18.4	1.3	(15.8, 20.9)
NEW YORK	1143	17.3	1.4	(14.6, 19.9)
NORTH CAROLINA	1173	15.5	1.2	(13.2, 17.7)
NORTH DAKOTA	669	15.7	1.6	(12.6, 18.7)
OHIO	1192	16.2	1.5	(13.3, 19.2)
OKLAHOMA	1352	22.2	1.3	(19.7, 24.6)
OREGON	1420	13.7	1.0	(11.7, 15.7)
PENNSYLVANIA	1354	16.2	1.1	(14.0, 18.4)
RHODE ISLAND	1329	13.7	1.1	(11.7, 15.8)
SOUTH CAROLINA	1209	17.5	1.3	(15.1, 20.0)
SOUTH DAKOTA	1981	16.8	0.9	(15.0, 18.6)
TENNESSEE	1238	21.7	1.3	(19.1, 24.3)
TEXAS	1688	23.0	1.2	(20.6, 25.3)
UTAH	928	16.5	1.6	(13.3, 19.7)
VERMONT	1344	18.2	1.2	(15.9, 20.5)
VIRGINIA	649	19.1	1.7	(15.8, 22.5)
WASHINGTON	1261	14.0	1.1	(11.9, 16.2)
WEST VIRGINIA	903	17.7	1.4	(15.0, 20.4)
WISCONSIN	949	15.6	1.3	(13.2, 18.1)
WYOMING	889	18.1	1.4	(15.3, 20.8)
PUERTO RICO	1728	23.7	1.3	(21.1, 26.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 18.1 RANGE = 11.7-24.4

* NEVER HAD A MAMMOGRAM AND BREAST EXAM

April 17, 2001

TABLE 34.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	22.9	18.6	-4.3
ALASKA	20.0	15.5	-4.5
ARIZONA	17.9	24.4	6.5
ARKANSAS	23.5	20.5	-3.0
COLORADO	20.4	18.8	-1.7
CONNECTICUT	16.7	13.4	-3.3
DELAWARE	15.0	13.3	-1.7
DISTRICT OF CO	15.6	18.2	2.6
FLORIDA	16.6	20.1	3.6
GEORGIA	21.4	19.7	-1.7
HAWAII	18.8	18.6	-0.2
IDAHO	20.4	20.9	0.5
ILLINOIS	20.8	19.8	-1.0
INDIANA	19.8	20.4	0.6
IOWA	18.3	17.8	-0.4
KANSAS	21.0	19.0	-2.0
KENTUCKY	19.0	18.7	-0.3
LOUISIANA	24.0	21.8	-2.2
MAINE	11.3	13.4	2.0
MARYLAND	15.7	15.0	-0.7
MASSACHUSETTS	15.3	12.5	-2.8
MICHIGAN	14.4	15.4	1.1
MINNESOTA	18.3	16.6	-1.7
MISSISSIPPI	26.7	22.3	-4.4
MISSOURI	23.2	21.8	-1.4
MONTANA	16.9	15.7	-1.2
NEBRASKA	19.1	17.8	-1.3
NEVADA	17.5	19.7	2.1
NEW HAMPSHIRE	14.8	11.7	-3.1
NEW JERSEY	24.4	22.1	-2.3
NEW MEXICO	21.3	18.4	-2.9
NEW YORK	21.1	17.3	-3.8
NORTH CAROLINA	15.6	15.5	-0.1
NORTH DAKOTA	16.4	15.7	-0.7
OHIO	15.2	16.2	1.0
OKLAHOMA	23.0	22.2	-0.8
OREGON	14.7	13.7	-1.0
PENNSYLVANIA	19.7	16.2	-3.5
RHODE ISLAND	18.4	13.7	-4.7
SOUTH CAROLINA	21.3	17.5	-3.7
SOUTH DAKOTA	17.8	16.8	-1.0
TENNESSEE	22.4	21.7	-0.6
TEXAS	21.6	23.0	1.4
UTAH	16.4	16.5	0.1
VERMONT	17.1	18.2	1.1
VIRGINIA	18.3	19.1	0.9
WASHINGTON	15.8	14.0	-1.8
WEST VIRGINIA	21.6	17.7	-3.9
WISCONSIN	15.3	15.6	0.3
WYOMING	21.5	18.1	-3.4
PUERTO RICO	26.0	23.7	-2.2

* NEVER HAD A MAMMOGRAM AND BREAST EXAM

April 17, 2001

TABLE 35.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	599	32.7	2.1	(28.6, 36.7)
ALASKA	307	19.0	3.0	(13.1, 24.9)
ARIZONA	701	29.6	2.6	(24.6, 34.6)
ARKANSAS	861	33.6	1.8	(30.2, 37.1)
COLORADO	691	29.7	2.6	(24.6, 34.7)
CONNECTICUT	920	21.4	1.6	(18.3, 24.5)
DELAWARE	678	18.4	1.8	(14.8, 22.0)
DISTRICT OF CO	366	26.0	2.7	(20.8, 31.2)
FLORIDA	1387	28.4	1.4	(25.7, 31.1)
GEORGIA	930	28.2	1.7	(24.8, 31.6)
HAWAII	1429	26.4	1.6	(23.2, 29.5)
IDAHO	1054	35.8	1.7	(32.4, 39.2)
ILLINOIS	887	30.0	1.7	(26.7, 33.4)
INDIANA	708	35.5	2.0	(31.6, 39.4)
IOWA	1014	28.9	1.5	(25.9, 31.9)
KANSAS	1074	30.5	1.5	(27.5, 33.5)
KENTUCKY	1807	30.2	1.4	(27.5, 33.0)
LOUISIANA	1203	32.2	1.5	(29.2, 35.2)
MAINE	1278	22.3	1.9	(18.6, 26.1)
MARYLAND	1033	22.3	1.6	(19.1, 25.5)
MASSACHUSETTS	1851	21.1	1.1	(19.0, 23.3)
MICHIGAN	606	27.1	2.1	(23.1, 31.2)
MINNESOTA	643	30.3	2.1	(26.3, 34.3)
MISSISSIPPI	545	37.9	2.3	(33.4, 42.4)
MISSOURI	1164	32.8	1.8	(29.3, 36.2)
MONTANA	785	25.6	1.9	(21.9, 29.3)
NEBRASKA	794	28.5	1.8	(25.0, 32.0)
NEVADA	394	37.2	3.6	(30.1, 44.3)
NEW HAMPSHIRE	422	18.2	2.0	(14.4, 22.1)
NEW JERSEY	888	33.1	1.8	(29.6, 36.7)
NEW MEXICO	758	29.0	1.9	(25.3, 32.7)
NEW YORK	723	27.7	2.0	(23.7, 31.6)
NORTH CAROLINA	812	24.9	1.7	(21.6, 28.2)
NORTH DAKOTA	431	26.5	2.4	(21.8, 31.1)
OHIO	786	26.8	2.2	(22.6, 31.1)
OKLAHOMA	965	33.0	1.7	(29.7, 36.2)
OREGON	951	26.1	1.6	(23.0, 29.2)
PENNSYLVANIA	895	29.0	1.7	(25.7, 32.2)
RHODE ISLAND	863	20.3	1.5	(17.4, 23.2)
SOUTH CAROLINA	786	24.3	1.7	(20.9, 27.7)
SOUTH DAKOTA	1342	27.8	1.4	(25.1, 30.4)
TENNESSEE	845	33.9	1.8	(30.3, 37.5)
TEXAS	1017	35.5	1.7	(32.2, 38.8)
UTAH	589	30.5	2.4	(25.7, 35.2)
VERMONT	860	28.2	1.7	(24.9, 31.5)
VIRGINIA	402	28.5	2.4	(23.8, 33.2)
WASHINGTON	808	27.9	1.8	(24.4, 31.3)
WEST VIRGINIA	636	29.6	2.0	(25.7, 33.4)
WISCONSIN	598	30.1	2.3	(25.7, 34.6)
WYOMING	570	35.9	2.3	(31.4, 40.3)
PUERTO RICO	1267	34.0	1.8	(30.6, 37.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 28.9 RANGE = 18.2-37.9 YEAR 2000 OBJ 16.11
 * DID NOT HAVE MAMMOGRAM AND BREAST EXAM IN PAST TWO YEARS April 17, 2001

TABLE 35.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 88
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	34.7	32.7	-2.0
ALASKA	26.9	19.0	-7.8
ARIZONA	23.1	29.6	6.5
ARKANSAS	38.9	33.6	-5.3
COLORADO	33.9	29.7	-4.3
CONNECTICUT	22.2	21.4	-0.9
DELAWARE	22.8	18.4	-4.4
DISTRICT OF CO	21.8	26.0	4.3
FLORIDA	23.9	28.4	4.6
GEORGIA	33.8	28.2	-5.6
HAWAII	30.5	26.4	-4.1
IDAHO	36.8	35.8	-1.0
ILLINOIS	32.1	30.0	-2.1
INDIANA	39.3	35.5	-3.8
IOWA	31.3	28.9	-2.4
KANSAS	28.5	30.5	2.0
KENTUCKY	31.7	30.2	-1.5
LOUISIANA	35.0	32.2	-2.8
MAINE	19.9	22.3	2.5
MARYLAND	24.4	22.3	-2.0
MASSACHUSETTS	22.3	21.1	-1.2
MICHIGAN	23.5	27.1	3.7
MINNESOTA	31.0	30.3	-0.7
MISSISSIPPI	40.7	37.9	-2.7
MISSOURI	35.1	32.8	-2.4
MONTANA	33.6	25.6	-8.0
NEBRASKA	31.1	28.5	-2.6
NEVADA	35.3	37.2	1.9
NEW HAMPSHIRE	19.6	18.2	-1.3
NEW JERSEY	32.0	33.1	1.2
NEW MEXICO	33.7	29.0	-4.7
NEW YORK	29.0	27.7	-1.3
NORTH CAROLINA	23.1	24.9	1.8
NORTH DAKOTA	28.7	26.5	-2.3
OHIO	29.0	26.8	-2.2
OKLAHOMA	35.3	33.0	-2.3
OREGON	30.7	26.1	-4.6
PENNSYLVANIA	32.1	29.0	-3.1
RHODE ISLAND	25.2	20.3	-4.9
SOUTH CAROLINA	27.4	24.3	-3.1
SOUTH DAKOTA	31.8	27.8	-4.1
TENNESSEE	32.1	33.9	1.8
TEXAS	34.7	35.5	0.8
UTAH	32.3	30.5	-1.9
VERMONT	30.0	28.2	-1.8
VIRGINIA	32.0	28.5	-3.5
WASHINGTON	32.6	27.9	-4.7
WEST VIRGINIA	34.3	29.6	-4.8
WISCONSIN	31.2	30.1	-1.1
WYOMING	35.9	35.9	-0.1
PUERTO RICO	38.8	34.0	-4.8

* DID NOT HAVE MAMMOGRAM AND BREAST EXAM IN PAST TWO YEARS

April 17, 2001

TABLE 36.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PAP SMEAR, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1392	3.7	0.6	(2.6, 4.8)
ALASKA	1082	2.5	0.7	(1.0, 3.9)
ARIZONA	1570	6.5	0.9	(4.8, 8.2)
ARKANSAS	1820	4.0	0.6	(2.8, 5.2)
COLORADO	1740	4.0	0.8	(2.5, 5.6)
CONNECTICUT	2327	6.1	0.6	(4.9, 7.3)
DELAWARE	1605	3.2	0.6	(2.0, 4.5)
DISTRICT OF CO	1019	5.3	0.8	(3.7, 6.9)
FLORIDA	3041	6.1	0.5	(5.1, 7.1)
GEORGIA	2490	3.4	0.5	(2.3, 4.4)
HAWAII	3299	6.1	0.6	(4.9, 7.3)
IDAHO	2833	5.0	0.5	(4.0, 6.0)
ILLINOIS	2208	6.9	0.7	(5.5, 8.2)
INDIANA	1689	5.9	0.8	(4.4, 7.4)
IOWA	2138	4.2	0.6	(3.1, 5.3)
KANSAS	2547	3.8	0.4	(2.9, 4.6)
KENTUCKY	3812	7.0	0.6	(5.9, 8.1)
LOUISIANA	3069	6.0	0.5	(5.0, 7.0)
MAINE	2678	5.4	0.8	(3.8, 7.0)
MARYLAND	2754	5.1	0.5	(4.0, 6.1)
MASSACHUSETTS	4821	6.3	0.4	(5.4, 7.2)
MICHIGAN	1468	5.6	0.9	(3.8, 7.3)
MINNESOTA	1592	4.0	0.6	(2.9, 5.2)
MISSISSIPPI	1330	5.0	0.7	(3.6, 6.4)
MISSOURI	2672	4.3	0.5	(3.3, 5.3)
MONTANA	1723	2.9	0.9	(1.1, 4.7)
NEBRASKA	1764	7.1	0.8	(5.5, 8.6)
NEVADA	1048	5.1	1.1	(2.9, 7.2)
NEW HAMPSHIRE	1118	3.3	0.7	(1.8, 4.7)
NEW JERSEY	2253	9.7	0.8	(8.1, 11.2)
NEW MEXICO	1762	5.4	0.7	(4.1, 6.8)
NEW YORK	1965	7.1	0.8	(5.6, 8.6)
NORTH CAROLINA	1853	3.6	0.5	(2.6, 4.7)
NORTH DAKOTA	1077	5.8	0.8	(4.2, 7.4)
OHIO	1950	4.3	0.7	(3.0, 5.6)
OKLAHOMA	2098	3.7	0.5	(2.7, 4.7)
OREGON	2226	3.5	0.5	(2.6, 4.4)
PENNSYLVANIA	2084	5.8	0.6	(4.6, 7.1)
RHODE ISLAND	2129	6.1	0.7	(4.8, 7.4)
SOUTH CAROLINA	1965	4.1	0.6	(3.0, 5.2)
SOUTH DAKOTA	3039	3.6	0.4	(2.8, 4.4)
TENNESSEE	2017	6.5	0.7	(5.2, 7.7)
TEXAS	2961	7.2	0.6	(5.9, 8.4)
UTAH	1611	5.7	0.9	(4.0, 7.5)
VERMONT	2088	6.3	0.8	(4.8, 7.8)
VIRGINIA	1112	5.9	1.3	(3.3, 8.4)
WASHINGTON	2081	3.5	0.5	(2.5, 4.4)
WEST VIRGINIA	1387	4.7	0.8	(3.3, 6.2)
WISCONSIN	1482	3.4	0.6	(2.3, 4.5)
WYOMING	1366	3.6	0.6	(2.4, 4.9)
PUERTO RICO	2636	15.3	1.1	(13.0, 17.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 5.1 RANGE = 2.5-15.3 YEAR 2000 OBJ 16.11
 * NEVER HAD PAP SMEAR April 17, 2001

TABLE 36.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO PAP SMEAR, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	3.5	3.7	0.3
ALASKA	2.9	2.5	-0.5
ARIZONA	8.1	6.5	-1.6
ARKANSAS	4.1	4.0	-0.2
COLORADO	4.3	4.0	-0.3
CONNECTICUT	5.7	6.1	0.4
DELAWARE	3.8	3.2	-0.5
DISTRICT OF CO	5.1	5.3	0.2
FLORIDA	5.0	6.1	1.1
GEORGIA	3.2	3.4	0.2
HAWAII	6.6	6.1	-0.4
IDAHO	4.8	5.0	0.1
ILLINOIS	5.9	6.9	1.0
INDIANA	4.7	5.9	1.2
IOWA	4.7	4.2	-0.5
KANSAS	4.9	3.8	-1.2
KENTUCKY	7.1	7.0	-0.2
LOUISIANA	8.1	6.0	-2.1
MAINE	2.4	5.4	3.1
MARYLAND	5.9	5.1	-0.8
MASSACHUSETTS	7.5	6.3	-1.2
MICHIGAN	5.3	5.6	0.2
MINNESOTA	4.6	4.0	-0.5
MISSISSIPPI	4.8	5.0	0.2
MISSOURI	5.1	4.3	-0.8
MONTANA	3.0	2.9	-0.1
NEBRASKA	5.7	7.1	1.4
NEVADA	4.9	5.1	0.1
NEW HAMPSHIRE	4.1	3.3	-0.8
NEW JERSEY	8.9	9.7	0.7
NEW MEXICO	4.9	5.4	0.5
NEW YORK	8.9	7.1	-1.8
NORTH CAROLINA	2.8	3.6	0.9
NORTH DAKOTA	3.6	5.8	2.2
OHIO	5.2	4.3	-0.9
OKLAHOMA	4.0	3.7	-0.3
OREGON	4.5	3.5	-1.0
PENNSYLVANIA	6.0	5.8	-0.2
RHODE ISLAND	7.3	6.1	-1.2
SOUTH CAROLINA	4.5	4.1	-0.4
SOUTH DAKOTA	4.6	3.6	-1.0
TENNESSEE	7.9	6.5	-1.5
TEXAS	7.1	7.2	0.1
UTAH	6.7	5.7	-1.0
VERMONT	4.9	6.3	1.4
VIRGINIA	5.7	5.9	0.1
WASHINGTON	3.5	3.5	-0.0
WEST VIRGINIA	6.3	4.7	-1.6
WISCONSIN	4.3	3.4	-0.9
WYOMING	3.1	3.6	0.5
PUERTO RICO	14.6	15.3	0.6

* NEVER HAD PAP SMEAR

TABLE 37.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	955	4.1	0.7	(2.7, 5.5)
ALASKA	907	2.9	0.9	(1.1, 4.6)
ARIZONA	1156	6.1	1.0	(4.2, 8.0)
ARKANSAS	1242	4.7	0.9	(3.1, 6.4)
COLORADO	1301	4.6	1.0	(2.6, 6.6)
CONNECTICUT	1967	5.9	0.7	(4.6, 7.2)
DELAWARE	1297	3.6	0.8	(2.1, 5.1)
DISTRICT OF CO	842	5.9	1.0	(4.0, 7.7)
FLORIDA	2274	6.7	0.6	(5.5, 8.0)
GEORGIA	1795	3.5	0.7	(2.2, 4.7)
HAWAII	2736	6.3	0.7	(5.0, 7.6)
IDAHO	2057	5.1	0.6	(3.9, 6.3)
ILLINOIS	1790	6.8	0.8	(5.3, 8.3)
INDIANA	1294	5.7	1.0	(3.9, 7.6)
IOWA	1675	4.0	0.7	(2.7, 5.3)
KANSAS	1866	3.9	0.5	(2.8, 4.9)
KENTUCKY	2692	5.8	0.6	(4.6, 7.0)
LOUISIANA	2162	6.0	0.6	(4.8, 7.2)
MAINE	2102	4.6	0.9	(2.8, 6.4)
MARYLAND	2263	5.1	0.6	(3.9, 6.3)
MASSACHUSETTS	4053	5.4	0.5	(4.5, 6.3)
MICHIGAN	1150	6.3	1.1	(4.2, 8.4)
MINNESOTA	1309	4.3	0.7	(2.9, 5.6)
MISSISSIPPI	885	4.5	0.8	(2.9, 6.2)
MISSOURI	1998	4.5	0.6	(3.3, 5.7)
MONTANA	1299	3.2	1.2	(0.8, 5.5)
NEBRASKA	1381	6.4	0.9	(4.6, 8.1)
NEVADA	765	5.6	1.3	(3.1, 8.1)
NEW HAMPSHIRE	957	3.1	0.8	(1.5, 4.7)
NEW JERSEY	1919	10.0	0.9	(8.2, 11.7)
NEW MEXICO	1327	6.1	0.8	(4.4, 7.7)
NEW YORK	1717	7.6	0.9	(5.9, 9.2)
NORTH CAROLINA	1357	4.3	0.7	(3.0, 5.7)
NORTH DAKOTA	860	5.4	1.0	(3.5, 7.3)
OHIO	1511	4.7	0.8	(3.1, 6.2)
OKLAHOMA	1468	4.4	0.7	(3.0, 5.8)
OREGON	1680	3.5	0.6	(2.5, 4.6)
PENNSYLVANIA	1665	5.6	0.7	(4.2, 7.0)
RHODE ISLAND	1759	5.5	0.7	(4.1, 6.9)
SOUTH CAROLINA	1405	4.3	0.7	(2.9, 5.6)
SOUTH DAKOTA	2385	3.5	0.5	(2.6, 4.4)
TENNESSEE	1458	5.6	0.8	(4.2, 7.1)
TEXAS	2239	8.6	0.8	(7.0, 10.2)
UTAH	1198	6.9	1.2	(4.6, 9.2)
VERMONT	1753	5.8	0.8	(4.2, 7.4)
VIRGINIA	887	6.4	1.6	(3.2, 9.5)
WASHINGTON	1600	3.8	0.6	(2.6, 4.9)
WEST VIRGINIA	1038	5.1	0.9	(3.3, 6.9)
WISCONSIN	1201	3.7	0.7	(2.5, 5.0)
WYOMING	977	4.7	0.9	(3.0, 6.3)
PUERTO RICO	2052	18.2	1.4	(15.5, 20.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 5.1 RANGE = 2.9-18.2

* NEVER HAD PAP SMEAR, WOMEN WITH INTACT CERVIX

April 17, 2001

TABLE 37.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	3.3	4.1	0.8
ALASKA	2.5	2.9	0.4
ARIZONA	9.1	6.1	-3.0
ARKANSAS	4.4	4.7	0.3
COLORADO	3.8	4.6	0.9
CONNECTICUT	5.7	5.9	0.1
DELAWARE	4.4	3.6	-0.8
DISTRICT OF CO	5.0	5.9	0.9
FLORIDA	5.2	6.7	1.5
GEORGIA	3.4	3.5	0.1
HAWAII	6.4	6.3	-0.1
IDAHO	5.5	5.1	-0.4
ILLINOIS	6.0	6.8	0.8
INDIANA	4.9	5.7	0.9
IOWA	4.6	4.0	-0.6
KANSAS	5.5	3.9	-1.6
KENTUCKY	6.8	5.8	-1.0
LOUISIANA	7.2	6.0	-1.2
MAINE	2.1	4.6	2.5
MARYLAND	6.0	5.1	-0.8
MASSACHUSETTS	7.3	5.4	-1.9
MICHIGAN	5.5	6.3	0.8
MINNESOTA	5.1	4.3	-0.8
MISSISSIPPI	5.4	4.5	-0.9
MISSOURI	4.3	4.5	0.2
MONTANA	3.0	3.2	0.1
NEBRASKA	5.6	6.4	0.8
NEVADA	5.6	5.6	-0.0
NEW HAMPSHIRE	4.2	3.1	-1.1
NEW JERSEY	9.3	10.0	0.7
NEW MEXICO	5.7	6.1	0.4
NEW YORK	9.2	7.6	-1.6
NORTH CAROLINA	2.8	4.3	1.6
NORTH DAKOTA	3.8	5.4	1.6
OHIO	5.6	4.7	-0.9
OKLAHOMA	4.4	4.4	-0.0
OREGON	4.4	3.5	-0.8
PENNSYLVANIA	6.4	5.6	-0.8
RHODE ISLAND	7.4	5.5	-1.9
SOUTH CAROLINA	5.3	4.3	-1.0
SOUTH DAKOTA	4.1	3.5	-0.6
TENNESSEE	7.3	5.6	-1.7
TEXAS	8.4	8.6	0.3
UTAH	8.3	6.9	-1.4
VERMONT	4.7	5.8	1.1
VIRGINIA	6.4	6.4	-0.1
WASHINGTON	3.7	3.8	0.1
WEST VIRGINIA	6.6	5.1	-1.5
WISCONSIN	4.5	3.7	-0.8
WYOMING	3.4	4.7	1.3
PUERTO RICO	17.4	18.2	0.7

* NEVER HAD PAP SMEAR, WOMEN WITH INTACT CERVIX

April 17, 2001

TABLE 38.1 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	949	13.7	1.4	(11.0, 16.3)
ALASKA	903	9.9	1.5	(7.0, 12.8)
ARIZONA	1155	12.1	1.4	(9.4, 14.9)
ARKANSAS	1231	16.4	1.2	(13.9, 18.8)
COLORADO	1300	12.4	1.4	(9.8, 15.1)
CONNECTICUT	1951	12.0	0.9	(10.2, 13.7)
DELAWARE	1288	8.0	0.9	(6.2, 9.8)
DISTRICT OF CO	832	11.3	1.3	(8.9, 13.8)
FLORIDA	2264	15.5	0.9	(13.7, 17.3)
GEORGIA	1781	10.9	1.0	(9.0, 12.8)
HAWAII	2724	12.2	0.8	(10.6, 13.8)
IDAHO	2053	16.9	1.0	(15.0, 18.8)
ILLINOIS	1774	15.5	1.0	(13.5, 17.4)
INDIANA	1287	15.3	1.2	(12.9, 17.7)
IOWA	1665	13.2	1.0	(11.1, 15.2)
KANSAS	1845	12.0	0.8	(10.4, 13.6)
KENTUCKY	2681	13.6	0.9	(11.9, 15.2)
LOUISIANA	2142	12.3	0.8	(10.7, 13.9)
MAINE	2086	11.0	1.2	(8.6, 13.3)
MARYLAND	2251	10.1	0.8	(8.5, 11.6)
MASSACHUSETTS	4033	10.4	0.6	(9.2, 11.6)
MICHIGAN	1143	13.3	1.3	(10.7, 15.8)
MINNESOTA	1303	14.0	1.1	(11.8, 16.2)
MISSISSIPPI	873	11.9	1.3	(9.5, 14.4)
MISSOURI	1986	15.3	1.1	(13.2, 17.4)
MONTANA	1292	11.2	1.5	(8.3, 14.1)
NEBRASKA	1372	14.0	1.2	(11.7, 16.2)
NEVADA	763	16.2	1.8	(12.7, 19.7)
NEW HAMPSHIRE	947	9.9	1.1	(7.7, 12.1)
NEW JERSEY	1910	18.1	1.1	(16.0, 20.2)
NEW MEXICO	1324	14.9	1.1	(12.7, 17.0)
NEW YORK	1711	14.4	1.1	(12.2, 16.5)
NORTH CAROLINA	1348	10.7	1.0	(8.8, 12.5)
NORTH DAKOTA	850	15.9	1.5	(13.1, 18.8)
OHIO	1495	13.5	1.3	(10.9, 16.1)
OKLAHOMA	1461	14.4	1.0	(12.4, 16.5)
OREGON	1671	12.0	0.9	(10.2, 13.7)
PENNSYLVANIA	1657	14.5	1.0	(12.5, 16.4)
RHODE ISLAND	1747	11.5	0.9	(9.7, 13.3)
SOUTH CAROLINA	1394	9.2	0.9	(7.5, 11.0)
SOUTH DAKOTA	2368	11.5	0.7	(10.1, 12.9)
TENNESSEE	1447	11.9	1.0	(10.0, 13.8)
TEXAS	2229	17.5	1.0	(15.5, 19.4)
UTAH	1195	16.6	1.5	(13.6, 19.7)
VERMONT	1743	11.9	1.0	(10.0, 13.8)
VIRGINIA	879	13.2	1.8	(9.8, 16.7)
WASHINGTON	1593	12.7	1.0	(10.8, 14.6)
WEST VIRGINIA	1035	17.9	1.4	(15.1, 20.7)
WISCONSIN	1197	13.2	1.2	(10.9, 15.5)
WYOMING	973	18.7	1.4	(15.9, 21.4)
PUERTO RICO	2035	27.3	1.5	(24.4, 30.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 13.2 RANGE = 8.0-27.3

* DID NOT HAVE A PAP SMEAR IN PAST THREE YEARS

April 17, 2001

TABLE 38.2 2000 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2000 NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2000 PERCENT	PERCENT DIFFERENCE
ALABAMA	13.4	13.7	0.3
ALASKA	9.1	9.9	0.8
ARIZONA	13.9	12.1	-1.8
ARKANSAS	17.4	16.4	-1.1
COLORADO	12.6	12.4	-0.1
CONNECTICUT	11.7	12.0	0.3
DELAWARE	12.3	8.0	-4.3
DISTRICT OF CO	9.2	11.3	2.2
FLORIDA	14.7	15.5	0.8
GEORGIA	12.0	10.9	-1.0
HAWAII	12.9	12.2	-0.7
IDAHO	20.7	16.9	-3.8
ILLINOIS	15.1	15.5	0.4
INDIANA	17.6	15.3	-2.4
IOWA	14.7	13.2	-1.5
KANSAS	12.6	12.0	-0.6
KENTUCKY	16.6	13.6	-3.0
LOUISIANA	14.6	12.3	-2.3
MAINE	10.5	11.0	0.5
MARYLAND	10.8	10.1	-0.8
MASSACHUSETTS	13.8	10.4	-3.4
MICHIGAN	14.2	13.3	-0.9
MINNESOTA	12.5	14.0	1.6
MISSISSIPPI	15.9	11.9	-3.9
MISSOURI	15.4	15.3	-0.0
MONTANA	14.8	11.2	-3.6
NEBRASKA	15.2	14.0	-1.2
NEVADA	21.5	16.2	-5.3
NEW HAMPSHIRE	11.6	9.9	-1.7
NEW JERSEY	15.7	18.1	2.4
NEW MEXICO	16.3	14.9	-1.4
NEW YORK	15.8	14.4	-1.4
NORTH CAROLINA	9.1	10.7	1.5
NORTH DAKOTA	13.4	15.9	2.5
OHIO	14.5	13.5	-0.9
OKLAHOMA	16.2	14.4	-1.8
OREGON	12.4	12.0	-0.5
PENNSYLVANIA	16.4	14.5	-2.0
RHODE ISLAND	14.0	11.5	-2.4
SOUTH CAROLINA	13.0	9.2	-3.7
SOUTH DAKOTA	14.6	11.5	-3.1
TENNESSEE	14.9	11.9	-3.0
TEXAS	17.3	17.5	0.1
UTAH	19.2	16.6	-2.5
VERMONT	11.6	11.9	0.3
VIRGINIA	14.5	13.2	-1.3
WASHINGTON	14.1	12.7	-1.4
WEST VIRGINIA	19.3	17.9	-1.4
WISCONSIN	15.3	13.2	-2.1
WYOMING	16.2	18.7	2.5
PUERTO RICO	26.9	27.3	0.4

* DID NOT HAVE A PAP SMEAR IN PAST THREE YEARS

April 17, 2001