

Mailstop K66
(770) 488-2466
August 9, 2002

Dear Colleague:

Information related to the 2001 BRFSS summary prevalence report of risk factors and preventive health measures from the core is attached.

This report focus on the following:

An introduction that contains:

An overview of the 2001 summary prevalence report.

An explanation of the tables for 2001 Behavioral Risk Factor Surveillance System (BRFSS) prevalence data.

Definitions of BRFSS risk factors and preventive health measures.

As soon as your schedule permits, please examine the enclosed materials closely.

Sincerely yours,

Ali Mokdad, Ph. D.
Chief, Behavioral Surveillance Branch
Division of Adult and Community Health
National Center for Chronic Disease
Prevention and Health Promotion

2001 BRFSS SUMMARY PREVALENCE REPORT

BEHAVIORAL SURVEILLANCE BRANCH
DIVISION OF ADULT AND COMMUNITY HEALTH
NATIONAL CENTER FOR CHRONIC DISEASE PREVENTION
AND HEALTH PROMOTION
CENTERS FOR DISEASE CONTROL AND PREVENTION

TABLE OF CONTENTS

OVERVIEW OF THE 2001 SUMMARY REPORT.....	I
EXPLANATION OF TABLES.....	II
DEFINITIONS OF RISK FACTORS & PREVENTIVE HEALTH MEASURES.....	IV
Limitation of Data.....	VI

SECTION I

PREVALENCE REPORT FOR CORE QUESTIONS

HEALTH INSURANCE

TABLE 1.1 HAVE NO HEALTH CARE COVERAGE, AGES 18-64.....	1
1.2 HAVE NO HEALTH CARE COVERAGE, AGES 18-64-BY SEX.....	2
1.3 HAVE NO HEALTH CARE COVERAGE, AGES 18-64, CHANGE 2000-2001.....	3

COLORECTAL CANCER SCREENING

TABLE 2.1 COLORECTAL SCREENING: NO FOBT IN PAST 2 YEARS, AGE 50 AND OLDER.....	4
2.2 COLORECTAL SCREENING: NO FOBT IN PAST 2 YEARS, AGE 50 AND OLDER-BY SEX.....	5
2.3 COLORECTAL SCREENING: NO FOBT IN PAST 2 YEARS, AGE 50 AND OLDER, CHANGE 1999-2001.....	6
TABLE 3.1 NEVER HAD A SIGMOIDOSCOPY/COLONOSCOPY, AGE 50 AND OLDER.....	7
3.2 NEVER HAD A SIGMOIDOSCOPY/COLONOSCOPY, AGE 50 AND OLDER-BY SEX.....	8
3.3 NEVER HAD A SIGMOIDOSCOPY/COLONOSCOPY, AGE 50 AND OLDER, CHANGE 1999-2001.....	9

DIABETES AWARENESS

TABLE 4.1 EVER TOLD YOU HAVE DIABETES.....	10
4.2 EVER TOLD YOU HAVE DIABETES-BY SEX.....	11
4.3 EVER TOLD YOU HAVE DIABETES, CHANGE 2000-2001.....	12

TABLE 5.1 EVER TOLD YOU HAVE DIABETES, AGE 20 AND OLDER.....	13
5.2 EVER TOLD YOU HAVE DIABETES, AGE 20 AND OLDER-BY SEX.....	14
5.3 EVER TOLD YOU HAVE DIABETES, AGE 20 AND OLDER, CHANGE 2000-2001.....	15

CHOLESTEROL AWARENESS

TABLE 6.1	CHOLESTEROL NOT CHECKED WITHIN PAST FIVE YEARS.....	16
6.2	CHOLESTEROL NOT CHECKED WITHIN PAST FIVE YEARS-BY SEX.....	17
6.3	CHOLESTEROL NOT CHECKED WITHIN PAST FIVE YEARS, CHANGE 1999-2001.....	18

IMMUNIZATION

TABLE 7.1	DIDN'T HAVE A FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OLDER.....	19
7.2	DIDN'T HAVE A FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OLDER-BY SEX.....	20
7.3	DIDN'T HAVE A FLU SHOT WITHIN PAST 12 MONTHS, AGE 65 AND OLDER, CHANGE 1999-2001.....	21

TABLE 8.1	NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65 AND OLDER.....	22
8.2	NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65 AND OLDER-BY SEX.....	23
8.3	NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65 AND OLDER, CHANGE 1999-2001.....	24

OBESITY

TABLE 9.1	OBSESE BASED ON BODY MASS INDEX (_BMI2 300 - 998).....	25
9.2	OBSESE BASED ON BODY MASS INDEX (_BMI2 300 - 998)-BY SEX.....	26
9.3	OBSESE BASED ON BODY MASS INDEX (_BMI2 300 - 998), CHANGE 2000-2001.....	27

TABLE 10.1	OBSESE BASED ON BODY MASS INDEX (_BMI2 300 - 998), AGE 20 AND OLDER.....	28
10.2	OBSESE BASED ON BODY MASS INDEX (_BMI2 300 - 998), AGE 20 AND OLDER BY-SEX.....	29
10.3	OBSESE BASED ON BODY MASS INDEX (_BMI2 300 - 998), AGE 20 AND OLDER, CHANGE 2000-2001....	30

EXERCISE

TABLE 11.1	NO LEISURE TIME EXERCISE OR PHYSICAL ACTIVITY.....	31
11.2	NO LEISURE TIME EXERCISE OR PHYSICAL ACTIVITY-BY SEX.....	32
11.3	NO LEISURE TIME EXERCISE OR PHYSICAL ACTIVITY, CHANGE 2000-2001.....	33

SMOKING STATUS

TABLE 12.1	CURRENT SMOKER.....	34
------------	------------------------	----

12.2	CURRENT SMOKER-BY	
SEX.....		35
12.3	CURRENT SMOKER, CHANGE	
2000-2001.....		36

ALCOHOL CONSUMPTION

TABLE 13.1	BINGE	
DRINKERS.....		37
13.2	BINGE DRINKERS-BY	
SEX.....		38
13.3	BINGE DRINKERS, CHANGE	
2000-2001.....		39

OVERVIEW OF THE 2001 SUMMARY REPORT

The report displays tables for selected variables from the survey core questions. We focused on core questions related to *Healthy People 2010 Objectives*. There are three tables for each risk factor or preventive measure unless otherwise noted. Estimated prevalence and relevant statistics are provided for the total population sampled and by sex. Prevalence estimates for both 2000 and 2001, or where appropriate 1999 and 2001, and the percentage point difference between the estimates are given for variables in the core.

Three tables (Tables 4.1–4.3) will be used as examples to explain the information contained in this report. Each table describes a behavioral risk factor or preventive health measure from the Behavioral Risk Factor Surveillance System (BRFSS) instrument.

Table 4.1 (page 10) shows the estimated prevalence and relevant statistics for each participant. (Participants refer to all states, and the District of Columbia and U.S. territories). The table shows the estimated prevalence and relevant statistics for "diabetes" in the population surveyed by each BRFSS participant. This prevalence estimate excludes respondents with unknown information on diabetes or respondents choosing not to answer the question.

The columns in Table 4.1 provide the following information:

COLUMN HEADINGCONTENTS

<u>Participant</u>	Name of the participant. Participants refers to all states including the District of Columbia and territories of the United States.
<u>Sample size</u>	The number of individuals who responded to the specific BRFSS question. These are unweighted numbers. These numbers <u>exclude</u> individuals who reported "not sure, don't know," those who refused to answer the question, or had no response coded (missing or skipped).
<u>Percent/Mean</u>	An estimate of the percent (or prevalence or mean number) of residents who report the behavioral risk factor of interest. This is weighted to the participant's population to estimate the true prevalence or mean in the population.
<u>Standard error(SE)</u>	A measure of the amount of variability that exists in the calculation of the prevalence estimate. SUDAAN was used to calculate SE
<u>95% confidence interval</u>	A range that contains the true population prevalence estimate 95% of the time, if repeated sampling of the population is performed. SUDAAN was used to calculate confidence intervals.

SUMMARY STATISTICS are given at the bottom of the table:

<u>Number of participants</u>	The total number of participants who asked question(s) regarding the respective "risk factor."
<u>Median</u>	The middle value of all estimated prevalences.
<u>Range</u>	The lowest prevalence and the highest prevalence among all participants.

For example, in Table 4.1, there were 2,795 individuals in the state of Alabama who responded to the "diabetes" question. Of those responding, the estimated prevalence of Alabama's population who reported that they had diabetes is 9.6%. This prevalence estimate has a standard error of 0.6. The 95% confidence interval for this estimate is (8.4, 10.8). Ninety-five percent (95%) of confidence intervals generated under repeated sampling will contain the true population prevalence of Alabama's population who "told they have diabetes."

Table 4.2 (page 11) shows the estimated prevalence and relevant statistics by sex for each state participant. The same statistics are displayed in Table 4.2 as in Table 4.1, except that Table 4.2 shows the estimated prevalence of “Diabetes” for males and females separately. This side by side display of the tables permits each participant to compare estimates for males and females in their respective state.

Table 4.3 (page 12) shows the percent difference between the prevalence estimates of diabetes in 2000 and 2001 survey years. This difference gives state health officials a quick method to examine changes that may have occurred in their respective state. In general, a negative change indicates a decrease in prevalence from 2000 to 2001, while a positive change indicates an increase in prevalence between the two years. Some of these changes may have been attributable to many factors such as change in the method of measurement, programs funded, or programs that came into existence because of results that states may have shown from the BRFSS. Therefore, conclusions drawn from this table should be interpreted with extreme caution.

Definitions of terms and risk factors in this report

Health Insurance

Health insurance – Respondents age 18–64 years who report they have no health care coverage. *Objective 2010– Increase the proportion of persons with health insurance (target 100%).*

Colorectal Cancer Exam

Sigmoidoscopy– Respondents, age 50 years and older, who report they never had a sigmoidoscopy. *Objective 2010–Increase colorectal cancer screening (target 50)*

Blood stool test– Respondents, age 50 years and older, who reported they did not use a home blood stool kit in the past two years. *Objective 2010– Increase colorectal cancer screening (target 50)*

Diabetes

Diabetes awareness– Respondents who report they were told by a doctor that they have diabetes. *Objective 2010– increase proportion of adults with diabetes whose condition has been diagnosed (target 80%).*

Cholesterol

Screening and awareness– Respondents who did not have their cholesterol checked within the past five years. *Objective 2010– increase the proportion of adults who have had their blood cholesterol checked within the preceding 5 years (target 80%).*

Immunization

Flu shot– Respondents, aged 65 years and older, who report they did not have a flu shot within the past 12 months. *Objective 2010– increase number of recipients in adults (target 90%).*

Pneumonia vaccination– Respondents, aged 65 years and older, who report they never had a pneumonia vaccination. *Objective 2010– increase number of recipients in adults (target 90%).*

Overweight

Overweight: by body mass index– Respondents with body mass index (BMI) equal to or greater than 30.0. BMI is defined as weight in kilograms divided by height in meters squared (w/h^{**2}).

Objective 2010– reduce the proportion of adults who are obese (target 15%).

Physical Activity

No leisure time physical activity– Respondents who report no leisure-time physical activity during the past month. *Objective 2010– reduce the proportion of adults who engage in no leisure time physical activity (target 20%).*

Smoking Status

Current smoker– Respondents who have smoked 100 cigarettes in their lifetime and reported smoking every day or some days. *Objective 2010– reduction in tobacco use by adults aged 18 years and older (target 12%).*

Binge Drinking (Acute drinking)– Respondents who report having five or more drinks on an occasion, one or more times in the past 30 days. *Objective 2010– reduction in adults and adolescents engaging in binge drinking during the past month (target 6%).*

Limitations of the Data

This report consolidates the most current data on selected risk factors and preventive health measures for states, the District of Columbia, and Puerto Rico. The information was obtained from the data files of individual states. Therefore, while all participants use telephone surveys to collect the data, the data in this report vary somewhat with respect to data collection, sampling design, and sample size.

A. Data Collection, Sample Design, and Sample Size

All states conduct interviews using a computer-assisted telephone interviewing (CATI) software program. All states use probability-based methods to select their sample. The BRFSS standard for sample designs is that sample records must be justifiable as a probability sample of all households with telephones in the state. All participating areas met this criterion in 2001. For data year 2001, fifty states and (one territory) used a disproportionate stratified sample (DSS) design. Puerto Rico, Guam, and Virgin Islands used a simple random sample design.

In a few states, a portion of sample records intended for use during one month may have been completed in another month. This deviation should only affect analyses based on monthly, rather than annual data. Illinois used a dual questionnaire or split sample survey design.

Several states did not collect data all 12 months of the year 2002. Hawaii, Texas, and Florida did not complete any interviews in January. New Jersey and Virgin Islands did not complete any interviews in January and February.

More information about the quality of the survey data can be found in the *2001 BRFSS Summary Data Quality Report*.

B. Sample Size, Estimates, and Data Limitations

Although almost all states collect data monthly, the annual number of interviews range from 871 in Guam to 8,628 in Massachusetts. Estimates for some states are therefore based upon smaller sample sizes, and for rare events, may yield unstable estimates. Some items such as immunizations are affected by seasonality, and estimates for these items may be biased for sub-populations.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. Respondents may not remember essential information, a question may not mean the same thing to different respondents, and some individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their impact on the data.

The data user must make his or her own evaluation of the data. Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. When the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the estimates or differences between groups and areas.

TABLE 1.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HEALTH CARE COVERAGE: AT RISK FOR NO HEALTH INSURANCE*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2221	17.8	1.0	(15.8, 19.8)
ALASKA	2593	20.3	1.3	(17.8, 22.8)
ARIZONA	2488	20.5	1.2	(18.1, 22.9)
ARKANSAS	2305	19.9	1.0	(17.9, 21.9)
CALIFORNIA	3487	17.3	0.8	(15.7, 18.9)
COLORADO	1722	17.9	1.1	(15.7, 20.1)
CONNECTICUT	6164	11.3	0.6	(10.1, 12.5)
DELAWARE	2744	10.1	0.8	(8.5, 11.7)
DISTRICT OF CO	1565	14.2	1.2	(11.8, 16.6)
FLORIDA	3459	21.5	0.9	(19.7, 23.3)
GEORGIA	3798	15.9	0.8	(14.3, 17.5)
HAWAII	3634	8.8	0.7	(7.4, 10.2)
IDAHO	3826	17.9	0.8	(16.3, 19.5)
ILLINOIS	3246	11.7	0.8	(10.1, 13.3)
INDIANA	3209	16.2	0.8	(14.6, 17.8)
IOWA	2699	10.1	0.7	(8.7, 11.5)
KANSAS	3673	12.1	0.6	(10.9, 13.3)
KENTUCKY	5882	18.0	0.7	(16.6, 19.4)
LOUISIANA	4060	25.3	0.8	(23.7, 26.9)
MAINE	1911	15.3	1.0	(13.3, 17.3)
MARYLAND	3639	11.8	0.8	(10.2, 13.4)
MASSACHUSETTS	7002	9.4	0.4	(8.6, 10.2)
MICHIGAN	3104	11.6	0.7	(10.2, 13.0)
MINNESOTA	3168	6.4	0.6	(5.2, 7.6)
MISSISSIPPI	2420	22.0	1.0	(20.0, 24.0)
MISSOURI	3243	12.9	0.8	(11.3, 14.5)
MONTANA	2589	20.4	1.1	(18.2, 22.6)
NEBRASKA	2797	16.5	0.9	(14.7, 18.3)
NEVADA	2108	20.2	1.5	(17.3, 23.1)
NEW HAMPSHIRE	3324	13.4	0.7	(12.0, 14.8)
NEW JERSEY	4801	13.5	0.7	(12.1, 14.9)
NEW MEXICO	2867	26.5	1.0	(24.5, 28.5)
NEW YORK	3221	19.5	0.9	(17.7, 21.3)
NORTH CAROLINA	4941	16.7	0.9	(14.9, 18.5)
NORTH DAKOTA	2007	14.2	0.9	(12.4, 16.0)
OHIO	2728	13.0	0.8	(11.4, 14.6)
OKLAHOMA	3464	25.1	1.0	(23.1, 27.1)
OREGON	2044	16.6	0.9	(14.8, 18.4)
PENNSYLVANIA	2836	11.3	0.7	(9.9, 12.7)
RHODE ISLAND	3245	10.5	0.8	(8.9, 12.1)
SOUTH CAROLINA	2610	19.2	1.0	(17.2, 21.2)
SOUTH DAKOTA	3794	12.4	0.7	(11.0, 13.8)
TENNESSEE	2390	12.4	0.9	(10.6, 14.2)
TEXAS	4874	26.6	0.8	(25.0, 28.2)
UTAH	3067	14.6	0.9	(12.8, 16.4)
VERMONT	3557	13.4	0.7	(12.0, 14.8)
VIRGINIA	2406	12.7	0.8	(11.1, 14.3)
WASHINGTON	3454	11.6	0.7	(10.2, 13.0)
WEST VIRGINIA	2323	23.8	1.1	(21.6, 26.0)
WISCONSIN	2753	11.7	0.7	(10.3, 13.1)
WYOMING	2487	19.5	0.9	(17.7, 21.3)
GUAM	801	20.0	1.7	(16.7, 23.3)
PUERTO RICO	3290	9.3	0.8	(7.7, 10.9)
VIRGIN ISLANDS	1931	31.8	1.4	(29.1, 34.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 15.6 RANGE = 6.4-31.8

HP2010: LHI, HP 1.1

*AT RISK FOR NO HEALTH INSURANCE, ADULTS AGE 18-64

August 8,

TABLE 1.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HEALTH CARE COVERAGE: AT RISK FOR NO HEALTH INSURANCE* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	865	16.9	1.5	(14.0, 19.8)	*	1356	18.6	1.2	(16.2, 21.0)	*
ALASKA	1231	20.3	1.8	(16.8, 23.8)	*	1362	20.4	2.0	(16.5, 24.3)	*
ARIZONA	1028	20.1	1.8	(16.6, 23.6)	*	1460	20.9	1.6	(17.8, 24.0)	*
ARKANSAS	923	19.0	1.5	(16.1, 21.9)	*	1382	20.8	1.3	(18.3, 23.3)	*
CALIFORNIA	1438	18.0	1.2	(15.6, 20.4)	*	2049	16.5	1.0	(14.5, 18.5)	*
COLORADO	732	18.2	1.7	(14.9, 21.5)	*	990	17.6	1.5	(14.7, 20.5)	*
CONNECTICUT	2613	12.6	0.9	(10.8, 14.4)	*	3551	10.0	0.7	(8.6, 11.4)	*
DELAWARE	1093	12.3	1.3	(9.8, 14.8)	*	1651	7.9	0.8	(6.3, 9.5)	*
DISTRICT OF CO	676	18.0	2.1	(13.9, 22.1)	*	889	10.7	1.4	(8.0, 13.4)	*
FLORIDA	1444	22.6	1.3	(20.1, 25.1)	*	2015	20.4	1.1	(18.2, 22.6)	*
GEORGIA	1484	17.3	1.4	(14.6, 20.0)	*	2314	14.6	1.0	(12.6, 16.6)	*
HAWAII	1604	10.5	1.1	(8.3, 12.7)	*	2030	6.9	0.7	(5.5, 8.3)	*
IDAHO	1670	17.9	1.2	(15.5, 20.3)	*	2156	17.9	1.0	(15.9, 19.9)	*
ILLINOIS	1376	11.9	1.2	(9.5, 14.3)	*	1870	11.6	0.9	(9.8, 13.4)	*
INDIANA	1351	17.5	1.2	(15.1, 19.9)	*	1858	14.9	1.0	(12.9, 16.9)	*
IOWA	1183	9.4	1.0	(7.4, 11.4)	*	1516	10.7	1.0	(8.7, 12.7)	*
KANSAS	1537	12.2	0.9	(10.4, 14.0)	*	2136	11.9	0.8	(10.3, 13.5)	*
KENTUCKY	2392	17.6	1.1	(15.4, 19.8)	*	3490	18.3	0.9	(16.5, 20.1)	*
LOUISIANA	1588	25.4	1.2	(23.0, 27.8)	*	2472	25.2	1.0	(23.2, 27.2)	*
MAINE	836	18.0	1.6	(14.9, 21.1)	*	1075	12.6	1.2	(10.2, 15.0)	*
MARYLAND	1502	12.7	1.5	(9.8, 15.6)	*	2137	10.8	0.9	(9.0, 12.6)	*
MASSACHUSETTS	2986	10.9	0.7	(9.5, 12.3)	*	4016	8.0	0.6	(6.8, 9.2)	*
MICHIGAN	1266	12.0	1.1	(9.8, 14.2)	*	1838	11.2	0.9	(9.4, 13.0)	*
MINNESOTA	1397	8.0	1.0	(6.0, 10.0)	*	1771	5.0	0.6	(3.8, 6.2)	*
MISSISSIPPI	953	22.1	1.6	(19.0, 25.2)	*	1467	22.0	1.2	(19.6, 24.4)	*
MISSOURI	1352	13.6	1.3	(11.1, 16.1)	*	1891	12.2	0.9	(10.4, 14.0)	*
MONTANA	1159	19.5	1.6	(16.4, 22.6)	*	1430	21.4	1.5	(18.5, 24.3)	*
NEBRASKA	1130	16.3	1.5	(13.4, 19.2)	*	1667	16.6	1.2	(14.2, 19.0)	*
NEVADA	1081	23.5	2.4	(18.8, 28.2)	*	1027	16.8	1.7	(13.5, 20.1)	*
NEW HAMPSHIRE	1442	15.6	1.1	(13.4, 17.8)	*	1882	11.3	0.9	(9.5, 13.1)	*
NEW JERSEY	2004	14.4	1.2	(12.0, 16.8)	*	2797	12.6	0.8	(11.0, 14.2)	*
NEW MEXICO	1224	28.3	1.5	(25.4, 31.2)	*	1643	24.8	1.2	(22.4, 27.2)	*
NEW YORK	1352	22.5	1.5	(19.6, 25.4)	*	1869	16.6	1.1	(14.4, 18.8)	*
NORTH CAROLINA	1961	17.7	1.4	(15.0, 20.4)	*	2980	15.7	1.1	(13.5, 17.9)	*
NORTH DAKOTA	898	16.1	1.4	(13.4, 18.8)	*	1109	12.3	1.1	(10.1, 14.5)	*
OHIO	1111	13.3	1.4	(10.6, 16.0)	*	1617	12.6	1.0	(10.6, 14.6)	*
OKLAHOMA	1402	26.1	1.6	(23.0, 29.2)	*	2062	24.1	1.3	(21.6, 26.6)	*
OREGON	870	18.2	1.5	(15.3, 21.1)	*	1174	15.0	1.2	(12.6, 17.4)	*
PENNSYLVANIA	1210	13.9	1.2	(11.5, 16.3)	*	1626	8.9	0.8	(7.3, 10.5)	*
RHODE ISLAND	1272	13.1	1.3	(10.6, 15.6)	*	1973	7.9	0.8	(6.3, 9.5)	*
SOUTH CAROLINA	1063	19.1	1.5	(16.2, 22.0)	*	1547	19.3	1.3	(16.8, 21.8)	*
SOUTH DAKOTA	1687	13.7	1.1	(11.5, 15.9)	*	2107	11.0	0.7	(9.6, 12.4)	*
TENNESSEE	813	15.6	1.5	(12.7, 18.5)	*	1577	9.4	0.8	(7.8, 11.0)	*
TEXAS	2061	25.8	1.1	(23.6, 28.0)	*	2813	27.5	1.0	(25.5, 29.5)	*
UTAH	1406	16.1	1.4	(13.4, 18.8)	*	1661	13.2	1.1	(11.0, 15.4)	*
VERMONT	1527	16.4	1.2	(14.0, 18.8)	*	2030	10.3	0.8	(8.7, 11.9)	*
VIRGINIA	1035	13.6	1.3	(11.1, 16.1)	*	1371	11.8	1.1	(9.6, 14.0)	*
WASHINGTON	1489	12.2	1.0	(10.2, 14.2)	*	1965	10.9	0.9	(9.1, 12.7)	*
WEST VIRGINIA	956	24.9	1.7	(21.6, 28.2)	*	1367	22.6	1.3	(20.1, 25.1)	*
WISCONSIN	1273	13.0	1.1	(10.8, 15.2)	*	1480	10.4	1.0	(8.4, 12.4)	*
WYOMING	1115	18.8	1.3	(16.3, 21.3)	*	1372	20.3	1.2	(17.9, 22.7)	*
GUAM	355	22.5	2.7	(17.2, 27.8)	*	446	17.2	2.2	(12.9, 21.5)	*
PUERTO RICO	1182	12.9	1.5	(10.0, 15.8)	*	2108	6.1	0.8	(4.5, 7.7)	*
VIRGIN ISLANDS	751	32.7	2.1	(28.6, 36.8)	*	1180	31.0	1.8	(27.5, 34.5)	*

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 17.1
 RANGE = 8.0-32.7

* AT RISK FOR NO HEALTH INSURANCE, ADULTS AGE 18-64

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 13.9
 RANGE = 5.0-31.0

August 8, 2002

TABLE 1.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 HEALTH CARE COVERAGE: AT RISK FOR NO HEALTH INSURANCE*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	19.4	17.8	-1.6	-8.4
ALASKA	19.1	20.3	1.2	6.5
ARIZONA	20.7	20.5	-0.2	-0.9
ARKANSAS	20.9	19.9	-1.0	-5.0
CALIFORNIA	21.3	17.3	-4.0	-18.6
COLORADO	15.8	17.9	2.1	13.4
CONNECTICUT	10.6	11.3	0.7	6.4
DELAWARE	9.7	10.1	0.4	3.8
DISTRICT OF CO	12.8	14.2	1.4	10.8
FLORIDA	21.6	21.5	-0.1	-0.4
GEORGIA	16.5	15.9	-0.6	-3.8
HAWAII	8.3	8.8	0.5	6.5
IDAHO	20.4	17.9	-2.5	-12.2
ILLINOIS	12.8	11.7	-1.1	-8.7
INDIANA	12.3	16.2	3.9	31.8
IOWA	10.9	10.1	-0.8	-7.1
KANSAS	12.9	12.1	-0.8	-6.5
KENTUCKY	16.6	18.0	1.4	8.3
LOUISIANA	25.6	25.3	-0.3	-1.1
MAINE	16.3	15.3	-1.0	-6.0
MARYLAND	11.2	11.8	0.7	5.8
MASSACHUSETTS	9.9	9.4	-0.5	-4.9
MICHIGAN	10.1	11.6	1.5	14.5
MINNESOTA	8.3	6.4	-1.9	-23.1
MISSISSIPPI	22.7	22.0	-0.7	-3.0
MISSOURI	13.4	12.9	-0.5	-3.7
MONTANA	18.0	20.4	2.4	13.1
NEBRASKA	11.1	16.5	5.4	48.1
NEVADA	16.0	20.2	4.2	26.2
NEW HAMPSHIRE	10.3	13.4	3.1	30.0
NEW JERSEY	15.4	13.5	-1.9	-12.1
NEW MEXICO	27.7	26.5	-1.2	-4.2
NEW YORK	15.3	19.5	4.2	27.1
NORTH CAROLINA	15.1	16.7	1.6	10.4
NORTH DAKOTA	14.3	14.2	-0.1	-0.4
OHIO	12.3	13.0	0.7	6.0
OKLAHOMA	20.1	25.1	5.0	24.6
OREGON	18.1	16.6	-1.5	-8.3
PENNSYLVANIA	11.1	11.3	0.2	1.7
RHODE ISLAND	13.6	10.5	-3.1	-22.5
SOUTH CAROLINA	16.9	19.2	2.3	13.5
SOUTH DAKOTA	12.7	12.4	-0.3	-2.5
TENNESSEE	13.2	12.4	-0.8	-5.8
TEXAS	26.9	26.6	-0.3	-1.0
UTAH	12.8	14.6	1.9	14.5
VERMONT	11.7	13.4	1.7	14.9
VIRGINIA	12.9	12.7	-0.2	-1.9
WASHINGTON	11.5	11.6	0.1	0.5
WEST VIRGINIA	23.5	23.8	0.3	1.3
WISCONSIN	8.9	11.7	2.8	31.6
WYOMING	20.2	19.5	-0.6	-3.2
GUAM	N/A	20.0	N/A	N/A
PUERTO RICO	10.1	9.3	-0.8	-8.2
VIRGIN ISLANDS	N/A	31.8	N/A	N/A
SUMMARY STATISTICS: yr2000				
NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	54	
MEDIAN =	13.9	MEDIAN =	15.6	
RANGE =	8.3-27.7	RANGE =	6.4-31.8	

* AT RISK FOR NO HEALTH INSURANCE, ADULTS AGE 18-64
 N/A - NOT AVAILABLE

August 8, 2002

TABLE 2.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BLDSTOOL: NO FECAL OCCULT BLOOD STOOL TEST IN PAST TWO YEARS, AGE 50 AND OLDER*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	918	88.1	1.2	(85.7, 90.5)
ALASKA	912	78.5	2.0	(74.6, 82.4)
ARIZONA	1433	65.6	1.8	(62.1, 69.1)
ARKANSAS	1270	74.1	1.4	(71.4, 76.8)
CALIFORNIA	1585	70.0	1.4	(67.3, 72.7)
COLORADO	686	63.8	2.0	(59.9, 67.7)
CONNECTICUT	3111	62.9	1.0	(60.9, 64.9)
DELAWARE	1465	70.5	1.5	(67.6, 73.4)
DISTRICT OF CO	673	62.1	2.1	(58.0, 66.2)
FLORIDA	2022	63.4	1.2	(61.0, 65.8)
GEORGIA	1608	67.6	1.4	(64.9, 70.3)
HAWAII	1830	61.1	1.6	(58.0, 64.2)
IDAHO	1931	73.5	1.2	(71.1, 75.9)
ILLINOIS	853	71.1	1.7	(67.8, 74.4)
INDIANA	1580	70.9	1.3	(68.4, 73.4)
IOWA	1585	66.2	1.3	(63.7, 68.7)
KANSAS	1713	65.6	1.3	(63.1, 68.1)
KENTUCKY	3314	70.8	1.1	(68.6, 73.0)
LOUISIANA	1835	71.4	1.2	(69.0, 73.8)
MAINE	1070	57.6	1.6	(54.5, 60.7)
MARYLAND	1721	57.4	1.5	(54.5, 60.3)
MASSACHUSETTS	3111	61.2	1.0	(59.2, 63.2)
MICHIGAN	1562	63.5	1.4	(60.8, 66.2)
MINNESOTA	1630	65.3	1.3	(62.8, 67.8)
MISSISSIPPI	1215	78.5	1.3	(76.0, 81.0)
MISSOURI	1873	69.1	1.5	(66.2, 72.0)
MONTANA	1566	70.1	1.6	(67.0, 73.2)
NEBRASKA	1562	65.4	1.3	(62.9, 67.9)
NEVADA	1050	71.2	2.6	(66.1, 76.3)
NEW HAMPSHIRE	1594	59.3	1.3	(56.8, 61.8)
NEW JERSEY	2267	68.4	1.3	(65.9, 70.9)
NEW MEXICO	1530	73.8	1.4	(71.1, 76.5)
NEW YORK	1402	66.5	1.5	(63.6, 69.4)
NORTH CAROLINA	2491	56.9	1.4	(54.2, 59.6)
NORTH DAKOTA	952	74.4	1.5	(71.5, 77.3)
OHIO	1340	67.8	1.5	(64.9, 70.7)
OKLAHOMA	2051	75.6	1.2	(73.2, 78.0)
OREGON	1016	61.3	1.7	(58.0, 64.6)
PENNSYLVANIA	1612	69.3	1.3	(66.8, 71.8)
RHODE ISLAND	1617	65.8	1.3	(63.3, 68.3)
SOUTH CAROLINA	1207	68.4	1.5	(65.5, 71.3)
SOUTH DAKOTA	2308	69.0	1.1	(66.8, 71.2)
TENNESSEE	1169	71.8	1.5	(68.9, 74.7)
TEXAS	2176	75.7	1.0	(73.7, 77.7)
UTAH	1253	79.1	1.6	(76.0, 82.2)
VERMONT	1769	56.8	1.3	(54.3, 59.3)
VIRGINIA	1140	66.7	1.5	(63.8, 69.6)
WASHINGTON	1531	59.0	1.4	(56.3, 61.7)
WEST VIRGINIA	1459	73.4	1.2	(71.0, 75.8)
WISCONSIN	1260	67.5	1.5	(64.6, 70.4)
WYOMING	1217	78.0	1.3	(75.5, 80.5)
GUAM	189	86.5	2.9	(80.8, 92.2)
PUERTO RICO	2008	99.9	0.1	(99.7, 100.1)
VIRGIN ISLANDS	801	86.2	1.6	(83.1, 89.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 68.7 RANGE = 56.8-99.9
 HP2010: LHI, HP 3.12a(approx.)

*AT RISK NO FECAL OCCULT BLOOD STOOL TEST IN PAST TWO YEARS, AGE 50 AND OLDER

August 8, 2002

TABLE 2.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BLDSTOOL: NO FECAL OCCULT BLOOD STOOL TEST IN PAST TWO YEARS, AGE 50 AND OLDER* -BY SEX

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	326	87.3	2.0	(83.4, 91.2)	*	592	88.8	1.5	(85.9, 91.7)
ALASKA	438	75.4	3.2	(69.1, 81.7)	*	474	81.7	2.4	(77.0, 86.4)
ARIZONA	574	67.5	2.6	(62.4, 72.6)	*	859	64.0	2.4	(59.3, 68.7)
ARKANSAS	465	72.2	2.3	(67.7, 76.7)	*	805	75.6	1.6	(72.5, 78.7)
CALIFORNIA	596	71.8	2.1	(67.7, 75.9)	*	989	68.5	1.7	(65.2, 71.8)
COLORADO	276	63.5	3.1	(57.4, 69.6)	*	410	64.1	2.6	(59.0, 69.2)
CONNECTICUT	1226	63.8	1.6	(60.7, 66.9)	*	1885	62.2	1.3	(59.7, 64.7)
DELAWARE	515	69.7	2.5	(64.8, 74.6)	*	950	71.1	1.9	(67.4, 74.8)
DISTRICT OF CO	271	59.4	3.4	(52.7, 66.1)	*	402	64.1	2.7	(58.8, 69.4)
FLORIDA	784	61.8	1.9	(58.1, 65.5)	*	1238	64.6	1.5	(61.7, 67.5)
GEORGIA	574	67.8	2.3	(63.3, 72.3)	*	1034	67.5	1.8	(64.0, 71.0)
HAWAII	750	62.3	2.4	(57.6, 67.0)	*	1080	60.0	2.2	(55.7, 64.3)
IDAHO	794	76.9	1.8	(73.4, 80.4)	*	1137	70.6	1.6	(67.5, 73.7)
ILLINOIS	317	71.9	2.8	(66.4, 77.4)	*	536	70.5	2.1	(66.4, 74.6)
INDIANA	593	70.5	2.0	(66.6, 74.4)	*	987	71.3	1.6	(68.2, 74.4)
IOWA	593	68.5	2.1	(64.4, 72.6)	*	992	64.2	1.7	(60.9, 67.5)
KANSAS	650	66.2	2.0	(62.3, 70.1)	*	1063	65.1	1.6	(62.0, 68.2)
KENTUCKY	1212	70.9	1.8	(67.4, 74.4)	*	2102	70.7	1.4	(68.0, 73.4)
LOUISIANA	618	71.5	2.0	(67.6, 75.4)	*	1217	71.3	1.5	(68.4, 74.2)
MAINE	429	59.1	2.5	(54.2, 64.0)	*	641	56.4	2.1	(52.3, 60.5)
MARYLAND	704	57.8	2.3	(53.3, 62.3)	*	1017	57.1	2.0	(53.2, 61.0)
MASSACHUSETTS	1212	60.4	1.6	(57.3, 63.5)	*	1899	61.9	1.3	(59.4, 64.4)
MICHIGAN	618	63.7	2.1	(59.6, 67.8)	*	944	63.3	1.8	(59.8, 66.8)
MINNESOTA	674	69.2	1.9	(65.5, 72.9)	*	956	62.0	1.7	(58.7, 65.3)
MISSISSIPPI	427	75.1	2.3	(70.6, 79.6)	*	788	81.2	1.5	(78.3, 84.1)
MISSOURI	681	66.9	2.5	(62.0, 71.8)	*	1192	70.9	1.8	(67.4, 74.4)
MONTANA	636	70.7	2.4	(66.0, 75.4)	*	930	69.7	2.1	(65.6, 73.8)
NEBRASKA	563	63.8	2.2	(59.5, 68.1)	*	999	66.7	1.7	(63.4, 70.0)
NEVADA	534	70.2	4.4	(61.6, 78.8)	*	516	72.0	2.7	(66.7, 77.3)
NEW HAMPSHIRE	654	59.0	2.1	(54.9, 63.1)	*	940	59.5	1.8	(56.0, 63.0)
NEW JERSEY	844	66.9	2.0	(63.0, 70.8)	*	1423	69.5	1.6	(66.4, 72.6)
NEW MEXICO	601	70.4	2.3	(65.9, 74.9)	*	929	76.7	1.6	(73.6, 79.8)
NEW YORK	579	65.5	2.3	(61.0, 70.0)	*	823	67.2	1.9	(63.5, 70.9)
NORTH CAROLINA	906	58.9	2.3	(54.4, 63.4)	*	1585	55.4	1.8	(51.9, 58.9)
NORTH DAKOTA	359	79.4	2.3	(74.9, 83.9)	*	593	70.2	2.0	(66.3, 74.1)
OHIO	499	68.3	2.3	(63.8, 72.8)	*	841	67.4	1.9	(63.7, 71.1)
OKLAHOMA	776	75.2	1.9	(71.5, 78.9)	*	1275	75.8	1.5	(72.9, 78.7)
OREGON	399	64.7	2.5	(59.8, 69.6)	*	617	58.6	2.2	(54.3, 62.9)
PENNSYLVANIA	609	68.4	2.0	(64.5, 72.3)	*	1003	70.0	1.6	(66.9, 73.1)
RHODE ISLAND	574	62.9	2.2	(58.6, 67.2)	*	1043	67.9	1.6	(64.8, 71.0)
SOUTH CAROLINA	464	66.6	2.4	(61.9, 71.3)	*	743	69.8	1.8	(66.3, 73.3)
SOUTH DAKOTA	943	68.5	1.7	(65.2, 71.8)	*	1365	69.4	1.4	(66.7, 72.1)
TENNESSEE	358	69.5	2.7	(64.2, 74.8)	*	811	73.7	1.7	(70.4, 77.0)
TEXAS	850	75.5	1.6	(72.4, 78.6)	*	1326	75.8	1.3	(73.3, 78.3)
UTAH	558	81.5	2.2	(77.2, 85.8)	*	695	77.2	2.2	(72.9, 81.5)
VERMONT	704	56.9	2.0	(53.0, 60.8)	*	1065	56.8	1.7	(53.5, 60.1)
VIRGINIA	481	65.5	2.3	(61.0, 70.0)	*	659	67.7	2.0	(63.8, 71.6)
WASHINGTON	610	63.0	2.1	(58.9, 67.1)	*	921	55.6	1.8	(52.1, 59.1)
WEST VIRGINIA	561	69.7	2.1	(65.6, 73.8)	*	898	76.3	1.5	(73.4, 79.2)
WISCONSIN	498	68.8	2.3	(64.3, 73.3)	*	762	66.4	2.0	(62.5, 70.3)
WYOMING	495	78.9	1.9	(75.2, 82.6)	*	722	77.2	1.7	(73.9, 80.5)
GUAM	91	87.2	4.4	(78.6, 95.8)	*	98	85.6	3.8	(78.2, 93.0)
PUERTO RICO	718	99.9	0.1	(99.7, 100.1)	*	1290	99.9	0.1	(99.7, 100.1)
VIRGIN ISLANDS	332	86.3	2.3	(81.8, 90.8)	*	469	86.1	2.1	(82.0, 90.2)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 68.5
 RANGE = 56.9-99.9

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 69.5
 RANGE = 55.4-99.9

*AT RISK NO FECAL OCCULT BLOOD STOOL TEST IN PAST TWO YEARS, AGE 50 AND OLDER

August 8, 2002

TABLE 2.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2001
 BLDSTOOL: NO FECAL OCCULT BLOOD STOOL TEST IN PAST TWO YEARS, AGE 50 AND OLDER*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	82.4	88.1	5.7	7.0
ALASKA	82.7	78.5	-4.2	-5.1
ARIZONA	67.1	65.6	-1.5	-2.2
ARKANSAS	77.1	74.1	-3.0	-3.9
CALIFORNIA	77.0	70.0	-7.0	-9.1
COLORADO	70.3	63.8	-6.5	-9.3
CONNECTICUT	63.2	62.9	-0.3	-0.5
DELAWARE	71.0	70.5	-0.5	-0.7
DISTRICT OF CO	57.0	62.1	5.1	8.9
FLORIDA	67.6	63.4	-4.2	-6.3
GEORGIA	78.3	67.6	-10.7	-13.6
HAWAII	71.3	61.1	-10.2	-14.3
IDAHO	76.9	73.5	-3.4	-4.4
ILLINOIS	74.2	71.1	-3.1	-4.1
INDIANA	77.6	70.9	-6.6	-8.6
IOWA	67.0	66.2	-0.8	-1.1
KANSAS	73.9	65.6	-8.3	-11.2
KENTUCKY	76.1	70.8	-5.3	-6.9
LOUISIANA	76.2	71.4	-4.8	-6.3
MAINE	64.1	57.6	-6.4	-10.1
MARYLAND	62.2	57.4	-4.8	-7.6
MASSACHUSETTS	64.8	61.2	-3.6	-5.5
MICHIGAN	66.9	63.5	-3.4	-5.1
MINNESOTA	73.8	65.3	-8.5	-11.5
MISSISSIPPI	82.2	78.5	-3.7	-4.5
MISSOURI	75.0	69.1	-5.9	-7.8
MONTANA	73.6	70.1	-3.5	-4.8
NEBRASKA	72.6	65.4	-7.2	-9.9
NEVADA	80.0	71.2	-8.8	-11.0
NEW HAMPSHIRE	59.3	59.3	0.0	0.0
NEW JERSEY	64.6	68.4	3.8	5.8
NEW MEXICO	76.1	73.8	-2.3	-3.0
NEW YORK	70.8	66.5	-4.3	-6.1
NORTH CAROLINA	61.5	56.9	-4.6	-7.4
NORTH DAKOTA	75.5	74.4	-1.1	-1.4
OHIO	69.1	67.8	-1.3	-1.9
OKLAHOMA	76.7	75.6	-1.1	-1.4
OREGON	67.8	61.3	-6.5	-9.6
PENNSYLVANIA	69.0	69.3	0.3	0.4
RHODE ISLAND	68.6	65.8	-2.8	-4.1
SOUTH CAROLINA	73.6	68.4	-5.2	-7.0
SOUTH DAKOTA	74.8	69.0	-5.8	-7.8
TENNESSEE	76.7	71.8	-4.9	-6.4
TEXAS	77.0	75.7	-1.3	-1.7
UTAH	80.2	79.1	-1.1	-1.4
VERMONT	62.6	56.8	-5.8	-9.3
VIRGINIA	73.8	66.7	-7.1	-9.6
WASHINGTON	64.3	59.0	-5.3	-8.3
WEST VIRGINIA	80.9	73.4	-7.5	-9.3
WISCONSIN	78.7	67.5	-11.2	-14.2
WYOMING	81.6	78.0	-3.6	-4.4
GUAM	N/A	86.5	N/A	N/A
PUERTO RICO	88.2	99.9	11.8	13.3
VIRGIN ISLANDS	N/A	86.2	N/A	N/A

SUMMARY STATISTICS: yr1999

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 73.8

RANGE

= 57.0-88.2

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 68.7

RANGE

= 56.8-99.9

*AT RISK NO FECAL OCCULT BLOOD STOOL TEST IN PAST TWO YEARS, AGE 50 AND OLDER

TABLE 3.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD SIGMOIDOSCOPY: NEVER HAD A SIGMOIDOSCOPY*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1146	55.3	1.7	(52.0, 58.6)
ALASKA	914	55.3	2.5	(50.4, 60.2)
ARIZONA	1436	50.0	1.9	(46.3, 53.7)
ARKANSAS	1281	55.4	1.5	(52.5, 58.3)
CALIFORNIA	1586	51.5	1.6	(48.4, 54.6)
COLORADO	693	52.9	2.1	(48.8, 57.0)
CONNECTICUT	3119	46.3	1.1	(44.1, 48.5)
DELAWARE	1478	41.6	1.6	(38.5, 44.7)
DISTRICT OF CO	669	44.0	2.2	(39.7, 48.3)
FLORIDA	2041	52.0	1.3	(49.5, 54.5)
GEORGIA	1631	51.1	1.5	(48.2, 54.0)
HAWAII	1834	50.2	1.6	(47.1, 53.3)
IDAHO	1931	56.6	1.4	(53.9, 59.3)
ILLINOIS	852	59.1	1.9	(55.4, 62.8)
INDIANA	1584	54.5	1.4	(51.8, 57.2)
IOWA	1613	54.5	1.4	(51.8, 57.2)
KANSAS	1723	55.6	1.3	(53.1, 58.1)
KENTUCKY	3346	60.7	1.2	(58.3, 63.1)
LOUISIANA	1847	60.6	1.3	(58.1, 63.1)
MAINE	1068	52.3	1.7	(49.0, 55.6)
MARYLAND	1719	47.6	1.5	(44.7, 50.5)
MASSACHUSETTS	3119	46.8	1.0	(44.8, 48.8)
MICHIGAN	1572	44.8	1.4	(42.1, 47.5)
MINNESOTA	1636	38.0	1.3	(35.5, 40.5)
MISSISSIPPI	1218	59.2	1.6	(56.1, 62.3)
MISSOURI	1874	57.0	1.6	(53.9, 60.1)
MONTANA	1570	56.8	1.7	(53.5, 60.1)
NEBRASKA	1575	61.6	1.4	(58.9, 64.3)
NEVADA	1046	54.7	2.3	(50.2, 59.2)
NEW HAMPSHIRE	1608	48.0	1.4	(45.3, 50.7)
NEW JERSEY	2283	53.7	1.3	(51.2, 56.2)
NEW MEXICO	1530	55.4	1.5	(52.5, 58.3)
NEW YORK	1412	52.7	1.5	(49.8, 55.6)
NORTH CAROLINA	2494	52.4	1.4	(49.7, 55.1)
NORTH DAKOTA	949	49.5	1.8	(46.0, 53.0)
OHIO	1340	54.2	1.6	(51.1, 57.3)
OKLAHOMA	2040	61.3	1.3	(58.8, 63.8)
OREGON	1014	46.5	1.7	(43.2, 49.8)
PENNSYLVANIA	1617	52.3	1.4	(49.6, 55.0)
RHODE ISLAND	1624	47.9	1.4	(45.2, 50.6)
SOUTH CAROLINA	1208	53.3	1.6	(50.2, 56.4)
SOUTH DAKOTA	2328	54.9	1.1	(52.7, 57.1)
TENNESSEE	1182	59.9	1.6	(56.8, 63.0)
TEXAS	2184	57.1	1.2	(54.7, 59.5)
UTAH	1252	53.8	1.9	(50.1, 57.5)
VERMONT	1780	50.8	1.3	(48.3, 53.3)
VIRGINIA	1146	48.5	1.6	(45.4, 51.6)
WASHINGTON	1533	48.0	1.4	(45.3, 50.7)
WEST VIRGINIA	1469	61.0	1.4	(58.3, 63.7)
WISCONSIN	1263	42.7	1.6	(39.6, 45.8)
WYOMING	1227	53.7	1.5	(50.8, 56.6)
GUAM	193	65.4	3.9	(57.8, 73.0)
PUERTO RICO	1982	67.8	1.4	(65.1, 70.5)
VIRGIN ISLANDS	800	69.5	2.0	(65.6, 73.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 53.7 RANGE = 38.0-69.5

HP2010: LHI, HP 3.12b

*AT RISK NEVER HAD A SIGMOIDOSCOPY, AGE 50 AND OLDER

August 8,

TABLE 3.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD SIGMOIDOSCOPY: NEVER HAD A SIGMOIDOSCOPY* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	413	57.8	2.7	(52.5, 63.1)	*	733	53.4	2.0	(49.5, 57.3)
ALASKA	438	52.4	3.6	(45.3, 59.5)	*	476	58.2	3.5	(51.3, 65.1)
ARIZONA	573	50.3	2.9	(44.6, 56.0)	*	863	49.9	2.5	(45.0, 54.8)
ARKANSAS	465	59.3	2.5	(54.4, 64.2)	*	816	52.2	1.9	(48.5, 55.9)
CALIFORNIA	596	53.1	2.5	(48.2, 58.0)	*	990	50.2	1.9	(46.5, 53.9)
COLORADO	282	52.2	3.2	(45.9, 58.5)	*	411	53.4	2.7	(48.1, 58.7)
CONNECTICUT	1224	43.1	1.7	(39.8, 46.4)	*	1895	48.8	1.4	(46.1, 51.5)
DELAWARE	521	37.4	2.6	(32.3, 42.5)	*	957	44.9	2.0	(41.0, 48.8)
DISTRICT OF CO	267	47.6	3.5	(40.7, 54.5)	*	402	41.6	2.8	(36.1, 47.1)
FLORIDA	790	52.2	2.0	(48.3, 56.1)	*	1251	51.9	1.6	(48.8, 55.0)
GEORGIA	584	50.9	2.4	(46.2, 55.6)	*	1047	51.3	1.9	(47.6, 55.0)
HAWAII	757	48.2	2.4	(43.5, 52.9)	*	1077	51.9	2.2	(47.6, 56.2)
IDAHO	791	57.7	2.2	(53.4, 62.0)	*	1140	55.6	1.7	(52.3, 58.9)
ILLINOIS	315	60.0	3.1	(53.9, 66.1)	*	537	58.5	2.3	(54.0, 63.0)
INDIANA	594	54.8	2.2	(50.5, 59.1)	*	990	54.2	1.7	(50.9, 57.5)
IOWA	601	55.9	2.3	(51.4, 60.4)	*	1012	53.4	1.8	(49.9, 56.9)
KANSAS	648	57.9	2.1	(53.8, 62.0)	*	1075	53.7	1.7	(50.4, 57.0)
KENTUCKY	1213	59.6	2.1	(55.5, 63.7)	*	2133	61.6	1.5	(58.7, 64.5)
LOUISIANA	619	64.0	2.1	(59.9, 68.1)	*	1228	58.1	1.6	(55.0, 61.2)
MAINE	428	54.8	2.6	(49.7, 59.9)	*	640	50.3	2.1	(46.2, 54.4)
MARYLAND	701	44.1	2.3	(39.6, 48.6)	*	1018	50.3	2.0	(46.4, 54.2)
MASSACHUSETTS	1217	44.7	1.7	(41.4, 48.0)	*	1902	48.4	1.3	(45.9, 50.9)
MICHIGAN	616	43.0	2.1	(38.9, 47.1)	*	956	46.3	1.8	(42.8, 49.8)
MINNESOTA	675	38.1	2.0	(34.2, 42.0)	*	961	38.0	1.7	(34.7, 41.3)
MISSISSIPPI	425	65.6	2.6	(60.5, 70.7)	*	793	54.4	2.0	(50.5, 58.3)
MISSOURI	677	57.1	2.6	(52.0, 62.2)	*	1197	57.0	2.0	(53.1, 60.9)
MONTANA	635	56.9	2.6	(51.8, 62.0)	*	935	56.8	2.2	(52.5, 61.1)
NEBRASKA	560	62.0	2.2	(57.7, 66.3)	*	1015	61.3	1.7	(58.0, 64.6)
NEVADA	529	55.9	3.5	(49.0, 62.8)	*	517	53.5	3.0	(47.6, 59.4)
NEW HAMPSHIRE	657	44.0	2.1	(39.9, 48.1)	*	951	51.4	1.8	(47.9, 54.9)
NEW JERSEY	856	49.2	2.1	(45.1, 53.3)	*	1427	57.2	1.7	(53.9, 60.5)
NEW MEXICO	597	54.6	2.4	(49.9, 59.3)	*	933	56.2	1.9	(52.5, 59.9)
NEW YORK	586	52.0	2.4	(47.3, 56.7)	*	826	53.3	2.0	(49.4, 57.2)
NORTH CAROLINA	901	51.4	2.4	(46.7, 56.1)	*	1593	53.2	1.8	(49.7, 56.7)
NORTH DAKOTA	356	49.2	2.8	(43.7, 54.7)	*	593	49.7	2.2	(45.4, 54.0)
OHIO	495	53.3	2.5	(48.4, 58.2)	*	845	54.9	2.0	(51.0, 58.8)
OKLAHOMA	765	65.1	2.2	(60.8, 69.4)	*	1275	58.3	1.8	(54.8, 61.8)
OREGON	400	47.0	2.7	(41.7, 52.3)	*	614	46.1	2.2	(41.8, 50.4)
PENNSYLVANIA	608	49.2	2.2	(44.9, 53.5)	*	1009	54.7	1.7	(51.4, 58.0)
RHODE ISLAND	575	44.4	2.3	(39.9, 48.9)	*	1049	50.5	1.8	(47.0, 54.0)
SOUTH CAROLINA	460	53.0	2.5	(48.1, 57.9)	*	748	53.5	2.0	(49.6, 57.4)
SOUTH DAKOTA	948	54.4	1.8	(50.9, 57.9)	*	1380	55.3	1.5	(52.4, 58.2)
TENNESSEE	360	62.7	2.8	(57.2, 68.2)	*	822	57.6	1.9	(53.9, 61.3)
TEXAS	852	58.2	1.8	(54.7, 61.7)	*	1332	56.3	1.5	(53.4, 59.2)
UTAH	560	52.5	2.9	(46.8, 58.2)	*	692	54.9	2.5	(50.0, 59.8)
VERMONT	705	47.5	2.0	(43.6, 51.4)	*	1075	53.5	1.7	(50.2, 56.8)
VIRGINIA	481	48.6	2.5	(43.7, 53.5)	*	665	48.5	2.1	(44.4, 52.6)
WASHINGTON	611	48.0	2.2	(43.7, 52.3)	*	922	47.9	1.8	(44.4, 51.4)
WEST VIRGINIA	561	61.5	2.2	(57.2, 65.8)	*	908	60.7	1.8	(57.2, 64.2)
WISCONSIN	496	43.4	2.5	(38.5, 48.3)	*	767	42.2	2.0	(38.3, 46.1)
WYOMING	499	51.4	2.4	(46.7, 56.1)	*	728	55.6	2.0	(51.7, 59.5)
GUAM	92	63.8	5.9	(52.2, 75.4)	*	101	67.2	5.1	(57.2, 77.2)
PUERTO RICO	708	65.5	2.3	(61.0, 70.0)	*	1274	69.7	1.7	(66.4, 73.0)
VIRGIN ISLANDS	330	71.2	3.0	(65.3, 77.1)	*	470	67.9	2.8	(62.4, 73.4)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
MEDIAN = 52.8
RANGE = 37.4-71.2

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
MEDIAN = 53.5
RANGE = 38.0-69.7

*AT RISK NEVER HAD A SIGMOIDOSCOPY, AGE 50 AND OLDER

August 8,

TABLE 3.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2001
 SIGMOIDOSCOPY: NEVER HAD A SIGMOIDOSCOPY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	58.8	55.3	-3.5	-6.0
ALASKA	49.6	55.3	5.7	11.5
ARIZONA	63.2	50.0	-13.2	-20.8
ARKANSAS	57.9	55.4	-2.5	-4.4
CALIFORNIA	51.8	51.5	-0.3	-0.6
COLORADO	55.8	52.9	-2.9	-5.2
CONNECTICUT	51.3	46.3	-5.0	-9.7
DELAWARE	44.0	41.6	-2.4	-5.4
DISTRICT OF CO	51.6	44.0	-7.6	-14.8
FLORIDA	54.9	52.0	-2.9	-5.2
GEORGIA	52.6	51.1	-1.5	-2.8
HAWAII	55.8	50.2	-5.6	-10.0
IDAHO	58.7	56.6	-2.1	-3.5
ILLINOIS	56.1	59.1	3.0	5.3
INDIANA	56.6	54.5	-2.1	-3.7
IOWA	53.6	54.5	0.9	1.7
KANSAS	61.2	55.6	-5.6	-9.2
KENTUCKY	65.3	60.7	-4.6	-7.0
LOUISIANA	63.3	60.6	-2.7	-4.3
MAINE	57.6	52.3	-5.3	-9.2
MARYLAND	49.5	47.6	-1.9	-3.8
MASSACHUSETTS	56.1	46.8	-9.3	-16.5
MICHIGAN	50.5	44.8	-5.7	-11.4
MINNESOTA	52.5	38.0	-14.5	-27.6
MISSISSIPPI	62.6	59.2	-3.4	-5.4
MISSOURI	62.0	57.0	-5.0	-8.0
MONTANA	56.8	56.8	0.0	0.0
NEBRASKA	66.3	61.6	-4.7	-7.1
NEVADA	56.9	54.7	-2.2	-3.9
NEW HAMPSHIRE	53.2	48.0	-5.2	-9.7
NEW JERSEY	56.0	53.7	-2.3	-4.1
NEW MEXICO	56.9	55.4	-1.5	-2.6
NEW YORK	57.8	52.7	-5.1	-8.8
NORTH CAROLINA	60.9	52.4	-8.5	-14.0
NORTH DAKOTA	58.0	49.5	-8.5	-14.6
OHIO	59.3	54.2	-5.1	-8.5
OKLAHOMA	62.3	61.3	-1.0	-1.5
OREGON	52.9	46.5	-6.4	-12.1
PENNSYLVANIA	60.3	52.3	-8.0	-13.3
RHODE ISLAND	51.3	47.9	-3.4	-6.7
SOUTH CAROLINA	58.2	53.3	-4.9	-8.5
SOUTH DAKOTA	54.5	54.9	0.4	0.7
TENNESSEE	60.7	59.9	-0.8	-1.3
TEXAS	55.7	57.1	1.4	2.5
UTAH	55.9	53.8	-2.1	-3.7
VERMONT	57.8	50.8	-7.0	-12.0
VIRGINIA	55.8	48.5	-7.3	-13.1
WASHINGTON	49.2	48.0	-1.2	-2.5
WEST VIRGINIA	65.1	61.0	-4.1	-6.3
WISCONSIN	47.6	42.7	-4.9	-10.3
WYOMING	53.5	53.7	0.2	0.4
GUAM	N/A	65.4	N/A	N/A
PUERTO RICO	71.8	67.8	-4.0	-5.6
VIRGIN ISLANDS	N/A	69.5	N/A	N/A

SUMMARY STATISTICS: yr1999

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 56.4
 RANGE = 44.0-71.8

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 53.7
 RANGE = 38.0-69.5

*AT RISK NEVER HAD A SIGMOIDOSCOPY, AGE 50 AND OLDER
 N/A - NOT AVAILABLE

August 8, 2002

TABLE 4.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES: TOLD THAT YOU HAVE DIABETES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2795	9.6	0.6	(8.4, 10.8)
ALASKA	2871	4.0	0.5	(3.0, 5.0)
ARIZONA	3253	6.1	0.5	(5.1, 7.1)
ARKANSAS	2925	7.8	0.6	(6.6, 9.0)
CALIFORNIA	4256	6.5	0.5	(5.5, 7.5)
COLORADO	2031	4.6	0.5	(3.6, 5.6)
CONNECTICUT	7736	6.3	0.3	(5.7, 6.9)
DELAWARE	3512	7.1	0.5	(6.1, 8.1)
DISTRICT OF CO	1885	8.3	0.8	(6.7, 9.9)
FLORIDA	4674	8.2	0.5	(7.2, 9.2)
GEORGIA	4525	6.9	0.4	(6.1, 7.7)
HAWAII	4500	6.2	0.6	(5.0, 7.4)
IDAHO	4830	5.4	0.4	(4.6, 6.2)
ILLINOIS	4007	6.6	0.4	(5.8, 7.4)
INDIANA	3989	6.5	0.4	(5.7, 7.3)
IOWA	3632	5.7	0.4	(4.9, 6.5)
KANSAS	4591	5.8	0.4	(5.0, 6.6)
KENTUCKY	7512	6.7	0.4	(5.9, 7.5)
LOUISIANA	4997	7.6	0.4	(6.8, 8.4)
MAINE	2417	6.7	0.6	(5.5, 7.9)
MARYLAND	4466	6.9	0.5	(5.9, 7.9)
MASSACHUSETTS	8619	5.6	0.3	(5.0, 6.2)
MICHIGAN	3824	7.2	0.5	(6.2, 8.2)
MINNESOTA	3960	4.4	0.3	(3.8, 5.0)
MISSISSIPPI	3037	9.3	0.6	(8.1, 10.5)
MISSOURI	4177	6.6	0.5	(5.6, 7.6)
MONTANA	3335	5.6	0.5	(4.6, 6.6)
NEBRASKA	3693	5.2	0.4	(4.4, 6.0)
NEVADA	2568	5.7	0.7	(4.3, 7.1)
NEW HAMPSHIRE	4062	5.4	0.4	(4.6, 6.2)
NEW JERSEY	6011	7.1	0.5	(6.1, 8.1)
NEW MEXICO	3619	6.2	0.5	(5.2, 7.2)
NEW YORK	3892	6.6	0.5	(5.6, 7.6)
NORTH CAROLINA	6199	6.7	0.5	(5.7, 7.7)
NORTH DAKOTA	2508	5.1	0.5	(4.1, 6.1)
OHIO	3427	7.2	0.5	(6.2, 8.2)
OKLAHOMA	4543	7.7	0.5	(6.7, 8.7)
OREGON	2533	5.7	0.5	(4.7, 6.7)
PENNSYLVANIA	3659	6.7	0.4	(5.9, 7.5)
RHODE ISLAND	4113	6.4	0.5	(5.4, 7.4)
SOUTH CAROLINA	3199	8.1	0.5	(7.1, 9.1)
SOUTH DAKOTA	5117	6.1	0.4	(5.3, 6.9)
TENNESSEE	2922	7.7	0.6	(6.5, 8.9)
TEXAS	5910	7.1	0.4	(6.3, 7.9)
UTAH	3654	4.3	0.4	(3.5, 5.1)
VERMONT	4298	5.1	0.4	(4.3, 5.9)
VIRGINIA	2934	6.0	0.5	(5.0, 7.0)
WASHINGTON	4205	5.7	0.4	(4.9, 6.5)
WEST VIRGINIA	3091	8.8	0.6	(7.6, 10.0)
WISCONSIN	3349	5.6	0.4	(4.8, 6.4)
WYOMING	3036	4.5	0.4	(3.7, 5.3)
GUAM	870	9.5	1.3	(7.0, 12.0)
PUERTO RICO	4234	9.8	0.6	(8.6, 11.0)
VIRGIN ISLANDS	2260	7.2	0.6	(6.0, 8.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 6.6 RANGE = 4.0-9.8

* TOLD THAT YOU HAVE DIABETES, AGE 18 AND OLDER

August 8, 2002

TABLE 4.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES: TOLD THAT YOU HAVE DIABETES* -BY SEX

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1050	10.5	1.0	(8.5, 12.5)	*	1745	8.8	0.7	(7.4, 10.2)
ALASKA	1356	4.0	0.7	(2.6, 5.4)	*	1515	4.0	0.7	(2.6, 5.4)
ARIZONA	1338	7.3	0.9	(5.5, 9.1)	*	1915	4.9	0.6	(3.7, 6.1)
ARKANSAS	1120	8.4	0.9	(6.6, 10.2)	*	1805	7.2	0.7	(5.8, 8.6)
CALIFORNIA	1723	5.8	0.6	(4.6, 7.0)	*	2533	7.1	0.7	(5.7, 8.5)
COLORADO	864	5.6	0.9	(3.8, 7.4)	*	1167	3.7	0.6	(2.5, 4.9)
CONNECTICUT	3198	6.5	0.5	(5.5, 7.5)	*	4538	6.0	0.4	(5.2, 6.8)
DELAWARE	1360	7.5	0.9	(5.7, 9.3)	*	2152	6.7	0.7	(5.3, 8.1)
DISTRICT OF CO	780	9.6	1.3	(7.1, 12.1)	*	1105	7.2	0.9	(5.4, 9.0)
FLORIDA	1910	9.1	0.8	(7.5, 10.7)	*	2764	7.3	0.6	(6.1, 8.5)
GEORGIA	1722	6.5	0.7	(5.1, 7.9)	*	2803	7.2	0.6	(6.0, 8.4)
HAWAII	1937	6.8	1.0	(4.8, 8.8)	*	2563	5.6	0.6	(4.4, 6.8)
IDAHO	2068	5.2	0.5	(4.2, 6.2)	*	2762	5.7	0.5	(4.7, 6.7)
ILLINOIS	1643	6.7	0.7	(5.3, 8.1)	*	2364	6.6	0.6	(5.4, 7.8)
INDIANA	1613	6.3	0.6	(5.1, 7.5)	*	2376	6.8	0.5	(5.8, 7.8)
IOWA	1483	5.8	0.6	(4.6, 7.0)	*	2149	5.6	0.5	(4.6, 6.6)
KANSAS	1854	5.5	0.5	(4.5, 6.5)	*	2737	6.1	0.5	(5.1, 7.1)
KENTUCKY	2916	6.9	0.6	(5.7, 8.1)	*	4596	6.4	0.4	(5.6, 7.2)
LOUISIANA	1886	6.9	0.6	(5.7, 8.1)	*	3111	8.2	0.5	(7.2, 9.2)
MAINE	1028	7.0	1.0	(5.0, 9.0)	*	1389	6.5	0.7	(5.1, 7.9)
MARYLAND	1822	7.1	0.8	(5.5, 8.7)	*	2644	6.6	0.7	(5.2, 8.0)
MASSACHUSETTS	3556	6.0	0.5	(5.0, 7.0)	*	5063	5.2	0.4	(4.4, 6.0)
MICHIGAN	1516	6.8	0.7	(5.4, 8.2)	*	2308	7.6	0.6	(6.4, 8.8)
MINNESOTA	1697	4.4	0.5	(3.4, 5.4)	*	2263	4.3	0.5	(3.3, 5.3)
MISSISSIPPI	1144	8.8	1.0	(6.8, 10.8)	*	1893	9.7	0.7	(8.3, 11.1)
MISSOURI	1652	7.0	0.8	(5.4, 8.6)	*	2525	6.2	0.6	(5.0, 7.4)
MONTANA	1422	4.9	0.6	(3.7, 6.1)	*	1913	6.2	0.8	(4.6, 7.8)
NEBRASKA	1422	4.9	0.6	(3.7, 6.1)	*	2271	5.5	0.5	(4.5, 6.5)
NEVADA	1308	5.2	0.7	(3.8, 6.6)	*	1260	6.2	1.1	(4.0, 8.4)
NEW HAMPSHIRE	1725	5.8	0.6	(4.6, 7.0)	*	2337	5.0	0.5	(4.0, 6.0)
NEW JERSEY	2443	7.3	0.7	(5.9, 8.7)	*	3568	7.0	0.6	(5.8, 8.2)
NEW MEXICO	1501	5.9	0.7	(4.5, 7.3)	*	2118	6.5	0.6	(5.3, 7.7)
NEW YORK	1610	6.4	0.7	(5.0, 7.8)	*	2282	6.8	0.7	(5.4, 8.2)
NORTH CAROLINA	2380	6.8	0.7	(5.4, 8.2)	*	3819	6.7	0.5	(5.7, 7.7)
NORTH DAKOTA	1066	4.7	0.7	(3.3, 6.1)	*	1442	5.6	0.6	(4.4, 6.8)
OHIO	1349	7.5	0.8	(5.9, 9.1)	*	2078	6.9	0.6	(5.7, 8.1)
OKLAHOMA	1769	8.3	0.8	(6.7, 9.9)	*	2774	7.2	0.6	(6.0, 8.4)
OREGON	1052	5.5	0.7	(4.1, 6.9)	*	1481	5.8	0.7	(4.4, 7.2)
PENNSYLVANIA	1481	6.6	0.7	(5.2, 8.0)	*	2178	6.7	0.6	(5.5, 7.9)
RHODE ISLAND	1546	7.4	0.7	(6.0, 8.8)	*	2567	5.6	0.6	(4.4, 6.8)
SOUTH CAROLINA	1267	8.4	0.9	(6.6, 10.2)	*	1932	7.7	0.7	(6.3, 9.1)
SOUTH DAKOTA	2165	6.6	0.6	(5.4, 7.8)	*	2952	5.6	0.4	(4.8, 6.4)
TENNESSEE	953	7.6	1.0	(5.6, 9.6)	*	1969	7.9	0.7	(6.5, 9.3)
TEXAS	2435	7.0	0.6	(5.8, 8.2)	*	3475	7.2	0.5	(6.2, 8.2)
UTAH	1652	4.3	0.7	(2.9, 5.7)	*	2002	4.2	0.6	(3.0, 5.4)
VERMONT	1789	4.6	0.5	(3.6, 5.6)	*	2509	5.5	0.5	(4.5, 6.5)
VIRGINIA	1237	6.2	0.7	(4.8, 7.6)	*	1697	5.8	0.7	(4.4, 7.2)
WASHINGTON	1756	6.2	0.6	(5.0, 7.4)	*	2449	5.2	0.5	(4.2, 6.2)
WEST VIRGINIA	1228	8.9	0.8	(7.3, 10.5)	*	1863	8.8	0.8	(7.2, 10.4)
WISCONSIN	1479	5.9	0.7	(4.5, 7.3)	*	1870	5.3	0.6	(4.1, 6.5)
WYOMING	1318	4.1	0.6	(2.9, 5.3)	*	1718	4.8	0.5	(3.8, 5.8)
GUAM	382	9.9	2.0	(6.0, 13.8)	*	488	9.1	1.6	(6.0, 12.2)
PUERTO RICO	1519	9.1	0.9	(7.3, 10.9)	*	2715	10.5	0.8	(8.9, 12.1)
VIRGIN ISLANDS	874	4.3	0.8	(2.7, 5.9)	*	1386	9.5	1.0	(7.5, 11.5)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 6.6
 RANGE = 4.0-10.5

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 6.5
 RANGE = 3.7-10.5

* TOLD THAT YOUR HAVE DIABETES, AGE 18 AND OLDER

August 8,

TABLE 4.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 DIABETES: TOLD THAT YOU HAVE DIABETES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	7.4	9.6	2.2	30.1
ALASKA	3.8	4.0	0.2	5.3
ARIZONA	5.9	6.1	0.2	2.9
ARKANSAS	6.2	7.8	1.6	25.2
CALIFORNIA	6.8	6.5	-0.3	-4.0
COLORADO	5.1	4.6	-0.5	-10.0
CONNECTICUT	5.5	6.3	0.8	15.4
DELAWARE	6.4	7.1	0.7	10.8
DISTRICT OF CO	7.2	8.3	1.1	15.1
FLORIDA	6.9	8.2	1.3	18.3
GEORGIA	6.8	6.9	0.1	1.5
HAWAII	5.2	6.2	1.0	18.8
IDAHO	4.9	5.4	0.5	10.7
ILLINOIS	6.2	6.6	0.4	6.5
INDIANA	6.0	6.5	0.5	7.6
IOWA	6.1	5.7	-0.4	-6.9
KANSAS	5.9	5.8	-0.1	-1.4
KENTUCKY	6.5	6.7	0.2	3.1
LOUISIANA	6.7	7.6	0.9	14.3
MAINE	6.0	6.7	0.7	12.0
MARYLAND	6.5	6.9	0.5	7.0
MASSACHUSETTS	5.8	5.6	-0.2	-3.1
MICHIGAN	7.0	7.2	0.2	2.4
MINNESOTA	4.9	4.4	-0.5	-10.2
MISSISSIPPI	7.6	9.3	1.7	22.7
MISSOURI	6.7	6.6	-0.1	-1.0
MONTANA	4.9	5.6	0.8	15.5
NEBRASKA	4.9	5.2	0.3	5.3
NEVADA	6.8	5.7	-1.1	-15.7
NEW HAMPSHIRE	4.4	5.4	1.0	22.4
NEW JERSEY	5.8	7.1	1.3	22.0
NEW MEXICO	6.5	6.2	-0.3	-4.8
NEW YORK	6.3	6.6	0.3	4.6
NORTH CAROLINA	6.4	6.7	0.3	4.2
NORTH DAKOTA	5.2	5.1	-0.1	-2.7
OHIO	6.4	7.2	0.9	13.4
OKLAHOMA	5.5	7.7	2.2	41.0
OREGON	6.0	5.7	-0.3	-5.6
PENNSYLVANIA	7.1	6.7	-0.4	-6.0
RHODE ISLAND	6.0	6.4	0.4	6.7
SOUTH CAROLINA	7.1	8.1	1.0	13.8
SOUTH DAKOTA	5.7	6.1	0.4	7.0
TENNESSEE	7.2	7.7	0.5	6.6
TEXAS	6.2	7.1	0.9	14.7
UTAH	5.4	4.3	-1.1	-19.8
VERMONT	4.4	5.1	0.7	15.4
VIRGINIA	6.2	6.0	-0.2	-2.4
WASHINGTON	5.5	5.7	0.2	3.3
WEST VIRGINIA	7.6	8.8	1.2	15.9
WISCONSIN	6.1	5.6	-0.5	-8.3
WYOMING	5.0	4.5	-0.5	-10.0
GUAM	N/A	9.5	N/A	N/A
PUERTO RICO	8.6	9.8	1.3	14.6
VIRGIN ISLANDS	N/A	7.2	N/A	N/A

SUMMARY STATISTICS: yr2000

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 6.1
 RANGE = 3.8-8.6

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 6.6
 RANGE = 4.0-9.8

* TOLD THAT YOU HAVE DIABETES, AGE 18 AND OLDER

N/A - NOT AVAILABLE

August 8, 2002

TABLE 5.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES: TOLD THAT YOU HAVE DIABETES, AGE 20+*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2717	10.0	0.6	(8.8, 11.2)
ALASKA	2749	4.2	0.5	(3.2, 5.2)
ARIZONA	3100	6.4	0.6	(5.2, 7.6)
ARKANSAS	2844	8.0	0.6	(6.8, 9.2)
CALIFORNIA	4145	6.7	0.5	(5.7, 7.7)
COLORADO	1962	4.8	0.6	(3.6, 6.0)
CONNECTICUT	7468	6.5	0.3	(5.9, 7.1)
DELAWARE	3406	7.3	0.6	(6.1, 8.5)
DISTRICT OF CO	1817	8.6	0.8	(7.0, 10.2)
FLORIDA	4493	8.4	0.5	(7.4, 9.4)
GEORGIA	4396	7.1	0.5	(6.1, 8.1)
HAWAII	4376	6.3	0.6	(5.1, 7.5)
IDAHO	4676	5.6	0.4	(4.8, 6.4)
ILLINOIS	3902	6.9	0.5	(5.9, 7.9)
INDIANA	3870	6.8	0.4	(6.0, 7.6)
IOWA	3516	5.9	0.4	(5.1, 6.7)
KANSAS	4462	5.9	0.4	(5.1, 6.7)
KENTUCKY	7326	7.0	0.4	(6.2, 7.8)
LOUISIANA	4774	8.0	0.4	(7.2, 8.8)
MAINE	2353	6.9	0.6	(5.7, 8.1)
MARYLAND	4306	7.1	0.5	(6.1, 8.1)
MASSACHUSETTS	8339	5.8	0.3	(5.2, 6.4)
MICHIGAN	3702	7.5	0.5	(6.5, 8.5)
MINNESOTA	3881	4.5	0.4	(3.7, 5.3)
MISSISSIPPI	2946	9.5	0.6	(8.3, 10.7)
MISSOURI	4046	6.7	0.5	(5.7, 7.7)
MONTANA	3275	5.8	0.5	(4.8, 6.8)
NEBRASKA	3570	5.5	0.4	(4.7, 6.3)
NEVADA	2493	5.9	0.7	(4.5, 7.3)
NEW HAMPSHIRE	3924	5.6	0.4	(4.8, 6.4)
NEW JERSEY	5826	7.3	0.5	(6.3, 8.3)
NEW MEXICO	3512	6.4	0.5	(5.4, 7.4)
NEW YORK	3756	6.8	0.5	(5.8, 7.8)
NORTH CAROLINA	6025	6.9	0.5	(5.9, 7.9)
NORTH DAKOTA	2433	5.2	0.5	(4.2, 6.2)
OHIO	3305	7.4	0.5	(6.4, 8.4)
OKLAHOMA	4422	8.0	0.5	(7.0, 9.0)
OREGON	2462	5.9	0.5	(4.9, 6.9)
PENNSYLVANIA	3557	6.9	0.5	(5.9, 7.9)
RHODE ISLAND	3985	6.7	0.5	(5.7, 7.7)
SOUTH CAROLINA	3091	8.4	0.6	(7.2, 9.6)
SOUTH DAKOTA	5004	6.3	0.4	(5.5, 7.1)
TENNESSEE	2843	8.0	0.6	(6.8, 9.2)
TEXAS	5744	7.3	0.4	(6.5, 8.1)
UTAH	3497	4.4	0.4	(3.6, 5.2)
VERMONT	4193	5.3	0.4	(4.5, 6.1)
VIRGINIA	2826	6.2	0.5	(5.2, 7.2)
WASHINGTON	4072	5.9	0.4	(5.1, 6.7)
WEST VIRGINIA	3014	9.2	0.6	(8.0, 10.4)
WISCONSIN	3261	5.8	0.5	(4.8, 6.8)
WYOMING	2944	4.6	0.4	(3.8, 5.4)
GUAM	834	10.0	1.3	(7.5, 12.5)
PUERTO RICO	4130	10.4	0.6	(9.2, 11.6)
VIRGIN ISLANDS	2140	7.7	0.7	(6.3, 9.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 6.8 RANGE = 4.2-10.4

HP2010: HP 5.4(approx.)

* TOLD THAT YOU HAVE DIABETES, AGE 20 AND OLDER

August 8,

TABLE 5.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES: TOLD THAT YOU HAVE DIABETES, AGE 20+* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1021	10.9	1.0	(8.9, 12.9)	*	1696	9.1	0.8 (7.5, 10.7)
ALASKA	1292	4.3	0.7	(2.9, 5.7)	*	1457	4.1	0.7 (2.7, 5.5)
ARIZONA	1274	7.7	1.0	(5.7, 9.7)	*	1826	5.1	0.6 (3.9, 6.3)
ARKANSAS	1091	8.8	0.9	(7.0, 10.6)	*	1753	7.4	0.8 (5.8, 9.0)
CALIFORNIA	1670	6.1	0.7	(4.7, 7.5)	*	2475	7.3	0.7 (5.9, 8.7)
COLORADO	836	5.9	0.9	(4.1, 7.7)	*	1126	3.8	0.7 (2.4, 5.2)
CONNECTICUT	3102	6.8	0.5	(5.8, 7.8)	*	4366	6.2	0.4 (5.4, 7.0)
DELAWARE	1321	7.8	0.9	(6.0, 9.6)	*	2085	6.8	0.7 (5.4, 8.2)
DISTRICT OF CO	769	9.9	1.3	(7.4, 12.4)	*	1048	7.4	0.9 (5.6, 9.2)
FLORIDA	1835	9.5	0.8	(7.9, 11.1)	*	2658	7.3	0.6 (6.1, 8.5)
GEORGIA	1675	6.8	0.7	(5.4, 8.2)	*	2721	7.5	0.6 (6.3, 8.7)
HAWAII	1878	6.9	1.0	(4.9, 8.9)	*	2498	5.7	0.6 (4.5, 6.9)
IDAHO	2000	5.4	0.6	(4.2, 6.6)	*	2676	5.9	0.5 (4.9, 6.9)
ILLINOIS	1597	6.9	0.7	(5.5, 8.3)	*	2305	6.9	0.6 (5.7, 8.1)
INDIANA	1565	6.6	0.7	(5.2, 8.0)	*	2305	6.9	0.6 (5.7, 8.1)
IOWA	1436	6.0	0.7	(4.6, 7.4)	*	2080	5.8	0.5 (4.8, 6.8)
KANSAS	1811	5.7	0.6	(4.5, 6.9)	*	2651	6.2	0.5 (5.2, 7.2)
KENTUCKY	2835	7.3	0.6	(6.1, 8.5)	*	4491	6.6	0.5 (5.6, 7.6)
LOUISIANA	1795	7.3	0.7	(5.9, 8.7)	*	2979	8.5	0.6 (7.3, 9.7)
MAINE	995	7.3	1.1	(5.1, 9.5)	*	1358	6.6	0.7 (5.2, 8.0)
MARYLAND	1760	7.4	0.8	(5.8, 9.0)	*	2546	6.9	0.7 (5.5, 8.3)
MASSACHUSETTS	3441	6.2	0.5	(5.2, 7.2)	*	4898	5.4	0.4 (4.6, 6.2)
MICHIGAN	1465	7.1	0.7	(5.7, 8.5)	*	2237	7.9	0.6 (6.7, 9.1)
MINNESOTA	1664	4.6	0.5	(3.6, 5.6)	*	2217	4.5	0.5 (3.5, 5.5)
MISSISSIPPI	1106	9.1	1.0	(7.1, 11.1)	*	1840	9.9	0.7 (8.5, 11.3)
MISSOURI	1595	7.1	0.9	(5.3, 8.9)	*	2451	6.4	0.6 (5.2, 7.6)
MONTANA	1390	5.2	0.7	(3.8, 6.6)	*	1885	6.3	0.8 (4.7, 7.9)
NEBRASKA	1364	5.2	0.6	(4.0, 6.4)	*	2206	5.7	0.5 (4.7, 6.7)
NEVADA	1281	5.3	0.8	(3.7, 6.9)	*	1212	6.5	1.1 (4.3, 8.7)
NEW HAMPSHIRE	1666	6.0	0.6	(4.8, 7.2)	*	2258	5.2	0.5 (4.2, 6.2)
NEW JERSEY	2368	7.3	0.7	(5.9, 8.7)	*	3458	7.3	0.6 (6.1, 8.5)
NEW MEXICO	1453	6.1	0.7	(4.7, 7.5)	*	2059	6.7	0.6 (5.5, 7.9)
NEW YORK	1557	6.6	0.7	(5.2, 8.0)	*	2199	7.0	0.7 (5.6, 8.4)
NORTH CAROLINA	2322	7.1	0.8	(5.5, 8.7)	*	3703	6.8	0.6 (5.6, 8.0)
NORTH DAKOTA	1038	4.8	0.7	(3.4, 6.2)	*	1395	5.6	0.6 (4.4, 6.8)
OHIO	1299	7.9	0.9	(6.1, 9.7)	*	2006	7.0	0.6 (5.8, 8.2)
OKLAHOMA	1725	8.6	0.8	(7.0, 10.2)	*	2697	7.5	0.7 (6.1, 8.9)
OREGON	1020	5.8	0.8	(4.2, 7.4)	*	1442	5.9	0.7 (4.5, 7.3)
PENNSYLVANIA	1443	6.8	0.7	(5.4, 8.2)	*	2114	7.0	0.6 (5.8, 8.2)
RHODE ISLAND	1499	7.7	0.8	(6.1, 9.3)	*	2486	5.9	0.6 (4.7, 7.1)
SOUTH CAROLINA	1224	8.6	0.9	(6.8, 10.4)	*	1867	8.1	0.7 (6.7, 9.5)
SOUTH DAKOTA	2120	6.8	0.6	(5.6, 8.0)	*	2884	5.7	0.4 (4.9, 6.5)
TENNESSEE	923	8.0	1.1	(5.8, 10.2)	*	1920	8.1	0.7 (6.7, 9.5)
TEXAS	2362	7.3	0.6	(6.1, 8.5)	*	3382	7.4	0.5 (6.4, 8.4)
UTAH	1582	4.5	0.7	(3.1, 5.9)	*	1915	4.3	0.6 (3.1, 5.5)
VERMONT	1752	4.8	0.5	(3.8, 5.8)	*	2441	5.7	0.5 (4.7, 6.7)
VIRGINIA	1200	6.5	0.7	(5.1, 7.9)	*	1626	6.0	0.7 (4.6, 7.4)
WASHINGTON	1700	6.4	0.6	(5.2, 7.6)	*	2372	5.5	0.5 (4.5, 6.5)
WEST VIRGINIA	1198	9.2	0.9	(7.4, 11.0)	*	1816	9.2	0.8 (7.6, 10.8)
WISCONSIN	1429	6.2	0.7	(4.8, 7.6)	*	1832	5.4	0.6 (4.2, 6.6)
WYOMING	1275	4.3	0.6	(3.1, 5.5)	*	1669	4.9	0.5 (3.9, 5.9)
GUAM	368	10.2	2.0	(6.3, 14.1)	*	466	9.7	1.7 (6.4, 13.0)
PUERTO RICO	1471	9.6	0.9	(7.8, 11.4)	*	2659	11.0	0.8 (9.4, 12.6)
VIRGIN ISLANDS	826	4.6	0.8	(3.0, 6.2)	*	1314	10.2	1.0 (8.2, 12.2)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 6.8
 RANGE = 4.3-10.9

* TOLD THAT YOU HAVE DIABETES, AGE 20 AND OLDER

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 6.7
 RANGE = 3.8-11.0

August 8,

TABLE 5.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 DIABETES: TOLD THAT YOU HAVE DIABETES, AGE 20+*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	7.7	10.0	2.3	29.7
ALASKA	3.9	4.2	0.3	7.7
ARIZONA	6.2	6.4	0.2	2.6
ARKANSAS	6.4	8.0	1.6	25.6
CALIFORNIA	6.9	6.7	-0.2	-3.0
COLORADO	5.2	4.8	-0.4	-6.8
CONNECTICUT	5.6	6.5	0.9	15.7
DELAWARE	6.6	7.3	0.7	10.3
DISTRICT OF CO	7.4	8.6	1.2	16.8
FLORIDA	7.2	8.4	1.3	17.5
GEORGIA	7.1	7.1	0.0	0.6
HAWAII	5.4	6.3	0.9	16.2
IDAHO	5.1	5.6	0.5	10.2
ILLINOIS	6.5	6.9	0.5	7.0
INDIANA	6.3	6.8	0.5	7.6
IOWA	6.4	5.9	-0.4	-7.1
KANSAS	6.1	5.9	-0.2	-2.6
KENTUCKY	6.7	7.0	0.3	4.8
LOUISIANA	7.0	8.0	1.0	14.4
MAINE	6.2	6.9	0.7	11.7
MARYLAND	6.7	7.1	0.4	5.8
MASSACHUSETTS	6.0	5.8	-0.2	-2.7
MICHIGAN	7.4	7.5	0.1	1.8
MINNESOTA	5.1	4.5	-0.5	-10.9
MISSISSIPPI	7.9	9.5	1.6	20.3
MISSOURI	6.9	6.7	-0.2	-2.5
MONTANA	5.0	5.8	0.8	15.1
NEBRASKA	5.2	5.5	0.3	6.4
NEVADA	6.9	5.9	-1.0	-15.0
NEW HAMPSHIRE	4.6	5.6	1.0	21.7
NEW JERSEY	6.0	7.3	1.3	21.5
NEW MEXICO	6.7	6.4	-0.3	-5.0
NEW YORK	6.5	6.8	0.3	4.0
NORTH CAROLINA	6.6	6.9	0.3	4.5
NORTH DAKOTA	5.5	5.2	-0.3	-5.1
OHIO	6.6	7.4	0.8	12.8
OKLAHOMA	5.7	8.0	2.3	41.3
OREGON	6.2	5.9	-0.3	-4.7
PENNSYLVANIA	7.3	6.9	-0.4	-5.9
RHODE ISLAND	6.2	6.7	0.5	7.7
SOUTH CAROLINA	7.3	8.4	1.1	14.8
SOUTH DAKOTA	5.8	6.3	0.5	9.4
TENNESSEE	7.4	8.0	0.6	7.5
TEXAS	6.4	7.3	0.9	14.4
UTAH	5.6	4.4	-1.2	-21.0
VERMONT	4.6	5.3	0.7	15.2
VIRGINIA	6.4	6.2	-0.2	-3.6
WASHINGTON	5.7	5.9	0.2	3.5
WEST VIRGINIA	7.9	9.2	1.3	16.6
WISCONSIN	6.3	5.8	-0.5	-7.3
WYOMING	5.1	4.6	-0.5	-10.0
GUAM	N/A	10.0	N/A	N/A
PUERTO RICO	9.1	10.4	1.3	14.3
VIRGIN ISLANDS	N/A	7.7	N/A	N/A

SUMMARY STATISTICS: yr2000

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 6.4
 RANGE = 3.9-9.1

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 6.8
 RANGE = 4.2-10.4

* TOLD YOU HAVE DIABETES, AGE 20 AND OLDER

N/A - NOT AVAILABLE

August 8, 2002

TABLE 6.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BLOOD CHOLESTEROL: NO CHOLESTEROL CHECK IN THE PAST FIVE YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2725	26.6	1.0	(24.6, 28.6)
ALASKA	2768	31.9	1.4	(29.2, 34.6)
ARIZONA	3186	27.5	1.2	(25.1, 29.9)
ARKANSAS	2825	29.3	1.0	(27.3, 31.3)
CALIFORNIA	4175	29.0	0.9	(27.2, 30.8)
COLORADO	1988	28.4	1.2	(26.0, 30.8)
CONNECTICUT	7519	20.5	0.6	(19.3, 21.7)
DELAWARE	3441	21.3	0.9	(19.5, 23.1)
DISTRICT OF CO	1843	19.9	1.2	(17.5, 22.3)
FLORIDA	4567	22.0	0.8	(20.4, 23.6)
GEORGIA	4354	25.5	0.9	(23.7, 27.3)
HAWAII	4409	24.3	0.9	(22.5, 26.1)
IDAHO	4674	34.1	0.9	(32.3, 35.9)
ILLINOIS	2118	29.4	1.2	(27.0, 31.8)
INDIANA	3867	27.6	0.8	(26.0, 29.2)
IOWA	3514	28.8	0.9	(27.0, 30.6)
KANSAS	4425	27.8	0.8	(26.2, 29.4)
KENTUCKY	7240	27.7	0.8	(26.1, 29.3)
LOUISIANA	4841	27.3	0.7	(25.9, 28.7)
MAINE	2343	23.1	1.0	(21.1, 25.1)
MARYLAND	4350	20.1	0.8	(18.5, 21.7)
MASSACHUSETTS	8354	18.6	0.5	(17.6, 19.6)
MICHIGAN	3712	25.7	0.9	(23.9, 27.5)
MINNESOTA	3813	23.8	0.8	(22.2, 25.4)
MISSISSIPPI	2897	29.8	1.0	(27.8, 31.8)
MISSOURI	4038	27.6	1.0	(25.6, 29.6)
MONTANA	3257	30.1	1.1	(27.9, 32.3)
NEBRASKA	3598	34.6	0.9	(32.8, 36.4)
NEVADA	2505	27.6	1.3	(25.1, 30.1)
NEW HAMPSHIRE	3946	22.8	0.8	(21.2, 24.4)
NEW JERSEY	5864	20.5	0.7	(19.1, 21.9)
NEW MEXICO	3540	31.8	0.9	(30.0, 33.6)
NEW YORK	3792	22.7	0.9	(20.9, 24.5)
NORTH CAROLINA	6049	24.9	0.9	(23.1, 26.7)
NORTH DAKOTA	2441	28.8	1.0	(26.8, 30.8)
OHIO	3329	28.1	1.0	(26.1, 30.1)
OKLAHOMA	4367	28.6	0.9	(26.8, 30.4)
OREGON	2443	29.1	1.0	(27.1, 31.1)
PENNSYLVANIA	3561	24.9	0.9	(23.1, 26.7)
RHODE ISLAND	4023	19.3	0.8	(17.7, 20.9)
SOUTH CAROLINA	3119	22.1	0.9	(20.3, 23.9)
SOUTH DAKOTA	4940	30.5	0.8	(28.9, 32.1)
TENNESSEE	2845	30.5	1.0	(28.5, 32.5)
TEXAS	5771	30.0	0.7	(28.6, 31.4)
UTAH	3550	31.5	1.0	(29.5, 33.5)
VERMONT	4178	24.4	0.8	(22.8, 26.0)
VIRGINIA	2860	23.6	1.0	(21.6, 25.6)
WASHINGTON	4058	28.5	0.8	(26.9, 30.1)
WEST VIRGINIA	2994	24.8	0.9	(23.0, 26.6)
WISCONSIN	3270	28.3	1.0	(26.3, 30.3)
WYOMING	2974	27.4	1.0	(25.4, 29.4)
GUAM	847	32.0	1.9	(28.3, 35.7)
PUERTO RICO	4096	24.4	1.0	(22.4, 26.4)
VIRGIN ISLANDS	2203	26.9	1.3	(24.4, 29.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 27.6 RANGE = 18.6-34.6

HP2010: HP 12.15

*AT RISK NO CHOLESTEROL CHECK IN PAST 5 YEARS

August 8,

2002

TABLE 6.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BLOOD CHOLESTEROL: NO CHOLESTEROL CHECK IN THE PAST FIVE YEARS*

-BY SEX

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1023	27.6	1.7	(24.3, 30.9)	*	1702	25.7	1.2	(23.3, 28.1)
ALASKA	1315	31.0	1.9	(27.3, 34.7)	*	1453	32.8	2.0	(28.9, 36.7)
ARIZONA	1314	29.8	1.8	(26.3, 33.3)	*	1872	25.3	1.6	(22.2, 28.4)
ARKANSAS	1087	30.9	1.6	(27.8, 34.0)	*	1738	27.8	1.2	(25.4, 30.2)
CALIFORNIA	1699	33.0	1.4	(30.3, 35.7)	*	2476	25.0	1.0	(23.0, 27.0)
COLORADO	843	30.9	1.8	(27.4, 34.4)	*	1145	26.0	1.4	(23.3, 28.7)
CONNECTICUT	3130	22.4	1.0	(20.4, 24.4)	*	4389	18.7	0.7	(17.3, 20.1)
DELAWARE	1333	23.4	1.5	(20.5, 26.3)	*	2108	19.4	1.2	(17.0, 21.8)
DISTRICT OF CO	764	23.5	2.0	(19.6, 27.4)	*	1079	16.9	1.3	(14.4, 19.4)
FLORIDA	1868	24.8	1.2	(22.4, 27.2)	*	2699	19.5	0.9	(17.7, 21.3)
GEORGIA	1660	26.9	1.5	(24.0, 29.8)	*	2694	24.2	1.1	(22.0, 26.4)
HAWAII	1903	25.8	1.4	(23.1, 28.5)	*	2506	22.8	1.2	(20.4, 25.2)
IDAHO	2013	36.4	1.3	(33.9, 38.9)	*	2661	31.9	1.1	(29.7, 34.1)
ILLINOIS	848	31.5	1.8	(28.0, 35.0)	*	1270	27.5	1.5	(24.6, 30.4)
INDIANA	1566	29.3	1.3	(26.8, 31.8)	*	2301	26.1	1.0	(24.1, 28.1)
IOWA	1446	32.0	1.5	(29.1, 34.9)	*	2068	25.8	1.2	(23.4, 28.2)
KANSAS	1797	29.6	1.2	(27.2, 32.0)	*	2628	26.0	1.0	(24.0, 28.0)
KENTUCKY	2820	29.3	1.3	(26.8, 31.8)	*	4420	26.2	1.0	(24.2, 28.2)
LOUISIANA	1816	29.0	1.2	(26.6, 31.4)	*	3025	25.8	0.9	(24.0, 27.6)
MAINE	1000	27.0	1.6	(23.9, 30.1)	*	1343	19.5	1.3	(17.0, 22.0)
MARYLAND	1784	22.1	1.3	(19.6, 24.6)	*	2566	18.2	1.0	(16.2, 20.2)
MASSACHUSETTS	3459	20.0	0.8	(18.4, 21.6)	*	4895	17.2	0.7	(15.8, 18.6)
MICHIGAN	1483	29.4	1.4	(26.7, 32.1)	*	2229	22.2	1.0	(20.2, 24.2)
MINNESOTA	1644	29.0	1.3	(26.5, 31.5)	*	2169	18.8	1.0	(16.8, 20.8)
MISSISSIPPI	1095	31.7	1.7	(28.4, 35.0)	*	1802	28.1	1.2	(25.7, 30.5)
MISSOURI	1603	29.1	1.5	(26.2, 32.0)	*	2435	26.3	1.2	(23.9, 28.7)
MONTANA	1391	31.9	1.7	(28.6, 35.2)	*	1866	28.3	1.4	(25.6, 31.0)
NEBRASKA	1384	38.2	1.6	(35.1, 41.3)	*	2214	31.2	1.2	(28.8, 33.6)
NEVADA	1283	31.1	1.9	(27.4, 34.8)	*	1222	23.9	1.8	(20.4, 27.4)
NEW HAMPSHIRE	1679	24.7	1.2	(22.3, 27.1)	*	2267	21.1	1.0	(19.1, 23.1)
NEW JERSEY	2387	21.8	1.2	(19.4, 24.2)	*	3477	19.3	0.9	(17.5, 21.1)
NEW MEXICO	1471	35.1	1.5	(32.2, 38.0)	*	2069	28.7	1.2	(26.3, 31.1)
NEW YORK	1575	24.7	1.4	(22.0, 27.4)	*	2217	20.9	1.1	(18.7, 23.1)
NORTH CAROLINA	2327	27.6	1.5	(24.7, 30.5)	*	3722	22.5	1.1	(20.3, 24.7)
NORTH DAKOTA	1038	32.2	1.6	(29.1, 35.3)	*	1403	25.5	1.3	(23.0, 28.0)
OHIO	1320	29.6	1.6	(26.5, 32.7)	*	2009	26.8	1.2	(24.4, 29.2)
OKLAHOMA	1705	30.2	1.5	(27.3, 33.1)	*	2662	27.0	1.1	(24.8, 29.2)
OREGON	1018	31.4	1.6	(28.3, 34.5)	*	1425	27.0	1.4	(24.3, 29.7)
PENNSYLVANIA	1445	24.4	1.3	(21.9, 26.9)	*	2116	25.3	1.2	(22.9, 27.7)
RHODE ISLAND	1509	22.5	1.4	(19.8, 25.2)	*	2514	16.4	0.9	(14.6, 18.2)
SOUTH CAROLINA	1231	23.4	1.4	(20.7, 26.1)	*	1888	21.0	1.2	(18.6, 23.4)
SOUTH DAKOTA	2101	33.5	1.2	(31.1, 35.9)	*	2839	27.6	1.0	(25.6, 29.6)
TENNESSEE	926	32.8	1.8	(29.3, 36.3)	*	1919	28.5	1.2	(26.1, 30.9)
TEXAS	2372	33.7	1.1	(31.5, 35.9)	*	3399	26.5	0.9	(24.7, 28.3)
UTAH	1622	33.1	1.6	(30.0, 36.2)	*	1928	29.9	1.3	(27.4, 32.4)
VERMONT	1738	27.2	1.2	(24.8, 29.6)	*	2440	21.9	1.0	(19.9, 23.9)
VIRGINIA	1208	25.5	1.4	(22.8, 28.2)	*	1652	21.8	1.3	(19.3, 24.3)
WASHINGTON	1700	31.9	1.3	(29.4, 34.4)	*	2358	25.4	1.1	(23.2, 27.6)
WEST VIRGINIA	1185	26.1	1.5	(23.2, 29.0)	*	1809	23.6	1.1	(21.4, 25.8)
WISCONSIN	1436	31.6	1.5	(28.7, 34.5)	*	1834	25.2	1.2	(22.8, 27.6)
WYOMING	1301	31.3	1.5	(28.4, 34.2)	*	1673	23.6	1.2	(21.2, 26.0)
GUAM	371	32.8	2.8	(27.3, 38.3)	*	476	31.2	2.4	(26.5, 35.9)
PUERTO RICO	1450	28.2	1.7	(24.9, 31.5)	*	2646	21.1	1.3	(18.6, 23.6)
VIRGIN ISLANDS	849	31.4	2.0	(27.5, 35.3)	*	1354	23.1	1.6	(20.0, 26.2)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 29.5
 RANGE = 20.0-38.2

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 25.3
 RANGE = 16.4-32.8

*AT RISK NO CHOLESTEROL CHECK IN PAST 5 YEARS

August 8,

TABLE 6.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1999 TO 2001
BLOOD CHOLESTEROL: NO CHOLESTEROL CHECK IN THE PAST FIVE YEARS*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	30.4	26.6	-3.8	-12.5
ALASKA	35.1	31.9	-3.1	-9.0
ARIZONA	24.6	27.5	2.9	11.8
ARKANSAS	35.4	29.3	-6.1	-17.3
CALIFORNIA	32.5	29.0	-3.5	-10.9
COLORADO	33.3	28.4	-4.9	-14.7
CONNECTICUT	24.8	20.5	-4.3	-17.3
DELAWARE	26.5	21.3	-5.2	-19.5
DISTRICT OF CO	19.7	19.9	0.2	0.9
FLORIDA	23.9	22.0	-1.9	-8.0
GEORGIA	26.6	25.5	-1.1	-4.2
HAWAII	32.7	24.3	-8.4	-25.8
IDAHO	36.2	34.1	-2.1	-5.8
ILLINOIS	31.5	29.4	-2.1	-6.8
INDIANA	29.5	27.6	-1.9	-6.3
IOWA	32.7	28.8	-3.9	-12.0
KANSAS	31.1	27.8	-3.3	-10.6
KENTUCKY	32.8	27.7	-5.1	-15.5
LOUISIANA	32.9	27.3	-5.6	-17.0
MAINE	26.8	23.1	-3.7	-13.7
MARYLAND	23.5	20.1	-3.4	-14.4
MASSACHUSETTS	22.8	18.6	-4.2	-18.4
MICHIGAN	28.6	25.7	-2.9	-10.1
MINNESOTA	40.2	23.8	-16.4	-40.8
MISSISSIPPI	33.5	29.8	-3.7	-11.1
MISSOURI	34.3	27.6	-6.7	-19.4
MONTANA	35.0	30.1	-4.9	-14.0
NEBRASKA	35.3	34.6	-0.7	-2.0
NEVADA	32.3	27.6	-4.7	-14.6
NEW HAMPSHIRE	25.7	22.8	-2.9	-11.4
NEW JERSEY	24.5	20.5	-4.0	-16.4
NEW MEXICO	37.9	31.8	-6.1	-16.0
NEW YORK	27.6	22.7	-4.9	-17.8
NORTH CAROLINA	25.7	24.9	-0.8	-3.0
NORTH DAKOTA	35.7	28.8	-6.9	-19.3
OHIO	31.3	28.1	-3.2	-10.1
OKLAHOMA	27.7	28.6	0.9	3.2
OREGON	32.9	29.1	-3.8	-11.6
PENNSYLVANIA	28.5	24.9	-3.6	-12.6
RHODE ISLAND	24.3	19.3	-5.0	-20.4
SOUTH CAROLINA	27.7	22.1	-5.6	-20.3
SOUTH DAKOTA	36.4	30.5	-5.9	-16.3
TENNESSEE	27.0	30.5	3.6	13.2
TEXAS	30.8	30.0	-0.8	-2.7
UTAH	36.4	31.5	-4.9	-13.6
VERMONT	29.6	24.4	-5.2	-17.6
VIRGINIA	27.1	23.6	-3.5	-12.9
WASHINGTON	31.6	28.5	-3.1	-9.8
WEST VIRGINIA	31.0	24.8	-6.2	-20.0
WISCONSIN	29.9	28.3	-1.6	-5.2
WYOMING	31.4	27.4	-4.0	-12.7
GUAM	N/A	32.0	N/A	N/A
PUERTO RICO	26.7	24.4	-2.3	-8.5
VIRGIN ISLANDS	N/A	26.9	N/A	N/A

SUMMARY STATISTICS: yr1999

NUMBER OF PARTICIPANTS = 52
MEDIAN = 30.9
RANGE = 19.7-40.2

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
MEDIAN = 27.6
RANGE = 18.6-34.6

*AT RISK NO CHOLESTEROL CHECK IN PAST 5 YEARS
N/A - NOT AVAILABLE

August 8, 2002

TABLE 7.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FLU: NO FLUSHOT IN THE PAST 12 MONTHS, AGE 65+*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	544	34.1	2.2	(29.8, 38.4)
ALASKA	232	37.2	4.5	(28.4, 46.0)
ARIZONA	699	38.2	2.6	(33.1, 43.3)
ARKANSAS	592	36.8	2.1	(32.7, 40.9)
CALIFORNIA	762	31.1	2.2	(26.8, 35.4)
COLORADO	279	22.6	2.7	(17.3, 27.9)
CONNECTICUT	1444	30.9	1.4	(28.2, 33.6)
DELAWARE	735	32.4	2.1	(28.3, 36.5)
DISTRICT OF CO	277	44.5	3.3	(38.0, 51.0)
FLORIDA	1120	45.1	1.6	(42.0, 48.2)
GEORGIA	682	37.8	2.2	(33.5, 42.1)
HAWAII	817	21.0	1.8	(17.5, 24.5)
IDAHO	949	34.9	1.8	(31.4, 38.4)
ILLINOIS	390	37.8	2.7	(32.5, 43.1)
INDIANA	737	34.3	1.9	(30.6, 38.0)
IOWA	882	27.2	1.7	(23.9, 30.5)
KANSAS	871	31.5	1.7	(28.2, 34.8)
KENTUCKY	1603	39.1	1.8	(35.6, 42.6)
LOUISIANA	853	43.9	1.9	(40.2, 47.6)
MAINE	475	28.5	2.2	(24.2, 32.8)
MARYLAND	741	32.7	2.2	(28.4, 37.0)
MASSACHUSETTS	1460	29.4	1.4	(26.7, 32.1)
MICHIGAN	677	39.6	2.1	(35.5, 43.7)
MINNESOTA	790	29.9	1.8	(26.4, 33.4)
MISSISSIPPI	589	38.2	2.2	(33.9, 42.5)
MISSOURI	895	32.5	2.1	(28.4, 36.6)
MONTANA	738	26.9	2.1	(22.8, 31.0)
NEBRASKA	859	29.9	1.8	(26.4, 33.4)
NEVADA	450	36.7	3.2	(30.4, 43.0)
NEW HAMPSHIRE	673	30.6	1.9	(26.9, 34.3)
NEW JERSEY	1112	35.5	1.7	(32.2, 38.8)
NEW MEXICO	720	30.0	1.8	(26.5, 33.5)
NEW YORK	596	37.5	2.3	(33.0, 42.0)
NORTH CAROLINA	1188	33.9	2.0	(30.0, 37.8)
NORTH DAKOTA	467	30.0	2.3	(25.5, 34.5)
OHIO	644	36.6	2.2	(32.3, 40.9)
OKLAHOMA	1031	27.3	1.8	(23.8, 30.8)
OREGON	470	28.3	2.2	(24.0, 32.6)
PENNSYLVANIA	788	36.2	1.9	(32.5, 39.9)
RHODE ISLAND	792	27.4	1.8	(23.9, 30.9)
SOUTH CAROLINA	524	33.8	2.3	(29.3, 38.3)
SOUTH DAKOTA	1261	25.9	1.3	(23.4, 28.4)
TENNESSEE	518	34.4	2.3	(29.9, 38.9)
TEXAS	995	38.2	1.7	(34.9, 41.5)
UTAH	553	31.3	2.8	(25.8, 36.8)
VERMONT	687	28.5	1.8	(25.0, 32.0)
VIRGINIA	472	34.7	2.4	(30.0, 39.4)
WASHINGTON	706	27.5	1.8	(24.0, 31.0)
WEST VIRGINIA	741	38.3	1.9	(34.6, 42.0)
WISCONSIN	578	29.6	2.2	(25.3, 33.9)
WYOMING	521	30.4	2.2	(26.1, 34.7)
GUAM	60	60.5	7.2	(46.4, 74.6)
PUERTO RICO	933	63.2	2.2	(58.9, 67.5)
VIRGIN ISLANDS	266	61.3	3.7	(54.0, 68.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 33.8 RANGE = 21.0-63.2

HP2010: LHI, HP 14.29a

*AT RISK HAVEN'T HAD A FLUSHOT IN THE PAST TWELVE MONTHS, AGE 65 AND OLDER

August 8, 2002

TABLE 7.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FLU: NO FLUSHOT IN THE PAST 12 MONTHS, AGE 65+* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	177	30.1	3.6	(23.0, 37.2)	*	367	36.7	2.7	(31.4, 42.0)
ALASKA	99	33.4	6.7	(20.3, 46.5)	*	133	40.8	5.9	(29.2, 52.4)
ARIZONA	288	37.5	3.8	(30.1, 44.9)	*	411	38.7	3.4	(32.0, 45.4)
ARKANSAS	190	31.9	3.6	(24.8, 39.0)	*	402	40.2	2.6	(35.1, 45.3)
CALIFORNIA	279	27.1	4.1	(19.1, 35.1)	*	483	33.9	2.6	(28.8, 39.0)
COLORADO	117	21.0	3.9	(13.4, 28.6)	*	162	23.8	3.7	(16.5, 31.1)
CONNECTICUT	558	28.5	2.2	(24.2, 32.8)	*	886	32.5	1.9	(28.8, 36.2)
DELAWARE	252	30.2	3.5	(23.3, 37.1)	*	483	33.8	2.7	(28.5, 39.1)
DISTRICT OF CO	99	40.1	5.5	(29.3, 50.9)	*	178	47.1	4.1	(39.1, 55.1)
FLORIDA	431	44.5	2.6	(39.4, 49.6)	*	689	45.5	2.1	(41.4, 49.6)
GEORGIA	225	33.3	3.5	(26.4, 40.2)	*	457	40.7	2.8	(35.2, 46.2)
HAWAII	311	22.3	2.8	(16.8, 27.8)	*	506	19.9	2.3	(15.4, 24.4)
IDAHO	374	34.8	2.9	(29.1, 40.5)	*	575	35.0	2.3	(30.5, 39.5)
ILLINOIS	124	35.4	4.7	(26.2, 44.6)	*	266	39.2	3.2	(32.9, 45.5)
INDIANA	246	35.6	3.2	(29.3, 41.9)	*	491	33.4	2.3	(28.9, 37.9)
IOWA	284	24.4	2.9	(18.7, 30.1)	*	598	29.1	2.1	(25.0, 33.2)
KANSAS	306	29.5	2.8	(24.0, 35.0)	*	565	32.9	2.2	(28.6, 37.2)
KENTUCKY	515	35.0	3.3	(28.5, 41.5)	*	1088	41.8	2.1	(37.7, 45.9)
LOUISIANA	265	46.0	3.3	(39.5, 52.5)	*	588	42.6	2.2	(38.3, 46.9)
MAINE	180	26.1	3.5	(19.2, 33.0)	*	295	30.2	2.9	(24.5, 35.9)
MARYLAND	288	31.0	3.2	(24.7, 37.3)	*	453	33.9	2.9	(28.2, 39.6)
MASSACHUSETTS	515	26.3	2.2	(22.0, 30.6)	*	945	31.4	1.8	(27.9, 34.9)
MICHIGAN	238	40.0	3.4	(33.3, 46.7)	*	439	39.3	2.7	(34.0, 44.6)
MINNESOTA	298	31.5	2.8	(26.0, 37.0)	*	492	28.8	2.2	(24.5, 33.1)
MISSISSIPPI	182	39.2	3.9	(31.6, 46.8)	*	407	37.6	2.7	(32.3, 42.9)
MISSOURI	289	29.6	3.4	(22.9, 36.3)	*	606	34.5	2.7	(29.2, 39.8)
MONTANA	261	27.8	3.3	(21.3, 34.3)	*	477	26.3	2.7	(21.0, 31.6)
NEBRASKA	275	30.0	3.2	(23.7, 36.3)	*	584	29.8	2.2	(25.5, 34.1)
NEVADA	221	33.7	4.1	(25.7, 41.7)	*	229	39.1	4.7	(29.9, 48.3)
NEW HAMPSHIRE	262	30.7	3.1	(24.6, 36.8)	*	411	30.4	2.4	(25.7, 35.1)
NEW JERSEY	407	35.1	2.9	(29.4, 40.8)	*	705	35.8	2.2	(31.5, 40.1)
NEW MEXICO	265	25.4	2.9	(19.7, 31.1)	*	455	33.5	2.4	(28.8, 38.2)
NEW YORK	234	34.5	3.5	(27.6, 41.4)	*	362	39.5	3.0	(33.6, 45.4)
NORTH CAROLINA	397	39.6	3.5	(32.7, 46.5)	*	791	30.1	2.3	(25.6, 34.6)
NORTH DAKOTA	154	32.0	4.0	(24.2, 39.8)	*	313	28.5	2.7	(23.2, 33.8)
OHIO	223	36.7	3.7	(29.4, 44.0)	*	421	36.4	2.8	(30.9, 41.9)
OKLAHOMA	352	24.0	2.9	(18.3, 29.7)	*	679	29.6	2.3	(25.1, 34.1)
OREGON	174	28.3	3.6	(21.2, 35.4)	*	296	28.4	2.7	(23.1, 33.7)
PENNSYLVANIA	262	34.4	3.1	(28.3, 40.5)	*	526	37.5	2.4	(32.8, 42.2)
RHODE ISLAND	253	25.1	3.1	(19.0, 31.2)	*	539	28.9	2.3	(24.4, 33.4)
SOUTH CAROLINA	185	30.5	3.7	(23.2, 37.8)	*	339	36.0	2.8	(30.5, 41.5)
SOUTH DAKOTA	456	23.3	2.1	(19.2, 27.4)	*	805	27.9	1.7	(24.6, 31.2)
TENNESSEE	137	32.2	4.3	(23.8, 40.6)	*	381	35.9	2.7	(30.6, 41.2)
TEXAS	363	35.7	2.7	(30.4, 41.0)	*	632	40.0	2.1	(35.9, 44.1)
UTAH	236	28.0	3.9	(20.4, 35.6)	*	317	33.9	3.8	(26.5, 41.3)
VERMONT	245	24.1	2.9	(18.4, 29.8)	*	442	31.5	2.4	(26.8, 36.2)
VIRGINIA	186	31.4	3.7	(24.1, 38.7)	*	286	36.9	3.2	(30.6, 43.2)
WASHINGTON	256	28.9	3.0	(23.0, 34.8)	*	450	26.5	2.2	(22.2, 30.8)
WEST VIRGINIA	262	34.2	3.1	(28.1, 40.3)	*	479	41.1	2.5	(36.2, 46.0)
WISCONSIN	199	24.8	3.2	(18.5, 31.1)	*	379	33.0	2.8	(27.5, 38.5)
WYOMING	195	25.3	3.3	(18.8, 31.8)	*	326	34.3	2.9	(28.6, 40.0)
PUERTO RICO	334	58.6	3.6	(51.5, 65.7)	*	599	66.8	2.6	(61.7, 71.9)
VIRGIN ISLANDS	105	63.0	5.7	(51.8, 74.2)	*	161	59.9	4.9	(50.3, 69.5)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 53
MEDIAN = 31.5
RANGE = 21.0-63.0

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 53
MEDIAN = 34.4
RANGE = 19.9-66.8

*AT RISK HAVEN'T HAD A FLUSHOT IN THE PAST TWELVE MONTHS, AGE 65 AND OLDER August 8, 2002
NOTE: GUAM WASN'T LISTED IN THE TABLE, DUE TO SAMPLE < 50, (RESPONDENTS = 24 MALES AND 36 FEMALES)

TABLE 7.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2001
 FLU: NO FLUSHOT IN THE PAST 12 MONTHS, AGE 65+*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	35.4	34.1	-1.3	-3.6
ALASKA	40.2	37.2	-3.0	-7.5
ARIZONA	28.7	38.2	9.5	33.1
ARKANSAS	32.7	36.8	4.1	12.4
CALIFORNIA	27.8	31.1	3.3	11.9
COLORADO	25.2	22.6	-2.6	-10.5
CONNECTICUT	35.2	30.9	-4.3	-12.3
DELAWARE	32.3	32.4	0.1	0.3
DISTRICT OF CO	44.2	44.5	0.4	0.8
FLORIDA	36.7	45.1	8.4	22.9
GEORGIA	43.1	37.8	-5.3	-12.2
HAWAII	25.9	21.0	-4.9	-19.0
IDAHO	31.0	34.9	3.9	12.5
ILLINOIS	32.5	37.8	5.3	16.3
INDIANA	33.9	34.3	0.4	1.3
IOWA	30.4	27.2	-3.2	-10.6
KANSAS	33.0	31.5	-1.5	-4.5
KENTUCKY	31.6	39.1	7.5	23.6
LOUISIANA	39.7	43.9	4.2	10.6
MAINE	26.3	28.5	2.2	8.3
MARYLAND	37.4	32.7	-4.7	-12.6
MASSACHUSETTS	30.6	29.4	-1.2	-4.0
MICHIGAN	30.0	39.6	9.6	32.0
MINNESOTA	36.0	29.9	-6.1	-16.9
MISSISSIPPI	37.2	38.2	1.0	2.7
MISSOURI	31.6	32.5	0.9	2.8
MONTANA	27.1	26.9	-0.2	-0.8
NEBRASKA	30.8	29.9	-0.9	-3.0
NEVADA	37.9	36.7	-1.1	-3.0
NEW HAMPSHIRE	34.9	30.6	-4.3	-12.3
NEW JERSEY	34.7	35.5	0.8	2.4
NEW MEXICO	31.2	30.0	-1.2	-3.9
NEW YORK	36.2	37.5	1.3	3.6
NORTH CAROLINA	35.8	33.9	-1.9	-5.4
NORTH DAKOTA	32.8	30.0	-2.8	-8.6
OHIO	31.2	36.6	5.5	17.5
OKLAHOMA	28.2	27.3	-0.9	-3.1
OREGON	34.8	28.3	-6.5	-18.8
PENNSYLVANIA	36.9	36.2	-0.7	-1.9
RHODE ISLAND	24.2	27.4	3.2	13.2
SOUTH CAROLINA	30.1	33.8	3.7	12.4
SOUTH DAKOTA	26.4	25.9	-0.5	-1.8
TENNESSEE	34.5	34.4	-0.1	-0.4
TEXAS	30.2	38.2	8.1	26.7
UTAH	24.9	31.3	6.4	25.9
VERMONT	26.6	28.5	1.9	7.3
VIRGINIA	34.3	34.7	0.4	1.1
WASHINGTON	31.1	27.5	-3.6	-11.5
WEST VIRGINIA	37.1	38.3	1.2	3.1
WISCONSIN	35.1	29.6	-5.5	-15.6
WYOMING	26.2	30.4	4.2	16.2
GUAM	N/A	60.5	N/A	N/A
PUERTO RICO	59.7	63.2	3.5	5.9
VIRGIN ISLANDS	N/A	61.3	N/A	N/A

SUMMARY STATISTICS: yr1999

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 32.6
 RANGE = 24.2-59.7

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 33.8
 RANGE = 21.0-63.2

*AT RISK HAVEN'T HAD A FLUSHOT IN THE PAST TWELVE MONTHS, AGE 65 AND OLDER

N/A - NOT AVAILABLE

August 8, 2002

TABLE 8.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PNEUMONIA: NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65+*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	535	39.7	2.3	(35.2, 44.2)
ALASKA	220	34.7	4.4	(26.1, 43.3)
ARIZONA	674	34.4	2.5	(29.5, 39.3)
ARKANSAS	582	41.0	2.2	(36.7, 45.3)
CALIFORNIA	740	40.4	2.4	(35.7, 45.1)
COLORADO	268	31.4	3.0	(25.5, 37.3)
CONNECTICUT	1394	36.7	1.5	(33.8, 39.6)
DELAWARE	730	31.1	2.1	(27.0, 35.2)
DISTRICT OF CO	263	51.0	3.4	(44.3, 57.7)
FLORIDA	1100	41.9	1.6	(38.8, 45.0)
GEORGIA	660	42.1	2.3	(37.6, 46.6)
HAWAII	802	36.3	2.3	(31.8, 40.8)
IDAHO	916	39.7	1.9	(36.0, 43.4)
ILLINOIS	388	43.3	2.8	(37.8, 48.8)
INDIANA	721	39.8	2.0	(35.9, 43.7)
IOWA	869	34.1	1.9	(30.4, 37.8)
KANSAS	860	37.1	1.8	(33.6, 40.6)
KENTUCKY	1585	44.9	1.8	(41.4, 48.4)
LOUISIANA	844	50.5	1.9	(46.8, 54.2)
MAINE	462	35.0	2.4	(30.3, 39.7)
MARYLAND	729	37.7	2.3	(33.2, 42.2)
MASSACHUSETTS	1431	36.5	1.5	(33.6, 39.4)
MICHIGAN	658	43.4	2.1	(39.3, 47.5)
MINNESOTA	770	37.1	1.9	(33.4, 40.8)
MISSISSIPPI	575	44.3	2.3	(39.8, 48.8)
MISSOURI	887	44.0	2.2	(39.7, 48.3)
MONTANA	731	32.1	2.3	(27.6, 36.6)
NEBRASKA	850	38.8	2.0	(34.9, 42.7)
NEVADA	431	33.7	3.2	(27.4, 40.0)
NEW HAMPSHIRE	655	37.3	2.0	(33.4, 41.2)
NEW JERSEY	1084	41.1	1.9	(37.4, 44.8)
NEW MEXICO	703	37.3	2.0	(33.4, 41.2)
NEW YORK	578	44.1	2.4	(39.4, 48.8)
NORTH CAROLINA	1167	34.2	2.0	(30.3, 38.1)
NORTH DAKOTA	456	35.8	2.4	(31.1, 40.5)
OHIO	617	40.7	2.3	(36.2, 45.2)
OKLAHOMA	1024	33.9	1.9	(30.2, 37.6)
OREGON	456	29.1	2.3	(24.6, 33.6)
PENNSYLVANIA	766	40.5	2.0	(36.6, 44.4)
RHODE ISLAND	774	33.0	1.9	(29.3, 36.7)
SOUTH CAROLINA	512	42.1	2.4	(37.4, 46.8)
SOUTH DAKOTA	1244	40.8	1.5	(37.9, 43.7)
TENNESSEE	514	44.6	2.5	(39.7, 49.5)
TEXAS	970	42.0	1.7	(38.7, 45.3)
UTAH	538	32.7	2.6	(27.6, 37.8)
VERMONT	669	32.7	2.0	(28.8, 36.6)
VIRGINIA	443	39.9	2.5	(35.0, 44.8)
WASHINGTON	680	33.2	2.0	(29.3, 37.1)
WEST VIRGINIA	732	38.7	1.9	(35.0, 42.4)
WISCONSIN	566	34.4	2.3	(29.9, 38.9)
WYOMING	512	31.6	2.2	(27.3, 35.9)
GUAM	55	66.9	7.0	(53.2, 80.6)
PUERTO RICO	902	75.9	2.0	(72.0, 79.8)
VIRGIN ISLANDS	250	69.3	3.5	(62.4, 76.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 38.8 RANGE = 29.1-75.9

HP2010: LHI, HP 14.29b

*AT RISK NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65 AND OLDER

August 8, 2002

**TABLE 8.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PNEUMONIA: NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65+* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED**

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	174	39.9	3.9	(32.3, 47.5)	*	361	39.5	2.8 (34.0, 45.0)
ALASKA	90	33.1	6.6	(20.2, 46.0)	*	130	36.1	5.9 (24.5, 47.7)
ARIZONA	273	37.2	3.9	(29.6, 44.8)	*	401	32.3	3.3 (25.8, 38.8)
ARKANSAS	185	33.7	3.7	(26.4, 41.0)	*	397	46.0	2.7 (40.7, 51.3)
CALIFORNIA	269	44.5	4.0	(36.7, 52.3)	*	471	37.5	2.8 (32.0, 43.0)
COLORADO	109	34.3	4.8	(24.9, 43.7)	*	159	29.4	4.0 (21.6, 37.2)
CONNECTICUT	530	35.7	2.3	(31.2, 40.2)	*	864	37.4	2.0 (33.5, 41.3)
DELAWARE	249	32.7	3.7	(25.4, 40.0)	*	481	29.9	2.6 (24.8, 35.0)
DISTRICT OF CO	91	57.4	5.7	(46.2, 68.6)	*	172	47.6	4.2 (39.4, 55.8)
FLORIDA	419	43.7	2.6	(38.6, 48.8)	*	681	40.6	2.1 (36.5, 44.7)
GEORGIA	213	42.8	3.9	(35.2, 50.4)	*	447	41.6	2.8 (36.1, 47.1)
HAWAII	304	37.6	3.4	(30.9, 44.3)	*	498	35.2	3.1 (29.1, 41.3)
IDAHO	350	42.8	3.1	(36.7, 48.9)	*	566	37.6	2.4 (32.9, 42.3)
ILLINOIS	122	49.3	4.9	(39.7, 58.9)	*	266	39.9	3.2 (33.6, 46.2)
INDIANA	239	45.7	3.4	(39.0, 52.4)	*	482	35.8	2.3 (31.3, 40.3)
IOWA	276	31.3	3.1	(25.2, 37.4)	*	593	36.0	2.3 (31.5, 40.5)
KANSAS	299	40.1	3.0	(34.2, 46.0)	*	561	35.2	2.2 (30.9, 39.5)
KENTUCKY	507	41.7	3.3	(35.2, 48.2)	*	1078	47.1	2.1 (43.0, 51.2)
LOUISIANA	261	53.8	3.4	(47.1, 60.5)	*	583	48.4	2.3 (43.9, 52.9)
MAINE	172	40.5	3.9	(32.9, 48.1)	*	290	31.4	2.9 (25.7, 37.1)
MARYLAND	282	37.0	3.4	(30.3, 43.7)	*	447	38.2	3.1 (32.1, 44.3)
MASSACHUSETTS	500	41.0	2.5	(36.1, 45.9)	*	931	33.6	1.8 (30.1, 37.1)
MICHIGAN	229	44.9	3.5	(38.0, 51.8)	*	429	42.4	2.6 (37.3, 47.5)
MINNESOTA	280	38.7	3.1	(32.6, 44.8)	*	490	36.0	2.4 (31.3, 40.7)
MISSISSIPPI	175	47.7	4.0	(39.9, 55.5)	*	400	42.2	2.7 (36.9, 47.5)
MISSOURI	283	43.5	3.8	(36.1, 50.9)	*	604	44.4	2.7 (39.1, 49.7)
MONTANA	258	32.7	3.7	(25.4, 40.0)	*	473	31.8	2.9 (26.1, 37.5)
NEBRASKA	272	40.9	3.3	(34.4, 47.4)	*	578	37.4	2.3 (32.9, 41.9)
NEVADA	205	30.9	4.0	(23.1, 38.7)	*	226	35.7	4.7 (26.5, 44.9)
NEW HAMPSHIRE	249	36.6	3.3	(30.1, 43.1)	*	406	37.8	2.6 (32.7, 42.9)
NEW JERSEY	396	44.0	3.0	(38.1, 49.9)	*	688	39.1	2.4 (34.4, 43.8)
NEW MEXICO	254	38.2	3.3	(31.7, 44.7)	*	449	36.6	2.5 (31.7, 41.5)
NEW YORK	224	43.9	3.8	(36.5, 51.3)	*	354	44.3	3.1 (38.2, 50.4)
NORTH CAROLINA	388	36.1	3.5	(29.2, 43.0)	*	779	33.0	2.4 (28.3, 37.7)
NORTH DAKOTA	151	35.2	4.1	(27.2, 43.2)	*	305	36.3	3.0 (30.4, 42.2)
OHIO	210	41.9	3.9	(34.3, 49.5)	*	407	39.7	2.9 (34.0, 45.4)
OKLAHOMA	349	34.3	3.1	(28.2, 40.4)	*	675	33.7	2.4 (29.0, 38.4)
OREGON	165	30.0	3.8	(22.6, 37.4)	*	291	28.6	2.8 (23.1, 34.1)
PENNSYLVANIA	250	40.1	3.3	(33.6, 46.6)	*	516	40.8	2.5 (35.9, 45.7)
RHODE ISLAND	243	32.6	3.3	(26.1, 39.1)	*	531	33.2	2.4 (28.5, 37.9)
SOUTH CAROLINA	178	41.6	4.1	(33.6, 49.6)	*	334	42.3	3.0 (36.4, 48.2)
SOUTH DAKOTA	447	40.5	2.5	(35.6, 45.4)	*	797	41.0	1.9 (37.3, 44.7)
TENNESSEE	136	45.3	4.6	(36.3, 54.3)	*	378	44.0	2.7 (38.7, 49.3)
TEXAS	351	40.5	2.8	(35.0, 46.0)	*	619	43.0	2.2 (38.7, 47.3)
UTAH	226	37.4	4.0	(29.6, 45.2)	*	312	29.0	3.2 (22.7, 35.3)
VERMONT	237	27.9	3.0	(22.0, 33.8)	*	432	36.0	2.5 (31.1, 40.9)
VIRGINIA	168	39.3	4.0	(31.5, 47.1)	*	275	40.2	3.3 (33.7, 46.7)
WASHINGTON	243	37.5	3.3	(31.0, 44.0)	*	437	30.2	2.4 (25.5, 34.9)
WEST VIRGINIA	257	34.1	3.1	(28.0, 40.2)	*	475	41.7	2.5 (36.8, 46.6)
WISCONSIN	191	37.2	3.9	(29.6, 44.8)	*	375	32.5	2.8 (27.0, 38.0)
WYOMING	191	26.0	3.4	(19.3, 32.7)	*	321	35.9	2.9 (30.2, 41.6)
PUERTO RICO	323	74.3	3.3	(67.8, 80.8)	*	579	77.1	2.5 (72.2, 82.0)
VIRGIN ISLANDS	98	66.9	5.9	(55.3, 78.5)	*	152	71.2	4.2 (63.0, 79.4)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 53

MEDIAN = 40.0
RANGE = 26.0-74.3

*AT RISK NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65 AND OLDER

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 53

MEDIAN = 37.6
RANGE = 28.6-77.1

August 8, 2002

NOTE: GUAM WASN'T LISTED IN THE TABLE, DUE TO SAMPLE < 50, (RESPONDENTS = 20 MALES AND 35 FEMALES)

TABLE 8.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2001
 PNEUMONIA: NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65+*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	46.1	39.7	-6.4	-13.9
ALASKA	56.2	34.7	-21.5	-38.3
ARIZONA	46.6	34.4	-12.2	-26.1
ARKANSAS	49.8	41.0	-8.8	-17.7
CALIFORNIA	43.0	40.4	-2.6	-6.1
COLORADO	37.3	31.4	-5.9	-15.7
CONNECTICUT	51.0	36.7	-14.3	-28.1
DELAWARE	33.5	31.1	-2.4	-7.0
DISTRICT OF CO	64.7	51.0	-13.7	-21.2
FLORIDA	46.5	41.9	-4.6	-9.9
GEORGIA	50.3	42.1	-8.2	-16.3
HAWAII	44.2	36.3	-7.9	-17.9
IDAHO	44.8	39.7	-5.1	-11.4
ILLINOIS	52.6	43.3	-9.3	-17.7
INDIANA	48.4	39.8	-8.6	-17.7
IOWA	38.8	34.1	-4.7	-12.0
KANSAS	44.9	37.1	-7.8	-17.4
KENTUCKY	48.0	44.9	-3.1	-6.5
LOUISIANA	59.6	50.5	-9.1	-15.3
MAINE	42.7	35.0	-7.7	-18.1
MARYLAND	45.9	37.7	-8.2	-17.8
MASSACHUSETTS	43.2	36.5	-6.7	-15.6
MICHIGAN	42.3	43.4	1.1	2.7
MINNESOTA	48.1	37.1	-11.0	-22.9
MISSISSIPPI	49.6	44.3	-5.3	-10.7
MISSOURI	47.2	44.0	-3.2	-6.8
MONTANA	38.8	32.1	-6.7	-17.4
NEBRASKA	45.2	38.8	-6.4	-14.1
NEVADA	38.3	33.7	-4.6	-11.9
NEW HAMPSHIRE	39.7	37.3	-2.4	-5.9
NEW JERSEY	44.9	41.1	-3.8	-8.5
NEW MEXICO	46.8	37.3	-9.5	-20.2
NEW YORK	50.0	44.1	-5.9	-11.9
NORTH CAROLINA	41.5	34.2	-7.3	-17.5
NORTH DAKOTA	45.0	35.8	-9.2	-20.4
OHIO	45.0	40.7	-4.3	-9.6
OKLAHOMA	46.3	33.9	-12.4	-26.8
OREGON	43.8	29.1	-14.7	-33.5
PENNSYLVANIA	47.8	40.5	-7.3	-15.2
RHODE ISLAND	43.1	33.0	-10.1	-23.4
SOUTH CAROLINA	43.9	42.1	-1.8	-4.0
SOUTH DAKOTA	49.6	40.8	-8.8	-17.8
TENNESSEE	45.7	44.6	-1.1	-2.4
TEXAS	44.1	42.0	-2.1	-4.8
UTAH	38.7	32.7	-6.0	-15.4
VERMONT	43.5	32.7	-10.8	-24.9
VIRGINIA	44.8	39.9	-4.9	-10.9
WASHINGTON	44.2	33.2	-11.0	-24.8
WEST VIRGINIA	45.7	38.7	-7.0	-15.3
WISCONSIN	46.3	34.4	-11.9	-25.8
WYOMING	38.5	31.6	-6.9	-18.0
GUAM	N/A	66.9	N/A	N/A
PUERTO RICO	78.2	75.9	-2.3	-2.9
VIRGIN ISLANDS	N/A	69.3	N/A	N/A

SUMMARY STATISTICS: yr1999

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 45.1
 RANGE = 33.5-78.2

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 38.8
 RANGE = 29.1-75.9

*AT RISK NEVER HAD A PNEUMOCOCCAL VACCINATION, AGE 65 AND OLDER
 N/A - NOT AVAILABLE

August 8, 2002

TABLE 9.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESITY: BODY MASS INDEX GE 30.0 AND LE 99.8*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2700	24.5	0.9	(22.7, 26.3)
ALASKA	2759	22.1	1.1	(19.9, 24.3)
ARIZONA	3055	18.5	1.0	(16.5, 20.5)
ARKANSAS	2814	22.4	0.9	(20.6, 24.2)
CALIFORNIA	4120	21.9	0.8	(20.3, 23.5)
COLORADO	1940	14.9	0.9	(13.1, 16.7)
CONNECTICUT	7296	17.9	0.5	(16.9, 18.9)
DELAWARE	3327	20.8	0.9	(19.0, 22.6)
DISTRICT OF CO	1794	20.0	1.1	(17.8, 22.2)
FLORIDA	4396	18.8	0.7	(17.4, 20.2)
GEORGIA	4286	22.7	0.8	(21.1, 24.3)
HAWAII	4403	17.9	0.9	(16.1, 19.7)
IDAHO	4588	20.5	0.7	(19.1, 21.9)
ILLINOIS	3807	21.0	0.8	(19.4, 22.6)
INDIANA	3861	24.5	0.8	(22.9, 26.1)
IOWA	3427	22.5	0.8	(20.9, 24.1)
KANSAS	4315	21.6	0.7	(20.2, 23.0)
KENTUCKY	7135	24.6	0.8	(23.0, 26.2)
LOUISIANA	4719	24.0	0.7	(22.6, 25.4)
MAINE	2292	19.5	0.9	(17.7, 21.3)
MARYLAND	4262	20.5	0.8	(18.9, 22.1)
MASSACHUSETTS	8081	16.6	0.5	(15.6, 17.6)
MICHIGAN	3677	25.0	0.8	(23.4, 26.6)
MINNESOTA	3811	19.9	0.7	(18.5, 21.3)
MISSISSIPPI	2888	26.5	0.9	(24.7, 28.3)
MISSOURI	3996	23.2	0.9	(21.4, 25.0)
MONTANA	3202	18.8	0.9	(17.0, 20.6)
NEBRASKA	3457	20.7	0.8	(19.1, 22.3)
NEVADA	2480	19.5	1.3	(17.0, 22.0)
NEW HAMPSHIRE	3830	19.4	0.7	(18.0, 20.8)
NEW JERSEY	5655	19.6	0.7	(18.2, 21.0)
NEW MEXICO	3472	19.7	0.8	(18.1, 21.3)
NEW YORK	3687	20.3	0.8	(18.7, 21.9)
NORTH CAROLINA	5852	22.9	0.9	(21.1, 24.7)
NORTH DAKOTA	2415	20.4	0.9	(18.6, 22.2)
OHIO	3239	22.4	0.9	(20.6, 24.2)
OKLAHOMA	4220	22.6	0.8	(21.0, 24.2)
OREGON	2433	21.1	0.9	(19.3, 22.9)
PENNSYLVANIA	3528	22.1	0.8	(20.5, 23.7)
RHODE ISLAND	3866	17.7	0.8	(16.1, 19.3)
SOUTH CAROLINA	3038	22.5	0.9	(20.7, 24.3)
SOUTH DAKOTA	4876	21.2	0.6	(20.0, 22.4)
TENNESSEE	2735	23.4	0.9	(21.6, 25.2)
TEXAS	5652	24.6	0.7	(23.2, 26.0)
UTAH	3534	19.1	0.9	(17.3, 20.9)
VERMONT	4119	17.6	0.7	(16.2, 19.0)
VIRGINIA	2819	20.9	0.9	(19.1, 22.7)
WASHINGTON	4026	19.3	0.7	(17.9, 20.7)
WEST VIRGINIA	2931	25.1	0.9	(23.3, 26.9)
WISCONSIN	3257	22.4	0.8	(20.8, 24.0)
WYOMING	2952	19.7	0.8	(18.1, 21.3)
GUAM	842	21.2	1.6	(18.1, 24.3)
PUERTO RICO	4062	22.2	0.9	(20.4, 24.0)
VIRGIN ISLANDS	2173	24.7	1.2	(22.3, 27.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 21.1 RANGE = 14.9-26.5

*AT RISK NEW BODY MASS INDEX(_BMI2 GE 30.0 AND LE 99.8), AGE 18 AND OLDER

August 8,

2002

TABLE 9.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESITY: BODY MASS INDEX GE 30.0 AND LE 99.8* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	1042	24.1	1.5	(21.2, 27.0)	*	1658	24.8	1.2	(22.4, 27.2)
ALASKA	1345	19.2	1.5	(16.3, 22.1)	*	1414	25.5	1.6	(22.4, 28.6)
ARIZONA	1297	19.6	1.5	(16.7, 22.5)	*	1758	17.2	1.3	(14.7, 19.7)
ARKANSAS	1114	22.8	1.4	(20.1, 25.5)	*	1700	22.1	1.1	(19.9, 24.3)
CALIFORNIA	1695	20.6	1.2	(18.2, 23.0)	*	2425	23.2	1.1	(21.0, 25.4)
COLORADO	849	15.1	1.3	(12.6, 17.6)	*	1091	14.8	1.2	(12.4, 17.2)
CONNECTICUT	3151	18.7	0.8	(17.1, 20.3)	*	4145	17.1	0.7	(15.7, 18.5)
DELAWARE	1340	21.2	1.4	(18.5, 23.9)	*	1987	20.3	1.1	(18.1, 22.5)
DISTRICT OF CO	771	16.3	1.6	(13.2, 19.4)	*	1023	23.5	1.6	(20.4, 26.6)
FLORIDA	1865	18.8	1.0	(16.8, 20.8)	*	2531	18.7	0.9	(16.9, 20.5)
GEORGIA	1692	21.8	1.3	(19.3, 24.3)	*	2594	23.5	1.0	(21.5, 25.5)
HAWAII	1919	19.5	1.3	(17.0, 22.0)	*	2484	16.2	1.1	(14.0, 18.4)
IDAHO	2047	21.3	1.1	(19.1, 23.5)	*	2541	19.7	0.9	(17.9, 21.5)
ILLINOIS	1604	20.3	1.1	(18.1, 22.5)	*	2203	21.7	1.0	(19.7, 23.7)
INDIANA	1598	24.9	1.2	(22.5, 27.3)	*	2263	24.2	1.0	(22.2, 26.2)
IOWA	1459	23.9	1.3	(21.4, 26.4)	*	1968	21.2	1.0	(19.2, 23.2)
KANSAS	1823	23.2	1.1	(21.0, 25.4)	*	2492	20.0	0.9	(18.2, 21.8)
KENTUCKY	2871	25.7	1.2	(23.3, 28.1)	*	4264	23.6	0.9	(21.8, 25.4)
LOUISIANA	1855	24.5	1.1	(22.3, 26.7)	*	2864	23.6	0.9	(21.8, 25.4)
MAINE	1015	19.9	1.4	(17.2, 22.6)	*	1277	19.1	1.2	(16.7, 21.5)
MARYLAND	1792	19.8	1.2	(17.4, 22.2)	*	2470	21.1	1.1	(18.9, 23.3)
MASSACHUSETTS	3476	17.4	0.8	(15.8, 19.0)	*	4605	15.8	0.6	(14.6, 17.0)
MICHIGAN	1503	25.2	1.3	(22.7, 27.7)	*	2174	24.7	1.0	(22.7, 26.7)
MINNESOTA	1686	20.4	1.1	(18.2, 22.6)	*	2125	19.4	0.9	(17.6, 21.2)
MISSISSIPPI	1128	25.2	1.4	(22.5, 27.9)	*	1760	27.7	1.2	(25.3, 30.1)
MISSOURI	1638	25.2	1.4	(22.5, 27.9)	*	2358	21.2	1.1	(19.0, 23.4)
MONTANA	1404	19.7	1.3	(17.2, 22.2)	*	1798	17.9	1.2	(15.5, 20.3)
NEBRASKA	1387	22.2	1.3	(19.7, 24.7)	*	2070	19.2	1.0	(17.2, 21.2)
NEVADA	1289	22.6	2.1	(18.5, 26.7)	*	1191	16.3	1.5	(13.4, 19.2)
NEW HAMPSHIRE	1694	21.3	1.1	(19.1, 23.5)	*	2136	17.4	0.9	(15.6, 19.2)
NEW JERSEY	2369	19.4	1.1	(17.2, 21.6)	*	3286	19.9	0.9	(18.1, 21.7)
NEW MEXICO	1471	20.7	1.3	(18.2, 23.2)	*	2001	18.7	1.0	(16.7, 20.7)
NEW YORK	1582	20.7	1.2	(18.3, 23.1)	*	2105	20.0	1.0	(18.0, 22.0)
NORTH CAROLINA	2337	22.6	1.3	(20.1, 25.1)	*	3515	23.2	1.1	(21.0, 25.4)
NORTH DAKOTA	1059	21.9	1.3	(19.4, 24.4)	*	1356	18.8	1.2	(16.4, 21.2)
OHIO	1335	23.1	1.4	(20.4, 25.8)	*	1904	21.7	1.1	(19.5, 23.9)
OKLAHOMA	1719	24.5	1.3	(22.0, 27.0)	*	2501	20.8	1.0	(18.8, 22.8)
OREGON	1039	19.7	1.3	(17.2, 22.2)	*	1394	22.4	1.2	(20.0, 24.8)
PENNSYLVANIA	1471	21.5	1.2	(19.1, 23.9)	*	2057	22.7	1.1	(20.5, 24.9)
RHODE ISLAND	1511	18.0	1.2	(15.6, 20.4)	*	2355	17.4	0.9	(15.6, 19.2)
SOUTH CAROLINA	1245	22.7	1.3	(20.2, 25.2)	*	1793	22.2	1.2	(19.8, 24.6)
SOUTH DAKOTA	2141	23.5	1.0	(21.5, 25.5)	*	2735	18.9	0.8	(17.3, 20.5)
TENNESSEE	918	24.2	1.5	(21.3, 27.1)	*	1817	22.8	1.1	(20.6, 25.0)
TEXAS	2389	24.5	1.0	(22.5, 26.5)	*	3263	24.7	0.9	(22.9, 26.5)
UTAH	1644	20.0	1.3	(17.5, 22.5)	*	1890	18.2	1.1	(16.0, 20.4)
VERMONT	1767	18.7	1.0	(16.7, 20.7)	*	2352	16.6	0.8	(15.0, 18.2)
VIRGINIA	1224	21.2	1.3	(18.7, 23.7)	*	1595	20.5	1.2	(18.1, 22.9)
WASHINGTON	1734	20.9	1.1	(18.7, 23.1)	*	2292	17.8	0.9	(16.0, 19.6)
WEST VIRGINIA	1208	25.4	1.4	(22.7, 28.1)	*	1723	24.8	1.2	(22.4, 27.2)
WISCONSIN	1475	22.9	1.3	(20.4, 25.4)	*	1782	21.9	1.1	(19.7, 24.1)
WYOMING	1312	20.1	1.2	(17.7, 22.5)	*	1640	19.2	1.1	(17.0, 21.4)
GUAM	375	20.5	2.4	(15.8, 25.2)	*	467	22.0	2.3	(17.5, 26.5)
PUERTO RICO	1496	23.6	1.5	(20.7, 26.5)	*	2566	20.9	1.1	(18.7, 23.1)
VIRGIN ISLANDS	860	20.5	1.7	(17.2, 23.8)	*	1313	28.2	1.6	(25.1, 31.3)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 21.3
 RANGE = 15.1-25.7

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 20.9
 RANGE = 14.8-28.2

*AT RISK BY NEW BODY MASS INDEX(_BMI2 GE 30.0 AND LE 99.8), AGE 18 AND OLDER

August 8,

TABLE 9.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 OBESITY: BODY MASS INDEX GE 30.0 AND LE 99.8*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	23.9	24.5	0.6	2.5
ALASKA	21.0	22.1	1.1	5.2
ARIZONA	19.2	18.5	-0.7	-3.5
ARKANSAS	23.3	22.4	-0.9	-3.8
CALIFORNIA	19.9	21.9	2.0	9.9
COLORADO	14.2	14.9	0.7	5.1
CONNECTICUT	17.4	17.9	0.5	3.0
DELAWARE	16.6	20.8	4.2	25.4
DISTRICT OF CO	21.5	20.0	-1.5	-7.1
FLORIDA	18.7	18.8	0.1	0.6
GEORGIA	21.5	22.7	1.2	5.5
HAWAII	15.8	17.9	2.1	13.7
IDAHO	18.9	20.5	1.6	8.5
ILLINOIS	21.7	21.0	-0.7	-3.4
INDIANA	21.8	24.5	2.7	12.2
IOWA	21.5	22.5	1.0	4.6
KANSAS	20.8	21.6	0.8	3.9
KENTUCKY	23.0	24.6	1.6	7.0
LOUISIANA	23.6	24.0	0.4	1.8
MAINE	20.0	19.5	-0.4	-2.3
MARYLAND	20.2	20.5	0.3	1.6
MASSACHUSETTS	16.8	16.6	-0.1	-0.9
MICHIGAN	22.4	25.0	2.6	11.9
MINNESOTA	17.4	19.9	2.5	14.2
MISSISSIPPI	25.0	26.5	1.5	6.1
MISSOURI	22.1	23.2	1.1	5.0
MONTANA	15.9	18.8	2.9	18.2
NEBRASKA	21.1	20.7	-0.4	-1.8
NEVADA	17.9	19.5	1.6	8.8
NEW HAMPSHIRE	18.1	19.4	1.3	7.1
NEW JERSEY	18.5	19.6	1.2	6.2
NEW MEXICO	19.4	19.7	0.3	1.8
NEW YORK	17.7	20.3	2.6	14.8
NORTH CAROLINA	21.8	22.9	1.1	5.1
NORTH DAKOTA	20.4	20.4	0.0	0.0
OHIO	21.5	22.4	0.9	4.1
OKLAHOMA	19.7	22.6	2.9	14.9
OREGON	21.5	21.1	-0.4	-1.8
PENNSYLVANIA	21.2	22.1	0.9	4.0
RHODE ISLAND	17.1	17.7	0.6	3.3
SOUTH CAROLINA	22.1	22.5	0.4	2.0
SOUTH DAKOTA	19.8	21.2	1.4	6.9
TENNESSEE	22.9	23.4	0.5	2.0
TEXAS	23.1	24.6	1.5	6.5
UTAH	19.1	19.1	-0.0	-0.2
VERMONT	18.2	17.6	-0.6	-3.2
VIRGINIA	18.2	20.9	2.7	15.1
WASHINGTON	18.8	19.3	0.5	2.6
WEST VIRGINIA	23.2	25.1	1.9	8.0
WISCONSIN	20.0	22.4	2.5	12.3
WYOMING	18.0	19.7	1.7	9.6
GUAM	N/A	21.2	N/A	N/A
PUERTO RICO	21.7	22.2	0.5	2.3
VIRGIN ISLANDS	N/A	24.7	N/A	N/A

SUMMARY STATISTICS: yr2000

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 20.1
 RANGE = 14.2-25.0

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 21.1
 RANGE = 14.9-26.5

*AT RISK BY NEW BODY MASS INDEX(_BMI2 GE 30.0 AND LE 99.8), AGE 18 AND OLDER

N/A - NOT AVAILABLE

August 8, 2002

TABLE 10.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESITY: BODY MASS INDEX GE 30.0 & LE 99.8, AGE 20+ *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2629	24.9	1.0	(22.9, 26.9)
ALASKA	2653	22.9	1.1	(20.7, 25.1)
ARIZONA	2922	19.3	1.0	(17.3, 21.3)
ARKANSAS	2741	23.0	0.9	(21.2, 24.8)
CALIFORNIA	4018	22.6	0.8	(21.0, 24.2)
COLORADO	1882	15.5	0.9	(13.7, 17.3)
CONNECTICUT	7075	18.2	0.6	(17.0, 19.4)
DELAWARE	3237	21.4	0.9	(19.6, 23.2)
DISTRICT OF CO	1744	20.2	1.2	(17.8, 22.6)
FLORIDA	4245	19.2	0.7	(17.8, 20.6)
GEORGIA	4179	23.1	0.8	(21.5, 24.7)
HAWAII	4293	18.3	0.9	(16.5, 20.1)
IDAHO	4462	21.0	0.7	(19.6, 22.4)
ILLINOIS	3712	21.7	0.8	(20.1, 23.3)
INDIANA	3756	25.1	0.8	(23.5, 26.7)
IOWA	3334	23.4	0.8	(21.8, 25.0)
KANSAS	4212	22.2	0.7	(20.8, 23.6)
KENTUCKY	6976	24.8	0.8	(23.2, 26.4)
LOUISIANA	4517	25.0	0.7	(23.6, 26.4)
MAINE	2239	20.0	0.9	(18.2, 21.8)
MARYLAND	4132	20.4	0.8	(18.8, 22.0)
MASSACHUSETTS	7845	17.1	0.5	(16.1, 18.1)
MICHIGAN	3574	25.8	0.8	(24.2, 27.4)
MINNESOTA	3735	20.4	0.7	(19.0, 21.8)
MISSISSIPPI	2808	27.1	1.0	(25.1, 29.1)
MISSOURI	3877	23.8	0.9	(22.0, 25.6)
MONTANA	3143	19.2	0.9	(17.4, 21.0)
NEBRASKA	3355	21.2	0.8	(19.6, 22.8)
NEVADA	2408	19.9	1.4	(17.2, 22.6)
NEW HAMPSHIRE	3716	19.9	0.7	(18.5, 21.3)
NEW JERSEY	5499	20.0	0.7	(18.6, 21.4)
NEW MEXICO	3373	20.3	0.8	(18.7, 21.9)
NEW YORK	3571	20.8	0.8	(19.2, 22.4)
NORTH CAROLINA	5710	23.1	0.9	(21.3, 24.9)
NORTH DAKOTA	2349	20.6	0.9	(18.8, 22.4)
OHIO	3138	22.8	0.9	(21.0, 24.6)
OKLAHOMA	4119	23.2	0.8	(21.6, 24.8)
OREGON	2368	21.6	0.9	(19.8, 23.4)
PENNSYLVANIA	3439	22.6	0.8	(21.0, 24.2)
RHODE ISLAND	3764	18.1	0.8	(16.5, 19.7)
SOUTH CAROLINA	2948	23.0	0.9	(21.2, 24.8)
SOUTH DAKOTA	4785	21.6	0.7	(20.2, 23.0)
TENNESSEE	2659	24.2	1.0	(22.2, 26.2)
TEXAS	5504	25.2	0.7	(23.8, 26.6)
UTAH	3390	19.9	0.9	(18.1, 21.7)
VERMONT	4031	18.1	0.7	(16.7, 19.5)
VIRGINIA	2728	21.4	0.9	(19.6, 23.2)
WASHINGTON	3904	19.9	0.7	(18.5, 21.3)
WEST VIRGINIA	2861	25.3	0.9	(23.5, 27.1)
WISCONSIN	3179	22.7	0.9	(20.9, 24.5)
WYOMING	2872	19.9	0.8	(18.3, 21.5)
GUAM	812	21.9	1.7	(18.6, 25.2)
PUERTO RICO	3962	22.9	0.9	(21.1, 24.7)
VIRGIN ISLANDS	2071	25.5	1.2	(23.1, 27.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 21.6 RANGE = 15.5-27.1

HP2010: LHI, HP 19.2(approx.)

*AT RISK BY NEW BODY MASS INDEX(_BMI2 30.0-99.8), AGE 20 AND OLDER

August 8,

TABLE 10.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OBESITY: BODY MASS INDEX GE 30.0 & LE 99.8, AGE 20+ * -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	1013	24.9	1.5	(22.0, 27.8)	*	1616	24.9	1.2	(22.5, 27.3)
ALASKA	1287	20.0	1.6	(16.9, 23.1)	*	1366	26.3	1.7	(23.0, 29.6)
ARIZONA	1235	20.5	1.6	(17.4, 23.6)	*	1687	18.0	1.3	(15.5, 20.5)
ARKANSAS	1085	23.5	1.4	(20.8, 26.2)	*	1656	22.6	1.1	(20.4, 24.8)
CALIFORNIA	1646	21.5	1.2	(19.1, 23.9)	*	2372	23.7	1.2	(21.3, 26.1)
COLORADO	823	15.8	1.4	(13.1, 18.5)	*	1059	15.1	1.2	(12.7, 17.5)
CONNECTICUT	3064	19.1	0.8	(17.5, 20.7)	*	4011	17.3	0.7	(15.9, 18.7)
DELAWARE	1305	21.6	1.4	(18.9, 24.3)	*	1932	21.1	1.2	(18.7, 23.5)
DISTRICT OF CO	761	16.3	1.6	(13.2, 19.4)	*	983	23.8	1.6	(20.7, 26.9)
FLORIDA	1800	19.3	1.0	(17.3, 21.3)	*	2445	19.2	0.9	(17.4, 21.0)
GEORGIA	1646	22.4	1.3	(19.9, 24.9)	*	2533	23.8	1.0	(21.8, 25.8)
HAWAII	1866	19.9	1.4	(17.2, 22.6)	*	2427	16.7	1.1	(14.5, 18.9)
IDAHO	1987	21.9	1.1	(19.7, 24.1)	*	2475	20.2	0.9	(18.4, 22.0)
ILLINOIS	1562	20.8	1.2	(18.4, 23.2)	*	2150	22.6	1.0	(20.6, 24.6)
INDIANA	1553	25.5	1.2	(23.1, 27.9)	*	2203	24.8	1.0	(22.8, 26.8)
IOWA	1415	24.7	1.3	(22.2, 27.2)	*	1919	22.0	1.1	(19.8, 24.2)
KANSAS	1783	23.9	1.1	(21.7, 26.1)	*	2429	20.5	0.9	(18.7, 22.3)
KENTUCKY	2795	26.1	1.2	(23.7, 28.5)	*	4181	23.6	0.9	(21.8, 25.4)
LOUISIANA	1768	25.6	1.1	(23.4, 27.8)	*	2749	24.4	0.9	(22.6, 26.2)
MAINE	984	20.7	1.4	(18.0, 23.4)	*	1255	19.3	1.2	(16.9, 21.7)
MARYLAND	1737	19.8	1.1	(17.6, 22.0)	*	2395	21.0	1.1	(18.8, 23.2)
MASSACHUSETTS	3366	18.1	0.8	(16.5, 19.7)	*	4479	16.1	0.6	(14.9, 17.3)
MICHIGAN	1454	26.2	1.3	(23.7, 28.7)	*	2120	25.5	1.0	(23.5, 27.5)
MINNESOTA	1654	21.0	1.1	(18.8, 23.2)	*	2081	19.7	1.0	(17.7, 21.7)
MISSISSIPPI	1091	26.1	1.5	(23.2, 29.0)	*	1717	28.0	1.2	(25.6, 30.4)
MISSOURI	1581	25.8	1.5	(22.9, 28.7)	*	2296	21.8	1.1	(19.6, 24.0)
MONTANA	1372	20.3	1.4	(17.6, 23.0)	*	1771	18.2	1.2	(15.8, 20.6)
NEBRASKA	1336	22.8	1.3	(20.3, 25.3)	*	2019	19.6	1.0	(17.6, 21.6)
NEVADA	1263	22.7	2.2	(18.4, 27.0)	*	1145	16.8	1.6	(13.7, 19.9)
NEW HAMPSHIRE	1642	21.9	1.2	(19.5, 24.3)	*	2074	17.9	0.9	(16.1, 19.7)
NEW JERSEY	2300	19.7	1.1	(17.5, 21.9)	*	3199	20.3	1.0	(18.3, 22.3)
NEW MEXICO	1426	21.5	1.3	(19.0, 24.0)	*	1947	19.1	1.0	(17.1, 21.1)
NEW YORK	1530	21.2	1.2	(18.8, 23.6)	*	2041	20.5	1.1	(18.3, 22.7)
NORTH CAROLINA	2284	22.8	1.3	(20.3, 25.3)	*	3426	23.3	1.1	(21.1, 25.5)
NORTH DAKOTA	1033	22.0	1.4	(19.3, 24.7)	*	1316	19.2	1.2	(16.8, 21.6)
OHIO	1289	23.2	1.4	(20.5, 25.9)	*	1849	22.4	1.1	(20.2, 24.6)
OKLAHOMA	1679	25.2	1.3	(22.7, 27.7)	*	2440	21.1	1.0	(19.1, 23.1)
OREGON	1008	20.1	1.3	(17.6, 22.6)	*	1360	23.0	1.3	(20.5, 25.5)
PENNSYLVANIA	1434	22.2	1.2	(19.8, 24.6)	*	2005	23.0	1.1	(20.8, 25.2)
RHODE ISLAND	1468	18.2	1.2	(15.8, 20.6)	*	2296	18.0	1.0	(16.0, 20.0)
SOUTH CAROLINA	1208	23.4	1.4	(20.7, 26.1)	*	1740	22.7	1.2	(20.3, 25.1)
SOUTH DAKOTA	2099	24.1	1.0	(22.1, 26.1)	*	2686	19.1	0.8	(17.5, 20.7)
TENNESSEE	889	25.2	1.6	(22.1, 28.3)	*	1770	23.3	1.1	(21.1, 25.5)
TEXAS	2320	25.5	1.0	(23.5, 27.5)	*	3184	24.9	0.9	(23.1, 26.7)
UTAH	1575	20.7	1.4	(18.0, 23.4)	*	1815	19.0	1.2	(16.6, 21.4)
VERMONT	1732	19.2	1.0	(17.2, 21.2)	*	2299	17.0	0.8	(15.4, 18.6)
VIRGINIA	1187	22.2	1.3	(19.7, 24.7)	*	1541	20.6	1.2	(18.2, 23.0)
WASHINGTON	1679	21.4	1.1	(19.2, 23.6)	*	2225	18.4	0.9	(16.6, 20.2)
WEST VIRGINIA	1180	25.6	1.4	(22.9, 28.3)	*	1681	25.0	1.2	(22.6, 27.4)
WISCONSIN	1427	22.9	1.3	(20.4, 25.4)	*	1752	22.5	1.1	(20.3, 24.7)
WYOMING	1271	20.6	1.3	(18.1, 23.1)	*	1601	19.3	1.1	(17.1, 21.5)
GUAM	363	21.2	2.4	(16.5, 25.9)	*	449	22.8	2.3	(18.3, 27.3)
PUERTO RICO	1448	23.9	1.5	(21.0, 26.8)	*	2514	22.0	1.1	(19.8, 24.2)
VIRGIN ISLANDS	816	20.7	1.8	(17.2, 24.2)	*	1255	29.5	1.6	(26.4, 32.6)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 21.9
 RANGE = 15.8-26.2

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 21.1
 RANGE = 15.1-29.5

*AT RISK BY NEW BODY MASS INDEX(_BMI2 30.0-99.8), AGE 20 AND OLDER

August

TABLE 10.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 OBESITY: BODY MASS INDEX GE 30.0 & LE 99.8, AGE 20+ *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	24.5	24.9	0.4	1.8
ALASKA	21.5	22.9	1.4	6.8
ARIZONA	20.1	19.3	-0.8	-4.1
ARKANSAS	23.7	23.0	-0.7	-3.1
CALIFORNIA	20.5	22.6	2.1	10.5
COLORADO	14.4	15.5	1.1	7.6
CONNECTICUT	17.9	18.2	0.3	1.7
DELAWARE	17.2	21.4	4.2	24.6
DISTRICT OF CO	21.9	20.2	-1.7	-7.6
FLORIDA	19.0	19.2	0.3	1.3
GEORGIA	21.9	23.1	1.2	5.4
HAWAII	16.0	18.3	2.3	14.2
IDAHO	19.5	21.0	1.5	7.7
ILLINOIS	22.3	21.7	-0.6	-2.5
INDIANA	22.3	25.1	2.9	12.8
IOWA	22.0	23.4	1.4	6.3
KANSAS	21.3	22.2	0.9	4.4
KENTUCKY	23.5	24.8	1.3	5.6
LOUISIANA	24.2	25.0	0.8	3.5
MAINE	20.7	20.0	-0.6	-3.1
MARYLAND	20.8	20.4	-0.4	-1.9
MASSACHUSETTS	16.9	17.1	0.2	1.4
MICHIGAN	23.0	25.8	2.8	12.0
MINNESOTA	17.8	20.4	2.6	14.9
MISSISSIPPI	25.3	27.1	1.8	7.0
MISSOURI	22.5	23.8	1.3	5.6
MONTANA	16.4	19.2	2.8	16.8
NEBRASKA	21.4	21.2	-0.2	-0.9
NEVADA	18.3	19.9	1.6	8.9
NEW HAMPSHIRE	18.5	19.9	1.4	7.4
NEW JERSEY	19.0	20.0	1.0	5.4
NEW MEXICO	19.7	20.3	0.7	3.3
NEW YORK	17.9	20.8	2.9	16.1
NORTH CAROLINA	22.0	23.1	1.1	4.9
NORTH DAKOTA	20.9	20.6	-0.3	-1.4
OHIO	22.1	22.8	0.7	3.1
OKLAHOMA	19.9	23.2	3.3	16.6
OREGON	22.0	21.6	-0.4	-2.0
PENNSYLVANIA	21.7	22.6	0.9	4.3
RHODE ISLAND	17.5	18.1	0.6	3.3
SOUTH CAROLINA	22.7	23.0	0.3	1.4
SOUTH DAKOTA	20.3	21.6	1.3	6.5
TENNESSEE	23.1	24.2	1.1	4.7
TEXAS	23.5	25.2	1.7	7.3
UTAH	19.8	19.9	0.1	0.8
VERMONT	18.4	18.1	-0.3	-1.4
VIRGINIA	18.7	21.4	2.7	14.4
WASHINGTON	19.1	19.9	0.8	4.2
WEST VIRGINIA	23.8	25.3	1.6	6.5
WISCONSIN	20.3	22.7	2.4	11.6
WYOMING	18.4	19.9	1.5	8.4
GUAM	N/A	21.9	N/A	N/A
PUERTO RICO	22.8	22.9	0.1	0.5
VIRGIN ISLANDS	N/A	25.5	N/A	N/A

SUMMARY STATISTICS: yr2000

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 20.7
 RANGE = 14.4-25.3
 MEAN = 20.5

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 21.6
 RANGE = 15.5-27.1
 MEAN = 21.7

*AT RISK BY NEW BODY MASS INDEX(_BMI2 30.0-99.8), AGE 20 AND OLDER
 N/A - NOT AVAILABLE

August 8, 2002

TABLE 11.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PHYSICAL ACTIVITYA: NO LEISURE TIME PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2794	31.2	1.0	(29.2, 33.2)
ALASKA	2874	21.1	1.1	(18.9, 23.3)
ARIZONA	3258	21.9	1.1	(19.7, 24.1)
ARKANSAS	2926	31.5	1.0	(29.5, 33.5)
CALIFORNIA	4145	26.6	0.9	(24.8, 28.4)
COLORADO	2031	19.2	1.0	(17.2, 21.2)
CONNECTICUT	7745	24.0	0.6	(22.8, 25.2)
DELAWARE	3513	25.7	0.9	(23.9, 27.5)
DISTRICT OF CO	1886	24.2	1.2	(21.8, 26.6)
FLORIDA	4678	27.7	0.8	(26.1, 29.3)
GEORGIA	4523	27.3	0.8	(25.7, 28.9)
HAWAII	4500	18.9	0.8	(17.3, 20.5)
IDAHO	4831	21.0	0.7	(19.6, 22.4)
ILLINOIS	4009	26.5	0.8	(24.9, 28.1)
INDIANA	3990	26.2	0.8	(24.6, 27.8)
IOWA	3631	25.9	0.8	(24.3, 27.5)
KANSAS	4598	26.7	0.7	(25.3, 28.1)
KENTUCKY	7526	33.4	0.7	(32.0, 34.8)
LOUISIANA	4994	35.6	0.8	(34.0, 37.2)
MAINE	2418	23.2	0.9	(21.4, 25.0)
MARYLAND	4471	24.2	0.8	(22.6, 25.8)
MASSACHUSETTS	8626	22.8	0.5	(21.8, 23.8)
MICHIGAN	3826	23.4	0.8	(21.8, 25.0)
MINNESOTA	3965	17.1	0.7	(15.7, 18.5)
MISSISSIPPI	3040	33.4	1.0	(31.4, 35.4)
MISSOURI	4175	27.5	0.9	(25.7, 29.3)
MONTANA	3337	21.9	0.9	(20.1, 23.7)
NEBRASKA	3697	31.4	0.9	(29.6, 33.2)
NEVADA	2569	22.6	1.2	(20.2, 25.0)
NEW HAMPSHIRE	4067	19.5	0.7	(18.1, 20.9)
NEW JERSEY	6012	26.6	0.8	(25.0, 28.2)
NEW MEXICO	3620	25.8	0.8	(24.2, 27.4)
NEW YORK	3899	28.7	0.9	(26.9, 30.5)
NORTH CAROLINA	6205	26.4	0.9	(24.6, 28.2)
NORTH DAKOTA	2508	23.2	0.9	(21.4, 25.0)
OHIO	3430	26.2	0.9	(24.4, 28.0)
OKLAHOMA	4548	32.8	0.9	(31.0, 34.6)
OREGON	2536	20.8	0.9	(19.0, 22.6)
PENNSYLVANIA	3659	24.7	0.8	(23.1, 26.3)
RHODE ISLAND	4113	24.9	0.8	(23.3, 26.5)
SOUTH CAROLINA	3199	26.4	0.9	(24.6, 28.2)
SOUTH DAKOTA	5116	25.4	0.7	(24.0, 26.8)
TENNESSEE	2922	35.1	1.0	(33.1, 37.1)
TEXAS	5916	27.1	0.7	(25.7, 28.5)
UTAH	3656	16.5	0.8	(14.9, 18.1)
VERMONT	4301	20.3	0.7	(18.9, 21.7)
VIRGINIA	2936	23.2	0.9	(21.4, 25.0)
WASHINGTON	4205	17.1	0.6	(15.9, 18.3)
WEST VIRGINIA	3091	31.7	0.9	(29.9, 33.5)
WISCONSIN	3350	20.7	0.8	(19.1, 22.3)
WYOMING	3039	21.2	0.8	(19.6, 22.8)
GUAM	869	27.4	1.8	(23.9, 30.9)
PUERTO RICO	4236	49.2	1.1	(47.0, 51.4)
VIRGIN ISLANDS	2263	29.2	1.2	(26.8, 31.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 25.8 RANGE = 16.5-49.2

HP2010: HP 22.1

*AT RISK NO LEISURE TIME PHYSICAL ACTIVITY, IN THE PAST MONTH

August 8,

2002

TABLE 11.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR PHYSICAL ACTIVITY: NO LEISURE TIME PHYSICAL ACTIVITY* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1050	27.6	1.5	(24.7, 30.5)	*	1744	34.4	1.3	(31.9, 36.9)
ALASKA	1359	18.5	1.5	(15.6, 21.4)	*	1515	23.9	1.6	(20.8, 27.0)
ARIZONA	1338	17.6	1.5	(14.7, 20.5)	*	1920	26.0	1.6	(22.9, 29.1)
ARKANSAS	1120	29.4	1.5	(26.5, 32.3)	*	1806	33.4	1.3	(30.9, 35.9)
CALIFORNIA	1676	23.9	1.4	(21.2, 26.6)	*	2469	29.2	1.2	(26.8, 31.6)
COLORADO	863	17.3	1.4	(14.6, 20.0)	*	1168	21.1	1.4	(18.4, 23.8)
CONNECTICUT	3200	21.1	0.9	(19.3, 22.9)	*	4545	26.6	0.8	(25.0, 28.2)
DELAWARE	1361	21.7	1.4	(19.0, 24.4)	*	2152	29.2	1.2	(26.8, 31.6)
DISTRICT OF CO	782	20.7	1.9	(17.0, 24.4)	*	1104	27.3	1.6	(24.2, 30.4)
FLORIDA	1910	25.5	1.2	(23.1, 27.9)	*	2768	29.7	1.0	(27.7, 31.7)
GEORGIA	1721	23.5	1.3	(21.0, 26.0)	*	2802	30.7	1.1	(28.5, 32.9)
HAWAII	1937	15.8	1.1	(13.6, 18.0)	*	2563	22.2	1.1	(20.0, 24.4)
IDAHO	2070	19.5	1.1	(17.3, 21.7)	*	2761	22.4	0.9	(20.6, 24.2)
ILLINOIS	1642	23.2	1.2	(20.8, 25.6)	*	2367	29.4	1.1	(27.2, 31.6)
INDIANA	1612	23.4	1.1	(21.2, 25.6)	*	2378	28.9	1.0	(26.9, 30.9)
IOWA	1484	23.9	1.3	(21.4, 26.4)	*	2147	27.9	1.1	(25.7, 30.1)
KANSAS	1856	24.2	1.1	(22.0, 26.4)	*	2742	29.0	0.9	(27.2, 30.8)
KENTUCKY	2926	30.3	1.1	(28.1, 32.5)	*	4600	36.3	1.0	(34.3, 38.3)
LOUISIANA	1883	32.1	1.2	(29.7, 34.5)	*	3111	38.8	1.0	(36.8, 40.8)
MAINE	1027	22.7	1.4	(20.0, 25.4)	*	1391	23.7	1.2	(21.3, 26.1)
MARYLAND	1822	21.1	1.2	(18.7, 23.5)	*	2649	27.1	1.1	(24.9, 29.3)
MASSACHUSETTS	3560	19.6	0.8	(18.0, 21.2)	*	5066	25.8	0.7	(24.4, 27.2)
MICHIGAN	1518	20.3	1.1	(18.1, 22.5)	*	2308	26.2	1.1	(24.0, 28.4)
MINNESOTA	1699	15.8	1.0	(13.8, 17.8)	*	2266	18.4	0.9	(16.6, 20.2)
MISSISSIPPI	1147	27.6	1.5	(24.7, 30.5)	*	1893	38.5	1.2	(36.1, 40.9)
MISSOURI	1652	24.3	1.4	(21.6, 27.0)	*	2523	30.5	1.2	(28.1, 32.9)
MONTANA	1422	20.0	1.3	(17.5, 22.5)	*	1915	23.6	1.3	(21.1, 26.1)
NEBRASKA	1422	32.0	1.5	(29.1, 34.9)	*	2275	30.7	1.1	(28.5, 32.9)
NEVADA	1307	19.1	1.4	(16.4, 21.8)	*	1262	26.2	1.8	(22.7, 29.7)
NEW HAMPSHIRE	1728	17.6	1.0	(15.6, 19.6)	*	2339	21.3	0.9	(19.5, 23.1)
NEW JERSEY	2442	23.8	1.2	(21.4, 26.2)	*	3570	29.2	1.0	(27.2, 31.2)
NEW MEXICO	1502	24.2	1.3	(21.7, 26.7)	*	2118	27.4	1.1	(25.2, 29.6)
NEW YORK	1612	26.2	1.4	(23.5, 28.9)	*	2287	31.0	1.2	(28.6, 33.4)
NORTH CAROLINA	2382	23.0	1.3	(20.5, 25.5)	*	3823	29.5	1.1	(27.3, 31.7)
NORTH DAKOTA	1065	23.7	1.4	(21.0, 26.4)	*	1443	22.8	1.2	(20.4, 25.2)
OHIO	1349	22.9	1.4	(20.2, 25.6)	*	2081	29.1	1.2	(26.7, 31.5)
OKLAHOMA	1775	30.8	1.5	(27.9, 33.7)	*	2773	34.6	1.2	(32.2, 37.0)
OREGON	1053	18.8	1.3	(16.3, 21.3)	*	1483	22.6	1.2	(20.2, 25.0)
PENNSYLVANIA	1481	22.4	1.3	(19.9, 24.9)	*	2178	26.7	1.1	(24.5, 28.9)
RHODE ISLAND	1546	21.1	1.2	(18.7, 23.5)	*	2567	28.3	1.0	(26.3, 30.3)
SOUTH CAROLINA	1269	23.2	1.4	(20.5, 25.9)	*	1930	29.4	1.2	(27.0, 31.8)
SOUTH DAKOTA	2162	24.5	1.0	(22.5, 26.5)	*	2954	26.3	0.9	(24.5, 28.1)
TENNESSEE	954	30.8	1.7	(27.5, 34.1)	*	1968	38.9	1.2	(36.5, 41.3)
TEXAS	2437	24.6	1.0	(22.6, 26.6)	*	3479	29.4	0.9	(27.6, 31.2)
UTAH	1653	15.4	1.1	(13.2, 17.6)	*	2003	17.5	1.0	(15.5, 19.5)
VERMONT	1790	20.2	1.0	(18.2, 22.2)	*	2511	20.5	0.9	(18.7, 22.3)
VIRGINIA	1239	20.0	1.2	(17.6, 22.4)	*	1697	26.2	1.2	(23.8, 28.6)
WASHINGTON	1757	14.9	0.9	(13.1, 16.7)	*	2448	19.2	0.9	(17.4, 21.0)
WEST VIRGINIA	1227	28.2	1.4	(25.5, 30.9)	*	1864	34.8	1.2	(32.4, 37.2)
WISCONSIN	1478	19.4	1.2	(17.0, 21.8)	*	1872	22.0	1.1	(19.8, 24.2)
WYOMING	1319	20.4	1.2	(18.0, 22.8)	*	1720	22.1	1.1	(19.9, 24.3)
GUAM	381	24.4	2.6	(19.3, 29.5)	*	488	30.6	2.4	(25.9, 35.3)
PUERTO RICO	1520	43.1	1.7	(39.8, 46.4)	*	2716	54.6	1.3	(52.1, 57.1)
VIRGIN ISLANDS	876	24.9	1.8	(21.4, 28.4)	*	1387	32.8	1.6	(29.7, 35.9)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 23.1
 RANGE = 14.9-43.1

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 28.1
 RANGE = 17.5-54.6

*AT RISK NO LEISURE TIME PHYSICAL ACTIVITY, IN THE PAST MONTH

August 8,

TABLE 11.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 PHYSICAL ACTIVITY: NO LEISURE TIME PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	31.6	31.2	-0.4	-1.2
ALASKA	20.0	21.1	1.1	5.4
ARIZONA	34.2	21.9	-12.3	-36.0
ARKANSAS	28.1	31.5	3.4	12.3
CALIFORNIA	26.5	26.6	0.1	0.5
COLORADO	19.8	19.2	-0.6	-2.9
CONNECTICUT	25.2	24.0	-1.2	-4.6
DELAWARE	28.0	25.7	-2.3	-8.3
DISTRICT OF CO	20.8	24.2	3.4	16.5
FLORIDA	28.8	27.7	-1.1	-3.9
GEORGIA	29.0	27.3	-1.7	-5.8
HAWAII	23.2	18.9	-4.3	-18.5
IDAHO	19.8	21.0	1.2	6.2
ILLINOIS	30.9	26.5	-4.4	-14.2
INDIANA	25.4	26.2	0.8	3.3
IOWA	27.3	25.9	-1.4	-5.1
KANSAS	30.4	26.7	-3.7	-12.2
KENTUCKY	41.1	33.4	-7.7	-18.7
LOUISIANA	36.2	35.6	-0.6	-1.7
MAINE	27.2	23.2	-4.0	-14.7
MARYLAND	24.2	24.2	0.0	-0.1
MASSACHUSETTS	24.6	22.8	-1.8	-7.3
MICHIGAN	22.9	23.4	0.5	2.3
MINNESOTA	24.8	17.1	-7.7	-31.0
MISSISSIPPI	33.3	33.4	0.1	0.2
MISSOURI	28.8	27.5	-1.3	-4.6
MONTANA	23.3	21.9	-1.4	-6.0
NEBRASKA	29.6	31.4	1.8	6.1
NEVADA	25.0	22.6	-2.3	-9.4
NEW HAMPSHIRE	26.7	19.5	-7.2	-26.8
NEW JERSEY	28.6	26.6	-2.0	-7.1
NEW MEXICO	24.4	25.8	1.4	5.8
NEW YORK	29.4	28.7	-0.7	-2.2
NORTH CAROLINA	30.4	26.4	-4.0	-13.3
NORTH DAKOTA	24.3	23.2	-1.1	-4.7
OHIO	31.3	26.2	-5.1	-16.2
OKLAHOMA	34.4	32.8	-1.6	-4.7
OREGON	20.1	20.8	0.7	3.7
PENNSYLVANIA	23.0	24.7	1.7	7.5
RHODE ISLAND	27.5	24.9	-2.6	-9.6
SOUTH CAROLINA	28.1	26.4	-1.7	-6.1
SOUTH DAKOTA	26.7	25.4	-1.3	-4.7
TENNESSEE	32.7	35.1	2.4	7.3
TEXAS	28.6	27.1	-1.4	-5.1
UTAH	15.5	16.5	1.0	6.3
VERMONT	23.2	20.3	-2.9	-12.6
VIRGINIA	25.0	23.2	-1.8	-7.2
WASHINGTON	16.9	17.1	0.2	1.2
WEST VIRGINIA	33.6	31.7	-1.9	-5.7
WISCONSIN	22.1	20.7	-1.4	-6.5
WYOMING	22.6	21.2	-1.4	-6.2
GUAM	N/A	27.4	N/A	N/A
PUERTO RICO	54.1	49.2	-4.9	-9.0
VIRGIN ISLANDS	N/A	29.2	N/A	N/A

SUMMARY STATISTICS: yr2000

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 26.9
 RANGE = 15.5-54.1

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 25.8
 RANGE = 16.5-49.2

*AT RISK NO LEISURE TIME PHYSICAL ACTIVITY, IN THE PAST MONTH
 N/A - NOT AVAILABLE

August 8, 2002

TABLE 12.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SMOKING: CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2789	23.8	0.9	(22.0, 25.6)
ALASKA	2866	26.2	1.2	(23.8, 28.6)
ARIZONA	3256	21.5	1.0	(19.5, 23.5)
ARKANSAS	2918	25.5	0.9	(23.7, 27.3)
CALIFORNIA	4247	17.2	0.7	(15.8, 18.6)
COLORADO	2025	22.3	1.0	(20.3, 24.3)
CONNECTICUT	7724	20.6	0.6	(19.4, 21.8)
DELAWARE	3508	25.0	1.0	(23.0, 27.0)
DISTRICT OF CO	1882	20.8	1.1	(18.6, 23.0)
FLORIDA	4668	22.4	0.7	(21.0, 23.8)
GEORGIA	4508	23.7	0.8	(22.1, 25.3)
HAWAII	4496	20.5	0.8	(18.9, 22.1)
IDAHO	4823	19.6	0.7	(18.2, 21.0)
ILLINOIS	4001	23.7	0.8	(22.1, 25.3)
INDIANA	3985	27.4	0.8	(25.8, 29.0)
IOWA	3621	22.1	0.8	(20.5, 23.7)
KANSAS	4592	22.2	0.7	(20.8, 23.6)
KENTUCKY	7505	30.9	0.8	(29.3, 32.5)
LOUISIANA	4973	24.6	0.7	(23.2, 26.0)
MAINE	2412	23.9	1.0	(21.9, 25.9)
MARYLAND	4450	21.1	0.9	(19.3, 22.9)
MASSACHUSETTS	8590	19.5	0.5	(18.5, 20.5)
MICHIGAN	3820	25.6	0.8	(24.0, 27.2)
MINNESOTA	3952	22.2	0.8	(20.6, 23.8)
MISSISSIPPI	3025	25.3	0.9	(23.5, 27.1)
MISSOURI	4164	25.9	0.9	(24.1, 27.7)
MONTANA	3330	21.9	1.0	(19.9, 23.9)
NEBRASKA	3688	20.2	0.8	(18.6, 21.8)
NEVADA	2567	26.9	1.2	(24.5, 29.3)
NEW HAMPSHIRE	4054	24.1	0.8	(22.5, 25.7)
NEW JERSEY	5995	21.1	0.7	(19.7, 22.5)
NEW MEXICO	3609	23.8	0.9	(22.0, 25.6)
NEW YORK	3882	23.2	0.8	(21.6, 24.8)
NORTH CAROLINA	6181	25.7	0.9	(23.9, 27.5)
NORTH DAKOTA	2506	22.1	0.9	(20.3, 23.9)
OHIO	3425	27.6	0.9	(25.8, 29.4)
OKLAHOMA	4539	28.7	0.9	(26.9, 30.5)
OREGON	2534	20.5	0.9	(18.7, 22.3)
PENNSYLVANIA	3652	24.5	0.9	(22.7, 26.3)
RHODE ISLAND	4096	23.9	0.8	(22.3, 25.5)
SOUTH CAROLINA	3190	26.0	0.9	(24.2, 27.8)
SOUTH DAKOTA	5103	22.3	0.7	(20.9, 23.7)
TENNESSEE	2916	24.4	1.0	(22.4, 26.4)
TEXAS	5906	22.4	0.6	(21.2, 23.6)
UTAH	3653	13.2	0.7	(11.8, 14.6)
VERMONT	4291	22.4	0.7	(21.0, 23.8)
VIRGINIA	2928	22.5	0.9	(20.7, 24.3)
WASHINGTON	4191	22.5	0.7	(21.1, 23.9)
WEST VIRGINIA	3083	28.2	0.9	(26.4, 30.0)
WISCONSIN	3344	23.6	0.9	(21.8, 25.4)
WYOMING	3032	22.2	0.8	(20.6, 23.8)
GUAM	870	31.2	1.9	(27.5, 34.9)
PUERTO RICO	4233	12.5	0.7	(11.1, 13.9)
VIRGIN ISLANDS	2259	9.6	0.7	(8.2, 11.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 22.9 RANGE = 9.6-31.2

HP2010: LHI, HP 27.1a

* AT RISK FOR CURRENT SMOKING

August 8, 2002

TABLE 12.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SMOKING: CURRENT SMOKERS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1048	25.8	1.5	(22.9, 28.7)	*	1741	22.1	1.1 (19.9, 24.3)
ALASKA	1356	26.4	1.8	(22.9, 29.9)	*	1510	25.9	1.6 (22.8, 29.0)
ARIZONA	1339	23.1	1.7	(19.8, 26.4)	*	1917	20.1	1.3 (17.6, 22.6)
ARKANSAS	1118	27.2	1.5	(24.3, 30.1)	*	1800	24.0	1.1 (21.8, 26.2)
CALIFORNIA	1719	20.6	1.2	(18.2, 23.0)	*	2528	14.0	0.8 (12.4, 15.6)
COLORADO	863	23.7	1.6	(20.6, 26.8)	*	1162	20.9	1.3 (18.4, 23.4)
CONNECTICUT	3195	21.2	0.9	(19.4, 23.0)	*	4529	20.1	0.7 (18.7, 21.5)
DELAWARE	1357	28.2	1.6	(25.1, 31.3)	*	2151	22.1	1.1 (19.9, 24.3)
DISTRICT OF CO	780	24.9	1.9	(21.2, 28.6)	*	1102	17.3	1.3 (14.8, 19.8)
FLORIDA	1910	25.6	1.2	(23.2, 28.0)	*	2758	19.4	0.9 (17.6, 21.2)
GEORGIA	1710	25.7	1.3	(23.2, 28.2)	*	2798	21.8	0.9 (20.0, 23.6)
HAWAII	1935	24.7	1.4	(22.0, 27.4)	*	2561	16.3	0.9 (14.5, 18.1)
IDAHO	2064	21.0	1.1	(18.8, 23.2)	*	2759	18.2	0.9 (16.4, 20.0)
ILLINOIS	1638	26.7	1.3	(24.2, 29.2)	*	2363	20.9	0.9 (19.1, 22.7)
INDIANA	1610	29.7	1.2	(27.3, 32.1)	*	2375	25.3	1.0 (23.3, 27.3)
IOWA	1479	24.2	1.3	(21.7, 26.7)	*	2142	20.3	1.0 (18.3, 22.3)
KANSAS	1853	22.5	1.1	(20.3, 24.7)	*	2739	22.0	0.9 (20.2, 23.8)
KENTUCKY	2917	31.9	1.3	(29.4, 34.4)	*	4588	30.0	1.0 (28.0, 32.0)
LOUISIANA	1878	28.7	1.2	(26.3, 31.1)	*	3095	21.0	0.8 (19.4, 22.6)
MAINE	1025	27.0	1.7	(23.7, 30.3)	*	1387	21.1	1.2 (18.7, 23.5)
MARYLAND	1814	24.6	1.5	(21.7, 27.5)	*	2636	17.9	1.0 (15.9, 19.9)
MASSACHUSETTS	3553	20.4	0.8	(18.8, 22.0)	*	5037	18.7	0.7 (17.3, 20.1)
MICHIGAN	1515	26.7	1.3	(24.2, 29.2)	*	2305	24.5	1.0 (22.5, 26.5)
MINNESOTA	1691	24.8	1.2	(22.4, 27.2)	*	2261	19.6	0.9 (17.8, 21.4)
MISSISSIPPI	1137	29.3	1.6	(26.2, 32.4)	*	1888	21.7	1.1 (19.5, 23.9)
MISSOURI	1645	27.6	1.4	(24.9, 30.3)	*	2519	24.4	1.1 (22.2, 26.6)
MONTANA	1419	21.7	1.5	(18.8, 24.6)	*	1911	22.1	1.3 (19.6, 24.6)
NEBRASKA	1417	20.6	1.2	(18.2, 23.0)	*	2271	19.8	1.0 (17.8, 21.8)
NEVADA	1307	27.8	1.8	(24.3, 31.3)	*	1260	26.0	1.7 (22.7, 29.3)
NEW HAMPSHIRE	1723	25.6	1.2	(23.2, 28.0)	*	2331	22.7	1.0 (20.7, 24.7)
NEW JERSEY	2436	21.6	1.1	(19.4, 23.8)	*	3559	20.7	0.9 (18.9, 22.5)
NEW MEXICO	1497	27.9	1.4	(25.2, 30.6)	*	2112	20.0	1.0 (18.0, 22.0)
NEW YORK	1607	26.1	1.3	(23.6, 28.6)	*	2275	20.7	1.0 (18.7, 22.7)
NORTH CAROLINA	2369	28.6	1.4	(25.9, 31.3)	*	3812	23.1	1.1 (20.9, 25.3)
NORTH DAKOTA	1064	24.6	1.4	(21.9, 27.3)	*	1442	19.7	1.1 (17.5, 21.9)
OHIO	1347	29.1	1.5	(26.2, 32.0)	*	2078	26.3	1.1 (24.1, 28.5)
OKLAHOMA	1768	31.1	1.5	(28.2, 34.0)	*	2771	26.4	1.1 (24.2, 28.6)
OREGON	1052	21.4	1.4	(18.7, 24.1)	*	1482	19.7	1.1 (17.5, 21.9)
PENNSYLVANIA	1479	26.3	1.4	(23.6, 29.0)	*	2173	22.8	1.0 (20.8, 24.8)
RHODE ISLAND	1544	25.8	1.4	(23.1, 28.5)	*	2552	22.1	1.0 (20.1, 24.1)
SOUTH CAROLINA	1261	27.9	1.4	(25.2, 30.6)	*	1929	24.3	1.1 (22.1, 26.5)
SOUTH DAKOTA	2158	23.2	1.1	(21.0, 25.4)	*	2945	21.4	0.9 (19.6, 23.2)
TENNESSEE	951	26.1	1.6	(23.0, 29.2)	*	1965	22.8	1.1 (20.6, 25.0)
TEXAS	2429	25.1	1.0	(23.1, 27.1)	*	3477	19.8	0.8 (18.2, 21.4)
UTAH	1651	14.6	1.1	(12.4, 16.8)	*	2002	12.0	0.9 (10.2, 13.8)
VERMONT	1786	24.4	1.2	(22.0, 26.8)	*	2505	20.5	0.9 (18.7, 22.3)
VIRGINIA	1235	23.5	1.3	(21.0, 26.0)	*	1693	21.6	1.2 (19.2, 24.0)
WASHINGTON	1752	24.5	1.1	(22.3, 26.7)	*	2439	20.5	0.9 (18.7, 22.3)
WEST VIRGINIA	1222	28.9	1.5	(26.0, 31.8)	*	1861	27.6	1.2 (25.2, 30.0)
WISCONSIN	1474	25.3	1.3	(22.8, 27.8)	*	1870	21.9	1.1 (19.7, 24.1)
WYOMING	1315	22.5	1.3	(20.0, 25.0)	*	1717	21.8	1.1 (19.6, 24.0)
GUAM	382	38.6	2.9	(32.9, 44.3)	*	488	23.0	2.2 (18.7, 27.3)
PUERTO RICO	1517	17.4	1.3	(14.9, 19.9)	*	2716	8.2	0.7 (6.8, 9.6)
VIRGIN ISLANDS	874	13.1	1.4	(10.4, 15.8)	*	1385	6.7	0.7 (5.3, 8.1)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 25.5
 RANGE = 13.1-38.6

* AT RISK FOR CURRENT SMOKING

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 21.3
 RANGE = 6.7-30.0

August 8, 2002

TABLE 12.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 2000 TO 2001
 SMOKING: CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	2000 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	25.2	23.8	-1.4	-5.7
ALASKA	25.0	26.2	1.2	4.7
ARIZONA	18.6	21.5	2.9	15.3
ARKANSAS	25.1	25.5	0.4	1.8
CALIFORNIA	17.2	17.2	-0.0	-0.2
COLORADO	20.0	22.3	2.3	11.5
CONNECTICUT	19.9	20.6	0.7	3.5
DELAWARE	22.9	25.0	2.1	9.4
DISTRICT OF CO	20.9	20.8	-0.1	-0.5
FLORIDA	23.2	22.4	-0.8	-3.4
GEORGIA	23.5	23.7	0.2	0.8
HAWAII	19.7	20.5	0.8	4.3
IDAHO	22.3	19.6	-2.7	-12.0
ILLINOIS	22.3	23.7	1.4	6.1
INDIANA	26.9	27.4	0.5	1.8
IOWA	23.2	22.1	-1.1	-4.7
KANSAS	21.1	22.2	1.1	5.5
KENTUCKY	30.5	30.9	0.4	1.3
LOUISIANA	24.1	24.6	0.5	2.2
MAINE	23.8	23.9	0.1	0.5
MARYLAND	20.5	21.1	0.6	3.0
MASSACHUSETTS	20.0	19.5	-0.4	-2.3
MICHIGAN	24.1	25.6	1.5	6.1
MINNESOTA	19.8	22.2	2.4	12.2
MISSISSIPPI	23.5	25.3	1.8	7.8
MISSOURI	27.2	25.9	-1.3	-4.7
MONTANA	18.8	21.9	3.1	16.2
NEBRASKA	21.2	20.2	-1.0	-4.9
NEVADA	29.0	26.9	-2.1	-7.3
NEW HAMPSHIRE	25.3	24.1	-1.2	-4.8
NEW JERSEY	21.0	21.1	0.1	0.6
NEW MEXICO	23.6	23.8	0.2	0.9
NEW YORK	21.6	23.2	1.6	7.5
NORTH CAROLINA	26.1	25.7	-0.4	-1.5
NORTH DAKOTA	23.2	22.1	-1.1	-4.8
OHIO	26.2	27.6	1.4	5.2
OKLAHOMA	23.3	28.7	5.4	23.1
OREGON	20.8	20.5	-0.3	-1.2
PENNSYLVANIA	24.3	24.5	0.2	0.7
RHODE ISLAND	23.4	23.9	0.5	2.1
SOUTH CAROLINA	24.9	26.0	1.1	4.4
SOUTH DAKOTA	21.9	22.3	0.4	1.9
TENNESSEE	25.7	24.4	-1.3	-4.9
TEXAS	21.9	22.4	0.5	2.3
UTAH	12.9	13.2	0.3	2.7
VERMONT	21.5	22.4	0.9	4.4
VIRGINIA	21.4	22.5	1.1	5.1
WASHINGTON	20.7	22.5	1.8	8.5
WEST VIRGINIA	26.1	28.2	2.1	8.0
WISCONSIN	24.1	23.6	-0.5	-2.2
WYOMING	23.8	22.2	-1.6	-6.6
GUAM	N/A	31.2	N/A	N/A
PUERTO RICO	13.1	12.5	-0.6	-4.9
VIRGIN ISLANDS	N/A	9.6	N/A	N/A

SUMMARY STATISTICS: yr2000

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 23.2
 RANGE = 12.9-30.5

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 22.9
 RANGE = 9.6-31.2

* AT RISK FOR CURRENT SMOKING
 N/A - NOT AVAILABLE

August 8, 2002

TABLE 13.1 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKING: BINGE DRINKING DURING THE PAST 30 DAYS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2739	11.6	0.8	(10.0, 13.2)
ALASKA	2812	18.2	1.2	(15.8, 20.6)
ARIZONA	3213	16.8	1.0	(14.8, 18.8)
ARKANSAS	2890	11.3	0.7	(9.9, 12.7)
CALIFORNIA	3919	15.5	0.8	(13.9, 17.1)
COLORADO	2005	16.7	1.0	(14.7, 18.7)
CONNECTICUT	7614	13.8	0.5	(12.8, 14.8)
DELAWARE	3447	15.7	0.9	(13.9, 17.5)
DISTRICT OF CO	1854	14.8	1.1	(12.6, 17.0)
FLORIDA	4587	12.0	0.6	(10.8, 13.2)
GEORGIA	4424	11.9	0.7	(10.5, 13.3)
HAWAII	4452	10.4	0.7	(9.0, 11.8)
IDAHO	4788	12.8	0.6	(11.6, 14.0)
ILLINOIS	3979	17.3	0.7	(15.9, 18.7)
INDIANA	3967	13.8	0.6	(12.6, 15.0)
IOWA	3580	16.2	0.8	(14.6, 17.8)
KANSAS	4493	14.7	0.6	(13.5, 15.9)
KENTUCKY	7356	8.7	0.6	(7.5, 9.9)
LOUISIANA	4812	13.8	0.6	(12.6, 15.0)
MAINE	2384	15.4	0.9	(13.6, 17.2)
MARYLAND	4409	11.9	0.6	(10.7, 13.1)
MASSACHUSETTS	8485	18.1	0.5	(17.1, 19.1)
MICHIGAN	3766	18.0	0.8	(16.4, 19.6)
MINNESOTA	3901	19.6	0.8	(18.0, 21.2)
MISSISSIPPI	2967	11.8	0.8	(10.2, 13.4)
MISSOURI	4129	14.1	0.8	(12.5, 15.7)
MONTANA	3264	16.7	1.0	(14.7, 18.7)
NEBRASKA	3604	14.6	0.8	(13.0, 16.2)
NEVADA	2537	16.7	1.1	(14.5, 18.9)
NEW HAMPSHIRE	4005	15.8	0.7	(14.4, 17.2)
NEW JERSEY	5903	13.5	0.6	(12.3, 14.7)
NEW MEXICO	3578	15.8	0.8	(14.2, 17.4)
NEW YORK	3833	14.4	0.7	(13.0, 15.8)
NORTH CAROLINA	6099	9.8	0.6	(8.6, 11.0)
NORTH DAKOTA	2482	22.3	1.0	(20.3, 24.3)
OHIO	3389	16.2	0.9	(14.4, 18.0)
OKLAHOMA	4428	11.0	0.8	(9.4, 12.6)
OREGON	2512	14.7	0.8	(13.1, 16.3)
PENNSYLVANIA	3625	15.6	0.8	(14.0, 17.2)
RHODE ISLAND	4063	15.1	0.7	(13.7, 16.5)
SOUTH CAROLINA	3155	12.3	0.7	(10.9, 13.7)
SOUTH DAKOTA	5060	18.5	0.7	(17.1, 19.9)
TENNESSEE	2817	6.8	0.7	(5.4, 8.2)
TEXAS	5837	15.1	0.6	(13.9, 16.3)
UTAH	3644	9.7	0.6	(8.5, 10.9)
VERMONT	4256	15.7	0.7	(14.3, 17.1)
VIRGINIA	2897	14.3	0.9	(12.5, 16.1)
WASHINGTON	4174	14.9	0.6	(13.7, 16.1)
WEST VIRGINIA	3061	9.4	0.7	(8.0, 10.8)
WISCONSIN	3324	25.7	0.9	(23.9, 27.5)
WYOMING	3016	16.0	0.8	(14.4, 17.6)
GUAM	855	18.1	1.6	(15.0, 21.2)
PUERTO RICO	4181	11.3	0.7	(9.9, 12.7)
VIRGIN ISLANDS	2201	11.1	0.8	(9.5, 12.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 54 MEDIAN = 14.8 RANGE = 6.8-25.7

HP2010: LHI, HP 26.11c

* AT RISK FOR BINGE DRINKING

August 8, 2002

**TABLE 13.2 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKING: BINGE DRINKING DURING THE PAST 30 DAYS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED**

PARTICIPANT	MALES				FEMALES			
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1026	20.0	1.5	(17.1, 22.9)	*	1713	4.2	0.5 (3.2, 5.2)
ALASKA	1325	24.8	1.9	(21.1, 28.5)	*	1487	11.0	1.5 (8.1, 13.9)
ARIZONA	1312	25.1	1.7	(21.8, 28.4)	*	1901	9.0	1.0 (7.0, 11.0)
ARKANSAS	1101	18.3	1.4	(15.6, 21.0)	*	1789	4.9	0.6 (3.7, 6.1)
CALIFORNIA	1572	24.4	1.4	(21.7, 27.1)	*	2347	6.9	0.6 (5.7, 8.1)
COLORADO	850	26.5	1.7	(23.2, 29.8)	*	1155	7.2	0.8 (5.6, 8.8)
CONNECTICUT	3132	21.1	0.9	(19.3, 22.9)	*	4482	7.2	0.5 (6.2, 8.2)
DELAWARE	1327	25.6	1.6	(22.5, 28.7)	*	2120	6.8	0.7 (5.4, 8.2)
DISTRICT OF CO	767	22.0	2.0	(18.1, 25.9)	*	1087	8.7	1.1 (6.5, 10.9)
FLORIDA	1863	17.9	1.0	(15.9, 19.9)	*	2724	6.6	0.6 (5.4, 7.8)
GEORGIA	1666	18.6	1.3	(16.1, 21.1)	*	2758	5.7	0.6 (4.5, 6.9)
HAWAII	1908	16.9	1.2	(14.5, 19.3)	*	2544	3.9	0.5 (2.9, 4.9)
IDAHO	2041	19.9	1.1	(17.7, 22.1)	*	2747	6.0	0.5 (5.0, 7.0)
ILLINOIS	1627	24.9	1.3	(22.4, 27.4)	*	2352	10.3	0.7 (8.9, 11.7)
INDIANA	1600	22.1	1.2	(19.7, 24.5)	*	2367	6.2	0.5 (5.2, 7.2)
IOWA	1452	24.7	1.3	(22.2, 27.2)	*	2128	8.6	0.8 (7.0, 10.2)
KANSAS	1805	23.1	1.1	(20.9, 25.3)	*	2688	6.9	0.6 (5.7, 8.1)
KENTUCKY	2832	14.8	1.0	(12.8, 16.8)	*	4524	3.2	0.5 (2.2, 4.2)
LOUISIANA	1781	22.7	1.1	(20.5, 24.9)	*	3031	6.0	0.5 (5.0, 7.0)
MAINE	1007	24.9	1.6	(21.8, 28.0)	*	1377	6.8	0.8 (5.2, 8.4)
MARYLAND	1792	18.8	1.2	(16.4, 21.2)	*	2617	5.6	0.5 (4.6, 6.6)
MASSACHUSETTS	3475	27.1	0.9	(25.3, 28.9)	*	5010	10.0	0.6 (8.8, 11.2)
MICHIGAN	1479	27.0	1.4	(24.3, 29.7)	*	2287	9.9	0.8 (8.3, 11.5)
MINNESOTA	1659	29.9	1.3	(27.4, 32.4)	*	2242	10.0	0.7 (8.6, 11.4)
MISSISSIPPI	1102	19.6	1.5	(16.7, 22.5)	*	1865	5.0	0.6 (3.8, 6.2)
MISSOURI	1619	22.1	1.4	(19.4, 24.8)	*	2510	6.9	0.7 (5.5, 8.3)
MONTANA	1379	25.6	1.7	(22.3, 28.9)	*	1885	8.3	1.0 (6.3, 10.3)
NEBRASKA	1374	22.1	1.4	(19.4, 24.8)	*	2230	7.7	0.7 (6.3, 9.1)
NEVADA	1284	25.3	1.8	(21.8, 28.8)	*	1253	8.1	1.0 (6.1, 10.1)
NEW HAMPSHIRE	1689	23.8	1.2	(21.4, 26.2)	*	2316	8.3	0.7 (6.9, 9.7)
NEW JERSEY	2381	20.5	1.1	(18.3, 22.7)	*	3522	7.3	0.6 (6.1, 8.5)
NEW MEXICO	1477	26.6	1.4	(23.9, 29.3)	*	2101	5.6	0.6 (4.4, 6.8)
NEW YORK	1577	21.2	1.2	(18.8, 23.6)	*	2256	8.2	0.8 (6.6, 9.8)
NORTH CAROLINA	2324	15.6	1.1	(13.4, 17.8)	*	3775	4.5	0.5 (3.5, 5.5)
NORTH DAKOTA	1049	34.1	1.6	(31.0, 37.2)	*	1433	10.9	0.9 (9.1, 12.7)
OHIO	1324	24.7	1.5	(21.8, 27.6)	*	2065	8.7	0.7 (7.3, 10.1)
OKLAHOMA	1701	18.5	1.4	(15.8, 21.2)	*	2727	4.2	0.5 (3.2, 5.2)
OREGON	1044	22.7	1.4	(20.0, 25.4)	*	1468	7.1	0.8 (5.5, 8.7)
PENNSYLVANIA	1462	25.2	1.4	(22.5, 27.9)	*	2163	7.0	0.7 (5.6, 8.4)
RHODE ISLAND	1519	21.4	1.3	(18.9, 23.9)	*	2544	9.6	0.8 (8.0, 11.2)
SOUTH CAROLINA	1244	19.5	1.2	(17.1, 21.9)	*	1911	5.9	0.7 (4.5, 7.3)
SOUTH DAKOTA	2129	27.6	1.2	(25.2, 30.0)	*	2931	10.0	0.7 (8.6, 11.4)
TENNESSEE	914	12.1	1.3	(9.6, 14.6)	*	1903	2.2	0.4 (1.4, 3.0)
TEXAS	2375	23.7	1.0	(21.7, 25.7)	*	3462	7.1	0.5 (6.1, 8.1)
UTAH	1643	15.2	1.1	(13.0, 17.4)	*	2001	4.6	0.6 (3.4, 5.8)
VERMONT	1767	22.8	1.2	(20.4, 25.2)	*	2489	9.2	0.7 (7.8, 10.6)
VIRGINIA	1216	20.9	1.4	(18.2, 23.6)	*	1681	8.1	1.0 (6.1, 10.1)
WASHINGTON	1738	23.3	1.1	(21.1, 25.5)	*	2436	7.0	0.6 (5.8, 8.2)
WEST VIRGINIA	1211	14.9	1.2	(12.5, 17.3)	*	1850	4.5	0.6 (3.3, 5.7)
WISCONSIN	1462	36.9	1.5	(34.0, 39.8)	*	1862	15.2	1.0 (13.2, 17.2)
WYOMING	1306	23.9	1.3	(21.4, 26.4)	*	1710	8.4	0.7 (7.0, 9.8)
GUAM	373	29.0	2.6	(23.9, 34.1)	*	482	6.1	1.3 (3.6, 8.6)
PUERTO RICO	1483	20.4	1.4	(17.7, 23.1)	*	2698	3.2	0.5 (2.2, 4.2)
VIRGIN ISLANDS	836	19.5	1.7	(16.2, 22.8)	*	1365	4.5	0.7 (3.1, 5.9)

SUMMARY STATISTICS: MALES

NUMBER OF PARTICIPANTS = 54
MEDIAN = 22.7
RANGE = 12.1-36.9

* AT RISK FOR BINGE DRINKING

SUMMARY STATISTICS: FEMALES

NUMBER OF PARTICIPANTS = 54
MEDIAN = 7.0
RANGE = 2.2-15.2

August 8, 2002

TABLE 13.3 2001 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1999 TO 2001
 BINGE DRINKING: BINGE DRINKING DURING THE PAST 30 DAYS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1999 PERCENT	2001 PERCENT	DIFFERENCE	PERCENT DIFFERENCE
ALABAMA	11.7	11.6	-0.1	-0.9
ALASKA	18.9	18.2	-0.7	-3.9
ARIZONA	8.8	16.8	8.0	90.7
ARKANSAS	10.3	11.3	1.0	9.5
CALIFORNIA	15.5	15.5	-0.0	-0.2
COLORADO	17.2	16.7	-0.5	-3.1
CONNECTICUT	14.0	13.8	-0.2	-1.6
DELAWARE	18.9	15.7	-3.2	-17.1
DISTRICT OF CO	13.0	14.8	1.9	14.3
FLORIDA	12.8	12.0	-0.8	-6.5
GEORGIA	12.5	11.9	-0.6	-4.6
HAWAII	14.0	10.4	-3.6	-25.7
IDAHO	14.7	12.8	-1.9	-13.0
ILLINOIS	19.7	17.3	-2.4	-12.0
INDIANA	19.1	13.8	-5.3	-27.9
IOWA	18.3	16.2	-2.1	-11.5
KANSAS	11.8	14.7	3.0	25.1
KENTUCKY	9.8	8.7	-1.1	-11.4
LOUISIANA	15.0	13.8	-1.2	-7.8
MAINE	14.8	15.4	0.6	4.1
MARYLAND	15.9	11.9	-4.0	-25.2
MASSACHUSETTS	17.4	18.1	0.7	3.9
MICHIGAN	19.0	18.0	-1.0	-5.4
MINNESOTA	16.3	19.6	3.3	20.2
MISSISSIPPI	12.1	11.8	-0.3	-2.2
MISSOURI	16.4	14.1	-2.3	-14.0
MONTANA	17.6	16.7	-0.9	-5.0
NEBRASKA	16.6	14.6	-2.0	-12.1
NEVADA	21.0	16.7	-4.3	-20.4
NEW HAMPSHIRE	20.0	15.8	-4.2	-21.1
NEW JERSEY	12.3	13.5	1.2	9.9
NEW MEXICO	15.0	15.8	0.9	5.7
NEW YORK	13.9	14.4	0.5	3.6
NORTH CAROLINA	12.0	9.8	-2.2	-18.1
NORTH DAKOTA	19.7	22.3	2.6	13.3
OHIO	12.1	16.2	4.1	33.8
OKLAHOMA	8.1	11.0	2.9	35.6
OREGON	14.9	14.7	-0.2	-1.0
PENNSYLVANIA	15.9	15.6	-0.3	-1.7
RHODE ISLAND	15.6	15.1	-0.5	-3.0
SOUTH CAROLINA	12.3	12.3	-0.0	-0.1
SOUTH DAKOTA	17.4	18.5	1.1	6.1
TENNESSEE	7.7	6.8	-0.9	-11.6
TEXAS	17.3	15.1	-2.2	-12.7
UTAH	10.2	9.7	-0.5	-5.3
VERMONT	17.4	15.7	-1.7	-9.5
VIRGINIA	12.7	14.3	1.6	12.2
WASHINGTON	14.4	14.9	0.5	3.7
WEST VIRGINIA	8.6	9.4	0.8	9.9
WISCONSIN	27.0	25.7	-1.3	-4.7
WYOMING	17.3	16.0	-1.3	-7.4
GUAM	N/A	18.1	N/A	N/A
PUERTO RICO	10.6	11.3	0.7	6.2
VIRGIN ISLANDS	N/A	11.1	N/A	N/A

SUMMARY STATISTICS: yr1999

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 14.9
 RANGE = 7.7-27.0

SUMMARY STATISTICS: yr2001

NUMBER OF PARTICIPANTS = 54
 MEDIAN = 14.8
 RANGE = 6.8-25.7

* AT RISK FOR BINGE DRINKING
 N/A - NOT AVAILABLE

August 8, 2002