

June 18, 1999

Dear Colleague:

Information related to the 1998 BRFSS summary prevalence report of selected risk factors and preventive health measures from the core and modules is attached.

This report consists of four parts:

An introduction that contains:

An overview of the 1998 summary prevalence report.

An explanation of the tables for 1998 Behavioral Risk Factor Surveillance System (BRFSS) prevalence data.

Definitions of BRFSS risk factors and preventive health measures. When appropriate, Year 2000 Objectives are identified.

Analysis of selected questions from the 1998 BRFSS questionnaire. The set of analyses is divided into three sections:

Section I - selected risk factors and preventive health measures in the BRFSS core.

Section II - selected risk factors and preventive health measures from modules used by at least 13 states

Section III - preventive health measures that concern women only.

As soon as your schedule permits, please examine the enclosed materials closely.

Sincerely yours,

David E. Nelson, M.D.,M.P.H.
Chief, Behavioral Surveillance Branch
Division of Adult and Community Health
National Center for Chronic Disease
Prevention and Health Promotion

1998 BRFSS SUMMARY PREVALENCE REPORT

BEHAVIORAL SURVEILLANCE BRANCH
DIVISION OF ADULT AND COMMUNITY HEALTH
NATIONAL CENTER FOR CHRONIC DISEASE PREVENTION
AND HEALTH PROMOTION
CENTERS FOR DISEASE CONTROL AND PREVENTION

TABLE OF CONTENTS

Overview of the 1998 Summary Report.....	ix
Explanation of Tables.....	x
Definitions of Terms and Risk Factors.....	xii
Limitations of the Data	xvi

SECTION I

PREVALENCE REPORT FOR CORE QUESTIONS

General Health Status

TABLE 1.1 Self reported health status: Fair or poor.....	1
1.2 Self reported health status: Fair or poor-by sex	2
1.3 Self reported health status: Fair or poor, change 1997-1998	3

Quality of Life: Physical and Mental Health

TABLE 2.1 Physical health not good.....	4
2.2 Physical health not good-by sex.....	5
2.3 Physical health not good, change 1997-1998.....	6

TABLE 3.1 Mental health not good.....	7
3.2 Mental health not good-by sex.....	8
3.3 Mental health not good, change 1997-1998.....	9

TABLE 4.1 Activities limited.....	10
4.2 Activities limited-by sex.....	11
4.3 Activities limited, change 1997-1998.....	12

Health Insurance

TABLE 5.1 No health insurance, age 18 and over.....	13
5.2 No health insurance, age 18 and over-by sex.....	14
5.3 No health insurance, age 18 and over, change 1997-1998.....	15

TABLE 6.1 No health insurance, age 18-64 years.....	16
6.2 No health insurance, age 18-64 year-by sex.....	17

6.3 No health insurance, age 18-64 years, change 1997- 1998.....	18
--	----

Diabetes Awareness

TABLE 7.1 Ever told you have diabetes.....	19
7.2 Ever told you have diabetes-by sex.....	20
7.3 Ever told you have diabetes, change 1997-1998.....	21

Physical Activity

TABLE 8.1 No leisure time physical activity.....	22
(Year 2000 Objective 1.5)	
8.2 No leisure time physical activity- by sex.....	23
8.3 No leisure time physical activity, change 1996-1998.....	24

TABLE 9.1 No regular or sustained physical activity.....	25
(Year 2000 Objective 1.3)	
9.2 No regular or sustained physical activity- by sex.....	26
9.3 No regular or sustained physical activity, change 1996-1998.....	27

TABLE 10.1 No regular or vigorous physical activity.....	28
(Year 2000 Objective 1.4)	
10.2 No regular or vigorous physical activity- by sex.....	29
10.3 No regular or vigorous physical activity, change 1996-1998.....	30

Fruits & Vegetables

TABLE 11.1 Did not consume five or more servings of fruits and vegetables per day.....	31
(Year 2000 Objective 2.6)	
11.2 Did not consume five or more servings of fruits and vegetables per day- by sex.....	32
11.3 Did not consume five or more servings of fruits and vegetables per day, change1996-1998.....	33

Weight Control

TABLE 12.1 Not eating fewer calories and/or less fat combined with increased physical activity.....	34
12.2 Not eating fewer calories and/or less fat combined with increased physical activity- by sex.....	35
12.3 Not eating fewer calories and/or less fat combined with increased physical activity, change 1996-1998.....	36

Smoking Status

TABLE 13.1 Ever smoked.....	37
13.2 Ever smoked-by sex.....	38
13.3 Ever smoked, change 1996-1997.....	39
TABLE 14.1 Current smoker.....	40
(Year 2000 Objective 3.4)	
14.2 Current smoker-by sex.....	41
14.3 Current smoker, change 1997-1998.....	42
 Overweight	
TABLE 15.1 Based on body mass index (BMI)	43
(Year 2000 Objective 2.3)	
15.2 Based on body mass index (BMI)-by sex.....	44
15.3 Based on body mass index (BMI), change 1997-1998	45
 Overweight	
TABLE 16.1 Based on percent of median.....	46
16.2 Based on percent of median-by sex.....	47
16.3 Based on percent of median, change 1997-1998.....	48
 AIDS Knowledge/Attitudes	
TABLE 17.1 Would not encourage teen condom use, age 18-64.....	49
17.2 Would not encourage teen condom use-by sex.....	50
17.3 Would not encourage teen condom use, change 1997-1998.....	51
TABLE 18.1 Chances of getting HIV medium or high, age 18-64.....	52
18.2 Chances of getting HIV medium or high, age 18-64 -by sex.....	53
18.3 Chances of getting HIV medium or high, change 1996-1997.....	54
TABLE 19.1 Not Tested for HIV, age 18-64.....	55
19.2 Not Tested for HIV, age 18-64 - by sex.....	56

SECTION II

PREVALENCE REPORT FOR MODULES

Diabetes

TABLE 20.1 Age at onset of diabetes.....	57
20.2 Age at onset of diabetes-by sex.....	58
TABLE 21.1 Diabetics using insulin.....	59
21.2 Diabetics using insulin-by sex.....	60
TABLE 22.1 Health professional visits in past year.....	61
22.2 Health professional visits in past year-by sex.....	62

Immunization

TABLE 23.1 Did not have a flu shot within past 12 months, age 65 and older.....	63
(Year 2000 Objective 20.11)	
23.2 Did not have a flu shot within past 12 months, age 65 and older-by sex.....	64
TABLE 24.1 Never had pneumonia vaccination, age 65 and older	65
(Year 2000 Objective 20.11)	
24.2 Never had pneumonia vaccination, age 65 and older-by sex.....	66

Smokeless tobacco

TABLE 25.1 Ever used smokeless tobacco.....	67
25.2 Ever used smokeless tobacco, change 1997-1998.....	68
TABLE 26.1 Currently uses smokeless tobacco.....	69
26.2 Currently uses smokeless tobacco, change 1997-1998.....	70

SECTION III

PREVALENCE REPORT FOR WOMEN'S HEALTH ISSUES

Mammography

TABLE 27.1 Never had a mammogram, women age 18 and older.....	71
27.2 Never had a mammogram, women age 18 and older, change 1997-1998.....	72
TABLE 28.1 Never had a mammogram, women age 40 and older.....	73
28.2 Never had a mammogram, women age 40 and older, change 1997-1998.....	74
TABLE 29.1 Did not have a mammogram within the past two years, women age 50 and older.....	75
29.2 Did not have a mammogram within the past two years, women age 50 and older, change 1997-1998.....	76

Breast Exam

TABLE 30.1 Never had a breast exam, women age 18 and older.....	77
30.2 Never had a breast exam, women age 18 and older, change 1997-1998.....	78
TABLE 31.1 Never had a breast exam, women age 40 and older.....	79
31.2 Never had a breast exam, women age 40 and older, change 1997-1998.....	80
TABLE 32.1 Did not have a breast exam within the past two years, women age 50 and older.....	81
32.2 Did not have a breast exam within the past two years, women age 50 and older, change 1997-1998.....	82

Mammogram and Breast Exam

TABLE 33.1 Never had a mammogram and breast exam, women age 40 and older.....	83
(Approximates Year 2000 Objective 16.11)	
33.2 Never had a mammogram and breast exam, women age 40 and older, change 1997-1998.....	84

TABLE 34.1 Did not have a mammogram and breast exam within the past two years, women age 50 and older.....	85
(Approximates Year 2000 Objective 16.11)	
34.2 Did not have a mammogram and breast exam within the past two years, women age 50 and older, change 1997-1998.....	86

Cervical Cancer

TABLE 35.1 Never had pap smear, women age 18 and older.....	87
(Year 2000 Objective 16.12)	
35.2 Never had pap smear, women age 18 and older, change 1997-1998.....	88
TABLE 36.1 Never had pap smear, women with uterine cervix.....	89
36.2 Never had pap smear, women with uterine cervix, change 1997-1998.....	90
TABLE 37.1 Did not have pap smear within the past three years, women with uterine cervix.....	91
37.2 Did not have pap smear within the past three years, women with uterine cervix, change 1997-1998.....	92

OVERVIEW OF THE 1998 SUMMARY REPORT

The report is divided into three sections. *Section I* displays tables for selected variables in the core. In *Section I*, there are three tables for each risk factor or preventive measure unless otherwise noted. Estimated prevalence and relevant statistics are provided for the total population sampled and by sex. Prevalence estimates for both 1997 and 1998, or where appropriate 1996 and 1998, and the percentage point difference between the estimates are given for variables in the core.

Section II shows tables for selected variables in modules used by at least 13 states. *Section II* contains two tables for each risk factor or preventive measure of interest in the modules. These tables show the estimated prevalence and relevant statistics for the total sampled population and both sex groups who responded to selected questions in the optional modules.

Section III contains tables for preventive health measures that are related to women's health issues. Estimated prevalence and relevant statistics are provided for the total population sampled. Prevalence estimates for both 1997 and 1998 and the percentage point difference between them are given.

Three tables (Tables 1.1-1.3) from Section I will be used as examples to explain the information contained in this report. Each table describes a behavioral risk factor or preventive health measure from the Behavioral Risk Factor Surveillance System (BRFSS) instrument.

Table 1.1 (page 1) shows the estimated prevalence and relevant statistics for each participant. (Participants refer to all states, and the District of Columbia and territories of the U.S.). The table shows the estimated prevalence and relevant statistics for "fair or poor health

status" in the population surveyed by each BRFSS participant. This prevalence estimate excludes respondents with unknown information on self reported health status.

The columns in Table 1.1 provide the following information:

<u>COLUMN HEADING</u>	<u>CONTENTS</u>
<u>Participant</u>	Name of the participant. Participants refers to all states including the District of Columbia and territories of the United States.
<u>Sample size</u>	The number of individuals who responded to the specific BRFSS question. These are unweighted numbers. These numbers exclude individuals who reported "not sure, don't know", those who refused to answer the question, or had no response coded (missing or skipped).
<u>Percent/Mean</u>	An estimate of the percent (or prevalence or mean number) of residents who report the behavioral risk factor of interest. This is weighted to the participant's population to estimate the true prevalence or mean in the population.
<u>Standard error</u>	A measure of the amount of variability that exists in the calculation of the prevalence estimate.
<u>95% confidence interval</u>	A range which contains the true population prevalence estimate 95% of the time, if repeated sampling of the population is performed.

SUMMARY STATISTICS are given at the bottom of the table:

<u>Number of participants</u>	The total number of participants who asked question(s) regarding the respective "risk factor".
<u>Median</u>	The middle value of all estimated prevalences.
<u>Range</u>	The lowest prevalence and the highest prevalence among all participants.

For example, in Table 1.1, there were 2234 individuals in the state of Alabama who responded to the "self reported health status" question. Of those responding, the estimated prevalence of Alabama's population who reported that they had fair or poor health status is 21.3 percent. This prevalence estimate has a standard error of 1.0. The 95% confidence interval for this estimate is (19.4, 23.2). Ninety-five percent (95%) of confidence intervals generated under repeated sampling will contain the true population prevalence of Alabama's population who "had fair or poor health status".

Table 1.2 (page 2) shows the estimated prevalence and relevant statistics by sex for each state participant. The same statistics are displayed in Table 1.2 as in Table 1.1, except that Table 1.2 shows the estimated prevalence of “ fair to poor health “ for males and females separately. An explanation of the terms was provided earlier. This side by side display of the tables permits each participant to compare estimates for males and females in their respective state.

Table 1.3 (page 3) shows the percent difference between the prevalence estimates, by participant, for the 1997 and 1998 survey years for having fair or poor health status. A negative change indicates a decrease in prevalence from 1997 to 1998, while a positive change indicates an increase in prevalence between the two years. This difference gives state health officials a quick method to examine changes that may have occurred in their respective state. Some of these changes may have resulted from legislative actions (or inactions) in the state, programs funded, or programs that came into existence because of results that states may have shown from the BRFSS. Since changes between two years may be attributable to many factors, conclusions drawn from this table should be interpreted with extreme caution.

Sample Size. Sample sizes vary by question. Some estimates may be based on responses from less than 50 respondents. Interpreting estimates that are based on a small number of respondents can mislead the reader into believing that a given finding is much more precise than it actually is. The BRFSS recommends not interpreting percentages where the denominator is based upon fewer than 50 unweighted respondents (sample size).

DEFINITIONS OF TERMS AND RISK FACTORS IN THIS REPORT

AIDS KNOWLEDGE/ATTITUDES

HIV TEST - Respondents, age 18-64, who have never had a test for HIV infection.

CHANCES OF GETTING HIV - Respondents, age 18-64, who believe that their chances of getting infected with HIV, the virus that causes AIDS are medium or high.

TEEN CONDOM USE - Respondents, age 18-64, who would not encourage sexually active teenagers to use a condom.

BREAST EXAM

BREAST EXAM (1)- Female respondents, age 18 and older, who report that they have never had a breast exam by a doctor.

BREAST EXAM (2)- Female respondents, age 40 and older, who report that they never had a breast exam by a doctor.

BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they did not have a breast exam within the past two years.

CERVICAL CANCER

PAP SMEAR (1)- Female respondents, age 18 and older who report that they have never had a pap smear. This measures Year 2000 Objective 16.12. - Target $\geq 95\%$.

PAP SMEAR, WITH UTERINE CERVIX (2)- Female respondents, with uterine cervix, age 18 and older, who report that they never had a pap smear.

PAP SMEAR WITHIN 3 YEARS - Female respondents, with uterine cervix, age 18 and older, who report that did not have a pap smear within the last three years.

CLINICAL BREAST EXAM See BREAST EXAM

DIABETES

AGE AT ONSET - Mean age at onset of diabetes: age when respondent was told that they have diabetes, among those who report they were told by a doctor that they have diabetes.

DIABETES AWARENESS - Respondents who report they were told by a doctor that they have diabetes.

NOW TAKING INSULIN - Respondents who are now taking insulin, among those who report they were told by a doctor that they have diabetes.

NUMBER OF TIMES SEEN HEALTH PROFESSIONAL - Mean number of times respondents report they have seen a doctor, nurse, or other health professional for their diabetes in the past year, among those who report they were told by a doctor that they have diabetes.

FRUITS AND VEGETABLES

FRUITS AND VEGETABLES - Respondents who report they are not consuming five or more servings of fruits and vegetables a day. This measures Year 2000 Objective 2.6 - Target is not specified.

HEALTH INSURANCE

HEALTH INSURANCE (1)- Respondents age 18 and older who report they have no health care plan.

HEALTH INSURANCE (2)- Respondents age 18-64 years who report they have no health care plan.

HEALTH STATUS

ACTIVITIES LIMITED - Mean number of days in past 30 days activities were limited as reported by respondents.

MENTAL HEALTH NOT GOOD - Mean number of days in past 30 days mental health was not good as reported by respondents.

PHYSICAL HEALTH NOT GOOD - Mean number of days in past 30 days physical health was not good as reported by respondents.

SELF REPORTED HEALTH STATUS - Respondents who report having a general health status of fair or poor.

IMMUNIZATION

FLU SHOT - Respondents, age 65 and older, who report that they did not have a flu shot within past 12 months. This measures Year 2000 Objective 20.11 - Target $\geq 80\%$.

PNEUMONIA VACCINATION - Respondents, age 65 and older, who report they have never had a pneumonia vaccination. This measures Year 2000 Objective 20.11 - Target \geq 80%.

MAMMOGRAPHY

MAMMOGRAM (1) - Female respondents age 18 and older, who report that they have never had a mammogram.

MAMMOGRAM (2) - Female respondents, age 40 and older, who report that they have never had a mammogram.

MAMMOGRAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they did not have a mammogram within the past two years.

MAMMOGRAPHY AND BREAST EXAM

MAMMOGRAM & BREAST EXAM - Female respondents, age 40 and older, who report that they have never had a mammogram and a breast exam. This measures Year 2000 Objective 16.11 - Target \geq 80%.

MAMMOGRAM & BREAST EXAM WITHIN 2 YEARS - Female respondents, age 50 and older, who report that they did not have a mammogram and a breast exam within the last two years. This measures Year 2000 Objective 16.11 - Target \geq 60%.

OVERWEIGHT

OVERWEIGHT: BY MEDIAN - Respondents at or above 120% of their ideal weight. Ideal weight is defined as the mid-value of a median frame person from the 1959 Metropolitan Life Insurance Corporation height-weight tables.

OVERWEIGHT: BY BODY MASS INDEX - Females with body mass index(BMI) ≥ 27.3 and males with BMI ≥ 27.8 . BMI is defined as weight in kilograms divided by height in meters squared (w/h^{**2}). This measures Year 2000 Objective 2.3 - Target \leq 20%. This should be used with extreme caution since people tend to underreport their weight. BRFSS may underestimate the prevalence of overweight.

PAP SMEAR See CERVICAL CANCER

PHYSICAL INACTIVITY

NO LEISURE TIME PHYSICAL ACTIVITY - Respondents who report no leisure-time physical activity during the past month. This measures Year 2000 Objective 1.5 - Target \leq 15%.

REGULAR AND SUSTAINED PHYSICAL ACTIVITY - Respondents who report no physical activity defined as: 5 or more sessions per week, 30 minutes or more per session, regardless of intensity. This measures Year 2000 Objective 1.3 - Target $\geq 30\%$.

REGULAR AND VIGOROUS PHYSICAL ACTIVITY - Respondents who report no physical activity or a pair of activities defined as: 3 or more sessions per week, 20 minutes or more per session, at 50% or more capacity. This measures Year 2000 Objective 1.4 - Target $\geq 20\%$.

SMOKELESS TOBACCO

EVER USED - Respondents who have ever used or tried smokeless tobacco.

CURRENT USE - Respondents who report that they currently use smokeless tobacco.

SMOKING STATUS

EVER SMOKED - Respondents who have ever smoked 100 cigarettes in their lifetime.

CURRENT SMOKER - Respondents who have ever smoked 100 cigarettes in their lifetime and reported smoking every day or some days. This measures Year 2000 Objective 3.4 - Target $\leq 15\%$.

WEIGHT CONTROL

CALORIES AND PHYSICAL ACTIVITY - Respondents at risk on BMI who report that they are not either eating fewer calories or less fat and not using physical activity or exercise to lose or maintain weight.

LIMITATIONS OF THE DATA

This report consolidates the most current data on selected risk factors and preventive health measures for states, the District of Columbia and Puerto Rico. The information was obtained from the data files of individual states. Therefore, while all participants use telephone surveys to collect the data, the data in this report vary somewhat with respect to data collection, sampling design and sample size.

Some states conduct interviews using a paper questionnaire, but most use a computer-assisted telephone interviewing (CATI) software program. All states use probability-based methods to select their sample. The BRFSS standard for sample designs is that sample records must be justifiable as a probability sample of all households with telephones in the state. All participating areas except Alaska, California, and Hawaii met this criterion in 1998.

For data year 1998, eleven states used the Mitofsky-Waksberg method of sampling. Thirty-seven states used a disproportionate stratified sample (DSS) design. Puerto Rico used a simple random sample design. The three remaining states—Alaska, California, and Hawaii—used a variety of other designs.

Alaska

Alaska's sample design is made up of three random digit-dialed strata and four listed strata, where phone numbers are selected from a list of putative household numbers. The sampling frame contains all of the household numbers in the random digit-dialed strata and an estimated 45% of all household numbers in the listed strata. This represents an exclusion of approximately 9% of all residential numbers in Alaska.

In 3 out of 4 geographic strata, prefixes are assigned to a high (RDD) or low (listed) density stratum depending largely on the number of active household numbers (about 2000 being the threshold) in the exchange (10,000 numbers). All numbers in the fourth geographic stratum are assigned to the listed density stratum. In the RDD strata, the probability that a number is selected depends on the number of active household numbers in its exchange. In the RDD strata, a number of prefixes equal to the target sample size are selected at random in proportion to their number of active HH's. (A prefix can be selected more than once.) Once a prefix is selected into the RDD sample, 48 suffixes are randomly generated. Once the entire sample for the stratum is generated duplicate numbers are deleted without replacement. The final result is h lists of phone numbers per stratum, each containing 48 or fewer unique numbers, where h is the target number of completes per stratum. Phone numbers are called sequentially from each list until one complete per list is obtained.

California

California, in July through December 1998, used a sample design in which only phone numbers from hundred blocks with one or more listed household numbers were included in the sampling

frame. Such hundred blocks are estimated to contain 97.8% of all household numbers in California. Numbers from this frame were selected with an equal probability of selection.

Hawaii

Hawaii uses a sampling frame obtained from the state's telephone company. The frame purportedly contains all working prefixes in Hawaii. The frame is divided into six strata, corresponding to the major islands of Hawaii. The working prefixes are used to randomly generate telephone numbers with a probability proportional to the number of known household numbers in the hundred block to which a number belongs. Telephone numbers from hundred blocks with no known household numbers have a zero probability of being included in the sample. The percentage of telephone households excluded from the sampling frame is unknown. The numbers generated are subsequently screened by GTE Hawaii, who screens out all but the listed and unlisted household numbers in the sample.

Other areas

In a few states, a portion of sample records intended for use during one month may have been completed in another month. This deviation should only affect analyses based on monthly, rather than annual, data.

B. Other 1998 limitations of the data

In addition to departing from the standard for sample designs, California modified the wording of mammography, Papanicolaou (PAP) smear questions, and are therefore excluded from those tables in the report. California also asked the HIV/AIDS section questions to persons 18-45 years of age rather than to those 18-64 years of age as specified, and are therefore not included in the tables for this topic.

Although almost all states collect data monthly, the annual number of interviews range from 1616 in Maine to 6005 in Texas. Estimates for some states are therefore based upon smaller sample sizes, and for rare events (e.g. diabetes), may yield unstable estimates. **Maryland** did not collect data in January, October, November and December. Some items, including drinking and driving and obesity, are affected by seasonality, and estimates for these items may be biased, especially for Maryland's subpopulations.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. People may not remember essential information, a question may not mean the same thing to different respondents, and some individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their impact on the data. The user must make his or her own evaluation of the data. Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. When the number of events is small and the probability of such an even is small, considerable caution must be observed in interpreting the estimates and/or differences between groups and areas.

TABLE 1.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2234	21.3	1.0	(19.4, 23.2)
ALASKA	1985	11.0	1.0	(9.1, 12.9)
ARIZONA	1915	10.3	1.0	(8.4, 12.1)
ARKANSAS	2987	20.4	0.8	(18.8, 22.0)
CALIFORNIA	4158	14.6	0.7	(13.3, 15.9)
COLORADO	1801	11.6	0.8	(9.9, 13.2)
CONNECTICUT	3122	11.7	0.7	(10.2, 13.1)
DELAWARE	2578	12.8	0.9	(11.0, 14.6)
DISTRICT OF CO	1448	12.4	1.2	(10.0, 14.8)
FLORIDA	4713	15.5	0.6	(14.3, 16.7)
GEORGIA	2366	15.9	0.9	(14.2, 17.6)
HAWAII	2148	12.3	0.9	(10.4, 14.1)
IDAHO	4921	12.2	0.5	(11.2, 13.3)
ILLINOIS	2877	12.5	0.7	(11.1, 13.9)
INDIANA	2422	13.3	0.8	(11.8, 14.7)
IOWA	3646	11.2	0.6	(10.0, 12.4)
KANSAS	3798	12.0	0.6	(10.9, 13.0)
KENTUCKY	3715	21.9	0.7	(20.4, 23.3)
LOUISIANA	1650	16.1	1.0	(14.1, 18.1)
MAINE	1615	12.5	0.9	(10.7, 14.3)
MARYLAND	3287	13.9	0.9	(12.3, 15.6)
MASSACHUSETTS	4933	10.9	0.6	(9.8, 12.0)
MICHIGAN	2532	14.5	0.8	(13.0, 16.0)
MINNESOTA	4217	10.4	0.5	(9.4, 11.4)
MISSISSIPPI	2297	21.0	1.0	(19.1, 22.9)
MISSOURI	3712	15.2	0.7	(13.8, 16.6)
MONTANA	1800	12.0	0.8	(10.4, 13.6)
NEBRASKA	2710	12.3	0.7	(11.0, 13.6)
NEVADA	1973	12.4	1.2	(10.2, 14.7)
NEW HAMPSHIRE	1493	9.9	0.8	(8.3, 11.6)
NEW JERSEY	2458	11.8	0.7	(10.3, 13.2)
NEW MEXICO	3636	15.0	0.7	(13.7, 16.3)
NEW YORK	2520	13.8	0.8	(12.3, 15.3)
NORTH CAROLINA	2397	16.6	0.9	(14.7, 18.4)
NORTH DAKOTA	1800	13.7	0.9	(11.9, 15.4)
OHIO	2986	16.1	0.9	(14.3, 17.9)
OKLAHOMA	2511	12.6	0.7	(11.2, 14.0)
OREGON	1817	13.4	0.9	(11.6, 15.2)
PENNSYLVANIA	3590	15.1	0.7	(13.7, 16.4)
RHODE ISLAND	3591	13.5	0.6	(12.3, 14.8)
SOUTH CAROLINA	3279	15.2	0.7	(13.8, 16.6)
SOUTH DAKOTA	2111	11.7	0.8	(10.1, 13.3)
TENNESSEE	2940	18.2	0.8	(16.6, 19.8)
TEXAS	5986	18.6	0.7	(17.3, 19.9)
UTAH	2859	10.8	0.7	(9.4, 12.2)
VERMONT	2995	10.0	0.6	(8.9, 11.2)
VIRGINIA	3489	12.3	0.8	(10.8, 13.8)
WASHINGTON	3594	11.1	0.6	(9.9, 12.3)
WEST VIRGINIA	2414	23.9	1.0	(22.0, 25.8)
WISCONSIN	2204	12.1	0.9	(10.4, 13.7)
WYOMING	2433	12.1	0.8	(10.5, 13.7)
PUERTO RICO	2803	32.1	1.0	(30.2, 34.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 12.7 RANGE = 9.9-32.1

* HEALTH STATUS IS FAIR OR POOR

June 18, 1999

TABLE 1.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

2

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	781	18.5	1.4	(15.7, 21.4)	*	1453	23.8	1.3	(21.3, 26.2)	*
ALASKA	921	11.0	1.5	(8.2, 13.9)	*	1064	10.9	1.3	(8.5, 13.4)	*
ARIZONA	824	9.5	1.3	(7.0, 12.0)	*	1091	11.0	1.4	(8.2, 13.7)	*
ARKANSAS	1142	18.8	1.3	(16.4, 21.3)	*	1845	21.7	1.1	(19.6, 23.8)	*
CALIFORNIA	1820	13.2	0.9	(11.4, 15.1)	*	2338	16.0	0.9	(14.2, 17.9)	*
COLORADO	810	9.5	1.1	(7.3, 11.7)	*	991	13.5	1.2	(11.1, 15.9)	*
CONNECTICUT	1284	11.1	1.1	(8.9, 13.4)	*	1838	12.2	0.9	(10.3, 14.0)	*
DELAWARE	1035	12.1	1.5	(9.2, 15.0)	*	1543	13.5	1.1	(11.4, 15.6)	*
DISTRICT OF CO	542	11.6	2.1	(7.4, 15.7)	*	906	13.1	1.3	(10.4, 15.7)	*
FLORIDA	1882	13.8	0.9	(12.0, 15.5)	*	2831	17.0	0.8	(15.5, 18.6)	*
GEORGIA	891	14.6	1.4	(12.0, 17.3)	*	1475	17.1	1.1	(14.9, 19.3)	*
HAWAII	931	9.9	1.2	(7.4, 12.3)	*	1217	14.7	1.4	(11.9, 17.4)	*
IDAHO	2078	12.5	0.9	(10.8, 14.2)	*	2843	11.9	0.7	(10.7, 13.2)	*
ILLINOIS	1231	12.4	1.1	(10.2, 14.6)	*	1646	12.6	0.9	(10.8, 14.4)	*
INDIANA	990	12.8	1.2	(10.5, 15.0)	*	1432	13.7	1.0	(11.8, 15.7)	*
IOWA	1447	11.0	1.0	(9.0, 12.9)	*	2199	11.4	0.8	(9.9, 12.9)	*
KANSAS	1484	10.3	0.8	(8.7, 11.8)	*	2314	13.6	0.8	(12.1, 15.1)	*
KENTUCKY	1363	20.5	1.2	(18.2, 22.9)	*	2352	23.1	0.9	(21.3, 24.9)	*
LOUISIANA	648	15.7	1.6	(12.6, 18.8)	*	1002	16.5	1.3	(13.9, 19.1)	*
MAINE	669	12.3	1.4	(9.6, 15.0)	*	946	12.7	1.2	(10.4, 15.0)	*
MARYLAND	1299	14.2	1.4	(11.5, 16.9)	*	1988	13.7	1.0	(11.6, 15.7)	*
MASSACHUSETTS	2014	10.7	0.9	(9.0, 12.4)	*	2919	11.0	0.7	(9.6, 12.4)	*
MICHIGAN	1037	12.6	1.1	(10.3, 14.8)	*	1495	16.2	1.0	(14.1, 18.2)	*
MINNESOTA	1989	9.5	0.8	(8.0, 11.0)	*	2228	11.3	0.7	(9.9, 12.7)	*
MISSISSIPPI	867	20.8	1.6	(17.8, 23.9)	*	1430	21.2	1.2	(19.0, 23.5)	*
MISSOURI	1429	13.0	1.0	(11.0, 15.0)	*	2283	17.2	1.0	(15.3, 19.1)	*
MONTANA	786	10.1	1.2	(7.8, 12.4)	*	1014	13.8	1.2	(11.5, 16.2)	*
NEBRASKA	1054	11.8	1.0	(9.8, 13.8)	*	1656	12.7	0.8	(11.1, 14.4)	*
NEVADA	896	11.9	1.7	(8.5, 15.2)	*	1077	12.9	1.5	(10.0, 15.9)	*
NEW HAMPSHIRE	613	8.3	1.2	(6.0, 10.5)	*	880	11.5	1.2	(9.1, 13.8)	*
NEW JERSEY	1008	10.9	1.1	(8.8, 13.0)	*	1450	12.5	1.0	(10.6, 14.5)	*
NEW MEXICO	1571	12.5	0.9	(10.7, 14.3)	*	2065	17.4	1.0	(15.4, 19.3)	*
NEW YORK	1052	12.3	1.1	(10.1, 14.4)	*	1468	15.2	1.1	(13.1, 17.4)	*
NORTH CAROLINA	919	14.2	1.4	(11.4, 17.0)	*	1478	18.8	1.2	(16.4, 21.2)	*
NORTH DAKOTA	805	13.8	1.3	(11.2, 16.4)	*	995	13.5	1.2	(11.2, 15.9)	*
OHIO	1182	14.5	1.4	(11.8, 17.2)	*	1804	17.6	1.3	(15.1, 20.0)	*
OKLAHOMA	1002	12.4	1.1	(10.3, 14.6)	*	1509	12.7	0.9	(11.0, 14.5)	*
OREGON	741	12.3	1.3	(9.8, 14.8)	*	1076	14.4	1.3	(11.9, 16.9)	*
PENNSYLVANIA	1425	14.0	1.1	(11.9, 16.0)	*	2165	16.1	0.9	(14.3, 17.8)	*
RHODE ISLAND	1409	12.3	1.0	(10.4, 14.3)	*	2182	14.6	0.8	(13.0, 16.3)	*
SOUTH CAROLINA	1367	14.3	1.1	(12.2, 16.5)	*	1912	16.0	1.0	(14.1, 17.9)	*
SOUTH DAKOTA	925	12.3	1.4	(9.6, 15.0)	*	1186	11.2	1.0	(9.3, 13.0)	*
TENNESSEE	1026	17.3	1.4	(14.7, 19.9)	*	1914	19.0	1.0	(17.2, 20.9)	*
TEXAS	2445	16.5	1.0	(14.6, 18.5)	*	3541	20.5	0.9	(18.8, 22.3)	*
UTAH	1234	10.7	1.2	(8.4, 13.0)	*	1625	10.9	0.9	(9.1, 12.7)	*
VERMONT	1279	8.6	0.8	(6.9, 10.2)	*	1716	11.4	0.8	(9.8, 13.0)	*
VIRGINIA	1478	11.4	1.0	(9.4, 13.5)	*	2011	13.1	1.1	(11.0, 15.3)	*
WASHINGTON	1586	11.0	0.9	(9.3, 12.6)	*	2008	11.2	0.9	(9.5, 12.9)	*
WEST VIRGINIA	942	24.6	1.5	(21.6, 27.5)	*	1472	23.3	1.2	(21.0, 25.7)	*
WISCONSIN	977	10.1	1.1	(7.9, 12.3)	*	1227	13.9	1.3	(11.4, 16.3)	*
WYOMING	1023	12.0	1.3	(9.4, 14.6)	*	1410	12.1	0.9	(10.3, 14.0)	*
PUERTO RICO	1073	26.3	1.4	(23.5, 29.1)	*	1730	37.2	1.3	(34.7, 39.7)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 12.3
 RANGE = 8.3-26.3

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 13.7
 RANGE = 10.9-37.2

June 18, 1999

* HEALTH STATUS IS FAIR OR POOR

TABLE 1.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 SELF REPORTED HEALTH STATUS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

3

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	19.0	21.3	2.3
ALASKA	11.0	11.0	-0.0
ARIZONA	11.4	10.3	-1.1
ARKANSAS	20.9	20.4	-0.5
CALIFORNIA	15.3	14.6	-0.7
COLORADO	10.0	11.6	1.6
CONNECTICUT	10.2	11.7	1.4
DELAWARE	13.8	12.8	-0.9
DISTRICT OF CO	9.7	12.4	2.7
FLORIDA	16.4	15.5	-0.9
GEORGIA	11.4	15.9	4.5
HAWAII	14.0	12.3	-1.8
IDAHO	12.4	12.2	-0.2
ILLINOIS	14.0	12.5	-1.5
INDIANA	14.1	13.3	-0.8
IOWA	11.3	11.2	-0.2
KANSAS	10.7	12.0	1.3
KENTUCKY	21.8	21.9	0.1
LOUISIANA	15.3	16.1	0.8
MAINE	13.9	12.5	-1.3
MARYLAND	11.9	13.9	2.0
MASSACHUSETTS	12.0	10.9	-1.1
MICHIGAN	12.7	14.5	1.7
MINNESOTA	9.5	10.4	1.0
MISSISSIPPI	21.9	21.0	-0.9
MISSOURI	15.0	15.2	0.2
MONTANA	10.1	12.0	1.9
NEBRASKA	11.4	12.3	0.9
NEVADA	14.4	12.4	-2.0
NEW HAMPSHIRE	9.6	9.9	0.3
NEW JERSEY	13.8	11.8	-2.1
NEW MEXICO	14.4	15.0	0.6
NEW YORK	13.3	13.8	0.5
NORTH CAROLINA	16.1	16.6	0.5
NORTH DAKOTA	12.1	13.7	1.6
OHIO	13.7	16.1	2.4
OKLAHOMA	13.4	12.6	-0.8
OREGON	12.8	13.4	0.6
PENNSYLVANIA	13.1	15.1	2.0
RHODE ISLAND	12.6	13.5	0.9
SOUTH CAROLINA	16.8	15.2	-1.6
SOUTH DAKOTA	11.3	11.7	0.4
TENNESSEE	18.1	18.2	0.1
TEXAS	16.9	18.6	1.7
UTAH	11.2	10.8	-0.4
VERMONT	11.4	10.0	-1.4
VIRGINIA	11.2	12.3	1.1
WASHINGTON	10.6	11.1	0.5
WEST VIRGINIA	21.7	23.9	2.2
WISCONSIN	9.8	12.1	2.3
WYOMING	11.3	12.1	0.7
PUERTO RICO	35.0	32.1	-2.9

* HEALTH STATUS IS FAIR OR POOR

June 18, 1999

TABLE 2.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD*

4

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2173	4.2	0.2	(3.8, 4.6)
ALASKA	1960	2.8	0.2	(2.4, 3.2)
ARIZONA	1911	1.1	0.1	(0.9, 1.4)
ARKANSAS	2957	3.6	0.2	(3.3, 4.0)
CALIFORNIA	4134	3.1	0.1	(2.8, 3.3)
COLORADO	1790	2.9	0.2	(2.6, 3.3)
CONNECTICUT	3081	2.9	0.2	(2.6, 3.2)
DELAWARE	2566	3.2	0.2	(2.8, 3.7)
DISTRICT OF CO	1423	2.2	0.2	(1.8, 2.5)
FLORIDA	4654	3.4	0.1	(3.1, 3.6)
GEORGIA	2316	3.0	0.2	(2.7, 3.3)
HAWAII	2132	2.2	0.2	(1.9, 2.6)
IDAHO	4892	3.1	0.1	(2.8, 3.3)
ILLINOIS	2856	2.8	0.2	(2.5, 3.1)
INDIANA	2406	3.3	0.2	(2.9, 3.6)
IOWA	3617	2.9	0.1	(2.7, 3.2)
KANSAS	3738	2.8	0.1	(2.5, 3.0)
KENTUCKY	3708	3.6	0.2	(3.3, 3.9)
LOUISIANA	1607	3.0	0.2	(2.6, 3.4)
MAINE	1603	3.3	0.2	(2.9, 3.7)
MARYLAND	3238	3.0	0.2	(2.6, 3.3)
MASSACHUSETTS	4867	2.8	0.1	(2.5, 3.0)
MICHIGAN	2503	3.7	0.2	(3.4, 4.1)
MINNESOTA	4167	3.2	0.1	(2.9, 3.5)
MISSISSIPPI	2204	3.7	0.2	(3.3, 4.1)
MISSOURI	3669	3.4	0.2	(3.1, 3.7)
MONTANA	1772	3.0	0.2	(2.6, 3.4)
NEBRASKA	2686	3.1	0.2	(2.8, 3.4)
NEVADA	1953	3.2	0.3	(2.7, 3.6)
NEW HAMPSHIRE	1470	3.0	0.2	(2.6, 3.4)
NEW JERSEY	2433	2.8	0.2	(2.5, 3.1)
NEW MEXICO	3623	3.5	0.2	(3.2, 3.8)
NEW YORK	2472	3.2	0.2	(2.9, 3.5)
NORTH CAROLINA	2354	3.7	0.2	(3.3, 4.1)
NORTH DAKOTA	1799	2.7	0.2	(2.4, 3.0)
OHIO	2976	3.1	0.2	(2.7, 3.4)
OKLAHOMA	2480	2.6	0.2	(2.3, 3.0)
OREGON	1804	3.3	0.2	(2.9, 3.7)
PENNSYLVANIA	3545	3.3	0.2	(3.0, 3.6)
RHODE ISLAND	3532	3.4	0.2	(3.1, 3.7)
SOUTH CAROLINA	3243	3.1	0.2	(2.8, 3.4)
SOUTH DAKOTA	2096	2.6	0.2	(2.3, 2.9)
TENNESSEE	2889	3.3	0.2	(2.9, 3.6)
TEXAS	5929	3.3	0.1	(3.0, 3.5)
UTAH	2843	3.4	0.2	(3.0, 3.8)
VERMONT	2950	3.1	0.2	(2.8, 3.4)
VIRGINIA	3402	2.8	0.2	(2.5, 3.1)
WASHINGTON	3527	3.1	0.1	(2.8, 3.4)
WEST VIRGINIA	2398	3.4	0.2	(3.1, 3.8)
WISCONSIN	2184	2.9	0.2	(2.6, 3.3)
WYOMING	2413	2.9	0.2	(2.5, 3.3)
PUERTO RICO	2761	3.6	0.2	(3.3, 3.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.1 RANGE = 1.1- 4.2

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

June 18, 1999

TABLE 2.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES					
	SAMPLE	MEAN	STANDARD	95% CONFIDENCE	INTERVAL	SAMPLE	MEAN	STANDARD	95% CONFIDENCE		
	SIZE		ERROR	*	*	SIZE		ERROR	INTERVAL		
ALABAMA	764	3.4	0.3	(2.8,	4.0)	*	1409	4.9	0.3	(4.3,	5.4)
ALASKA	912	2.6	0.3	(2.0,	3.2)	*	1048	3.0	0.3	(2.4,	3.5)
ARIZONA	823	1.1	0.2	(0.7,	1.5)	*	1088	1.2	0.2	(0.9,	1.5)
ARKANSAS	1137	3.2	0.3	(2.7,	3.7)	*	1820	4.1	0.3	(3.6,	4.6)
CALIFORNIA	1811	2.7	0.2	(2.4,	3.1)	*	2323	3.4	0.2	(3.1,	3.8)
COLORADO	806	2.0	0.2	(1.6,	2.5)	*	984	3.8	0.3	(3.3,	4.3)
CONNECTICUT	1273	2.6	0.3	(2.1,	3.1)	*	1808	3.2	0.2	(2.7,	3.6)
DELAWARE	1030	2.9	0.4	(2.2,	3.6)	*	1536	3.6	0.3	(3.0,	4.1)
DISTRICT OF CO	539	1.8	0.3	(1.3,	2.3)	*	884	2.5	0.3	(2.0,	3.0)
FLORIDA	1858	2.8	0.2	(2.5,	3.2)	*	2796	3.8	0.2	(3.5,	4.2)
GEORGIA	884	2.4	0.2	(1.9,	2.8)	*	1432	3.5	0.2	(3.1,	4.0)
HAWAII	925	1.9	0.3	(1.4,	2.4)	*	1207	2.6	0.2	(2.1,	3.0)
IDAHO	2073	2.5	0.2	(2.2,	2.9)	*	2819	3.6	0.2	(3.2,	3.9)
ILLINOIS	1227	2.3	0.2	(1.8,	2.7)	*	1629	3.3	0.2	(2.9,	3.7)
INDIANA	987	3.0	0.2	(2.5,	3.4)	*	1419	3.6	0.2	(3.1,	4.0)
IOWA	1439	2.6	0.2	(2.2,	3.1)	*	2178	3.2	0.2	(2.9,	3.6)
KANSAS	1474	2.5	0.2	(2.2,	2.9)	*	2264	3.0	0.2	(2.7,	3.4)
KENTUCKY	1361	2.9	0.2	(2.5,	3.4)	*	2347	4.3	0.2	(3.9,	4.7)
LOUISIANA	636	3.2	0.4	(2.5,	3.9)	*	971	2.8	0.3	(2.3,	3.3)
MAINE	666	2.8	0.3	(2.2,	3.4)	*	937	3.8	0.3	(3.2,	4.3)
MARYLAND	1285	2.5	0.2	(2.1,	2.9)	*	1953	3.4	0.2	(2.9,	3.9)
MASSACHUSETTS	1995	2.6	0.2	(2.2,	2.9)	*	2872	3.0	0.2	(2.7,	3.3)
MICHIGAN	1028	3.1	0.3	(2.6,	3.6)	*	1475	4.3	0.3	(3.8,	4.8)
MINNESOTA	1970	2.9	0.2	(2.5,	3.2)	*	2197	3.5	0.2	(3.1,	3.9)
MISSISSIPPI	844	3.6	0.4	(2.9,	4.3)	*	1360	3.8	0.2	(3.3,	4.2)
MISSOURI	1421	2.8	0.2	(2.4,	3.2)	*	2248	3.9	0.2	(3.5,	4.3)
MONTANA	780	2.5	0.2	(2.0,	2.9)	*	992	3.5	0.3	(2.9,	4.0)
NEBRASKA	1050	2.6	0.2	(2.1,	3.1)	*	1636	3.6	0.2	(3.2,	4.0)
NEVADA	889	2.3	0.3	(1.8,	2.8)	*	1064	4.0	0.4	(3.2,	4.8)
NEW HAMPSHIRE	608	2.5	0.3	(2.0,	3.1)	*	862	3.5	0.3	(3.0,	4.0)
NEW JERSEY	1000	2.4	0.2	(1.9,	2.8)	*	1433	3.1	0.2	(2.7,	3.6)
NEW MEXICO	1567	2.8	0.2	(2.4,	3.2)	*	2056	4.2	0.2	(3.8,	4.6)
NEW YORK	1036	2.8	0.3	(2.3,	3.3)	*	1436	3.6	0.2	(3.2,	4.1)
NORTH CAROLINA	912	2.9	0.3	(2.3,	3.4)	*	1442	4.5	0.3	(3.9,	5.1)
NORTH DAKOTA	802	2.3	0.2	(1.8,	2.7)	*	997	3.1	0.3	(2.6,	3.6)
OHIO	1182	2.6	0.3	(2.1,	3.1)	*	1794	3.5	0.3	(3.0,	4.0)
OKLAHOMA	991	2.4	0.3	(1.9,	2.9)	*	1489	2.8	0.2	(2.4,	3.2)
OREGON	738	3.0	0.3	(2.4,	3.5)	*	1066	3.6	0.3	(3.1,	4.2)
PENNSYLVANIA	1414	2.9	0.2	(2.5,	3.3)	*	2131	3.8	0.2	(3.4,	4.1)
RHODE ISLAND	1397	3.0	0.2	(2.6,	3.5)	*	2135	3.7	0.2	(3.3,	4.0)
SOUTH CAROLINA	1362	2.8	0.2	(2.4,	3.2)	*	1881	3.3	0.2	(2.9,	3.7)
SOUTH DAKOTA	921	2.2	0.2	(1.7,	2.7)	*	1175	3.0	0.2	(2.5,	3.4)
TENNESSEE	1013	2.9	0.3	(2.4,	3.5)	*	1876	3.6	0.2	(3.2,	4.0)
TEXAS	2432	2.6	0.2	(2.2,	2.9)	*	3497	4.0	0.2	(3.6,	4.3)
UTAH	1229	2.5	0.3	(2.0,	3.0)	*	1614	4.2	0.3	(3.7,	4.8)
VERMONT	1263	2.7	0.2	(2.3,	3.2)	*	1687	3.4	0.2	(3.0,	3.8)
VIRGINIA	1452	2.3	0.2	(1.9,	2.6)	*	1950	3.4	0.3	(2.9,	3.9)
WASHINGTON	1564	2.6	0.2	(2.3,	3.0)	*	1963	3.5	0.2	(3.1,	3.9)
WEST VIRGINIA	939	3.2	0.3	(2.6,	3.7)	*	1459	3.6	0.2	(3.2,	4.1)
WISCONSIN	973	2.5	0.2	(2.1,	3.0)	*	1211	3.3	0.3	(2.8,	3.8)
WYOMING	1016	2.6	0.3	(2.0,	3.3)	*	1397	3.1	0.2	(2.7,	3.5)
PUERTO RICO	1062	3.0	0.2	(2.6,	3.5)	*	1699	4.1	0.2	(3.6,	4.5)

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 2.6
 RANGE = 1.1- 3.6

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.5
 RANGE = 1.2- 4.9

June 18, 1999

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

TABLE 2.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

6

MEAN DIFFERENCES FROM 1997 TO 1998

PHYSICAL HEALTH NOT GOOD*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997	1998	DIFFERENCE
	MEAN	MEAN	
ALABAMA	3.8	4.2	0.4
ALASKA	2.9	2.8	-0.1
ARIZONA	1.1	1.1	0.1
ARKANSAS	3.4	3.6	0.2
CALIFORNIA	3.1	3.1	-0.1
COLORADO	2.8	2.9	0.1
CONNECTICUT	3.1	2.9	-0.2
DELAWARE	3.7	3.2	-0.5
DISTRICT OF CO	1.5	2.2	0.7
FLORIDA	3.5	3.4	-0.2
GEORGIA	2.2	3.0	0.8
HAWAII	3.0	2.2	-0.8
IDAHO	3.4	3.1	-0.4
ILLINOIS	3.3	2.8	-0.5
INDIANA	3.4	3.3	-0.1
IOWA	2.9	2.9	0.0
KANSAS	2.1	2.8	0.7
KENTUCKY	4.1	3.6	-0.5
LOUISIANA	3.2	3.0	-0.2
MAINE	3.5	3.3	-0.2
MARYLAND	2.4	3.0	0.6
MASSACHUSETTS	3.3	2.8	-0.5
MICHIGAN	3.3	3.7	0.5
MINNESOTA	3.0	3.2	0.2
MISSISSIPPI	3.1	3.7	0.6
MISSOURI	3.5	3.4	-0.1
MONTANA	2.7	3.0	0.3
NEBRASKA	2.9	3.1	0.2
NEVADA	3.5	3.2	-0.4
NEW HAMPSHIRE	3.1	3.0	-0.0
NEW JERSEY	3.0	2.8	-0.3
NEW MEXICO	3.2	3.5	0.3
NEW YORK	3.1	3.2	0.1
NORTH CAROLINA	2.9	3.7	0.8
NORTH DAKOTA	2.9	2.7	-0.2
OHIO	2.6	3.1	0.5
OKLAHOMA	2.3	2.6	0.4
OREGON	3.2	3.3	0.1
PENNSYLVANIA	3.2	3.3	0.2
RHODE ISLAND	3.0	3.4	0.3
SOUTH CAROLINA	2.9	3.1	0.2
SOUTH DAKOTA	2.6	2.6	-0.1
TENNESSEE	3.4	3.3	-0.1
TEXAS	3.2	3.3	0.1
UTAH	3.6	3.4	-0.2
VERMONT	2.7	3.1	0.3
VIRGINIA	2.4	2.8	0.4
WASHINGTON	3.1	3.1	0.0
WEST VIRGINIA	3.4	3.4	0.1
WISCONSIN	2.8	2.9	0.2
WYOMING	2.6	2.9	0.2
PUERTO RICO	2.8	3.6	0.7

* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

June 18, 1999

TABLE 3.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD*

7

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2181	3.7	0.2	(3.3, 4.0)
ALASKA	1956	2.3	0.2	(2.0, 2.6)
ARIZONA	1913	0.8	0.1	(0.5, 1.0)
ARKANSAS	2937	3.6	0.2	(3.3, 4.0)
CALIFORNIA	4126	3.0	0.1	(2.8, 3.3)
COLORADO	1785	3.0	0.2	(2.7, 3.4)
CONNECTICUT	3075	2.8	0.2	(2.5, 3.1)
DELAWARE	2548	3.1	0.2	(2.6, 3.6)
DISTRICT OF CO	1436	2.2	0.2	(1.8, 2.5)
FLORIDA	4671	3.0	0.1	(2.7, 3.2)
GEORGIA	2315	3.4	0.2	(3.0, 3.7)
HAWAII	2132	2.2	0.2	(1.8, 2.6)
IDAHO	4892	2.8	0.1	(2.6, 3.1)
ILLINOIS	2860	2.8	0.1	(2.5, 3.1)
INDIANA	2398	3.2	0.2	(2.9, 3.5)
IOWA	3628	3.0	0.1	(2.7, 3.2)
KANSAS	3738	2.7	0.1	(2.4, 3.0)
KENTUCKY	3680	4.5	0.2	(4.1, 4.8)
LOUISIANA	1608	2.8	0.2	(2.5, 3.2)
MAINE	1603	2.7	0.2	(2.2, 3.1)
MARYLAND	3239	3.4	0.2	(3.0, 3.7)
MASSACHUSETTS	4864	2.8	0.1	(2.5, 3.0)
MICHIGAN	2501	3.6	0.2	(3.3, 4.0)
MINNESOTA	4173	3.2	0.1	(2.9, 3.4)
MISSISSIPPI	2210	3.3	0.2	(2.9, 3.7)
MISSOURI	3675	3.1	0.2	(2.8, 3.4)
MONTANA	1757	2.6	0.2	(2.3, 3.0)
NEBRASKA	2689	3.0	0.2	(2.7, 3.2)
NEVADA	1948	3.5	0.2	(3.0, 4.0)
NEW HAMPSHIRE	1462	3.0	0.2	(2.5, 3.4)
NEW JERSEY	2436	2.7	0.2	(2.4, 3.1)
NEW MEXICO	3614	3.3	0.1	(3.1, 3.6)
NEW YORK	2466	3.0	0.2	(2.7, 3.2)
NORTH CAROLINA	2367	2.7	0.2	(2.3, 3.1)
NORTH DAKOTA	1799	2.5	0.1	(2.2, 2.8)
OHIO	2960	2.9	0.2	(2.5, 3.2)
OKLAHOMA	2489	1.9	0.1	(1.7, 2.2)
OREGON	1806	3.3	0.2	(2.9, 3.7)
PENNSYLVANIA	3544	2.9	0.1	(2.7, 3.2)
RHODE ISLAND	3524	3.0	0.1	(2.8, 3.3)
SOUTH CAROLINA	3233	3.0	0.2	(2.6, 3.3)
SOUTH DAKOTA	2101	2.5	0.2	(2.1, 2.8)
TENNESSEE	2876	3.5	0.2	(3.2, 3.8)
TEXAS	5944	3.5	0.1	(3.2, 3.7)
UTAH	2843	3.7	0.2	(3.3, 4.0)
VERMONT	2946	2.8	0.2	(2.5, 3.1)
VIRGINIA	3397	3.0	0.2	(2.6, 3.3)
WASHINGTON	3524	3.1	0.1	(2.9, 3.4)
WEST VIRGINIA	2401	2.3	0.2	(2.0, 2.6)
WISCONSIN	2179	2.9	0.2	(2.6, 3.3)
WYOMING	2411	3.1	0.1	(2.8, 3.3)
PUERTO RICO	2735	2.6	0.1	(2.3, 2.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.0 RANGE = 0.8- 4.5

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

June 18, 1999

TABLE 3.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE	MEAN	STANDARD	95% CONFIDENCE	*	SAMPLE	MEAN	STANDARD	95% CONFIDENCE	*
	SIZE		ERROR	INTERVAL	*	SIZE		ERROR	INTERVAL	*
ALABAMA	762	2.9	0.3	(2.4, 3.5)	*	1419	4.3	0.3	(3.8, 4.8)	*
ALASKA	910	1.7	0.2	(1.3, 2.1)	*	1046	3.0	0.3	(2.5, 3.5)	*
ARIZONA	823	0.4	0.1	(0.2, 0.5)	*	1090	1.2	0.2	(0.7, 1.7)	*
ARKANSAS	1131	3.4	0.3	(2.9, 3.9)	*	1806	3.8	0.2	(3.4, 4.3)	*
CALIFORNIA	1806	2.7	0.2	(2.3, 3.0)	*	2320	3.4	0.2	(3.1, 3.7)	*
COLORADO	800	2.3	0.2	(1.8, 2.7)	*	985	3.7	0.3	(3.2, 4.3)	*
CONNECTICUT	1264	2.4	0.3	(1.9, 2.9)	*	1811	3.2	0.2	(2.8, 3.6)	*
DELAWARE	1029	2.5	0.3	(1.9, 3.2)	*	1519	3.7	0.3	(3.0, 4.3)	*
DISTRICT OF CO	538	1.9	0.3	(1.4, 2.4)	*	898	2.4	0.3	(1.9, 2.9)	*
FLORIDA	1871	2.1	0.2	(1.8, 2.4)	*	2800	3.7	0.2	(3.4, 4.1)	*
GEORGIA	878	2.7	0.3	(2.2, 3.2)	*	1437	4.0	0.2	(3.5, 4.4)	*
HAWAII	925	1.8	0.3	(1.2, 2.4)	*	1207	2.6	0.2	(2.1, 3.0)	*
IDAHO	2067	2.2	0.2	(1.9, 2.6)	*	2825	3.4	0.2	(3.1, 3.7)	*
ILLINOIS	1228	2.1	0.2	(1.7, 2.4)	*	1632	3.5	0.2	(3.1, 3.9)	*
INDIANA	980	2.9	0.2	(2.4, 3.4)	*	1418	3.5	0.2	(3.1, 3.9)	*
IOWA	1441	2.3	0.2	(1.9, 2.6)	*	2187	3.6	0.2	(3.3, 4.0)	*
KANSAS	1464	2.4	0.2	(2.0, 2.7)	*	2274	3.0	0.2	(2.7, 3.4)	*
KENTUCKY	1350	3.3	0.2	(2.8, 3.8)	*	2330	5.5	0.2	(5.0, 6.0)	*
LOUISIANA	640	2.4	0.3	(1.8, 2.9)	*	968	3.3	0.3	(2.8, 3.7)	*
MAINE	665	1.8	0.2	(1.4, 2.3)	*	938	3.4	0.3	(2.8, 4.1)	*
MARYLAND	1277	2.5	0.2	(2.0, 3.0)	*	1962	4.1	0.2	(3.7, 4.6)	*
MASSACHUSETTS	1989	2.2	0.2	(1.9, 2.5)	*	2875	3.2	0.2	(2.9, 3.6)	*
MICHIGAN	1026	2.6	0.2	(2.2, 3.1)	*	1475	4.6	0.2	(4.1, 5.0)	*
MINNESOTA	1975	2.6	0.2	(2.3, 2.9)	*	2198	3.7	0.2	(3.4, 4.1)	*
MISSISSIPPI	842	2.7	0.3	(2.1, 3.3)	*	1368	3.8	0.2	(3.3, 4.2)	*
MISSOURI	1420	2.3	0.2	(1.8, 2.7)	*	2255	3.9	0.2	(3.5, 4.3)	*
MONTANA	772	1.8	0.2	(1.4, 2.3)	*	985	3.4	0.3	(2.9, 3.9)	*
NEBRASKA	1048	2.5	0.2	(2.0, 3.0)	*	1641	3.3	0.2	(3.0, 3.7)	*
NEVADA	889	2.9	0.3	(2.3, 3.5)	*	1059	4.1	0.4	(3.4, 4.8)	*
NEW HAMPSHIRE	605	2.1	0.3	(1.6, 2.6)	*	857	3.8	0.3	(3.2, 4.4)	*
NEW JERSEY	1002	2.5	0.3	(2.0, 3.0)	*	1434	2.9	0.2	(2.5, 3.3)	*
NEW MEXICO	1565	2.3	0.2	(2.0, 2.6)	*	2049	4.3	0.2	(3.9, 4.7)	*
NEW YORK	1031	2.3	0.2	(1.9, 2.7)	*	1435	3.6	0.2	(3.1, 4.0)	*
NORTH CAROLINA	908	1.6	0.2	(1.2, 2.0)	*	1459	3.7	0.3	(3.2, 4.3)	*
NORTH DAKOTA	805	2.0	0.2	(1.6, 2.3)	*	994	3.0	0.2	(2.6, 3.4)	*
OHIO	1173	2.4	0.3	(1.9, 2.9)	*	1787	3.3	0.3	(2.8, 3.8)	*
OKLAHOMA	996	1.5	0.2	(1.1, 1.8)	*	1493	2.4	0.2	(2.0, 2.7)	*
OREGON	737	3.0	0.3	(2.4, 3.6)	*	1069	3.6	0.2	(3.1, 4.0)	*
PENNSYLVANIA	1405	2.7	0.2	(2.3, 3.1)	*	2139	3.1	0.2	(2.8, 3.5)	*
RHODE ISLAND	1389	2.6	0.2	(2.2, 3.0)	*	2135	3.5	0.2	(3.1, 3.8)	*
SOUTH CAROLINA	1348	2.3	0.2	(1.9, 2.7)	*	1885	3.6	0.3	(3.1, 4.1)	*
SOUTH DAKOTA	923	2.3	0.3	(1.7, 2.9)	*	1178	2.7	0.2	(2.3, 3.1)	*
TENNESSEE	1007	2.7	0.3	(2.2, 3.2)	*	1869	4.2	0.2	(3.8, 4.7)	*
TEXAS	2431	2.8	0.2	(2.3, 3.2)	*	3513	4.1	0.2	(3.8, 4.5)	*
UTAH	1231	2.9	0.3	(2.4, 3.4)	*	1612	4.4	0.2	(4.0, 4.9)	*
VERMONT	1262	2.5	0.3	(2.0, 3.0)	*	1684	3.1	0.2	(2.7, 3.4)	*
VIRGINIA	1438	2.0	0.2	(1.7, 2.3)	*	1959	3.9	0.3	(3.3, 4.4)	*
WASHINGTON	1565	2.6	0.2	(2.2, 3.0)	*	1959	3.7	0.2	(3.3, 4.1)	*
WEST VIRGINIA	938	1.6	0.2	(1.2, 2.0)	*	1463	2.9	0.2	(2.4, 3.3)	*
WISCONSIN	968	2.5	0.2	(2.0, 2.9)	*	1211	3.4	0.3	(2.8, 3.9)	*
WYOMING	1016	2.2	0.2	(1.9, 2.6)	*	1395	3.9	0.2	(3.5, 4.3)	*
PUERTO RICO	1050	2.2	0.2	(1.8, 2.6)	*	1685	2.9	0.2	(2.6, 3.3)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 2.4
 RANGE = 0.4- 3.4

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 3.5
 RANGE = 1.2- 5.5

June 18, 1999

TABLE 3.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

9

MEAN DIFFERENCES FROM 1997 TO 1998
 MENTAL HEALTH NOT GOOD*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997	1998	DIFFERENCE
	MEAN	MEAN	
ALABAMA	3.3	3.7	0.4
ALASKA	3.2	2.3	-0.9
ARIZONA	1.5	0.8	-0.7
ARKANSAS	3.0	3.6	0.7
CALIFORNIA	3.4	3.0	-0.4
COLORADO	3.2	3.0	-0.1
CONNECTICUT	2.8	2.8	0.0
DELAWARE	3.2	3.1	-0.0
DISTRICT OF CO	2.1	2.2	0.1
FLORIDA	3.4	3.0	-0.5
GEORGIA	2.8	3.4	0.5
HAWAII	2.6	2.2	-0.4
IDAHO	3.1	2.8	-0.3
ILLINOIS	3.1	2.8	-0.3
INDIANA	3.6	3.2	-0.4
IOWA	2.9	3.0	0.1
KANSAS	1.7	2.7	1.0
KENTUCKY	4.9	4.5	-0.5
LOUISIANA	3.5	2.8	-0.7
MAINE	2.9	2.7	-0.2
MARYLAND	1.8	3.4	1.5
MASSACHUSETTS	3.1	2.8	-0.4
MICHIGAN	3.3	3.6	0.4
MINNESOTA	3.0	3.2	0.2
MISSISSIPPI	2.5	3.3	0.8
MISSOURI	3.2	3.1	-0.1
MONTANA	2.5	2.6	0.2
NEBRASKA	2.5	3.0	0.4
NEVADA	3.7	3.5	-0.2
NEW HAMPSHIRE	3.3	3.0	-0.3
NEW JERSEY	2.9	2.7	-0.2
NEW MEXICO	3.3	3.3	0.1
NEW YORK	3.0	3.0	-0.1
NORTH CAROLINA	2.5	2.7	0.3
NORTH DAKOTA	3.0	2.5	-0.5
OHIO	2.2	2.9	0.7
OKLAHOMA	1.8	1.9	0.2
OREGON	3.2	3.3	0.1
PENNSYLVANIA	2.9	2.9	-0.0
RHODE ISLAND	2.7	3.0	0.3
SOUTH CAROLINA	1.8	3.0	1.2
SOUTH DAKOTA	2.7	2.5	-0.3
TENNESSEE	3.5	3.5	-0.0
TEXAS	2.9	3.5	0.6
UTAH	3.5	3.7	0.2
VERMONT	2.7	2.8	0.1
VIRGINIA	2.7	3.0	0.3
WASHINGTON	3.1	3.1	0.0
WEST VIRGINIA	2.3	2.3	-0.0
WISCONSIN	2.5	2.9	0.4
WYOMING	2.7	3.1	0.4
PUERTO RICO	1.7	2.6	0.9

* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD
 June 18, 1999

TABLE 4.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED*

10

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1240	4.9	0.3	(4.3, 5.5)
ALASKA	1002	2.8	0.3	(2.3, 3.3)
ARIZONA	317	4.2	0.6	(3.1, 5.3)
ARKANSAS	1266	6.1	0.3	(5.4, 6.7)
CALIFORNIA	2128	3.7	0.2	(3.3, 4.0)
COLORADO	947	2.9	0.2	(2.4, 3.4)
CONNECTICUT	1456	3.8	0.3	(3.2, 4.3)
DELAWARE	874	6.3	0.6	(5.2, 7.4)
DISTRICT OF CO	534	5.5	0.4	(4.7, 6.3)
FLORIDA	2086	4.8	0.2	(4.3, 5.3)
GEORGIA	1184	3.3	0.2	(2.9, 3.8)
HAWAII	937	3.2	0.3	(2.7, 3.8)
IDAHO	2606	2.9	0.2	(2.6, 3.2)
ILLINOIS	1518	2.8	0.2	(2.4, 3.2)
INDIANA	1389	3.1	0.2	(2.7, 3.6)
IOWA	1873	2.9	0.2	(2.5, 3.3)
KANSAS	1707	3.3	0.2	(3.0, 3.7)
KENTUCKY	1485	6.6	0.3	(6.0, 7.2)
LOUISIANA	739	4.1	0.3	(3.5, 4.8)
MAINE	747	4.1	0.3	(3.4, 4.8)
MARYLAND	1687	3.6	0.3	(3.1, 4.1)
MASSACHUSETTS	2357	3.5	0.2	(3.1, 3.9)
MICHIGAN	1441	3.6	0.2	(3.2, 4.1)
MINNESOTA	2267	4.2	0.2	(3.8, 4.5)
MISSISSIPPI	1072	4.1	0.3	(3.6, 4.7)
MISSOURI	1911	3.4	0.2	(3.0, 3.8)
MONTANA	814	3.3	0.3	(2.7, 3.8)
NEBRASKA	1338	3.6	0.2	(3.1, 4.0)
NEVADA	1147	3.0	0.3	(2.4, 3.5)
NEW HAMPSHIRE	775	3.2	0.3	(2.6, 3.7)
NEW JERSEY	1087	4.0	0.3	(3.4, 4.6)
NEW MEXICO	1924	3.8	0.2	(3.4, 4.2)
NEW YORK	1244	4.0	0.3	(3.4, 4.5)
NORTH CAROLINA	1119	4.7	0.4	(3.9, 5.4)
NORTH DAKOTA	839	3.0	0.2	(2.5, 3.4)
OHIO	1335	3.8	0.3	(3.2, 4.4)
OKLAHOMA	826	4.9	0.3	(4.3, 5.6)
OREGON	1009	3.4	0.3	(2.9, 3.9)
PENNSYLVANIA	1754	4.0	0.2	(3.5, 4.4)
RHODE ISLAND	1783	3.7	0.2	(3.3, 4.2)
SOUTH CAROLINA	1471	4.1	0.3	(3.5, 4.6)
SOUTH DAKOTA	930	3.1	0.3	(2.6, 3.6)
TENNESSEE	1438	4.1	0.3	(3.6, 4.6)
TEXAS	3093	4.0	0.2	(3.5, 4.4)
UTAH	1726	3.1	0.2	(2.6, 3.5)
VERMONT	1463	2.9	0.2	(2.5, 3.3)
VIRGINIA	1837	3.1	0.2	(2.7, 3.5)
WASHINGTON	2036	3.1	0.2	(2.7, 3.4)
WEST VIRGINIA	940	6.8	0.4	(6.1, 7.5)
WISCONSIN	1257	2.9	0.2	(2.4, 3.4)
WYOMING	1311	2.9	0.2	(2.5, 3.3)
PUERTO RICO	1074	5.6	0.3	(5.0, 6.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.7 RANGE = 2.8- 6.8

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

June 18, 1999

TABLE 4.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

11

PARTICIPANT	MALES					FEMALES					
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL		
	*	*	*	*	*	*	*	*	*	*	
ALABAMA	369	4.6	0.6	(3.5,	5.7)	*	871	5.1	0.3	(4.4,	5.7)
ALASKA	398	3.1	0.5	(2.2,	4.0)	*	604	2.6	0.3	(2.0,	3.2)
ARIZONA	107	5.2	0.9	(3.5,	6.9)	*	210	3.7	0.7	(2.3,	5.0)
ARKANSAS	421	6.7	0.6	(5.5,	7.8)	*	845	5.7	0.4	(4.9,	6.4)
CALIFORNIA	813	3.2	0.3	(2.7,	3.7)	*	1315	4.0	0.3	(3.5,	4.5)
COLORADO	353	2.7	0.4	(2.0,	3.4)	*	594	3.1	0.3	(2.5,	3.7)
CONNECTICUT	529	4.4	0.5	(3.4,	5.4)	*	927	3.2	0.3	(2.6,	3.8)
DELAWARE	291	6.7	1.0	(4.7,	8.7)	*	583	6.0	0.6	(4.8,	7.2)
DISTRICT OF CO	181	5.0	0.7	(3.6,	6.3)	*	353	5.9	0.5	(4.9,	7.0)
FLORIDA	709	4.8	0.4	(4.0,	5.6)	*	1377	4.8	0.3	(4.2,	5.4)
GEORGIA	383	3.3	0.4	(2.4,	4.1)	*	801	3.4	0.3	(2.8,	3.9)
HAWAII	348	3.5	0.5	(2.4,	4.5)	*	589	3.0	0.3	(2.4,	3.6)
IDAHO	935	3.0	0.3	(2.4,	3.5)	*	1671	2.8	0.2	(2.5,	3.2)
ILLINOIS	548	3.1	0.4	(2.3,	3.8)	*	970	2.7	0.2	(2.2,	3.1)
INDIANA	507	3.5	0.4	(2.8,	4.2)	*	882	2.9	0.3	(2.4,	3.4)
IOWA	619	2.9	0.3	(2.3,	3.6)	*	1254	2.8	0.2	(2.4,	3.3)
KANSAS	575	3.3	0.3	(2.7,	3.9)	*	1132	3.3	0.2	(2.9,	3.8)
KENTUCKY	454	6.5	0.6	(5.5,	7.6)	*	1031	6.7	0.4	(5.9,	7.4)
LOUISIANA	254	5.2	0.7	(3.9,	6.5)	*	485	3.4	0.4	(2.7,	4.1)
MAINE	265	4.0	0.6	(2.9,	5.1)	*	482	4.2	0.4	(3.3,	5.0)
MARYLAND	596	3.4	0.4	(2.5,	4.2)	*	1091	3.8	0.3	(3.3,	4.4)
MASSACHUSETTS	888	3.4	0.3	(2.7,	4.0)	*	1469	3.6	0.3	(3.1,	4.1)
MICHIGAN	492	3.7	0.4	(2.9,	4.4)	*	949	3.6	0.3	(3.1,	4.2)
MINNESOTA	980	4.1	0.3	(3.6,	4.7)	*	1287	4.2	0.2	(3.7,	4.6)
MISSISSIPPI	333	4.3	0.5	(3.3,	5.4)	*	739	4.0	0.3	(3.4,	4.6)
MISSOURI	605	3.0	0.3	(2.5,	3.6)	*	1306	3.7	0.3	(3.2,	4.2)
MONTANA	310	3.3	0.5	(2.4,	4.3)	*	504	3.2	0.4	(2.5,	3.9)
NEBRASKA	446	3.3	0.4	(2.6,	4.1)	*	892	3.7	0.3	(3.2,	4.2)
NEVADA	461	3.1	0.5	(2.1,	4.0)	*	686	2.9	0.4	(2.1,	3.6)
NEW HAMPSHIRE	272	2.8	0.5	(1.9,	3.7)	*	503	3.4	0.4	(2.8,	4.1)
NEW JERSEY	384	4.5	0.6	(3.4,	5.6)	*	703	3.7	0.4	(3.0,	4.3)
NEW MEXICO	728	3.6	0.3	(2.9,	4.2)	*	1196	3.9	0.3	(3.4,	4.4)
NEW YORK	469	3.8	0.4	(2.9,	4.6)	*	775	4.1	0.3	(3.4,	4.8)
NORTH CAROLINA	337	5.0	0.7	(3.7,	6.3)	*	782	4.4	0.4	(3.7,	5.1)
NORTH DAKOTA	335	2.3	0.3	(1.7,	2.9)	*	504	3.5	0.4	(2.8,	4.2)
OHIO	443	3.1	0.5	(2.3,	4.0)	*	892	4.3	0.4	(3.4,	5.2)
OKLAHOMA	271	4.8	0.6	(3.6,	5.9)	*	555	5.1	0.4	(4.3,	5.9)
OREGON	367	3.5	0.4	(2.6,	4.3)	*	642	3.4	0.3	(2.7,	4.0)
PENNSYLVANIA	612	4.2	0.4	(3.4,	5.1)	*	1142	3.8	0.3	(3.3,	4.3)
RHODE ISLAND	635	3.8	0.4	(3.1,	4.5)	*	1148	3.7	0.3	(3.2,	4.2)
SOUTH CAROLINA	528	4.5	0.4	(3.7,	5.4)	*	943	3.7	0.4	(2.9,	4.5)
SOUTH DAKOTA	354	3.4	0.4	(2.6,	4.3)	*	576	2.9	0.3	(2.4,	3.5)
TENNESSEE	415	4.5	0.5	(3.5,	5.6)	*	1023	3.8	0.3	(3.3,	4.3)
TEXAS	1059	4.5	0.4	(3.7,	5.4)	*	2034	3.6	0.2	(3.2,	4.0)
UTAH	641	2.7	0.4	(2.0,	3.4)	*	1085	3.3	0.3	(2.8,	3.9)
VERMONT	546	3.2	0.4	(2.4,	4.0)	*	917	2.7	0.2	(2.3,	3.1)
VIRGINIA	659	3.1	0.3	(2.5,	3.6)	*	1178	3.1	0.3	(2.5,	3.6)
WASHINGTON	782	3.4	0.3	(2.8,	4.0)	*	1254	2.8	0.2	(2.4,	3.2)
WEST VIRGINIA	299	7.8	0.7	(6.4,	9.1)	*	641	6.1	0.4	(5.3,	6.9)
WISCONSIN	494	2.9	0.4	(2.2,	3.6)	*	763	2.9	0.3	(2.3,	3.5)
WYOMING	470	3.0	0.4	(2.2,	3.7)	*	841	2.9	0.3	(2.4,	3.3)
PUERTO RICO	335	6.2	0.6	(5.1,	7.3)	*	739	5.2	0.4	(4.5,	5.9)

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 3.5
RANGE = 2.3- 7.8

NUMBER OF PARTICIPANTS = 52
MEDIAN = 3.6
RANGE = 2.6- 6.7

June 18, 1999

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

TABLE 4.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
MEAN DIFFERENCES FROM 1997 TO 1998

12

PARTICIPANT	ACTIVITIES LIMITED*		
	1997 MEAN	1998 MEAN	DIFFERENCE
ALABAMA	4.2	4.9	0.7
ALASKA	2.3	2.8	0.5
ARIZONA	5.2	4.2	-0.9
ARKANSAS	5.9	6.1	0.2
CALIFORNIA	3.4	3.7	0.3
COLORADO	2.9	2.9	0.0
CONNECTICUT	3.7	3.8	0.0
DELAWARE	5.2	6.3	1.1
DISTRICT OF CO	2.0	5.5	3.5
FLORIDA	4.1	4.8	0.7
GEORGIA	3.2	3.3	0.1
HAWAII	3.2	3.2	0.1
IDAHO	3.5	2.9	-0.6
ILLINOIS	3.1	2.8	-0.3
INDIANA	3.1	3.1	0.1
IOWA	3.0	2.9	-0.1
KANSAS	2.3	3.3	1.0
KENTUCKY	7.2	6.6	-0.5
LOUISIANA	4.3	4.1	-0.2
MAINE	4.1	4.1	0.0
MARYLAND	3.5	3.6	0.1
MASSACHUSETTS	3.7	3.5	-0.2
MICHIGAN	3.2	3.6	0.5
MINNESOTA	3.7	4.2	0.4
MISSISSIPPI	3.6	4.1	0.5
MISSOURI	3.7	3.4	-0.3
MONTANA	2.8	3.3	0.5
NEBRASKA	3.6	3.6	-0.0
NEVADA	3.5	3.0	-0.6
NEW HAMPSHIRE	3.5	3.2	-0.3
NEW JERSEY	3.7	4.0	0.3
NEW MEXICO	3.8	3.8	0.0
NEW YORK	3.8	4.0	0.2
NORTH CAROLINA	4.0	4.7	0.6
NORTH DAKOTA	3.0	3.0	-0.1
OHIO	3.9	3.8	-0.1
OKLAHOMA	3.6	4.9	1.4
OREGON	2.9	3.4	0.5
PENNSYLVANIA	3.5	4.0	0.5
RHODE ISLAND	3.9	3.7	-0.2
SOUTH CAROLINA	4.9	4.1	-0.8
SOUTH DAKOTA	2.8	3.1	0.3
TENNESSEE	3.9	4.1	0.2
TEXAS	3.7	4.0	0.3
UTAH	3.2	3.1	-0.2
VERMONT	3.0	2.9	-0.1
VIRGINIA	2.8	3.1	0.2
WASHINGTON	3.0	3.1	0.1
WEST VIRGINIA	5.9	6.8	0.8
WISCONSIN	2.6	2.9	0.3
WYOMING	2.7	2.9	0.2
PUERTO RICO	4.0	5.6	1.6

* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

June 18, 1999

TABLE 5.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

13

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2244	16.4	1.0	(14.5, 18.2)
ALASKA	1982	20.8	1.3	(18.3, 23.2)
ARIZONA	1911	13.2	1.2	(10.7, 15.6)
ARKANSAS	2999	15.5	0.8	(14.0, 17.0)
CALIFORNIA	4151	18.1	0.7	(16.7, 19.6)
COLORADO	1799	14.9	1.0	(12.9, 16.9)
CONNECTICUT	3121	8.8	0.7	(7.5, 10.1)
DELAWARE	2580	8.0	0.7	(6.5, 9.4)
DISTRICT OF CO	1449	11.6	1.0	(9.6, 13.5)
FLORIDA	4721	17.6	0.7	(16.3, 18.9)
GEORGIA	2371	14.7	0.9	(12.9, 16.5)
HAWAII	2150	5.9	0.6	(4.6, 7.1)
IDAHO	4919	15.2	0.6	(14.0, 16.5)
ILLINOIS	2875	10.9	0.7	(9.6, 12.3)
INDIANA	2420	13.3	0.8	(11.8, 14.9)
IOWA	3645	8.7	0.6	(7.5, 9.8)
KANSAS	3797	10.8	0.6	(9.5, 12.0)
KENTUCKY	3708	14.4	0.7	(12.9, 15.8)
LOUISIANA	1650	22.2	1.3	(19.8, 24.7)
MAINE	1615	13.2	1.0	(11.2, 15.2)
MARYLAND	3286	13.6	0.9	(11.9, 15.4)
MASSACHUSETTS	4929	9.0	0.6	(7.8, 10.1)
MICHIGAN	2533	10.1	0.7	(8.7, 11.5)
MINNESOTA	4219	8.6	0.5	(7.6, 9.5)
MISSISSIPPI	2300	18.7	1.0	(16.8, 20.6)
MISSOURI	3709	13.0	0.7	(11.5, 14.4)
MONTANA	1794	17.2	1.0	(15.2, 19.2)
NEBRASKA	2703	8.0	0.6	(6.8, 9.1)
NEVADA	1970	16.2	1.2	(13.8, 18.6)
NEW HAMPSHIRE	1497	11.7	1.0	(9.8, 13.6)
NEW JERSEY	2457	10.2	0.8	(8.7, 11.7)
NEW MEXICO	3631	21.8	0.8	(20.2, 23.3)
NEW YORK	2519	13.7	0.8	(12.1, 15.4)
NORTH CAROLINA	2395	12.7	1.1	(10.6, 14.8)
NORTH DAKOTA	1800	10.3	0.8	(8.7, 11.9)
OHIO	2986	8.7	0.7	(7.2, 10.1)
OKLAHOMA	2513	18.4	1.0	(16.5, 20.3)
OREGON	1815	13.2	1.0	(11.3, 15.1)
PENNSYLVANIA	3591	10.9	0.6	(9.6, 12.1)
RHODE ISLAND	3600	9.7	0.6	(8.5, 10.9)
SOUTH CAROLINA	3290	16.3	0.9	(14.6, 18.0)
SOUTH DAKOTA	2107	12.3	0.8	(10.7, 14.0)
TENNESSEE	2943	13.0	0.8	(11.5, 14.5)
TEXAS	5982	23.6	0.8	(22.0, 25.3)
UTAH	2851	11.9	0.8	(10.3, 13.5)
VERMONT	2991	10.7	0.7	(9.4, 11.9)
VIRGINIA	3491	12.2	0.8	(10.6, 13.7)
WASHINGTON	3583	10.9	0.7	(9.6, 12.2)
WEST VIRGINIA	2409	17.0	0.9	(15.2, 18.7)
WISCONSIN	2202	8.3	0.7	(6.8, 9.7)
WYOMING	2428	17.1	0.9	(15.4, 18.8)
PUERTO RICO	2812	11.1	0.7	(9.7, 12.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 13.0 RANGE = 5.9-23.6

* ANSWERED "NO" TO: "Do you have any kind of health care coverage, including health insurance,
 prepaid plans such as HMOs, or government plans such as Medicare."

June 18, 1999

TABLE 5.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18 AND OVER* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

14

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	781	17.4	1.6	(14.2, 20.5)	*	1463	15.5	1.1	(13.3, 17.6)	*
ALASKA	919	22.9	2.0	(19.1, 26.7)	*	1063	18.4	1.6	(15.4, 21.5)	*
ARIZONA	821	13.3	1.9	(9.5, 17.1)	*	1090	13.0	1.5	(10.0, 16.1)	*
ARKANSAS	1145	16.0	1.3	(13.5, 18.5)	*	1854	15.1	1.0	(13.2, 16.9)	*
CALIFORNIA	1816	20.2	1.1	(18.0, 22.4)	*	2335	16.1	0.9	(14.3, 17.8)	*
COLORADO	809	17.1	1.7	(13.8, 20.3)	*	990	12.9	1.2	(10.5, 15.3)	*
CONNECTICUT	1282	10.0	1.1	(7.8, 12.1)	*	1839	7.8	0.8	(6.2, 9.4)	*
DELAWARE	1035	9.3	1.3	(6.8, 11.8)	*	1545	6.8	0.8	(5.2, 8.3)	*
DISTRICT OF CO	543	15.2	1.8	(11.7, 18.7)	*	906	8.4	1.0	(6.4, 10.4)	*
FLORIDA	1884	19.1	1.1	(16.9, 21.3)	*	2837	16.2	0.8	(14.6, 17.8)	*
GEORGIA	892	14.6	1.5	(11.6, 17.5)	*	1479	14.8	1.1	(12.6, 17.0)	*
HAWAII	931	5.7	0.9	(3.9, 7.4)	*	1219	6.1	0.9	(4.3, 7.9)	*
IDAHO	2074	17.0	1.0	(15.0, 19.0)	*	2845	13.6	0.7	(12.2, 15.0)	*
ILLINOIS	1228	11.8	1.1	(9.7, 13.9)	*	1647	10.1	0.9	(8.5, 11.8)	*
INDIANA	989	14.3	1.3	(11.9, 16.8)	*	1431	12.5	1.0	(10.5, 14.4)	*
IOWA	1446	9.4	1.0	(7.4, 11.3)	*	2199	8.0	0.7	(6.6, 9.5)	*
KANSAS	1484	12.5	1.1	(10.4, 14.5)	*	2313	9.2	0.7	(7.8, 10.6)	*
KENTUCKY	1358	15.2	1.2	(12.8, 17.7)	*	2350	13.6	0.8	(11.9, 15.2)	*
LOUISIANA	646	21.2	2.0	(17.2, 25.1)	*	1004	23.2	1.6	(20.0, 26.3)	*
MAINE	668	15.6	1.7	(12.3, 18.9)	*	947	11.0	1.2	(8.7, 13.3)	*
MARYLAND	1298	14.3	1.4	(11.5, 17.1)	*	1988	13.0	1.1	(11.0, 15.1)	*
MASSACHUSETTS	2008	11.6	1.0	(9.7, 13.5)	*	2921	6.6	0.6	(5.4, 7.9)	*
MICHIGAN	1036	11.8	1.2	(9.5, 14.1)	*	1497	8.6	0.8	(7.0, 10.2)	*
MINNESOTA	1989	10.4	0.8	(8.9, 11.9)	*	2230	6.8	0.6	(5.6, 8.1)	*
MISSISSIPPI	867	18.6	1.6	(15.5, 21.7)	*	1433	18.8	1.2	(16.5, 21.2)	*
MISSOURI	1430	13.3	1.2	(11.0, 15.5)	*	2279	12.7	0.9	(10.9, 14.6)	*
MONTANA	781	19.9	1.7	(16.6, 23.1)	*	1013	14.7	1.2	(12.3, 17.0)	*
NEBRASKA	1049	9.1	1.0	(7.1, 11.1)	*	1654	6.9	0.7	(5.5, 8.2)	*
NEVADA	891	15.3	1.7	(12.1, 18.6)	*	1079	17.1	1.8	(13.5, 20.7)	*
NEW HAMPSHIRE	616	14.3	1.6	(11.2, 17.4)	*	881	9.2	1.1	(7.0, 11.4)	*
NEW JERSEY	1004	10.5	1.2	(8.2, 12.7)	*	1453	9.9	1.0	(8.0, 11.9)	*
NEW MEXICO	1566	20.3	1.2	(18.0, 22.6)	*	2065	23.1	1.1	(20.9, 25.3)	*
NEW YORK	1049	16.8	1.4	(14.0, 19.6)	*	1470	11.0	0.9	(9.1, 12.8)	*
NORTH CAROLINA	918	14.0	1.9	(10.2, 17.8)	*	1477	11.5	1.0	(9.5, 13.6)	*
NORTH DAKOTA	803	11.5	1.2	(9.1, 14.0)	*	997	9.1	1.0	(7.1, 11.0)	*
OHIO	1184	7.8	1.1	(5.7, 9.9)	*	1802	9.4	1.0	(7.4, 11.4)	*
OKLAHOMA	1004	18.8	1.6	(15.7, 21.8)	*	1509	18.0	1.2	(15.8, 20.3)	*
OREGON	740	13.1	1.4	(10.4, 15.9)	*	1075	13.2	1.4	(10.6, 15.9)	*
PENNSYLVANIA	1422	12.4	1.1	(10.2, 14.5)	*	2169	9.5	0.7	(8.1, 10.9)	*
RHODE ISLAND	1413	11.7	1.0	(9.7, 13.6)	*	2187	8.0	0.7	(6.6, 9.3)	*
SOUTH CAROLINA	1371	17.1	1.4	(14.5, 19.8)	*	1919	15.5	1.1	(13.3, 17.6)	*
SOUTH DAKOTA	924	14.4	1.4	(11.8, 17.1)	*	1183	10.3	1.0	(8.4, 12.3)	*
TENNESSEE	1027	14.7	1.3	(12.2, 17.2)	*	1916	11.5	0.9	(9.7, 13.3)	*
TEXAS	2439	24.1	1.3	(21.5, 26.7)	*	3543	23.2	1.0	(21.2, 25.2)	*
UTAH	1231	11.9	1.3	(9.4, 14.3)	*	1620	11.9	1.1	(9.8, 14.0)	*
VERMONT	1276	12.6	1.1	(10.5, 14.6)	*	1715	8.9	0.8	(7.3, 10.5)	*
VIRGINIA	1478	12.7	1.1	(10.5, 14.8)	*	2013	11.7	1.1	(9.5, 13.9)	*
WASHINGTON	1575	12.6	1.1	(10.5, 14.8)	*	2008	9.3	0.8	(7.8, 10.8)	*
WEST VIRGINIA	937	18.1	1.4	(15.3, 20.8)	*	1472	16.0	1.1	(13.8, 18.2)	*
WISCONSIN	977	8.5	1.2	(6.2, 10.7)	*	1225	8.1	1.0	(6.3, 10.0)	*
WYOMING	1019	17.1	1.4	(14.5, 19.8)	*	1409	17.1	1.1	(15.0, 19.2)	*
PUERTO RICO	1077	13.2	1.2	(10.8, 15.6)	*	1735	9.3	0.8	(7.8, 10.9)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 14.3
RANGE = 5.7-24.1

NUMBER OF PARTICIPANTS = 52
MEDIAN = 11.6
RANGE = 6.1-23.2

* ANSWERED "NO" TO: "Do you have any kind of health care coverage, including health insurance, prepaid plans such as HMOs, or government plans such as Medicare."

June 18, 1999

TABLE 5.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NO HEALTH INSURANCE, AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

15

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.4	16.4	2.0
ALASKA	21.3	20.8	-0.5
ARIZONA	14.7	13.2	-1.5
ARKANSAS	17.3	15.5	-1.8
CALIFORNIA	19.3	18.1	-1.2
COLORADO	11.9	14.9	3.1
CONNECTICUT	9.4	8.8	-0.6
DELAWARE	10.8	8.0	-2.8
DISTRICT OF CO	11.6	11.6	-0.0
FLORIDA	17.6	17.6	0.0
GEORGIA	12.0	14.7	2.7
HAWAII	6.4	5.9	-0.5
IDAHO	16.6	15.2	-1.4
ILLINOIS	11.3	10.9	-0.4
INDIANA	12.4	13.3	1.0
IOWA	9.6	8.7	-0.9
KANSAS	9.5	10.8	1.3
KENTUCKY	13.8	14.4	0.6
LOUISIANA	20.4	22.2	1.8
MAINE	12.0	13.2	1.2
MARYLAND	10.2	13.6	3.4
MASSACHUSETTS	9.3	9.0	-0.3
MICHIGAN	9.9	10.1	0.2
MINNESOTA	8.4	8.6	0.2
MISSISSIPPI	15.2	18.7	3.6
MISSOURI	12.2	13.0	0.8
MONTANA	14.6	17.2	2.6
NEBRASKA	7.6	8.0	0.4
NEVADA	14.1	16.2	2.1
NEW HAMPSHIRE	10.3	11.7	1.4
NEW JERSEY	11.7	10.2	-1.5
NEW MEXICO	22.2	21.8	-0.4
NEW YORK	14.0	13.7	-0.3
NORTH CAROLINA	14.7	12.7	-2.0
NORTH DAKOTA	11.7	10.3	-1.4
OHIO	10.5	8.7	-1.8
OKLAHOMA	17.0	18.4	1.4
OREGON	12.5	13.2	0.6
PENNSYLVANIA	9.4	10.9	1.4
RHODE ISLAND	10.7	9.7	-1.0
SOUTH CAROLINA	15.0	16.3	1.2
SOUTH DAKOTA	13.2	12.3	-0.9
TENNESSEE	11.7	13.0	1.3
TEXAS	24.2	23.6	-0.6
UTAH	11.2	11.9	0.7
VERMONT	14.4	10.7	-3.7
VIRGINIA	11.3	12.2	0.9
WASHINGTON	10.9	10.9	0.0
WEST VIRGINIA	18.1	17.0	-1.1
WISCONSIN	9.3	8.3	-1.0
WYOMING	19.0	17.1	-1.8
PUERTO RICO	10.3	11.1	0.9

ANSWERED "NO" TO: "Do you have any kind of health care coverage, including health insurance,
 prepaid plans such as HMOs, or government plans such as Medicare."

June 18, 1999

TABLE 6.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

16

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1737	19.5	1.1	(17.3, 21.7)
ALASKA	1800	21.8	1.3	(19.2, 24.4)
ARIZONA	1382	15.8	1.5	(12.9, 18.7)
ARKANSAS	2315	19.1	1.0	(17.2, 21.0)
CALIFORNIA	3525	21.2	0.8	(19.6, 22.9)
COLORADO	1510	17.3	1.2	(15.0, 19.6)
CONNECTICUT	2484	10.6	0.8	(9.0, 12.2)
DELAWARE	2031	9.7	0.9	(7.9, 11.4)
DISTRICT OF CO	1167	13.0	1.2	(10.7, 15.2)
FLORIDA	3493	22.4	0.8	(20.8, 24.0)
GEORGIA	1992	16.9	1.1	(14.8, 19.0)
HAWAII	1772	7.0	0.8	(5.5, 8.5)
IDAHO	3976	18.3	0.7	(16.8, 19.7)
ILLINOIS	2404	13.0	0.8	(11.4, 14.6)
INDIANA	1964	15.9	0.9	(14.1, 17.8)
IOWA	2792	10.8	0.7	(9.3, 12.2)
KANSAS	2850	13.0	0.8	(11.5, 14.5)
KENTUCKY	2894	17.3	0.9	(15.6, 19.1)
LOUISIANA	1341	25.9	1.5	(23.0, 28.7)
MAINE	1254	15.7	1.2	(13.3, 18.1)
MARYLAND	2714	15.7	1.0	(13.7, 17.7)
MASSACHUSETTS	3983	10.5	0.7	(9.1, 11.9)
MICHIGAN	2101	11.9	0.8	(10.3, 13.5)
MINNESOTA	3502	9.9	0.6	(8.8, 11.0)
MISSISSIPPI	1802	22.5	1.2	(20.2, 24.7)
MISSOURI	2842	15.9	0.9	(14.1, 17.6)
MONTANA	1403	21.3	1.3	(18.9, 23.8)
NEBRASKA	1955	9.8	0.8	(8.3, 11.3)
NEVADA	1634	19.2	1.5	(16.3, 22.0)
NEW HAMPSHIRE	1208	13.7	1.2	(11.4, 15.9)
NEW JERSEY	1969	11.8	0.9	(10.0, 13.6)
NEW MEXICO	2951	25.7	0.9	(23.8, 27.5)
NEW YORK	2020	16.6	1.0	(14.6, 18.5)
NORTH CAROLINA	1873	15.2	1.3	(12.7, 17.7)
NORTH DAKOTA	1356	12.7	1.0	(10.7, 14.6)
OHIO	2276	10.4	0.9	(8.7, 12.2)
OKLAHOMA	1876	22.4	1.2	(20.1, 24.6)
OREGON	1510	15.8	1.2	(13.5, 18.1)
PENNSYLVANIA	2835	12.9	0.8	(11.4, 14.4)
RHODE ISLAND	2877	11.7	0.7	(10.2, 13.1)
SOUTH CAROLINA	2719	18.7	1.0	(16.8, 20.7)
SOUTH DAKOTA	1549	15.2	1.0	(13.1, 17.2)
TENNESSEE	2363	14.8	0.9	(13.1, 16.6)
TEXAS	5087	27.5	0.9	(25.7, 29.4)
UTAH	2378	13.8	0.9	(12.0, 15.6)
VERMONT	2393	12.4	0.8	(10.9, 13.9)
VIRGINIA	2871	13.8	0.9	(12.1, 15.6)
WASHINGTON	3014	12.8	0.8	(11.3, 14.3)
WEST VIRGINIA	1838	20.6	1.1	(18.5, 22.7)
WISCONSIN	1800	10.0	0.9	(8.2, 11.8)
WYOMING	2011	20.2	1.0	(18.2, 22.2)
PUERTO RICO	2307	12.7	0.8	(11.1, 14.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 15.4 RANGE = 7.0-27.5 YEAR 2000 OBJECTIVE 21.4

* ANSWERED "NO" TO: "Do you have any kind of health care coverage, including health insurance,
 prepaid plans such as HMOs, or government plans such as Medicare" AMONG RESPONDENTS 18-64 YEARS OF AGE

June

TABLE 6.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18-64 YEARS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

17

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	630	20.1	1.9	(16.4, 23.8)	*	1107	19.0	1.3	(16.3, 21.6)	*
ALASKA	839	23.7	2.1	(19.7, 27.7)	*	961	19.7	1.7	(16.4, 23.0)	*
ARIZONA	613	15.3	2.2	(11.0, 19.7)	*	769	16.3	1.9	(12.5, 20.0)	*
ARKANSAS	929	19.1	1.5	(16.1, 22.0)	*	1386	19.2	1.2	(16.8, 21.5)	*
CALIFORNIA	1581	23.1	1.3	(20.6, 25.6)	*	1944	19.3	1.1	(17.2, 21.3)	*
COLORADO	690	19.4	1.9	(15.7, 23.1)	*	820	15.2	1.4	(12.4, 18.0)	*
CONNECTICUT	1039	11.5	1.3	(9.0, 14.0)	*	1445	9.7	1.1	(7.6, 11.7)	*
DELAWARE	853	10.9	1.5	(8.0, 13.8)	*	1178	8.5	1.0	(6.6, 10.4)	*
DISTRICT OF CO	462	16.1	2.0	(12.2, 19.9)	*	705	10.1	1.2	(7.7, 12.6)	*
FLORIDA	1443	23.6	1.3	(21.0, 26.2)	*	2050	21.3	1.0	(19.2, 23.3)	*
GEORGIA	805	16.4	1.7	(13.1, 19.7)	*	1187	17.4	1.3	(14.8, 20.0)	*
HAWAII	798	6.5	1.0	(4.5, 8.5)	*	974	7.5	1.1	(5.3, 9.7)	*
IDAHO	1745	19.8	1.2	(17.4, 22.1)	*	2231	16.8	0.9	(15.0, 18.5)	*
ILLINOIS	1061	13.4	1.2	(11.0, 15.8)	*	1343	12.5	1.0	(10.5, 14.6)	*
INDIANA	838	16.5	1.5	(13.6, 19.3)	*	1126	15.4	1.2	(13.0, 17.8)	*
IOWA	1175	11.1	1.2	(8.9, 13.4)	*	1617	10.4	0.9	(8.6, 12.2)	*
KANSAS	1166	14.4	1.2	(12.0, 16.9)	*	1684	11.5	0.9	(9.7, 13.3)	*
KENTUCKY	1107	17.8	1.4	(15.0, 20.6)	*	1787	16.9	1.0	(14.9, 18.9)	*
LOUISIANA	555	23.7	2.2	(19.4, 28.1)	*	786	27.9	1.9	(24.2, 31.7)	*
MAINE	539	17.7	1.9	(13.9, 21.4)	*	715	13.8	1.5	(10.9, 16.7)	*
MARYLAND	1111	16.3	1.6	(13.1, 19.5)	*	1603	15.2	1.3	(12.7, 17.6)	*
MASSACHUSETTS	1699	13.2	1.1	(10.9, 15.4)	*	2284	7.9	0.8	(6.4, 9.5)	*
MICHIGAN	906	13.3	1.3	(10.7, 15.9)	*	1195	10.5	1.0	(8.5, 12.4)	*
MINNESOTA	1740	11.7	0.9	(10.0, 13.4)	*	1762	8.0	0.8	(6.6, 9.5)	*
MISSISSIPPI	733	21.6	1.8	(18.0, 25.2)	*	1069	23.3	1.5	(20.4, 26.1)	*
MISSOURI	1155	15.4	1.4	(12.7, 18.0)	*	1687	16.3	1.2	(14.0, 18.6)	*
MONTANA	626	24.0	2.0	(20.1, 27.8)	*	777	18.7	1.5	(15.7, 21.6)	*
NEBRASKA	792	10.9	1.2	(8.5, 13.3)	*	1163	8.8	0.9	(7.0, 10.5)	*
NEVADA	771	17.6	1.9	(13.9, 21.4)	*	863	20.8	2.2	(16.4, 25.2)	*
NEW HAMPSHIRE	507	16.0	1.8	(12.5, 19.6)	*	701	11.3	1.4	(8.5, 14.1)	*
NEW JERSEY	827	11.9	1.4	(9.2, 14.5)	*	1142	11.7	1.2	(9.2, 14.1)	*
NEW MEXICO	1333	23.4	1.3	(20.7, 26.0)	*	1618	27.9	1.3	(25.3, 30.5)	*
NEW YORK	857	19.9	1.7	(16.7, 23.2)	*	1163	13.3	1.2	(11.1, 15.6)	*
NORTH CAROLINA	732	16.2	2.2	(11.8, 20.5)	*	1141	14.2	1.3	(11.7, 16.7)	*
NORTH DAKOTA	636	13.9	1.5	(11.0, 16.8)	*	720	11.3	1.3	(8.8, 13.8)	*
OHIO	947	9.2	1.3	(6.7, 11.6)	*	1329	11.7	1.3	(9.2, 14.1)	*
OKLAHOMA	777	22.0	1.8	(18.4, 25.5)	*	1099	22.7	1.5	(19.9, 25.6)	*
OREGON	636	15.3	1.6	(12.1, 18.4)	*	874	16.4	1.7	(13.1, 19.6)	*
PENNSYLVANIA	1197	13.8	1.3	(11.4, 16.3)	*	1638	12.0	0.9	(10.2, 13.7)	*
RHODE ISLAND	1197	13.6	1.2	(11.3, 15.9)	*	1680	9.8	0.9	(8.0, 11.5)	*
SOUTH CAROLINA	1138	19.2	1.6	(16.1, 22.2)	*	1581	18.3	1.3	(15.8, 20.9)	*
SOUTH DAKOTA	693	16.8	1.6	(13.7, 19.9)	*	856	13.5	1.3	(11.0, 16.0)	*
TENNESSEE	873	16.0	1.4	(13.2, 18.7)	*	1490	13.8	1.1	(11.6, 15.9)	*
TEXAS	2142	27.1	1.5	(24.2, 30.0)	*	2945	28.0	1.2	(25.6, 30.3)	*
UTAH	1053	13.5	1.4	(10.7, 16.3)	*	1325	14.1	1.3	(11.6, 16.6)	*
VERMONT	1062	14.2	1.2	(11.8, 16.5)	*	1331	10.6	1.0	(8.6, 12.5)	*
VIRGINIA	1270	14.1	1.3	(11.6, 16.5)	*	1601	13.6	1.3	(11.0, 16.1)	*
WASHINGTON	1368	14.4	1.2	(12.0, 16.9)	*	1646	11.2	0.9	(9.3, 13.0)	*
WEST VIRGINIA	748	21.1	1.6	(17.9, 24.3)	*	1090	20.0	1.4	(17.3, 22.7)	*
WISCONSIN	818	10.1	1.4	(7.4, 12.8)	*	982	9.9	1.2	(7.6, 12.2)	*
WYOMING	880	19.4	1.6	(16.3, 22.5)	*	1131	21.0	1.3	(18.4, 23.5)	*
PUERTO RICO	891	15.1	1.4	(12.4, 17.9)	*	1416	10.5	0.9	(8.7, 12.3)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS = 52	NUMBER OF PARTICIPANTS = 52
MEDIAN	= 16.0	MEDIAN = 14.0
RANGE	= 6.5-27.1	RANGE = 7.5-28.0

* ANSWERED "NO" TO: "Do you have any kind of health care coverage, including health insurance, prepaid plans such as HMOs, or government plans such as Medicare" AMONG RESPONDENTS 18-64 YEARS OF AGE

June 18, 1999

TABLE 6.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NO HEALTH INSURANCE, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

18

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	17.3	19.5	2.2
ALASKA	22.8	21.8	-1.0
ARIZONA	17.9	15.8	-2.1
ARKANSAS	21.6	19.1	-2.5
CALIFORNIA	22.7	21.2	-1.5
COLORADO	14.0	17.3	3.3
CONNECTICUT	10.8	10.6	-0.3
DELAWARE	13.0	9.7	-3.4
DISTRICT OF CO	13.2	13.0	-0.3
FLORIDA	22.6	22.4	-0.1
GEORGIA	13.7	16.9	3.2
HAWAII	7.4	7.0	-0.4
IDAHO	20.1	18.3	-1.9
ILLINOIS	13.5	13.0	-0.5
INDIANA	14.6	15.9	1.4
IOWA	12.0	10.8	-1.3
KANSAS	11.5	13.0	1.5
KENTUCKY	16.8	17.3	0.5
LOUISIANA	24.4	25.9	1.5
MAINE	14.3	15.7	1.4
MARYLAND	11.7	15.7	4.0
MASSACHUSETTS	11.0	10.5	-0.5
MICHIGAN	11.8	11.9	0.0
MINNESOTA	9.6	9.9	0.3
MISSISSIPPI	18.3	22.5	4.2
MISSOURI	15.0	15.9	0.8
MONTANA	17.9	21.3	3.5
NEBRASKA	9.5	9.8	0.3
NEVADA	16.7	19.2	2.4
NEW HAMPSHIRE	12.3	13.7	1.3
NEW JERSEY	14.0	11.8	-2.2
NEW MEXICO	25.7	25.7	-0.0
NEW YORK	16.9	16.6	-0.4
NORTH CAROLINA	17.2	15.2	-2.0
NORTH DAKOTA	14.4	12.7	-1.7
OHIO	12.7	10.4	-2.3
OKLAHOMA	20.9	22.4	1.4
OREGON	15.2	15.8	0.6
PENNSYLVANIA	11.5	12.9	1.4
RHODE ISLAND	13.1	11.7	-1.4
SOUTH CAROLINA	17.6	18.7	1.1
SOUTH DAKOTA	16.5	15.2	-1.3
TENNESSEE	13.9	14.8	1.0
TEXAS	28.2	27.5	-0.6
UTAH	12.8	13.8	1.0
VERMONT	16.5	12.4	-4.1
VIRGINIA	12.8	13.8	1.1
WASHINGTON	13.0	12.8	-0.3
WEST VIRGINIA	22.4	20.6	-1.9
WISCONSIN	11.2	10.0	-1.2
WYOMING	22.4	20.2	-2.3
PUERTO RICO	11.8	12.7	0.9

* ANSWERED "NO" TO: "Do you have any kind of health care coverage, including health insurance, prepaid plans such as HMOs, or government plans such as Medicare" AMONG RESPONDENTS 18-64 YEARS OF AGE

June 18,

TABLE 7.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

19

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2250	7.0	0.6	(5.9, 8.1)
ALASKA	1988	3.0	0.5	(2.0, 3.9)
ARIZONA	1916	2.8	0.6	(1.7, 4.0)
ARKANSAS	2998	6.7	0.5	(5.6, 7.7)
CALIFORNIA	4152	5.5	0.4	(4.6, 6.3)
COLORADO	1799	4.6	0.5	(3.6, 5.6)
CONNECTICUT	3125	4.5	0.5	(3.6, 5.4)
DELAWARE	2579	4.3	0.4	(3.5, 5.2)
DISTRICT OF CO	1452	7.1	1.0	(5.1, 9.1)
FLORIDA	4717	6.4	0.4	(5.6, 7.1)
GEORGIA	2371	5.9	0.6	(4.8, 7.0)
HAWAII	2153	5.6	0.7	(4.2, 7.0)
IDAHO	4928	4.3	0.4	(3.6, 5.0)
ILLINOIS	2879	6.2	0.5	(5.2, 7.2)
INDIANA	2422	6.0	0.6	(4.9, 7.1)
IOWA	3653	5.2	0.4	(4.4, 6.1)
KANSAS	3800	4.1	0.4	(3.4, 4.7)
KENTUCKY	3718	5.6	0.4	(4.8, 6.4)
LOUISIANA	1653	6.4	0.7	(5.1, 7.7)
MAINE	1615	3.6	0.5	(2.7, 4.5)
MARYLAND	3294	5.4	0.5	(4.3, 6.5)
MASSACHUSETTS	4936	3.9	0.4	(3.2, 4.5)
MICHIGAN	2533	7.0	0.6	(5.9, 8.1)
MINNESOTA	4220	4.8	0.4	(4.1, 5.4)
MISSISSIPPI	2302	7.6	0.7	(6.4, 8.9)
MISSOURI	3719	5.8	0.5	(4.8, 6.7)
MONTANA	1803	3.6	0.5	(2.7, 4.4)
NEBRASKA	2711	5.2	0.4	(4.3, 6.0)
NEVADA	1979	4.4	0.6	(3.2, 5.7)
NEW HAMPSHIRE	1496	3.9	0.6	(2.8, 5.1)
NEW JERSEY	2462	5.4	0.5	(4.4, 6.4)
NEW MEXICO	3635	5.0	0.4	(4.2, 5.7)
NEW YORK	2522	6.0	0.6	(4.9, 7.1)
NORTH CAROLINA	2401	6.4	0.6	(5.3, 7.4)
NORTH DAKOTA	1802	4.2	0.5	(3.1, 5.2)
OHIO	2981	5.8	0.6	(4.6, 6.9)
OKLAHOMA	2514	7.8	0.6	(6.7, 8.9)
OREGON	1817	5.3	0.6	(4.1, 6.5)
PENNSYLVANIA	3595	5.6	0.4	(4.8, 6.5)
RHODE ISLAND	3599	6.0	0.4	(5.1, 6.8)
SOUTH CAROLINA	3290	5.7	0.5	(4.8, 6.6)
SOUTH DAKOTA	2109	3.1	0.4	(2.3, 3.8)
TENNESSEE	2943	5.9	0.5	(4.9, 7.0)
TEXAS	6001	5.5	0.3	(4.8, 6.2)
UTAH	2863	4.4	0.5	(3.4, 5.4)
VERMONT	2997	4.4	0.4	(3.6, 5.2)
VIRGINIA	3500	4.5	0.4	(3.6, 5.4)
WASHINGTON	3600	4.9	0.4	(4.1, 5.8)
WEST VIRGINIA	2414	6.3	0.5	(5.2, 7.3)
WISCONSIN	2202	4.4	0.5	(3.4, 5.4)
WYOMING	2433	3.7	0.4	(2.9, 4.5)
PUERTO RICO	2805	9.4	0.6	(8.3, 10.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 5.4 RANGE = 2.8- 9.4

* EVER TOLD YOU HAVE DIABETES

June 18, 1999

TABLE 7.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

20

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	784	6.0	0.9	(4.4, 7.7)	*	1466	7.9	0.8	(6.4, 9.4)	*
ALASKA	923	3.3	0.8	(1.7, 4.8)	*	1065	2.6	0.6	(1.6, 3.7)	*
ARIZONA	825	2.7	0.9	(1.0, 4.5)	*	1091	2.9	0.8	(1.4, 4.4)	*
ARKANSAS	1144	7.0	0.8	(5.4, 8.5)	*	1854	6.4	0.7	(5.0, 7.8)	*
CALIFORNIA	1820	5.4	0.6	(4.2, 6.6)	*	2332	5.5	0.6	(4.3, 6.7)	*
COLORADO	810	4.6	0.8	(3.1, 6.1)	*	989	4.6	0.7	(3.2, 6.0)	*
CONNECTICUT	1284	5.2	0.8	(3.7, 6.7)	*	1841	3.8	0.5	(2.8, 4.8)	*
DELAWARE	1037	4.1	0.7	(2.8, 5.4)	*	1542	4.5	0.6	(3.4, 5.7)	*
DISTRICT OF CO	544	7.0	1.9	(3.3, 10.6)	*	908	7.2	1.0	(5.2, 9.1)	*
FLORIDA	1881	6.8	0.6	(5.6, 8.1)	*	2836	5.9	0.5	(4.9, 6.9)	*
GEORGIA	892	6.8	1.0	(4.9, 8.7)	*	1479	5.1	0.6	(3.9, 6.3)	*
HAWAII	933	5.5	1.0	(3.5, 7.4)	*	1220	5.7	1.0	(3.7, 7.6)	*
IDAHO	2082	4.7	0.6	(3.6, 5.8)	*	2846	3.9	0.4	(3.1, 4.8)	*
ILLINOIS	1232	6.5	0.8	(4.9, 8.0)	*	1647	5.9	0.7	(4.6, 7.2)	*
INDIANA	990	6.3	0.8	(4.7, 7.9)	*	1432	5.8	0.8	(4.3, 7.3)	*
IOWA	1450	5.1	0.7	(3.8, 6.4)	*	2203	5.3	0.6	(4.2, 6.5)	*
KANSAS	1485	4.5	0.6	(3.3, 5.7)	*	2315	3.6	0.4	(2.9, 4.4)	*
KENTUCKY	1363	5.4	0.6	(4.2, 6.6)	*	2355	5.7	0.5	(4.8, 6.7)	*
LOUISIANA	647	5.9	1.0	(4.0, 7.8)	*	1006	6.8	0.9	(5.1, 8.5)	*
MAINE	668	2.2	0.6	(1.1, 3.3)	*	947	4.9	0.7	(3.5, 6.2)	*
MARYLAND	1301	5.1	0.8	(3.6, 6.6)	*	1993	5.7	0.8	(4.2, 7.1)	*
MASSACHUSETTS	2014	4.7	0.6	(3.5, 6.0)	*	2922	3.1	0.4	(2.4, 3.8)	*
MICHIGAN	1035	6.3	0.9	(4.6, 8.0)	*	1498	7.6	0.8	(6.1, 9.1)	*
MINNESOTA	1989	4.4	0.5	(3.4, 5.4)	*	2231	5.1	0.5	(4.1, 6.1)	*
MISSISSIPPI	866	7.0	1.0	(5.1, 9.0)	*	1436	8.2	0.9	(6.5, 9.9)	*
MISSOURI	1433	5.9	0.8	(4.3, 7.5)	*	2286	5.6	0.6	(4.5, 6.7)	*
MONTANA	786	3.0	0.6	(1.8, 4.2)	*	1017	4.1	0.6	(2.8, 5.4)	*
NEBRASKA	1054	5.4	0.7	(4.0, 6.8)	*	1657	5.0	0.5	(4.0, 6.0)	*
NEVADA	897	5.2	1.0	(3.3, 7.1)	*	1082	3.6	0.8	(2.2, 5.1)	*
NEW HAMPSHIRE	613	4.2	0.9	(2.3, 6.0)	*	883	3.7	0.8	(2.2, 5.2)	*
NEW JERSEY	1007	5.9	0.8	(4.3, 7.5)	*	1455	4.9	0.6	(3.7, 6.2)	*
NEW MEXICO	1569	5.1	0.6	(3.9, 6.3)	*	2066	4.8	0.5	(3.8, 5.8)	*
NEW YORK	1051	5.4	0.8	(3.8, 6.9)	*	1471	6.7	0.8	(5.1, 8.2)	*
NORTH CAROLINA	921	5.1	0.8	(3.6, 6.6)	*	1480	7.5	0.8	(6.0, 9.1)	*
NORTH DAKOTA	806	4.2	0.9	(2.5, 5.8)	*	996	4.2	0.7	(2.9, 5.5)	*
OHIO	1180	6.4	1.0	(4.5, 8.3)	*	1801	5.3	0.7	(3.9, 6.6)	*
OKLAHOMA	1003	7.8	0.9	(6.1, 9.5)	*	1511	7.8	0.8	(6.3, 9.3)	*
OREGON	741	6.0	1.0	(4.1, 7.9)	*	1076	4.7	0.8	(3.2, 6.2)	*
PENNSYLVANIA	1424	5.2	0.7	(3.9, 6.4)	*	2171	6.0	0.6	(4.9, 7.1)	*
RHODE ISLAND	1412	6.1	0.7	(4.7, 7.4)	*	2187	5.9	0.6	(4.8, 7.0)	*
SOUTH CAROLINA	1373	5.1	0.7	(3.8, 6.4)	*	1917	6.3	0.6	(5.0, 7.6)	*
SOUTH DAKOTA	925	2.6	0.5	(1.6, 3.5)	*	1184	3.6	0.6	(2.5, 4.7)	*
TENNESSEE	1027	5.4	0.9	(3.7, 7.1)	*	1916	6.4	0.6	(5.2, 7.6)	*
TEXAS	2452	4.8	0.5	(3.8, 5.7)	*	3549	6.2	0.5	(5.3, 7.1)	*
UTAH	1238	5.1	0.8	(3.5, 6.6)	*	1625	3.8	0.6	(2.7, 4.8)	*
VERMONT	1279	4.2	0.6	(3.0, 5.3)	*	1718	4.6	0.5	(3.5, 5.6)	*
VIRGINIA	1481	4.1	0.6	(3.0, 5.2)	*	2019	4.8	0.7	(3.5, 6.2)	*
WASHINGTON	1589	4.6	0.6	(3.5, 5.7)	*	2011	5.3	0.7	(4.0, 6.5)	*
WEST VIRGINIA	942	5.8	0.9	(4.1, 7.5)	*	1472	6.7	0.7	(5.4, 8.0)	*
WISCONSIN	976	4.8	0.8	(3.3, 6.3)	*	1226	4.0	0.7	(2.8, 5.3)	*
WYOMING	1022	4.4	0.7	(3.0, 5.7)	*	1411	3.1	0.5	(2.1, 4.0)	*
PUERTO RICO	1074	8.7	0.9	(7.0, 10.4)	*	1731	10.0	0.7	(8.6, 11.4)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 5.2
 RANGE = 2.2- 8.7

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 5.3
 RANGE = 2.6-10.0

June 18, 1999

* EVER TOLD YOU HAVE DIABETES

TABLE 7.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 DIABETES AWARENESS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

21

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	7.0	7.0	0.1
ALASKA	3.3	3.0	-0.4
ARIZONA	3.1	2.8	-0.3
ARKANSAS	5.1	6.7	1.5
CALIFORNIA	5.6	5.5	-0.1
COLORADO	3.9	4.6	0.7
CONNECTICUT	5.2	4.5	-0.7
DELAWARE	6.4	4.3	-2.1
DISTRICT OF CO	4.6	7.1	2.5
FLORIDA	5.7	6.4	0.6
GEORGIA	4.1	5.9	1.8
HAWAII	5.0	5.6	0.6
IDAHO	4.0	4.3	0.3
ILLINOIS	7.0	6.2	-0.9
INDIANA	5.3	6.0	0.8
IOWA	4.6	5.2	0.6
KANSAS	3.0	4.1	1.0
KENTUCKY	5.3	5.6	0.3
LOUISIANA	5.5	6.4	0.9
MAINE	4.9	3.6	-1.3
MARYLAND	5.9	5.4	-0.5
MASSACHUSETTS	4.7	3.9	-0.8
MICHIGAN	5.8	7.0	1.2
MINNESOTA	3.9	4.8	0.9
MISSISSIPPI	6.1	7.6	1.5
MISSOURI	4.8	5.8	1.0
MONTANA	3.2	3.6	0.4
NEBRASKA	4.2	5.2	1.0
NEVADA	4.0	4.4	0.5
NEW HAMPSHIRE	3.9	3.9	-0.0
NEW JERSEY	5.3	5.4	0.2
NEW MEXICO	4.9	5.0	0.0
NEW YORK	4.8	6.0	1.2
NORTH CAROLINA	5.0	6.4	1.4
NORTH DAKOTA	3.5	4.2	0.7
OHIO	4.7	5.8	1.0
OKLAHOMA	5.9	7.8	1.9
OREGON	4.7	5.3	0.6
PENNSYLVANIA	5.1	5.6	0.5
RHODE ISLAND	4.9	6.0	1.1
SOUTH CAROLINA	4.9	5.7	0.8
SOUTH DAKOTA	3.8	3.1	-0.7
TENNESSEE	4.4	5.9	1.6
TEXAS	5.9	5.5	-0.4
UTAH	4.1	4.4	0.3
VERMONT	4.7	4.4	-0.3
VIRGINIA	4.2	4.5	0.3
WASHINGTON	4.1	4.9	0.8
WEST VIRGINIA	6.3	6.3	-0.0
WISCONSIN	4.7	4.4	-0.3
WYOMING	3.0	3.7	0.7
PUERTO RICO	10.5	9.4	-1.1

* EVER TOLD YOU HAVE DIABETES

June 18, 1999

TABLE 8.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

22

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2252	29.7	1.1	(27.5, 32.0)
ALASKA	1988	23.5	1.3	(20.9, 26.1)
ARIZONA	1914	51.3	1.8	(47.9, 54.8)
ARKANSAS	2997	35.9	1.0	(33.9, 37.9)
CALIFORNIA	4161	25.5	0.8	(24.0, 27.1)
COLORADO	1802	21.3	1.1	(19.1, 23.5)
CONNECTICUT	3126	27.1	1.0	(25.1, 29.0)
DELAWARE	2582	35.4	1.3	(32.8, 38.0)
DISTRICT OF CO	1452	38.5	1.5	(35.5, 41.5)
FLORIDA	4722	31.1	0.8	(29.5, 32.7)
GEORGIA	2375	29.6	1.2	(27.4, 31.9)
HAWAII	2153	18.0	1.1	(15.8, 20.1)
IDAHO	4930	20.4	0.7	(19.1, 21.8)
ILLINOIS	1440	27.1	1.3	(24.5, 29.7)
INDIANA	2425	27.1	1.0	(25.1, 29.0)
IOWA	3654	26.7	0.9	(25.0, 28.3)
KANSAS	3802	38.3	0.9	(36.5, 40.1)
KENTUCKY	3717	42.7	0.9	(40.8, 44.5)
LOUISIANA	1655	32.2	1.3	(29.7, 34.7)
MAINE	1615	27.7	1.3	(25.2, 30.2)
MARYLAND	3295	20.3	1.0	(18.4, 22.2)
MASSACHUSETTS	4940	25.4	0.8	(23.8, 26.9)
MICHIGAN	2534	21.4	0.9	(19.6, 23.1)
MINNESOTA	4222	25.5	0.7	(24.0, 26.9)
MISSISSIPPI	2306	33.8	1.1	(31.6, 36.0)
MISSOURI	3720	27.9	1.0	(26.0, 29.8)
MONTANA	1799	25.2	1.1	(23.0, 27.4)
NEBRASKA	2708	26.1	1.0	(24.2, 28.0)
NEVADA	1979	24.1	1.6	(20.9, 27.2)
NEW HAMPSHIRE	1498	24.9	1.3	(22.4, 27.3)
NEW JERSEY	2458	32.6	1.1	(30.4, 34.8)
NEW MEXICO	3637	23.0	0.8	(21.5, 24.6)
NEW YORK	2524	31.0	1.1	(28.9, 33.1)
NORTH CAROLINA	2400	27.7	1.2	(25.4, 30.1)
NORTH DAKOTA	1803	33.1	1.2	(30.7, 35.5)
OHIO	2985	29.8	1.2	(27.5, 32.1)
OKLAHOMA	2516	43.0	1.1	(40.7, 45.2)
OREGON	1818	18.9	1.0	(16.9, 20.9)
PENNSYLVANIA	3597	32.7	0.9	(31.0, 34.5)
RHODE ISLAND	3595	29.9	0.9	(28.2, 31.6)
SOUTH CAROLINA	3293	33.7	1.0	(31.7, 35.6)
SOUTH DAKOTA	2111	33.3	1.2	(30.9, 35.7)
TENNESSEE	2944	35.8	1.0	(33.7, 37.8)
TEXAS	6004	27.9	0.8	(26.4, 29.4)
UTAH	2862	17.1	0.9	(15.3, 19.0)
VERMONT	2998	26.0	0.9	(24.2, 27.8)
VIRGINIA	3504	24.8	1.1	(22.7, 26.8)
WASHINGTON	3600	17.6	0.7	(16.2, 19.0)
WEST VIRGINIA	2414	43.7	1.2	(41.4, 46.0)
WISCONSIN	2204	23.4	1.1	(21.3, 25.6)
WYOMING	2433	21.1	1.0	(19.1, 23.0)
PUERTO RICO	2812	60.6	1.1	(58.5, 62.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 27.7 RANGE = 17.1-60.6

* NO PHYSICAL ACTIVITY

June 18, 1999

TABLE 8.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PHYSICAL ACTIVITY* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

23

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	786	25.5	1.8	(22.1, 29.0)	*	1466	33.5	1.4	(30.7, 36.2)	*
ALASKA	924	23.5	1.9	(19.7, 27.3)	*	1064	23.5	1.8	(20.0, 27.0)	*
ARIZONA	824	52.3	2.5	(47.5, 57.1)	*	1090	50.5	2.6	(45.5, 55.5)	*
ARKANSAS	1143	33.7	1.5	(30.7, 36.7)	*	1854	37.9	1.3	(35.3, 40.5)	*
CALIFORNIA	1822	23.6	1.1	(21.4, 25.9)	*	2339	27.4	1.1	(25.3, 29.5)	*
COLORADO	811	20.9	1.7	(17.4, 24.3)	*	991	21.7	1.5	(18.8, 24.5)	*
CONNECTICUT	1285	24.1	1.5	(21.1, 27.0)	*	1841	29.8	1.3	(27.2, 32.4)	*
DELAWARE	1037	34.3	2.1	(30.3, 38.4)	*	1545	36.4	1.7	(33.1, 39.6)	*
DISTRICT OF CO	544	31.6	2.3	(27.1, 36.2)	*	908	44.3	1.9	(40.5, 48.1)	*
FLORIDA	1887	29.3	1.2	(27.0, 31.7)	*	2835	32.7	1.0	(30.7, 34.8)	*
GEORGIA	895	26.6	1.9	(22.8, 30.3)	*	1480	32.5	1.4	(29.7, 35.2)	*
HAWAII	933	13.5	1.5	(10.5, 16.4)	*	1220	22.5	1.6	(19.5, 25.6)	*
IDAHO	2083	19.4	1.0	(17.3, 21.4)	*	2847	21.4	0.9	(19.7, 23.2)	*
ILLINOIS	615	25.1	1.9	(21.3, 28.8)	*	825	28.9	1.8	(25.3, 32.5)	*
INDIANA	991	23.1	1.4	(20.2, 25.9)	*	1434	30.8	1.4	(28.1, 33.4)	*
IOWA	1450	26.4	1.3	(23.7, 29.0)	*	2204	26.9	1.1	(24.7, 29.1)	*
KANSAS	1486	38.8	1.4	(36.0, 41.7)	*	2316	37.7	1.2	(35.5, 40.0)	*
KENTUCKY	1363	40.0	1.5	(37.1, 42.8)	*	2354	45.1	1.1	(42.8, 47.3)	*
LOUISIANA	648	28.8	1.9	(25.0, 32.6)	*	1007	35.3	1.7	(32.0, 38.5)	*
MAINE	668	26.6	1.9	(22.9, 30.3)	*	947	28.7	1.7	(25.3, 32.0)	*
MARYLAND	1301	19.3	1.5	(16.4, 22.2)	*	1994	21.2	1.2	(18.8, 23.6)	*
MASSACHUSETTS	2015	23.7	1.2	(21.4, 26.1)	*	2925	26.8	1.1	(24.8, 28.9)	*
MICHIGAN	1037	19.1	1.3	(16.5, 21.7)	*	1497	23.4	1.2	(21.1, 25.8)	*
MINNESOTA	1991	25.8	1.1	(23.7, 28.0)	*	2231	25.1	1.0	(23.1, 27.1)	*
MISSISSIPPI	870	30.8	1.8	(27.3, 34.3)	*	1436	36.5	1.4	(33.6, 39.3)	*
MISSOURI	1434	25.6	1.5	(22.7, 28.6)	*	2286	30.0	1.2	(27.7, 32.3)	*
MONTANA	784	23.2	1.6	(20.0, 26.4)	*	1015	27.1	1.5	(24.1, 30.1)	*
NEBRASKA	1052	26.6	1.6	(23.5, 29.7)	*	1656	25.6	1.2	(23.3, 27.9)	*
NEVADA	897	20.3	2.1	(16.3, 24.3)	*	1082	27.9	2.5	(23.1, 32.7)	*
NEW HAMPSHIRE	616	23.9	2.0	(20.0, 27.7)	*	882	25.8	1.7	(22.5, 29.1)	*
NEW JERSEY	1005	27.6	1.6	(24.4, 30.8)	*	1453	37.2	1.5	(34.3, 40.1)	*
NEW MEXICO	1570	19.5	1.1	(17.3, 21.7)	*	2067	26.4	1.1	(24.2, 28.6)	*
NEW YORK	1052	26.4	1.6	(23.3, 29.5)	*	1472	35.1	1.4	(32.4, 37.9)	*
NORTH CAROLINA	920	24.1	1.9	(20.3, 27.8)	*	1480	31.1	1.5	(28.3, 34.0)	*
NORTH DAKOTA	806	32.9	1.8	(29.4, 36.4)	*	997	33.4	1.6	(30.2, 36.5)	*
OHIO	1184	25.8	1.7	(22.5, 29.2)	*	1801	33.4	1.6	(30.3, 36.5)	*
OKLAHOMA	1004	40.2	1.8	(36.8, 43.7)	*	1512	45.4	1.4	(42.6, 48.2)	*
OREGON	741	16.8	1.6	(13.8, 19.8)	*	1077	20.9	1.4	(18.2, 23.5)	*
PENNSYLVANIA	1426	29.2	1.3	(26.5, 31.8)	*	2171	35.9	1.2	(33.6, 38.2)	*
RHODE ISLAND	1412	26.5	1.3	(24.0, 29.1)	*	2183	32.9	1.1	(30.7, 35.2)	*
SOUTH CAROLINA	1373	32.7	1.5	(29.7, 35.6)	*	1920	34.6	1.3	(32.1, 37.0)	*
SOUTH DAKOTA	924	40.1	1.9	(36.4, 43.8)	*	1187	26.9	1.4	(24.1, 29.7)	*
TENNESSEE	1027	33.3	1.7	(30.0, 36.6)	*	1917	38.0	1.3	(35.5, 40.4)	*
TEXAS	2452	24.3	1.1	(22.2, 26.5)	*	3552	31.2	1.1	(29.2, 33.3)	*
UTAH	1237	15.4	1.3	(12.8, 17.9)	*	1625	18.8	1.3	(16.2, 21.3)	*
VERMONT	1281	25.0	1.4	(22.4, 27.7)	*	1717	26.9	1.3	(24.4, 29.4)	*
VIRGINIA	1485	22.0	1.5	(19.1, 24.9)	*	2019	27.4	1.5	(24.5, 30.3)	*
WASHINGTON	1587	17.6	1.1	(15.6, 19.7)	*	2013	17.6	1.0	(15.7, 19.5)	*
WEST VIRGINIA	942	43.0	1.7	(39.6, 46.3)	*	1472	44.4	1.5	(41.5, 47.3)	*
WISCONSIN	978	23.8	1.7	(20.5, 27.0)	*	1226	23.1	1.5	(20.3, 26.0)	*
WYOMING	1022	20.0	1.6	(16.9, 23.1)	*	1411	22.1	1.2	(19.8, 24.4)	*
PUERTO RICO	1077	56.9	1.7	(53.6, 60.2)	*	1735	63.9	1.3	(61.3, 66.4)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 25.7
 RANGE = 13.5-56.9

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 29.9
 RANGE = 17.6-63.9

June 18, 1999

* NO PHYSICAL ACTIVITY

TABLE 8.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1998
 NO PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

24

PARTICIPANT	1996 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	32.4	29.7	-2.7
ALASKA	25.4	23.5	-1.9
ARIZONA	33.3	51.3	18.1
ARKANSAS	37.4	35.9	-1.5
CALIFORNIA	23.6	25.5	1.9
COLORADO	20.2	21.3	1.1
CONNECTICUT	25.6	27.1	1.5
DELAWARE	36.1	35.4	-0.7
DISTRICT OF CO	30.3	38.5	8.2
FLORIDA	27.1	31.1	4.0
GEORGIA	51.4	29.6	-21.7
HAWAII	21.0	18.0	-3.0
IDAHO	20.6	20.4	-0.2
ILLINOIS	24.9	27.1	2.3
INDIANA	31.0	27.1	-3.9
IOWA	26.9	26.7	-0.3
KANSAS	36.4	38.3	1.9
KENTUCKY	45.6	42.7	-2.9
LOUISIANA	34.9	32.2	-2.7
MAINE	34.0	27.7	-6.3
MARYLAND	33.8	20.3	-13.5
MASSACHUSETTS	23.0	25.4	2.3
MICHIGAN	23.3	21.4	-1.9
MINNESOTA	23.6	25.5	1.8
MISSISSIPPI	39.5	33.8	-5.7
MISSOURI	30.2	27.9	-2.3
MONTANA	21.2	25.2	4.0
NEBRASKA	22.9	26.1	3.2
NEVADA	22.7	24.1	1.3
NEW HAMPSHIRE	25.5	24.9	-0.6
NEW JERSEY	26.3	32.6	6.4
NEW MEXICO	27.7	23.0	-4.7
NEW YORK	30.4	31.0	0.6
NORTH CAROLINA	40.7	27.7	-12.9
NORTH DAKOTA	33.9	33.1	-0.7
OHIO	42.6	29.8	-12.8
OKLAHOMA	38.2	43.0	4.8
OREGON	19.6	18.9	-0.7
PENNSYLVANIA	26.3	32.7	6.4
RHODE ISLAND	26.7	29.9	3.3
SOUTH CAROLINA	29.7	33.7	4.0
SOUTH DAKOTA	34.8	33.3	-1.5
TENNESSEE	40.8	35.8	-5.0
TEXAS	27.9	27.9	0.0
UTAH	17.1	17.1	-0.0
VERMONT	21.5	26.0	4.6
VIRGINIA	29.2	24.8	-4.5
WASHINGTON	19.1	17.6	-1.5
WEST VIRGINIA	42.7	43.7	1.0
WISCONSIN	22.1	23.4	1.3
WYOMING	20.4	21.1	0.7
PUERTO RICO	48.1	60.6	12.5

* NO PHYSICAL ACTIVITY

June 18, 1999

TABLE 9.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND SUSTAINED PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

25

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2252	78.6	1.0	(76.6, 80.6)
ALASKA	1988	75.2	1.4	(72.5, 77.9)
ARIZONA	1914	86.5	1.2	(84.1, 88.9)
ARKANSAS	2997	84.6	0.8	(83.1, 86.1)
CALIFORNIA	4161	75.3	0.8	(73.8, 76.8)
COLORADO	1802	75.9	1.1	(73.7, 78.0)
CONNECTICUT	3126	79.6	0.9	(77.8, 81.4)
DELAWARE	2582	82.8	1.1	(80.6, 85.0)
DISTRICT OF CO	1452	79.7	1.4	(76.9, 82.4)
FLORIDA	4722	79.5	0.7	(78.2, 80.8)
GEORGIA	2375	82.0	1.0	(80.1, 83.8)
HAWAII	2153	69.7	1.4	(67.0, 72.3)
IDAHO	4930	72.9	0.8	(71.4, 74.3)
ILLINOIS	1440	78.1	1.3	(75.6, 80.5)
INDIANA	2425	80.5	0.9	(78.7, 82.2)
IOWA	3654	80.2	0.8	(78.6, 81.8)
KANSAS	3802	85.7	0.7	(84.4, 87.0)
KENTUCKY	3717	87.0	0.7	(85.7, 88.3)
LOUISIANA	1655	83.9	1.1	(81.8, 86.0)
MAINE	1615	75.9	1.2	(73.4, 78.3)
MARYLAND	3295	75.0	1.0	(73.1, 77.0)
MASSACHUSETTS	4940	78.0	0.8	(76.5, 79.5)
MICHIGAN	2534	76.4	1.0	(74.5, 78.2)
MINNESOTA	4222	78.3	0.7	(76.9, 79.6)
MISSISSIPPI	2306	80.8	1.0	(78.9, 82.6)
MISSOURI	3720	80.9	0.8	(79.3, 82.5)
MONTANA	1799	77.5	1.1	(75.3, 79.7)
NEBRASKA	2708	80.9	0.9	(79.1, 82.6)
NEVADA	1979	74.4	1.4	(71.7, 77.2)
NEW HAMPSHIRE	1498	76.3	1.3	(73.9, 78.8)
NEW JERSEY	2458	80.9	0.9	(79.1, 82.7)
NEW MEXICO	3637	74.2	0.8	(72.6, 75.9)
NEW YORK	2524	80.3	0.9	(78.5, 82.0)
NORTH CAROLINA	2400	81.5	1.0	(79.5, 83.5)
NORTH DAKOTA	1803	81.3	1.1	(79.2, 83.4)
OHIO	2985	81.0	1.0	(79.0, 83.0)
OKLAHOMA	2516	84.3	0.9	(82.6, 86.0)
OREGON	1818	73.3	1.2	(71.0, 75.7)
PENNSYLVANIA	3597	80.6	0.8	(79.1, 82.2)
RHODE ISLAND	3595	79.6	0.8	(78.1, 81.0)
SOUTH CAROLINA	3293	82.2	0.8	(80.6, 83.7)
SOUTH DAKOTA	2111	81.8	1.0	(79.9, 83.8)
TENNESSEE	2944	83.5	0.8	(81.9, 85.0)
TEXAS	6004	79.2	0.8	(77.7, 80.7)
UTAH	2862	72.9	1.2	(70.6, 75.1)
VERMONT	2998	74.5	1.0	(72.4, 76.5)
VIRGINIA	3504	78.7	1.1	(76.7, 80.8)
WASHINGTON	3600	74.5	0.9	(72.9, 76.2)
WEST VIRGINIA	2414	86.5	0.8	(85.0, 88.0)
WISCONSIN	2204	79.8	1.0	(77.8, 81.8)
WYOMING	2433	71.8	1.1	(69.7, 73.9)
PUERTO RICO	2812	85.6	0.8	(84.1, 87.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 79.6 RANGE = 69.7-87.0

*5+ times/week, 30+ minutes/session, regardless of intensity

June 18, 1999

TABLE 9.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND SUSTAINED PHYSICAL ACTIVITY* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

26

PARTICIPANT	MALES					FEMALES				
	SAMPLE	PERCENT	STANDARD	95% CONFIDENCE	*	SAMPLE	PERCENT	STANDARD	95% CONFIDENCE	*
	SIZE		ERROR	INTERVAL	*	SIZE		ERROR	INTERVAL	*
ALABAMA	786	74.5	1.8	(71.1, 77.9)	*	1466	82.2	1.1	(80.0, 84.4)	*
ALASKA	924	74.0	2.1	(69.9, 78.1)	*	1064	76.6	1.7	(73.2, 79.9)	*
ARIZONA	824	86.2	1.7	(82.8, 89.6)	*	1090	86.7	1.7	(83.4, 90.1)	*
ARKANSAS	1143	84.7	1.2	(82.4, 87.0)	*	1854	84.4	1.0	(82.5, 86.4)	*
CALIFORNIA	1822	74.5	1.2	(72.2, 76.7)	*	2339	76.2	1.0	(74.2, 78.1)	*
COLORADO	811	75.0	1.7	(71.7, 78.3)	*	991	76.7	1.4	(73.9, 79.5)	*
CONNECTICUT	1285	78.0	1.5	(75.2, 80.9)	*	1841	81.0	1.2	(78.7, 83.2)	*
DELAWARE	1037	82.0	1.9	(78.4, 85.6)	*	1545	83.5	1.3	(81.0, 86.0)	*
DISTRICT OF CO	544	76.8	2.4	(72.1, 81.6)	*	908	82.1	1.5	(79.1, 85.0)	*
FLORIDA	1887	79.1	1.0	(77.1, 81.2)	*	2835	79.8	0.8	(78.1, 81.4)	*
GEORGIA	895	82.0	1.6	(78.9, 85.0)	*	1480	82.0	1.2	(79.7, 84.3)	*
HAWAII	933	66.3	2.1	(62.2, 70.5)	*	1220	73.1	1.7	(69.8, 76.3)	*
IDAHO	2083	72.4	1.2	(70.1, 74.6)	*	2847	73.4	1.0	(71.4, 75.3)	*
ILLINOIS	615	78.2	1.9	(74.5, 81.9)	*	825	78.0	1.7	(74.6, 81.3)	*
INDIANA	991	79.7	1.4	(76.9, 82.5)	*	1434	81.2	1.1	(79.0, 83.4)	*
IOWA	1450	79.2	1.3	(76.7, 81.7)	*	2204	81.1	1.0	(79.1, 83.1)	*
KANSAS	1486	86.3	1.0	(84.3, 88.4)	*	2316	85.1	0.9	(83.4, 86.8)	*
KENTUCKY	1363	85.0	1.1	(82.8, 87.1)	*	2354	88.9	0.7	(87.4, 90.3)	*
LOUISIANA	648	82.2	1.8	(78.6, 85.7)	*	1007	85.4	1.2	(83.0, 87.8)	*
MAINE	668	74.7	2.0	(70.8, 78.5)	*	947	76.9	1.5	(74.0, 79.9)	*
MARYLAND	1301	77.3	1.5	(74.4, 80.1)	*	1994	72.9	1.4	(70.3, 75.6)	*
MASSACHUSETTS	2015	77.5	1.2	(75.1, 79.8)	*	2925	78.5	1.0	(76.6, 80.5)	*
MICHIGAN	1037	74.2	1.5	(71.2, 77.2)	*	1497	78.3	1.2	(76.0, 80.7)	*
MINNESOTA	1991	77.6	1.0	(75.6, 79.6)	*	2231	78.9	0.9	(77.0, 80.7)	*
MISSISSIPPI	870	78.9	1.6	(75.8, 82.0)	*	1436	82.4	1.1	(80.3, 84.6)	*
MISSOURI	1434	80.1	1.3	(77.5, 82.6)	*	2286	81.6	1.0	(79.6, 83.5)	*
MONTANA	784	78.2	1.6	(75.0, 81.4)	*	1015	76.8	1.5	(73.9, 79.8)	*
NEBRASKA	1052	80.1	1.5	(77.1, 83.1)	*	1656	81.5	1.1	(79.5, 83.6)	*
NEVADA	897	74.9	2.1	(70.8, 78.9)	*	1082	74.0	1.9	(70.2, 77.7)	*
NEW HAMPSHIRE	616	76.1	1.9	(72.3, 79.9)	*	882	76.6	1.6	(73.4, 79.8)	*
NEW JERSEY	1005	78.2	1.5	(75.3, 81.2)	*	1453	83.4	1.1	(81.2, 85.6)	*
NEW MEXICO	1570	72.1	1.3	(69.5, 74.7)	*	2067	76.2	1.1	(74.1, 78.2)	*
NEW YORK	1052	77.3	1.5	(74.4, 80.2)	*	1472	82.9	1.1	(80.7, 85.1)	*
NORTH CAROLINA	920	79.7	1.7	(76.4, 82.9)	*	1480	83.1	1.2	(80.8, 85.4)	*
NORTH DAKOTA	806	79.7	1.7	(76.4, 83.0)	*	997	82.8	1.3	(80.2, 85.4)	*
OHIO	1184	77.7	1.7	(74.4, 81.0)	*	1801	83.9	1.2	(81.5, 86.3)	*
OKLAHOMA	1004	83.7	1.4	(81.0, 86.4)	*	1512	84.9	1.1	(82.8, 87.0)	*
OREGON	741	73.3	1.9	(69.6, 76.9)	*	1077	73.4	1.5	(70.4, 76.3)	*
PENNSYLVANIA	1426	78.7	1.3	(76.1, 81.3)	*	2171	82.4	0.9	(80.5, 84.2)	*
RHODE ISLAND	1412	79.5	1.2	(77.1, 81.8)	*	2183	79.7	1.0	(77.8, 81.5)	*
SOUTH CAROLINA	1373	81.3	1.2	(78.9, 83.7)	*	1920	82.9	1.1	(80.8, 85.1)	*
SOUTH DAKOTA	924	85.4	1.3	(82.8, 88.0)	*	1187	78.5	1.5	(75.6, 81.4)	*
TENNESSEE	1027	81.5	1.3	(79.0, 84.0)	*	1917	85.2	0.9	(83.4, 87.0)	*
TEXAS	2452	78.2	1.3	(75.8, 80.7)	*	3552	80.1	0.9	(78.3, 81.9)	*
UTAH	1237	72.9	1.8	(69.5, 76.4)	*	1625	72.8	1.5	(69.8, 75.7)	*
VERMONT	1281	74.1	1.7	(70.8, 77.5)	*	1717	74.8	1.2	(72.4, 77.2)	*
VIRGINIA	1485	78.2	1.5	(75.3, 81.2)	*	2019	79.2	1.5	(76.3, 82.1)	*
WASHINGTON	1587	75.7	1.2	(73.3, 78.1)	*	2013	73.4	1.2	(71.1, 75.7)	*
WEST VIRGINIA	942	85.0	1.3	(82.3, 87.6)	*	1472	87.9	0.9	(86.0, 89.7)	*
WISCONSIN	978	81.4	1.4	(78.6, 84.2)	*	1226	78.3	1.5	(75.4, 81.2)	*
WYOMING	1022	71.7	1.7	(68.4, 74.9)	*	1411	71.9	1.4	(69.2, 74.6)	*
PUERTO RICO	1077	84.2	1.3	(81.7, 86.7)	*	1735	86.9	0.9	(85.1, 88.6)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52
MEDIAN = 78.2
RANGE = 66.3-86.3

NUMBER OF PARTICIPANTS = 52
MEDIAN = 80.5
RANGE = 71.9-88.9

*5+ times/week, 30+ minutes/session, regardless of intensity

June 18, 1999

TABLE 9.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1998
 NO REGULAR AND SUSTAINED PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

27

PARTICIPANT	1996 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	82.4	78.6	-3.8
ALASKA	76.2	75.2	-1.0
ARIZONA	79.0	86.5	7.5
ARKANSAS	81.9	84.6	2.7
CALIFORNIA	78.6	75.3	-3.3
COLORADO	74.9	75.9	0.9
CONNECTICUT	78.6	79.6	1.0
DELAWARE	83.9	82.8	-1.1
DISTRICT OF CO	80.0	79.7	-0.4
FLORIDA	79.0	79.5	0.5
GEORGIA	88.4	82.0	-6.4
HAWAII	71.9	69.7	-2.2
IDAHO	74.9	72.9	-2.0
ILLINOIS	78.9	78.1	-0.8
INDIANA	86.5	80.5	-6.0
IOWA	81.1	80.2	-0.9
KANSAS	82.1	85.7	3.6
KENTUCKY	87.6	87.0	-0.6
LOUISIANA	84.1	83.9	-0.2
MAINE	79.0	75.9	-3.1
MARYLAND	86.7	75.0	-11.7
MASSACHUSETTS	75.8	78.0	2.2
MICHIGAN	77.2	76.4	-0.8
MINNESOTA	75.9	78.3	2.3
MISSISSIPPI	83.0	80.8	-2.3
MISSOURI	82.2	80.9	-1.3
MONTANA	75.8	77.5	1.7
NEBRASKA	79.8	80.9	1.1
NEVADA	75.3	74.4	-0.9
NEW HAMPSHIRE	77.6	76.3	-1.2
NEW JERSEY	77.6	80.9	3.3
NEW MEXICO	76.1	74.2	-1.9
NEW YORK	80.0	80.3	0.2
NORTH CAROLINA	86.4	81.5	-4.9
NORTH DAKOTA	83.2	81.3	-2.0
OHIO	89.8	81.0	-8.8
OKLAHOMA	83.6	84.3	0.7
OREGON	74.3	73.3	-1.0
PENNSYLVANIA	79.6	80.6	1.0
RHODE ISLAND	76.6	79.6	2.9
SOUTH CAROLINA	85.7	82.2	-3.6
SOUTH DAKOTA	81.4	81.8	0.4
TENNESSEE	85.7	83.5	-2.3
TEXAS	78.5	79.2	0.7
UTAH	72.8	72.9	0.0
VERMONT	75.3	74.5	-0.9
VIRGINIA	78.6	78.7	0.2
WASHINGTON	73.7	74.5	0.8
WEST VIRGINIA	87.3	86.5	-0.8
WISCONSIN	78.0	79.8	1.8
WYOMING	71.6	71.8	0.2
PUERTO RICO	79.4	85.6	6.2

*5+ times/week, 30+ minutes/session, regardless of intensity

June 18, 1999

TABLE 10.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

28

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2252	90.0	0.7	(88.6, 91.4)
ALASKA	1988	82.5	1.2	(80.1, 84.9)
ARIZONA	1914	90.2	1.1	(88.0, 92.3)
ARKANSAS	2997	89.7	0.7	(88.4, 91.0)
CALIFORNIA	4161	83.6	0.6	(82.3, 84.8)
COLORADO	1802	85.2	0.9	(83.5, 86.9)
CONNECTICUT	3126	85.9	0.9	(84.1, 87.7)
DELAWARE	2582	87.6	1.0	(85.7, 89.5)
DISTRICT OF CO	1452	83.4	1.2	(81.1, 85.7)
FLORIDA	4722	84.4	0.6	(83.2, 85.6)
GEORGIA	2375	87.4	0.8	(85.7, 89.0)
HAWAII	2153	82.3	1.0	(80.3, 84.3)
IDAHO	4930	84.1	0.6	(82.9, 85.3)
ILLINOIS	1440	85.8	1.0	(83.8, 87.8)
INDIANA	2425	87.1	0.8	(85.7, 88.6)
IOWA	3654	89.4	0.6	(88.2, 90.6)
KANSAS	3802	91.2	0.5	(90.2, 92.3)
KENTUCKY	3717	92.9	0.5	(92.0, 93.9)
LOUISIANA	1655	89.3	0.9	(87.6, 91.0)
MAINE	1615	85.7	1.0	(83.8, 87.6)
MARYLAND	3295	83.6	0.9	(82.0, 85.3)
MASSACHUSETTS	4940	84.4	0.7	(83.0, 85.7)
MICHIGAN	2534	85.9	0.8	(84.3, 87.5)
MINNESOTA	4222	84.5	0.6	(83.4, 85.7)
MISSISSIPPI	2306	89.6	0.7	(88.2, 91.0)
MISSOURI	3720	88.6	0.7	(87.3, 90.0)
MONTANA	1799	86.7	0.9	(85.0, 88.3)
NEBRASKA	2708	89.5	0.7	(88.2, 90.7)
NEVADA	1979	85.7	1.2	(83.3, 88.1)
NEW HAMPSHIRE	1498	87.7	0.9	(85.9, 89.5)
NEW JERSEY	2458	87.6	0.8	(86.1, 89.1)
NEW MEXICO	3637	85.1	0.7	(83.8, 86.5)
NEW YORK	2524	85.2	0.8	(83.6, 86.8)
NORTH CAROLINA	2400	88.4	0.8	(86.8, 89.9)
NORTH DAKOTA	1803	89.0	0.8	(87.4, 90.6)
OHIO	2985	88.4	0.8	(86.8, 90.0)
OKLAHOMA	2516	88.8	0.7	(87.4, 90.2)
OREGON	1818	84.9	0.9	(83.1, 86.7)
PENNSYLVANIA	3597	86.8	0.7	(85.5, 88.1)
RHODE ISLAND	3595	84.5	0.7	(83.1, 85.8)
SOUTH CAROLINA	3293	88.9	0.6	(87.6, 90.1)
SOUTH DAKOTA	2111	90.3	0.8	(88.7, 91.8)
TENNESSEE	2944	90.4	0.6	(89.1, 91.6)
TEXAS	6004	86.2	0.7	(84.8, 87.6)
UTAH	2862	85.0	0.9	(83.3, 86.7)
VERMONT	2998	83.0	1.0	(81.1, 84.9)
VIRGINIA	3504	84.6	0.9	(82.8, 86.5)
WASHINGTON	3600	85.5	0.7	(84.2, 86.8)
WEST VIRGINIA	2414	92.7	0.6	(91.5, 93.8)
WISCONSIN	2204	86.8	0.9	(84.9, 88.6)
WYOMING	2433	84.4	0.9	(82.6, 86.1)
PUERTO RICO	2812	92.1	0.6	(90.9, 93.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 86.7 RANGE = 82.3-92.9

* 3+ TIMES/WEEK 20+ MINUTE/SESSIONS, 50% OF CAPACITY

June 18, 1999

TABLE 10.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

29

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	786	88.8	1.2	(86.4, 91.1)	*	1466	91.1	0.8	(89.5, 92.6)	*
ALASKA	924	83.6	1.8	(80.2, 87.1)	*	1064	81.3	1.7	(78.1, 84.6)	*
ARIZONA	824	91.0	1.4	(88.2, 93.8)	*	1090	89.4	1.7	(86.1, 92.6)	*
ARKANSAS	1143	89.9	1.0	(88.0, 91.9)	*	1854	89.5	0.9	(87.7, 91.2)	*
CALIFORNIA	1822	85.2	0.9	(83.4, 87.0)	*	2339	82.0	0.9	(80.2, 83.7)	*
COLORADO	811	85.5	1.3	(83.0, 88.1)	*	991	84.9	1.2	(82.5, 87.3)	*
CONNECTICUT	1285	86.3	1.5	(83.4, 89.2)	*	1841	85.5	1.0	(83.5, 87.5)	*
DELAWARE	1037	87.0	1.6	(83.8, 90.2)	*	1545	88.2	1.1	(85.9, 90.4)	*
DISTRICT OF CO	544	82.0	2.0	(78.1, 85.8)	*	908	84.7	1.4	(81.9, 87.5)	*
FLORIDA	1887	84.2	0.9	(82.4, 86.1)	*	2835	84.5	0.8	(83.0, 86.0)	*
GEORGIA	895	85.3	1.5	(82.5, 88.2)	*	1480	89.3	0.9	(87.5, 91.0)	*
HAWAII	933	80.2	1.6	(77.0, 83.5)	*	1220	84.4	1.2	(82.0, 86.8)	*
IDAHO	2083	83.7	1.0	(81.8, 85.6)	*	2847	84.5	0.8	(83.0, 86.0)	*
ILLINOIS	615	86.5	1.5	(83.5, 89.5)	*	825	85.3	1.4	(82.5, 88.0)	*
INDIANA	991	87.5	1.1	(85.3, 89.7)	*	1434	86.8	1.0	(84.8, 88.7)	*
IOWA	1450	89.6	0.9	(87.8, 91.3)	*	2204	89.3	0.8	(87.7, 91.0)	*
KANSAS	1486	91.4	0.8	(89.8, 92.9)	*	2316	91.1	0.7	(89.8, 92.5)	*
KENTUCKY	1363	92.2	0.8	(90.6, 93.8)	*	2354	93.6	0.5	(92.5, 94.6)	*
LOUISIANA	648	88.4	1.5	(85.6, 91.3)	*	1007	90.1	1.1	(88.0, 92.2)	*
MAINE	668	85.3	1.5	(82.3, 88.2)	*	947	86.0	1.3	(83.6, 88.5)	*
MARYLAND	1301	84.8	1.3	(82.3, 87.3)	*	1994	82.6	1.2	(80.3, 84.9)	*
MASSACHUSETTS	2015	86.0	1.0	(84.1, 87.8)	*	2925	83.0	1.0	(81.0, 84.9)	*
MICHIGAN	1037	84.5	1.3	(82.0, 87.0)	*	1497	87.2	1.0	(85.3, 89.1)	*
MINNESOTA	1991	85.1	0.9	(83.5, 86.8)	*	2231	83.9	0.8	(82.3, 85.5)	*
MISSISSIPPI	870	88.3	1.2	(85.9, 90.6)	*	1436	90.8	0.8	(89.2, 92.4)	*
MISSOURI	1434	89.4	1.1	(87.3, 91.5)	*	2286	87.9	0.9	(86.2, 89.7)	*
MONTANA	784	87.4	1.2	(85.0, 89.8)	*	1015	85.9	1.2	(83.6, 88.2)	*
NEBRASKA	1052	89.7	1.0	(87.8, 91.6)	*	1656	89.3	0.8	(87.6, 90.9)	*
NEVADA	897	85.2	1.9	(81.4, 89.0)	*	1082	86.2	1.4	(83.4, 88.9)	*
NEW HAMPSHIRE	616	87.8	1.4	(84.9, 90.6)	*	882	87.6	1.2	(85.2, 90.0)	*
NEW JERSEY	1005	88.7	1.2	(86.4, 91.0)	*	1453	86.6	1.0	(84.7, 88.5)	*
NEW MEXICO	1570	85.1	1.0	(83.1, 87.2)	*	2067	85.2	0.9	(83.4, 86.9)	*
NEW YORK	1052	85.4	1.3	(82.9, 87.9)	*	1472	85.0	1.1	(82.9, 87.0)	*
NORTH CAROLINA	920	89.1	1.3	(86.6, 91.5)	*	1480	87.7	1.0	(85.7, 89.7)	*
NORTH DAKOTA	806	89.5	1.2	(87.2, 91.8)	*	997	88.5	1.2	(86.2, 90.8)	*
OHIO	1184	88.2	1.3	(85.7, 90.7)	*	1801	88.6	1.0	(86.5, 90.6)	*
OKLAHOMA	1004	89.5	1.1	(87.4, 91.6)	*	1512	88.1	0.9	(86.3, 89.9)	*
OREGON	741	86.0	1.4	(83.3, 88.8)	*	1077	83.8	1.2	(81.4, 86.2)	*
PENNSYLVANIA	1426	86.5	1.1	(84.4, 88.7)	*	2171	87.0	0.8	(85.5, 88.5)	*
RHODE ISLAND	1412	83.0	1.1	(80.7, 85.2)	*	2183	85.8	0.8	(84.2, 87.5)	*
SOUTH CAROLINA	1373	88.6	1.0	(86.6, 90.6)	*	1920	89.1	0.8	(87.5, 90.7)	*
SOUTH DAKOTA	924	92.9	0.9	(91.2, 94.6)	*	1187	87.8	1.3	(85.3, 90.2)	*
TENNESSEE	1027	90.0	1.0	(87.9, 92.0)	*	1917	90.7	0.7	(89.3, 92.2)	*
TEXAS	2452	86.2	1.1	(84.0, 88.4)	*	3552	86.2	0.9	(84.4, 88.0)	*
UTAH	1237	86.1	1.4	(83.4, 88.8)	*	1625	84.0	1.1	(81.7, 86.2)	*
VERMONT	1281	82.0	1.7	(78.8, 85.3)	*	1717	84.0	1.0	(82.0, 85.9)	*
VIRGINIA	1485	85.6	1.3	(83.0, 88.1)	*	2019	83.8	1.4	(81.1, 86.4)	*
WASHINGTON	1587	85.5	1.0	(83.6, 87.4)	*	2013	85.5	0.9	(83.8, 87.3)	*
WEST VIRGINIA	942	92.5	1.0	(90.6, 94.4)	*	1472	92.8	0.8	(91.3, 94.3)	*
WISCONSIN	978	88.4	1.2	(86.0, 90.8)	*	1226	85.3	1.4	(82.5, 88.0)	*
WYOMING	1022	85.4	1.4	(82.7, 88.1)	*	1411	83.4	1.1	(81.2, 85.5)	*
PUERTO RICO	1077	90.7	1.0	(88.8, 92.7)	*	1735	93.3	0.7	(92.0, 94.7)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 86.5
 RANGE = 80.2-92.9

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 86.4
 RANGE = 81.3-93.6

June 18, 1999

* 3+ TIMES/WEEK 20+ MINUTE/SESSIONS, 50% OF CAPACITY

TABLE 10.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1998
 NO REGULAR AND VIGOROUS PHYSICAL ACTIVITY*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

30

PARTICIPANT	1996 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	89.9	90.0	0.1
ALASKA	85.9	82.5	-3.4
ARIZONA	84.2	90.2	6.0
ARKANSAS	88.5	89.7	1.2
CALIFORNIA	92.1	83.6	-8.5
COLORADO	83.4	85.2	1.8
CONNECTICUT	85.7	85.9	0.2
DELAWARE	87.3	87.6	0.4
DISTRICT OF CO	85.3	83.4	-1.8
FLORIDA	82.4	84.4	2.0
GEORGIA	93.5	87.4	-6.2
HAWAII	81.6	82.3	0.7
IDAHO	85.8	84.1	-1.7
ILLINOIS	83.2	85.8	2.6
INDIANA	88.6	87.1	-1.4
IOWA	86.5	89.4	2.9
KANSAS	84.1	91.2	7.1
KENTUCKY	89.3	92.9	3.6
LOUISIANA	88.7	89.3	0.6
MAINE	85.9	85.7	-0.3
MARYLAND	87.6	83.6	-4.0
MASSACHUSETTS	82.1	84.4	2.3
MICHIGAN	83.3	85.9	2.6
MINNESOTA	82.9	84.5	1.6
MISSISSIPPI	88.8	89.6	0.8
MISSOURI	88.1	88.6	0.5
MONTANA	83.5	86.7	3.2
NEBRASKA	87.5	89.5	2.0
NEVADA	84.7	85.7	1.0
NEW HAMPSHIRE	84.4	87.7	3.3
NEW JERSEY	85.9	87.6	1.7
NEW MEXICO	85.1	85.1	0.1
NEW YORK	86.1	85.2	-0.9
NORTH CAROLINA	90.9	88.4	-2.6
NORTH DAKOTA	88.9	89.0	0.1
OHIO	93.6	88.4	-5.2
OKLAHOMA	87.9	88.8	0.9
OREGON	82.7	84.9	2.2
PENNSYLVANIA	87.4	86.8	-0.6
RHODE ISLAND	84.7	84.5	-0.3
SOUTH CAROLINA	87.3	88.9	1.6
SOUTH DAKOTA	86.5	90.3	3.7
TENNESSEE	88.8	90.4	1.5
TEXAS	85.6	86.2	0.6
UTAH	83.9	85.0	1.1
VERMONT	85.6	83.0	-2.5
VIRGINIA	85.0	84.6	-0.3
WASHINGTON	83.9	85.5	1.6
WEST VIRGINIA	89.6	92.7	3.1
WISCONSIN	85.3	86.8	1.5
WYOMING	84.4	84.4	0.0
PUERTO RICO	93.1	92.1	-1.0

* 3+ TIMES/WEEK 20+ MINUTE/SESSIONS, 50% OF CAPACITY

June 18, 1999

TABLE 11.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FRUITS AND VEGETABLES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

31

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2251	76.1	1.1	(74.0, 78.1)
ALASKA	1989	76.9	1.4	(74.2, 79.5)
ARIZONA	1915	90.9	1.0	(88.9, 92.9)
ARKANSAS	2999	72.1	1.0	(70.2, 74.0)
CALIFORNIA	4161	72.5	0.8	(70.9, 74.0)
COLORADO	1801	74.0	1.2	(71.7, 76.3)
CONNECTICUT	3123	72.1	1.0	(70.1, 74.1)
DELAWARE	2582	73.3	1.2	(71.0, 75.7)
DISTRICT OF CO	1451	82.3	1.3	(79.8, 84.8)
FLORIDA	4723	75.1	0.7	(73.7, 76.5)
GEORGIA	2374	79.3	1.0	(77.5, 81.2)
HAWAII	2152	72.5	1.2	(70.1, 74.9)
IDAHO	4931	76.2	0.7	(74.7, 77.6)
ILLINOIS	1440	77.3	1.3	(74.8, 79.7)
INDIANA	2425	76.5	1.0	(74.7, 78.4)
IOWA	3654	81.4	0.8	(79.9, 82.9)
KANSAS	3799	76.5	0.8	(74.9, 78.0)
KENTUCKY	3717	84.4	0.7	(83.0, 85.7)
LOUISIANA	1654	82.7	1.0	(80.6, 84.7)
MAINE	1615	73.6	1.3	(71.1, 76.0)
MARYLAND	3297	69.9	1.1	(67.9, 72.0)
MASSACHUSETTS	4939	69.0	0.9	(67.3, 70.6)
MICHIGAN	2535	72.6	1.0	(70.6, 74.5)
MINNESOTA	4224	68.1	0.8	(66.5, 69.6)
MISSISSIPPI	2305	84.4	0.9	(82.7, 86.1)
MISSOURI	3720	80.0	0.8	(78.4, 81.6)
MONTANA	1794	76.2	1.1	(74.0, 78.4)
NEBRASKA	2707	82.3	0.9	(80.6, 84.0)
NEVADA	1980	77.9	1.4	(75.1, 80.7)
NEW HAMPSHIRE	1496	72.1	1.3	(69.6, 74.7)
NEW JERSEY	2461	73.9	1.1	(71.8, 76.0)
NEW MEXICO	3638	79.3	0.7	(77.9, 80.8)
NEW YORK	2519	74.5	1.0	(72.5, 76.5)
NORTH CAROLINA	2400	78.6	1.0	(76.6, 80.6)
NORTH DAKOTA	1802	77.2	1.1	(75.1, 79.4)
OHIO	2979	84.0	0.9	(82.1, 85.8)
OKLAHOMA	2515	79.7	0.9	(77.9, 81.5)
OREGON	1818	75.1	1.2	(72.8, 77.4)
PENNSYLVANIA	3595	75.2	0.8	(73.5, 76.8)
RHODE ISLAND	3600	75.4	0.8	(73.9, 77.0)
SOUTH CAROLINA	3293	78.2	0.8	(76.6, 79.8)
SOUTH DAKOTA	2110	80.0	1.0	(78.0, 82.0)
TENNESSEE	2940	70.3	1.0	(68.3, 72.2)
TEXAS	6005	77.5	0.8	(76.1, 79.0)
UTAH	2864	73.8	1.1	(71.7, 75.9)
VERMONT	2996	70.8	1.0	(68.9, 72.7)
VIRGINIA	3504	73.1	1.1	(70.8, 75.3)
WASHINGTON	3601	74.2	0.9	(72.5, 75.9)
WEST VIRGINIA	2414	81.4	0.9	(79.6, 83.1)
WISCONSIN	2205	74.3	1.2	(72.0, 76.6)
WYOMING	2434	78.8	0.9	(77.0, 80.5)
PUERTO RICO	2752	91.7	0.6	(90.5, 92.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 76.2 RANGE = 68.1-91.7

* NOT EATING 5 SERVINGS A DAY

June 18, 1999

TABLE 11.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR FRUITS AND VEGETABLES* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

32

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	786	76.5	1.8	(73.0, 79.9)	*	1465	75.7	1.2	(73.3, 78.1)	*
ALASKA	924	79.0	2.1	(74.9, 83.0)	*	1065	74.5	1.8	(71.0, 78.0)	*
ARIZONA	825	92.3	1.4	(89.6, 95.0)	*	1090	89.6	1.6	(86.6, 92.7)	*
ARKANSAS	1145	75.6	1.5	(72.7, 78.5)	*	1854	68.9	1.3	(66.4, 71.5)	*
CALIFORNIA	1822	77.7	1.1	(75.5, 79.9)	*	2339	67.4	1.1	(65.2, 69.5)	*
COLORADO	810	78.3	1.7	(74.9, 81.6)	*	991	69.9	1.6	(66.7, 73.1)	*
CONNECTICUT	1285	77.4	1.5	(74.5, 80.3)	*	1838	67.3	1.4	(64.6, 70.0)	*
DELAWARE	1037	78.3	1.8	(74.7, 81.8)	*	1545	68.9	1.6	(65.7, 72.0)	*
DISTRICT OF CO	544	82.5	2.2	(78.2, 86.8)	*	907	82.1	1.5	(79.2, 85.0)	*
FLORIDA	1887	79.7	1.1	(77.6, 81.8)	*	2836	71.0	0.9	(69.1, 72.8)	*
GEORGIA	895	82.3	1.4	(79.5, 85.1)	*	1479	76.6	1.2	(74.2, 79.0)	*
HAWAII	932	78.2	1.7	(74.8, 81.6)	*	1220	66.7	1.7	(63.3, 70.0)	*
IDAHO	2083	81.7	1.1	(79.6, 83.7)	*	2848	70.9	1.0	(69.0, 72.9)	*
ILLINOIS	615	82.9	1.7	(79.6, 86.1)	*	825	72.4	1.8	(68.8, 76.0)	*
INDIANA	991	79.7	1.4	(76.9, 82.5)	*	1434	73.7	1.3	(71.2, 76.2)	*
IOWA	1450	84.7	1.1	(82.5, 86.9)	*	2204	78.3	1.0	(76.3, 80.4)	*
KANSAS	1484	80.1	1.2	(77.8, 82.4)	*	2315	73.1	1.1	(71.0, 75.1)	*
KENTUCKY	1362	87.9	1.0	(86.0, 89.8)	*	2355	81.1	0.9	(79.4, 82.9)	*
LOUISIANA	649	84.2	1.5	(81.2, 87.1)	*	1005	81.3	1.4	(78.7, 83.9)	*
MAINE	668	78.4	1.8	(74.9, 82.0)	*	947	69.1	1.7	(65.7, 72.5)	*
MARYLAND	1301	74.5	1.6	(71.4, 77.7)	*	1996	65.8	1.4	(63.0, 68.5)	*
MASSACHUSETTS	2015	74.8	1.2	(72.4, 77.2)	*	2924	63.7	1.2	(61.4, 66.1)	*
MICHIGAN	1037	77.9	1.5	(75.0, 80.8)	*	1498	67.7	1.3	(65.1, 70.3)	*
MINNESOTA	1991	70.8	1.1	(68.6, 73.0)	*	2233	65.5	1.1	(63.4, 67.7)	*
MISSISSIPPI	869	86.9	1.3	(84.3, 89.4)	*	1436	82.2	1.1	(80.0, 84.4)	*
MISSOURI	1434	85.1	1.2	(82.8, 87.3)	*	2286	75.5	1.1	(73.3, 77.7)	*
MONTANA	780	81.6	1.5	(78.7, 84.5)	*	1014	71.1	1.6	(68.0, 74.2)	*
NEBRASKA	1053	87.1	1.4	(84.4, 89.7)	*	1654	78.0	1.1	(75.8, 80.1)	*
NEVADA	898	80.8	2.0	(76.9, 84.8)	*	1082	74.9	2.0	(71.1, 78.8)	*
NEW HAMPSHIRE	615	78.6	1.9	(75.0, 82.2)	*	881	66.1	1.9	(62.4, 69.8)	*
NEW JERSEY	1005	77.9	1.6	(74.8, 81.0)	*	1456	70.3	1.4	(67.6, 73.0)	*
NEW MEXICO	1571	83.0	1.0	(80.9, 85.0)	*	2067	75.9	1.0	(73.9, 77.9)	*
NEW YORK	1050	78.7	1.6	(75.6, 81.7)	*	1469	70.8	1.3	(68.2, 73.4)	*
NORTH CAROLINA	921	83.6	1.4	(80.7, 86.4)	*	1479	74.0	1.4	(71.3, 76.8)	*
NORTH DAKOTA	805	84.1	1.5	(81.3, 87.0)	*	997	70.6	1.6	(67.5, 73.6)	*
OHIO	1182	88.5	1.3	(86.0, 91.0)	*	1797	79.9	1.3	(77.3, 82.5)	*
OKLAHOMA	1004	82.7	1.4	(80.0, 85.5)	*	1511	76.9	1.2	(74.5, 79.3)	*
OREGON	741	80.5	1.7	(77.2, 83.9)	*	1077	70.0	1.7	(66.8, 73.3)	*
PENNSYLVANIA	1425	77.7	1.3	(75.1, 80.2)	*	2170	72.9	1.1	(70.8, 75.0)	*
RHODE ISLAND	1413	79.4	1.2	(77.0, 81.8)	*	2187	71.9	1.1	(69.8, 74.0)	*
SOUTH CAROLINA	1374	81.8	1.2	(79.5, 84.1)	*	1919	75.0	1.1	(72.8, 77.2)	*
SOUTH DAKOTA	925	86.7	1.2	(84.4, 88.9)	*	1185	73.7	1.6	(70.6, 76.8)	*
TENNESSEE	1025	73.6	1.5	(70.6, 76.5)	*	1915	67.3	1.3	(64.9, 69.8)	*
TEXAS	2453	81.1	1.2	(78.8, 83.3)	*	3552	74.2	1.0	(72.3, 76.1)	*
UTAH	1238	78.1	1.6	(75.0, 81.3)	*	1626	69.6	1.4	(66.9, 72.3)	*
VERMONT	1279	76.8	1.4	(74.0, 79.6)	*	1717	65.2	1.3	(62.7, 67.8)	*
VIRGINIA	1486	77.6	1.5	(74.6, 80.6)	*	2018	68.8	1.6	(65.6, 72.0)	*
WASHINGTON	1588	80.6	1.2	(78.3, 82.9)	*	2013	68.1	1.2	(65.7, 70.5)	*
WEST VIRGINIA	942	85.0	1.3	(82.4, 87.6)	*	1472	78.1	1.2	(75.8, 80.4)	*
WISCONSIN	978	82.8	1.5	(79.9, 85.7)	*	1227	66.4	1.7	(63.0, 69.8)	*
WYOMING	1023	84.2	1.2	(81.7, 86.6)	*	1411	73.5	1.3	(71.0, 76.0)	*
PUERTO RICO	1053	92.7	0.9	(90.9, 94.5)	*	1699	90.8	0.8	(89.2, 92.3)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
MEDIAN = 80.5
RANGE = 70.8-92.7

NUMBER OF PARTICIPANTS = 52
MEDIAN = 72.2
RANGE = 63.7-90.8

June 18, 1999

* NOT EATING 5 SERVINGS A DAY

TABLE 11.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1998
 FRUITS AND VEGETABLES*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

33

PARTICIPANT	1996 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	78.6	76.1	-2.5
ALASKA	73.8	76.9	3.0
ARIZONA	75.5	90.9	15.4
ARKANSAS	65.7	72.1	6.4
CALIFORNIA	68.0	72.5	4.4
COLORADO	73.8	74.0	0.2
CONNECTICUT	72.8	72.1	-0.7
DELAWARE	83.9	73.3	-10.6
DISTRICT OF CO	67.6	82.3	14.7
FLORIDA	78.1	75.1	-2.9
GEORGIA	77.3	79.3	2.0
HAWAII	74.9	72.5	-2.4
IDAHO	76.8	76.2	-0.7
ILLINOIS	78.9	77.3	-1.6
INDIANA	79.2	76.5	-2.6
IOWA	84.9	81.4	-3.5
KANSAS	71.7	76.5	4.8
KENTUCKY	82.6	84.4	1.7
LOUISIANA	82.0	82.7	0.6
MAINE	73.8	73.6	-0.3
MARYLAND	75.3	69.9	-5.4
MASSACHUSETTS	74.2	69.0	-5.2
MICHIGAN	76.6	72.6	-4.0
MINNESOTA	69.8	68.1	-1.7
MISSISSIPPI	81.8	84.4	2.6
MISSOURI	77.4	80.0	2.7
MONTANA	76.3	76.2	-0.1
NEBRASKA	77.5	82.3	4.8
NEVADA	78.1	77.9	-0.2
NEW HAMPSHIRE	73.7	72.1	-1.5
NEW JERSEY	73.1	73.9	0.8
NEW MEXICO	78.9	79.3	0.4
NEW YORK	74.6	74.5	-0.1
NORTH CAROLINA	83.5	78.6	-4.9
NORTH DAKOTA	82.2	77.2	-4.9
OHIO	86.1	84.0	-2.1
OKLAHOMA	82.8	79.7	-3.1
OREGON	74.9	75.1	0.3
PENNSYLVANIA	76.3	75.2	-1.2
RHODE ISLAND	76.3	75.4	-0.8
SOUTH CAROLINA	75.8	78.2	2.4
SOUTH DAKOTA	76.1	80.0	3.9
TENNESSEE	74.2	70.3	-3.9
TEXAS	71.8	77.5	5.7
UTAH	78.5	73.8	-4.8
VERMONT	71.2	70.8	-0.4
VIRGINIA	74.3	73.1	-1.3
WASHINGTON	77.1	74.2	-2.9
WEST VIRGINIA	79.9	81.4	1.4
WISCONSIN	74.8	74.3	-0.5
WYOMING	77.9	78.8	0.9
PUERTO RICO	79.7	91.7	12.0

* NOT EATING 5 SERVINGS A DAY

June 18, 1999

TABLE 12.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LOSEWT: NOT EATING FEWER CALORIES AND EXERCISING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

34

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	670	52.1	2.3	(47.5, 56.6)
ALASKA	559	43.4	2.9	(37.7, 49.2)
ARIZONA	238	58.7	5.2	(48.5, 69.0)
ARKANSAS	681	61.5	2.2	(57.2, 65.8)
CALIFORNIA	1037	48.1	1.8	(44.6, 51.6)
COLORADO	394	50.3	2.9	(44.7, 55.9)
CONNECTICUT	717	46.1	2.4	(41.3, 50.8)
DELAWARE	620	55.3	2.8	(49.7, 60.8)
DISTRICT OF CO	369	62.9	3.4	(56.2, 69.6)
FLORIDA	1173	59.6	1.7	(56.3, 62.8)
GEORGIA	694	51.8	2.2	(47.5, 56.0)
HAWAII	478	44.6	3.1	(38.6, 50.6)
IDAHO	1247	46.2	1.7	(42.9, 49.5)
ILLINOIS	385	54.0	2.8	(48.5, 59.6)
INDIANA	702	48.9	2.1	(44.7, 53.0)
IOWA	1047	51.8	1.8	(48.3, 55.4)
KANSAS	838	53.1	1.9	(49.4, 56.9)
KENTUCKY	851	61.0	1.9	(57.3, 64.7)
LOUISIANA	444	55.5	2.6	(50.5, 60.5)
MAINE	399	53.9	2.9	(48.2, 59.5)
MARYLAND	912	48.0	2.3	(43.6, 52.4)
MASSACHUSETTS	1081	46.3	2.0	(42.4, 50.3)
MICHIGAN	740	52.1	2.0	(48.1, 56.0)
MINNESOTA	859	48.8	1.8	(45.2, 52.3)
MISSISSIPPI	693	55.3	2.2	(51.1, 59.6)
MISSOURI	1098	52.5	1.9	(48.8, 56.2)
MONTANA	395	52.6	2.7	(47.3, 57.9)
NEBRASKA	700	49.6	2.1	(45.5, 53.7)
NEVADA	462	52.3	3.2	(46.0, 58.7)
NEW HAMPSHIRE	354	54.9	3.0	(49.0, 60.8)
NEW JERSEY	560	53.4	2.5	(48.5, 58.3)
NEW MEXICO	819	47.4	2.0	(43.5, 51.2)
NEW YORK	570	51.6	2.4	(46.9, 56.2)
NORTH CAROLINA	622	60.4	2.4	(55.7, 65.2)
NORTH DAKOTA	408	55.3	2.7	(50.0, 60.6)
OHIO	796	52.5	2.4	(47.8, 57.3)
OKLAHOMA	577	63.1	2.3	(58.5, 67.6)
OREGON	500	45.1	2.5	(40.2, 50.0)
PENNSYLVANIA	930	54.6	1.9	(51.0, 58.2)
RHODE ISLAND	865	53.4	1.9	(49.6, 57.1)
SOUTH CAROLINA	866	56.4	2.0	(52.5, 60.3)
SOUTH DAKOTA	482	50.6	2.6	(45.5, 55.6)
TENNESSEE	688	51.0	2.2	(46.7, 55.2)
TEXAS	1726	53.3	1.6	(50.2, 56.3)
UTAH	705	47.8	2.6	(42.7, 52.8)
VERMONT	629	52.6	2.3	(48.1, 57.1)
VIRGINIA	1001	47.2	2.2	(42.8, 51.5)
WASHINGTON	856	47.2	2.1	(43.0, 51.3)
WEST VIRGINIA	588	60.2	2.3	(55.6, 64.8)
WISCONSIN	657	45.5	2.4	(40.8, 50.1)
WYOMING	589	47.3	2.3	(42.9, 51.7)
PUERTO RICO	681	70.3	1.9	(66.5, 74.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 52.4 RANGE = 43.4-70.3

* NOT EATING FEWER CALORIES AND EXERCISING TO LOSE WEIGHT

June 18, 1999

TABLE 12.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR LOSEWT: NOT EATING FEWER CALORIES AND EXERCISING* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

35

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	242	53.0	3.7	(45.8, 60.3)	*	428	51.1	2.7	(45.7, 56.5)	*
ALASKA	266	41.8	4.3	(33.4, 50.1)	*	293	45.4	3.9	(37.8, 53.0)	*
ARIZONA	100	63.1	6.9	(49.7, 76.6)	*	138	54.0	7.4	(39.5, 68.5)	*
ARKANSAS	295	59.2	3.3	(52.7, 65.7)	*	386	64.0	2.7	(58.7, 69.4)	*
CALIFORNIA	450	48.1	2.6	(43.0, 53.2)	*	587	48.1	2.4	(43.4, 52.8)	*
COLORADO	175	53.1	4.5	(44.3, 61.9)	*	219	47.6	3.7	(40.3, 54.9)	*
CONNECTICUT	329	44.2	3.5	(37.3, 51.0)	*	388	48.3	3.2	(41.9, 54.6)	*
DELAWARE	251	61.8	4.2	(53.6, 70.1)	*	369	49.0	3.5	(42.2, 55.9)	*
DISTRICT OF CO	112	60.6	6.6	(47.7, 73.6)	*	257	64.4	3.5	(57.5, 71.3)	*
FLORIDA	473	58.6	2.6	(53.6, 63.6)	*	700	60.5	2.1	(56.4, 64.5)	*
GEORGIA	251	51.4	3.5	(44.5, 58.2)	*	443	52.1	2.7	(46.8, 57.4)	*
HAWAII	222	43.9	4.3	(35.4, 52.4)	*	256	45.5	4.2	(37.3, 53.7)	*
IDAHO	534	47.9	2.6	(42.8, 53.0)	*	713	44.4	2.1	(40.2, 48.6)	*
ILLINOIS	164	52.9	4.3	(44.5, 61.3)	*	221	55.0	3.8	(47.6, 62.5)	*
INDIANA	294	50.7	3.2	(44.4, 57.0)	*	408	47.0	2.7	(41.6, 52.4)	*
IOWA	419	53.9	2.9	(48.3, 59.5)	*	628	49.9	2.4	(45.3, 54.6)	*
KANSAS	358	55.2	2.9	(49.7, 60.8)	*	480	51.0	2.5	(46.1, 55.8)	*
KENTUCKY	313	62.0	2.9	(56.3, 67.8)	*	538	60.1	2.4	(55.5, 64.8)	*
LOUISIANA	182	61.1	4.0	(53.3, 68.8)	*	262	50.4	3.5	(43.6, 57.1)	*
MAINE	164	56.1	4.3	(47.7, 64.4)	*	235	52.0	3.9	(44.3, 59.6)	*
MARYLAND	378	50.4	3.5	(43.6, 57.2)	*	534	45.6	2.9	(39.9, 51.4)	*
MASSACHUSETTS	514	48.4	2.9	(42.8, 54.0)	*	567	43.6	2.7	(38.3, 48.9)	*
MICHIGAN	288	52.5	3.2	(46.2, 58.7)	*	452	51.7	2.6	(46.6, 56.8)	*
MINNESOTA	392	47.8	2.6	(42.6, 52.9)	*	467	49.7	2.5	(44.7, 54.6)	*
MISSISSIPPI	258	54.3	3.4	(47.5, 61.0)	*	435	56.2	2.7	(51.0, 61.5)	*
MISSOURI	424	55.3	3.0	(49.3, 61.2)	*	674	50.2	2.5	(45.4, 55.0)	*
MONTANA	166	52.2	4.1	(44.1, 60.3)	*	229	53.1	3.6	(46.1, 60.0)	*
NEBRASKA	260	54.8	3.3	(48.2, 61.3)	*	440	45.0	2.6	(39.9, 50.1)	*
NEVADA	233	49.9	4.3	(41.4, 58.3)	*	229	55.1	4.8	(45.6, 64.5)	*
NEW HAMPSHIRE	160	60.1	4.1	(52.0, 68.2)	*	194	49.2	4.0	(41.3, 57.0)	*
NEW JERSEY	259	50.9	3.6	(43.9, 58.0)	*	301	56.0	3.4	(49.3, 62.7)	*
NEW MEXICO	339	46.4	3.0	(40.6, 52.2)	*	480	48.1	2.6	(43.0, 53.2)	*
NEW YORK	231	52.0	3.6	(44.8, 59.1)	*	339	51.3	3.1	(45.1, 57.4)	*
NORTH CAROLINA	229	64.7	3.9	(57.0, 72.3)	*	393	57.0	3.0	(51.1, 62.9)	*
NORTH DAKOTA	186	62.0	3.8	(54.5, 69.5)	*	222	48.6	3.8	(41.2, 56.0)	*
OHIO	328	53.2	3.7	(46.1, 60.4)	*	468	51.8	3.2	(45.6, 58.1)	*
OKLAHOMA	228	61.3	3.7	(54.1, 68.5)	*	349	64.9	2.8	(59.3, 70.4)	*
OREGON	205	42.1	3.8	(34.6, 49.6)	*	295	48.0	3.2	(41.9, 54.2)	*
PENNSYLVANIA	381	55.9	2.8	(50.3, 61.4)	*	549	53.4	2.4	(48.7, 58.1)	*
RHODE ISLAND	380	51.9	2.8	(46.4, 57.5)	*	485	54.9	2.5	(50.0, 59.8)	*
SOUTH CAROLINA	386	59.5	2.9	(53.8, 65.2)	*	480	53.4	2.7	(48.1, 58.6)	*
SOUTH DAKOTA	201	54.4	3.8	(47.0, 61.9)	*	281	47.3	3.4	(40.5, 54.0)	*
TENNESSEE	219	48.7	3.8	(41.3, 56.1)	*	469	52.8	2.5	(47.8, 57.8)	*
TEXAS	691	53.4	2.3	(48.8, 58.0)	*	1035	53.1	2.1	(49.1, 57.2)	*
UTAH	328	51.2	3.8	(43.8, 58.5)	*	377	44.0	3.5	(37.2, 50.8)	*
VERMONT	277	52.7	3.6	(45.7, 59.6)	*	352	52.6	2.9	(46.8, 58.3)	*
VIRGINIA	430	44.8	3.3	(38.4, 51.2)	*	571	49.3	3.1	(43.3, 55.3)	*
WASHINGTON	410	46.2	3.1	(40.2, 52.1)	*	446	48.2	2.9	(42.6, 53.8)	*
WEST VIRGINIA	228	62.6	3.6	(55.6, 69.6)	*	360	58.1	3.0	(52.2, 63.9)	*
WISCONSIN	292	46.6	3.4	(39.9, 53.2)	*	365	44.3	3.4	(37.8, 50.9)	*
WYOMING	239	50.4	3.5	(43.4, 57.3)	*	350	44.3	2.8	(38.8, 49.9)	*
PUERTO RICO	249	69.0	3.2	(62.8, 75.2)	*	432	71.4	2.4	(66.8, 76.1)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
MEDIAN = 53.0
RANGE = 41.8-69.0

NUMBER OF PARTICIPANTS = 52
MEDIAN = 51.0
RANGE = 43.6-71.4

June 18, 1999

* NOT EATING FEWER CALORIES AND EXERCISING TO LOSE WEIGHT

TABLE 12.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1996 TO 1998
 LOSEWT: NOT EATING FEWER CALORIES AND EXERCISING*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

36

PARTICIPANT	1996 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	56.1	52.1	-4.1
ALASKA	43.9	43.4	-0.5
ARIZONA	52.1	58.7	6.6
ARKANSAS	57.9	61.5	3.7
CALIFORNIA	46.1	48.1	2.0
COLORADO	44.5	50.3	5.8
CONNECTICUT	46.7	46.1	-0.6
DELAWARE	58.9	55.3	-3.6
DISTRICT OF CO	54.7	62.9	8.2
FLORIDA	51.4	59.6	8.2
GEORGIA	61.3	51.8	-9.6
HAWAII	46.3	44.6	-1.8
IDAHO	47.8	46.2	-1.6
ILLINOIS	51.2	54.0	2.9
INDIANA	53.1	48.9	-4.3
IOWA	53.4	51.8	-1.5
KANSAS	50.1	53.1	3.1
KENTUCKY	58.8	61.0	2.2
LOUISIANA	76.0	55.5	-20.5
MAINE	53.8	53.9	0.0
MARYLAND	49.7	48.0	-1.7
MASSACHUSETTS	46.4	46.3	-0.1
MICHIGAN	51.2	52.1	0.9
MINNESOTA	45.1	48.8	3.6
MISSISSIPPI	63.9	55.3	-8.5
MISSOURI	48.4	52.5	4.1
MONTANA	45.5	52.6	7.2
NEBRASKA	43.4	49.6	6.2
NEVADA	48.3	52.3	4.0
NEW HAMPSHIRE	47.3	54.9	7.6
NEW JERSEY	50.7	53.4	2.7
NEW MEXICO	51.0	47.4	-3.7
NEW YORK	51.9	51.6	-0.4
NORTH CAROLINA	63.7	60.4	-3.3
NORTH DAKOTA	52.1	55.3	3.3
OHIO	47.7	52.5	4.8
OKLAHOMA	45.1	63.1	18.0
OREGON	45.0	45.1	0.1
PENNSYLVANIA	51.2	54.6	3.4
RHODE ISLAND	49.5	53.4	3.9
SOUTH CAROLINA	53.7	56.4	2.7
SOUTH DAKOTA	48.9	50.6	1.7
TENNESSEE	52.2	51.0	-1.2
TEXAS	49.8	53.3	3.5
UTAH	50.0	47.8	-2.2
VERMONT	46.5	52.6	6.2
VIRGINIA	48.5	47.2	-1.3
WASHINGTON	48.1	47.2	-0.9
WEST VIRGINIA	61.3	60.2	-1.0
WISCONSIN	41.1	45.5	4.4
WYOMING	45.6	47.3	1.7
PUERTO RICO	66.2	70.3	4.1

* NOT EATING FEWER CALORIES AND EXERCISING TO LOSE WEIGHT

June 18, 1999

TABLE 13.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

37

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2245	46.2	1.3	(43.7, 48.6)
ALASKA	1987	51.6	1.6	(48.5, 54.7)
ARIZONA	1912	39.1	1.8	(35.7, 42.6)
ARKANSAS	2994	50.2	1.1	(48.1, 52.2)
CALIFORNIA	4154	45.4	0.9	(43.7, 47.1)
COLORADO	1800	47.1	1.3	(44.5, 49.7)
CONNECTICUT	3118	49.2	1.2	(46.9, 51.5)
DELAWARE	2579	48.9	1.4	(46.2, 51.7)
DISTRICT OF CO	1450	38.7	1.6	(35.6, 41.7)
FLORIDA	4716	48.1	0.8	(46.4, 49.7)
GEORGIA	2369	45.6	1.2	(43.2, 47.9)
HAWAII	2149	45.6	1.4	(42.8, 48.4)
IDAHO	4925	45.4	0.8	(43.8, 47.1)
ILLINOIS	2876	48.1	1.1	(46.1, 50.2)
INDIANA	2422	52.0	1.1	(49.8, 54.2)
IOWA	3649	47.1	1.0	(45.1, 49.0)
KANSAS	3795	42.9	0.9	(41.1, 44.8)
KENTUCKY	3706	53.7	0.9	(51.9, 55.5)
LOUISIANA	1652	45.4	1.4	(42.7, 48.1)
MAINE	1615	53.7	1.4	(50.9, 56.4)
MARYLAND	3289	46.6	1.2	(44.3, 48.9)
MASSACHUSETTS	4914	48.9	1.0	(47.0, 50.7)
MICHIGAN	2530	52.8	1.1	(50.7, 55.0)
MINNESOTA	4220	36.6	0.8	(35.0, 38.2)
MISSISSIPPI	2294	46.1	1.2	(43.7, 48.4)
MISSOURI	3715	50.0	1.1	(47.9, 52.1)
MONTANA	1801	48.2	1.3	(45.5, 50.8)
NEBRASKA	2704	46.5	1.1	(44.3, 48.7)
NEVADA	1974	54.4	1.7	(51.1, 57.7)
NEW HAMPSHIRE	1495	51.8	1.5	(48.9, 54.6)
NEW JERSEY	2455	45.5	1.2	(43.3, 47.8)
NEW MEXICO	3632	46.9	0.9	(45.0, 48.7)
NEW YORK	2515	48.4	1.2	(46.1, 50.6)
NORTH CAROLINA	2393	48.1	1.3	(45.5, 50.7)
NORTH DAKOTA	1801	48.2	1.3	(45.6, 50.7)
OHIO	2980	49.7	1.3	(47.2, 52.2)
OKLAHOMA	2514	43.5	1.1	(41.2, 45.7)
OREGON	1813	48.4	1.4	(45.8, 51.1)
PENNSYLVANIA	3587	46.7	1.0	(44.9, 48.6)
RHODE ISLAND	3592	51.6	0.9	(49.8, 53.5)
SOUTH CAROLINA	3288	45.4	1.0	(43.4, 47.4)
SOUTH DAKOTA	2108	47.6	1.2	(45.2, 50.0)
TENNESSEE	2942	47.6	1.1	(45.4, 49.7)
TEXAS	6000	44.5	0.9	(42.8, 46.2)
UTAH	2864	32.9	1.2	(30.5, 35.3)
VERMONT	2988	50.9	1.1	(48.7, 53.1)
VIRGINIA	3494	46.1	1.3	(43.6, 48.6)
WASHINGTON	3584	48.9	1.0	(47.0, 50.8)
WEST VIRGINIA	2411	51.7	1.1	(49.5, 53.9)
WISCONSIN	2202	52.8	1.3	(50.2, 55.4)
WYOMING	2429	48.0	1.2	(45.7, 50.4)
PUERTO RICO	2807	31.7	1.0	(29.7, 33.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 47.8 RANGE = 31.7-54.4

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

June 18, 1999

TABLE 13.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

38

PARTICIPANT	MALES					FEMALES				
	SAMPLE	PERCENT	STANDARD	95% CONFIDENCE	*	SAMPLE	PERCENT	STANDARD	95% CONFIDENCE	*
	SIZE		ERROR	INTERVAL	*	SIZE		ERROR	INTERVAL	*
ALABAMA	785	56.2	2.1	(52.2, 60.2)	*	1460	37.2	1.5	(34.3, 40.1)	*
ALASKA	923	56.4	2.3	(51.8, 60.9)	*	1064	46.3	2.1	(42.3, 50.4)	*
ARIZONA	824	42.1	2.4	(37.4, 46.7)	*	1088	36.4	2.6	(31.3, 41.4)	*
ARKANSAS	1140	60.8	1.6	(57.6, 64.0)	*	1854	40.7	1.3	(38.1, 43.2)	*
CALIFORNIA	1818	50.7	1.3	(48.2, 53.3)	*	2336	40.2	1.1	(38.0, 42.5)	*
COLORADO	810	53.1	2.0	(49.2, 56.9)	*	990	41.4	1.7	(38.1, 44.7)	*
CONNECTICUT	1283	52.3	1.8	(48.8, 55.9)	*	1835	46.3	1.5	(43.4, 49.2)	*
DELAWARE	1036	55.6	2.3	(51.1, 60.0)	*	1543	42.9	1.7	(39.6, 46.2)	*
DISTRICT OF CO	544	41.7	2.6	(36.7, 46.8)	*	906	36.1	1.9	(32.4, 39.7)	*
FLORIDA	1883	53.8	1.3	(51.2, 56.3)	*	2833	42.9	1.0	(40.9, 45.0)	*
GEORGIA	891	54.6	2.0	(50.7, 58.6)	*	1478	37.3	1.4	(34.5, 40.1)	*
HAWAII	933	52.1	2.2	(47.9, 56.4)	*	1216	39.0	1.8	(35.5, 42.4)	*
IDAHO	2083	52.4	1.3	(49.9, 55.0)	*	2842	38.7	1.1	(36.7, 40.8)	*
ILLINOIS	1231	53.9	1.6	(50.7, 57.0)	*	1645	42.9	1.4	(40.1, 45.6)	*
INDIANA	990	60.2	1.7	(56.8, 63.6)	*	1432	44.5	1.4	(41.7, 47.3)	*
IOWA	1448	55.3	1.5	(52.3, 58.3)	*	2201	39.5	1.2	(37.1, 41.9)	*
KANSAS	1483	48.8	1.5	(45.9, 51.8)	*	2312	37.4	1.2	(35.1, 39.7)	*
KENTUCKY	1359	62.8	1.5	(59.9, 65.7)	*	2347	45.5	1.1	(43.2, 47.7)	*
LOUISIANA	647	52.6	2.2	(48.3, 56.8)	*	1005	39.0	1.7	(35.7, 42.2)	*
MAINE	668	58.4	2.2	(54.2, 62.7)	*	947	49.3	1.9	(45.6, 52.9)	*
MARYLAND	1298	49.8	1.8	(46.2, 53.4)	*	1991	43.6	1.5	(40.7, 46.5)	*
MASSACHUSETTS	2006	51.7	1.5	(48.8, 54.6)	*	2908	46.3	1.2	(43.9, 48.8)	*
MICHIGAN	1035	59.7	1.7	(56.4, 63.0)	*	1495	46.6	1.4	(43.8, 49.4)	*
MINNESOTA	1987	39.5	1.2	(37.1, 41.9)	*	2233	33.9	1.1	(31.7, 36.0)	*
MISSISSIPPI	865	56.2	1.9	(52.5, 60.0)	*	1429	37.1	1.4	(34.3, 39.9)	*
MISSOURI	1432	57.2	1.7	(53.8, 60.5)	*	2283	43.6	1.3	(41.1, 46.2)	*
MONTANA	786	52.4	2.0	(48.5, 56.3)	*	1015	44.1	1.7	(40.8, 47.4)	*
NEBRASKA	1050	54.9	1.8	(51.3, 58.5)	*	1654	38.7	1.3	(36.2, 41.3)	*
NEVADA	894	60.2	2.4	(55.5, 64.9)	*	1080	48.5	2.4	(43.7, 53.2)	*
NEW HAMPSHIRE	616	56.9	2.2	(52.6, 61.2)	*	879	46.9	1.9	(43.1, 50.7)	*
NEW JERSEY	1003	50.3	1.9	(46.7, 54.0)	*	1452	41.2	1.5	(38.3, 44.0)	*
NEW MEXICO	1566	54.4	1.4	(51.6, 57.2)	*	2066	39.8	1.2	(37.4, 42.2)	*
NEW YORK	1048	52.5	1.8	(48.9, 56.0)	*	1467	44.7	1.5	(41.8, 47.6)	*
NORTH CAROLINA	916	55.0	2.2	(50.8, 59.2)	*	1477	41.8	1.6	(38.8, 44.9)	*
NORTH DAKOTA	806	55.7	2.0	(51.9, 59.6)	*	995	40.8	1.7	(37.4, 44.2)	*
OHIO	1182	59.4	1.9	(55.6, 63.2)	*	1798	40.9	1.6	(37.7, 44.1)	*
OKLAHOMA	1003	50.0	1.8	(46.5, 53.5)	*	1511	37.5	1.4	(34.8, 40.2)	*
OREGON	739	53.5	2.1	(49.4, 57.6)	*	1074	43.6	1.7	(40.3, 47.0)	*
PENNSYLVANIA	1421	50.7	1.5	(47.7, 53.6)	*	2166	43.3	1.2	(40.9, 45.6)	*
RHODE ISLAND	1408	56.9	1.5	(54.0, 59.8)	*	2184	47.0	1.2	(44.7, 49.3)	*
SOUTH CAROLINA	1373	55.4	1.6	(52.3, 58.6)	*	1915	36.4	1.3	(33.9, 38.8)	*
SOUTH DAKOTA	925	59.8	1.8	(56.2, 63.3)	*	1183	36.1	1.6	(33.1, 39.2)	*
TENNESSEE	1026	57.0	1.7	(53.5, 60.4)	*	1916	39.2	1.3	(36.6, 41.7)	*
TEXAS	2450	54.8	1.3	(52.1, 57.4)	*	3550	34.9	1.0	(32.9, 36.8)	*
UTAH	1238	40.0	1.9	(36.3, 43.7)	*	1626	26.2	1.4	(23.5, 28.9)	*
VERMONT	1276	53.3	1.7	(49.9, 56.7)	*	1712	48.7	1.4	(45.9, 51.4)	*
VIRGINIA	1482	52.7	1.9	(48.9, 56.4)	*	2012	40.0	1.7	(36.7, 43.2)	*
WASHINGTON	1582	53.3	1.5	(50.4, 56.2)	*	2002	44.7	1.3	(42.2, 47.2)	*
WEST VIRGINIA	941	61.0	1.7	(57.6, 64.4)	*	1470	43.4	1.5	(40.5, 46.3)	*
WISCONSIN	975	58.6	1.9	(54.8, 62.3)	*	1227	47.5	1.8	(43.9, 51.0)	*
WYOMING	1020	53.3	1.8	(49.7, 56.9)	*	1409	42.9	1.4	(40.1, 45.7)	*
PUERTO RICO	1075	44.4	1.7	(41.1, 47.7)	*	1732	20.4	1.1	(18.4, 22.5)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
MEDIAN = 54.1
RANGE = 39.5-62.8

NUMBER OF PARTICIPANTS = 52
MEDIAN = 41.3
RANGE = 20.4-49.3

June 18, 1999

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

TABLE 13.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
PREVALENCE DIFFERENCES FROM 1997 TO 1998
EVER SMOKED*
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

39

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	46.7	46.2	-0.5
ALASKA	51.2	51.6	0.4
ARIZONA	34.7	39.1	4.5
ARKANSAS	48.6	50.2	1.5
CALIFORNIA	45.4	45.4	0.1
COLORADO	48.1	47.1	-1.0
CONNECTICUT	50.7	49.2	-1.5
DELAWARE	53.1	48.9	-4.1
DISTRICT OF CO	34.6	38.7	4.0
FLORIDA	50.0	48.1	-1.9
GEORGIA	37.7	45.6	7.8
HAWAII	45.1	45.6	0.5
IDAHO	44.2	45.4	1.2
ILLINOIS	49.0	48.1	-0.8
INDIANA	49.8	52.0	2.1
IOWA	47.2	47.1	-0.1
KANSAS	42.0	42.9	0.9
KENTUCKY	54.2	53.7	-0.5
LOUISIANA	46.4	45.4	-1.0
MAINE	53.7	53.7	-0.1
MARYLAND	43.3	46.6	3.4
MASSACHUSETTS	50.2	48.9	-1.4
MICHIGAN	51.1	52.8	1.8
MINNESOTA	44.3	36.6	-7.7
MISSISSIPPI	44.4	46.1	1.7
MISSOURI	50.3	50.0	-0.2
MONTANA	47.1	48.2	1.0
NEBRASKA	44.9	46.5	1.6
NEVADA	53.4	54.4	1.0
NEW HAMPSHIRE	56.0	51.8	-4.3
NEW JERSEY	47.6	45.5	-2.1
NEW MEXICO	47.4	46.9	-0.5
NEW YORK	46.8	48.4	1.5
NORTH CAROLINA	48.2	48.1	-0.1
NORTH DAKOTA	46.7	48.2	1.5
OHIO	47.9	49.7	1.8
OKLAHOMA	43.7	43.5	-0.3
OREGON	51.1	48.4	-2.7
PENNSYLVANIA	46.9	46.7	-0.2
RHODE ISLAND	51.0	51.6	0.7
SOUTH CAROLINA	43.5	45.4	2.0
SOUTH DAKOTA	50.1	47.6	-2.5
TENNESSEE	48.3	47.6	-0.7
TEXAS	42.9	44.5	1.6
UTAH	31.0	32.9	2.0
VERMONT	52.4	50.9	-1.5
VIRGINIA	50.3	46.1	-4.1
WASHINGTON	50.7	48.9	-1.8
WEST VIRGINIA	51.8	51.7	-0.1
WISCONSIN	48.5	52.8	4.3
WYOMING	49.4	48.0	-1.3
PUERTO RICO	30.1	31.7	1.6

* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

June 18, 1999

TABLE 14.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

40

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2244	24.6	1.1	(22.5, 26.7)
ALASKA	1986	26.1	1.3	(23.5, 28.7)
ARIZONA	1912	21.9	1.3	(19.3, 24.4)
ARKANSAS	2993	25.9	0.9	(24.1, 27.8)
CALIFORNIA	4154	19.2	0.7	(17.8, 20.6)
COLORADO	1800	22.8	1.1	(20.6, 25.1)
CONNECTICUT	3118	21.2	1.0	(19.2, 23.2)
DELAWARE	2579	24.4	1.2	(22.0, 26.9)
DISTRICT OF CO	1450	21.6	1.3	(19.0, 24.3)
FLORIDA	4716	22.0	0.7	(20.6, 23.3)
GEORGIA	2369	23.6	1.0	(21.6, 25.7)
HAWAII	2149	19.5	1.2	(17.2, 21.7)
IDAHO	4925	20.3	0.7	(18.9, 21.6)
ILLINOIS	2876	23.1	0.9	(21.3, 24.9)
INDIANA	2422	26.0	1.0	(24.0, 28.0)
IOWA	3648	23.4	0.9	(21.7, 25.0)
KANSAS	3794	21.1	0.8	(19.6, 22.7)
KENTUCKY	3706	30.8	0.9	(29.1, 32.5)
LOUISIANA	1651	25.5	1.2	(23.1, 27.9)
MAINE	1614	22.4	1.2	(20.0, 24.7)
MARYLAND	3288	22.4	1.0	(20.4, 24.4)
MASSACHUSETTS	4914	20.9	0.8	(19.3, 22.5)
MICHIGAN	2529	27.4	1.0	(25.4, 29.3)
MINNESOTA	4220	18.0	0.7	(16.7, 19.3)
MISSISSIPPI	2292	24.1	1.0	(22.1, 26.1)
MISSOURI	3714	26.4	1.0	(24.4, 28.3)
MONTANA	1801	21.5	1.1	(19.3, 23.6)
NEBRASKA	2704	22.0	0.9	(20.3, 23.8)
NEVADA	1972	30.4	1.6	(27.1, 33.6)
NEW HAMPSHIRE	1495	23.3	1.3	(20.7, 25.8)
NEW JERSEY	2453	19.1	0.9	(17.2, 20.9)
NEW MEXICO	3631	22.6	0.8	(21.0, 24.1)
NEW YORK	2515	24.1	1.0	(22.2, 26.1)
NORTH CAROLINA	2390	24.6	1.1	(22.4, 26.8)
NORTH DAKOTA	1801	20.0	1.0	(18.0, 22.0)
OHIO	2980	26.1	1.2	(23.8, 28.3)
OKLAHOMA	2514	23.9	1.0	(21.9, 25.8)
OREGON	1813	21.1	1.1	(18.9, 23.2)
PENNSYLVANIA	3586	23.8	0.8	(22.2, 25.4)
RHODE ISLAND	3591	22.6	0.8	(21.1, 24.1)
SOUTH CAROLINA	3287	24.7	0.9	(22.9, 26.5)
SOUTH DAKOTA	2108	27.2	1.2	(25.0, 29.5)
TENNESSEE	2942	26.1	1.0	(24.1, 28.0)
TEXAS	6000	21.9	0.7	(20.5, 23.4)
UTAH	2864	14.2	0.8	(12.6, 15.8)
VERMONT	2987	22.3	0.9	(20.5, 24.0)
VIRGINIA	3494	22.9	1.0	(21.0, 24.8)
WASHINGTON	3583	21.4	0.8	(19.8, 22.9)
WEST VIRGINIA	2411	27.9	1.0	(25.9, 30.0)
WISCONSIN	2202	23.4	1.2	(21.1, 25.7)
WYOMING	2428	22.8	1.0	(20.9, 24.8)
PUERTO RICO	2807	15.7	0.8	(14.1, 17.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 22.9 RANGE = 14.2-30.8 YEAR 2000 OBJ. 3.4

*SMOKED 100 CIGARETTES IN LIFETIME AND NOW SMOKE EVERYDAY OR SOME DAYS June 18, 1999

TABLE 14.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

41

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	784	27.2	1.8	(23.7, 30.7)	*	1460	22.3	1.3	(19.8, 24.8)	*
ALASKA	922	28.3	2.0	(24.4, 32.2)	*	1064	23.7	1.7	(20.4, 27.1)	*
ARIZONA	824	24.6	2.0	(20.7, 28.6)	*	1088	19.2	1.7	(15.9, 22.5)	*
ARKANSAS	1139	28.5	1.5	(25.5, 31.5)	*	1854	23.7	1.1	(21.4, 25.9)	*
CALIFORNIA	1818	21.9	1.1	(19.7, 24.1)	*	2336	16.6	0.9	(14.9, 18.3)	*
COLORADO	810	26.4	1.8	(22.8, 29.9)	*	990	19.4	1.3	(16.8, 22.1)	*
CONNECTICUT	1283	22.1	1.7	(18.7, 25.4)	*	1835	20.4	1.1	(18.1, 22.6)	*
DELAWARE	1036	27.3	2.1	(23.3, 31.4)	*	1543	21.8	1.4	(19.0, 24.7)	*
DISTRICT OF CO	544	24.4	2.2	(20.0, 28.8)	*	906	19.3	1.6	(16.1, 22.4)	*
FLORIDA	1883	23.4	1.1	(21.3, 25.6)	*	2833	20.6	0.8	(19.0, 22.2)	*
GEORGIA	891	28.0	1.7	(24.5, 31.4)	*	1478	19.7	1.2	(17.3, 22.0)	*
HAWAII	933	22.2	1.9	(18.6, 25.8)	*	1216	16.7	1.4	(14.0, 19.4)	*
IDAHO	2083	21.9	1.1	(19.7, 24.0)	*	2842	18.7	0.9	(17.0, 20.4)	*
ILLINOIS	1231	25.9	1.4	(23.2, 28.7)	*	1645	20.6	1.2	(18.3, 22.9)	*
INDIANA	990	29.6	1.6	(26.5, 32.8)	*	1432	22.7	1.2	(20.3, 25.0)	*
IOWA	1447	25.9	1.4	(23.2, 28.5)	*	2201	21.1	1.0	(19.1, 23.2)	*
KANSAS	1482	23.0	1.3	(20.5, 25.4)	*	2312	19.4	1.0	(17.6, 21.3)	*
KENTUCKY	1359	33.4	1.5	(30.5, 36.2)	*	2347	28.5	1.0	(26.4, 30.5)	*
LOUISIANA	646	28.2	2.0	(24.3, 32.1)	*	1005	23.1	1.5	(20.2, 26.1)	*
MAINE	667	21.2	1.8	(17.6, 24.7)	*	947	23.5	1.6	(20.3, 26.7)	*
MARYLAND	1298	24.3	1.6	(21.1, 27.5)	*	1990	20.7	1.2	(18.3, 23.0)	*
MASSACHUSETTS	2006	22.5	1.3	(19.9, 25.0)	*	2908	19.5	1.0	(17.6, 21.4)	*
MICHIGAN	1034	30.3	1.6	(27.1, 33.4)	*	1495	24.8	1.2	(22.3, 27.2)	*
MINNESOTA	1987	19.7	1.0	(17.8, 21.6)	*	2233	16.4	0.8	(14.8, 18.1)	*
MISSISSIPPI	865	26.9	1.7	(23.5, 30.3)	*	1427	21.6	1.2	(19.2, 24.0)	*
MISSOURI	1432	29.4	1.7	(26.2, 32.6)	*	2282	23.6	1.2	(21.4, 25.9)	*
MONTANA	786	21.5	1.6	(18.4, 24.5)	*	1015	21.4	1.5	(18.6, 24.3)	*
NEBRASKA	1050	25.3	1.5	(22.4, 28.1)	*	1654	19.1	1.1	(17.0, 21.2)	*
NEVADA	893	32.6	2.4	(28.0, 37.3)	*	1079	28.0	2.4	(23.3, 32.7)	*
NEW HAMPSHIRE	616	25.7	2.0	(21.7, 29.7)	*	879	21.0	1.7	(17.7, 24.2)	*
NEW JERSEY	1002	20.9	1.6	(17.8, 23.9)	*	1451	17.4	1.1	(15.3, 19.6)	*
NEW MEXICO	1565	25.1	1.2	(22.7, 27.5)	*	2066	20.2	1.0	(18.2, 22.2)	*
NEW YORK	1048	25.6	1.6	(22.6, 28.7)	*	1467	22.8	1.3	(20.3, 25.2)	*
NORTH CAROLINA	915	27.4	1.8	(23.8, 31.0)	*	1475	22.2	1.3	(19.5, 24.8)	*
NORTH DAKOTA	806	21.8	1.6	(18.7, 24.8)	*	995	18.3	1.4	(15.6, 20.9)	*
OHIO	1182	29.7	1.9	(26.0, 33.3)	*	1798	22.8	1.4	(20.1, 25.6)	*
OKLAHOMA	1003	26.7	1.7	(23.5, 30.0)	*	1511	21.2	1.2	(18.9, 23.6)	*
OREGON	739	21.6	1.7	(18.2, 25.0)	*	1074	20.5	1.4	(17.8, 23.2)	*
PENNSYLVANIA	1420	24.1	1.3	(21.6, 26.6)	*	2166	23.5	1.0	(21.5, 25.6)	*
RHODE ISLAND	1408	24.0	1.3	(21.5, 26.6)	*	2183	21.3	1.0	(19.4, 23.2)	*
SOUTH CAROLINA	1372	29.8	1.5	(26.8, 32.7)	*	1915	20.1	1.0	(18.1, 22.2)	*
SOUTH DAKOTA	925	36.5	1.9	(32.9, 40.2)	*	1183	18.4	1.2	(16.1, 20.8)	*
TENNESSEE	1026	30.2	1.6	(27.0, 33.4)	*	1916	22.3	1.1	(20.1, 24.5)	*
TEXAS	2450	25.3	1.2	(22.9, 27.7)	*	3550	18.8	0.8	(17.2, 20.4)	*
UTAH	1238	16.0	1.3	(13.4, 18.5)	*	1626	12.6	1.0	(10.6, 14.5)	*
VERMONT	1275	23.7	1.4	(21.0, 26.4)	*	1712	21.0	1.2	(18.7, 23.2)	*
VIRGINIA	1482	25.8	1.6	(22.6, 28.9)	*	2012	20.2	1.2	(17.9, 22.5)	*
WASHINGTON	1582	22.4	1.3	(19.9, 24.8)	*	2001	20.4	1.1	(18.3, 22.4)	*
WEST VIRGINIA	941	29.7	1.7	(26.4, 33.0)	*	1470	26.4	1.3	(23.9, 28.8)	*
WISCONSIN	975	24.0	1.7	(20.6, 27.4)	*	1227	22.8	1.6	(19.7, 26.0)	*
WYOMING	1020	23.9	1.6	(20.8, 27.0)	*	1408	21.8	1.2	(19.5, 24.1)	*
PUERTO RICO	1075	21.9	1.4	(19.1, 24.7)	*	1732	10.1	0.8	(8.5, 11.7)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52

MEDIAN = 25.3

RANGE = 16.0-36.5

NUMBER OF PARTICIPANTS = 52

MEDIAN = 20.8

RANGE = 10.1-28.5

*SMOKED 100 CIGARETTES IN LIFETIME AND NOW SMOKE EVERYDAY OR SOME DAYS June 18, 1999

TABLE 14.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 CURRENT SMOKERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

42

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	24.7	24.6	-0.0
ALASKA	26.5	26.1	-0.4
ARIZONA	21.1	21.9	0.8
ARKANSAS	28.4	25.9	-2.5
CALIFORNIA	18.4	19.2	0.8
COLORADO	22.6	22.8	0.3
CONNECTICUT	21.6	21.2	-0.4
DELAWARE	26.6	24.4	-2.1
DISTRICT OF CO	18.8	21.6	2.8
FLORIDA	23.6	22.0	-1.6
GEORGIA	22.4	23.6	1.2
HAWAII	18.7	19.5	0.8
IDAHO	19.9	20.3	0.3
ILLINOIS	23.2	23.1	-0.1
INDIANA	26.4	26.0	-0.4
IOWA	23.1	23.4	0.3
KANSAS	22.6	21.1	-1.5
KENTUCKY	30.7	30.8	0.1
LOUISIANA	24.5	25.5	1.1
MAINE	22.7	22.4	-0.3
MARYLAND	20.4	22.4	1.9
MASSACHUSETTS	20.5	20.9	0.4
MICHIGAN	26.0	27.4	1.3
MINNESOTA	21.9	18.0	-3.9
MISSISSIPPI	23.1	24.1	1.0
MISSOURI	28.6	26.4	-2.3
MONTANA	20.5	21.5	1.0
NEBRASKA	22.1	22.0	-0.1
NEVADA	28.0	30.4	2.3
NEW HAMPSHIRE	24.7	23.3	-1.4
NEW JERSEY	21.4	19.1	-2.3
NEW MEXICO	22.1	22.6	0.5
NEW YORK	23.1	24.1	1.0
NORTH CAROLINA	25.9	24.6	-1.2
NORTH DAKOTA	22.3	20.0	-2.3
OHIO	25.1	26.1	1.0
OKLAHOMA	24.6	23.9	-0.7
OREGON	20.7	21.1	0.4
PENNSYLVANIA	24.2	23.8	-0.4
RHODE ISLAND	24.3	22.6	-1.7
SOUTH CAROLINA	23.4	24.7	1.3
SOUTH DAKOTA	24.3	27.2	2.9
TENNESSEE	26.9	26.1	-0.9
TEXAS	22.6	21.9	-0.6
UTAH	13.8	14.2	0.5
VERMONT	23.3	22.3	-1.0
VIRGINIA	24.4	22.9	-1.5
WASHINGTON	23.8	21.4	-2.5
WEST VIRGINIA	27.4	27.9	0.5
WISCONSIN	23.2	23.4	0.2
WYOMING	24.0	22.8	-1.1
PUERTO RICO	14.5	15.7	1.2

*SMOKED 100 CIGARETTES IN LIFETIME AND NOW SMOKE EVERYDAY OR SOME DAYS June 18, 1999

TABLE 15.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

43

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2149	35.8	1.2	(33.4, 38.3)
ALASKA	1912	36.2	1.6	(33.2, 39.3)
ARIZONA	1850	22.5	1.7	(19.1, 25.8)
ARKANSAS	2837	33.4	1.0	(31.4, 35.5)
CALIFORNIA	4046	31.0	0.8	(29.3, 32.6)
COLORADO	1770	27.2	1.2	(24.8, 29.5)
CONNECTICUT	2922	27.4	1.1	(25.4, 29.5)
DELAWARE	2492	32.8	1.4	(30.2, 35.5)
DISTRICT OF CO	1431	31.8	1.5	(28.8, 34.7)
FLORIDA	4584	32.3	0.8	(30.7, 33.9)
GEORGIA	2297	33.3	1.1	(31.1, 35.5)
HAWAII	2122	27.8	1.3	(25.2, 30.4)
IDAHO	4782	29.9	0.8	(28.4, 31.4)
ILLINOIS	2790	33.5	1.0	(31.5, 35.5)
INDIANA	2373	34.5	1.1	(32.4, 36.6)
IOWA	3569	34.6	0.9	(32.7, 36.4)
KANSAS	3549	31.7	0.9	(30.0, 33.5)
KENTUCKY	3593	35.8	0.9	(34.0, 37.6)
LOUISIANA	1587	36.1	1.4	(33.4, 38.8)
MAINE	1550	31.2	1.3	(28.6, 33.8)
MARYLAND	3161	34.5	1.2	(32.2, 36.7)
MASSACHUSETTS	4617	26.9	0.9	(25.2, 28.5)
MICHIGAN	2457	34.6	1.1	(32.5, 36.7)
MINNESOTA	4120	32.5	0.8	(30.9, 34.0)
MISSISSIPPI	2183	37.5	1.2	(35.2, 39.8)
MISSOURI	3621	35.6	1.1	(33.4, 37.7)
MONTANA	1728	29.1	1.2	(26.8, 31.4)
NEBRASKA	2594	32.3	1.0	(30.3, 34.3)
NEVADA	1909	27.0	1.4	(24.2, 29.7)
NEW HAMPSHIRE	1430	28.7	1.4	(25.9, 31.4)
NEW JERSEY	2313	29.7	1.1	(27.5, 31.9)
NEW MEXICO	3547	29.9	0.9	(28.2, 31.6)
NEW YORK	2383	28.6	1.1	(26.5, 30.8)
NORTH CAROLINA	2258	32.7	1.3	(30.3, 35.2)
NORTH DAKOTA	1767	33.5	1.3	(31.1, 36.0)
OHIO	2835	33.6	1.2	(31.2, 36.0)
OKLAHOMA	2414	32.4	1.1	(30.3, 34.5)
OREGON	1768	32.8	1.3	(30.3, 35.3)
PENNSYLVANIA	3410	34.8	0.9	(33.0, 36.7)
RHODE ISLAND	3404	30.3	0.9	(28.5, 32.1)
SOUTH CAROLINA	3139	34.6	1.0	(32.6, 36.5)
SOUTH DAKOTA	2055	30.5	1.2	(28.2, 32.8)
TENNESSEE	2818	32.4	1.0	(30.4, 34.4)
TEXAS	5768	34.9	0.8	(33.3, 36.5)
UTAH	2775	28.8	1.2	(26.5, 31.1)
VERMONT	2840	27.4	1.0	(25.4, 29.3)
VIRGINIA	3353	31.8	1.2	(29.6, 34.1)
WASHINGTON	3421	32.3	1.0	(30.5, 34.2)
WEST VIRGINIA	2338	37.8	1.2	(35.5, 40.1)
WISCONSIN	2170	34.0	1.3	(31.6, 36.5)
WYOMING	2371	28.1	1.0	(26.1, 30.1)
PUERTO RICO	2643	33.6	1.0	(31.6, 35.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 32.4 RANGE = 22.5-37.8 YEAR 2000 OBJ. 2.3
 * BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES

June 18, 1999

TABLE 15.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

44

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	778	37.7	2.0	(33.8, 41.7)	*	1371	34.0	1.4	(31.2, 36.8)	*
ALASKA	904	38.8	2.3	(34.2, 43.4)	*	1008	33.3	2.0	(29.4, 37.2)	*
ARIZONA	809	23.0	2.2	(18.8, 27.3)	*	1041	21.9	2.6	(16.8, 27.1)	*
ARKANSAS	1114	35.9	1.6	(32.8, 39.1)	*	1723	31.1	1.4	(28.4, 33.7)	*
CALIFORNIA	1794	31.1	1.2	(28.7, 33.5)	*	2252	30.8	1.1	(28.6, 33.0)	*
COLORADO	805	28.3	1.9	(24.7, 31.9)	*	965	26.1	1.6	(23.0, 29.2)	*
CONNECTICUT	1259	30.2	1.7	(27.0, 33.5)	*	1663	24.7	1.3	(22.2, 27.2)	*
DELAWARE	1024	34.2	2.2	(29.9, 38.6)	*	1468	31.5	1.6	(28.4, 34.6)	*
DISTRICT OF CO	542	28.9	2.5	(24.0, 33.7)	*	889	34.2	1.8	(30.6, 37.8)	*
FLORIDA	1867	33.8	1.3	(31.3, 36.2)	*	2717	30.9	1.0	(28.9, 32.9)	*
GEORGIA	882	32.4	1.8	(28.9, 35.8)	*	1415	34.2	1.4	(31.5, 37.0)	*
HAWAII	930	31.5	2.1	(27.5, 35.6)	*	1192	24.0	1.6	(20.8, 27.2)	*
IDAHO	2070	31.4	1.2	(29.1, 33.8)	*	2712	28.4	1.0	(26.5, 30.3)	*
ILLINOIS	1220	33.9	1.5	(30.9, 36.9)	*	1570	33.1	1.4	(30.4, 35.7)	*
INDIANA	986	36.6	1.7	(33.2, 39.9)	*	1387	32.5	1.4	(29.8, 35.2)	*
IOWA	1437	35.8	1.5	(33.0, 38.7)	*	2132	33.3	1.2	(31.0, 35.7)	*
KANSAS	1456	33.8	1.4	(31.2, 36.5)	*	2093	29.6	1.1	(27.4, 31.9)	*
KENTUCKY	1351	37.2	1.5	(34.3, 40.0)	*	2242	34.6	1.1	(32.4, 36.7)	*
LOUISIANA	644	36.9	2.2	(32.5, 41.3)	*	943	35.3	1.8	(31.8, 38.8)	*
MAINE	661	31.3	2.0	(27.4, 35.2)	*	889	31.1	1.8	(27.5, 34.6)	*
MARYLAND	1288	35.6	1.8	(32.0, 39.1)	*	1873	33.4	1.5	(30.5, 36.3)	*
MASSACHUSETTS	1973	30.9	1.4	(28.2, 33.5)	*	2644	23.0	1.1	(20.9, 25.0)	*
MICHIGAN	1029	33.3	1.6	(30.1, 36.4)	*	1428	35.8	1.4	(33.1, 38.6)	*
MINNESOTA	1971	36.6	1.2	(34.2, 38.9)	*	2149	28.5	1.0	(26.5, 30.5)	*
MISSISSIPPI	850	36.6	1.9	(32.9, 40.2)	*	1333	38.4	1.5	(35.4, 41.4)	*
MISSOURI	1424	34.9	1.7	(31.6, 38.2)	*	2197	36.2	1.4	(33.6, 38.8)	*
MONTANA	777	29.9	1.8	(26.5, 33.4)	*	951	28.2	1.5	(25.3, 31.1)	*
NEBRASKA	1039	33.3	1.6	(30.2, 36.4)	*	1555	31.3	1.3	(28.8, 33.8)	*
NEVADA	889	28.2	2.0	(24.3, 32.0)	*	1020	25.7	2.1	(21.6, 29.8)	*
NEW HAMPSHIRE	607	31.4	2.1	(27.3, 35.4)	*	823	26.0	1.7	(22.6, 29.3)	*
NEW JERSEY	990	32.5	1.8	(29.0, 35.9)	*	1323	27.0	1.5	(24.1, 29.8)	*
NEW MEXICO	1559	29.2	1.3	(26.7, 31.7)	*	1988	30.6	1.2	(28.3, 32.9)	*
NEW YORK	1029	27.7	1.6	(24.5, 30.9)	*	1354	29.5	1.4	(26.7, 32.3)	*
NORTH CAROLINA	902	32.0	2.0	(28.1, 35.8)	*	1356	33.5	1.6	(30.5, 36.5)	*
NORTH DAKOTA	804	36.3	1.9	(32.6, 40.0)	*	963	30.8	1.6	(27.6, 34.0)	*
OHIO	1169	35.4	1.9	(31.7, 39.1)	*	1666	31.9	1.6	(28.7, 35.0)	*
OKLAHOMA	980	33.2	1.7	(29.9, 36.5)	*	1434	31.7	1.4	(29.0, 34.3)	*
OREGON	738	34.1	2.0	(30.3, 38.0)	*	1030	31.5	1.6	(28.4, 34.6)	*
PENNSYLVANIA	1401	35.8	1.5	(32.9, 38.6)	*	2009	33.9	1.2	(31.6, 36.3)	*
RHODE ISLAND	1392	32.8	1.4	(30.0, 35.5)	*	2012	28.0	1.1	(25.8, 30.2)	*
SOUTH CAROLINA	1353	36.1	1.5	(33.1, 39.1)	*	1786	33.1	1.3	(30.5, 35.7)	*
SOUTH DAKOTA	922	32.9	1.8	(29.4, 36.5)	*	1133	28.1	1.5	(25.2, 31.1)	*
TENNESSEE	1008	31.9	1.7	(28.7, 35.2)	*	1810	32.9	1.2	(30.4, 35.3)	*
TEXAS	2414	34.9	1.3	(32.5, 37.4)	*	3354	34.9	1.1	(32.7, 37.1)	*
UTAH	1229	30.8	1.8	(27.3, 34.4)	*	1546	26.8	1.5	(23.9, 29.6)	*
VERMONT	1263	28.5	1.5	(25.5, 31.4)	*	1577	26.3	1.2	(23.8, 28.7)	*
VIRGINIA	1466	31.0	1.6	(27.8, 34.2)	*	1887	32.7	1.7	(29.4, 35.9)	*
WASHINGTON	1563	34.8	1.4	(31.9, 37.6)	*	1858	29.8	1.3	(27.3, 32.3)	*
WEST VIRGINIA	936	39.6	1.8	(36.1, 43.1)	*	1402	36.1	1.4	(33.4, 38.8)	*
WISCONSIN	976	36.4	1.9	(32.7, 40.0)	*	1194	31.8	1.7	(28.5, 35.2)	*
WYOMING	1014	28.2	1.6	(25.2, 31.3)	*	1357	27.9	1.3	(25.4, 30.5)	*
PUERTO RICO	1048	33.5	1.6	(30.4, 36.6)	*	1595	33.7	1.3	(31.1, 36.2)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 33.3
 RANGE = 23.0-39.6

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 31.4
 RANGE = 21.9-38.4

June 18, 1999

* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES

TABLE 15.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 OVERWEIGHT: BY BODY MASS INDEX*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

45

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	32.7	35.8	3.1
ALASKA	34.1	36.2	2.1
ARIZONA	25.4	22.5	-2.9
ARKANSAS	32.1	33.4	1.3
CALIFORNIA	29.1	31.0	1.8
COLORADO	25.1	27.2	2.0
CONNECTICUT	29.1	27.4	-1.6
DELAWARE	32.5	32.8	0.3
DISTRICT OF CO	28.0	31.8	3.7
FLORIDA	30.4	32.3	1.9
GEORGIA	30.8	33.3	2.6
HAWAII	26.6	27.8	1.3
IDAHO	29.6	29.9	0.4
ILLINOIS	32.0	33.5	1.5
INDIANA	34.6	34.5	-0.2
IOWA	33.3	34.6	1.2
KANSAS	31.5	31.7	0.2
KENTUCKY	35.6	35.8	0.2
LOUISIANA	33.5	36.1	2.6
MAINE	30.2	31.2	1.0
MARYLAND	31.2	34.5	3.2
MASSACHUSETTS	25.5	26.9	1.4
MICHIGAN	34.9	34.6	-0.3
MINNESOTA	29.7	32.5	2.7
MISSISSIPPI	35.1	37.5	2.4
MISSOURI	31.6	35.6	4.0
MONTANA	27.9	29.1	1.2
NEBRASKA	31.1	32.3	1.2
NEVADA	26.2	27.0	0.8
NEW HAMPSHIRE	27.5	28.7	1.1
NEW JERSEY	28.2	29.7	1.5
NEW MEXICO	29.8	29.9	0.1
NEW YORK	28.4	28.6	0.3
NORTH CAROLINA	32.4	32.7	0.3
NORTH DAKOTA	32.9	33.5	0.6
OHIO	32.6	33.6	1.0
OKLAHOMA	28.4	32.4	4.0
OREGON	32.4	32.8	0.4
PENNSYLVANIA	32.2	34.8	2.6
RHODE ISLAND	28.1	30.3	2.2
SOUTH CAROLINA	31.8	34.6	2.8
SOUTH DAKOTA	31.1	30.5	-0.6
TENNESSEE	31.6	32.4	0.8
TEXAS	32.8	34.9	2.1
UTAH	27.0	28.8	1.8
VERMONT	29.9	27.4	-2.5
VIRGINIA	30.0	31.8	1.8
WASHINGTON	29.1	32.3	3.2
WEST VIRGINIA	36.3	37.8	1.5
WISCONSIN	32.8	34.0	1.2
WYOMING	27.9	28.1	0.2
PUERTO RICO	34.9	33.6	-1.3

* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES

June 18, 1999

TABLE 16.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY MEDIAN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

46

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2146	41.6	1.3	(39.1, 44.1)
ALASKA	1907	40.8	1.6	(37.7, 43.9)
ARIZONA	1848	26.6	1.8	(23.2, 30.1)
ARKANSAS	2836	38.3	1.1	(36.2, 40.4)
CALIFORNIA	4038	35.6	0.9	(33.9, 37.3)
COLORADO	1766	32.0	1.3	(29.6, 34.5)
CONNECTICUT	2920	31.9	1.1	(29.7, 34.0)
DELAWARE	2491	36.4	1.4	(33.7, 39.1)
DISTRICT OF CO	1430	35.7	1.6	(32.7, 38.7)
FLORIDA	4579	37.2	0.9	(35.6, 38.9)
GEORGIA	2292	37.1	1.2	(34.8, 39.4)
HAWAII	2121	31.2	1.4	(28.5, 33.8)
IDAHO	4779	34.9	0.8	(33.3, 36.5)
ILLINOIS	2782	38.2	1.0	(36.2, 40.2)
INDIANA	2371	38.5	1.1	(36.3, 40.6)
IOWA	3567	39.7	1.0	(37.8, 41.6)
KANSAS	3545	36.6	0.9	(34.8, 38.4)
KENTUCKY	3581	40.6	0.9	(38.7, 42.4)
LOUISIANA	1588	41.7	1.4	(38.9, 44.4)
MAINE	1547	36.0	1.4	(33.3, 38.8)
MARYLAND	3156	38.7	1.2	(36.4, 41.0)
MASSACHUSETTS	4610	32.0	0.9	(30.2, 33.8)
MICHIGAN	2454	39.0	1.1	(36.9, 41.2)
MINNESOTA	4117	38.8	0.8	(37.1, 40.4)
MISSISSIPPI	2180	42.2	1.2	(39.9, 44.6)
MISSOURI	3619	39.6	1.1	(37.4, 41.8)
MONTANA	1724	33.1	1.2	(30.7, 35.5)
NEBRASKA	2593	37.2	1.1	(35.1, 39.3)
NEVADA	1903	31.2	1.5	(28.2, 34.1)
NEW HAMPSHIRE	1429	34.0	1.4	(31.2, 36.8)
NEW JERSEY	2311	35.2	1.2	(32.9, 37.6)
NEW MEXICO	3544	34.4	0.9	(32.6, 36.1)
NEW YORK	2383	33.2	1.1	(31.0, 35.4)
NORTH CAROLINA	2256	38.4	1.3	(35.9, 40.9)
NORTH DAKOTA	1766	38.9	1.3	(36.3, 41.5)
OHIO	2836	38.3	1.3	(35.8, 40.8)
OKLAHOMA	2410	36.6	1.1	(34.4, 38.8)
OREGON	1764	38.3	1.3	(35.7, 40.8)
PENNSYLVANIA	3412	39.7	1.0	(37.8, 41.5)
RHODE ISLAND	3402	35.3	0.9	(33.5, 37.1)
SOUTH CAROLINA	3134	39.5	1.0	(37.5, 41.6)
SOUTH DAKOTA	2052	34.8	1.2	(32.4, 37.1)
TENNESSEE	2817	37.1	1.1	(35.0, 39.1)
TEXAS	5762	40.0	0.9	(38.3, 41.7)
UTAH	2772	33.6	1.2	(31.2, 36.0)
VERMONT	2838	32.2	1.0	(30.2, 34.2)
VIRGINIA	3346	36.2	1.2	(33.8, 38.5)
WASHINGTON	3419	37.6	1.0	(35.7, 39.5)
WEST VIRGINIA	2334	43.0	1.2	(40.7, 45.3)
WISCONSIN	2168	39.2	1.3	(36.6, 41.7)
WYOMING	2367	33.0	1.1	(30.9, 35.0)
PUERTO RICO	2641	38.5	1.1	(36.4, 40.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 37.2 RANGE = 26.6-43.0

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

June 18, 1999

TABLE 16.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY MEDIAN* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

47

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	778	43.8	2.1	(39.8, 47.8)	*	1368	39.5	1.5	(36.5, 42.4)	*
ALASKA	900	44.3	2.4	(39.6, 48.9)	*	1007	36.9	2.0	(32.9, 40.8)	*
ARIZONA	809	26.5	2.3	(22.1, 31.0)	*	1039	26.7	2.7	(21.5, 31.9)	*
ARKANSAS	1114	39.1	1.6	(35.9, 42.2)	*	1722	37.7	1.4	(34.9, 40.4)	*
CALIFORNIA	1788	35.0	1.3	(32.5, 37.5)	*	2250	36.1	1.2	(33.8, 38.4)	*
COLORADO	804	34.3	1.9	(30.5, 38.0)	*	962	29.9	1.6	(26.7, 33.1)	*
CONNECTICUT	1259	35.5	1.8	(32.0, 39.0)	*	1661	28.3	1.4	(25.7, 31.0)	*
DELAWARE	1023	37.4	2.2	(33.0, 41.8)	*	1468	35.3	1.6	(32.2, 38.5)	*
DISTRICT OF CO	541	32.1	2.5	(27.1, 37.0)	*	889	38.8	1.9	(35.1, 42.5)	*
FLORIDA	1865	38.1	1.3	(35.5, 40.6)	*	2714	36.4	1.1	(34.3, 38.5)	*
GEORGIA	878	35.4	1.8	(31.8, 38.9)	*	1414	38.8	1.5	(35.9, 41.6)	*
HAWAII	929	34.5	2.1	(30.4, 38.6)	*	1192	27.8	1.7	(24.4, 31.1)	*
IDAHO	2068	36.5	1.3	(34.0, 38.9)	*	2711	33.3	1.0	(31.3, 35.4)	*
ILLINOIS	1215	37.7	1.6	(34.7, 40.8)	*	1567	38.7	1.4	(35.9, 41.4)	*
INDIANA	986	40.2	1.7	(36.9, 43.6)	*	1385	36.8	1.4	(34.1, 39.6)	*
IOWA	1436	40.6	1.5	(37.7, 43.6)	*	2131	38.8	1.3	(36.4, 41.3)	*
KANSAS	1453	38.8	1.4	(35.9, 41.6)	*	2092	34.5	1.2	(32.1, 36.8)	*
KENTUCKY	1345	42.8	1.5	(39.8, 45.7)	*	2236	38.5	1.1	(36.3, 40.8)	*
LOUISIANA	644	43.0	2.3	(38.5, 47.5)	*	944	40.4	1.8	(36.9, 43.9)	*
MAINE	658	35.8	2.1	(31.7, 39.9)	*	889	36.2	1.9	(32.6, 39.9)	*
MARYLAND	1287	39.2	1.8	(35.6, 42.7)	*	1869	38.3	1.5	(35.4, 41.3)	*
MASSACHUSETTS	1968	36.1	1.4	(33.4, 38.9)	*	2642	28.1	1.2	(25.8, 30.3)	*
MICHIGAN	1025	37.6	1.7	(34.3, 40.8)	*	1429	40.5	1.4	(37.6, 43.3)	*
MINNESOTA	1969	42.2	1.2	(39.8, 44.6)	*	2148	35.4	1.1	(33.3, 37.6)	*
MISSISSIPPI	848	40.4	1.9	(36.7, 44.1)	*	1332	43.9	1.5	(40.9, 46.9)	*
MISSOURI	1422	38.9	1.7	(35.5, 42.3)	*	2197	40.2	1.4	(37.6, 42.9)	*
MONTANA	775	33.0	1.8	(29.4, 36.6)	*	949	33.1	1.5	(30.1, 36.1)	*
NEBRASKA	1038	38.1	1.8	(34.7, 41.6)	*	1555	36.3	1.3	(33.7, 38.9)	*
NEVADA	887	32.2	2.1	(28.1, 36.3)	*	1016	30.0	2.2	(25.7, 34.3)	*
NEW HAMPSHIRE	606	36.5	2.2	(32.2, 40.7)	*	823	31.6	1.8	(28.1, 35.1)	*
NEW JERSEY	990	37.3	1.8	(33.7, 40.9)	*	1321	33.2	1.5	(30.2, 36.2)	*
NEW MEXICO	1558	33.8	1.3	(31.2, 36.4)	*	1986	34.9	1.2	(32.5, 37.3)	*
NEW YORK	1029	32.2	1.7	(28.8, 35.5)	*	1354	34.2	1.5	(31.3, 37.1)	*
NORTH CAROLINA	902	37.8	2.1	(33.8, 41.9)	*	1354	39.0	1.6	(35.9, 42.1)	*
NORTH DAKOTA	804	42.5	2.0	(38.6, 46.3)	*	962	35.3	1.7	(32.0, 38.7)	*
OHIO	1170	39.1	1.9	(35.3, 42.9)	*	1666	37.5	1.7	(34.2, 40.8)	*
OKLAHOMA	979	37.2	1.7	(33.8, 40.6)	*	1431	36.1	1.4	(33.3, 38.9)	*
OREGON	735	39.6	2.0	(35.7, 43.6)	*	1029	37.0	1.7	(33.7, 40.2)	*
PENNSYLVANIA	1402	40.1	1.5	(37.2, 43.0)	*	2010	39.2	1.2	(36.8, 41.7)	*
RHODE ISLAND	1392	38.0	1.5	(35.2, 40.9)	*	2010	32.7	1.2	(30.4, 35.0)	*
SOUTH CAROLINA	1350	40.9	1.6	(37.8, 44.0)	*	1784	38.2	1.4	(35.5, 40.8)	*
SOUTH DAKOTA	920	37.7	1.8	(34.1, 41.3)	*	1132	31.9	1.6	(28.9, 35.0)	*
TENNESSEE	1007	36.7	1.7	(33.4, 40.0)	*	1810	37.4	1.3	(34.9, 39.9)	*
TEXAS	2411	40.2	1.3	(37.6, 42.7)	*	3351	39.8	1.1	(37.6, 42.0)	*
UTAH	1228	33.9	1.9	(30.3, 37.5)	*	1544	33.3	1.6	(30.1, 36.4)	*
VERMONT	1263	33.5	1.6	(30.4, 36.7)	*	1575	30.9	1.3	(28.3, 33.5)	*
VIRGINIA	1462	34.9	1.7	(31.6, 38.3)	*	1884	37.3	1.7	(34.0, 40.7)	*
WASHINGTON	1562	40.3	1.5	(37.4, 43.2)	*	1857	34.8	1.3	(32.3, 37.4)	*
WEST VIRGINIA	935	44.4	1.8	(40.8, 47.9)	*	1399	41.8	1.4	(39.0, 44.6)	*
WISCONSIN	974	40.6	1.9	(36.9, 44.3)	*	1194	37.8	1.8	(34.3, 41.3)	*
WYOMING	1012	33.4	1.7	(30.1, 36.6)	*	1355	32.5	1.4	(29.9, 35.2)	*
PUERTO RICO	1049	37.8	1.6	(34.6, 41.1)	*	1592	39.1	1.4	(36.5, 41.8)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52
 MEDIAN = 37.8
 RANGE = 26.5-44.4

NUMBER OF PARTICIPANTS = 52
 MEDIAN = 36.4
 RANGE = 26.7-43.9

June 18, 1999

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

TABLE 16.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 OVERWEIGHT: BY MEDIAN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

48

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	38.3	41.6	3.3
ALASKA	36.6	40.8	4.2
ARIZONA	29.4	26.6	-2.8
ARKANSAS	36.7	38.3	1.7
CALIFORNIA	33.6	35.6	2.0
COLORADO	29.4	32.0	2.6
CONNECTICUT	34.1	31.9	-2.2
DELAWARE	37.4	36.4	-1.1
DISTRICT OF CO	32.8	35.7	2.9
FLORIDA	35.4	37.2	1.9
GEORGIA	37.8	37.1	-0.6
HAWAII	31.3	31.2	-0.1
IDAHO	34.8	34.9	0.1
ILLINOIS	36.4	38.2	1.8
INDIANA	39.2	38.5	-0.7
IOWA	38.0	39.7	1.7
KANSAS	36.0	36.6	0.6
KENTUCKY	40.8	40.6	-0.2
LOUISIANA	38.4	41.7	3.3
MAINE	36.0	36.0	0.0
MARYLAND	35.0	38.7	3.7
MASSACHUSETTS	30.3	32.0	1.7
MICHIGAN	39.9	39.0	-0.9
MINNESOTA	35.3	38.8	3.5
MISSISSIPPI	40.4	42.2	1.9
MISSOURI	36.6	39.6	3.0
MONTANA	33.2	33.1	-0.2
NEBRASKA	35.9	37.2	1.4
NEVADA	31.6	31.2	-0.5
NEW HAMPSHIRE	32.8	34.0	1.2
NEW JERSEY	33.0	35.2	2.2
NEW MEXICO	34.6	34.4	-0.2
NEW YORK	32.5	33.2	0.7
NORTH CAROLINA	36.6	38.4	1.8
NORTH DAKOTA	38.7	38.9	0.2
OHIO	37.9	38.3	0.4
OKLAHOMA	32.6	36.6	4.1
OREGON	37.2	38.3	1.1
PENNSYLVANIA	36.7	39.7	3.0
RHODE ISLAND	33.4	35.3	1.9
SOUTH CAROLINA	37.7	39.5	1.8
SOUTH DAKOTA	36.2	34.8	-1.4
TENNESSEE	36.7	37.1	0.4
TEXAS	37.6	40.0	2.4
UTAH	31.2	33.6	2.4
VERMONT	34.6	32.2	-2.4
VIRGINIA	34.6	36.2	1.5
WASHINGTON	34.1	37.6	3.5
WEST VIRGINIA	41.2	43.0	1.8
WISCONSIN	37.1	39.2	2.0
WYOMING	32.7	33.0	0.2
PUERTO RICO	40.4	38.5	-1.9

* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

June 18, 1999

TABLE 17.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV: WOULD NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64

49

YEARS*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1707	9.4	0.9	(7.8, 11.1)
ALASKA	1741	10.2	1.1	(8.1, 12.3)
ARIZONA	1366	5.4	0.7	(4.0, 6.7)
ARKANSAS	2160	21.7	1.1	(19.6, 23.8)
COLORADO	1473	6.2	0.7	(4.8, 7.5)
CONNECTICUT	2392	9.7	0.8	(8.2, 11.2)
DELAWARE	1996	7.6	0.8	(6.0, 9.2)
DISTRICT OF CO	1159	7.6	0.9	(5.9, 9.4)
FLORIDA	3424	10.5	0.6	(9.3, 11.6)
GEORGIA	1942	12.4	0.8	(10.8, 14.1)
HAWAII	1724	8.6	0.9	(6.8, 10.4)
IDAHO	3908	12.1	0.6	(10.9, 13.3)
ILLINOIS	2368	10.8	0.7	(9.4, 12.1)
INDIANA	1930	10.5	0.8	(9.0, 12.0)
IOWA	2719	8.7	0.6	(7.6, 9.9)
KANSAS	2736	11.6	0.7	(10.2, 12.9)
KENTUCKY	2824	13.9	0.7	(12.5, 15.3)
LOUISIANA	1319	11.9	1.0	(10.0, 13.8)
MAINE	1215	8.3	0.9	(6.6, 10.0)
MARYLAND	2656	11.3	0.8	(9.7, 12.8)
MASSACHUSETTS	3868	7.5	0.5	(6.4, 8.5)
MICHIGAN	2066	8.5	0.7	(7.1, 9.9)
MINNESOTA	3458	41.5	1.0	(39.5, 43.4)
MISSISSIPPI	1750	14.8	0.9	(12.9, 16.6)
MISSOURI	2791	11.2	0.8	(9.7, 12.7)
MONTANA	1371	16.8	1.1	(14.6, 19.1)
NEBRASKA	1881	14.0	1.0	(12.0, 16.0)
NEVADA	1620	8.3	0.9	(6.6, 10.0)
NEW HAMPSHIRE	1153	7.9	1.0	(6.0, 9.7)
NEW JERSEY	1902	11.4	0.9	(9.6, 13.1)
NEW MEXICO	2892	9.1	0.6	(7.9, 10.2)
NEW YORK	1950	8.3	0.7	(6.9, 9.8)
NORTH CAROLINA	1814	11.6	1.0	(9.7, 13.5)
NORTH DAKOTA	1326	9.8	0.9	(8.1, 11.5)
OHIO	2229	9.6	0.9	(7.9, 11.4)
OKLAHOMA	1841	11.1	0.9	(9.3, 12.8)
OREGON	1490	10.7	0.9	(8.9, 12.5)
PENNSYLVANIA	2773	11.2	0.7	(9.7, 12.6)
RHODE ISLAND	2789	9.1	0.6	(7.9, 10.3)
SOUTH CAROLINA	2674	13.1	0.7	(11.7, 14.6)
SOUTH DAKOTA	1513	11.1	0.9	(9.4, 12.8)
TENNESSEE	2198	21.1	1.1	(19.0, 23.1)
TEXAS	4972	9.5	0.6	(8.3, 10.7)
UTAH	2327	11.2	0.9	(9.5, 12.9)
VERMONT	2348	7.0	1.0	(5.1, 8.9)
VIRGINIA	2822	9.1	0.8	(7.5, 10.8)
WASHINGTON	2943	8.7	0.7	(7.4, 10.0)
WEST VIRGINIA	1811	12.2	0.8	(10.6, 13.8)
WISCONSIN	1771	8.8	0.8	(7.2, 10.4)
WYOMING	1986	9.5	0.8	(8.0, 11.0)
PUERTO RICO	2210	12.0	0.8	(10.5, 13.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 10.5 RANGE = 5.4-41.5

* WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

June 18, 1999

TABLE 17.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HIV: WOULD NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64 YEARS*

50

-BY SEX

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES				FEMALES			
	SAMPLE	STANDARD	95% CONFIDENCE	*	SAMPLE	STANDARD	95% CONFIDENCE	*
	SIZE	PERCENT	ERROR	INTERVAL	SIZE	PERCENT	ERROR	INTERVAL
ALABAMA	625	10.3	1.3	(7.7, 12.8)	*	1082	8.6	1.1 (6.5, 10.7)
ALASKA	807	9.3	1.5	(6.3, 12.2)	*	934	11.2	1.5 (8.2, 14.2)
ARIZONA	609	6.5	1.2	(4.2, 8.8)	*	757	4.2	0.8 (2.6, 5.8)
ARKANSAS	865	22.9	1.6	(19.8, 26.0)	*	1295	20.5	1.4 (17.7, 23.2)
COLORADO	672	5.8	0.9	(3.9, 7.6)	*	801	6.6	1.0 (4.6, 8.5)
CONNECTICUT	1004	10.6	1.2	(8.3, 12.9)	*	1388	8.9	1.0 (7.0, 10.8)
DELAWARE	837	8.3	1.2	(5.9, 10.7)	*	1159	7.0	1.1 (4.8, 9.1)
DISTRICT OF CO	458	8.6	1.6	(5.6, 11.7)	*	701	6.7	0.9 (4.9, 8.5)
FLORIDA	1409	11.0	1.0	(9.1, 12.8)	*	2015	10.0	0.8 (8.5, 11.5)
GEORGIA	778	13.9	1.3	(11.3, 16.5)	*	1164	11.0	1.0 (9.1, 13.0)
HAWAII	776	8.8	1.4	(6.1, 11.5)	*	948	8.4	1.2 (6.0, 10.8)
IDAHO	1712	13.3	0.9	(11.5, 15.2)	*	2196	10.8	0.8 (9.4, 12.3)
ILLINOIS	1048	11.7	1.1	(9.7, 13.8)	*	1320	9.8	0.9 (8.0, 11.6)
INDIANA	818	10.0	1.1	(7.7, 12.2)	*	1112	11.0	1.0 (9.0, 13.0)
IOWA	1134	9.1	0.9	(7.3, 10.9)	*	1585	8.4	0.8 (6.8, 9.9)
KANSAS	1117	13.4	1.2	(11.2, 15.7)	*	1619	9.6	0.8 (8.1, 11.2)
KENTUCKY	1082	15.2	1.2	(13.0, 17.5)	*	1742	12.6	0.9 (10.9, 14.3)
LOUISIANA	540	11.3	1.5	(8.5, 14.2)	*	779	12.5	1.3 (10.0, 14.9)
MAINE	520	10.3	1.4	(7.6, 13.1)	*	695	6.2	1.0 (4.3, 8.2)
MARYLAND	1077	11.8	1.3	(9.3, 14.3)	*	1579	10.7	1.0 (8.8, 12.7)
MASSACHUSETTS	1659	8.7	0.9	(7.0, 10.4)	*	2209	6.3	0.7 (5.0, 7.5)
MICHIGAN	890	9.4	1.2	(7.1, 11.7)	*	1176	7.6	0.8 (6.0, 9.2)
MINNESOTA	1716	43.9	1.4	(41.2, 46.5)	*	1742	39.1	1.3 (36.5, 41.7)
MISSISSIPPI	706	16.3	1.5	(13.4, 19.2)	*	1044	13.3	1.2 (11.0, 15.6)
MISSOURI	1135	12.1	1.2	(9.7, 14.5)	*	1656	10.4	1.0 (8.4, 12.3)
MONTANA	613	17.9	1.8	(14.4, 21.3)	*	758	15.8	1.5 (12.9, 18.7)
NEBRASKA	754	15.4	1.8	(11.9, 19.0)	*	1127	12.6	1.1 (10.5, 14.6)
NEVADA	765	8.4	1.2	(6.0, 10.7)	*	855	8.2	1.3 (5.7, 10.6)
NEW HAMPSHIRE	479	9.5	1.7	(6.2, 12.7)	*	674	6.3	1.0 (4.3, 8.2)
NEW JERSEY	800	13.7	1.5	(10.8, 16.6)	*	1102	9.1	1.0 (7.1, 11.0)
NEW MEXICO	1301	9.9	0.9	(8.1, 11.6)	*	1591	8.4	0.8 (6.8, 9.9)
NEW YORK	822	9.1	1.2	(6.9, 11.4)	*	1128	7.6	0.9 (5.9, 9.3)
NORTH CAROLINA	705	13.7	1.6	(10.5, 16.9)	*	1109	9.6	1.0 (7.5, 11.6)
NORTH DAKOTA	622	9.8	1.3	(7.4, 12.3)	*	704	9.8	1.2 (7.5, 12.1)
OHIO	924	9.6	1.3	(7.0, 12.1)	*	1305	9.7	1.2 (7.3, 12.1)
OKLAHOMA	756	12.4	1.5	(9.5, 15.3)	*	1085	9.8	1.0 (7.8, 11.7)
OREGON	628	11.3	1.5	(8.4, 14.1)	*	862	10.2	1.1 (8.0, 12.4)
PENNSYLVANIA	1173	11.7	1.0	(9.7, 13.7)	*	1600	10.6	1.0 (8.6, 12.6)
RHODE ISLAND	1152	9.2	1.0	(7.3, 11.1)	*	1637	9.0	0.8 (7.4, 10.5)
SOUTH CAROLINA	1119	12.9	1.1	(10.8, 15.0)	*	1555	13.4	1.0 (11.4, 15.4)
SOUTH DAKOTA	676	11.3	1.3	(8.8, 13.7)	*	837	10.9	1.2 (8.6, 13.2)
TENNESSEE	801	22.1	1.7	(18.8, 25.5)	*	1397	20.0	1.2 (17.7, 22.4)
TEXAS	2091	11.0	1.1	(8.8, 13.1)	*	2881	8.1	0.5 (7.0, 9.1)
UTAH	1031	13.7	1.4	(10.9, 16.5)	*	1296	8.8	1.0 (6.9, 10.8)
VERMONT	1044	8.3	1.8	(4.9, 11.8)	*	1304	5.7	0.8 (4.1, 7.2)
VIRGINIA	1248	8.7	1.1	(6.5, 11.0)	*	1574	9.5	1.2 (7.1, 11.9)
WASHINGTON	1336	9.1	1.1	(7.0, 11.2)	*	1607	8.3	0.8 (6.9, 9.8)
WEST VIRGINIA	737	12.9	1.3	(10.3, 15.5)	*	1074	11.5	1.0 (9.6, 13.5)
WISCONSIN	804	10.3	1.4	(7.7, 13.0)	*	967	7.3	1.0 (5.4, 9.2)
WYOMING	868	10.3	1.2	(7.9, 12.8)	*	1118	8.7	0.9 (6.9, 10.5)
PUERTO RICO	863	11.7	1.2	(9.3, 14.0)	*	1347	12.3	1.0 (10.4, 14.3)

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 51
 MEDIAN = 11.0
 RANGE = 5.8-43.9

NUMBER OF PARTICIPANTS = 51
 MEDIAN = 9.6
 RANGE = 4.2-39.1

* WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

June 18, 1999

TABLE 17.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA

51

PREVALENCE DIFFERENCES FROM 1997 TO 1998

HIV: WOULD NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64 YEARS*

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	8.4	9.4	1.0
ALASKA	10.2	10.2	0.0
ARIZONA	5.0	5.4	0.3
ARKANSAS	16.3	21.7	5.4
COLORADO	6.8	6.2	-0.7
CONNECTICUT	6.8	9.7	2.9
DELAWARE	7.2	7.6	0.5
DISTRICT OF CO	9.3	7.6	-1.7
FLORIDA	10.7	10.5	-0.2
GEORGIA	9.4	12.4	3.1
HAWAII	10.1	8.6	-1.5
IDAHO	11.5	12.1	0.6
ILLINOIS	9.3	10.8	1.5
INDIANA	9.7	10.5	0.8
IOWA	8.4	8.7	0.3
KANSAS	13.1	11.6	-1.5
KENTUCKY	10.5	13.9	3.4
LOUISIANA	11.5	11.9	0.4
MAINE	7.7	8.3	0.5
MARYLAND	13.3	11.3	-2.1
MASSACHUSETTS	4.0	7.5	3.5
MICHIGAN	7.1	8.5	1.4
MINNESOTA	21.6	41.5	19.9
MISSISSIPPI	13.9	14.8	0.9
MISSOURI	10.9	11.2	0.4
MONTANA	16.0	16.8	0.8
NEBRASKA	11.6	14.0	2.3
NEVADA	8.2	8.3	0.1
NEW HAMPSHIRE	5.8	7.9	2.1
NEW JERSEY	7.4	11.4	3.9
NEW MEXICO	9.2	9.1	-0.1
NEW YORK	10.9	8.3	-2.6
NORTH CAROLINA	11.3	11.6	0.3
NORTH DAKOTA	10.3	9.8	-0.4
OHIO	9.9	9.6	-0.3
OKLAHOMA	10.7	11.1	0.4
OREGON	8.4	10.7	2.3
PENNSYLVANIA	8.7	11.2	2.5
RHODE ISLAND	8.1	9.1	1.0
SOUTH CAROLINA	17.5	13.1	-4.4
SOUTH DAKOTA	13.2	11.1	-2.1
TENNESSEE	20.2	21.1	0.9
TEXAS	13.9	9.5	-4.4
UTAH	11.6	11.2	-0.4
VERMONT	5.7	7.0	1.3
VIRGINIA	10.4	9.1	-1.3
WASHINGTON	7.6	8.7	1.2
WEST VIRGINIA	12.2	12.2	0.0
WISCONSIN	6.8	8.8	2.1
WYOMING	6.5	9.5	3.1
PUERTO RICO	12.1	12.0	-0.1

* WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

June 18, 1999

TABLE 18.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING HIV MEDIUM/HIGH, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

52

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1703	6.1	0.7	(4.8, 7.4)
ALASKA	1761	5.6	0.8	(4.0, 7.2)
ARIZONA	1376	1.9	0.5	(1.0, 2.8)
ARKANSAS	2181	5.5	0.7	(4.2, 6.9)
COLORADO	1486	6.2	0.8	(4.7, 7.7)
CONNECTICUT	2400	6.7	0.7	(5.4, 8.1)
DELAWARE	2009	4.7	0.9	(3.0, 6.4)
DISTRICT OF CO	1148	8.5	1.1	(6.4, 10.6)
FLORIDA	3426	6.1	0.5	(5.2, 7.0)
GEORGIA	1944	5.7	0.6	(4.5, 6.9)
HAWAII	1730	7.7	0.9	(6.0, 9.4)
IDAHO	3958	6.9	0.5	(6.0, 7.9)
ILLINOIS	2369	6.7	0.6	(5.5, 7.8)
INDIANA	1946	6.5	0.6	(5.3, 7.7)
IOWA	2738	5.4	0.5	(4.4, 6.4)
KANSAS	2772	5.0	0.5	(4.0, 6.0)
KENTUCKY	2844	5.2	0.5	(4.2, 6.2)
LOUISIANA	1317	7.3	0.9	(5.6, 8.9)
MAINE	1234	5.8	0.9	(4.1, 7.5)
MARYLAND	2668	6.0	0.6	(4.8, 7.2)
MASSACHUSETTS	3880	7.3	0.6	(6.1, 8.5)
MICHIGAN	2072	4.1	0.5	(3.2, 5.0)
MINNESOTA	3479	9.3	0.6	(8.2, 10.4)
MISSISSIPPI	1749	7.8	0.8	(6.2, 9.3)
MISSOURI	2825	6.5	0.6	(5.3, 7.7)
MONTANA	1388	3.8	0.5	(2.8, 4.9)
NEBRASKA	1914	7.7	0.7	(6.3, 9.0)
NEVADA	1624	6.5	0.9	(4.9, 8.2)
NEW HAMPSHIRE	1160	4.2	0.7	(2.9, 5.5)
NEW JERSEY	1916	7.4	0.7	(6.0, 8.8)
NEW MEXICO	2898	5.5	0.5	(4.5, 6.4)
NEW YORK	1946	7.3	0.7	(5.9, 8.8)
NORTH CAROLINA	1811	6.6	0.7	(5.2, 8.1)
NORTH DAKOTA	1341	4.7	0.6	(3.4, 5.9)
OHIO	2239	6.0	0.7	(4.6, 7.4)
OKLAHOMA	1869	6.2	0.7	(4.9, 7.5)
OREGON	1498	5.9	0.7	(4.5, 7.4)
PENNSYLVANIA	2790	6.6	0.6	(5.5, 7.6)
RHODE ISLAND	2800	6.8	0.6	(5.6, 7.9)
SOUTH CAROLINA	2661	6.8	0.6	(5.5, 8.0)
SOUTH DAKOTA	1524	11.0	1.0	(9.0, 13.0)
TENNESSEE	2269	6.7	0.6	(5.5, 8.0)
TEXAS	5026	7.0	0.5	(6.0, 7.9)
UTAH	2379	4.8	0.6	(3.7, 5.9)
VERMONT	2360	5.3	0.6	(4.2, 6.3)
VIRGINIA	2818	5.9	0.5	(4.9, 7.0)
WASHINGTON	2986	3.8	0.4	(3.0, 4.7)
WEST VIRGINIA	1815	11.4	0.9	(9.7, 13.1)
WISCONSIN	1777	5.6	0.6	(4.3, 6.8)
WYOMING	2001	6.3	0.6	(5.1, 7.5)
PUERTO RICO	2206	9.1	0.7	(7.7, 10.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 6.2 RANGE = 1.9-11.4

* CHANCES OF GETTING HIV MEDIUM/HIGH

June 18, 1999

TABLE 18.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING HIV MEDIUM/HIGH, AGE 18-64 YEARS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

53

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	624	6.8	1.1	(4.7, 8.9)	*	1079	5.4	0.8	(3.9, 6.9)	*
ALASKA	817	5.4	1.1	(3.4, 7.5)	*	944	5.8	1.3	(3.3, 8.2)	*
ARIZONA	614	2.4	0.7	(1.0, 3.7)	*	762	1.5	0.6	(0.3, 2.7)	*
ARKANSAS	870	5.4	1.1	(3.3, 7.5)	*	1311	5.6	0.9	(3.9, 7.4)	*
COLORADO	682	6.8	1.3	(4.4, 9.3)	*	804	5.5	0.9	(3.7, 7.3)	*
CONNECTICUT	1005	7.0	1.0	(4.9, 9.0)	*	1395	6.5	0.9	(4.7, 8.3)	*
DELAWARE	844	5.9	1.6	(2.8, 9.0)	*	1165	3.5	0.7	(2.1, 4.9)	*
DISTRICT OF CO	452	12.0	2.0	(8.2, 15.8)	*	696	5.3	0.9	(3.6, 7.0)	*
FLORIDA	1408	6.7	0.8	(5.2, 8.2)	*	2018	5.5	0.6	(4.4, 6.6)	*
GEORGIA	781	6.4	1.0	(4.5, 8.4)	*	1163	5.0	0.7	(3.6, 6.4)	*
HAWAII	776	8.7	1.4	(6.0, 11.5)	*	954	6.6	1.0	(4.7, 8.6)	*
IDAHO	1739	7.4	0.7	(5.9, 8.8)	*	2219	6.5	0.6	(5.3, 7.7)	*
ILLINOIS	1047	7.4	1.0	(5.5, 9.3)	*	1322	6.0	0.7	(4.6, 7.4)	*
INDIANA	832	6.9	1.0	(5.0, 8.9)	*	1114	6.1	0.8	(4.6, 7.6)	*
IOWA	1146	5.7	0.8	(4.1, 7.2)	*	1592	5.2	0.6	(3.9, 6.4)	*
KANSAS	1133	6.3	0.8	(4.6, 7.9)	*	1639	3.7	0.6	(2.7, 4.8)	*
KENTUCKY	1087	5.5	0.8	(4.0, 7.0)	*	1757	4.9	0.6	(3.6, 6.1)	*
LOUISIANA	544	9.2	1.4	(6.6, 11.9)	*	773	5.4	1.0	(3.5, 7.3)	*
MAINE	530	5.6	1.2	(3.3, 8.0)	*	704	5.9	1.3	(3.4, 8.5)	*
MARYLAND	1089	6.1	0.9	(4.2, 7.9)	*	1579	6.0	0.8	(4.4, 7.5)	*
MASSACHUSETTS	1659	9.2	1.1	(7.1, 11.3)	*	2221	5.5	0.6	(4.3, 6.6)	*
MICHIGAN	893	3.9	0.7	(2.6, 5.2)	*	1179	4.2	0.6	(3.0, 5.4)	*
MINNESOTA	1724	10.9	0.9	(9.2, 12.5)	*	1755	7.8	0.7	(6.4, 9.1)	*
MISSISSIPPI	704	7.7	1.3	(5.2, 10.2)	*	1045	7.8	1.0	(5.9, 9.8)	*
MISSOURI	1151	7.7	1.1	(5.6, 9.9)	*	1674	5.3	0.7	(3.9, 6.6)	*
MONTANA	622	3.8	0.8	(2.1, 5.4)	*	766	3.9	0.7	(2.6, 5.2)	*
NEBRASKA	769	8.5	1.1	(6.5, 10.6)	*	1145	6.9	0.9	(5.1, 8.6)	*
NEVADA	763	7.6	1.3	(5.1, 10.0)	*	861	5.4	1.1	(3.2, 7.7)	*
NEW HAMPSHIRE	481	4.2	1.0	(2.2, 6.2)	*	679	4.3	0.9	(2.6, 6.0)	*
NEW JERSEY	808	8.4	1.1	(6.2, 10.6)	*	1108	6.4	0.9	(4.6, 8.1)	*
NEW MEXICO	1301	6.0	0.7	(4.6, 7.5)	*	1597	4.9	0.7	(3.5, 6.3)	*
NEW YORK	821	8.4	1.2	(6.0, 10.7)	*	1125	6.3	0.8	(4.7, 8.0)	*
NORTH CAROLINA	703	6.6	1.1	(4.5, 8.6)	*	1108	6.7	1.0	(4.7, 8.7)	*
NORTH DAKOTA	630	4.0	0.9	(2.2, 5.7)	*	711	5.4	0.9	(3.6, 7.2)	*
OHIO	929	5.3	1.1	(3.2, 7.4)	*	1310	6.7	1.0	(4.8, 8.6)	*
OKLAHOMA	770	7.1	1.1	(4.9, 9.2)	*	1099	5.4	0.8	(3.9, 7.0)	*
OREGON	635	7.3	1.2	(4.9, 9.6)	*	863	4.6	0.8	(3.0, 6.2)	*
PENNSYLVANIA	1181	7.3	0.9	(5.6, 9.0)	*	1609	5.8	0.7	(4.4, 7.2)	*
RHODE ISLAND	1167	8.3	1.0	(6.4, 10.2)	*	1633	5.3	0.7	(4.0, 6.6)	*
SOUTH CAROLINA	1110	8.0	1.1	(5.9, 10.1)	*	1551	5.6	0.7	(4.2, 6.9)	*
SOUTH DAKOTA	680	11.2	1.4	(8.5, 13.9)	*	844	10.8	1.6	(7.8, 13.9)	*
TENNESSEE	828	8.2	1.1	(6.0, 10.5)	*	1441	5.3	0.7	(4.0, 6.6)	*
TEXAS	2112	8.5	0.8	(6.9, 10.1)	*	2914	5.4	0.5	(4.5, 6.3)	*
UTAH	1055	5.6	1.0	(3.7, 7.5)	*	1324	3.9	0.6	(2.7, 5.2)	*
VERMONT	1047	5.7	0.8	(4.2, 7.2)	*	1313	4.8	0.8	(3.4, 6.3)	*
VIRGINIA	1244	7.5	0.9	(5.6, 9.3)	*	1574	4.4	0.5	(3.4, 5.5)	*
WASHINGTON	1356	3.3	0.5	(2.3, 4.3)	*	1630	4.4	0.7	(3.0, 5.8)	*
WEST VIRGINIA	737	10.4	1.3	(7.8, 12.9)	*	1078	12.3	1.1	(10.2, 14.4)	*
WISCONSIN	810	5.6	1.0	(3.7, 7.5)	*	967	5.6	0.9	(3.9, 7.2)	*
WYOMING	876	6.4	0.9	(4.7, 8.1)	*	1125	6.2	0.9	(4.6, 7.9)	*
PUERTO RICO	859	10.2	1.1	(7.9, 12.4)	*	1347	8.0	0.8	(6.4, 9.7)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 51
MEDIAN = 6.9
RANGE = 2.4-12.0

NUMBER OF PARTICIPANTS = 51
MEDIAN = 5.5
RANGE = 1.5-12.3

* CHANCES OF GETTING HIV MEDIUM/HIGH

June 18, 1999

TABLE 18.3 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 CHANCE OF GETTING HIV MEDIUM/HIGH, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

54

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	5.8	6.1	0.2
ALASKA	5.7	5.6	-0.1
ARIZONA	3.3	1.9	-1.4
ARKANSAS	7.2	5.5	-1.7
COLORADO	4.6	6.2	1.6
CONNECTICUT	6.1	6.7	0.7
DELAWARE	7.0	4.7	-2.3
DISTRICT OF CO	7.2	8.5	1.3
FLORIDA	7.9	6.1	-1.8
GEORGIA	4.9	5.7	0.8
HAWAII	7.0	7.7	0.7
IDAHO	5.2	6.9	1.8
ILLINOIS	7.6	6.7	-0.9
INDIANA	6.1	6.5	0.4
IOWA	5.5	5.4	-0.1
KANSAS	8.7	5.0	-3.7
KENTUCKY	5.8	5.2	-0.6
LOUISIANA	4.9	7.3	2.4
MAINE	4.9	5.8	0.9
MARYLAND	5.0	6.0	1.0
MASSACHUSETTS	6.1	7.3	1.2
MICHIGAN	5.4	4.1	-1.3
MINNESOTA	5.7	9.3	3.6
MISSISSIPPI	6.8	7.8	1.0
MISSOURI	8.0	6.5	-1.5
MONTANA	4.3	3.8	-0.5
NEBRASKA	8.3	7.7	-0.6
NEVADA	6.3	6.5	0.2
NEW HAMPSHIRE	4.2	4.2	0.0
NEW JERSEY	6.4	7.4	1.0
NEW MEXICO	7.9	5.5	-2.4
NEW YORK	7.3	7.3	-0.0
NORTH CAROLINA	6.7	6.6	-0.1
NORTH DAKOTA	4.9	4.7	-0.2
OHIO	6.2	6.0	-0.2
OKLAHOMA	4.9	6.2	1.3
OREGON	5.5	5.9	0.4
PENNSYLVANIA	6.8	6.6	-0.3
RHODE ISLAND	7.3	6.8	-0.5
SOUTH CAROLINA	5.5	6.8	1.2
SOUTH DAKOTA	8.9	11.0	2.1
TENNESSEE	7.0	6.7	-0.3
TEXAS	12.1	7.0	-5.1
UTAH	5.2	4.8	-0.4
VERMONT	6.6	5.3	-1.3
VIRGINIA	5.0	5.9	0.9
WASHINGTON	4.5	3.8	-0.7
WEST VIRGINIA	9.1	11.4	2.2
WISCONSIN	5.6	5.6	-0.0
WYOMING	7.9	6.3	-1.5
PUERTO RICO	10.2	9.1	-1.1

* CHANCES OF GETTING HIV MEDIUM/HIGH

June 18, 1999

TABLE 19.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAVE NOT BEEN TESTED FOR HIV, AGE 18-64 YEARS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

55

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	264	56.4	3.5	(49.5, 63.2)
ALASKA	202	58.8	4.8	(49.4, 68.3)
ARIZONA	155	69.4	6.5	(56.8, 82.1)
ARKANSAS	255	61.2	3.8	(53.8, 68.6)
COLORADO	210	68.4	3.5	(61.7, 75.2)
CONNECTICUT	260	59.1	4.1	(51.1, 67.2)
DELAWARE	254	57.0	4.3	(48.6, 65.4)
DISTRICT OF CO	207	58.8	4.5	(49.9, 67.7)
FLORIDA	559	56.4	2.4	(51.7, 61.2)
GEORGIA	349	51.2	3.5	(44.3, 58.1)
HAWAII	209	45.4	4.6	(36.5, 54.3)
IDAHO	413	64.5	2.8	(59.0, 70.1)
ILLINOIS	262	66.2	3.4	(59.4, 72.9)
INDIANA	171	61.6	4.2	(53.4, 69.8)
IOWA	250	66.1	3.6	(59.1, 73.1)
KANSAS	234	53.7	3.8	(46.3, 61.2)
KENTUCKY	244	61.8	3.6	(54.8, 68.7)
LOUISIANA	214	53.3	4.2	(45.2, 61.5)
MAINE	121	62.2	5.0	(52.3, 72.0)
MARYLAND	411	57.1	3.3	(50.5, 63.6)
MASSACHUSETTS	515	64.7	2.7	(59.3, 70.1)
MICHIGAN	286	63.2	3.2	(56.9, 69.4)
MINNESOTA	243	61.1	3.3	(54.6, 67.6)
MISSISSIPPI	225	49.8	3.7	(42.4, 57.1)
MISSOURI	334	62.5	3.7	(55.2, 69.8)
MONTANA	124	59.9	5.0	(50.2, 69.6)
NEBRASKA	150	63.1	4.3	(54.7, 71.6)
NEVADA	249	59.5	4.2	(51.2, 67.8)
NEW HAMPSHIRE	145	61.2	4.5	(52.4, 70.0)
NEW JERSEY	228	63.2	3.9	(55.4, 70.9)
NEW MEXICO	383	63.0	3.1	(56.9, 69.1)
NEW YORK	223	57.2	4.0	(49.4, 64.9)
NORTH CAROLINA	255	53.6	3.9	(46.1, 61.2)
NORTH DAKOTA	110	70.7	4.7	(61.5, 79.8)
OHIO	198	66.4	4.6	(57.4, 75.4)
OKLAHOMA	176	58.4	4.3	(50.1, 66.8)
OREGON	187	61.8	4.0	(53.9, 69.7)
PENNSYLVANIA	341	56.8	3.1	(50.6, 62.9)
RHODE ISLAND	365	62.7	3.0	(56.9, 68.5)
SOUTH CAROLINA	377	58.7	3.1	(52.6, 64.7)
SOUTH DAKOTA	122	56.7	5.0	(46.8, 66.5)
TENNESSEE	312	55.7	3.3	(49.2, 62.1)
TEXAS	703	59.4	2.5	(54.5, 64.3)
UTAH	256	69.0	4.0	(61.2, 76.8)
VERMONT	274	63.9	3.6	(56.9, 70.8)
VIRGINIA	439	56.4	3.4	(49.8, 63.0)
WASHINGTON	343	63.6	2.9	(57.8, 69.3)
WEST VIRGINIA	127	51.9	5.0	(42.2, 61.6)
WISCONSIN	191	63.6	4.7	(54.3, 72.9)
WYOMING	204	67.3	4.0	(59.4, 75.2)
PUERTO RICO	248	56.5	3.6	(49.6, 63.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 61.1 RANGE = 45.4-70.7

* HAVE NOT BEEN TESTED FOR HIV, IN THE PAST 12 MONTHS

June 18, 1999

TABLE 19.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAVE NOT BEEN TESTED FOR HIV, AGE 18-64 YEARS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

56

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	116	53.2	5.0	(43.3, 63.1)	*	148	60.6	4.8	(51.2, 69.9)	*
ALASKA	115	57.1	6.5	(44.3, 69.9)	*	87	62.0	6.8	(48.6, 75.4)	*
ARIZONA	81	72.9	7.6	(58.1, 87.7)	*	74	64.6	10.7	(43.5, 85.6)	*
ARKANSAS	109	60.8	5.7	(49.7, 71.9)	*	146	61.7	4.8	(52.3, 71.0)	*
COLORADO	107	62.3	5.0	(52.5, 72.2)	*	103	76.4	4.5	(67.6, 85.1)	*
CONNECTICUT	144	59.4	5.7	(48.2, 70.5)	*	116	58.7	5.4	(48.2, 69.2)	*
DELAWARE	131	51.5	5.9	(40.0, 63.0)	*	123	64.2	5.8	(52.9, 75.6)	*
DISTRICT OF CO	102	60.0	6.6	(47.0, 72.9)	*	105	57.1	5.6	(46.2, 68.1)	*
FLORIDA	276	59.0	3.5	(52.2, 65.8)	*	283	52.9	3.3	(46.5, 59.3)	*
GEORGIA	183	45.2	4.8	(35.9, 54.5)	*	166	61.7	4.4	(53.2, 70.2)	*
HAWAII	122	42.3	5.8	(30.8, 53.7)	*	87	52.5	6.6	(39.6, 65.4)	*
IDAHO	213	63.7	3.9	(56.2, 71.3)	*	200	65.7	4.1	(57.8, 73.7)	*
ILLINOIS	142	61.7	4.8	(52.4, 71.1)	*	120	72.6	4.6	(63.6, 81.7)	*
INDIANA	85	64.6	5.9	(53.0, 76.2)	*	86	57.8	5.7	(46.5, 69.0)	*
IOWA	131	69.0	4.8	(59.6, 78.5)	*	119	61.9	5.2	(51.7, 72.0)	*
KANSAS	110	53.1	5.5	(42.4, 63.9)	*	124	54.4	5.2	(44.3, 64.6)	*
KENTUCKY	108	61.0	5.1	(51.0, 71.0)	*	136	62.7	5.0	(53.0, 72.4)	*
LOUISIANA	108	55.6	5.7	(44.4, 66.8)	*	106	50.0	5.4	(39.4, 60.7)	*
MAINE	64	60.7	6.9	(47.1, 74.3)	*	57	64.3	7.0	(50.5, 78.0)	*
MARYLAND	185	59.5	4.9	(49.9, 69.1)	*	226	54.0	4.4	(45.5, 62.5)	*
MASSACHUSETTS	298	65.3	3.6	(58.3, 72.3)	*	217	63.5	3.9	(55.9, 71.2)	*
MICHIGAN	138	59.4	4.7	(50.3, 68.6)	*	148	67.4	4.2	(59.2, 75.6)	*
MINNESOTA	145	62.8	4.3	(54.4, 71.3)	*	98	58.5	5.6	(47.6, 69.4)	*
MISSISSIPPI	110	49.5	5.2	(39.2, 59.8)	*	115	50.1	5.2	(39.9, 60.2)	*
MISSOURI	169	62.0	5.4	(51.4, 72.5)	*	165	63.4	4.6	(54.3, 72.4)	*
MONTANA	74	58.8	6.2	(46.5, 71.0)	*	50	61.6	7.9	(46.1, 77.1)	*
NEBRASKA	77	63.3	6.0	(51.6, 75.0)	*	73	62.9	6.0	(51.3, 74.6)	*
NEVADA	121	55.3	6.1	(43.4, 67.2)	*	128	65.1	5.6	(54.2, 75.9)	*
NEW HAMPSHIRE	71	57.7	6.5	(44.9, 70.5)	*	74	66.3	6.1	(54.3, 78.2)	*
NEW JERSEY	124	62.5	5.3	(52.1, 72.9)	*	104	64.3	5.7	(53.1, 75.5)	*
NEW MEXICO	201	60.1	4.4	(51.6, 68.7)	*	182	67.1	4.2	(58.8, 75.4)	*
NEW YORK	116	56.0	5.6	(45.1, 67.0)	*	107	58.8	5.4	(48.2, 69.3)	*
NORTH CAROLINA	123	50.8	5.4	(40.2, 61.4)	*	132	57.4	5.4	(46.9, 67.9)	*
NORTH DAKOTA	58	67.8	6.5	(55.0, 80.5)	*	52	74.6	6.2	(62.4, 86.9)	*
OHIO	100	73.6	5.7	(62.5, 84.8)	*	98	56.1	7.1	(42.2, 70.1)	*
OKLAHOMA	82	60.0	6.3	(47.6, 72.4)	*	94	56.5	5.5	(45.8, 67.2)	*
OREGON	87	63.0	5.8	(51.7, 74.3)	*	100	60.4	5.5	(49.7, 71.2)	*
PENNSYLVANIA	166	54.6	4.5	(45.8, 63.4)	*	175	59.7	4.1	(51.6, 67.8)	*
RHODE ISLAND	177	57.9	4.2	(49.6, 66.1)	*	188	69.3	3.8	(62.0, 76.7)	*
SOUTH CAROLINA	202	60.5	4.2	(52.3, 68.7)	*	175	55.6	4.3	(47.2, 64.0)	*
SOUTH DAKOTA	66	56.7	6.6	(43.9, 69.6)	*	56	56.6	7.7	(41.4, 71.7)	*
TENNESSEE	146	49.0	4.6	(40.1, 58.0)	*	166	65.6	4.1	(57.5, 73.7)	*
TEXAS	339	61.2	3.6	(54.2, 68.2)	*	364	57.0	3.3	(50.5, 63.5)	*
UTAH	142	65.7	5.5	(55.0, 76.5)	*	114	74.2	5.3	(63.8, 84.5)	*
VERMONT	151	59.8	4.8	(50.4, 69.2)	*	123	69.4	5.1	(59.4, 79.4)	*
VIRGINIA	221	53.7	4.7	(44.5, 62.8)	*	218	60.2	4.7	(51.0, 69.3)	*
WASHINGTON	174	60.1	4.2	(51.9, 68.2)	*	169	67.9	4.1	(59.9, 75.9)	*
WEST VIRGINIA	70	57.4	6.7	(44.4, 70.5)	*	57	43.4	7.1	(29.6, 57.3)	*
WISCONSIN	91	63.2	5.8	(51.8, 74.6)	*	100	63.9	7.2	(49.8, 78.0)	*
WYOMING	112	65.4	5.7	(54.1, 76.6)	*	92	70.5	5.1	(60.5, 80.4)	*
PUERTO RICO	156	55.6	4.4	(47.0, 64.1)	*	92	59.1	5.9	(47.5, 70.6)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 51
MEDIAN = 60.0
RANGE = 42.3-73.6

NUMBER OF PARTICIPANTS = 51
MEDIAN = 61.7
RANGE = 43.4-76.4

June 18, 1999

* HAVE NOT BEEN TESTED FOR HIV, IN THE PAST 12 MONTHS

TABLE 20.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED MEAN AGE AT ONSET* AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

57

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	165	47.6	1.4	(45.0, 50.2)
ALASKA	74	45.9	3.6	(38.9, 52.9)
ARIZONA	49	46.7	3.1	(40.7, 52.7)
ARKANSAS	203	46.7	1.5	(43.8, 49.6)
CALIFORNIA	199	46.8	1.2	(44.5, 49.2)
COLORADO	85	52.0	1.7	(48.7, 55.3)
CONNECTICUT	151	48.3	1.7	(45.0, 51.6)
DELAWARE	141	50.0	2.0	(46.2, 53.8)
FLORIDA	292	50.6	1.0	(48.6, 52.5)
GEORGIA	127	48.8	1.7	(45.5, 52.1)
HAWAII	108	50.0	2.2	(45.7, 54.3)
IDAHO	203	49.6	1.4	(46.8, 52.3)
IOWA	187	49.8	1.4	(47.1, 52.6)
KANSAS	162	47.9	1.5	(45.0, 50.8)
KENTUCKY	219	48.8	1.4	(46.0, 51.6)
MAINE	70	49.2	2.3	(44.7, 53.6)
MASSACHUSETTS	196	46.4	1.9	(42.7, 50.2)
MICHIGAN	161	44.6	1.6	(41.6, 47.6)
MINNESOTA	185	49.2	1.6	(46.1, 52.2)
MISSISSIPPI	168	43.6	1.7	(40.3, 47.0)
MONTANA	67	50.4	2.2	(46.2, 54.6)
NEBRASKA	152	49.0	1.6	(45.8, 52.1)
NEVADA	84	47.8	2.5	(43.0, 52.7)
NEW HAMPSHIRE	58	50.2	2.7	(45.0, 55.4)
NEW JERSEY	122	53.1	1.2	(50.8, 55.5)
NEW MEXICO	181	47.4	1.4	(44.7, 50.1)
NORTH CAROLINA	140	48.8	1.7	(45.4, 52.2)
NORTH DAKOTA	75	52.7	2.2	(48.4, 57.1)
OHIO	184	47.0	1.8	(43.5, 50.5)
PENNSYLVANIA	193	46.9	1.5	(44.0, 49.9)
RHODE ISLAND	197	49.7	1.3	(47.3, 52.2)
TENNESSEE	161	46.0	1.2	(43.6, 48.3)
TEXAS	327	46.2	0.9	(44.4, 47.9)
UTAH	127	48.8	2.2	(44.6, 53.1)
VERMONT	132	49.7	1.6	(46.6, 52.8)
VIRGINIA	180	47.3	1.2	(45.1, 49.6)
WEST VIRGINIA	160	51.7	1.2	(49.3, 54.2)
WISCONSIN	101	48.6	2.3	(44.0, 53.2)
WYOMING	86	52.6	1.9	(48.8, 56.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 39 MEDIAN = 48.8 RANGE = 43.6-53.1

* MEAN AGE AT ONSET OF DIABETES

June 18, 1999

TABLE 20.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN AGE AT ONSET* AMONG RESPONDENTS TOLD BY DOCTOR THAT THEY HAVE DIABETES -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

58

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	53	49.1	2.0	(45.1, 53.0)	*	112	46.6	1.8	(43.1, 50.1)	*
ALASKA	39	44.7	5.1	(34.8, 54.6)	*	35	47.6	4.6	(38.5, 56.7)	*
ARIZONA	20	47.3	4.9	(37.6, 57.0)	*	29	46.1	3.4	(39.5, 52.7)	*
ARKANSAS	81	49.3	1.8	(45.7, 52.9)	*	122	44.2	2.1	(40.0, 48.3)	*
CALIFORNIA	91	47.2	1.7	(43.8, 50.7)	*	108	46.5	1.6	(43.4, 49.6)	*
COLORADO	37	52.2	2.4	(47.6, 56.8)	*	48	51.9	2.4	(47.2, 56.6)	*
CONNECTICUT	74	48.3	2.3	(43.8, 52.7)	*	77	48.4	2.5	(43.6, 53.3)	*
DELAWARE	59	49.5	3.2	(43.2, 55.7)	*	82	50.5	2.3	(45.9, 55.1)	*
FLORIDA	135	52.3	1.4	(49.7, 55.0)	*	157	48.5	1.3	(45.9, 51.1)	*
GEORGIA	52	47.9	2.6	(42.9, 52.9)	*	75	50.0	2.0	(46.1, 53.9)	*
HAWAII	51	49.4	3.7	(42.2, 56.6)	*	57	50.6	2.4	(46.0, 55.2)	*
IDAHO	92	50.8	1.9	(47.1, 54.5)	*	111	48.1	2.1	(44.0, 52.1)	*
IAOWA	74	50.2	1.9	(46.6, 53.9)	*	113	49.5	2.1	(45.4, 53.5)	*
KANSAS	70	46.5	2.2	(42.2, 50.8)	*	92	49.6	1.9	(45.8, 53.3)	*
KENTUCKY	80	47.5	2.2	(43.2, 51.8)	*	139	49.9	1.8	(46.4, 53.4)	*
MAINE	16	50.0	4.6	(41.1, 59.0)	*	54	48.8	2.6	(43.6, 53.9)	*
MASSACHUSETTS	89	45.2	2.9	(39.4, 50.9)	*	107	48.2	2.0	(44.3, 52.1)	*
MICHIGAN	56	49.1	2.1	(44.9, 53.2)	*	105	41.1	2.0	(37.1, 45.1)	*
MINNESOTA	79	51.5	2.2	(47.2, 55.8)	*	106	47.3	2.1	(43.1, 51.4)	*
MISSISSIPPI	60	43.1	3.1	(37.0, 49.3)	*	108	44.0	1.8	(40.5, 47.5)	*
MONTANA	25	51.2	2.8	(45.8, 56.6)	*	42	49.8	3.0	(43.9, 55.8)	*
NEBRASKA	64	48.0	2.3	(43.5, 52.4)	*	88	50.0	2.2	(45.7, 54.4)	*
NEVADA	42	49.3	3.2	(43.1, 55.5)	*	42	45.7	3.9	(38.0, 53.3)	*
NEW HAMPSHIRE	30	53.5	2.1	(49.5, 57.5)	*	28	46.4	4.8	(36.9, 55.8)	*
NEW JERSEY	56	53.1	1.7	(49.8, 56.4)	*	66	53.2	1.7	(49.9, 56.5)	*
NEW MEXICO	80	48.9	2.0	(44.9, 52.8)	*	101	45.9	1.9	(42.2, 49.6)	*
NORTH CAROLINA	47	52.8	2.4	(48.2, 57.4)	*	93	46.3	2.3	(41.7, 50.8)	*
NORTH DAKOTA	33	51.6	3.4	(45.0, 58.2)	*	42	53.8	3.0	(48.0, 59.6)	*
OHIO	68	43.7	2.9	(37.9, 49.4)	*	116	50.6	1.7	(47.3, 53.8)	*
PENNSYLVANIA	70	46.6	2.9	(40.9, 52.3)	*	123	47.2	1.5	(44.3, 50.0)	*
RHODE ISLAND	80	49.7	1.7	(46.3, 53.0)	*	117	49.8	1.8	(46.2, 53.3)	*
TENNESSEE	46	49.3	1.9	(45.5, 53.0)	*	115	43.4	1.6	(40.3, 46.5)	*
TEXAS	117	44.8	1.4	(42.0, 47.6)	*	210	47.2	1.1	(45.0, 49.4)	*
UTAH	59	52.6	2.8	(47.1, 58.2)	*	68	44.0	2.9	(38.3, 49.7)	*
VERMONT	58	47.2	2.3	(42.7, 51.6)	*	74	52.2	2.2	(47.9, 56.5)	*
VIRGINIA	80	48.4	1.6	(45.2, 51.6)	*	100	46.4	1.6	(43.2, 49.6)	*
WEST VIRGINIA	56	51.6	2.0	(47.7, 55.4)	*	104	51.9	1.6	(48.7, 55.0)	*
WISCONSIN	51	48.6	3.2	(42.3, 54.9)	*	50	48.6	3.4	(42.0, 55.2)	*
WYOMING	42	51.9	2.7	(46.6, 57.2)	*	44	53.6	2.6	(48.4, 58.8)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 39
MEDIAN = 49.3
RANGE = 43.1-53.5

NUMBER OF PARTICIPANTS = 39
MEDIAN = 48.4
RANGE = 41.1-53.8

June 18, 1999

* MEAN AGE AT ONSET OF DIABETES

TABLE 21.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NOW TAKING INSULIN AMONG RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	182	38.3	4.1	(30.2, 46.3)
ALASKA	75	28.2	6.7	(15.1, 41.3)
ARIZONA	55	17.8	6.6	(4.8, 30.8)
ARKANSAS	209	38.8	4.2	(30.6, 47.0)
CALIFORNIA	203	29.3	3.7	(22.1, 36.5)
COLORADO	87	34.0	5.4	(23.4, 44.6)
CONNECTICUT	164	32.1	4.8	(22.6, 41.5)
DELAWARE	144	25.4	4.5	(16.5, 34.2)
FLORIDA	302	32.2	3.1	(26.1, 38.3)
GEORGIA	140	26.1	4.1	(18.0, 34.1)
HAWAII	113	18.5	5.1	(8.6, 28.4)
IDAHO	208	34.0	4.0	(26.1, 41.8)
IOWA	195	24.4	3.6	(17.3, 31.4)
KANSAS	170	36.8	4.4	(28.2, 45.5)
KENTUCKY	231	31.8	3.4	(25.1, 38.5)
MAINE	73	37.8	6.2	(25.8, 49.9)
MASSACHUSETTS	212	32.5	4.4	(23.9, 41.1)
MICHIGAN	168	36.0	4.3	(27.6, 44.3)
MINNESOTA	189	35.9	3.8	(28.5, 43.3)
MISSISSIPPI	185	39.8	4.6	(30.8, 48.8)
MONTANA	70	37.6	6.6	(24.6, 50.5)
NEBRASKA	163	40.0	4.2	(31.7, 48.3)
NEVADA	85	26.9	6.3	(14.4, 39.3)
NEW HAMPSHIRE	61	30.7	7.8	(15.5, 46.0)
NEW JERSEY	128	28.7	4.2	(20.4, 37.0)
NEW MEXICO	186	36.5	3.9	(28.8, 44.2)
NORTH CAROLINA	152	35.6	4.5	(26.8, 44.3)
NORTH DAKOTA	78	37.5	6.3	(25.2, 49.7)
OHIO	195	28.1	4.8	(18.6, 37.5)
PENNSYLVANIA	202	38.7	3.9	(31.0, 46.4)
RHODE ISLAND	216	31.0	3.5	(24.2, 37.8)
TENNESSEE	171	42.1	4.6	(33.0, 51.2)
TEXAS	339	31.2	3.0	(25.4, 37.0)
UTAH	127	35.1	5.4	(24.5, 45.8)
VERMONT	144	34.5	4.5	(25.7, 43.2)
VIRGINIA	196	40.0	5.1	(30.1, 50.0)
WEST VIRGINIA	168	33.4	4.2	(25.2, 41.6)
WISCONSIN	103	29.8	5.4	(19.3, 40.3)
WYOMING	89	32.8	5.5	(22.1, 43.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 39 MEDIAN = 33.4 RANGE = 17.8-42.1
 June 18, 1999

TABLE 21.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE NOW TAKING INSULIN AMONG RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES-BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

60

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	59	35.8	7.0	(22.0, 49.5)	*	123	40.0	5.0	(30.2, 49.8)	*
ALASKA	39	24.6	8.1	(8.7, 40.4)	*	36	33.1	10.5	(12.6, 53.7)	*
ARIZONA	21	12.8	9.8	(-6.4, 32.0)	*	34	22.2	9.0	(4.6, 39.9)	*
ARKANSAS	85	35.2	5.6	(24.2, 46.2)	*	124	42.3	6.1	(30.4, 54.2)	*
CALIFORNIA	94	30.6	5.5	(19.9, 41.2)	*	109	28.1	4.9	(18.5, 37.7)	*
COLORADO	39	27.9	7.4	(13.4, 42.5)	*	48	39.7	7.6	(24.7, 54.6)	*
CONNECTICUT	78	35.3	7.1	(21.3, 49.2)	*	86	28.1	6.1	(16.2, 39.9)	*
DELAWARE	60	30.0	7.3	(15.8, 44.2)	*	84	21.5	5.6	(10.5, 32.5)	*
FLORIDA	138	27.9	4.2	(19.7, 36.1)	*	164	36.9	4.4	(28.3, 45.4)	*
GEORGIA	55	26.6	6.3	(14.4, 38.9)	*	85	25.4	5.0	(15.6, 35.1)	*
HAWAII	52	15.4	7.5	(0.8, 30.1)	*	61	21.5	6.9	(7.9, 35.0)	*
IDAHO	93	31.2	5.9	(19.7, 42.7)	*	115	37.2	5.4	(26.7, 47.7)	*
IAOWA	75	23.8	5.6	(12.8, 34.9)	*	120	24.8	4.6	(15.8, 33.9)	*
KANSAS	71	38.5	6.8	(25.0, 51.9)	*	99	35.0	5.3	(24.6, 45.3)	*
KENTUCKY	82	28.6	5.4	(18.0, 39.2)	*	149	34.5	4.3	(26.2, 42.9)	*
MAINE	17	38.2	12.5	(13.6, 62.7)	*	56	37.7	7.0	(24.0, 51.4)	*
MASSACHUSETTS	95	36.7	6.6	(23.7, 49.6)	*	117	26.7	4.9	(17.1, 36.4)	*
MICHIGAN	58	35.3	7.0	(21.6, 48.9)	*	110	36.5	5.3	(26.1, 46.9)	*
MINNESOTA	80	30.3	5.4	(19.7, 41.0)	*	109	40.4	5.2	(30.2, 50.6)	*
MISSISSIPPI	63	41.3	7.5	(26.5, 56.0)	*	122	38.7	5.7	(27.5, 49.9)	*
MONTANA	26	45.0	10.9	(23.6, 66.3)	*	44	32.4	8.0	(16.8, 48.0)	*
NEBRASKA	66	41.0	6.6	(28.1, 53.9)	*	97	39.0	5.4	(28.5, 49.5)	*
NEVADA	42	36.0	9.4	(17.7, 54.3)	*	43	13.5	4.9	(3.9, 23.0)	*
NEW HAMPSHIRE	30	32.8	11.5	(10.3, 55.4)	*	31	28.5	10.4	(8.1, 48.8)	*
NEW JERSEY	60	28.1	5.9	(16.4, 39.7)	*	68	29.5	6.1	(17.6, 41.3)	*
NEW MEXICO	82	36.8	5.8	(25.4, 48.1)	*	104	36.2	5.3	(25.8, 46.6)	*
NORTH CAROLINA	52	27.4	6.9	(13.9, 40.9)	*	100	40.9	5.7	(29.7, 52.1)	*
NORTH DAKOTA	34	37.0	9.9	(17.6, 56.4)	*	44	38.0	7.9	(22.4, 53.5)	*
OHIO	70	23.6	6.9	(10.0, 37.1)	*	125	33.0	6.5	(20.2, 45.8)	*
PENNSYLVANIA	73	43.4	6.5	(30.8, 56.1)	*	129	35.1	4.8	(25.7, 44.5)	*
RHODE ISLAND	87	29.3	5.3	(18.8, 39.7)	*	129	32.7	4.5	(23.9, 41.5)	*
TENNESSEE	47	43.8	8.6	(27.0, 60.6)	*	124	40.8	5.1	(30.8, 50.8)	*
TEXAS	118	26.3	4.5	(17.4, 35.2)	*	221	34.8	3.9	(27.2, 42.4)	*
UTAH	59	29.7	7.8	(14.5, 45.0)	*	68	42.1	7.7	(26.9, 57.2)	*
VERMONT	60	40.0	7.1	(26.1, 53.8)	*	84	29.8	5.6	(18.9, 40.8)	*
VIRGINIA	83	31.7	5.8	(20.4, 43.1)	*	113	46.6	7.2	(32.4, 60.8)	*
WEST VIRGINIA	59	24.2	6.2	(12.1, 36.3)	*	109	40.5	5.3	(30.1, 50.8)	*
WISCONSIN	52	25.0	7.0	(11.3, 38.7)	*	51	35.3	8.1	(19.5, 51.1)	*
WYOMING	43	31.1	7.5	(16.4, 45.7)	*	46	35.1	7.8	(19.8, 50.4)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 39
MEDIAN = 31.1

NUMBER OF PARTICIPANTS = 39
MEDIAN = 35.1

June 18, 1999

TABLE 22.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN PROFESSIONAL VISITS IN PAST YEAR AMONG RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES*

61

DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	158	4.3	0.4	(3.6, 5.0)
ALASKA	69	4.6	0.6	(3.4, 5.8)
ARIZONA	49	3.7	0.5	(2.7, 4.8)
ARKANSAS	200	6.4	0.9	(4.6, 8.1)
CALIFORNIA	197	4.7	0.4	(3.9, 5.5)
COLORADO	83	3.8	0.4	(3.1, 4.5)
CONNECTICUT	152	3.9	0.3	(3.3, 4.6)
DELAWARE	141	5.3	0.8	(3.7, 6.8)
FLORIDA	294	4.7	0.3	(4.1, 5.4)
GEORGIA	126	4.0	0.4	(3.3, 4.7)
HAWAII	109	7.0	1.0	(5.0, 9.0)
IDAHO	201	3.7	0.3	(3.1, 4.3)
IOWA	189	4.1	0.5	(3.2, 4.9)
KANSAS	160	5.0	0.7	(3.7, 6.3)
KENTUCKY	221	5.1	0.4	(4.4, 5.8)
MAINE	70	3.9	0.5	(3.0, 4.8)
MASSACHUSETTS	204	3.6	0.3	(3.0, 4.1)
MICHIGAN	159	4.4	0.5	(3.4, 5.3)
MINNESOTA	189	3.5	0.3	(2.9, 4.0)
MISSISSIPPI	167	4.4	0.5	(3.5, 5.3)
MONTANA	65	4.6	0.7	(3.1, 6.0)
NEBRASKA	158	4.7	0.6	(3.6, 5.7)
NEVADA	83	4.0	0.8	(2.3, 5.6)
NEW HAMPSHIRE	56	3.8	0.6	(2.6, 5.0)
NEW JERSEY	126	5.2	0.6	(4.0, 6.4)
NEW MEXICO	180	5.6	0.6	(4.4, 6.9)
NORTH CAROLINA	140	4.5	0.4	(3.7, 5.3)
NORTH DAKOTA	77	4.1	0.4	(3.4, 4.9)
OHIO	188	4.4	0.4	(3.6, 5.2)
PENNSYLVANIA	199	4.6	0.3	(4.1, 5.2)
RHODE ISLAND	208	3.8	0.3	(3.3, 4.3)
TENNESSEE	161	4.5	0.4	(3.7, 5.3)
TEXAS	329	4.8	0.3	(4.2, 5.4)
UTAH	125	4.2	0.5	(3.2, 5.1)
VERMONT	133	4.1	0.4	(3.3, 4.8)
VIRGINIA	181	3.9	0.3	(3.3, 4.5)
WEST VIRGINIA	167	4.8	0.3	(4.1, 5.5)
WISCONSIN	104	3.8	0.6	(2.7, 5.0)
WYOMING	84	3.2	0.4	(2.4, 4.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 39 MEDIAN = 4.4 RANGE = 3.2- 7.0

* MEAN TIMES SEEN HEALTH PROFESSIONALS FOR DIABETES IN PAST YEAR

June 18, 1999

TABLE 22.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED MEAN PROFESSIONAL VISITS IN PAST YEAR AMONG RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

62

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALABAMA	52	4.2	0.6	(3.0, 5.3)	*	106	4.5	0.5	(3.6, 5.4)	*
ALASKA	36	4.1	0.7	(2.7, 5.6)	*	33	5.3	1.1	(3.2, 7.4)	*
ARIZONA	20	4.5	0.8	(2.9, 6.0)	*	29	2.9	0.5	(2.0, 3.8)	*
ARKANSAS	80	7.0	1.6	(3.9, 10.1)	*	120	5.8	0.8	(4.1, 7.4)	*
CALIFORNIA	93	4.6	0.5	(3.6, 5.6)	*	104	4.8	0.6	(3.7, 6.0)	*
COLORADO	38	3.6	0.4	(2.8, 4.4)	*	45	4.0	0.6	(2.9, 5.2)	*
CONNECTICUT	73	3.6	0.5	(2.7, 4.5)	*	79	4.4	0.5	(3.4, 5.4)	*
DELAWARE	59	5.6	1.5	(2.6, 8.5)	*	82	5.0	0.7	(3.6, 6.5)	*
FLORIDA	135	4.8	0.5	(3.9, 5.8)	*	159	4.6	0.4	(3.7, 5.4)	*
GEORGIA	51	3.7	0.5	(2.7, 4.7)	*	75	4.4	0.5	(3.3, 5.4)	*
HAWAII	49	5.5	0.6	(4.3, 6.8)	*	60	8.3	1.8	(4.8, 11.9)	*
IDAHO	91	3.7	0.4	(2.8, 4.5)	*	110	3.8	0.4	(2.9, 4.6)	*
IAOWA	73	4.1	0.9	(2.4, 5.8)	*	116	4.0	0.4	(3.3, 4.7)	*
KANSAS	68	5.6	1.2	(3.3, 7.9)	*	92	4.3	0.5	(3.4, 5.3)	*
KENTUCKY	77	4.3	0.4	(3.5, 5.1)	*	144	5.8	0.6	(4.7, 6.8)	*
MAINE	16	4.0	1.0	(2.0, 6.0)	*	54	3.8	0.5	(2.9, 4.8)	*
MASSACHUSETTS	92	2.9	0.3	(2.3, 3.6)	*	112	4.4	0.5	(3.5, 5.3)	*
MICHIGAN	57	3.2	0.4	(2.3, 4.0)	*	102	5.3	0.8	(3.7, 6.9)	*
MINNESOTA	80	3.6	0.5	(2.6, 4.5)	*	109	3.4	0.4	(2.7, 4.1)	*
MISSISSIPPI	62	4.8	1.0	(2.9, 6.6)	*	105	4.1	0.4	(3.3, 4.8)	*
MONTANA	24	4.7	1.0	(2.7, 6.7)	*	41	4.5	1.0	(2.5, 6.5)	*
NEBRASKA	64	4.0	0.8	(2.4, 5.6)	*	94	5.3	0.7	(3.9, 6.8)	*
NEVADA	41	4.3	1.3	(1.8, 6.8)	*	42	3.5	0.9	(1.8, 5.2)	*
NEW HAMPSHIRE	28	2.3	0.3	(1.8, 2.8)	*	28	5.5	1.0	(3.5, 7.5)	*
NEW JERSEY	59	4.2	0.4	(3.4, 5.0)	*	67	6.3	1.2	(4.1, 8.6)	*
NEW MEXICO	81	4.6	0.7	(3.3, 5.8)	*	99	6.7	1.1	(4.6, 8.9)	*
NORTH CAROLINA	48	4.0	0.5	(2.9, 5.0)	*	92	4.9	0.6	(3.8, 6.0)	*
NORTH DAKOTA	34	4.8	0.6	(3.5, 6.1)	*	43	3.4	0.4	(2.6, 4.3)	*
OHIO	65	4.2	0.7	(2.9, 5.5)	*	123	4.6	0.4	(3.7, 5.4)	*
PENNSYLVANIA	72	4.2	0.4	(3.4, 5.0)	*	127	4.9	0.4	(4.1, 5.7)	*
RHODE ISLAND	85	3.7	0.4	(3.0, 4.5)	*	123	3.9	0.3	(3.2, 4.5)	*
TENNESSEE	45	3.9	0.5	(2.9, 4.9)	*	116	5.0	0.6	(3.9, 6.1)	*
TEXAS	118	4.1	0.4	(3.2, 4.9)	*	211	5.3	0.4	(4.5, 6.1)	*
UTAH	59	3.9	0.7	(2.6, 5.3)	*	66	4.4	0.6	(3.2, 5.6)	*
VERMONT	59	3.5	0.4	(2.7, 4.3)	*	74	4.6	0.6	(3.4, 5.8)	*
VIRGINIA	80	3.4	0.4	(2.5, 4.3)	*	101	4.4	0.5	(3.4, 5.3)	*
WEST VIRGINIA	59	4.8	0.6	(3.7, 5.9)	*	108	4.8	0.4	(4.0, 5.6)	*
WISCONSIN	52	3.4	0.9	(1.6, 5.2)	*	52	4.3	0.8	(2.8, 5.9)	*
WYOMING	42	3.2	0.5	(2.2, 4.3)	*	42	3.2	0.6	(2.0, 4.3)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 39
MEDIAN = 4.1
RANGE = 2.3- 7.0

NUMBER OF PARTICIPANTS = 39
MEDIAN = 4.5
RANGE = 2.9- 8.3

* MEAN TIMES SEEN HEALTH PROFESSIONALS FOR DIABETES IN PAST YEAR

June 18, 1999

TABLE 23.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DID NOT HAVE FLU SHOT IN PAST 12 MONTHS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

63

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
GEORGIA	370	36.9	2.9	(31.1, 42.7)
ILLINOIS	237	35.3	3.6	(28.2, 42.4)
INDIANA	444	33.7	2.4	(29.0, 38.5)
LOUISIANA	298	40.3	3.2	(34.1, 46.5)
MINNESOTA	716	36.4	1.9	(32.7, 40.1)
MONTANA	388	27.1	2.4	(22.5, 31.7)
NEW JERSEY	444	33.3	2.4	(28.6, 38.1)
OHIO	681	31.0	2.4	(26.3, 35.7)
SOUTH CAROLINA	537	37.5	2.3	(33.1, 42.0)
TENNESSEE	560	30.4	2.2	(26.1, 34.7)
TEXAS	884	35.3	2.1	(31.2, 39.5)
WEST VIRGINIA	562	37.1	2.2	(32.7, 41.5)
WYOMING	413	28.6	2.4	(23.9, 33.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 13 MEDIAN = 35.3 RANGE = 27.1-40.3 YEAR 2000 OBJ. 20.11

* AMONG RESPONDENTS 65 YEARS OF AGE AND OLDER

June 29, 1999

TABLE 23.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DID NOT HAVE FLU SHOT IN PAST 12 MONTHS -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

64

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
GEORGIA	85	37.6	5.7	(26.5, 48.7)	*	285	36.4	3.2	(30.2, 42.6)	
ILLINOIS	77	43.4	6.7	(30.2, 56.5)	*	160	30.4	4.0	(22.7, 38.2)	
INDIANA	148	33.6	4.1	(25.6, 41.5)	*	296	33.8	3.0	(27.9, 39.7)	
LOUISIANA	91	41.2	5.7	(30.1, 52.2)	*	207	39.7	3.8	(32.3, 47.2)	
MINNESOTA	249	35.9	3.2	(29.6, 42.1)	*	467	36.7	2.3	(32.2, 41.2)	
MONTANA	154	26.8	3.6	(19.7, 33.9)	*	234	27.3	3.2	(21.1, 33.5)	
NEW JERSEY	168	33.1	3.9	(25.5, 40.7)	*	276	33.5	3.1	(27.4, 39.6)	
OHIO	229	30.0	4.1	(21.9, 38.0)	*	452	31.7	2.9	(26.0, 37.5)	
SOUTH CAROLINA	222	28.8	3.3	(22.2, 35.3)	*	315	43.4	3.1	(37.4, 49.3)	
TENNESSEE	147	28.9	4.0	(21.0, 36.7)	*	413	31.4	2.5	(26.5, 36.4)	
TEXAS	296	39.7	3.8	(32.3, 47.2)	*	588	32.3	2.3	(27.9, 36.8)	
WEST VIRGINIA	188	40.4	3.7	(33.1, 47.8)	*	374	34.9	2.7	(29.7, 40.2)	
WYOMING	139	26.9	3.9	(19.3, 34.5)	*	274	29.9	3.0	(24.0, 35.7)	

SUMMARY STATISTICS:
NUMBER OF PARTICIPANTS = 13
MEDIAN = 33.6
RANGE = 26.8-43.4

NUMBER OF PARTICIPANTS = 13
MEDIAN = 33.5
RANGE = 27.3-43.4

* AMONG RESPONDENTS 65 YEARS OF AGE AND OLDER

June 29, 1999

TABLE 24.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NEVER HAD PNEUMONIA VACCINATION*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

65

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
GEORGIA	361	49.6	3.1	(43.6, 55.6)
ILLINOIS	232	45.3	3.7	(38.0, 52.6)
INDIANA	432	53.2	2.6	(48.0, 58.4)
LOUISIANA	289	60.4	3.3	(53.9, 66.9)
MINNESOTA	703	53.9	2.0	(50.0, 57.8)
MONTANA	378	44.1	2.7	(38.8, 49.4)
NEW JERSEY	432	52.8	2.6	(47.7, 57.8)
OHIO	661	51.4	2.7	(46.1, 56.7)
SOUTH CAROLINA	527	49.3	2.4	(44.6, 53.9)
TENNESSEE	554	48.9	2.5	(44.0, 53.8)
TEXAS	871	50.1	2.1	(46.0, 54.1)
WEST VIRGINIA	556	54.9	2.3	(50.3, 59.5)
WYOMING	407	44.1	2.7	(38.9, 49.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 13 MEDIAN = 50.1 RANGE = 44.1-60.4 YEAR 2000 OBJ. 20.11

* AMONG RESPONDENTS 65 YEARS OF AGE AND OLDER

June 29, 1999

TABLE 24.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NEVER HAD PNEUMONIA VACCINATION* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

66

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
GEORGIA	81	50.6	6.0	(38.8, 62.3)	*	280	49.0	3.3	(42.6, 55.4)	
ILLINOIS	75	58.0	6.4	(45.4, 70.5)	*	157	37.8	4.2	(29.5, 46.0)	
INDIANA	141	55.9	4.5	(47.1, 64.6)	*	291	51.5	3.2	(45.2, 57.8)	
LOUISIANA	88	56.5	5.9	(45.0, 68.0)	*	201	62.9	3.9	(55.4, 70.5)	
MINNESOTA	243	58.2	3.2	(51.9, 64.5)	*	460	50.9	2.6	(45.8, 56.0)	
MONTANA	146	44.8	4.3	(36.4, 53.1)	*	232	43.6	3.4	(36.9, 50.3)	
NEW JERSEY	162	54.9	4.2	(46.8, 63.0)	*	270	51.3	3.3	(44.9, 57.8)	
OHIO	219	54.1	4.6	(45.1, 63.1)	*	442	49.6	3.3	(43.2, 56.1)	
SOUTH CAROLINA	217	46.2	3.7	(38.9, 53.4)	*	310	51.4	3.1	(45.3, 57.4)	
TENNESSEE	146	45.6	4.6	(36.6, 54.6)	*	408	51.1	2.7	(45.8, 56.4)	
TEXAS	291	54.7	3.4	(48.1, 61.2)	*	580	47.0	2.5	(42.1, 52.0)	
WEST VIRGINIA	183	57.9	4.0	(50.1, 65.6)	*	373	53.0	2.9	(47.2, 58.7)	
WYOMING	137	49.5	4.5	(40.6, 58.3)	*	270	40.1	3.2	(33.9, 46.3)	

SUMMARY STATISTICS:
 NUMBER OF PARTICIPANTS = 13
 MEDIAN = 54.7
 RANGE = 44.8-58.2

NUMBER OF PARTICIPANTS = 13
 MEDIAN = 50.9
 RANGE = 37.8-62.9

* AMONG RESPONDENTS 65 YEARS OF AGE AND OLDER

June 29, 1999

TABLE 25.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

67

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	1989	24.7	1.4	(22.0, 27.4)
DELAWARE	2572	9.3	0.8	(7.7, 10.8)
IDAHO	4926	21.1	0.7	(19.7, 22.5)
INDIANA	2423	16.8	0.9	(15.0, 18.5)
MONTANA	1798	27.9	1.2	(25.6, 30.2)
NORTH DAKOTA	1794	20.6	1.1	(18.5, 22.8)
OHIO	2889	18.5	1.1	(16.4, 20.7)
OKLAHOMA	2513	11.6	0.8	(10.1, 13.1)
SOUTH CAROLINA	3209	18.3	0.8	(16.7, 19.8)
VIRGINIA	3504	14.1	0.8	(12.6, 15.6)
WASHINGTON	3570	17.8	0.8	(16.3, 19.3)
WEST VIRGINIA	2405	29.1	1.1	(26.9, 31.2)
WYOMING	2431	32.2	1.1	(30.0, 34.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 13 MEDIAN = 18.5 RANGE = 9.3-32.2

* EVER USED OR TRIED SMOKELESS TOBACCO

June 18, 1999

TABLE 25.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO* -BY SEX
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

68

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALASKA	924	41.4	2.3	(36.9, 45.9)	*	1065	6.5	0.9	(4.8, 8.2)	*
DELAWARE	1032	17.7	1.6	(14.6, 20.8)	*	1540	1.6	0.4	(0.9, 2.3)	*
IDAHO	2079	37.6	1.3	(35.1, 40.1)	*	2847	5.3	0.5	(4.3, 6.4)	*
INDIANA	991	31.6	1.7	(28.4, 34.9)	*	1432	3.3	0.5	(2.2, 4.3)	*
MONTANA	783	50.2	1.9	(46.4, 53.9)	*	1015	6.8	0.9	(5.0, 8.5)	*
NORTH DAKOTA	801	36.4	1.8	(32.8, 40.0)	*	993	5.4	0.9	(3.6, 7.1)	*
OHIO	1148	36.0	2.0	(32.1, 39.8)	*	1741	3.0	0.6	(1.9, 4.1)	*
OKLAHOMA	1001	20.2	1.4	(17.4, 23.0)	*	1512	3.8	0.6	(2.7, 4.9)	*
SOUTH CAROLINA	1340	33.5	1.5	(30.6, 36.4)	*	1869	4.4	0.6	(3.4, 5.5)	*
VIRGINIA	1484	27.3	1.5	(24.4, 30.2)	*	2020	1.9	0.3	(1.2, 2.5)	*
WASHINGTON	1574	32.1	1.4	(29.5, 34.8)	*	1996	4.0	0.5	(3.0, 4.9)	*
WEST VIRGINIA	940	56.9	1.8	(53.4, 60.4)	*	1465	4.2	0.6	(3.0, 5.4)	*
WYOMING	1023	56.5	1.7	(53.1, 60.0)	*	1408	8.4	0.8	(6.8, 9.9)	*

SUMMARY STATISTICS:
 NUMBER OF PARTICIPANTS = 13
 MEDIAN = 36.0
 RANGE = 17.7-56.9

NUMBER OF PARTICIPANTS = 13
 MEDIAN = 4.2
 RANGE = 1.6- 8.4

* EVER USED OR TRIED SMOKELESS TOBACCO

June 18, 1999

TABLE 26.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

69

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	1989	5.4	0.7	(4.0, 6.8)
DELAWARE	2572	1.1	0.3	(0.6, 1.6)
IDAHO	4926	3.6	0.3	(2.9, 4.2)
INDIANA	2423	2.6	0.4	(1.8, 3.3)
MONTANA	1796	6.8	0.7	(5.4, 8.3)
NORTH DAKOTA	1794	4.0	0.5	(3.0, 5.1)
OHIO	2889	3.6	0.6	(2.5, 4.6)
OKLAHOMA	2513	3.8	0.5	(2.9, 4.7)
SOUTH CAROLINA	3208	3.1	0.4	(2.4, 3.9)
VIRGINIA	3502	3.0	0.4	(2.3, 3.7)
WASHINGTON	3569	3.0	0.4	(2.2, 3.7)
WEST VIRGINIA	2405	8.4	0.7	(7.0, 9.8)
WYOMING	2431	6.9	0.6	(5.7, 8.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 13 MEDIAN = 3.6 RANGE = 1.1- 8.4

* CURRENT USERS OF SMOKELESS TOBACCO

June 18, 1999

TABLE 26.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS* -BY SEX
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

70

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
	*	*	*	*	*	*	*	*	*	*
ALASKA	924	8.6	1.3	(6.0, 11.1)	*	1065	1.9	0.5	(1.0, 2.8)	*
DELAWARE	1032	2.2	0.6	(1.1, 3.3)	*	1540	0.1	0.1	(-0.1, 0.3)	*
IDAHO	2079	6.9	0.6	(5.6, 8.1)	*	2847	0.4	0.1	(0.1, 0.7)	*
INDIANA	991	5.1	0.8	(3.5, 6.6)	*	1432	0.3	0.1	(0.0, 0.6)	*
MONTANA	781	13.7	1.4	(10.9, 16.4)	*	1015	0.4	0.2	(-0.0, 0.8)	*
NORTH DAKOTA	801	8.0	1.1	(5.9, 10.0)	*	993	0.2	0.1	(-0.1, 0.4)	*
OHIO	1148	7.4	1.2	(5.1, 9.7)	*	1741	0.1	0.1	(-0.1, 0.4)	*
OKLAHOMA	1001	6.9	0.9	(5.2, 8.7)	*	1512	0.9	0.3	(0.3, 1.6)	*
SOUTH CAROLINA	1339	5.9	0.8	(4.4, 7.5)	*	1869	0.6	0.3	(0.1, 1.1)	*
VIRGINIA	1483	5.9	0.7	(4.5, 7.2)	*	2019	0.3	0.1	(0.1, 0.6)	*
WASHINGTON	1573	6.1	0.8	(4.6, 7.6)	*	1996	0.0	0.0	(0.0, 0.0)	*
WEST VIRGINIA	940	17.5	1.4	(14.7, 20.3)	*	1465	0.3	0.2	(-0.1, 0.6)	*
WYOMING	1023	13.6	1.2	(11.3, 16.0)	*	1408	0.3	0.2	(-0.0, 0.6)	*

SUMMARY STATISTICS:
 NUMBER OF PARTICIPANTS = 13
 MEDIAN = 6.9
 RANGE = 2.2-17.5

NUMBER OF PARTICIPANTS = 13
 MEDIAN = 0.3
 RANGE = 0.0- 1.9

* CURRENT USERS OF SMOKELESS TOBACCO

June 18, 1999

TABLE 27.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NEVER HAD MAMMOGRAM FOR WOMEN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

71

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1457	38.0	1.5	(35.1, 41.0)
ALASKA	1058	44.3	2.1	(40.2, 48.3)
ARIZONA	1091	43.8	2.5	(38.8, 48.7)
ARKANSAS	1796	40.7	1.4	(38.0, 43.4)
COLORADO	983	39.9	1.7	(36.4, 43.3)
CONNECTICUT	1813	36.8	1.4	(34.0, 39.6)
DELAWARE	1541	35.8	1.7	(32.5, 39.1)
DISTRICT OF CO	906	36.3	1.9	(32.7, 40.0)
FLORIDA	2824	35.0	1.0	(32.9, 37.0)
GEORGIA	1470	42.2	1.5	(39.4, 45.1)
HAWAII	1219	38.0	1.8	(34.5, 41.5)
IDAHO	2844	43.6	1.1	(41.5, 45.7)
ILLINOIS	1643	41.5	1.4	(38.9, 44.1)
INDIANA	1425	42.3	1.4	(39.5, 45.2)
IOWA	2168	39.5	1.3	(37.0, 41.9)
KANSAS	2286	40.9	1.2	(38.6, 43.3)
KENTUCKY	2328	40.2	1.2	(37.9, 42.4)
LOUISIANA	1004	42.3	1.8	(38.8, 45.7)
MAINE	946	40.5	1.9	(36.7, 44.2)
MARYLAND	1979	36.1	1.5	(33.3, 39.0)
MASSACHUSETTS	2884	36.7	1.2	(34.3, 39.1)
MIAMI GAN	1495	38.2	1.4	(35.4, 40.9)
MINNESOTA	2227	51.4	1.2	(49.0, 53.7)
MISSISSIPPI	1419	43.7	1.5	(40.8, 46.6)
MISSOURI	2273	40.3	1.3	(37.9, 42.8)
MONTANA	1014	42.2	1.7	(39.0, 45.5)
NEBRASKA	1644	44.0	1.4	(41.3, 46.7)
NEVADA	1079	39.9	2.4	(35.3, 44.5)
NEW HAMPSHIRE	856	42.6	2.0	(38.7, 46.6)
NEW JERSEY	1440	39.9	1.5	(36.9, 42.8)
NEW MEXICO	2061	44.3	1.2	(42.0, 46.7)
NEW YORK	1459	38.1	1.4	(35.3, 40.8)
NORTH CAROLINA	1460	40.0	1.6	(37.0, 43.1)
NORTH DAKOTA	996	41.8	1.8	(38.3, 45.3)
OHIO	1793	41.7	1.7	(38.3, 45.0)
OKLAHOMA	1510	47.7	1.4	(44.9, 50.6)
OREGON	1076	40.3	1.7	(36.9, 43.7)
PENNSYLVANIA	2144	37.5	1.2	(35.2, 39.9)
RHODE ISLAND	2172	38.5	1.2	(36.2, 40.9)
SOUTH CAROLINA	1910	39.6	1.3	(37.0, 42.1)
SOUTH DAKOTA	1182	42.0	1.7	(38.7, 45.3)
TENNESSEE	1903	38.5	1.3	(36.0, 41.1)
TEXAS	3549	43.8	1.1	(41.7, 46.0)
UTAH	1620	47.2	1.7	(43.9, 50.4)
VERMONT	1698	43.7	1.4	(40.8, 46.5)
VIRGINIA	2015	41.2	1.7	(37.8, 44.6)
WASHINGTON	2006	39.5	1.3	(37.0, 42.0)
WEST VIRGINIA	1465	40.1	1.4	(37.3, 42.9)
WISCONSIN	1224	41.2	1.8	(37.6, 44.8)
WYOMING	1408	43.5	1.5	(40.6, 46.4)
PUERTO RICO	1732	49.0	1.4	(46.4, 51.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 40.7 RANGE = 35.0-51.4

* NEVER HAD A MAMMOGRAM, AGE 18 AND OVER

June 18, 1999

TABLE 27.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NEVER HAD MAMMOGRAM FOR WOMEN*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

72

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	35.0	38.0	3.0
ALASKA	49.8	44.3	-5.6
ARIZONA	45.4	43.8	-1.6
ARKANSAS	44.0	40.7	-3.3
COLORADO	42.7	39.9	-2.9
CONNECTICUT	38.3	36.8	-1.5
DELAWARE	36.9	35.8	-1.1
DISTRICT OF CO	40.1	36.3	-3.8
FLORIDA	33.3	35.0	1.6
GEORGIA	41.0	42.2	1.2
HAWAII	37.7	38.0	0.3
IDAHO	44.1	43.6	-0.6
ILLINOIS	42.4	41.5	-0.9
INDIANA	42.9	42.3	-0.5
IOWA	41.7	39.5	-2.2
KANSAS	45.5	40.9	-4.5
KENTUCKY	42.0	40.2	-1.8
LOUISIANA	42.3	42.3	0.0
MAINE	40.7	40.5	-0.2
MARYLAND	40.3	36.1	-4.1
MASSACHUSETTS	39.4	36.7	-2.7
MICHIGAN	37.7	38.2	0.5
MINNESOTA	43.9	51.4	7.5
MISSISSIPPI	45.9	43.7	-2.2
MISSOURI	41.3	40.3	-1.0
MONTANA	40.4	42.2	1.8
NEBRASKA	44.5	44.0	-0.4
NEVADA	39.7	39.9	0.2
NEW HAMPSHIRE	40.9	42.6	1.8
NEW JERSEY	37.2	39.9	2.6
NEW MEXICO	45.8	44.3	-1.5
NEW YORK	39.4	38.1	-1.4
NORTH CAROLINA	39.1	40.0	1.0
NORTH DAKOTA	41.6	41.8	0.2
OHIO	39.6	41.7	2.1
OKLAHOMA	47.3	47.7	0.4
OREGON	38.9	40.3	1.4
PENNSYLVANIA	38.8	37.5	-1.3
RHODE ISLAND	37.8	38.5	0.8
SOUTH CAROLINA	39.1	39.6	0.5
SOUTH DAKOTA	40.9	42.0	1.1
TENNESSEE	41.6	38.5	-3.0
TEXAS	44.3	43.8	-0.5
UTAH	48.7	47.2	-1.6
VERMONT	45.6	43.7	-1.9
VIRGINIA	42.6	41.2	-1.4
WASHINGTON	39.9	39.5	-0.4
WEST VIRGINIA	41.1	40.1	-1.0
WISCONSIN	43.3	41.2	-2.2
WYOMING	40.3	43.5	3.2
PUERTO RICO	44.9	49.0	4.1

* AMONG WOMEN, AGE 18 AND OVER

June 18, 1999

TABLE 28.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NEVER HAD MAMMOGRAM FOR WOMEN, AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

73

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	956	14.3	1.2	(11.9, 16.6)
ALASKA	548	10.6	1.7	(7.3, 14.0)
ARIZONA	686	18.0	3.8	(10.5, 25.5)
ARKANSAS	1161	17.5	1.3	(15.0, 19.9)
COLORADO	580	11.0	1.4	(8.3, 13.7)
CONNECTICUT	1083	12.9	1.3	(10.4, 15.5)
DELAWARE	964	9.9	1.2	(7.5, 12.3)
DISTRICT OF CO	534	8.3	1.3	(5.8, 10.8)
FLORIDA	1839	13.0	0.9	(11.3, 14.8)
GEORGIA	861	15.6	1.4	(12.9, 18.4)
HAWAII	766	11.1	1.4	(8.3, 13.9)
IDAHO	1797	18.2	1.0	(16.2, 20.3)
ILLINOIS	983	16.5	1.3	(13.9, 19.1)
INDIANA	861	15.9	1.4	(13.1, 18.7)
IOWA	1422	14.9	1.1	(12.8, 17.0)
KANSAS	1490	16.1	1.1	(14.1, 18.2)
KENTUCKY	1502	17.2	1.1	(15.0, 19.3)
LOUISIANA	596	19.4	1.8	(16.0, 22.9)
MAINE	642	13.1	1.4	(10.3, 16.0)
MARYLAND	1159	10.9	1.1	(8.7, 13.1)
MASSACHUSETTS	1635	8.4	0.9	(6.7, 10.0)
MICHIGAN	898	10.7	1.2	(8.4, 13.0)
MINNESOTA	1288	27.5	1.4	(24.8, 30.1)
MISSISSIPPI	876	19.5	1.5	(16.5, 22.5)
MISSOURI	1458	15.3	1.0	(13.3, 17.3)
MONTANA	643	17.3	1.5	(14.3, 20.2)
NEBRASKA	1086	18.6	1.3	(16.0, 21.2)
NEVADA	656	13.9	2.0	(9.9, 17.8)
NEW HAMPSHIRE	505	15.1	1.7	(11.8, 18.4)
NEW JERSEY	875	16.8	1.4	(14.1, 19.5)
NEW MEXICO	1293	16.0	1.2	(13.7, 18.2)
NEW YORK	858	13.3	1.3	(10.8, 15.8)
NORTH CAROLINA	900	15.2	1.5	(12.2, 18.2)
NORTH DAKOTA	659	14.6	1.5	(11.7, 17.4)
OHIO	1182	15.4	1.5	(12.4, 18.4)
OKLAHOMA	998	24.9	1.5	(21.9, 27.8)
OREGON	656	13.8	1.5	(10.8, 16.7)
PENNSYLVANIA	1334	15.2	1.2	(12.9, 17.4)
RHODE ISLAND	1318	11.7	1.0	(9.7, 13.7)
SOUTH CAROLINA	1161	15.8	1.2	(13.3, 18.2)
SOUTH DAKOTA	753	15.9	1.4	(13.2, 18.6)
TENNESSEE	1223	15.0	1.1	(12.8, 17.3)
TEXAS	2063	18.1	1.0	(16.1, 20.2)
UTAH	902	15.9	1.7	(12.6, 19.2)
VERMONT	1104	15.8	1.2	(13.4, 18.2)
VIRGINIA	1199	13.6	1.3	(11.0, 16.1)
WASHINGTON	1207	11.0	0.9	(9.2, 12.9)
WEST VIRGINIA	992	17.0	1.3	(14.5, 19.5)
WISCONSIN	743	14.6	1.5	(11.8, 17.5)
WYOMING	888	17.5	1.3	(14.8, 20.1)
PUERTO RICO	1040	24.7	1.5	(21.8, 27.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 15.3 RANGE = 8.3-27.5

*AMONG WOMEN AGE 40 AND OLDER

June 18, 1999

TABLE 28.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NEVER HAD MAMMOGRAM FOR WOMEN, AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

74

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	13.0	14.3	1.3
ALASKA	15.8	10.6	-5.2
ARIZONA	21.6	18.0	-3.6
ARKANSAS	23.5	17.5	-6.0
COLORADO	15.0	11.0	-3.9
CONNECTICUT	12.5	12.9	0.5
DELAWARE	10.7	9.9	-0.8
DISTRICT OF CO	12.3	8.3	-4.0
FLORIDA	12.9	13.0	0.1
GEORGIA	12.7	15.6	2.9
HAWAII	11.6	11.1	-0.5
IDAHO	18.9	18.2	-0.7
ILLINOIS	15.5	16.5	1.0
INDIANA	19.2	15.9	-3.3
IOWA	17.9	14.9	-2.9
KANSAS	20.6	16.1	-4.4
KENTUCKY	19.7	17.2	-2.6
LOUISIANA	19.6	19.4	-0.2
MAINE	11.7	13.1	1.4
MARYLAND	12.2	10.9	-1.3
MASSACHUSETTS	11.4	8.4	-3.0
MICHIGAN	11.2	10.7	-0.5
MINNESOTA	16.7	27.5	10.7
MISSISSIPPI	24.8	19.5	-5.2
MISSOURI	17.3	15.3	-1.9
MONTANA	17.3	17.3	-0.1
NEBRASKA	19.9	18.6	-1.3
NEVADA	14.8	13.9	-1.0
NEW HAMPSHIRE	11.4	15.1	3.8
NEW JERSEY	14.1	16.8	2.8
NEW MEXICO	18.6	16.0	-2.7
NEW YORK	13.3	13.3	0.0
NORTH CAROLINA	13.5	15.2	1.7
NORTH DAKOTA	15.5	14.6	-0.9
OHIO	16.2	15.4	-0.8
OKLAHOMA	25.6	24.9	-0.7
OREGON	12.7	13.8	1.1
PENNSYLVANIA	16.1	15.2	-1.0
RHODE ISLAND	11.0	11.7	0.7
SOUTH CAROLINA	10.8	15.8	5.0
SOUTH DAKOTA	15.2	15.9	0.7
TENNESSEE	19.0	15.0	-4.0
TEXAS	18.0	18.1	0.2
UTAH	18.8	15.9	-2.9
VERMONT	16.1	15.8	-0.3
VIRGINIA	14.8	13.6	-1.2
WASHINGTON	12.4	11.0	-1.4
WEST VIRGINIA	19.0	17.0	-2.0
WISCONSIN	17.4	14.6	-2.8
WYOMING	15.6	17.5	1.8
PUERTO RICO	23.1	24.7	1.7

* AMONG WOMEN, AGE 40 AND OVER

June 18, 1999

TABLE 29.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE DID NOT HAVE MAMMOGRAM WITH PAST TWO YEARS, WOMEN, AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

75

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	655	23.9	1.8	(20.4, 27.4)
ALASKA	272	20.9	3.7	(13.6, 28.1)
ARIZONA	509	17.4	2.2	(13.0, 21.7)
ARKANSAS	815	34.1	1.9	(30.3, 37.9)
COLORADO	356	22.6	2.4	(17.9, 27.2)
CONNECTICUT	715	22.0	1.9	(18.2, 25.8)
DELAWARE	661	18.7	1.8	(15.1, 22.3)
DISTRICT OF CO	373	10.6	1.7	(7.2, 14.0)
FLORIDA	1328	22.0	1.3	(19.4, 24.5)
GEORGIA	569	27.5	2.1	(23.3, 31.6)
HAWAII	496	22.9	2.8	(17.5, 28.3)
IDAHO	1193	29.1	1.5	(26.3, 32.0)
ILLINOIS	631	28.6	1.9	(24.9, 32.3)
INDIANA	565	28.5	2.0	(24.5, 32.5)
IOWA	965	28.6	1.7	(25.2, 31.9)
KANSAS	1033	24.7	1.5	(21.7, 27.6)
KENTUCKY	1057	31.6	1.6	(28.6, 34.7)
LOUISIANA	401	29.8	2.5	(24.9, 34.6)
MAINE	445	22.9	2.1	(18.7, 27.1)
MARYLAND	739	19.2	1.8	(15.6, 22.8)
MASSACHUSETTS	1069	15.8	1.4	(13.2, 18.5)
MIAMI GAN	583	21.1	1.8	(17.5, 24.6)
MINNESOTA	872	35.1	1.8	(31.6, 38.5)
MISSISSIPPI	615	32.1	2.1	(28.1, 36.1)
MISSOURI	1035	26.0	1.5	(23.1, 29.0)
MONTANA	446	27.7	2.2	(23.4, 31.9)
NEBRASKA	778	30.3	1.8	(26.8, 33.7)
NEVADA	426	28.4	3.3	(21.9, 34.8)
NEW HAMPSHIRE	320	22.9	2.5	(18.0, 27.8)
NEW JERSEY	579	24.5	1.9	(20.7, 28.3)
NEW MEXICO	866	25.0	1.6	(21.9, 28.0)
NEW YORK	539	22.9	2.0	(19.0, 26.8)
NORTH CAROLINA	608	21.0	1.8	(17.4, 24.5)
NORTH DAKOTA	449	25.1	2.2	(20.8, 29.4)
OHIO	836	26.6	2.1	(22.4, 30.8)
OKLAHOMA	722	29.1	1.8	(25.6, 32.7)
OREGON	440	21.3	2.2	(17.0, 25.6)
PENNSYLVANIA	929	22.7	1.5	(19.8, 25.6)
RHODE ISLAND	864	18.8	1.5	(15.9, 21.7)
SOUTH CAROLINA	728	22.8	1.8	(19.3, 26.4)
SOUTH DAKOTA	508	25.0	2.1	(20.9, 29.0)
TENNESSEE	822	24.8	1.7	(21.5, 28.0)
TEXAS	1303	27.3	1.3	(24.7, 29.9)
UTAH	587	28.8	2.5	(23.9, 33.7)
VERMONT	708	23.0	1.7	(19.6, 26.3)
VIRGINIA	786	22.2	2.0	(18.2, 26.2)
WASHINGTON	735	21.6	1.6	(18.5, 24.7)
WEST VIRGINIA	705	25.0	1.8	(21.6, 28.5)
WISCONSIN	482	26.0	2.3	(21.5, 30.5)
WYOMING	567	31.3	2.1	(27.2, 35.4)
PUERTO RICO	717	43.2	2.1	(39.2, 47.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 24.8 RANGE = 10.6-43.2

* AMONG WOMEN, AGE 50 AND OVER

June 18, 1999

PREVALENCE DIFFERENCES FROM 1997 TO 1998
 DID NOT HAVE MAMMOGRAM WITH PAST TWO YEARS, WOMEN, AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	24.8	23.9	-0.9
ALASKA	21.8	20.9	-1.0
ARIZONA	26.2	17.4	-8.8
ARKANSAS	44.1	34.1	-10.0
COLORADO	23.9	22.6	-1.3
CONNECTICUT	22.3	22.0	-0.3
DELAWARE	20.3	18.7	-1.5
DISTRICT OF CO	16.4	10.6	-5.8
FLORIDA	21.1	22.0	0.9
GEORGIA	24.6	27.5	2.8
HAWAII	17.6	22.9	5.3
IDAHO	35.1	29.1	-6.0
ILLINOIS	27.0	28.6	1.7
INDIANA	34.5	28.5	-6.0
IOWA	32.1	28.6	-3.6
KANSAS	30.0	24.7	-5.3
KENTUCKY	30.4	31.6	1.3
LOUISIANA	30.1	29.8	-0.4
MAINE	22.4	22.9	0.5
MARYLAND	19.5	19.2	-0.3
MASSACHUSETTS	19.1	15.8	-3.2
MICHIGAN	20.1	21.1	1.0
MINNESOTA	26.3	35.1	8.8
MISSISSIPPI	35.3	32.1	-3.2
MISSOURI	31.0	26.0	-4.9
MONTANA	29.6	27.7	-2.0
NEBRASKA	32.5	30.3	-2.2
NEVADA	32.3	28.4	-4.0
NEW HAMPSHIRE	20.3	22.9	2.6
NEW JERSEY	26.3	24.5	-1.8
NEW MEXICO	32.3	25.0	-7.4
NEW YORK	21.9	22.9	1.0
NORTH CAROLINA	25.6	21.0	-4.6
NORTH DAKOTA	27.4	25.1	-2.3
OHIO	25.1	26.6	1.5
OKLAHOMA	34.8	29.1	-5.7
OREGON	20.7	21.3	0.6
PENNSYLVANIA	26.2	22.7	-3.5
RHODE ISLAND	16.9	18.8	2.0
SOUTH CAROLINA	24.5	22.8	-1.6
SOUTH DAKOTA	29.2	25.0	-4.3
TENNESSEE	26.0	24.8	-1.2
TEXAS	31.9	27.3	-4.6
UTAH	29.8	28.8	-1.0
VERMONT	26.5	23.0	-3.5
VIRGINIA	23.7	22.2	-1.5
WASHINGTON	26.6	21.6	-5.0
WEST VIRGINIA	30.7	25.0	-5.6
WISCONSIN	29.1	26.0	-3.1
WYOMING	30.7	31.3	0.6
PUERTO RICO	38.1	43.2	5.1

* AMONG WOMEN, AGE 50 AND OVER

June 18, 1999

TABLE 30.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NEVER HAD BREAST EXAM, FOR WOMEN AGE 18 AND OLDER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

77

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1459	13.3	1.2	(11.0, 15.6)
ALASKA	1058	8.4	1.2	(5.9, 10.8)
ARIZONA	1087	18.7	2.7	(13.4, 23.9)
ARKANSAS	1794	17.5	1.1	(15.2, 19.7)
COLORADO	979	8.1	1.1	(6.0, 10.2)
CONNECTICUT	1803	13.7	1.1	(11.6, 15.9)
DELAWARE	1537	5.5	0.8	(3.9, 7.1)
DISTRICT OF CO	906	6.1	0.8	(4.5, 7.7)
FLORIDA	2823	14.8	0.8	(13.3, 16.3)
GEORGIA	1466	11.2	1.0	(9.3, 13.1)
HAWAII	1214	10.9	1.3	(8.4, 13.3)
IDAHO	2838	8.1	0.6	(6.9, 9.2)
ILLINOIS	1643	14.3	1.1	(12.2, 16.5)
INDIANA	1418	12.6	1.0	(10.6, 14.7)
IOWA	2165	9.6	0.8	(7.9, 11.2)
KANSAS	2273	13.3	0.9	(11.6, 14.9)
KENTUCKY	2323	10.1	0.8	(8.7, 11.6)
LOUISIANA	1001	13.2	1.2	(10.8, 15.6)
MAINE	945	9.0	1.2	(6.6, 11.4)
MARYLAND	1975	10.9	0.9	(9.1, 12.7)
MASSACHUSETTS	2864	11.2	0.9	(9.6, 12.9)
MIAMI GAN	1488	11.2	0.9	(9.4, 13.0)
MINNESOTA	2226	7.1	0.6	(5.9, 8.2)
MISSISSIPPI	1416	14.9	1.1	(12.7, 17.0)
MISSOURI	2257	11.1	0.9	(9.4, 12.8)
MONTANA	1009	7.2	0.9	(5.4, 9.0)
NEBRASKA	1634	9.7	0.8	(8.1, 11.2)
NEVADA	1077	11.1	1.4	(8.4, 13.8)
NEW HAMPSHIRE	852	8.9	1.2	(6.6, 11.2)
NEW JERSEY	1433	17.3	1.2	(14.8, 19.7)
NEW MEXICO	2050	12.8	0.9	(11.1, 14.5)
NEW YORK	1453	15.3	1.1	(13.1, 17.5)
NORTH CAROLINA	1458	8.0	1.0	(6.1, 10.0)
NORTH DAKOTA	995	8.3	1.0	(6.3, 10.3)
OHIO	1791	8.9	1.0	(7.0, 10.7)
OKLAHOMA	1507	20.5	1.2	(18.1, 22.8)
OREGON	1073	9.3	1.3	(6.8, 11.9)
PENNSYLVANIA	2136	14.9	0.9	(13.2, 16.7)
RHODE ISLAND	2163	14.1	0.9	(12.3, 15.8)
SOUTH CAROLINA	1904	15.7	1.0	(13.6, 17.7)
SOUTH DAKOTA	1178	8.5	1.2	(6.1, 10.9)
TENNESSEE	1902	11.6	0.9	(9.9, 13.3)
TEXAS	3542	17.4	0.9	(15.6, 19.2)
UTAH	1618	12.8	1.1	(10.7, 15.0)
VERMONT	1692	12.8	1.0	(10.9, 14.8)
VIRGINIA	2004	11.3	1.0	(9.3, 13.4)
WASHINGTON	1992	7.7	0.7	(6.4, 9.1)
WEST VIRGINIA	1464	8.8	0.9	(7.0, 10.5)
WISCONSIN	1217	8.7	1.1	(6.5, 10.9)
WYOMING	1405	8.2	0.8	(6.7, 9.7)
PUERTO RICO	1720	25.6	1.3	(23.2, 28.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 11.2 RANGE = 5.5-25.6

* NEVER HAD A BREAST EXAM BY A DOCTOR, AMONG WOMEN AGE 18 AND OVER

June 18, 1999

TABLE 30.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NEVER HAD BREAST EXAM, FOR WOMEN AGE 18 AND OLDER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

78

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	11.3	13.3	2.0
ALASKA	8.6	8.4	-0.2
ARIZONA	19.8	18.7	-1.2
ARKANSAS	17.1	17.5	0.3
COLORADO	7.7	8.1	0.4
CONNECTICUT	12.3	13.7	1.5
DELAWARE	6.4	5.5	-0.9
DISTRICT OF CO	7.1	6.1	-1.0
FLORIDA	11.5	14.8	3.3
GEORGIA	6.7	11.2	4.5
HAWAII	10.2	10.9	0.7
IDAHO	7.6	8.1	0.4
ILLINOIS	13.7	14.3	0.6
INDIANA	11.1	12.6	1.5
IOWA	10.1	9.6	-0.5
KANSAS	8.7	13.3	4.5
KENTUCKY	12.0	10.1	-1.8
LOUISIANA	15.1	13.2	-1.9
MAINE	6.6	9.0	2.4
MARYLAND	5.2	10.9	5.7
MASSACHUSETTS	10.1	11.2	1.1
MICHIGAN	10.7	11.2	0.5
MINNESOTA	8.2	7.1	-1.2
MISSISSIPPI	11.5	14.9	3.4
MISSOURI	7.4	11.1	3.7
MONTANA	6.9	7.2	0.3
NEBRASKA	9.6	9.7	0.1
NEVADA	8.0	11.1	3.1
NEW HAMPSHIRE	6.7	8.9	2.2
NEW JERSEY	13.7	17.3	3.6
NEW MEXICO	13.2	12.8	-0.4
NEW YORK	7.2	15.3	8.2
NORTH CAROLINA	6.7	8.0	1.3
NORTH DAKOTA	8.6	8.3	-0.4
OHIO	9.3	8.9	-0.4
OKLAHOMA	7.5	20.5	13.0
OREGON	5.8	9.3	3.5
PENNSYLVANIA	15.5	14.9	-0.5
RHODE ISLAND	13.2	14.1	0.9
SOUTH CAROLINA	6.9	15.7	8.8
SOUTH DAKOTA	8.8	8.5	-0.2
TENNESSEE	10.8	11.6	0.8
TEXAS	16.6	17.4	0.7
UTAH	7.8	12.8	5.0
VERMONT	11.6	12.8	1.2
VIRGINIA	12.2	11.3	-0.9
WASHINGTON	8.7	7.7	-1.0
WEST VIRGINIA	9.9	8.8	-1.1
WISCONSIN	6.5	8.7	2.2
WYOMING	8.8	8.2	-0.6
PUERTO RICO	18.6	25.6	7.0

* NEVER HAD A BREAST EXAM BY A DOCTOR, WOMEN AGE 18 AND OVER

June 18, 1999

TABLE 31.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NEVER HAD A BREAST EXAM FOR WOMEN, AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

79

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	956	13.2	1.2	(10.9, 15.5)
ALASKA	548	7.1	1.7	(3.8, 10.4)
ARIZONA	682	16.8	3.9	(9.3, 24.4)
ARKANSAS	1159	14.6	1.3	(12.1, 17.1)
COLORADO	576	5.4	1.0	(3.6, 7.3)
CONNECTICUT	1077	12.4	1.4	(9.7, 15.1)
DELAWARE	960	5.1	0.9	(3.3, 6.8)
DISTRICT OF CO	533	4.6	0.9	(2.8, 6.4)
FLORIDA	1837	13.5	0.9	(11.8, 15.3)
GEORGIA	858	11.6	1.2	(9.2, 14.0)
HAWAII	761	8.5	1.5	(5.6, 11.3)
IDAHO	1791	7.1	0.7	(5.8, 8.4)
ILLINOIS	983	13.6	1.3	(11.1, 16.1)
INDIANA	853	11.7	1.3	(9.2, 14.2)
IOWA	1419	9.3	1.0	(7.4, 11.2)
KANSAS	1479	12.0	0.9	(10.2, 13.8)
KENTUCKY	1499	9.9	0.9	(8.2, 11.6)
LOUISIANA	594	13.8	1.5	(10.8, 16.8)
MAINE	641	6.8	1.1	(4.7, 8.9)
MARYLAND	1156	11.8	1.3	(9.4, 14.3)
MASSACHUSETTS	1621	10.2	1.0	(8.3, 12.0)
MICHIGAN	893	11.2	1.3	(8.7, 13.7)
MINNESOTA	1286	7.6	0.8	(6.2, 9.1)
MISSISSIPPI	874	15.4	1.3	(12.7, 18.0)
MISSOURI	1441	10.8	1.0	(8.9, 12.7)
MONTANA	639	4.8	0.9	(3.1, 6.6)
NEBRASKA	1077	10.0	1.0	(8.1, 11.9)
NEVADA	654	12.7	2.0	(8.9, 16.5)
NEW HAMPSHIRE	501	9.0	1.5	(6.1, 11.9)
NEW JERSEY	871	15.5	1.4	(12.8, 18.1)
NEW MEXICO	1282	10.9	1.0	(8.9, 12.8)
NEW YORK	852	13.9	1.4	(11.2, 16.6)
NORTH CAROLINA	899	7.2	1.2	(4.8, 9.6)
NORTH DAKOTA	658	7.5	1.1	(5.4, 9.6)
OHIO	1180	7.8	1.0	(5.8, 9.8)
OKLAHOMA	995	15.8	1.3	(13.2, 18.3)
OREGON	655	7.5	1.2	(5.2, 9.9)
PENNSYLVANIA	1327	15.1	1.1	(13.0, 17.1)
RHODE ISLAND	1312	12.4	1.0	(10.4, 14.3)
SOUTH CAROLINA	1156	14.8	1.3	(12.3, 17.2)
SOUTH DAKOTA	749	6.0	0.9	(4.3, 7.7)
TENNESSEE	1222	9.4	0.9	(7.6, 11.1)
TEXAS	2057	14.5	0.9	(12.8, 16.2)
UTAH	899	9.1	1.2	(6.7, 11.5)
VERMONT	1100	12.9	1.1	(10.7, 15.0)
VIRGINIA	1187	11.1	1.1	(8.9, 13.3)
WASHINGTON	1195	5.6	0.7	(4.2, 6.9)
WEST VIRGINIA	992	7.5	0.9	(5.7, 9.3)
WISCONSIN	736	7.0	1.1	(4.9, 9.1)
WYOMING	884	8.5	1.0	(6.5, 10.4)
PUERTO RICO	1030	19.3	1.4	(16.5, 22.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 10.8 RANGE = 4.6-19.3

* NEVER HAD A BREAST EXAM BY A DOCTOR, WOMEN AGE 40 AND OVER

June 18, 1999

TABLE 31.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 ESTIMATED PREVALENCE NEVER HAD A BREAST EXAM FOR WOMEN, AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

80

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	12.2	13.2	1.0
ALASKA	7.3	7.1	-0.2
ARIZONA	18.8	16.8	-1.9
ARKANSAS	16.7	14.6	-2.1
COLORADO	6.9	5.4	-1.5
CONNECTICUT	12.3	12.4	0.1
DELAWARE	6.1	5.1	-1.0
DISTRICT OF CO	7.0	4.6	-2.4
FLORIDA	9.7	13.5	3.8
GEORGIA	7.4	11.6	4.3
HAWAII	8.6	8.5	-0.1
IDAHO	7.2	7.1	-0.0
ILLINOIS	11.7	13.6	1.9
INDIANA	12.8	11.7	-1.1
IOWA	9.4	9.3	-0.1
KANSAS	9.5	12.0	2.5
KENTUCKY	10.9	9.9	-1.0
LOUISIANA	16.7	13.8	-2.9
MAINE	6.8	6.8	-0.0
MARYLAND	4.2	11.8	7.7
MASSACHUSETTS	9.4	10.2	0.8
MICHIGAN	8.8	11.2	2.4
MINNESOTA	6.1	7.6	1.5
MISSISSIPPI	9.8	15.4	5.6
MISSOURI	8.2	10.8	2.6
MONTANA	5.9	4.8	-1.1
NEBRASKA	9.8	10.0	0.3
NEVADA	8.3	12.7	4.4
NEW HAMPSHIRE	5.7	9.0	3.3
NEW JERSEY	13.5	15.5	1.9
NEW MEXICO	11.9	10.9	-1.0
NEW YORK	5.5	13.9	8.4
NORTH CAROLINA	5.7	7.2	1.5
NORTH DAKOTA	7.1	7.5	0.4
OHIO	7.4	7.8	0.4
OKLAHOMA	5.7	15.8	10.1
OREGON	5.1	7.5	2.5
PENNSYLVANIA	15.8	15.1	-0.7
RHODE ISLAND	11.8	12.4	0.5
SOUTH CAROLINA	5.0	14.8	9.7
SOUTH DAKOTA	8.9	6.0	-3.0
TENNESSEE	10.7	9.4	-1.4
TEXAS	11.6	14.5	2.9
UTAH	4.2	9.1	4.9
VERMONT	10.2	12.9	2.7
VIRGINIA	10.6	11.1	0.5
WASHINGTON	6.2	5.6	-0.6
WEST VIRGINIA	9.4	7.5	-1.9
WISCONSIN	6.4	7.0	0.6
WYOMING	7.4	8.5	1.0
PUERTO RICO	11.2	19.3	8.0

* NEVER HAD A BREAST EXAM BY A DOCTOR, WOMEN AGE 40 AND OVER

June 18, 1999

TABLE 32.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE DID NOT HAVE BREAST EXAM WITHIN PAST TWO YEARS FOR WOMEN, AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

81

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	639	22.0	1.7	(18.6, 25.4)
ALASKA	270	20.2	3.6	(13.1, 27.4)
ARIZONA	505	17.3	2.2	(13.0, 21.6)
ARKANSAS	814	34.7	1.9	(30.9, 38.5)
COLORADO	349	22.0	2.4	(17.4, 26.6)
CONNECTICUT	707	22.2	2.0	(18.3, 26.1)
DELAWARE	650	16.5	1.7	(13.2, 19.8)
DISTRICT OF CO	370	9.8	1.8	(6.4, 13.3)
FLORIDA	1316	24.0	1.3	(21.4, 26.6)
GEORGIA	563	27.1	2.1	(22.9, 31.3)
HAWAII	482	20.4	2.6	(15.3, 25.6)
IDAHO	1181	26.6	1.4	(23.9, 29.4)
ILLINOIS	624	27.6	2.1	(23.6, 31.6)
INDIANA	556	26.6	2.0	(22.6, 30.5)
IOWA	958	27.7	1.7	(24.3, 31.1)
KANSAS	1013	24.2	1.5	(21.3, 27.1)
KENTUCKY	1056	27.5	1.5	(24.6, 30.4)
LOUISIANA	392	28.5	2.4	(23.7, 33.2)
MAINE	444	18.5	2.2	(14.2, 22.9)
MARYLAND	730	22.5	2.0	(18.5, 26.4)
MASSACHUSETTS	1062	20.6	1.6	(17.5, 23.7)
MICHIGAN	577	24.6	2.0	(20.7, 28.5)
MINNESOTA	863	27.8	1.7	(24.5, 31.1)
MISSISSIPPI	602	29.9	2.0	(25.9, 33.8)
MISSOURI	1016	26.0	1.6	(22.9, 29.1)
MONTANA	437	22.2	2.2	(18.0, 26.5)
NEBRASKA	763	30.1	1.8	(26.7, 33.6)
NEVADA	424	29.5	3.3	(23.1, 36.0)
NEW HAMPSHIRE	316	20.1	2.5	(15.1, 25.0)
NEW JERSEY	571	28.3	2.0	(24.3, 32.3)
NEW MEXICO	854	27.1	1.7	(23.8, 30.4)
NEW YORK	537	23.5	2.0	(19.5, 27.5)
NORTH CAROLINA	603	18.4	1.8	(14.9, 21.9)
NORTH DAKOTA	447	23.3	2.2	(18.9, 27.6)
OHIO	828	24.7	2.1	(20.6, 28.8)
OKLAHOMA	720	25.5	1.8	(22.0, 29.1)
OREGON	437	25.2	2.4	(20.6, 29.9)
PENNSYLVANIA	920	27.0	1.6	(23.9, 30.1)
RHODE ISLAND	863	22.2	1.6	(19.1, 25.2)
SOUTH CAROLINA	718	24.8	1.8	(21.2, 28.3)
SOUTH DAKOTA	499	19.2	1.8	(15.6, 22.7)
TENNESSEE	824	20.5	1.5	(17.5, 23.5)
TEXAS	1297	29.3	1.4	(26.5, 32.0)
UTAH	580	30.1	2.4	(25.5, 34.8)
VERMONT	701	24.7	1.8	(21.3, 28.2)
VIRGINIA	775	24.4	2.0	(20.5, 28.4)
WASHINGTON	721	22.3	1.9	(18.6, 26.0)
WEST VIRGINIA	703	22.8	1.7	(19.4, 26.1)
WISCONSIN	469	25.2	2.4	(20.6, 29.8)
WYOMING	555	32.2	2.1	(28.0, 36.4)
PUERTO RICO	709	41.0	2.1	(36.9, 45.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 24.7 RANGE = 9.8-41.0

* AMONG WOMEN, AGE 50 AND OVER

June 18, 1999

TABLE 32.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 DID NOT HAVE BREAST EXAM WITHIN PAST TWO YEARS FOR WOMEN, AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

82

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	26.0	22.0	-4.0
ALASKA	20.3	20.2	-0.1
ARIZONA	24.4	17.3	-7.1
ARKANSAS	36.6	34.7	-1.9
COLORADO	21.5	22.0	0.5
CONNECTICUT	23.7	22.2	-1.5
DELAWARE	18.7	16.5	-2.1
DISTRICT OF CO	16.7	9.8	-6.9
FLORIDA	21.8	24.0	2.2
GEORGIA	17.2	27.1	9.9
HAWAII	19.1	20.4	1.3
IDAHO	28.3	26.6	-1.6
ILLINOIS	27.6	27.6	0.0
INDIANA	30.6	26.6	-4.0
IOWA	26.3	27.7	1.5
KANSAS	23.6	24.2	0.6
KENTUCKY	26.7	27.5	0.8
LOUISIANA	35.2	28.5	-6.7
MAINE	17.0	18.5	1.5
MARYLAND	15.6	22.5	6.9
MASSACHUSETTS	16.3	20.6	4.4
MICHIGAN	22.4	24.6	2.2
MINNESOTA	21.3	27.8	6.5
MISSISSIPPI	26.7	29.9	3.2
MISSOURI	25.5	26.0	0.5
MONTANA	21.6	22.2	0.6
NEBRASKA	28.9	30.1	1.3
NEVADA	30.3	29.5	-0.7
NEW HAMPSHIRE	14.7	20.1	5.3
NEW JERSEY	26.8	28.3	1.5
NEW MEXICO	27.5	27.1	-0.4
NEW YORK	13.6	23.5	9.9
NORTH CAROLINA	17.6	18.4	0.8
NORTH DAKOTA	24.4	23.3	-1.1
OHIO	18.3	24.7	6.4
OKLAHOMA	18.3	25.5	7.2
OREGON	19.9	25.2	5.3
PENNSYLVANIA	29.7	27.0	-2.7
RHODE ISLAND	20.4	22.2	1.8
SOUTH CAROLINA	17.0	24.8	7.7
SOUTH DAKOTA	24.1	19.2	-4.9
TENNESSEE	20.2	20.5	0.3
TEXAS	28.1	29.3	1.2
UTAH	22.8	30.1	7.4
VERMONT	24.5	24.7	0.2
VIRGINIA	21.5	24.4	3.0
WASHINGTON	22.3	22.3	-0.0
WEST VIRGINIA	27.3	22.8	-4.5
WISCONSIN	24.8	25.2	0.4
WYOMING	29.2	32.2	3.1
PUERTO RICO	23.2	41.0	17.8

* WOMEN, AGE 50 AND OVER

June 18, 1999

TABLE 33.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NEVER HAD MAMMOGRAM AND BREAST EXAM FOR WOMEN, AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

83

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	952	21.5	1.4	(18.7, 24.3)
ALASKA	547	15.0	2.1	(10.9, 19.0)
ARIZONA	682	21.2	3.9	(13.7, 28.8)
ARKANSAS	1156	25.2	1.5	(22.2, 28.2)
COLORADO	576	15.0	1.6	(11.9, 18.0)
CONNECTICUT	1077	20.4	1.6	(17.2, 23.5)
DELAWARE	958	12.1	1.3	(9.5, 14.7)
DISTRICT OF CO	533	11.6	1.5	(8.7, 14.6)
FLORIDA	1836	21.2	1.1	(19.1, 23.2)
GEORGIA	857	21.9	1.6	(18.8, 25.0)
HAWAII	760	16.7	1.8	(13.1, 20.2)
IDAHO	1788	22.8	1.1	(20.6, 25.0)
ILLINOIS	983	24.0	1.5	(21.0, 27.0)
INDIANA	853	23.0	1.6	(19.8, 26.1)
IOWA	1418	19.2	1.2	(16.8, 21.6)
KANSAS	1478	22.4	1.2	(20.0, 24.7)
KENTUCKY	1495	19.8	1.1	(17.6, 22.0)
LOUISIANA	593	26.3	1.9	(22.5, 30.0)
MAINE	641	15.4	1.5	(12.4, 18.4)
MARYLAND	1155	19.3	1.5	(16.4, 22.2)
MASSACHUSETTS	1621	15.2	1.1	(13.0, 17.5)
MICHIGAN	893	18.3	1.5	(15.4, 21.2)
MINNESOTA	1285	30.4	1.4	(27.6, 33.1)
MISSISSIPPI	870	26.9	1.7	(23.6, 30.2)
MISSOURI	1441	21.4	1.3	(19.0, 23.9)
MONTANA	637	18.7	1.6	(15.6, 21.8)
NEBRASKA	1077	24.4	1.5	(21.5, 27.2)
NEVADA	653	21.8	2.4	(17.0, 26.6)
NEW HAMPSHIRE	501	19.5	2.0	(15.6, 23.4)
NEW JERSEY	871	25.4	1.6	(22.2, 28.6)
NEW MEXICO	1281	22.2	1.3	(19.6, 24.8)
NEW YORK	851	21.4	1.6	(18.3, 24.5)
NORTH CAROLINA	898	18.4	1.6	(15.2, 21.6)
NORTH DAKOTA	657	18.7	1.6	(15.5, 21.8)
OHIO	1179	18.3	1.6	(15.1, 21.5)
OKLAHOMA	995	31.5	1.6	(28.3, 34.7)
OREGON	655	19.0	1.7	(15.6, 22.4)
PENNSYLVANIA	1327	24.0	1.3	(21.4, 26.6)
RHODE ISLAND	1311	19.5	1.2	(17.1, 22.0)
SOUTH CAROLINA	1155	24.3	1.5	(21.4, 27.1)
SOUTH DAKOTA	749	18.8	1.5	(15.8, 21.7)
TENNESSEE	1221	19.5	1.2	(17.1, 22.0)
TEXAS	2055	25.9	1.2	(23.6, 28.2)
UTAH	899	22.1	1.8	(18.5, 25.7)
VERMONT	1098	24.0	1.4	(21.2, 26.7)
VIRGINIA	1186	21.3	1.6	(18.2, 24.3)
WASHINGTON	1195	14.9	1.1	(12.8, 17.0)
WEST VIRGINIA	989	19.6	1.4	(16.9, 22.2)
WISCONSIN	735	17.7	1.6	(14.6, 20.8)
WYOMING	884	22.2	1.5	(19.3, 25.1)
PUERTO RICO	1028	32.9	1.7	(29.6, 36.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 21.2 RANGE = 11.6-32.9 YEAR 2000 OBJ 16.11

* AMONG WOMEN, AGE 40 AND OVER

June 18, 1999

TABLE 33.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NEVER HAD MAMMOGRAM AND BREAST EXAM FOR WOMEN, AGE 40 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

84

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	22.0	21.5	-0.5
ALASKA	19.9	15.0	-4.9
ARIZONA	26.7	21.2	-5.5
ARKANSAS	32.0	25.2	-6.8
COLORADO	17.9	15.0	-3.0
CONNECTICUT	19.5	20.4	0.9
DELAWARE	14.6	12.1	-2.5
DISTRICT OF CO	17.0	11.6	-5.4
FLORIDA	18.5	21.2	2.6
GEORGIA	16.3	21.9	5.7
HAWAII	17.5	16.7	-0.9
IDAHO	23.0	22.8	-0.2
ILLINOIS	22.8	24.0	1.3
INDIANA	24.5	23.0	-1.5
IOWA	22.3	19.2	-3.1
KANSAS	25.0	22.4	-2.6
KENTUCKY	22.3	19.8	-2.5
LOUISIANA	29.1	26.3	-2.9
MAINE	15.4	15.4	0.0
MARYLAND	13.7	19.3	5.6
MASSACHUSETTS	16.8	15.2	-1.6
MICHIGAN	16.8	18.3	1.5
MINNESOTA	19.2	30.4	11.1
MISSISSIPPI	28.2	26.9	-1.3
MISSOURI	20.8	21.4	0.6
MONTANA	20.6	18.7	-1.9
NEBRASKA	25.0	24.4	-0.6
NEVADA	19.0	21.8	2.8
NEW HAMPSHIRE	14.5	19.5	5.0
NEW JERSEY	23.2	25.4	2.2
NEW MEXICO	24.1	22.2	-1.8
NEW YORK	15.7	21.4	5.7
NORTH CAROLINA	15.6	18.4	2.7
NORTH DAKOTA	18.0	18.7	0.7
OHIO	18.3	18.3	0.0
OKLAHOMA	27.4	31.5	4.1
OREGON	15.0	19.0	4.0
PENNSYLVANIA	25.9	24.0	-1.8
RHODE ISLAND	19.3	19.5	0.3
SOUTH CAROLINA	13.6	24.3	10.7
SOUTH DAKOTA	20.2	18.8	-1.4
TENNESSEE	24.9	19.5	-5.4
TEXAS	23.0	25.9	2.9
UTAH	20.4	22.1	1.6
VERMONT	21.4	24.0	2.6
VIRGINIA	23.1	21.3	-1.8
WASHINGTON	16.7	14.9	-1.8
WEST VIRGINIA	22.0	19.6	-2.4
WISCONSIN	19.8	17.7	-2.1
WYOMING	20.0	22.2	2.3
PUERTO RICO	26.8	32.9	6.0

* AMONG WOMEN, AGE 40 AND OVER

June 18, 1999

TABLE 34.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE DID NOT HAVE MAMMOGRAM AND BREAST EXAM WITHIN LAST TWO YEARS FOR WOMEN, AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

85

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	637	30.1	2.0	(26.3, 34.0)
ALASKA	270	25.9	3.8	(18.5, 33.4)
ARIZONA	507	19.0	2.2	(14.6, 23.4)
ARKANSAS	812	43.0	2.0	(39.0, 46.9)
COLORADO	349	29.9	2.6	(24.7, 35.1)
CONNECTICUT	706	30.6	2.2	(26.3, 34.9)
DELAWARE	650	22.9	2.0	(19.0, 26.8)
DISTRICT OF CO	371	14.8	2.1	(10.7, 18.9)
FLORIDA	1316	31.1	1.4	(28.3, 33.9)
GEORGIA	566	35.9	2.3	(31.4, 40.3)
HAWAII	486	30.3	3.0	(24.4, 36.2)
IDAHO	1182	36.5	1.6	(33.5, 39.6)
ILLINOIS	625	37.9	2.1	(33.8, 42.0)
INDIANA	558	36.1	2.2	(31.8, 40.4)
IOWA	960	36.0	1.8	(32.4, 39.5)
KANSAS	1018	33.7	1.7	(30.4, 36.9)
KENTUCKY	1053	36.3	1.6	(33.1, 39.4)
LOUISIANA	395	36.5	2.5	(31.6, 41.4)
MAINE	444	26.4	2.4	(21.6, 31.1)
MARYLAND	729	29.3	2.2	(25.0, 33.5)
MASSACHUSETTS	1054	25.2	1.7	(21.9, 28.5)
MICHIGAN	579	30.2	2.1	(26.1, 34.3)
MINNESOTA	863	40.0	1.8	(36.4, 43.6)
MISSISSIPPI	601	39.9	2.2	(35.6, 44.2)
MISSOURI	1024	33.4	1.7	(30.0, 36.7)
MONTANA	439	31.0	2.3	(26.6, 35.5)
NEBRASKA	769	40.5	1.9	(36.7, 44.3)
NEVADA	423	37.6	3.5	(30.7, 44.5)
NEW HAMPSHIRE	318	28.4	2.8	(23.0, 33.8)
NEW JERSEY	574	34.9	2.2	(30.6, 39.1)
NEW MEXICO	857	33.7	1.8	(30.3, 37.2)
NEW YORK	534	31.3	2.3	(26.9, 35.8)
NORTH CAROLINA	605	25.9	2.0	(21.9, 29.8)
NORTH DAKOTA	447	31.7	2.4	(27.0, 36.4)
OHIO	831	31.9	2.3	(27.5, 36.3)
OKLAHOMA	719	37.5	2.0	(33.6, 41.4)
OREGON	437	32.2	2.5	(27.3, 37.1)
PENNSYLVANIA	922	33.5	1.7	(30.2, 36.7)
RHODE ISLAND	856	29.8	1.7	(26.4, 33.1)
SOUTH CAROLINA	722	33.1	2.0	(29.1, 37.1)
SOUTH DAKOTA	501	30.5	2.2	(26.3, 34.8)
TENNESSEE	821	29.1	1.8	(25.7, 32.6)
TEXAS	1296	37.0	1.5	(34.1, 39.9)
UTAH	580	38.5	2.6	(33.4, 43.5)
VERMONT	700	32.9	1.9	(29.1, 36.7)
VIRGINIA	775	31.9	2.3	(27.4, 36.3)
WASHINGTON	718	29.8	2.1	(25.8, 33.8)
WEST VIRGINIA	701	31.1	1.9	(27.4, 34.8)
WISCONSIN	472	33.2	2.5	(28.2, 38.1)
WYOMING	560	39.5	2.2	(35.2, 43.9)
PUERTO RICO	715	50.9	2.1	(46.8, 55.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 32.2 RANGE = 14.8-50.9 YEAR 2000 OJB 16.11

* AMONG WOMEN AGE 50 AND OLDER

June 18, 1999

TABLE 34.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 DID NOT HAVE MAMMOGRAM AND BREAST EXAM WITHIN LAST TWO YEARS FOR WOMEN, AGE 50 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	33.7	30.1	-3.5
ALASKA	29.1	25.9	-3.1
ARIZONA	30.5	19.0	-11.5
ARKANSAS	51.7	43.0	-8.7
COLORADO	31.2	29.9	-1.3
CONNECTICUT	31.0	30.6	-0.4
DELAWARE	26.7	22.9	-3.8
DISTRICT OF CO	25.4	14.8	-10.6
FLORIDA	29.8	31.1	1.3
GEORGIA	29.3	35.9	6.5
HAWAII	24.9	30.3	5.4
IDAHO	41.7	36.5	-5.2
ILLINOIS	35.8	37.9	2.1
INDIANA	42.8	36.1	-6.7
IOWA	37.8	36.0	-1.8
KANSAS	37.0	33.7	-3.3
KENTUCKY	33.6	36.3	2.7
LOUISIANA	41.9	36.5	-5.4
MAINE	26.9	26.4	-0.5
MARYLAND	24.0	29.3	5.2
MASSACHUSETTS	25.2	25.2	0.1
MICHIGAN	28.4	30.2	1.8
MINNESOTA	31.0	40.0	9.0
MISSISSIPPI	39.6	39.9	0.3
MISSOURI	36.2	33.4	-2.8
MONTANA	33.1	31.0	-2.0
NEBRASKA	40.0	40.5	0.5
NEVADA	41.0	37.6	-3.4
NEW HAMPSHIRE	24.4	28.4	4.0
NEW JERSEY	36.1	34.9	-1.3
NEW MEXICO	38.3	33.7	-4.6
NEW YORK	25.4	31.3	5.9
NORTH CAROLINA	29.3	25.9	-3.4
NORTH DAKOTA	33.7	31.7	-2.0
OHIO	28.6	31.9	3.3
OKLAHOMA	38.3	37.5	-0.9
OREGON	26.8	32.2	5.4
PENNSYLVANIA	36.5	33.5	-3.0
RHODE ISLAND	27.0	29.8	2.8
SOUTH CAROLINA	28.5	33.1	4.5
SOUTH DAKOTA	36.7	30.5	-6.1
TENNESSEE	32.5	29.1	-3.3
TEXAS	37.6	37.0	-0.6
UTAH	36.3	38.5	2.2
VERMONT	34.3	32.9	-1.3
VIRGINIA	31.3	31.9	0.6
WASHINGTON	33.0	29.8	-3.2
WEST VIRGINIA	36.0	31.1	-4.9
WISCONSIN	35.2	33.2	-2.0
WYOMING	38.2	39.5	1.3
PUERTO RICO	41.2	50.9	9.7

* AMONG WOMEN AGE 50 AND OLDER

June 18, 1999

TABLE 35.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NEVER HAD PAP SMEAR, WOMEN AGE 18 AND OVER
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

87

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1448	4.6	0.8	(3.0, 6.1)
ALASKA	1050	3.1	0.8	(1.5, 4.7)
ARIZONA	1086	13.1	2.0	(9.3, 17.0)
ARKANSAS	1781	7.0	0.8	(5.4, 8.6)
COLORADO	977	4.2	0.8	(2.5, 5.8)
CONNECTICUT	1768	7.7	0.8	(6.1, 9.3)
DELAWARE	1537	4.6	0.9	(2.9, 6.3)
DISTRICT OF CO	899	3.0	0.6	(1.8, 4.1)
FLORIDA	2804	4.6	0.5	(3.7, 5.6)
GEORGIA	1463	4.2	0.6	(3.0, 5.4)
HAWAII	1208	4.8	0.9	(3.1, 6.6)
IDAHO	2837	5.3	0.6	(4.2, 6.3)
ILLINOIS	1638	5.2	0.7	(3.8, 6.6)
INDIANA	1413	5.9	0.8	(4.3, 7.5)
IOWA	2164	5.2	0.7	(3.8, 6.6)
KANSAS	2254	6.2	0.6	(5.0, 7.5)
KENTUCKY	2325	5.4	0.6	(4.3, 6.5)
LOUISIANA	989	7.3	1.0	(5.4, 9.2)
MAINE	941	5.3	1.0	(3.5, 7.2)
MARYLAND	1959	5.0	0.6	(3.8, 6.2)
MASSACHUSETTS	2816	6.6	0.7	(5.2, 8.0)
MICHIGAN	1480	4.5	0.6	(3.4, 5.7)
MINNESOTA	2229	5.5	0.5	(4.4, 6.5)
MISSISSIPPI	1406	6.2	0.8	(4.6, 7.7)
MISSOURI	2258	4.5	0.6	(3.3, 5.7)
MONTANA	1007	3.6	0.7	(2.3, 5.0)
NEBRASKA	1626	5.3	0.6	(4.0, 6.6)
NEVADA	1076	4.9	1.0	(3.0, 6.8)
NEW HAMPSHIRE	841	5.6	1.0	(3.7, 7.5)
NEW JERSEY	1411	9.9	1.0	(7.9, 11.9)
NEW MEXICO	2046	6.0	0.7	(4.7, 7.2)
NEW YORK	1422	6.8	0.8	(5.2, 8.4)
NORTH CAROLINA	1450	4.1	0.8	(2.6, 5.6)
NORTH DAKOTA	995	4.3	0.8	(2.6, 5.9)
OHIO	1785	5.3	0.8	(3.7, 6.8)
OKLAHOMA	1506	7.4	0.8	(5.9, 8.9)
OREGON	1072	5.6	1.2	(3.3, 7.9)
PENNSYLVANIA	2114	6.7	0.6	(5.5, 7.9)
RHODE ISLAND	2138	9.3	0.8	(7.7, 10.8)
SOUTH CAROLINA	1893	6.0	0.9	(4.2, 7.7)
SOUTH DAKOTA	1172	5.8	1.2	(3.5, 8.1)
TENNESSEE	1886	7.2	0.7	(5.8, 8.6)
TEXAS	3533	7.1	0.6	(6.0, 8.2)
UTAH	1610	8.6	1.0	(6.6, 10.6)
VERMONT	1677	6.5	0.8	(5.0, 8.1)
VIRGINIA	1992	5.1	0.7	(3.6, 6.5)
WASHINGTON	1997	4.4	0.6	(3.3, 5.5)
WEST VIRGINIA	1460	6.3	0.8	(4.7, 7.8)
WISCONSIN	1215	5.0	0.8	(3.4, 6.5)
WYOMING	1405	3.0	0.5	(2.0, 4.0)
PUERTO RICO	1727	16.8	1.2	(14.5, 19.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 5.4 RANGE = 3.0-16.8 YEAR 2000 OBJ 16.11
 * AMONG WOMEN, AGE 18 AND OVER June 18, 1999

TABLE 35.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NEVER HAD PAP SMEAR, WOMEN AGE 18 AND OVER*
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

88

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	3.5	4.6	1.1
ALASKA	3.8	3.1	-0.7
ARIZONA	15.3	13.1	-2.2
ARKANSAS	8.0	7.0	-0.9
COLORADO	3.5	4.2	0.7
CONNECTICUT	8.2	7.7	-0.4
DELAWARE	3.7	4.6	0.9
DISTRICT OF CO	4.0	3.0	-1.1
FLORIDA	6.4	4.6	-1.7
GEORGIA	3.2	4.2	1.0
HAWAII	4.1	4.8	0.8
IDAHO	3.9	5.3	1.4
ILLINOIS	6.5	5.2	-1.3
INDIANA	4.1	5.9	1.8
IOWA	5.7	5.2	-0.5
KANSAS	5.9	6.2	0.3
KENTUCKY	7.2	5.4	-1.8
LOUISIANA	6.6	7.3	0.7
MAINE	4.3	5.3	1.1
MARYLAND	5.4	5.0	-0.4
MASSACHUSETTS	6.5	6.6	0.1
MICHIGAN	5.3	4.5	-0.8
MINNESOTA	7.1	5.5	-1.6
MISSISSIPPI	4.2	6.2	2.0
MISSOURI	3.5	4.5	1.0
MONTANA	3.8	3.6	-0.2
NEBRASKA	4.9	5.3	0.4
NEVADA	4.5	4.9	0.4
NEW HAMPSHIRE	5.5	5.6	0.0
NEW JERSEY	8.6	9.9	1.3
NEW MEXICO	6.5	6.0	-0.5
NEW YORK	7.0	6.8	-0.2
NORTH CAROLINA	4.7	4.1	-0.6
NORTH DAKOTA	5.7	4.3	-1.4
OHIO	5.2	5.3	0.1
OKLAHOMA	4.8	7.4	2.6
OREGON	3.8	5.6	1.8
PENNSYLVANIA	8.3	6.7	-1.6
RHODE ISLAND	6.2	9.3	3.0
SOUTH CAROLINA	3.5	6.0	2.4
SOUTH DAKOTA	4.4	5.8	1.4
TENNESSEE	5.3	7.2	2.0
TEXAS	6.9	7.1	0.1
UTAH	7.6	8.6	1.1
VERMONT	6.8	6.5	-0.3
VIRGINIA	4.5	5.1	0.5
WASHINGTON	3.6	4.4	0.8
WEST VIRGINIA	6.3	6.3	-0.1
WISCONSIN	4.4	5.0	0.6
WYOMING	4.3	3.0	-1.3
PUERTO RICO	15.6	16.8	1.3

* AMONG WOMEN, AGE 18 AND OVER

June 18, 1999

TABLE 36.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE NEVER HAD PAP SMEAR, FOR WOMEN AGE 18 AND OLDER WITH INTACT CERVIX *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

89

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	964	5.7	1.1	(3.5, 7.9)
ALASKA	900	3.3	0.9	(1.5, 5.1)
ARIZONA	838	15.1	2.4	(10.4, 19.8)
ARKANSAS	1218	7.9	1.1	(5.8, 10.0)
COLORADO	746	4.5	1.0	(2.5, 6.5)
CONNECTICUT	1495	7.3	0.9	(5.6, 9.0)
DELAWARE	1209	5.1	1.0	(3.1, 7.1)
DISTRICT OF CO	748	3.0	0.7	(1.7, 4.2)
FLORIDA	2045	5.2	0.6	(4.1, 6.4)
GEORGIA	1070	4.3	0.8	(2.8, 5.7)
HAWAII	1001	5.5	1.0	(3.5, 7.6)
IDAHO	2023	4.8	0.7	(3.5, 6.1)
ILLINOIS	1311	5.6	0.8	(4.0, 7.2)
INDIANA	1121	6.3	0.9	(4.6, 8.1)
IOWA	1679	5.2	0.8	(3.5, 6.8)
KANSAS	1646	6.5	0.8	(5.0, 8.0)
KENTUCKY	1697	5.0	0.7	(3.7, 6.4)
LOUISIANA	673	6.4	1.1	(4.2, 8.6)
MAINE	737	4.7	1.1	(2.6, 6.8)
MARYLAND	1571	4.8	0.7	(3.4, 6.2)
MASSACHUSETTS	2435	6.2	0.8	(4.6, 7.7)
MICHIGAN	1163	4.6	0.7	(3.2, 6.0)
MINNESOTA	1858	6.0	0.6	(4.8, 7.3)
MISSISSIPPI	947	6.7	1.0	(4.7, 8.7)
MISSOURI	1680	4.3	0.7	(2.8, 5.7)
MONTANA	767	3.5	0.8	(2.0, 5.0)
NEBRASKA	1197	5.4	0.8	(3.9, 6.9)
NEVADA	771	4.7	1.0	(2.8, 6.6)
NEW HAMPSHIRE	711	5.6	1.1	(3.4, 7.7)
NEW JERSEY	1214	10.3	1.1	(8.0, 12.5)
NEW MEXICO	1515	6.7	0.8	(5.1, 8.3)
NEW YORK	1234	7.4	0.9	(5.6, 9.1)
NORTH CAROLINA	1080	4.6	1.0	(2.7, 6.4)
NORTH DAKOTA	771	4.5	1.0	(2.5, 6.4)
OHIO	1311	5.4	0.9	(3.6, 7.2)
OKLAHOMA	1053	7.3	0.9	(5.4, 9.1)
OREGON	803	6.3	1.5	(3.4, 9.3)
PENNSYLVANIA	1679	6.7	0.7	(5.3, 8.0)
RHODE ISLAND	1750	8.8	0.9	(7.1, 10.4)
SOUTH CAROLINA	1375	6.9	1.2	(4.6, 9.2)
SOUTH DAKOTA	946	5.9	1.4	(3.2, 8.6)
TENNESSEE	1327	7.1	0.9	(5.4, 8.8)
TEXAS	2646	7.8	0.7	(6.4, 9.1)
UTAH	1223	9.7	1.3	(7.3, 12.2)
VERMONT	1388	6.0	0.9	(4.2, 7.8)
VIRGINIA	1544	5.6	0.9	(3.8, 7.3)
WASHINGTON	1547	5.2	0.7	(3.8, 6.6)
WEST VIRGINIA	1045	6.7	1.0	(4.7, 8.6)
WISCONSIN	1012	5.4	0.9	(3.5, 7.2)
WYOMING	997	2.7	0.6	(1.6, 3.8)
PUERTO RICO	1389	19.7	1.4	(17.0, 22.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 5.6 RANGE = 2.7-19.7

* AMONG WOMEN WITH INTACT CERVIX, AGE 18 AND OVER

June 18, 1999

TABLE 36.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 NEVER HAD PAP SMEAR, FOR WOMEN AGE 18 AND OLDER WITH INTACT CERVIX *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

90

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	3.6	5.7	2.1
ALASKA	4.1	3.3	-0.8
ARIZONA	17.1	15.1	-2.0
ARKANSAS	9.2	7.9	-1.3
COLORADO	4.2	4.5	0.3
CONNECTICUT	7.8	7.3	-0.5
DELAWARE	3.5	5.1	1.6
DISTRICT OF CO	3.7	3.0	-0.7
FLORIDA	7.4	5.2	-2.2
GEORGIA	3.5	4.3	0.8
HAWAII	4.7	5.5	0.8
IDAHO	4.3	4.8	0.5
ILLINOIS	6.6	5.6	-1.0
INDIANA	3.8	6.3	2.6
IOWA	5.4	5.2	-0.2
KANSAS	5.9	6.5	0.6
KENTUCKY	7.5	5.0	-2.5
LOUISIANA	7.0	6.4	-0.6
MAINE	4.3	4.7	0.5
MARYLAND	5.6	4.8	-0.7
MASSACHUSETTS	6.3	6.2	-0.2
MICHIGAN	5.5	4.6	-0.9
MINNESOTA	7.6	6.0	-1.5
MISSISSIPPI	4.2	6.7	2.5
MISSOURI	4.1	4.3	0.1
MONTANA	3.6	3.5	-0.1
NEBRASKA	5.1	5.4	0.3
NEVADA	5.0	4.7	-0.2
NEW HAMPSHIRE	5.1	5.6	0.5
NEW JERSEY	8.5	10.3	1.8
NEW MEXICO	7.5	6.7	-0.8
NEW YORK	6.9	7.4	0.5
NORTH CAROLINA	5.1	4.6	-0.6
NORTH DAKOTA	6.1	4.5	-1.7
OHIO	4.9	5.4	0.5
OKLAHOMA	4.9	7.3	2.4
OREGON	4.6	6.3	1.8
PENNSYLVANIA	7.5	6.7	-0.8
RHODE ISLAND	5.1	8.8	3.7
SOUTH CAROLINA	3.7	6.9	3.2
SOUTH DAKOTA	4.9	5.9	1.0
TENNESSEE	5.2	7.1	1.9
TEXAS	8.0	7.8	-0.2
UTAH	9.0	9.7	0.7
VERMONT	6.3	6.0	-0.3
VIRGINIA	5.3	5.6	0.3
WASHINGTON	3.8	5.2	1.4
WEST VIRGINIA	6.6	6.7	0.0
WISCONSIN	4.9	5.4	0.4
WYOMING	5.0	2.7	-2.3
PUERTO RICO	18.2	19.7	1.5

* AMONG WOMEN WITH INTACT CERVIX, AGE 18 AND OVER

June 18, 1999

TABLE 37.1 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 ESTIMATED PREVALENCE DID NOT HAVE PAP SMEAR WITHIN PAST 3 YEARS FOR WOMEN WITH INTACT CERVIX *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

91

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	951	14.7	1.4	(12.0, 17.5)
ALASKA	896	10.5	1.5	(7.6, 13.3)
ARIZONA	835	18.3	2.4	(13.5, 23.0)
ARKANSAS	1201	21.1	1.5	(18.2, 24.0)
COLORADO	743	13.0	1.5	(10.0, 15.9)
CONNECTICUT	1483	13.8	1.2	(11.5, 16.1)
DELAWARE	1200	12.9	1.4	(10.3, 15.6)
DISTRICT OF CO	747	6.1	1.0	(4.1, 8.0)
FLORIDA	2028	14.1	0.9	(12.3, 15.8)
GEORGIA	1064	13.9	1.2	(11.4, 16.3)
HAWAII	992	13.7	1.5	(10.8, 16.7)
IDAHO	2011	17.5	1.0	(15.5, 19.4)
ILLINOIS	1305	15.5	1.2	(13.1, 17.8)
INDIANA	1120	19.1	1.3	(16.5, 21.7)
IOWA	1672	15.5	1.1	(13.4, 17.7)
KANSAS	1626	14.4	1.0	(12.4, 16.4)
KENTUCKY	1686	15.7	1.0	(13.7, 17.7)
LOUISIANA	671	15.2	1.5	(12.2, 18.1)
MAINE	734	15.0	1.6	(11.8, 18.1)
MARYLAND	1560	10.8	1.1	(8.7, 12.8)
MASSACHUSETTS	2420	12.3	1.0	(10.4, 14.1)
MICHIGAN	1157	11.9	1.0	(10.0, 13.9)
MINNESOTA	1842	15.0	0.9	(13.1, 16.8)
MISSISSIPPI	932	15.6	1.4	(12.9, 18.3)
MISSOURI	1668	14.6	1.1	(12.5, 16.7)
MONTANA	761	19.4	1.6	(16.3, 22.5)
NEBRASKA	1185	17.6	1.2	(15.2, 19.9)
NEVADA	770	14.9	1.8	(11.5, 18.3)
NEW HAMPSHIRE	709	15.8	1.5	(12.8, 18.8)
NEW JERSEY	1207	18.6	1.4	(16.0, 21.3)
NEW MEXICO	1506	17.4	1.1	(15.2, 19.6)
NEW YORK	1224	16.2	1.2	(13.8, 18.5)
NORTH CAROLINA	1073	12.3	1.3	(9.8, 14.9)
NORTH DAKOTA	767	14.5	1.5	(11.6, 17.3)
OHIO	1306	15.8	1.4	(13.0, 18.6)
OKLAHOMA	1046	15.2	1.2	(12.8, 17.6)
OREGON	798	16.3	1.8	(12.8, 19.7)
PENNSYLVANIA	1669	16.2	1.0	(14.2, 18.2)
RHODE ISLAND	1734	14.8	1.0	(12.8, 16.8)
SOUTH CAROLINA	1365	15.0	1.4	(12.3, 17.7)
SOUTH DAKOTA	934	13.7	1.5	(10.7, 16.7)
TENNESSEE	1324	13.9	1.2	(11.6, 16.2)
TEXAS	2637	18.5	1.0	(16.4, 20.5)
UTAH	1220	21.6	1.7	(18.3, 24.8)
VERMONT	1371	14.0	1.1	(11.8, 16.2)
VIRGINIA	1530	15.1	1.5	(12.2, 18.0)
WASHINGTON	1542	14.6	1.1	(12.4, 16.8)
WEST VIRGINIA	1043	19.0	1.3	(16.4, 21.6)
WISCONSIN	1005	16.6	1.4	(13.7, 19.4)
WYOMING	991	18.0	1.3	(15.4, 20.5)
PUERTO RICO	1375	32.4	1.5	(29.5, 35.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 15.1 RANGE = 6.1-32.4

* AMONG WOMEN WITH INTACT CERVIX, AGE 18 AND OVER

June 18, 1999

TABLE 37.2 1998 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA
 PREVALENCE DIFFERENCES FROM 1997 TO 1998
 DID NOT HAVE PAP SMEAR WITHIN PAST 3 YEARS FOR WOMEN WITH INTACT CERVIX *
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

92

PARTICIPANT	1997 PERCENT	1998 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.0	14.7	0.7
ALASKA	9.7	10.5	0.7
ARIZONA	20.7	18.3	-2.4
ARKANSAS	21.8	21.1	-0.7
COLORADO	12.4	13.0	0.6
CONNECTICUT	17.0	13.8	-3.2
DELAWARE	10.2	12.9	2.8
DISTRICT OF CO	7.9	6.1	-1.8
FLORIDA	15.3	14.1	-1.2
GEORGIA	7.7	13.9	6.1
HAWAII	13.2	13.7	0.6
IDAHO	18.4	17.5	-1.0
ILLINOIS	17.3	15.5	-1.8
INDIANA	15.5	19.1	3.5
IOWA	19.5	15.5	-4.0
KANSAS	14.1	14.4	0.3
KENTUCKY	18.3	15.7	-2.6
LOUISIANA	16.2	15.2	-1.1
MAINE	11.9	15.0	3.1
MARYLAND	11.3	10.8	-0.5
MASSACHUSETTS	11.6	12.3	0.6
MICHIGAN	13.7	11.9	-1.8
MINNESOTA	16.6	15.0	-1.7
MISSISSIPPI	14.1	15.6	1.5
MISSOURI	16.1	14.6	-1.6
MONTANA	15.4	19.4	4.0
NEBRASKA	15.9	17.6	1.6
NEVADA	14.4	14.9	0.5
NEW HAMPSHIRE	13.0	15.8	2.8
NEW JERSEY	18.7	18.6	-0.1
NEW MEXICO	19.4	17.4	-2.1
NEW YORK	13.0	16.2	3.1
NORTH CAROLINA	12.7	12.3	-0.3
NORTH DAKOTA	17.6	14.5	-3.2
OHIO	12.7	15.8	3.1
OKLAHOMA	15.6	15.2	-0.4
OREGON	14.0	16.3	2.3
PENNSYLVANIA	17.7	16.2	-1.5
RHODE ISLAND	12.6	14.8	2.2
SOUTH CAROLINA	10.3	15.0	4.7
SOUTH DAKOTA	15.0	13.7	-1.4
TENNESSEE	12.3	13.9	1.7
TEXAS	19.1	18.5	-0.7
UTAH	21.5	21.6	0.1
VERMONT	16.0	14.0	-2.0
VIRGINIA	12.1	15.1	3.0
WASHINGTON	13.0	14.6	1.6
WEST VIRGINIA	21.1	19.0	-2.1
WISCONSIN	16.2	16.6	0.4
WYOMING	16.3	18.0	1.7
PUERTO RICO	28.2	32.4	4.2

* AMONG WOMEN WITH INTACT CERVIX, AGE 18 AND OVER

June 18, 1999