

June 23, 2000

Dear Colleague:

Information related to the 1999 BRFSS summary prevalence report of selected risk factors and preventive health measures from the core and modules is attached.

This report consists of four parts:

An introduction that contains:

An overview of the 1999 summary prevalence report.

An explanation of the tables for 1999 Behavioral Risk Factor Surveillance System (BRFSS) prevalence data.

Definitions of BRFSS risk factors and preventive health measures. When appropriate, Year 2000 Objectives are identified.

Analysis of selected questions from the 1999 BRFSS questionnaire. The set of analyses is divided into three sections:

Section I - selected risk factors and preventive health measures in the BRFSS core.

Section II - selected risk factors and preventive health measures from modules used by at least 13 states

Section III - preventive health measures that concern women only.

As soon as your schedule permits, please examine the enclosed materials closely.

Sincerely yours,

David E. Nelson, M.D.,M.P.H.  
Chief, Behavioral Surveillance Branch  
Division of Adult and Community Health  
National Center for Chronic Disease  
Prevention and Health Promotion

# **1999 BRFSS SUMMARY PREVALENCE REPORT**

BEHAVIORAL SURVEILLANCE BRANCH  
DIVISION OF ADULT AND COMMUNITY HEALTH  
NATIONAL CENTER FOR CHRONIC DISEASE PREVENTION  
AND HEALTH PROMOTION  
CENTERS FOR DISEASE CONTROL AND PREVENTION

# TABLE OF CONTENTS

Overview of the 1999 Summary Report.....	ix
Explanation of Tables.....	x
Definitions of Terms and Risk Factors.....	xii
Limitations of the Data .....	xvii

## SECTION I

### PREVALENCE REPORT FOR CORE QUESTIONS

#### General Health Status

TABLE 1.1	Self reported health status: Fair or poor.....	1
1.2	Self reported health status: Fair or poor-by sex .....	2
1.3	Self reported health status: Fair or poor, change 1998-1999.....	3

#### Quality of Life: Physical and Mental Health

TABLE 2.1	Physical health not good.....	4
2.2	Physical health not good-by sex.....	5
2.3	Physical health not good, change 1998-1999.....	6
TABLE 3.1	Mental health not good.....	7
3.2	Mental health not good-by sex.....	8
3.3	Mental health not good, change 1998-1999.....	9
TABLE 4.1	Activities limited.....	10
4.2	Activities limited-by sex.....	11
4.3	Activities limited, change 1998-1999.....	12

#### Health Insurance

TABLE 5.1	No health insurance, age 18 and over.....	13
5.2	No health insurance, age 18 and over-by sex.....	14
5.3	No health insurance, age 18 and over, change 1998-1999.....	15

TABLE 6.1	No health insurance, age 18-64 years.....	16
6.2	No health insurance, age 18-64 year-by sex.....	17
6.3	No health insurance, age 18-64 years, change 1998-1999.....	18

**Diabetes Awareness**

TABLE 7.1	Ever told you have diabetes.....	19
7.2	Ever told you have diabetes-by sex.....	20
7.3	Ever told you have diabetes, change 1998-1999.....	21

**Hypertension Awareness and Screening**

TABLE 8.1	Ever told blood pressure high.....	22
8.2	Ever told blood pressure high-by sex.....	23
8.3	Ever told blood pressure high change 1997-1999.....	24

TABLE 9.1	Blood pressure not checked within past two years.....	25
9.2	Blood pressure not checked within past two years-by sex.....	26
9.3	Blood pressure not checked within past two years, change 1997-1999.....	27

**Cholesterol Screening and Awareness**

TABLE 10.1	Never had cholesterol check.....	28
10.2	Never had cholesterol check-by sex.....	29
10.3	Never had cholesterol check, change 1997-1999.....	30

TABLE 11.1	Cholesterol not checked within past five years.....	31
11.2	Cholesterol not checked within past five years-by sex.....	32
11.3	Cholesterol not checked within past five years, change 1997-1999.....	33

TABLE 12.1	Ever told cholesterol high.....	34
12.2	Ever told cholesterol high-by sex.....	35
12.3	Ever told cholesterol high, change 1997-1999.....	36

**Smoking Status**

TABLE 13.1	Ever smoked.....	37
13.2	Ever smoked-by sex.....	38
13.3	Ever smoked, change 1998-1999.....	39
TABLE 14.1	Current smoker.....	40
14.2	Current smoker-by sex.....	41
14.3	Current smoker, change 1998-1999.....	42

## Alcohol Consumption

TABLE 15.1	Current drinkers.....	43
15.2	Current drinkers-by sex.....	44
15.3	Current drinkers, change 1997-1999.....	45
TABLE 16.1	Binge drinkers .....	46
16.2	Binge drinkers-by sex.....	47
16.3	Binge drinkers, change 1997-1999.....	48
TABLE 17.1	Chronic drinkers.....	49
17.2	Chronic drinkers-by sex.....	50
17.3	Chronic drinkers, change 1997-1999.....	51
TABLE 18.1	Drinking and driving.....	52
18.2	Drinking and driving-by sex.....	53
18.3	Drinking and driving, change 1997-1999.....	54

## Immunization

TABLE 19.1	Did not have flu shot within past 12 months, age 65 and older.....	55
19.2	Did not have flu shot within past 12 months, age 65 and older-by sex.....	56
19.3	Did not have flu shot within past 12 months, age 65 and older, change 1997- 1999.....	57
TABLE 20.1	Never had pneumonia vaccination, age 65 and older.....	58
20.2	Never had pneumonia vaccination, age 65 and older-by sex.....	59
20.3	Never had pneumonia vaccination, age 65 and older, change 1997-1999.....	60

## Colorectal Cancer Screening

TABLE 21.1	Never had sigmoidoscopy, age 50 and older.....	61
21.2	Never had sigmoidoscopy, age 50 and older-by sex.....	62
TABLE 22.1	Did not use home blood stool test kit within past two years, age 50 and older.....	63
22.2	Did not use home blood stool test kit within past two years, age 50 and older- by sex.....	64
22.3	Did not use home blood stool test kit within past two years, age 50 and older- change 1997-1999.....	65

## Overweight

TABLE 23.1	Overweight based on body mass index (BMI).....	66
23.2	Overweight based on body mass index (BMI)-by sex.....	67
23.3	Overweight based on body mass index (BMI), change 1998-1999.....	68
TABLE 24.1	Overweight based on percent of median.....	69
24.2	Overweight based on percent of median-by sex.....	70
24.3	Overweight based on percent of median, change 1998-1999.....	71

## HIV/AIDS

TABLE 25.1	Would not encourage teen condom use, age 18-64.....	72
25.2	Would not encourage teen condom use-by sex.....	73
25.3	Would not encourage teen condom use, change 1998-1999.....	74
TABLE 26.1	Chances of getting HIV medium or high, age 18-64.....	75
26.2	Chances of getting HIV medium or high, age 18-64 -by sex.....	76
26.3	Chances of getting HIV medium or high, change 1998-1999.....	77
TABLE 27.1	Not Tested for HIV, age 18-64.....	78
27.2	Not Tested for HIV, age 18-64 - by sex.....	79
27.3	Not Tested for HIV, age 18-64 - change 1998-1999.....	80

## SECTION II

### PREVALENCE REPORT FOR MODULES

## Diabetes

TABLE 28.1	Age at onset of diabetes.....	81
28.2	Age at onset of diabetes-by sex.....	82
TABLE 29.1	Diabetics using insulin.....	83
29.2	Diabetics using insulin-by sex.....	84
TABLE 30.1	Health professional visits in past year.....	85
30.2	Health professional visits in past year-by sex.....	86

## Smokeless tobacco

TABLE 31.1	Ever used smokeless tobacco.....	87
31.2	Ever used smokeless tobacco, by sex.....	88

TABLE 32.1	Currently uses smokeless tobacco.....	89
32.2	Currently uses smokeless tobacco, by sex.....	90

**ASTHMA**

TABLE 33.1	Currently has asthma.....	91
33.2	Currently has asthma, by sex.....	92

**CARDIOVASCULAR DISEASE**

TABLE 34.1	Ever told you had a heart attack or myocardial infarction.....	93
34.2	Ever told you had a heart attack or myocardial infarction- by sex.....	94

TABLE 35.1	Ever told you had a stroke.....	95
35.2	Ever told you had a stroke- by sex.....	96

**SECTION III**

**PREVALENCE REPORT FOR WOMEN'S HEALTH ISSUES**

**Mammography**

TABLE 36.1	Never had a mammogram, women age 18 and older.....	97
36.2	Never had a mammogram, women age 18 and older, change 1998-1999.....	98
TABLE 37.1	Never had a mammogram, women age 40 and older.....	99
37.2	Never had a mammogram, women age 40 and older, change 1998-1999.....	100
TABLE 38.1	Did not have a mammogram within the past two years, women age 50 and older...101	
38.2	Did not have a mammogram within the past two years, women age 50 and older, change 1998-1999.....	102

**Breast Exam**

TABLE 39.1	Never had a breast exam, women age 18 and older.....	103
39.2	Never had a breast exam, women age 18 and older, change 1998-1999.....	104
TABLE 40.1	Never had a breast exam, women age 40 and older.....	105
40.2	Never had a breast exam, women age 40 and older, change 1998-1999.....	106
TABLE 41.1	Did not have a breast exam within the past two years, women age 50 and older.....	107
41.2	Did not have a breast exam within the past two years, women age 50 and older, change 1998-1999.....	108

## **Mammogram and Breast Exam**

TABLE 42.1	Never had a mammogram and breast exam, women age 40 and older.....	109
42.2	Never had a mammogram and breast exam, women age 40 and older, change 1998-1999.....	110
TABLE 43.1	Did not have a mammogram and breast exam within the past two years, women age 50 and older.....	111
43.2	Did not have a mammogram and breast exam within the past two years, women age 50 and older, change 1998-1999.....	112

## **Cervical Cancer**

TABLE 44.1	Never had pap smear, women age 18 and older.....	113
44.2	Never had pap smear, women age 18 and older, change 1998-1999.....	114
TABLE 45.1	Never had pap smear, women with intact cervix.....	115
45.2	Never had pap smear, women with intact cervix, change 1998-1999.....	116
TABLE 46.1	Did not have pap smear within the past three years, women with intact cervix.....	117
46.2	Did not have pap smear within the past three years, women with intact cervix, change 1998-1999.....	118

## OVERVIEW OF THE 1999 SUMMARY REPORT

The report is divided into three sections. *Section I* displays tables for selected variables in the core. In *Section I*, there are three tables for each risk factor or preventive measure unless otherwise noted. Estimated prevalence and relevant statistics are provided for the total population sampled and by sex. Prevalence estimates for both 1998 and 1999, or where appropriate 1997 and 1999, and the percentage point difference between the estimates are given for variables in the core.

*Section II* shows tables for selected variables in modules used by at least 13 states. *Section II* contains two tables for each risk factor or preventive measure of interest in the modules. These tables show the estimated prevalence and relevant statistics for the total sampled population and both sex groups who responded to selected questions in the optional modules.

*Section III* contains tables for preventive health measures that are related to women's health issues. Estimated prevalence and relevant statistics are provided for the total population sampled. Prevalence estimates for both 1998 and 1999 and the percentage point difference between them are given.

Three tables (Tables 1.1-1.3) from Section I will be used as examples to explain the information contained in this report. Each table describes a behavioral risk factor or preventive health measure from the Behavioral Risk Factor Surveillance System (BRFSS) instrument.

Table 1.1 (page 1) shows the estimated prevalence and relevant statistics for each participant. (Participants refer to all states, and the District of Columbia and territories of the U.S.). The table shows the estimated prevalence and relevant statistics for "fair or poor health status" in the population surveyed by each BRFSS participant. This prevalence estimate excludes respondents with unknown information on self reported health status or respondents choosing not to answer the question.

The columns in Table 1.1 provide the following information:

COLUMN HEADING

CONTENTS

<u>Participant</u>	Name of the participant. Participants refers to all states including the District of Columbia and territories of the United States.
<u>Sample size</u>	The number of individuals who responded to the specific BRFSS question. These are unweighted numbers. These numbers <u>exclude</u> individuals who reported "not sure, don't know", those who refused to answer the question, or had no response coded (missing or skipped).
<u>Percent/Mean</u>	An estimate of the percent (or prevalence or mean number) of residents who report the behavioral risk factor of interest. This is weighted to the participant's population to estimate the true prevalence or mean in the population.
<u>Standard error</u>	A measure of the amount of variability that exists in the calculation of the prevalence estimate.
<u>95% confidence interval</u>	A range which contains the true population prevalence estimate 95% of the time, if repeated sampling of the population is performed.
<u>SUMMARY STATISTICS</u> are given at the bottom of the table:	
<u>Number of participants</u>	The total number of participants who asked question(s) regarding the respective "risk factor".
<u>Median</u>	The middle value of all estimated prevalences.
<u>Range</u>	The lowest prevalence and the highest prevalence among all participants.

For example, in Table 1.1, there were 2092 individuals in the state of Alabama who responded to the "self reported health status" question. Of those responding, the estimated prevalence of Alabama's population who reported that they had fair or poor health status is 18.4%. This prevalence estimate has a standard error of 0.9. The 95% confidence interval for this estimate is (16.6, 20.2). Ninety-five percent (95%) of confidence intervals generated under repeated sampling will contain the true population prevalence of Alabama's population who "had fair or poor health status".

Table 1.2 (page 2) shows the estimated prevalence and relevant statistics by sex for each state participant. The same statistics are displayed in Table 1.2 as in Table 1.1, except that Table 1.2 shows the estimated prevalence of “ fair to poor health “ for males and females separately. This side by side display of the tables permits each participant to compare estimates for males and females in their respective state.

Table 1.3 (page 3) shows the percent difference between the prevalence estimates of poor health in 1998 and 1999 survey years. A negative change indicates a decrease in prevalence from 1998 to 1999, while a positive change indicates an increase in prevalence between the two years. This difference gives state health officials a quick method to examine changes that may have occurred in their respective state. Some of these changes may have resulted from legislative actions (or inactions) in the state, programs funded, or programs that came into existence because of results that states may have shown from the BRFSS. Since changes between two years may be attributable to many factors, conclusions drawn from this table should be interpreted with extreme caution.

**Sample Size.** Sample sizes vary by question. Some estimates may be based on responses from less than 50 respondents. Interpreting estimates that are based on a small number of respondents can mislead the reader into believing that a given finding is much more precise than it actually is. The BRFSS recommends not interpreting percentages where the denominator is based upon fewer than 50 unweighted respondents (sample size).

# **DEFINITIONS OF TERMS AND RISK FACTORS IN THIS REPORT**

## **AIDS KNOWLEDGE/ATTITUDES**

**HIV TEST** - Respondents, age 18-64, who have never had a test for HIV infection.

**CHANCES OF GETTING HIV** - Respondents, age 18-64, who believe that their chances of getting infected with HIV, the virus that causes AIDS are medium or high.

**TEEN CONDOM USE** - Respondents, age 18-64, who would not encourage sexually active teenagers to use a condom.

## **ASTHMA**

**CURRENTLY HAS ASTHMA** - Respondents who report they currently have asthma.

## **BREAST EXAM**

**BREAST EXAM (1)**- Female respondents, age 18 and older, who report they never had a breast exam by a doctor.

**BREAST EXAM (2)**- Female respondents, age 40 and older, who report they never had a breast exam by a doctor.

**BREAST EXAM WITHIN 2 YEARS** - Female respondents, age 50 and older, who report that they did not have a breast exam within the past two years.

## **CARDIOVASCULAR DISEASE**

**HEART ATTACK OR MYOCARDIAL INFARCTION** - Respondents who report they were told by a doctor they had a heart attack or myocardial infarction.

**STROKE** - Respondents who report they were told by a doctor they had a stroke.

## **CERVICAL CANCER**

**PAP SMEAR (1)**- Female respondents, age 18 and older who report they never had a pap smear. Year 2000 Objective 16.12. - Target  $\geq 95\%$  who have ever had a pap smear.

**PAP SMEAR, WITH UTERINE CERVIX (2)**- Female respondents, with uterine cervix, age 18 and older, who report they never had a pap smear.

**PAP SMEAR WITHIN 3 YEARS** - Female respondents, with uterine cervix, age 18 and older, who report they did not have a pap smear within the last three years.

### **CHOLESTEROL SCREENING & AWARENESS**

**CHOLESTEROL SCREENING (1)** - Respondents who never had their cholesterol checked.

**CHOLESTEROL SCREENING (2)** - Respondents who did not have their cholesterol checked within the past five years. Year 2000 Objective 15.14.- Target  $\geq 75\%$  who had their cholesterol checked within the past five years.

**CHOLESTEROL AWARENESS** - Respondents who report their cholesterol has been checked and they have been told their cholesterol is high by a health professional.

**CLINICAL BREAST EXAM** See BREAST EXAM

### **COLORECTAL CANCER SCREENING**

**SIGMOIDOSCOPY** - Respondents, age 50 and over, who report they never had a sigmoidoscopy. Year 2000 Objective 16.13.- Target  $\geq 40\%$  age 50 and over who ever had a sigmoidoscopy.

**BLOOD STOOL TEST** - Respondents, age 50 and over, who reported they did not use a home blood stool kit in the past two years.

### **DIABETES**

**AGE AT ONSET** - Mean age when respondent was told that they have diabetes, among those who report they were told by a doctor that they have diabetes.

**DIABETES AWARENESS** - Respondents who report they were told by a doctor that they have diabetes.

**NOW TAKING INSULIN** - Respondents who are now taking insulin, among those who report they were told by a doctor that they have diabetes.

**NUMBER OF TIMES SEEN HEALTH PROFESSIONAL** - Mean number of times respondents report they have seen a doctor, nurse, or other health professional for their diabetes in the past year, among those who report they were told by a doctor that they have diabetes.

### **DRINKING STATUS**

**CURRENT DRINKERS** - Respondents who report they had alcoholic beverages in the

past month.

**BINGE DRINKING (ACUTE DRINKING)** - Respondents who report having five or more drinks on an occasion, one or more times in the past month.

**CHRONIC DRINKING** - Respondents who report drinking 60 or more alcohol beverages a month.

**DRINKING & DRIVING** - Respondents who report having driven after having too much to drink, one or more times in the past month.

### **FRUITS AND VEGETABLES**

**FRUITS AND VEGETABLES** - Respondents who report they are not consuming five or more servings of fruits and vegetables a day. Year 2000 Objective 2.6. Target  $\geq 50\%$  who are consuming five or more servings of fruits or vegetables a day.

### **HEALTH INSURANCE**

**HEALTH INSURANCE (1)**- Respondents age 18 and older who report they have no health care coverage.

**HEALTH INSURANCE (2)**- Respondents age 18-64 years who report they have no health care coverage.

### **HEALTH STATUS**

**ACTIVITIES LIMITED** - Mean number of days in past 30 days activities were limited among respondents reporting days when physical health or mental health was not good.

**MENTAL HEALTH NOT GOOD** - Mean number of days in past 30 days mental health was not good as reported by respondents.

**PHYSICAL HEALTH NOT GOOD** - Mean number of days in past 30 days physical health was not good as reported by respondents.

**SELF REPORTED HEALTH STATUS** - Respondents who report having a general health status of fair or poor.

### **HYPERTENSION**

**HYPERTENSION AWARENESS** - Respondents who report that their blood pressure has been checked and they have been told they have high blood pressure.

**HYPERTENSION SCREENING** - Respondents who report their blood pressure has not been checked within the past two years.

## **IMMUNIZATION**

**FLU SHOT** - Respondents, age 65 and older, who report they did not have a flu shot within the past 12 months. Year 2000 Objective 20.11 - Target  $\geq 80\%$  who had a flu shot in the past 12 months.

**PNEUMONIA VACCINATION** - Respondents, age 65 and older, who report they never had a pneumonia vaccination. Year 2000 Objective 20.11 - Target  $\geq 80\%$  who have had a pneumonia vaccination.

## **MAMMOGRAPHY**

**MAMMOGRAM (1)** - Female respondents age 18 and older, who report that they have never had a mammogram.

**MAMMOGRAM (2)** - Female respondents, age 40 and older, who report that they have never had a mammogram.

**MAMMOGRAM WITHIN 2 YEARS** - Female respondents, age 50 and older, who report that they did not have a mammogram within the past two years.

## **MAMMOGRAPHY AND BREAST EXAM**

**MAMMOGRAM & BREAST EXAM** - Female respondents, age 40 and older, who report that they have never had a mammogram and a breast exam.

**MAMMOGRAM & BREAST EXAM WITHIN 2 YEARS** - Female respondents, age 50 and older, who report that they did not have a mammogram and a breast exam within the last two years. Year 2000 Objective 16.11 - Target  $\geq 60\%$  women, age 50 and over who had a mammogram and a breast exam in the preceding 1-2 years.

## **OVERWEIGHT**

**OVERWEIGHT: BY MEDIAN** - Respondents at or above 120% of their ideal weight. Ideal weight is defined as the mid-value of a median frame person from the 1959 Metropolitan Life Insurance Corporation height-weight tables.

**OVERWEIGHT: BY BODY MASS INDEX** - Females with body mass index (BMI)  $\geq 27.3$  and males with BMI  $\geq 27.8$ . BMI is defined as weight in kilograms divided by height in meters squared ( $w/h^{**2}$ ). Year 2000 Objective 2.3 - Target  $\leq 20\%$  at or above

the BMI level. *This should be used with extreme caution since people tend to under report their weight. BRFSS may underestimate the prevalence of overweight.*

**PAP SMEAR** See CERVICAL CANCER

### **PHYSICAL INACTIVITY**

**NO LEISURE TIME PHYSICAL ACTIVITY** - Respondents who report no leisure-time physical activity during the past month. Year 2000 Objective 1.5 - Target  $\leq 15\%$  with no-leisure time activity.

**REGULAR AND SUSTAINED PHYSICAL ACTIVITY** - Respondents who report no physical activity defined as: 5 or more sessions per week, 30 minutes or more per session, regardless of intensity. Year 2000 Objective 1.3 - Target  $\geq 30\%$  who engage in physical activity for at least 30 minutes per day.

**REGULAR AND VIGOROUS PHYSICAL ACTIVITY** - Respondents who report no physical activity or a pair of activities defined as: 3 or more sessions per week, 20 minutes or more per session, at 50% or more capacity. Year 2000 Objective 1.4 - Target  $\geq 20\%$  who engage in physical activity 3 or more days per week for 20 or more minutes.

**SIGMOIDOSCOPY** - See Colorectal Cancer Screening

### **SMOKELESS TOBACCO**

**EVER USED** - Respondents who have ever used or tried smokeless tobacco.

**CURRENT USE** - Respondents who report that they currently use smokeless tobacco.

### **SMOKING STATUS**

**EVER SMOKED** - Respondents who have ever smoked 100 cigarettes in their lifetime.

**CURRENT SMOKER** - Respondents who have ever smoked 100 cigarettes in their lifetime and reported smoking every day or some days. Year 2000 Objective 3.4 - Target  $\leq 15\%$  who smoke cigarettes..

### **WEIGHT CONTROL**

**CALORIES AND PHYSICAL ACTIVITY** - Respondents at risk on BMI who report that they are not either eating fewer calories or less fat and not using physical activity or exercise to lose or maintain weight.

## **LIMITATIONS OF THE DATA**

This report consolidates the most current data on selected risk factors and preventive health measures for states, the District of Columbia and Puerto Rico. The information was obtained from the data files of individual states. Therefore, while all participants use telephone surveys to collect the data, the data in this report vary somewhat with respect to data collection, sampling design and sample size.

All states conduct interviews using a computer-assisted telephone interviewing (CATI) software program. All states use probability-based methods to select their sample. The BRFSS standard for sample designs is that sample records must be justifiable as a probability sample of all households with telephones in the state. All participating areas except California, and Hawaii met this criterion in 1999. For data year 1999, two states, Tennessee and Minnesota used the Mitofsky-Waksberg method of sampling. Forty-six states and the District of Columbia used a disproportionate stratified sample (DSS) design. Puerto Rico used a simple random sample design. The two remaining states— California, and Hawaii—used other designs.

### **California**

California, in 1999, used a sample design in which only phone numbers from hundred blocks with one or more listed household numbers were included in the sampling frame. Such hundred blocks are estimated to contain 97.8% of all household numbers in California. Numbers from this frame were selected with an equal probability of selection.

### **Hawaii**

Hawaii uses a sampling frame obtained from the state's telephone company. The frame purportedly contains all working prefixes in Hawaii. The frame is divided into six strata, corresponding to the major islands of Hawaii. The working prefixes are used to randomly generate telephone numbers with a probability proportional to the number of known household numbers in the hundred block to which a number belongs. Telephone numbers from hundred blocks with no known household numbers have a zero probability of being included in the sample. The percentage of telephone households excluded from the sampling frame is unknown. The numbers generated are subsequently screened by GTE Hawaii, who screens out all but the listed and unlisted household numbers in the sample.

### **Other areas**

In a few states, a portion of sample records intended for use during one month may have been completed in another month. This deviation should only affect analyses based on monthly, rather than annual, data. Illinois completed their 1999 survey in January 2000. One hundred eighteen records have a January interview date.

### **B. Other 1999 limitations of the data**

In addition to departing from the standard for sample designs, California modified the wording of mammography, breast exam and Papanicolaou (PAP) smear questions, and are therefore

excluded from those tables in the report. California also asked the HIV/AIDS section questions to persons 18-45 years of age rather than to those 18-64 years of age as specified, and are therefore not included in the tables for this topic. Illinois used a skip pattern in the HIV/AIDS section which affected the HIV testing questions and are therefore excluded from Table 27.

Although almost all states collect data monthly, the annual number of interviews range from 1267 in the District of Columbia to 7543 in Kentucky. Estimates for some states are therefore based upon smaller sample sizes, and for rare events (e.g. diabetes), may yield unstable estimates. Some items, including drinking and driving and obesity, are affected by seasonality, and estimates for these items may be biased for subpopulations.

The 1998 estimates presented for Puerto Rico in the 1999 summary report do not match the estimates presented in the 1998 Summary Prevalence Report. The original 1998 estimates were based on mixed mode (personal interview and telephone survey data). The 1999 estimates were calculated using telephone data only.

All data collection systems are subject to error, and records may be incomplete or contain inaccurate information. Respondents may not remember essential information, a question may not mean the same thing to different respondents, and some individuals may not respond at all. It is not always possible to measure the magnitude of these errors or their impact on the data. The user must make his or her own evaluation of the data. Overall estimates generally have relatively small sampling errors, but estimates for certain population subgroups may be based on small numbers and have relatively large sampling errors. When the number of events is small and the probability of such an event is small, considerable caution must be observed in interpreting the estimates and/or differences between groups and areas.

TABLE 1.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR SELF REPORTED HEALTH STATUS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2092	18.4	0.9	( 16.6, 20.2)
ALASKA	2045	10.7	0.9	( 8.8, 12.5)
ARIZONA	1743	8.4	0.8	( 6.8, 10.1)
ARKANSAS	2993	19.7	0.8	( 18.1, 21.3)
CALIFORNIA	4302	15.8	0.7	( 14.5, 17.1)
COLORADO	3053	11.5	0.8	( 9.9, 13.1)
CONNECTICUT	3507	11.6	0.7	( 10.2, 13.0)
DELAWARE	2550	12.4	0.9	( 10.6, 14.2)
DISTRICT OF CO	1264	13.0	1.0	( 11.1, 15.0)
FLORIDA	5166	15.2	0.6	( 13.9, 16.5)
GEORGIA	2266	15.0	0.9	( 13.3, 16.7)
HAWAII	2156	14.3	1.0	( 12.3, 16.3)
IDAHO	4939	12.9	0.6	( 11.8, 14.0)
ILLINOIS	2909	14.7	0.8	( 13.1, 16.3)
INDIANA	2441	12.8	1.1	( 10.8, 14.9)
IOWA	3608	12.1	0.6	( 11.0, 13.3)
KANSAS	3863	12.9	0.6	( 11.7, 14.0)
KENTUCKY	7528	21.6	0.6	( 20.4, 22.8)
LOUISIANA	1663	16.9	1.0	( 14.9, 18.9)
MAINE	1670	12.9	0.9	( 11.1, 14.6)
MARYLAND	3953	14.2	0.7	( 12.8, 15.6)
MASSACHUSETTS	5011	11.6	0.6	( 10.5, 12.7)
MICHIGAN	2521	11.5	0.7	( 10.2, 12.9)
MINNESOTA	5337	10.0	0.4	( 9.1, 10.8)
MISSISSIPPI	2172	20.9	1.0	( 18.8, 22.9)
MISSOURI	4271	15.7	0.7	( 14.3, 17.2)
MONTANA	1792	10.9	0.8	( 9.4, 12.4)
NEBRASKA	2819	12.4	0.7	( 11.1, 13.7)
NEVADA	2186	13.8	1.1	( 11.6, 16.0)
NEW HAMPSHIRE	1246	10.6	1.0	( 8.7, 12.5)
NEW JERSEY	2815	12.7	0.7	( 11.2, 14.1)
NEW MEXICO	3487	16.9	0.7	( 15.4, 18.3)
NEW YORK	2641	13.8	0.8	( 12.2, 15.3)
NORTH CAROLINA	2441	17.9	0.9	( 16.1, 19.7)
NORTH DAKOTA	1979	12.2	0.8	( 10.7, 13.7)
OHIO	1633	13.8	0.9	( 11.9, 15.6)
OKLAHOMA	2964	17.4	0.8	( 15.8, 19.0)
OREGON	1802	13.7	0.9	( 11.9, 15.5)
PENNSYLVANIA	3552	13.7	0.6	( 12.5, 15.0)
RHODE ISLAND	3992	12.6	0.6	( 11.4, 13.8)
SOUTH CAROLINA	3461	13.9	0.7	( 12.6, 15.2)
SOUTH DAKOTA	4141	13.1	0.6	( 12.0, 14.2)
TENNESSEE	3009	19.9	0.8	( 18.3, 21.6)
TEXAS	4970	19.2	0.7	( 17.7, 20.6)
UTAH	3192	10.0	0.7	( 8.7, 11.4)
VERMONT	3156	9.9	0.6	( 8.7, 11.0)
VIRGINIA	3488	11.7	0.7	( 10.3, 13.0)
WASHINGTON	3601	12.0	0.7	( 10.5, 13.4)
WEST VIRGINIA	2586	24.0	0.9	( 22.1, 25.8)
WISCONSIN	2174	11.9	0.8	( 10.3, 13.5)
WYOMING	2401	11.3	0.7	( 9.9, 12.8)
PUERTO RICO	3049	33.0	1.0	( 31.1, 34.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 13.1 RANGE = 8.4-33.0

\* HEALTH STATUS IS FAIR OR POOR

June 20, 2000



TABLE 1.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999  
 SELF REPORTED HEALTH STATUS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	21.3	18.4	-2.9
ALASKA	11.0	10.7	-0.3
ARIZONA	10.3	8.4	-1.8
ARKANSAS	20.4	19.7	-0.7
CALIFORNIA	14.6	15.8	1.2
COLORADO	11.6	11.5	-0.1
CONNECTICUT	11.7	11.6	-0.1
DELAWARE	12.8	12.4	-0.4
DISTRICT OF CO	12.4	13.0	0.6
FLORIDA	15.5	15.2	-0.3
GEORGIA	15.9	15.0	-0.9
HAWAII	12.3	14.3	2.1
IDAHO	12.2	12.9	0.6
ILLINOIS	12.5	14.7	2.2
INDIANA	13.3	12.8	-0.4
IOWA	11.2	12.1	0.9
KANSAS	12.0	12.9	0.9
KENTUCKY	21.9	21.6	-0.2
LOUISIANA	16.1	16.9	0.8
MAINE	12.5	12.9	0.3
MARYLAND	13.9	14.2	0.3
MASSACHUSETTS	10.9	11.6	0.8
MICHIGAN	14.5	11.5	-3.0
MINNESOTA	10.4	10.0	-0.4
MISSISSIPPI	21.0	20.9	-0.2
MISSOURI	15.2	15.7	0.5
MONTANA	12.0	10.9	-1.1
NEBRASKA	12.3	12.4	0.1
NEVADA	12.4	13.8	1.4
NEW HAMPSHIRE	9.9	10.6	0.7
NEW JERSEY	11.8	12.7	0.9
NEW MEXICO	15.0	16.9	1.9
NEW YORK	13.8	13.8	-0.1
NORTH CAROLINA	16.6	17.9	1.4
NORTH DAKOTA	13.7	12.2	-1.5
OHIO	16.1	13.8	-2.4
OKLAHOMA	12.6	17.4	4.8
OREGON	13.4	13.7	0.3
PENNSYLVANIA	15.1	13.7	-1.4
RHODE ISLAND	13.5	12.6	-0.9
SOUTH CAROLINA	15.2	13.9	-1.3
SOUTH DAKOTA	11.7	13.1	1.4
TENNESSEE	18.2	19.9	1.7
TEXAS	18.6	19.2	0.6
UTAH	10.8	10.0	-0.8
VERMONT	10.0	9.9	-0.2
VIRGINIA	12.3	11.7	-0.6
WASHINGTON	11.1	12.0	0.9
WEST VIRGINIA	23.9	24.0	0.1
WISCONSIN	12.1	11.9	-0.2
WYOMING	12.1	11.3	-0.7
PUERTO RICO	32.5	33.0	0.6

\* HEALTH STATUS IS FAIR OR POOR

June 20, 2000

TABLE 2.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2034	4.0	0.2	( 3.6, 4.4)
ALASKA	2018	2.8	0.2	( 2.3, 3.2)
ARIZONA	1740	0.9	0.1	( 0.7, 1.1)
ARKANSAS	2961	4.0	0.2	( 3.7, 4.3)
CALIFORNIA	4285	3.3	0.1	( 3.0, 3.5)
COLORADO	3030	3.3	0.2	( 2.9, 3.7)
CONNECTICUT	3453	3.2	0.3	( 2.7, 3.7)
DELAWARE	2521	3.3	0.2	( 2.9, 3.7)
DISTRICT OF CO	1240	2.7	0.2	( 2.3, 3.1)
FLORIDA	5095	3.4	0.1	( 3.1, 3.6)
GEORGIA	2222	2.9	0.2	( 2.5, 3.2)
HAWAII	2141	2.4	0.2	( 2.1, 2.8)
IDAHO	4876	3.2	0.1	( 3.0, 3.5)
ILLINOIS	2887	3.2	0.2	( 2.9, 3.6)
INDIANA	2422	3.1	0.2	( 2.6, 3.5)
IOWA	3585	3.4	0.1	( 3.1, 3.7)
KANSAS	3783	2.6	0.1	( 2.4, 2.9)
KENTUCKY	7470	4.1	0.1	( 3.8, 4.3)
LOUISIANA	1613	3.3	0.2	( 2.8, 3.7)
MAINE	1652	3.5	0.2	( 3.1, 3.9)
MARYLAND	3916	3.1	0.2	( 2.8, 3.4)
MASSACHUSETTS	4935	3.2	0.1	( 3.0, 3.5)
MICHIGAN	2491	3.4	0.2	( 3.0, 3.7)
MINNESOTA	5271	2.6	0.1	( 2.4, 2.8)
MISSISSIPPI	2109	3.5	0.2	( 3.1, 3.9)
MISSOURI	4233	3.5	0.2	( 3.2, 3.8)
MONTANA	1770	3.1	0.2	( 2.8, 3.5)
NEBRASKA	2800	3.1	0.2	( 2.8, 3.4)
NEVADA	2170	3.5	0.3	( 3.0, 4.1)
NEW HAMPSHIRE	1234	3.1	0.2	( 2.7, 3.6)
NEW JERSEY	2771	3.2	0.2	( 2.8, 3.5)
NEW MEXICO	3445	3.7	0.2	( 3.4, 4.0)
NEW YORK	2582	3.3	0.2	( 3.0, 3.7)
NORTH CAROLINA	2407	3.7	0.2	( 3.3, 4.1)
NORTH DAKOTA	1975	2.9	0.2	( 2.6, 3.2)
OHIO	1617	3.1	0.2	( 2.7, 3.5)
OKLAHOMA	2935	3.0	0.2	( 2.7, 3.3)
OREGON	1774	3.6	0.2	( 3.2, 4.1)
PENNSYLVANIA	3492	3.2	0.1	( 2.9, 3.4)
RHODE ISLAND	3923	3.0	0.1	( 2.7, 3.2)
SOUTH CAROLINA	3388	3.2	0.2	( 2.9, 3.4)
SOUTH DAKOTA	4106	3.0	0.1	( 2.7, 3.2)
TENNESSEE	2933	3.4	0.2	( 3.0, 3.7)
TEXAS	4909	3.3	0.1	( 3.1, 3.6)
UTAH	3159	3.2	0.2	( 2.9, 3.5)
VERMONT	3101	2.9	0.1	( 2.6, 3.1)
VIRGINIA	3394	2.7	0.1	( 2.4, 2.9)
WASHINGTON	3525	3.5	0.2	( 3.2, 3.8)
WEST VIRGINIA	2572	4.0	0.2	( 3.7, 4.4)
WISCONSIN	2162	3.6	0.2	( 3.2, 4.0)
WYOMING	2387	2.9	0.2	( 2.6, 3.1)
PUERTO RICO	2954	3.8	0.2	( 3.5, 4.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.2 RANGE = 0.9- 4.1

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

June 20, 2000

TABLE 2.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR PHYSICAL HEALTH NOT GOOD\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*				*-----FEMALES-----*			
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	835	3.5	0.3	( 3.0, 4.1)	1199	4.4	0.3	( 3.9, 5.0)
ALASKA	988	2.3	0.3	( 1.7, 2.8)	1030	3.3	0.3	( 2.7, 4.0)
ARIZONA	712	1.0	0.2	( 0.7, 1.4)	1028	0.8	0.1	( 0.5, 1.0)
ARKANSAS	1158	3.4	0.3	( 2.9, 3.9)	1803	4.6	0.2	( 4.1, 5.0)
CALIFORNIA	1827	2.8	0.2	( 2.5, 3.2)	2458	3.7	0.2	( 3.4, 4.1)
COLORADO	1367	3.2	0.3	( 2.6, 3.8)	1663	3.4	0.3	( 2.9, 3.9)
CONNECTICUT	1354	2.7	0.2	( 2.2, 3.2)	2099	3.7	0.4	( 2.9, 4.5)
DELAWARE	1015	3.1	0.3	( 2.4, 3.7)	1506	3.6	0.3	( 3.1, 4.1)
DISTRICT OF CO	452	2.5	0.3	( 1.9, 3.1)	788	2.9	0.3	( 2.4, 3.3)
FLORIDA	2027	2.8	0.2	( 2.4, 3.1)	3068	3.9	0.2	( 3.6, 4.2)
GEORGIA	868	2.4	0.3	( 1.9, 3.0)	1354	3.3	0.2	( 2.9, 3.8)
HAWAII	990	1.8	0.2	( 1.4, 2.2)	1151	3.0	0.3	( 2.5, 3.6)
IDAHO	2072	2.7	0.2	( 2.3, 3.1)	2804	3.8	0.2	( 3.4, 4.1)
ILLINOIS	1105	2.6	0.3	( 2.1, 3.1)	1782	3.8	0.2	( 3.4, 4.2)
INDIANA	1012	2.5	0.3	( 1.9, 3.2)	1410	3.6	0.3	( 3.0, 4.2)
IOWA	1415	2.9	0.2	( 2.4, 3.3)	2170	3.9	0.2	( 3.5, 4.3)
KANSAS	1542	2.2	0.2	( 1.9, 2.6)	2241	3.0	0.2	( 2.7, 3.3)
KENTUCKY	2732	3.8	0.2	( 3.4, 4.2)	4738	4.3	0.2	( 4.0, 4.6)
LOUISIANA	644	2.5	0.3	( 1.9, 3.1)	969	3.9	0.4	( 3.3, 4.6)
MAINE	733	3.3	0.3	( 2.7, 3.9)	919	3.7	0.3	( 3.1, 4.2)
MARYLAND	1576	2.8	0.2	( 2.4, 3.3)	2340	3.4	0.2	( 3.0, 3.8)
MASSACHUSETTS	1979	3.2	0.2	( 2.8, 3.6)	2956	3.2	0.2	( 2.9, 3.5)
MICHIGAN	1031	2.7	0.2	( 2.2, 3.2)	1460	4.0	0.2	( 3.5, 4.4)
MINNESOTA	2416	2.5	0.2	( 2.2, 2.8)	2855	2.8	0.1	( 2.5, 3.0)
MISSISSIPPI	828	3.0	0.3	( 2.5, 3.5)	1281	4.0	0.2	( 3.5, 4.4)
MISSOURI	1709	2.9	0.2	( 2.5, 3.3)	2524	4.1	0.2	( 3.6, 4.5)
MONTANA	758	2.6	0.3	( 2.1, 3.1)	1012	3.6	0.3	( 3.1, 4.1)
NEBRASKA	1118	3.1	0.3	( 2.7, 3.6)	1682	3.1	0.2	( 2.8, 3.5)
NEVADA	1092	2.9	0.3	( 2.3, 3.5)	1078	4.2	0.5	( 3.3, 5.1)
NEW HAMPSHIRE	499	2.7	0.3	( 2.0, 3.3)	735	3.5	0.3	( 2.9, 4.2)
NEW JERSEY	1111	3.1	0.3	( 2.5, 3.6)	1660	3.3	0.2	( 2.8, 3.7)
NEW MEXICO	1521	3.3	0.2	( 2.9, 3.7)	1924	4.1	0.2	( 3.7, 4.5)
NEW YORK	1044	3.2	0.3	( 2.7, 3.7)	1538	3.5	0.2	( 3.1, 3.9)
NORTH CAROLINA	955	3.3	0.3	( 2.7, 3.9)	1452	4.1	0.3	( 3.6, 4.6)
NORTH DAKOTA	859	2.7	0.2	( 2.2, 3.2)	1116	3.1	0.2	( 2.6, 3.5)
OHIO	631	2.7	0.3	( 2.1, 3.3)	986	3.4	0.3	( 2.9, 4.0)
OKLAHOMA	1147	2.7	0.2	( 2.2, 3.1)	1788	3.3	0.2	( 2.9, 3.6)
OREGON	760	3.3	0.3	( 2.7, 3.9)	1014	4.0	0.3	( 3.4, 4.5)
PENNSYLVANIA	1404	2.8	0.2	( 2.4, 3.2)	2088	3.5	0.2	( 3.2, 3.9)
RHODE ISLAND	1536	2.6	0.2	( 2.2, 3.0)	2387	3.3	0.2	( 2.9, 3.6)
SOUTH CAROLINA	1376	2.7	0.2	( 2.3, 3.1)	2012	3.6	0.2	( 3.2, 4.0)
SOUTH DAKOTA	1671	2.6	0.2	( 2.3, 3.0)	2435	3.3	0.2	( 3.0, 3.7)
TENNESSEE	1012	3.2	0.3	( 2.6, 3.7)	1921	3.5	0.2	( 3.1, 3.9)
TEXAS	1976	2.7	0.2	( 2.3, 3.1)	2933	3.9	0.2	( 3.5, 4.4)
UTAH	1362	2.7	0.2	( 2.2, 3.1)	1797	3.7	0.2	( 3.3, 4.2)
VERMONT	1255	2.6	0.2	( 2.1, 3.0)	1846	3.1	0.2	( 2.8, 3.5)
VIRGINIA	1377	2.3	0.2	( 1.9, 2.6)	2017	3.1	0.2	( 2.7, 3.5)
WASHINGTON	1578	3.1	0.2	( 2.6, 3.5)	1947	3.9	0.2	( 3.5, 4.3)
WEST VIRGINIA	1062	3.9	0.3	( 3.3, 4.5)	1510	4.2	0.3	( 3.7, 4.7)
WISCONSIN	917	3.1	0.3	( 2.6, 3.6)	1245	4.0	0.3	( 3.4, 4.6)
WYOMING	1061	2.5	0.2	( 2.1, 3.0)	1326	3.2	0.2	( 2.8, 3.6)
PUERTO RICO	1095	3.1	0.2	( 2.6, 3.5)	1859	4.5	0.2	( 4.1, 5.0)

SUMMARY STATISTICS:      NUMBER OF PARTICIPANTS = 52  
 MEDIAN                      = 2.7  
 RANGE                        = 1.0- 3.9

NUMBER OF PARTICIPANTS = 52  
 MEDIAN                      = 3.6  
 RANGE                        = 0.8- 4.6

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

June 20, 2000

TABLE 2.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 MEAN DIFFERENCES FROM 1998 TO 1999 PHYSICAL HEALTH NOT GOOD\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 MEAN	1999 MEAN	DIFFERENCE
ALABAMA	4.2	4.0	-0.2
ALASKA	2.8	2.8	-0.0
ARIZONA	1.1	0.9	-0.2
ARKANSAS	3.6	4.0	0.4
CALIFORNIA	3.1	3.3	0.2
COLORADO	2.9	3.3	0.4
CONNECTICUT	2.9	3.2	0.3
DELAWARE	3.2	3.3	0.1
DISTRICT OF CO	2.2	2.7	0.5
FLORIDA	3.4	3.4	0.0
GEORGIA	3.0	2.9	-0.1
HAWAII	2.2	2.4	0.2
IDAHO	3.1	3.2	0.2
ILLINOIS	2.8	3.2	0.4
INDIANA	3.3	3.1	-0.2
IOWA	2.9	3.4	0.4
KANSAS	2.8	2.6	-0.1
KENTUCKY	3.6	4.1	0.5
LOUISIANA	3.0	3.3	0.3
MAINE	3.3	3.5	0.2
MARYLAND	3.0	3.1	0.1
MASSACHUSETTS	2.8	3.2	0.4
MICHIGAN	3.7	3.4	-0.4
MINNESOTA	3.2	2.6	-0.6
MISSISSIPPI	3.7	3.5	-0.2
MISSOURI	3.4	3.5	0.1
MONTANA	3.0	3.1	0.2
NEBRASKA	3.1	3.1	0.0
NEVADA	3.2	3.5	0.4
NEW HAMPSHIRE	3.0	3.1	0.1
NEW JERSEY	2.8	3.2	0.4
NEW MEXICO	3.5	3.7	0.2
NEW YORK	3.2	3.3	0.1
NORTH CAROLINA	3.7	3.7	0.0
NORTH DAKOTA	2.7	2.9	0.2
OHIO	3.1	3.1	0.0
OKLAHOMA	2.6	3.0	0.3
OREGON	3.3	3.6	0.3
PENNSYLVANIA	3.3	3.2	-0.2
RHODE ISLAND	3.4	3.0	-0.4
SOUTH CAROLINA	3.1	3.2	0.1
SOUTH DAKOTA	2.6	3.0	0.4
TENNESSEE	3.3	3.4	0.1
TEXAS	3.3	3.3	0.1
UTAH	3.4	3.2	-0.2
VERMONT	3.1	2.9	-0.2
VIRGINIA	2.8	2.7	-0.1
WASHINGTON	3.1	3.5	0.4
WEST VIRGINIA	3.4	4.0	0.6
WISCONSIN	2.9	3.6	0.7
WYOMING	2.9	2.9	-0.0
PUERTO RICO	3.6	3.8	0.2

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS PHYSICAL HEALTH NOT GOOD

June 20, 2000

TABLE 3.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2046	3.6	0.2	( 3.2, 4.0)
ALASKA	2010	3.2	0.2	( 2.8, 3.7)
ARIZONA	1736	0.7	0.1	( 0.4, 0.9)
ARKANSAS	2950	3.2	0.2	( 2.9, 3.5)
CALIFORNIA	4278	3.3	0.1	( 3.0, 3.5)
COLORADO	3029	3.3	0.2	( 2.9, 3.7)
CONNECTICUT	3452	2.8	0.2	( 2.5, 3.1)
DELAWARE	2527	3.3	0.2	( 2.9, 3.8)
DISTRICT OF CO	1257	2.8	0.2	( 2.3, 3.2)
FLORIDA	5097	2.9	0.1	( 2.7, 3.2)
GEORGIA	2217	3.1	0.2	( 2.7, 3.5)
HAWAII	2135	2.2	0.2	( 1.9, 2.5)
IDAHO	4894	3.1	0.1	( 2.8, 3.3)
ILLINOIS	2881	2.8	0.1	( 2.5, 3.0)
INDIANA	2420	3.1	0.2	( 2.7, 3.6)
IOWA	3586	2.8	0.1	( 2.5, 3.0)
KANSAS	3796	2.4	0.1	( 2.1, 2.6)
KENTUCKY	7448	4.7	0.2	( 4.4, 5.0)
LOUISIANA	1619	3.2	0.2	( 2.7, 3.6)
MAINE	1647	2.7	0.2	( 2.3, 3.1)
MARYLAND	3892	3.7	0.2	( 3.4, 4.0)
MASSACHUSETTS	4918	2.8	0.1	( 2.5, 3.0)
MICHIGAN	2478	3.4	0.2	( 3.1, 3.7)
MINNESOTA	5283	2.8	0.1	( 2.6, 3.0)
MISSISSIPPI	2105	3.5	0.2	( 3.1, 3.9)
MISSOURI	4195	2.9	0.1	( 2.7, 3.2)
MONTANA	1744	2.7	0.2	( 2.3, 3.0)
NEBRASKA	2794	2.8	0.2	( 2.5, 3.1)
NEVADA	2160	4.1	0.3	( 3.6, 4.6)
NEW HAMPSHIRE	1222	3.3	0.3	( 2.8, 3.8)
NEW JERSEY	2782	2.9	0.2	( 2.6, 3.3)
NEW MEXICO	3432	3.8	0.2	( 3.5, 4.1)
NEW YORK	2602	2.9	0.2	( 2.6, 3.2)
NORTH CAROLINA	2414	3.0	0.2	( 2.6, 3.4)
NORTH DAKOTA	1971	2.5	0.2	( 2.2, 2.8)
OHIO	1624	2.3	0.2	( 2.0, 2.6)
OKLAHOMA	2941	2.4	0.1	( 2.2, 2.7)
OREGON	1786	3.9	0.2	( 3.5, 4.3)
PENNSYLVANIA	3505	2.7	0.1	( 2.5, 3.0)
RHODE ISLAND	3920	2.7	0.1	( 2.4, 2.9)
SOUTH CAROLINA	3413	2.7	0.1	( 2.4, 3.0)
SOUTH DAKOTA	4105	2.8	0.1	( 2.5, 3.0)
TENNESSEE	2934	3.0	0.1	( 2.7, 3.3)
TEXAS	4921	3.4	0.1	( 3.1, 3.7)
UTAH	3169	3.3	0.2	( 3.0, 3.6)
VERMONT	3091	2.9	0.1	( 2.6, 3.1)
VIRGINIA	3399	3.1	0.2	( 2.8, 3.4)
WASHINGTON	3515	3.1	0.2	( 2.8, 3.4)
WEST VIRGINIA	2569	3.0	0.2	( 2.6, 3.3)
WISCONSIN	2148	3.2	0.2	( 2.8, 3.5)
WYOMING	2388	2.8	0.2	( 2.5, 3.1)
PUERTO RICO	2948	2.4	0.1	( 2.2, 2.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 2.9 RANGE = 0.7- 4.7

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

June 20, 2000

TABLE 3.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR MENTAL HEALTH NOT GOOD\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	835	2.7	0.2	( 2.2, 3.1)	*	1211	4.4	0.3	( 3.9, 4.9)	*
ALASKA	986	2.5	0.3	( 1.9, 3.2)	*	1024	4.0	0.3	( 3.3, 4.6)	*
ARIZONA	711	0.7	0.2	( 0.3, 1.1)	*	1025	0.7	0.1	( 0.4, 0.9)	*
ARKANSAS	1152	2.5	0.2	( 2.1, 2.9)	*	1798	3.8	0.2	( 3.4, 4.2)	*
CALIFORNIA	1828	2.8	0.2	( 2.5, 3.2)	*	2450	3.7	0.2	( 3.3, 4.0)	*
COLORADO	1362	3.0	0.3	( 2.4, 3.5)	*	1667	3.6	0.2	( 3.1, 4.1)	*
CONNECTICUT	1349	2.9	0.3	( 2.3, 3.4)	*	2103	2.8	0.2	( 2.4, 3.1)	*
DELAWARE	1011	2.7	0.3	( 2.1, 3.2)	*	1516	4.0	0.4	( 3.2, 4.7)	*
DISTRICT OF CO	459	2.6	0.4	( 1.8, 3.3)	*	798	2.9	0.3	( 2.4, 3.4)	*
FLORIDA	2022	2.4	0.2	( 2.0, 2.8)	*	3075	3.4	0.2	( 3.1, 3.7)	*
GEORGIA	868	2.4	0.3	( 1.9, 2.9)	*	1349	3.8	0.3	( 3.3, 4.3)	*
HAWAII	988	1.7	0.2	( 1.3, 2.1)	*	1147	2.7	0.3	( 2.2, 3.2)	*
IDAHO	2074	2.3	0.2	( 2.0, 2.7)	*	2820	3.7	0.2	( 3.4, 4.1)	*
ILLINOIS	1099	2.3	0.2	( 1.9, 2.7)	*	1782	3.2	0.2	( 2.8, 3.5)	*
INDIANA	1013	2.6	0.3	( 2.0, 3.2)	*	1407	3.6	0.3	( 3.0, 4.2)	*
IOWA	1416	2.4	0.2	( 2.0, 2.7)	*	2170	3.2	0.2	( 2.8, 3.5)	*
KANSAS	1537	2.0	0.2	( 1.6, 2.4)	*	2259	2.7	0.2	( 2.4, 3.1)	*
KENTUCKY	2724	3.9	0.2	( 3.5, 4.3)	*	4724	5.4	0.2	( 5.0, 5.8)	*
LOUISIANA	642	2.3	0.3	( 1.8, 2.9)	*	977	3.9	0.3	( 3.3, 4.6)	*
MAINE	728	2.0	0.2	( 1.6, 2.5)	*	919	3.3	0.3	( 2.7, 3.9)	*
MARYLAND	1567	3.2	0.2	( 2.7, 3.7)	*	2325	4.2	0.2	( 3.8, 4.6)	*
MASSACHUSETTS	1975	2.5	0.2	( 2.1, 2.9)	*	2943	3.0	0.2	( 2.7, 3.3)	*
MICHIGAN	1023	2.6	0.2	( 2.1, 3.0)	*	1455	4.1	0.2	( 3.7, 4.6)	*
MINNESOTA	2416	2.5	0.2	( 2.2, 2.8)	*	2867	3.1	0.1	( 2.9, 3.4)	*
MISSISSIPPI	827	2.7	0.3	( 2.1, 3.2)	*	1278	4.2	0.3	( 3.6, 4.7)	*
MISSOURI	1692	2.2	0.2	( 1.8, 2.6)	*	2503	3.6	0.2	( 3.2, 4.0)	*
MONTANA	748	2.1	0.2	( 1.7, 2.6)	*	996	3.2	0.2	( 2.7, 3.7)	*
NEBRASKA	1110	2.6	0.2	( 2.1, 3.0)	*	1684	3.0	0.2	( 2.6, 3.3)	*
NEVADA	1089	3.2	0.3	( 2.7, 3.8)	*	1071	5.0	0.4	( 4.3, 5.8)	*
NEW HAMPSHIRE	497	2.9	0.4	( 2.1, 3.6)	*	725	3.7	0.3	( 3.1, 4.4)	*
NEW JERSEY	1114	3.1	0.3	( 2.6, 3.6)	*	1668	2.8	0.2	( 2.4, 3.2)	*
NEW MEXICO	1512	3.0	0.2	( 2.6, 3.4)	*	1920	4.5	0.2	( 4.1, 4.9)	*
NEW YORK	1043	2.8	0.3	( 2.3, 3.3)	*	1559	3.0	0.2	( 2.7, 3.4)	*
NORTH CAROLINA	951	2.5	0.3	( 1.9, 3.1)	*	1463	3.5	0.3	( 3.0, 4.0)	*
NORTH DAKOTA	856	2.1	0.2	( 1.7, 2.5)	*	1115	2.9	0.2	( 2.5, 3.4)	*
OHIO	632	2.1	0.3	( 1.6, 2.6)	*	992	2.5	0.2	( 2.1, 2.9)	*
OKLAHOMA	1145	1.9	0.2	( 1.5, 2.2)	*	1796	3.0	0.2	( 2.6, 3.3)	*
OREGON	762	3.3	0.3	( 2.8, 3.9)	*	1024	4.5	0.3	( 3.9, 5.1)	*
PENNSYLVANIA	1405	2.1	0.2	( 1.8, 2.5)	*	2100	3.2	0.2	( 2.9, 3.6)	*
RHODE ISLAND	1539	2.4	0.2	( 2.0, 2.8)	*	2381	2.9	0.2	( 2.6, 3.2)	*
SOUTH CAROLINA	1380	2.4	0.2	( 2.0, 2.8)	*	2033	3.0	0.2	( 2.7, 3.4)	*
SOUTH DAKOTA	1675	2.3	0.2	( 1.9, 2.7)	*	2430	3.2	0.2	( 2.9, 3.5)	*
TENNESSEE	1020	2.4	0.2	( 2.0, 2.8)	*	1914	3.5	0.2	( 3.1, 3.9)	*
TEXAS	1980	2.8	0.2	( 2.4, 3.2)	*	2941	4.0	0.2	( 3.6, 4.3)	*
UTAH	1361	2.5	0.2	( 2.1, 2.9)	*	1808	4.1	0.3	( 3.6, 4.6)	*
VERMONT	1253	2.4	0.2	( 2.0, 2.8)	*	1838	3.3	0.2	( 2.9, 3.7)	*
VIRGINIA	1372	2.6	0.2	( 2.2, 3.1)	*	2027	3.5	0.2	( 3.0, 3.9)	*
WASHINGTON	1562	2.7	0.3	( 2.2, 3.2)	*	1953	3.5	0.2	( 3.2, 3.9)	*
WEST VIRGINIA	1054	2.5	0.3	( 2.0, 3.0)	*	1515	3.3	0.2	( 2.8, 3.8)	*
WISCONSIN	918	2.7	0.2	( 2.2, 3.1)	*	1230	3.6	0.2	( 3.1, 4.1)	*
WYOMING	1058	2.4	0.2	( 2.0, 2.8)	*	1330	3.2	0.2	( 2.8, 3.6)	*
PUERTO RICO	1092	2.1	0.2	( 1.7, 2.5)	*	1856	2.7	0.2	( 2.4, 3.1)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 2.5	MEDIAN	= 3.4
	RANGE	= 0.7- 3.9	RANGE	= 0.7- 5.4

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

June 20, 2000

TABLE 3.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 MEAN DIFFERENCES FROM 1998 TO 1999 MENTAL HEALTH NOT GOOD\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 MEAN	1999 MEAN	DIFFERENCE
ALABAMA	3.7	3.6	-0.1
ALASKA	2.3	3.2	0.9
ARIZONA	0.8	0.7	-0.1
ARKANSAS	3.6	3.2	-0.4
CALIFORNIA	3.0	3.3	0.2
COLORADO	3.0	3.3	0.3
CONNECTICUT	2.8	2.8	0.0
DELAWARE	3.1	3.3	0.2
DISTRICT OF CO	2.2	2.8	0.6
FLORIDA	3.0	2.9	-0.0
GEORGIA	3.4	3.1	-0.3
HAWAII	2.2	2.2	-0.0
IDAHO	2.8	3.1	0.2
ILLINOIS	2.8	2.8	-0.0
INDIANA	3.2	3.1	-0.1
IOWA	3.0	2.8	-0.2
KANSAS	2.7	2.4	-0.3
KENTUCKY	4.5	4.7	0.2
LOUISIANA	2.8	3.2	0.4
MAINE	2.7	2.7	0.0
MARYLAND	3.4	3.7	0.4
MASSACHUSETTS	2.8	2.8	0.0
MICHIGAN	3.6	3.4	-0.3
MINNESOTA	3.2	2.8	-0.4
MISSISSIPPI	3.3	3.5	0.2
MISSOURI	3.1	2.9	-0.2
MONTANA	2.6	2.7	0.1
NEBRASKA	3.0	2.8	-0.2
NEVADA	3.5	4.1	0.6
NEW HAMPSHIRE	3.0	3.3	0.3
NEW JERSEY	2.7	2.9	0.2
NEW MEXICO	3.3	3.8	0.4
NEW YORK	3.0	2.9	-0.0
NORTH CAROLINA	2.7	3.0	0.3
NORTH DAKOTA	2.5	2.5	0.0
OHIO	2.9	2.3	-0.6
OKLAHOMA	1.9	2.4	0.5
OREGON	3.3	3.9	0.6
PENNSYLVANIA	2.9	2.7	-0.2
RHODE ISLAND	3.0	2.7	-0.4
SOUTH CAROLINA	3.0	2.7	-0.3
SOUTH DAKOTA	2.5	2.8	0.3
TENNESSEE	3.5	3.0	-0.5
TEXAS	3.5	3.4	-0.1
UTAH	3.7	3.3	-0.4
VERMONT	2.8	2.9	0.1
VIRGINIA	3.0	3.1	0.1
WASHINGTON	3.1	3.1	-0.0
WEST VIRGINIA	2.3	3.0	0.7
WISCONSIN	2.9	3.2	0.2
WYOMING	3.1	2.8	-0.3
PUERTO RICO	2.6	2.4	-0.2

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS MENTAL HEALTH NOT GOOD

June 20, 2000

TABLE 4.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR ACTIVITIES LIMITED \*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1153	4.3	0.3	( 3.8, 4.8)
ALASKA	1039	3.8	0.4	( 3.0, 4.6)
ARIZONA	295	2.9	0.3	( 2.2, 3.6)
ARKANSAS	1558	4.2	0.2	( 3.7, 4.7)
CALIFORNIA	2341	4.0	0.2	( 3.6, 4.5)
COLORADO	1654	3.1	0.3	( 2.6, 3.6)
CONNECTICUT	1668	3.5	0.3	( 2.9, 4.0)
DELAWARE	1339	3.3	0.3	( 2.8, 3.8)
DISTRICT OF CO	598	2.7	0.3	( 2.2, 3.2)
FLORIDA	2370	4.6	0.2	( 4.2, 5.1)
GEORGIA	1106	3.4	0.3	( 2.9, 4.0)
HAWAII	929	2.8	0.3	( 2.3, 3.3)
IDAHO	2743	3.2	0.2	( 2.8, 3.6)
ILLINOIS	1510	3.5	0.3	( 3.0, 4.0)
INDIANA	1382	3.5	0.4	( 2.8, 4.3)
IOWA	1890	3.1	0.2	( 2.8, 3.5)
KANSAS	1726	2.8	0.2	( 2.5, 3.2)
KENTUCKY	3477	6.2	0.2	( 5.8, 6.7)
LOUISIANA	739	4.5	0.4	( 3.7, 5.3)
MAINE	767	4.3	0.3	( 3.7, 5.0)
MARYLAND	2134	3.2	0.2	( 2.8, 3.6)
MASSACHUSETTS	2563	3.8	0.2	( 3.4, 4.1)
MICHIGAN	1420	3.1	0.2	( 2.7, 3.5)
MINNESOTA	2470	4.3	0.2	( 4.0, 4.7)
MISSISSIPPI	1097	4.2	0.3	( 3.6, 4.7)
MISSOURI	2199	3.7	0.2	( 3.2, 4.1)
MONTANA	880	3.4	0.3	( 2.9, 3.9)
NEBRASKA	1316	3.8	0.2	( 3.3, 4.3)
NEVADA	1260	3.2	0.3	( 2.7, 3.7)
NEW HAMPSHIRE	619	2.8	0.3	( 2.3, 3.3)
NEW JERSEY	1385	3.6	0.3	( 3.0, 4.1)
NEW MEXICO	1935	3.8	0.2	( 3.4, 4.3)
NEW YORK	1355	3.3	0.2	( 2.8, 3.7)
NORTH CAROLINA	1126	4.6	0.3	( 4.0, 5.2)
NORTH DAKOTA	940	2.9	0.2	( 2.4, 3.3)
OHIO	738	3.5	0.3	( 2.8, 4.1)
OKLAHOMA	1183	4.1	0.3	( 3.6, 4.6)
OREGON	1061	3.6	0.3	( 3.0, 4.1)
PENNSYLVANIA	1669	3.9	0.2	( 3.4, 4.3)
RHODE ISLAND	1913	3.7	0.2	( 3.2, 4.1)
SOUTH CAROLINA	1499	4.4	0.3	( 3.9, 4.9)
SOUTH DAKOTA	2033	3.0	0.2	( 2.6, 3.3)
TENNESSEE	1525	3.9	0.3	( 3.4, 4.4)
TEXAS	2657	3.7	0.2	( 3.3, 4.2)
UTAH	1916	2.7	0.2	( 2.3, 3.0)
VERMONT	1571	3.0	0.2	( 2.6, 3.3)
VIRGINIA	1839	3.0	0.2	( 2.7, 3.4)
WASHINGTON	2066	3.4	0.2	( 2.9, 3.8)
WEST VIRGINIA	1053	7.2	0.4	( 6.5, 7.9)
WISCONSIN	1279	2.9	0.3	( 2.5, 3.4)
WYOMING	1195	3.1	0.2	( 2.7, 3.6)
PUERTO RICO	1185	6.4	0.3	( 5.8, 7.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 3.5 RANGE = 2.7- 7.2

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

June 20, 2000



TABLE 4.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 MEAN DIFFERENCES FROM 1998 TO 1999 ACTIVITIES LIMITED \*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 MEAN	1999 MEAN	DIFFERENCE
ALABAMA	4.9	4.3	-0.6
ALASKA	2.8	3.8	1.0
ARIZONA	4.2	2.9	-1.3
ARKANSAS	6.1	4.2	-1.9
CALIFORNIA	3.7	4.0	0.4
COLORADO	2.9	3.1	0.2
CONNECTICUT	3.8	3.5	-0.3
DELAWARE	6.3	3.3	-3.0
DISTRICT OF CO	5.5	2.7	-2.8
FLORIDA	4.8	4.6	-0.2
GEORGIA	3.3	3.4	0.1
HAWAII	3.2	2.8	-0.4
IDAHO	2.9	3.2	0.4
ILLINOIS	2.8	3.5	0.7
INDIANA	3.1	3.5	0.4
IOWA	2.9	3.1	0.2
KANSAS	3.3	2.8	-0.5
KENTUCKY	6.6	6.2	-0.4
LOUISIANA	4.1	4.5	0.4
MAINE	4.1	4.3	0.2
MARYLAND	3.6	3.2	-0.4
MASSACHUSETTS	3.5	3.8	0.3
MICHIGAN	3.6	3.1	-0.5
MINNESOTA	4.2	4.3	0.2
MISSISSIPPI	4.1	4.2	0.0
MISSOURI	3.4	3.7	0.3
MONTANA	3.3	3.4	0.1
NEBRASKA	3.6	3.8	0.3
NEVADA	3.0	3.2	0.2
NEW HAMPSHIRE	3.2	2.8	-0.4
NEW JERSEY	4.0	3.6	-0.4
NEW MEXICO	3.8	3.8	0.0
NEW YORK	4.0	3.3	-0.7
NORTH CAROLINA	4.7	4.6	-0.0
NORTH DAKOTA	3.0	2.9	-0.1
OHIO	3.8	3.5	-0.3
OKLAHOMA	4.9	4.1	-0.9
OREGON	3.4	3.6	0.1
PENNSYLVANIA	4.0	3.9	-0.1
RHODE ISLAND	3.7	3.7	-0.1
SOUTH CAROLINA	4.1	4.4	0.3
SOUTH DAKOTA	3.1	3.0	-0.1
TENNESSEE	4.1	3.9	-0.2
TEXAS	4.0	3.7	-0.3
UTAH	3.1	2.7	-0.4
VERMONT	2.9	3.0	0.1
VIRGINIA	3.1	3.0	-0.0
WASHINGTON	3.1	3.4	0.3
WEST VIRGINIA	6.8	7.2	0.4
WISCONSIN	2.9	2.9	0.0
WYOMING	2.9	3.1	0.2
PUERTO RICO	5.4	6.4	1.1

\* MEAN NUMBER OF DAYS IN PAST 30 DAYS ACTIVITIES LIMITED

June 20, 2000

TABLE 5.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2091	15.1	0.9	( 13.2, 16.9)
ALASKA	2043	23.3	1.4	( 20.5, 26.1)
ARIZONA	1736	14.2	1.3	( 11.7, 16.7)
ARKANSAS	2995	16.1	0.8	( 14.5, 17.6)
CALIFORNIA	4299	19.3	0.7	( 17.9, 20.8)
COLORADO	3049	13.9	0.9	( 12.2, 15.7)
CONNECTICUT	3508	10.1	0.8	( 8.7, 11.6)
DELAWARE	2555	9.6	0.8	( 8.0, 11.2)
DISTRICT OF CO	1266	13.2	1.1	( 11.0, 15.3)
FLORIDA	5167	16.2	0.6	( 14.9, 17.4)
GEORGIA	2269	13.6	0.9	( 11.8, 15.3)
HAWAII	2153	9.1	0.8	( 7.5, 10.7)
IDAHO	4948	16.7	0.6	( 15.4, 18.0)
ILLINOIS	2900	11.8	0.8	( 10.3, 13.2)
INDIANA	2434	12.4	1.1	( 10.3, 14.5)
IOWA	3611	8.8	0.6	( 7.6, 9.9)
KANSAS	3867	10.4	0.6	( 9.2, 11.5)
KENTUCKY	7524	14.5	0.6	( 13.4, 15.6)
LOUISIANA	1658	22.4	1.2	( 20.0, 24.8)
MAINE	1672	13.2	1.0	( 11.3, 15.1)
MARYLAND	3948	9.9	0.6	( 8.7, 11.0)
MASSACHUSETTS	5007	7.2	0.5	( 6.2, 8.1)
MICHIGAN	2513	9.8	0.7	( 8.5, 11.1)
MINNESOTA	5334	5.8	0.4	( 5.1, 6.5)
MISSISSIPPI	2171	17.1	0.9	( 15.3, 18.8)
MISSOURI	4266	10.9	0.6	( 9.7, 12.1)
MONTANA	1796	17.0	1.1	( 14.9, 19.1)
NEBRASKA	2815	8.1	0.6	( 6.9, 9.2)
NEVADA	2184	18.1	1.4	( 15.4, 20.8)
NEW HAMPSHIRE	1248	11.1	1.2	( 8.8, 13.3)
NEW JERSEY	2813	11.9	0.8	( 10.4, 13.5)
NEW MEXICO	3482	22.0	0.8	( 20.4, 23.6)
NEW YORK	2643	14.6	0.8	( 12.9, 16.2)
NORTH CAROLINA	2442	11.5	0.8	( 9.9, 13.0)
NORTH DAKOTA	1978	11.1	0.8	( 9.6, 12.7)
OHIO	1629	10.0	0.9	( 8.2, 11.9)
OKLAHOMA	2960	17.3	0.9	( 15.6, 19.0)
OREGON	1806	15.4	1.0	( 13.4, 17.3)
PENNSYLVANIA	3550	10.2	0.6	( 9.0, 11.3)
RHODE ISLAND	3992	10.6	0.6	( 9.3, 11.8)
SOUTH CAROLINA	3451	15.6	0.8	( 14.1, 17.1)
SOUTH DAKOTA	4140	10.5	0.6	( 9.3, 11.6)
TENNESSEE	3011	12.4	0.7	( 10.9, 13.8)
TEXAS	4985	22.6	0.8	( 21.0, 24.2)
UTAH	3186	12.4	0.8	( 10.9, 13.9)
VERMONT	3154	12.8	0.7	( 11.4, 14.2)
VIRGINIA	3494	10.0	0.7	( 8.6, 11.3)
WASHINGTON	3599	11.9	0.7	( 10.5, 13.4)
WEST VIRGINIA	2584	18.1	0.9	( 16.3, 19.8)
WISCONSIN	2172	10.2	0.9	( 8.5, 11.9)
WYOMING	2399	16.9	0.9	( 15.1, 18.6)
PUERTO RICO	3048	11.7	0.8	( 10.2, 13.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 12.4 RANGE = 5.8-23.3

\* RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"



TABLE 5.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO HEALTH INSURANCE, AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	16.4	15.1	-1.3
ALASKA	20.8	23.3	2.5
ARIZONA	13.2	14.2	1.1
ARKANSAS	15.5	16.1	0.6
CALIFORNIA	18.1	19.3	1.2
COLORADO	14.9	13.9	-1.0
CONNECTICUT	8.8	10.1	1.3
DELAWARE	8.0	9.6	1.6
DISTRICT OF CO	11.6	13.2	1.6
FLORIDA	17.6	16.2	-1.4
GEORGIA	14.7	13.6	-1.2
HAWAII	5.9	9.1	3.2
IDAHO	15.2	16.7	1.5
ILLINOIS	10.9	11.8	0.8
INDIANA	13.3	12.4	-1.0
IOWA	8.7	8.8	0.1
KANSAS	10.8	10.4	-0.4
KENTUCKY	14.4	14.5	0.2
LOUISIANA	22.2	22.4	0.2
MAINE	13.2	13.2	-0.0
MARYLAND	13.6	9.9	-3.8
MASSACHUSETTS	9.0	7.2	-1.8
MICHIGAN	10.1	9.8	-0.3
MINNESOTA	8.6	5.8	-2.7
MISSISSIPPI	18.7	17.1	-1.7
MISSOURI	13.0	10.9	-2.1
MONTANA	17.2	17.0	-0.2
NEBRASKA	8.0	8.1	0.1
NEVADA	16.2	18.1	1.9
NEW HAMPSHIRE	11.7	11.1	-0.6
NEW JERSEY	10.2	11.9	1.7
NEW MEXICO	21.8	22.0	0.3
NEW YORK	13.7	14.6	0.8
NORTH CAROLINA	12.7	11.5	-1.3
NORTH DAKOTA	10.3	11.1	0.9
OHIO	8.7	10.0	1.4
OKLAHOMA	18.4	17.3	-1.1
OREGON	13.2	15.4	2.2
PENNSYLVANIA	10.9	10.2	-0.7
RHODE ISLAND	9.7	10.6	0.9
SOUTH CAROLINA	16.3	15.6	-0.6
SOUTH DAKOTA	12.3	10.5	-1.9
TENNESSEE	13.0	12.4	-0.7
TEXAS	23.6	22.6	-1.1
UTAH	11.9	12.4	0.5
VERMONT	10.7	12.8	2.1
VIRGINIA	12.2	10.0	-2.2
WASHINGTON	10.9	11.9	1.0
WEST VIRGINIA	17.0	18.1	1.1
WISCONSIN	8.3	10.2	1.9
WYOMING	17.1	16.9	-0.3
PUERTO RICO	11.0	11.7	0.7

\* RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

June 20, 2000

TABLE 6.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HEALTH INSURANCE, AGE 18-64 YEARS\*  
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1650	18.0	1.1	( 15.8, 20.2)
ALASKA	1872	24.8	1.5	( 21.8, 27.7)
ARIZONA	1262	16.7	1.5	( 13.7, 19.7)
ARKANSAS	2409	19.8	1.0	( 17.9, 21.8)
CALIFORNIA	3657	22.2	0.8	( 20.6, 23.9)
COLORADO	2551	16.3	1.1	( 14.2, 18.4)
CONNECTICUT	2849	12.3	0.9	( 10.5, 14.1)
DELAWARE	2095	11.3	1.0	( 9.4, 13.3)
DISTRICT OF CO	1011	15.2	1.3	( 12.7, 17.8)
FLORIDA	3905	20.5	0.8	( 18.9, 22.0)
GEORGIA	1936	15.6	1.0	( 13.6, 17.6)
HAWAII	1741	10.3	1.0	( 8.4, 12.1)
IDAHO	4046	20.1	0.8	( 18.6, 21.5)
ILLINOIS	2419	13.6	0.9	( 11.8, 15.3)
INDIANA	2000	15.1	1.3	( 12.5, 17.7)
IOWA	2693	10.9	0.7	( 9.5, 12.4)
KANSAS	2967	12.5	0.7	( 11.1, 13.9)
KENTUCKY	5767	17.3	0.7	( 15.9, 18.6)
LOUISIANA	1333	25.8	1.4	( 23.0, 28.6)
MAINE	1312	16.1	1.2	( 13.8, 18.5)
MARYLAND	3313	11.1	0.7	( 9.7, 12.5)
MASSACHUSETTS	4048	8.3	0.6	( 7.2, 9.5)
MICHIGAN	2043	11.4	0.8	( 9.8, 13.0)
MINNESOTA	4433	6.8	0.4	( 6.0, 7.6)
MISSISSIPPI	1779	20.3	1.1	( 18.2, 22.4)
MISSOURI	3296	13.1	0.8	( 11.6, 14.6)
MONTANA	1408	20.9	1.3	( 18.4, 23.4)
NEBRASKA	2061	9.8	0.7	( 8.3, 11.2)
NEVADA	1841	21.3	1.6	( 18.1, 24.4)
NEW HAMPSHIRE	1016	13.2	1.4	( 10.5, 15.8)
NEW JERSEY	2233	14.1	0.9	( 12.3, 16.0)
NEW MEXICO	2838	25.8	1.0	( 24.0, 27.7)
NEW YORK	2168	17.4	1.0	( 15.4, 19.3)
NORTH CAROLINA	1893	13.6	1.0	( 11.7, 15.5)
NORTH DAKOTA	1492	13.8	1.0	( 11.9, 15.8)
OHIO	1224	12.2	1.2	( 9.9, 14.4)
OKLAHOMA	2285	20.5	1.0	( 18.4, 22.5)
OREGON	1446	18.6	1.2	( 16.2, 21.0)
PENNSYLVANIA	2808	12.5	0.8	( 11.1, 14.0)
RHODE ISLAND	3173	12.6	0.8	( 11.1, 14.1)
SOUTH CAROLINA	2897	18.2	0.9	( 16.4, 19.9)
SOUTH DAKOTA	3171	13.0	0.7	( 11.6, 14.4)
TENNESSEE	2409	14.2	0.9	( 12.5, 15.9)
TEXAS	4260	26.3	0.9	( 24.5, 28.1)
UTAH	2628	14.2	0.9	( 12.4, 16.0)
VERMONT	2517	14.8	0.9	( 13.1, 16.5)
VIRGINIA	2900	11.2	0.8	( 9.6, 12.7)
WASHINGTON	2983	14.1	0.9	( 12.4, 15.8)
WEST VIRGINIA	1963	22.3	1.1	( 20.2, 24.4)
WISCONSIN	1763	12.1	1.0	( 10.1, 14.1)
WYOMING	1990	19.9	1.0	( 17.9, 21.9)
PUERTO RICO	2358	13.2	0.9	( 11.4, 14.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 14.5 RANGE = 6.8-26.3 YEAR 2000 OBJECTIVE 21.4

\* RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

June 20, 2000



TABLE 6.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO HEALTH INSURANCE, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	19.5	18.0	-1.5
ALASKA	21.8	24.8	3.0
ARIZONA	15.8	16.7	0.9
ARKANSAS	19.1	19.8	0.7
CALIFORNIA	21.2	22.2	1.0
COLORADO	17.3	16.3	-1.0
CONNECTICUT	10.6	12.3	1.7
DELAWARE	9.7	11.3	1.7
DISTRICT OF CO	13.0	15.2	2.2
FLORIDA	22.4	20.5	-2.0
GEORGIA	16.9	15.6	-1.3
HAWAII	7.0	10.3	3.3
IDAHO	18.3	20.1	1.8
ILLINOIS	13.0	13.6	0.6
INDIANA	15.9	15.1	-0.8
IOWA	10.8	10.9	0.2
KANSAS	13.0	12.5	-0.4
KENTUCKY	17.3	17.3	-0.0
LOUISIANA	25.9	25.8	-0.1
MAINE	15.7	16.1	0.4
MARYLAND	15.7	11.1	-4.6
MASSACHUSETTS	10.5	8.3	-2.2
MICHIGAN	11.9	11.4	-0.5
MINNESOTA	9.9	6.8	-3.1
MISSISSIPPI	22.5	20.3	-2.1
MISSOURI	15.9	13.1	-2.7
MONTANA	21.3	20.9	-0.4
NEBRASKA	9.8	9.8	-0.1
NEVADA	19.2	21.3	2.1
NEW HAMPSHIRE	13.7	13.2	-0.5
NEW JERSEY	11.8	14.1	2.3
NEW MEXICO	25.7	25.8	0.2
NEW YORK	16.6	17.4	0.8
NORTH CAROLINA	15.2	13.6	-1.5
NORTH DAKOTA	12.7	13.8	1.2
OHIO	10.4	12.2	1.8
OKLAHOMA	22.4	20.5	-1.9
OREGON	15.8	18.6	2.8
PENNSYLVANIA	12.9	12.5	-0.3
RHODE ISLAND	11.7	12.6	0.9
SOUTH CAROLINA	18.7	18.2	-0.6
SOUTH DAKOTA	15.2	13.0	-2.1
TENNESSEE	14.8	14.2	-0.6
TEXAS	27.5	26.3	-1.3
UTAH	13.8	14.2	0.4
VERMONT	12.4	14.8	2.4
VIRGINIA	13.8	11.2	-2.7
WASHINGTON	12.8	14.1	1.3
WEST VIRGINIA	20.6	22.3	1.7
WISCONSIN	10.0	12.1	2.1
WYOMING	20.2	19.9	-0.3
PUERTO RICO	12.6	13.2	0.6

\* RESPONDENTS ANSWERING "NO" TO "DO YOU HAVE ANY KIND OF HEALTH CARE COVERAGE, INCLUDING HEALTH INSURANCE, PREPAID PLANS SUCH AS HMOs OR GOVERNMENT PLANS SUCH AS MEDICARE"

TABLE 7.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS\*  
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2093	7.4	0.6	( 6.2, 8.6)
ALASKA	2048	3.5	0.6	( 2.4, 4.6)
ARIZONA	1739	4.3	0.6	( 3.1, 5.5)
ARKANSAS	2995	6.6	0.5	( 5.6, 7.6)
CALIFORNIA	4300	6.1	0.5	( 5.2, 7.0)
COLORADO	3052	3.8	0.5	( 2.9, 4.7)
CONNECTICUT	3513	4.3	0.4	( 3.5, 5.2)
DELAWARE	2559	6.0	0.6	( 4.9, 7.2)
DISTRICT OF CO	1267	6.5	0.7	( 5.1, 7.9)
FLORIDA	5174	6.8	0.4	( 6.0, 7.7)
GEORGIA	2273	5.6	0.5	( 4.5, 6.6)
HAWAII	2150	5.2	0.6	( 3.9, 6.4)
IDAHO	4953	4.8	0.4	( 4.1, 5.5)
ILLINOIS	2910	6.4	0.6	( 5.2, 7.6)
INDIANA	2440	6.6	0.9	( 4.9, 8.4)
IOWA	3613	5.2	0.4	( 4.5, 6.0)
KANSAS	3872	5.4	0.4	( 4.7, 6.1)
KENTUCKY	7535	6.4	0.4	( 5.7, 7.1)
LOUISIANA	1668	6.1	0.6	( 4.9, 7.3)
MAINE	1671	5.4	0.6	( 4.3, 6.5)
MARYLAND	3955	6.8	0.5	( 5.8, 7.8)
MASSACHUSETTS	5019	5.0	0.4	( 4.2, 5.8)
MICHIGAN	2519	5.5	0.5	( 4.5, 6.4)
MINNESOTA	5336	4.8	0.3	( 4.1, 5.4)
MISSISSIPPI	2179	7.9	0.6	( 6.6, 9.1)
MISSOURI	4274	6.1	0.5	( 5.2, 7.0)
MONTANA	1795	5.9	0.6	( 4.7, 7.1)
NEBRASKA	2818	4.3	0.4	( 3.5, 5.1)
NEVADA	2186	5.8	0.7	( 4.5, 7.1)
NEW HAMPSHIRE	1245	4.3	0.7	( 3.0, 5.6)
NEW JERSEY	2819	5.4	0.5	( 4.4, 6.4)
NEW MEXICO	3488	5.5	0.5	( 4.6, 6.4)
NEW YORK	2647	5.7	0.5	( 4.7, 6.7)
NORTH CAROLINA	2442	6.2	0.5	( 5.2, 7.1)
NORTH DAKOTA	1979	5.0	0.5	( 4.0, 6.0)
OHIO	1638	6.1	0.6	( 4.8, 7.3)
OKLAHOMA	2963	5.8	0.5	( 4.8, 6.7)
OREGON	1808	4.6	0.6	( 3.5, 5.7)
PENNSYLVANIA	3554	6.4	0.5	( 5.4, 7.4)
RHODE ISLAND	3997	5.3	0.4	( 4.4, 6.1)
SOUTH CAROLINA	3462	6.4	0.5	( 5.5, 7.4)
SOUTH DAKOTA	4145	4.9	0.4	( 4.2, 5.7)
TENNESSEE	3012	6.0	0.5	( 5.0, 7.0)
TEXAS	4989	6.2	0.5	( 5.3, 7.1)
UTAH	3194	4.2	0.5	( 3.3, 5.1)
VERMONT	3159	4.3	0.4	( 3.5, 5.0)
VIRGINIA	3496	6.1	0.6	( 4.9, 7.3)
WASHINGTON	3606	5.2	0.4	( 4.3, 6.0)
WEST VIRGINIA	2588	7.3	0.5	( 6.3, 8.3)
WISCONSIN	2175	5.3	0.5	( 4.3, 6.4)
WYOMING	2402	4.6	0.5	( 3.7, 5.5)
PUERTO RICO	3045	9.6	0.6	( 8.5, 10.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 5.6 RANGE = 3.5- 9.6

\* EVER TOLD YOU HAVE DIABETES

June 20, 2000

TABLE 7.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETES AWARENESS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	853	7.5	0.9	( 5.6,	9.3)	1240	7.3	0.8	( 5.8,	8.8)
ALASKA	1000	3.9	0.8	( 2.3,	5.5)	1048	3.1	0.8	( 1.7,	4.6)
ARIZONA	710	5.1	1.1	( 3.0,	7.2)	1029	3.5	0.6	( 2.3,	4.7)
ARKANSAS	1166	6.1	0.8	( 4.6,	7.6)	1829	7.1	0.7	( 5.7,	8.4)
CALIFORNIA	1835	6.0	0.7	( 4.7,	7.3)	2465	6.2	0.6	( 5.0,	7.4)
COLORADO	1371	3.4	0.7	( 2.1,	4.7)	1681	4.2	0.6	( 3.0,	5.5)
CONNECTICUT	1372	4.5	0.7	( 3.1,	5.8)	2141	4.2	0.5	( 3.2,	5.2)
DELAWARE	1027	7.0	1.0	( 5.1,	8.9)	1532	5.2	0.7	( 3.7,	6.6)
DISTRICT OF CO	459	6.3	1.1	( 4.1,	8.4)	808	6.8	0.9	( 5.0,	8.6)
FLORIDA	2054	6.9	0.7	( 5.6,	8.2)	3120	6.8	0.5	( 5.8,	7.8)
GEORGIA	886	5.1	0.8	( 3.6,	6.7)	1387	5.9	0.7	( 4.5,	7.4)
HAWAII	991	6.4	1.0	( 4.3,	8.4)	1159	3.9	0.7	( 2.6,	5.3)
IDAHO	2096	4.8	0.5	( 3.8,	5.9)	2857	4.8	0.5	( 3.8,	5.7)
ILLINOIS	1111	6.9	1.0	( 4.9,	9.0)	1799	5.9	0.7	( 4.6,	7.2)
INDIANA	1019	6.1	1.3	( 3.5,	8.7)	1421	7.1	1.2	( 4.8,	9.4)
IOWA	1423	5.8	0.6	( 4.6,	7.0)	2190	4.7	0.5	( 3.7,	5.7)
KANSAS	1562	5.3	0.6	( 4.1,	6.4)	2310	5.5	0.5	( 4.6,	6.5)
KENTUCKY	2749	6.1	0.6	( 5.0,	7.3)	4786	6.6	0.5	( 5.7,	7.5)
LOUISIANA	662	6.0	1.0	( 4.1,	7.8)	1006	6.2	0.8	( 4.6,	7.9)
MAINE	737	4.7	0.8	( 3.1,	6.2)	934	6.1	0.8	( 4.4,	7.7)
MARYLAND	1587	6.0	0.7	( 4.6,	7.4)	2368	7.5	0.7	( 6.0,	8.9)
MASSACHUSETTS	2006	6.5	0.7	( 5.1,	7.9)	3013	3.7	0.4	( 2.9,	4.6)
MICHIGAN	1040	6.3	0.8	( 4.7,	7.8)	1479	4.7	0.6	( 3.6,	5.9)
MINNESOTA	2436	5.3	0.5	( 4.3,	6.3)	2900	4.3	0.4	( 3.5,	5.1)
MISSISSIPPI	852	7.6	1.0	( 5.7,	9.5)	1327	8.2	0.8	( 6.6,	9.8)
MISSOURI	1721	6.2	0.8	( 4.7,	7.7)	2553	6.0	0.6	( 4.9,	7.2)
MONTANA	767	6.5	1.0	( 4.6,	8.4)	1028	5.4	0.8	( 3.9,	6.8)
NEBRASKA	1117	4.5	0.6	( 3.3,	5.8)	1701	4.1	0.5	( 3.1,	5.1)
NEVADA	1100	6.7	1.0	( 4.6,	8.7)	1086	4.9	0.9	( 3.2,	6.6)
NEW HAMPSHIRE	501	4.5	1.0	( 2.4,	6.5)	744	4.2	0.8	( 2.7,	5.7)
NEW JERSEY	1123	4.9	0.8	( 3.4,	6.4)	1696	5.9	0.7	( 4.5,	7.2)
NEW MEXICO	1532	4.6	0.6	( 3.5,	5.7)	1956	6.2	0.7	( 4.8,	7.6)
NEW YORK	1067	5.7	0.8	( 4.2,	7.3)	1580	5.7	0.7	( 4.3,	7.0)
NORTH CAROLINA	964	5.6	0.7	( 4.2,	7.0)	1478	6.7	0.7	( 5.3,	8.0)
NORTH DAKOTA	860	4.4	0.7	( 3.0,	5.8)	1119	5.6	0.7	( 4.2,	6.9)
OHIO	636	5.0	0.9	( 3.2,	6.7)	1002	7.1	0.9	( 5.3,	8.8)
OKLAHOMA	1152	6.4	0.8	( 5.0,	7.9)	1811	5.1	0.5	( 4.1,	6.2)
OREGON	769	4.1	0.8	( 2.6,	5.6)	1039	5.1	0.8	( 3.5,	6.7)
PENNSYLVANIA	1416	6.7	0.7	( 5.2,	8.1)	2138	6.1	0.7	( 4.9,	7.4)
RHODE ISLAND	1558	5.4	0.7	( 4.2,	6.7)	2439	5.1	0.6	( 4.0,	6.2)
SOUTH CAROLINA	1394	6.6	0.7	( 5.2,	8.1)	2068	6.2	0.6	( 5.0,	7.4)
SOUTH DAKOTA	1683	4.7	0.5	( 3.7,	5.8)	2462	5.1	0.5	( 4.2,	6.1)
TENNESSEE	1035	5.9	0.8	( 4.2,	7.5)	1977	6.2	0.6	( 5.0,	7.4)
TEXAS	2004	6.3	0.6	( 5.0,	7.5)	2985	6.2	0.7	( 4.9,	7.5)
UTAH	1371	4.2	0.7	( 2.8,	5.5)	1823	4.2	0.6	( 3.0,	5.4)
VERMONT	1272	4.5	0.6	( 3.3,	5.7)	1887	4.1	0.5	( 3.2,	5.0)
VIRGINIA	1405	6.9	1.1	( 4.7,	9.1)	2091	5.4	0.6	( 4.2,	6.5)
WASHINGTON	1601	5.2	0.7	( 3.8,	6.5)	2005	5.2	0.6	( 4.1,	6.3)
WEST VIRGINIA	1065	6.9	0.8	( 5.3,	8.4)	1523	7.7	0.7	( 6.3,	9.1)
WISCONSIN	919	6.0	0.9	( 4.3,	7.6)	1256	4.7	0.7	( 3.4,	6.0)
WYOMING	1063	4.1	0.6	( 2.9,	5.3)	1339	5.0	0.7	( 3.7,	6.3)
PUERTO RICO	1119	8.2	0.8	( 6.5,	9.8)	1926	10.9	0.8	( 9.4,	12.4)

SUMMARY STATISTICS:      NUMBER OF PARTICIPANTS = 52      NUMBER OF PARTICIPANTS = 52  
 MEDIAN                      = 5.9      MEDIAN                      = 5.5  
 RANGE                        = 3.4- 8.2      RANGE                        = 3.1-10.9

\* EVER TOLD YOU HAVE DIABETES

June 20, 2000

TABLE 7.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 DIABETES AWARENESS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	7.0	7.4	0.4
ALASKA	3.0	3.5	0.6
ARIZONA	2.8	4.3	1.4
ARKANSAS	6.7	6.6	-0.1
CALIFORNIA	5.5	6.1	0.6
COLORADO	4.6	3.8	-0.8
CONNECTICUT	4.5	4.3	-0.2
DELAWARE	4.4	6.0	1.7
DISTRICT OF CO	7.1	6.5	-0.5
FLORIDA	6.4	6.8	0.5
GEORGIA	5.9	5.6	-0.4
HAWAII	5.6	5.2	-0.4
IDAHO	4.3	4.8	0.5
ILLINOIS	6.2	6.4	0.2
INDIANA	6.0	6.6	0.6
IOWA	5.2	5.2	0.0
KANSAS	4.1	5.4	1.4
KENTUCKY	5.6	6.4	0.8
LOUISIANA	6.4	6.1	-0.3
MAINE	3.6	5.4	1.8
MARYLAND	5.4	6.8	1.4
MASSACHUSETTS	3.9	5.0	1.2
MICHIGAN	7.0	5.5	-1.6
MINNESOTA	4.8	4.8	0.0
MISSISSIPPI	7.6	7.9	0.3
MISSOURI	5.8	6.1	0.3
MONTANA	3.6	5.9	2.4
NEBRASKA	5.2	4.3	-0.8
NEVADA	4.4	5.8	1.4
NEW HAMPSHIRE	3.9	4.3	0.4
NEW JERSEY	5.4	5.4	-0.0
NEW MEXICO	5.0	5.5	0.5
NEW YORK	6.0	5.7	-0.3
NORTH CAROLINA	6.4	6.2	-0.2
NORTH DAKOTA	4.2	5.0	0.8
OHIO	5.8	6.1	0.3
OKLAHOMA	7.8	5.8	-2.0
OREGON	5.3	4.6	-0.7
PENNSYLVANIA	5.6	6.4	0.8
RHODE ISLAND	6.0	5.3	-0.7
SOUTH CAROLINA	5.7	6.4	0.7
SOUTH DAKOTA	3.1	4.9	1.9
TENNESSEE	5.9	6.0	0.1
TEXAS	5.5	6.2	0.7
UTAH	4.4	4.2	-0.2
VERMONT	4.4	4.3	-0.1
VIRGINIA	4.5	6.1	1.6
WASHINGTON	4.9	5.2	0.3
WEST VIRGINIA	6.3	7.3	1.0
WISCONSIN	4.4	5.3	0.9
WYOMING	3.7	4.6	0.9
PUERTO RICO	9.6	9.6	0.0

\* EVER TOLD YOU HAVE DIABETES

June 20, 2000

TABLE 8.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD BLOOD PRESSURE IS HIGH\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2090	31.2	1.1	( 29.0, 33.4)
ALASKA	2046	21.3	1.4	( 18.6, 24.0)
ARIZONA	1710	14.2	1.1	( 12.1, 16.3)
ARKANSAS	2991	28.4	0.9	( 26.6, 30.2)
CALIFORNIA	4281	23.0	0.7	( 21.5, 24.4)
COLORADO	3036	22.2	1.0	( 20.1, 24.2)
CONNECTICUT	3494	20.4	0.9	( 18.7, 22.2)
DELAWARE	2545	25.5	1.3	( 23.1, 28.0)
DISTRICT OF CO	1261	24.7	1.3	( 22.2, 27.2)
FLORIDA	5140	27.7	0.8	( 26.2, 29.3)
GEORGIA	2262	26.3	1.1	( 24.2, 28.3)
HAWAII	2146	22.7	1.2	( 20.3, 25.1)
IDAHO	4929	23.0	0.7	( 21.6, 24.4)
ILLINOIS	1438	26.7	1.4	( 24.0, 29.4)
INDIANA	2427	25.7	1.5	( 22.8, 28.6)
IOWA	3610	24.2	0.8	( 22.6, 25.9)
KANSAS	3860	21.4	0.7	( 20.0, 22.8)
KENTUCKY	7487	27.5	0.7	( 26.2, 28.9)
LOUISIANA	1662	26.0	1.2	( 23.7, 28.3)
MAINE	1669	26.7	1.2	( 24.3, 29.0)
MARYLAND	3956	24.5	0.9	( 22.8, 26.2)
MASSACHUSETTS	4981	21.8	0.8	( 20.3, 23.3)
MICHIGAN	2508	25.2	0.9	( 23.3, 27.0)
MINNESOTA	5303	22.0	0.6	( 20.8, 23.2)
MISSISSIPPI	2174	33.5	1.2	( 31.2, 35.8)
MISSOURI	4262	24.6	0.9	( 22.9, 26.3)
MONTANA	1791	23.2	1.0	( 21.2, 25.2)
NEBRASKA	2815	22.0	0.9	( 20.3, 23.6)
NEVADA	2178	29.1	1.6	( 26.0, 32.1)
NEW HAMPSHIRE	1232	23.4	1.4	( 20.8, 26.1)
NEW JERSEY	2786	23.5	1.0	( 21.6, 25.4)
NEW MEXICO	3469	21.0	0.8	( 19.4, 22.5)
NEW YORK	2625	22.9	0.9	( 21.0, 24.7)
NORTH CAROLINA	2434	24.1	1.0	( 22.1, 26.0)
NORTH DAKOTA	1980	26.1	1.1	( 24.0, 28.1)
OHIO	1634	27.4	1.3	( 24.9, 29.9)
OKLAHOMA	2953	20.9	0.8	( 19.4, 22.5)
OREGON	1795	22.3	1.1	( 20.2, 24.4)
PENNSYLVANIA	3546	23.9	0.8	( 22.3, 25.5)
RHODE ISLAND	3982	22.9	0.8	( 21.4, 24.3)
SOUTH CAROLINA	3454	25.2	0.8	( 23.6, 26.8)
SOUTH DAKOTA	4136	23.8	0.7	( 22.4, 25.3)
TENNESSEE	3005	28.6	1.0	( 26.7, 30.4)
TEXAS	4943	24.2	0.8	( 22.7, 25.7)
UTAH	3184	21.3	1.0	( 19.4, 23.2)
VERMONT	3146	21.0	0.8	( 19.4, 22.5)
VIRGINIA	3488	23.5	1.0	( 21.5, 25.5)
WASHINGTON	3590	22.1	0.8	( 20.5, 23.7)
WEST VIRGINIA	2578	31.0	1.0	( 29.1, 32.9)
WISCONSIN	2160	25.0	1.0	( 23.0, 27.0)
WYOMING	2392	22.0	0.9	( 20.2, 23.8)
PUERTO RICO	3017	26.2	0.9	( 24.5, 28.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 24.0 RANGE = 14.2-33.5

\* AMONG RESPONDENTS HAVING THEIR BLOOD PRESSURE CHECKED

June 20, 2000

TABLE 8.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD BLOOD PRESSURE IS HIGH\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	851	28.6	1.7	( 25.3, 31.9)	*	1239	33.4	1.5	( 30.5, 36.3)	*
ALASKA	997	19.4	1.8	( 15.8, 22.9)	*	1049	23.4	2.0	( 19.4, 27.4)	*
ARIZONA	695	14.9	1.7	( 11.7, 18.2)	*	1015	13.6	1.4	( 10.8, 16.3)	*
ARKANSAS	1162	26.1	1.4	( 23.3, 28.8)	*	1829	30.4	1.2	( 28.1, 32.8)	*
CALIFORNIA	1819	22.0	1.1	( 19.8, 24.1)	*	2462	24.0	1.0	( 22.1, 25.9)	*
COLORADO	1360	21.9	1.6	( 18.8, 24.9)	*	1676	22.5	1.4	( 19.7, 25.2)	*
CONNECTICUT	1358	19.4	1.4	( 16.7, 22.1)	*	2136	21.4	1.2	( 19.0, 23.7)	*
DELAWARE	1017	26.1	1.8	( 22.5, 29.7)	*	1528	25.0	1.7	( 21.7, 28.3)	*
DISTRICT OF CO	456	24.8	2.1	( 20.6, 28.9)	*	805	24.7	1.6	( 21.5, 27.8)	*
FLORIDA	2035	27.4	1.2	( 24.9, 29.8)	*	3105	28.1	1.0	( 26.2, 30.0)	*
GEORGIA	881	25.4	1.6	( 22.2, 28.7)	*	1381	27.0	1.3	( 24.4, 29.6)	*
HAWAII	987	22.6	1.7	( 19.2, 26.0)	*	1159	22.8	1.7	( 19.4, 26.2)	*
IDAHO	2081	22.9	1.1	( 20.7, 25.2)	*	2848	23.0	0.9	( 21.2, 24.8)	*
ILLINOIS	569	26.9	2.2	( 22.5, 31.2)	*	869	26.6	1.7	( 23.3, 29.8)	*
INDIANA	1007	25.0	2.3	( 20.5, 29.4)	*	1420	26.3	1.9	( 22.6, 30.0)	*
IOWA	1419	22.2	1.3	( 19.7, 24.7)	*	2191	26.1	1.1	( 24.0, 28.2)	*
KANSAS	1551	20.1	1.1	( 18.0, 22.2)	*	2309	22.6	0.9	( 20.8, 24.5)	*
KENTUCKY	2720	24.7	1.1	( 22.6, 26.8)	*	4767	30.1	0.9	( 28.3, 31.8)	*
LOUISIANA	656	25.0	1.8	( 21.5, 28.5)	*	1006	26.9	1.5	( 23.9, 29.9)	*
MAINE	735	23.5	1.7	( 20.2, 26.9)	*	934	29.5	1.6	( 26.3, 32.7)	*
MARYLAND	1586	23.5	1.3	( 20.9, 26.0)	*	2370	25.5	1.1	( 23.3, 27.6)	*
MASSACHUSETTS	1984	22.9	1.3	( 20.3, 25.4)	*	2997	20.9	0.9	( 19.1, 22.7)	*
MICHIGAN	1031	23.9	1.4	( 21.1, 26.7)	*	1477	26.3	1.2	( 23.9, 28.8)	*
MINNESOTA	2419	20.8	0.9	( 19.0, 22.5)	*	2884	23.2	0.9	( 21.5, 24.8)	*
MISSISSIPPI	850	30.9	1.9	( 27.1, 34.6)	*	1324	35.8	1.4	( 33.0, 38.7)	*
MISSOURI	1715	22.1	1.3	( 19.6, 24.6)	*	2547	26.9	1.2	( 24.6, 29.2)	*
MONTANA	762	22.1	1.6	( 19.0, 25.1)	*	1029	24.3	1.4	( 21.5, 27.0)	*
NEBRASKA	1114	20.4	1.3	( 17.9, 22.9)	*	1701	23.4	1.1	( 21.2, 25.6)	*
NEVADA	1094	28.3	2.2	( 24.0, 32.7)	*	1084	29.8	2.3	( 25.3, 34.3)	*
NEW HAMPSHIRE	495	24.0	2.1	( 19.9, 28.0)	*	737	22.9	1.7	( 19.5, 26.3)	*
NEW JERSEY	1107	23.6	1.5	( 20.7, 26.6)	*	1679	23.3	1.2	( 20.9, 25.7)	*
NEW MEXICO	1521	20.2	1.1	( 18.0, 22.4)	*	1948	21.6	1.1	( 19.5, 23.7)	*
NEW YORK	1056	21.9	1.4	( 19.1, 24.6)	*	1569	23.8	1.2	( 21.4, 26.2)	*
NORTH CAROLINA	958	23.1	1.6	( 20.0, 26.1)	*	1476	25.0	1.2	( 22.5, 27.4)	*
NORTH DAKOTA	861	25.4	1.6	( 22.3, 28.5)	*	1119	26.7	1.4	( 23.9, 29.4)	*
OHIO	636	27.8	2.1	( 23.7, 31.8)	*	998	27.1	1.6	( 24.0, 30.2)	*
OKLAHOMA	1148	17.5	1.2	( 15.2, 19.8)	*	1805	24.1	1.1	( 21.9, 26.3)	*
OREGON	762	21.4	1.6	( 18.2, 24.6)	*	1033	23.1	1.4	( 20.4, 25.8)	*
PENNSYLVANIA	1411	23.9	1.3	( 21.4, 26.4)	*	2135	23.9	1.1	( 21.8, 25.9)	*
RHODE ISLAND	1551	21.5	1.2	( 19.2, 23.7)	*	2431	24.1	1.0	( 22.1, 26.0)	*
SOUTH CAROLINA	1384	23.3	1.2	( 20.8, 25.7)	*	2070	26.9	1.1	( 24.8, 29.1)	*
SOUTH DAKOTA	1675	22.9	1.1	( 20.8, 25.0)	*	2461	24.7	0.9	( 22.9, 26.6)	*
TENNESSEE	1032	27.2	1.5	( 24.3, 30.2)	*	1973	29.7	1.2	( 27.5, 32.0)	*
TEXAS	1971	23.8	1.1	( 21.6, 26.0)	*	2972	24.5	1.1	( 22.5, 26.6)	*
UTAH	1363	22.3	1.5	( 19.3, 25.2)	*	1821	20.3	1.3	( 17.9, 22.8)	*
VERMONT	1260	20.5	1.2	( 18.1, 22.9)	*	1886	21.4	1.0	( 19.4, 23.4)	*
VIRGINIA	1403	22.2	1.6	( 19.2, 25.2)	*	2085	24.8	1.3	( 22.1, 27.4)	*
WASHINGTON	1589	21.1	1.2	( 18.8, 23.3)	*	2001	23.1	1.2	( 20.8, 25.4)	*
WEST VIRGINIA	1060	32.0	1.6	( 28.9, 35.0)	*	1518	30.2	1.3	( 27.7, 32.6)	*
WISCONSIN	912	24.3	1.5	( 21.3, 27.3)	*	1248	25.5	1.4	( 22.9, 28.2)	*
WYOMING	1056	22.4	1.4	( 19.7, 25.2)	*	1336	21.6	1.2	( 19.3, 24.0)	*
PUERTO RICO	1099	25.2	1.4	( 22.5, 27.9)	*	1918	27.1	1.1	( 25.0, 29.3)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 23.2	MEDIAN	= 24.7
	RANGE	= 14.9-32.0	RANGE	= 13.6-35.8

\* AMONG RESPONDENTS HAVING THEIR BLOOD PRESSURE CHECKED

TABLE 8.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 EVER TOLD BLOOD PRESSURE IS HIGH\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	28.9	31.2	2.2
ALASKA	22.6	21.3	-1.3
ARIZONA	16.3	14.2	-2.1
ARKANSAS	26.3	28.4	2.0
CALIFORNIA	21.3	23.0	1.7
COLORADO	20.4	22.2	1.8
CONNECTICUT	20.6	20.4	-0.2
DELAWARE	25.5	25.5	-0.0
DISTRICT OF CO	19.4	24.7	5.3
FLORIDA	26.1	27.7	1.7
GEORGIA	21.4	26.3	4.8
HAWAII	24.0	22.7	-1.3
IDAHO	24.1	23.0	-1.1
ILLINOIS	24.3	26.7	2.4
INDIANA	25.2	25.7	0.5
IOWA	23.4	24.2	0.9
KANSAS	20.9	21.4	0.6
KENTUCKY	27.1	27.5	0.4
LOUISIANA	25.1	26.0	0.9
MAINE	22.8	26.7	3.8
MARYLAND	23.8	24.5	0.7
MASSACHUSETTS	19.8	21.8	2.0
MICHIGAN	23.3	25.2	1.9
MINNESOTA	21.2	22.0	0.8
MISSISSIPPI	34.4	33.5	-0.9
MISSOURI	27.3	24.6	-2.7
MONTANA	22.9	23.2	0.3
NEBRASKA	22.4	22.0	-0.4
NEVADA	24.1	29.1	5.0
NEW HAMPSHIRE	22.6	23.4	0.8
NEW JERSEY	23.6	23.5	-0.2
NEW MEXICO	21.3	21.0	-0.4
NEW YORK	22.7	22.9	0.2
NORTH CAROLINA	23.3	24.1	0.7
NORTH DAKOTA	25.5	26.1	0.6
OHIO	22.0	27.4	5.4
OKLAHOMA	21.7	20.9	-0.8
OREGON	22.8	22.3	-0.5
PENNSYLVANIA	21.7	23.9	2.2
RHODE ISLAND	22.5	22.9	0.4
SOUTH CAROLINA	26.9	25.2	-1.7
SOUTH DAKOTA	20.6	23.8	3.2
TENNESSEE	27.8	28.6	0.8
TEXAS	23.1	24.2	1.1
UTAH	22.5	21.3	-1.3
VERMONT	20.9	21.0	0.0
VIRGINIA	24.5	23.5	-0.9
WASHINGTON	23.2	22.1	-1.1
WEST VIRGINIA	28.3	31.0	2.7
WISCONSIN	23.1	25.0	1.8
WYOMING	22.1	22.0	-0.1
PUERTO RICO	20.9	26.2	5.4

\* AMONG RESPONDENTS HAVING THEIR BLOOD PRESSURE CHECKED

June 20, 2000

TABLE 9.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HYPERTENSION SCREENING IN PAST TWO YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2079	4.8	0.6	( 3.7, 5.9)
ALASKA	2018	6.2	0.8	( 4.6, 7.7)
ARIZONA	1717	7.2	1.0	( 5.3, 9.1)
ARKANSAS	2981	6.7	0.6	( 5.7, 7.8)
CALIFORNIA	4284	8.0	0.5	( 7.0, 8.9)
COLORADO	3038	7.8	0.7	( 6.4, 9.3)
CONNECTICUT	3484	4.8	0.5	( 3.8, 5.9)
DELAWARE	2549	3.8	0.5	( 2.8, 4.9)
DISTRICT OF CO	1254	3.9	0.7	( 2.6, 5.1)
FLORIDA	5133	5.4	0.4	( 4.7, 6.2)
GEORGIA	2248	4.4	0.5	( 3.5, 5.3)
HAWAII	2132	3.5	0.5	( 2.5, 4.5)
IDAHO	4908	9.4	0.5	( 8.3, 10.4)
ILLINOIS	1439	6.5	0.8	( 4.9, 8.0)
INDIANA	2443	7.7	0.9	( 5.9, 9.5)
IOWA	3590	5.4	0.5	( 4.5, 6.3)
KANSAS	3825	4.5	0.4	( 3.8, 5.3)
KENTUCKY	7441	5.6	0.4	( 4.8, 6.4)
LOUISIANA	1650	4.2	0.6	( 3.1, 5.4)
MAINE	1663	5.2	0.6	( 4.0, 6.4)
MARYLAND	3937	3.4	0.4	( 2.6, 4.1)
MASSACHUSETTS	4978	4.1	0.4	( 3.3, 4.9)
MICHIGAN	2496	4.9	0.5	( 3.9, 5.8)
MINNESOTA	5325	5.6	0.4	( 4.9, 6.3)
MISSISSIPPI	2154	4.7	0.6	( 3.6, 5.7)
MISSOURI	4240	4.9	0.5	( 4.0, 5.8)
MONTANA	1768	6.7	0.7	( 5.4, 8.1)
NEBRASKA	2802	5.4	0.5	( 4.4, 6.5)
NEVADA	2183	8.1	0.8	( 6.5, 9.6)
NEW HAMPSHIRE	1243	5.5	0.8	( 3.9, 7.0)
NEW JERSEY	2795	5.0	0.6	( 3.8, 6.2)
NEW MEXICO	3449	9.7	0.6	( 8.6, 10.8)
NEW YORK	2619	6.4	0.6	( 5.1, 7.6)
NORTH CAROLINA	2424	3.8	0.5	( 2.9, 4.7)
NORTH DAKOTA	1966	4.5	0.5	( 3.5, 5.6)
OHIO	1630	3.9	0.5	( 2.8, 4.9)
OKLAHOMA	2942	4.8	0.5	( 3.8, 5.7)
OREGON	1795	8.5	0.8	( 7.0, 10.0)
PENNSYLVANIA	3515	3.9	0.4	( 3.2, 4.7)
RHODE ISLAND	3964	3.3	0.3	( 2.6, 4.0)
SOUTH CAROLINA	3416	4.8	0.5	( 3.9, 5.7)
SOUTH DAKOTA	4118	5.7	0.4	( 4.8, 6.5)
TENNESSEE	2988	4.1	0.4	( 3.2, 4.9)
TEXAS	4931	8.7	0.6	( 7.4, 9.9)
UTAH	3177	7.6	0.7	( 6.4, 8.9)
VERMONT	3119	5.7	0.5	( 4.7, 6.7)
VIRGINIA	3477	6.4	0.7	( 5.0, 7.8)
WASHINGTON	3596	7.1	0.6	( 5.9, 8.4)
WEST VIRGINIA	2569	6.2	0.5	( 5.1, 7.2)
WISCONSIN	2177	7.4	0.6	( 6.1, 8.6)
WYOMING	2388	7.7	0.6	( 6.4, 9.0)
PUERTO RICO	2918	5.1	0.6	( 4.1, 6.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 5.4 RANGE = 3.3- 9.7

\* BLOOD PRESSURE NOT CHECKED BY DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL June 20, 2000

TABLE 9.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HYPERTENSION SCREENING IN PAST TWO YEARS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	847	6.5	0.9	( 4.7, 8.3)	*	1232	3.3	0.7	( 2.0, 4.6)	*
ALASKA	981	8.7	1.3	( 6.2, 11.3)	*	1037	3.3	0.8	( 1.8, 4.8)	*
ARIZONA	698	8.2	1.5	( 5.3, 11.0)	*	1019	6.3	1.3	( 3.9, 8.8)	*
ARKANSAS	1158	8.9	1.0	( 7.0, 10.8)	*	1823	4.8	0.6	( 3.7, 5.9)	*
CALIFORNIA	1825	11.4	0.9	( 9.7, 13.0)	*	2459	4.7	0.5	( 3.7, 5.7)	*
COLORADO	1365	11.3	1.3	( 8.8, 13.8)	*	1673	4.5	0.8	( 3.0, 6.1)	*
CONNECTICUT	1355	7.5	1.0	( 5.6, 9.5)	*	2129	2.4	0.5	( 1.5, 3.3)	*
DELAWARE	1020	5.9	1.0	( 4.0, 7.8)	*	1529	1.9	0.4	( 1.1, 2.8)	*
DISTRICT OF CO	451	5.1	1.2	( 2.8, 7.4)	*	803	2.8	0.7	( 1.5, 4.1)	*
FLORIDA	2034	6.7	0.6	( 5.5, 8.0)	*	3099	4.2	0.4	( 3.3, 5.1)	*
GEORGIA	875	6.2	0.9	( 4.5, 7.9)	*	1373	2.7	0.4	( 1.8, 3.5)	*
HAWAII	982	5.0	0.9	( 3.2, 6.7)	*	1150	2.0	0.5	( 0.9, 3.0)	*
IDAHO	2073	12.6	0.9	( 10.8, 14.4)	*	2835	6.3	0.5	( 5.3, 7.3)	*
ILLINOIS	568	10.0	1.5	( 7.1, 12.9)	*	871	3.0	0.6	( 1.8, 4.2)	*
INDIANA	1019	9.5	1.5	( 6.5, 12.5)	*	1424	6.1	1.1	( 3.9, 8.3)	*
IOWA	1412	7.7	0.8	( 6.1, 9.3)	*	2178	3.4	0.4	( 2.6, 4.2)	*
KANSAS	1540	6.7	0.7	( 5.3, 8.0)	*	2285	2.6	0.4	( 1.9, 3.3)	*
KENTUCKY	2703	7.5	0.7	( 6.1, 8.9)	*	4738	3.9	0.4	( 3.2, 4.6)	*
LOUISIANA	651	5.4	1.0	( 3.4, 7.4)	*	999	3.2	0.6	( 2.0, 4.4)	*
MAINE	733	8.2	1.2	( 5.9, 10.4)	*	930	2.4	0.5	( 1.5, 3.4)	*
MARYLAND	1582	5.0	0.7	( 3.6, 6.3)	*	2355	1.9	0.3	( 1.2, 2.5)	*
MASSACHUSETTS	1983	5.6	0.7	( 4.3, 7.0)	*	2995	2.8	0.5	( 1.9, 3.7)	*
MICHIGAN	1028	6.5	0.8	( 4.9, 8.0)	*	1468	3.4	0.6	( 2.3, 4.6)	*
MINNESOTA	2430	7.4	0.6	( 6.2, 8.5)	*	2895	3.9	0.4	( 3.1, 4.7)	*
MISSISSIPPI	839	6.7	1.0	( 4.7, 8.6)	*	1315	2.9	0.6	( 1.8, 3.9)	*
MISSOURI	1699	6.9	0.8	( 5.4, 8.5)	*	2541	3.1	0.5	( 2.2, 3.9)	*
MONTANA	755	9.2	1.2	( 6.9, 11.5)	*	1013	4.4	0.7	( 3.0, 5.7)	*
NEBRASKA	1109	8.2	0.9	( 6.4, 10.0)	*	1693	2.9	0.6	( 1.8, 4.0)	*
NEVADA	1100	11.6	1.4	( 8.9, 14.2)	*	1083	4.5	0.8	( 3.0, 6.0)	*
NEW HAMPSHIRE	500	7.1	1.4	( 4.4, 9.8)	*	743	3.9	0.8	( 2.4, 5.4)	*
NEW JERSEY	1107	6.9	1.1	( 4.7, 9.1)	*	1688	3.3	0.6	( 2.2, 4.4)	*
NEW MEXICO	1514	12.5	0.9	( 10.7, 14.3)	*	1935	7.0	0.7	( 5.7, 8.3)	*
NEW YORK	1052	9.6	1.1	( 7.4, 11.7)	*	1567	3.6	0.7	( 2.2, 5.0)	*
NORTH CAROLINA	953	5.0	0.8	( 3.3, 6.6)	*	1471	2.8	0.5	( 1.9, 3.7)	*
NORTH DAKOTA	850	6.2	0.9	( 4.4, 7.9)	*	1116	3.0	0.6	( 1.9, 4.1)	*
OHIO	629	4.4	0.8	( 2.8, 5.9)	*	1001	3.5	0.8	( 2.0, 5.0)	*
OKLAHOMA	1138	6.4	0.8	( 4.9, 8.0)	*	1804	3.2	0.5	( 2.2, 4.2)	*
OREGON	761	12.8	1.4	( 10.1, 15.5)	*	1034	4.5	0.8	( 3.0, 5.9)	*
PENNSYLVANIA	1401	5.3	0.7	( 4.0, 6.7)	*	2114	2.7	0.4	( 1.9, 3.5)	*
RHODE ISLAND	1537	4.2	0.6	( 3.1, 5.3)	*	2427	2.5	0.4	( 1.8, 3.3)	*
SOUTH CAROLINA	1372	7.3	0.9	( 5.5, 9.0)	*	2044	2.6	0.4	( 1.8, 3.3)	*
SOUTH DAKOTA	1670	8.0	0.7	( 6.6, 9.4)	*	2448	3.5	0.4	( 2.6, 4.3)	*
TENNESSEE	1024	5.9	0.8	( 4.3, 7.4)	*	1964	2.4	0.4	( 1.7, 3.2)	*
TEXAS	1966	13.2	1.2	( 10.8, 15.5)	*	2965	4.5	0.4	( 3.6, 5.3)	*
UTAH	1362	10.8	1.2	( 8.5, 13.0)	*	1815	4.7	0.6	( 3.4, 5.9)	*
VERMONT	1253	8.2	0.9	( 6.4, 10.0)	*	1866	3.4	0.4	( 2.6, 4.2)	*
VIRGINIA	1401	8.7	1.2	( 6.5, 11.0)	*	2076	4.2	0.9	( 2.5, 5.8)	*
WASHINGTON	1594	10.4	1.1	( 8.2, 12.6)	*	2002	4.0	0.5	( 3.1, 4.9)	*
WEST VIRGINIA	1057	8.0	0.9	( 6.3, 9.7)	*	1512	4.5	0.6	( 3.4, 5.7)	*
WISCONSIN	921	10.5	1.1	( 8.3, 12.7)	*	1256	4.4	0.6	( 3.2, 5.6)	*
WYOMING	1053	11.1	1.1	( 8.9, 13.2)	*	1335	4.5	0.6	( 3.3, 5.6)	*
PUERTO RICO	1073	7.2	1.0	( 5.2, 9.2)	*	1845	3.3	0.5	( 2.3, 4.3)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 7.5	MEDIAN	= 3.4
	RANGE	= 4.2-13.2	RANGE	= 1.9- 7.0

\* BLOOD PRESSURE NOT CHECKED BY DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

June 20, 2000

TABLE 9.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 NO HYPERTENSION SCREENING IN PAST TWO YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	4.6	4.8	0.3
ALASKA	8.2	6.2	-2.0
ARIZONA	6.5	7.2	0.8
ARKANSAS	6.7	6.7	0.1
CALIFORNIA	9.3	8.0	-1.3
COLORADO	6.9	7.8	1.0
CONNECTICUT	4.9	4.8	-0.1
DELAWARE	5.7	3.8	-1.9
DISTRICT OF CO	2.7	3.9	1.1
FLORIDA	5.8	5.4	-0.3
GEORGIA	3.6	4.4	0.8
HAWAII	4.5	3.5	-1.1
IDAHO	8.8	9.4	0.6
ILLINOIS	5.7	6.5	0.8
INDIANA	6.9	7.7	0.8
IOWA	7.4	5.4	-1.9
KANSAS	5.3	4.5	-0.7
KENTUCKY	6.2	5.6	-0.6
LOUISIANA	5.6	4.2	-1.4
MAINE	5.5	5.2	-0.3
MARYLAND	3.6	3.4	-0.3
MASSACHUSETTS	4.0	4.1	0.1
MICHIGAN	5.1	4.9	-0.2
MINNESOTA	7.5	5.6	-1.9
MISSISSIPPI	4.5	4.7	0.2
MISSOURI	5.3	4.9	-0.4
MONTANA	7.7	6.7	-1.0
NEBRASKA	6.9	5.4	-1.4
NEVADA	6.7	8.1	1.3
NEW HAMPSHIRE	6.0	5.5	-0.5
NEW JERSEY	5.6	5.0	-0.6
NEW MEXICO	9.2	9.7	0.4
NEW YORK	5.2	6.4	1.2
NORTH CAROLINA	6.1	3.8	-2.2
NORTH DAKOTA	7.0	4.5	-2.5
OHIO	4.5	3.9	-0.6
OKLAHOMA	5.8	4.8	-1.1
OREGON	7.8	8.5	0.7
PENNSYLVANIA	4.4	3.9	-0.5
RHODE ISLAND	4.3	3.3	-1.0
SOUTH CAROLINA	3.5	4.8	1.3
SOUTH DAKOTA	7.3	5.7	-1.7
TENNESSEE	4.9	4.1	-0.8
TEXAS	7.6	8.7	1.1
UTAH	7.9	7.6	-0.2
VERMONT	6.5	5.7	-0.8
VIRGINIA	6.1	6.4	0.3
WASHINGTON	7.1	7.1	0.0
WEST VIRGINIA	6.4	6.2	-0.3
WISCONSIN	7.4	7.4	-0.0
WYOMING	9.3	7.7	-1.6
PUERTO RICO	5.1	5.1	0.0

\* BLOOD PRESSURE NOT CHECKED BY DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL  
 June 20, 2000

TABLE 10.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CHOLESTEROL SCREENING\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2035	26.3	1.2	( 24.0, 28.5)
ALASKA	2001	30.0	1.6	( 26.9, 33.0)
ARIZONA	1694	23.2	1.6	( 20.1, 26.2)
ARKANSAS	2915	30.0	1.0	( 28.0, 31.9)
CALIFORNIA	4230	28.2	0.8	( 26.6, 29.8)
COLORADO	2987	27.9	1.2	( 25.5, 30.3)
CONNECTICUT	3427	20.7	1.0	( 18.7, 22.6)
DELAWARE	2505	23.2	1.3	( 20.7, 25.7)
DISTRICT OF CO	1253	19.3	1.4	( 16.5, 22.0)
FLORIDA	5061	20.3	0.7	( 18.9, 21.7)
GEORGIA	2198	22.5	1.1	( 20.4, 24.7)
HAWAII	2095	29.8	1.4	( 27.1, 32.4)
IDAHO	4808	29.0	0.8	( 27.4, 30.6)
ILLINOIS	1431	27.3	1.4	( 24.6, 30.0)
INDIANA	2378	24.6	1.5	( 21.6, 27.6)
IOWA	3535	26.8	0.9	( 25.0, 28.6)
KANSAS	3776	27.4	0.9	( 25.7, 29.0)
KENTUCKY	7268	27.5	0.8	( 26.0, 29.0)
LOUISIANA	1617	29.7	1.3	( 27.1, 32.3)
MAINE	1633	22.7	1.3	( 20.2, 25.1)
MARYLAND	3859	19.5	0.8	( 17.9, 21.1)
MASSACHUSETTS	4894	19.4	0.8	( 17.9, 20.9)
MICHIGAN	2469	23.3	1.0	( 21.3, 25.2)
MINNESOTA	5249	35.9	0.8	( 34.5, 37.4)
MISSISSIPPI	2073	29.2	1.2	( 26.8, 31.5)
MISSOURI	4165	29.7	1.0	( 27.7, 31.6)
MONTANA	1741	28.7	1.3	( 26.2, 31.1)
NEBRASKA	2746	29.8	1.0	( 27.8, 31.8)
NEVADA	2144	26.5	1.5	( 23.6, 29.5)
NEW HAMPSHIRE	1207	20.9	1.5	( 18.0, 23.8)
NEW JERSEY	2769	21.2	1.0	( 19.2, 23.2)
NEW MEXICO	3368	32.8	0.9	( 31.0, 34.6)
NEW YORK	2590	23.2	1.0	( 21.2, 25.1)
NORTH CAROLINA	2374	22.8	1.1	( 20.7, 24.9)
NORTH DAKOTA	1930	29.6	1.2	( 27.4, 31.9)
OHIO	1608	26.6	1.4	( 23.9, 29.3)
OKLAHOMA	2874	24.9	1.0	( 23.0, 26.8)
OREGON	1752	26.6	1.2	( 24.2, 29.0)
PENNSYLVANIA	3476	24.0	0.8	( 22.3, 25.6)
RHODE ISLAND	3911	21.3	0.8	( 19.7, 22.8)
SOUTH CAROLINA	3391	23.9	0.9	( 22.2, 25.6)
SOUTH DAKOTA	3994	29.6	0.9	( 28.0, 31.3)
TENNESSEE	2926	23.5	0.9	( 21.7, 25.3)
TEXAS	4865	27.4	0.9	( 25.7, 29.1)
UTAH	3128	30.3	1.2	( 28.0, 32.5)
VERMONT	3081	24.4	1.0	( 22.5, 26.2)
VIRGINIA	3402	21.4	1.0	( 19.5, 23.4)
WASHINGTON	3498	24.9	1.0	( 23.0, 26.7)
WEST VIRGINIA	2525	27.2	1.0	( 25.2, 29.3)
WISCONSIN	2106	23.4	1.1	( 21.3, 25.5)
WYOMING	2358	26.8	1.1	( 24.8, 28.9)
PUERTO RICO	3020	24.1	1.0	( 22.2, 26.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 26.4 RANGE = 19.3-35.9

\* NEVER HAD CHOLESTEROL SCREENING

June 20, 2000



TABLE 10.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 NO CHOLESTEROL SCREENING\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	25.3	26.3	1.0
ALASKA	32.6	30.0	-2.6
ARIZONA	29.1	23.2	-5.9
ARKANSAS	36.5	30.0	-6.6
CALIFORNIA	30.1	28.2	-1.9
COLORADO	24.4	27.9	3.6
CONNECTICUT	22.1	20.7	-1.4
DELAWARE	25.9	23.2	-2.7
DISTRICT OF CO	18.7	19.3	0.6
FLORIDA	21.0	20.3	-0.7
GEORGIA	24.2	22.5	-1.6
HAWAII	25.3	29.8	4.5
IDAHO	27.3	29.0	1.8
ILLINOIS	28.1	27.3	-0.8
INDIANA	28.1	24.6	-3.5
IOWA	27.1	26.8	-0.3
KANSAS	41.6	27.4	-14.2
KENTUCKY	28.9	27.5	-1.4
LOUISIANA	30.2	29.7	-0.5
MAINE	22.5	22.7	0.1
MARYLAND	22.7	19.5	-3.1
MASSACHUSETTS	20.9	19.4	-1.5
MICHIGAN	24.4	23.3	-1.1
MINNESOTA	34.2	35.9	1.8
MISSISSIPPI	33.6	29.2	-4.4
MISSOURI	25.6	29.7	4.1
MONTANA	28.6	28.7	0.1
NEBRASKA	26.7	29.8	3.1
NEVADA	28.3	26.5	-1.8
NEW HAMPSHIRE	22.3	20.9	-1.4
NEW JERSEY	21.9	21.2	-0.7
NEW MEXICO	32.0	32.8	0.8
NEW YORK	24.1	23.2	-0.9
NORTH CAROLINA	24.4	22.8	-1.6
NORTH DAKOTA	28.2	29.6	1.4
OHIO	30.0	26.6	-3.4
OKLAHOMA	22.6	24.9	2.4
OREGON	26.1	26.6	0.5
PENNSYLVANIA	27.8	24.0	-3.8
RHODE ISLAND	21.7	21.3	-0.4
SOUTH CAROLINA	25.1	23.9	-1.2
SOUTH DAKOTA	31.4	29.6	-1.8
TENNESSEE	26.9	23.5	-3.4
TEXAS	29.0	27.4	-1.7
UTAH	28.1	30.3	2.2
VERMONT	25.5	24.4	-1.1
VIRGINIA	21.7	21.4	-0.3
WASHINGTON	24.2	24.9	0.7
WEST VIRGINIA	29.5	27.2	-2.3
WISCONSIN	24.4	23.4	-0.9
WYOMING	25.0	26.8	1.9
PUERTO RICO	19.1	24.1	5.1

\* NEVER HAD CHOLESTEROL SCREENING

June 20, 2000

TABLE 11.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CHOLESTEROL SCREENING IN PAST FIVE YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1993	30.4	1.2	( 28.0, 32.8)
ALASKA	1981	35.1	1.6	( 31.9, 38.2)
ARIZONA	1679	24.6	1.6	( 21.5, 27.7)
ARKANSAS	2892	35.4	1.0	( 33.4, 37.4)
CALIFORNIA	4217	32.5	0.8	( 30.9, 34.2)
COLORADO	2962	33.3	1.3	( 30.8, 35.8)
CONNECTICUT	3375	24.8	1.1	( 22.7, 26.8)
DELAWARE	2485	26.5	1.3	( 23.9, 29.0)
DISTRICT OF CO	1243	19.7	1.4	( 16.9, 22.5)
FLORIDA	5002	24.0	0.8	( 22.5, 25.5)
GEORGIA	2173	26.6	1.1	( 24.4, 28.8)
HAWAII	2075	32.7	1.4	( 30.0, 35.5)
IDAHO	4755	36.2	0.9	( 34.5, 37.9)
ILLINOIS	1418	31.5	1.4	( 28.7, 34.3)
INDIANA	2357	29.5	1.6	( 26.4, 32.6)
IOWA	3468	32.7	1.0	( 30.8, 34.6)
KANSAS	3705	31.1	0.9	( 29.4, 32.8)
KENTUCKY	7162	32.8	0.8	( 31.2, 34.4)
LOUISIANA	1597	32.9	1.4	( 30.2, 35.6)
MAINE	1616	26.8	1.3	( 24.2, 29.4)
MARYLAND	3830	23.5	0.9	( 21.7, 25.2)
MASSACHUSETTS	4817	22.8	0.8	( 21.2, 24.4)
MICHIGAN	2441	28.6	1.0	( 26.6, 30.6)
MINNESOTA	5196	40.2	0.8	( 38.7, 41.7)
MISSISSIPPI	2032	33.5	1.3	( 31.1, 36.0)
MISSOURI	4113	34.3	1.0	( 32.2, 36.3)
MONTANA	1708	35.0	1.3	( 32.4, 37.6)
NEBRASKA	2733	35.3	1.0	( 33.3, 37.4)
NEVADA	2137	32.3	1.6	( 29.3, 35.4)
NEW HAMPSHIRE	1192	25.7	1.6	( 22.7, 28.8)
NEW JERSEY	2741	24.5	1.0	( 22.5, 26.6)
NEW MEXICO	3320	37.9	1.0	( 36.0, 39.7)
NEW YORK	2557	27.6	1.1	( 25.6, 29.7)
NORTH CAROLINA	2357	25.7	1.1	( 23.5, 27.8)
NORTH DAKOTA	1913	35.7	1.2	( 33.3, 38.1)
OHIO	1584	31.3	1.4	( 28.5, 34.1)
OKLAHOMA	2835	27.7	1.0	( 25.7, 29.7)
OREGON	1725	32.9	1.3	( 30.4, 35.4)
PENNSYLVANIA	3429	28.5	0.9	( 26.7, 30.2)
RHODE ISLAND	3860	24.3	0.8	( 22.6, 25.9)
SOUTH CAROLINA	3333	27.7	0.9	( 25.9, 29.6)
SOUTH DAKOTA	3922	36.4	0.9	( 34.7, 38.2)
TENNESSEE	2888	27.0	1.0	( 25.1, 28.8)
TEXAS	4825	30.8	0.9	( 29.1, 32.5)
UTAH	3104	36.4	1.3	( 34.0, 38.9)
VERMONT	3028	29.6	1.0	( 27.7, 31.5)
VIRGINIA	3360	27.1	1.1	( 24.9, 29.3)
WASHINGTON	3459	31.6	1.0	( 29.6, 33.6)
WEST VIRGINIA	2499	31.0	1.1	( 28.9, 33.1)
WISCONSIN	2088	29.9	1.1	( 27.6, 32.1)
WYOMING	2338	31.4	1.1	( 29.2, 33.5)
PUERTO RICO	2958	26.7	1.0	( 24.7, 28.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 30.9 RANGE = 19.7-40.2 YEAR 2000 OBJ. 15.14  
 \* DID NOT HAVE A CHOLESTEROL SCREENING June 20, 2000

TABLE 11.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CHOLESTEROL SCREENING IN PAST FIVE YEARS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	MALES					FEMALES				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	821	30.8	1.9	( 27.1, 34.5)	*	1172	30.1	1.6	( 27.0, 33.2)	*
ALASKA	972	37.1	2.2	( 32.7, 41.5)	*	1009	32.7	2.4	( 28.1, 37.4)	*
ARIZONA	696	25.1	2.4	( 20.4, 29.7)	*	983	24.1	2.1	( 20.1, 28.2)	*
ARKANSAS	1136	38.8	1.6	( 35.6, 42.0)	*	1756	32.3	1.3	( 29.8, 34.8)	*
CALIFORNIA	1800	36.6	1.3	( 34.0, 39.1)	*	2417	28.6	1.1	( 26.5, 30.7)	*
COLORADO	1336	35.2	2.0	( 31.3, 39.1)	*	1626	31.5	1.7	( 28.2, 34.7)	*
CONNECTICUT	1323	27.3	1.7	( 24.1, 30.6)	*	2052	22.5	1.3	( 20.0, 24.9)	*
DELAWARE	998	27.4	2.0	( 23.4, 31.3)	*	1487	25.6	1.8	( 22.2, 29.1)	*
DISTRICT OF CO	449	21.1	2.4	( 16.4, 25.9)	*	794	18.6	1.7	( 15.2, 21.9)	*
FLORIDA	1993	27.6	1.2	( 25.2, 30.0)	*	3009	20.7	0.9	( 19.0, 22.4)	*
GEORGIA	858	27.7	1.8	( 24.2, 31.1)	*	1315	25.6	1.5	( 22.7, 28.5)	*
HAWAII	958	34.0	2.1	( 29.9, 38.2)	*	1117	31.5	1.8	( 27.9, 35.0)	*
IDAHO	2015	38.8	1.3	( 36.2, 41.4)	*	2740	33.7	1.1	( 31.7, 35.8)	*
ILLINOIS	562	34.1	2.3	( 29.6, 38.6)	*	856	29.1	1.7	( 25.7, 32.4)	*
INDIANA	984	30.6	2.5	( 25.8, 35.5)	*	1373	28.4	2.0	( 24.4, 32.4)	*
IOWA	1367	35.2	1.6	( 32.2, 38.3)	*	2101	30.5	1.2	( 28.2, 32.8)	*
KANSAS	1494	32.2	1.4	( 29.5, 34.9)	*	2211	30.1	1.1	( 27.9, 32.2)	*
KENTUCKY	2607	35.6	1.3	( 33.0, 38.2)	*	4555	30.3	1.0	( 28.4, 32.2)	*
LOUISIANA	627	35.1	2.2	( 30.8, 39.4)	*	970	31.0	1.7	( 27.6, 34.4)	*
MAINE	709	29.5	2.1	( 25.4, 33.6)	*	907	24.3	1.6	( 21.0, 27.5)	*
MARYLAND	1545	26.1	1.4	( 23.3, 28.9)	*	2285	21.1	1.1	( 19.0, 23.3)	*
MASSACHUSETTS	1920	25.1	1.4	( 22.4, 27.8)	*	2897	20.7	1.0	( 18.8, 22.6)	*
MICHIGAN	1009	33.2	1.7	( 29.9, 36.4)	*	1432	24.5	1.3	( 22.0, 27.0)	*
MINNESOTA	2380	44.6	1.2	( 42.3, 46.8)	*	2816	36.1	1.0	( 34.1, 38.0)	*
MISSISSIPPI	806	36.5	2.1	( 32.4, 40.5)	*	1226	30.9	1.5	( 28.0, 33.8)	*
MISSOURI	1655	36.9	1.7	( 33.6, 40.1)	*	2458	31.9	1.3	( 29.4, 34.5)	*
MONTANA	732	36.3	2.0	( 32.4, 40.2)	*	976	33.8	1.7	( 30.4, 37.1)	*
NEBRASKA	1089	38.2	1.6	( 35.0, 41.4)	*	1644	32.7	1.3	( 30.1, 35.3)	*
NEVADA	1079	35.9	2.3	( 31.4, 40.5)	*	1058	28.7	2.1	( 24.6, 32.8)	*
NEW HAMPSHIRE	477	28.2	2.5	( 23.3, 33.0)	*	715	23.4	1.9	( 19.7, 27.1)	*
NEW JERSEY	1091	26.0	1.7	( 22.7, 29.3)	*	1650	23.2	1.3	( 20.7, 25.8)	*
NEW MEXICO	1471	40.8	1.4	( 38.0, 43.6)	*	1849	35.0	1.3	( 32.6, 37.5)	*
NEW YORK	1042	29.4	1.7	( 26.2, 32.6)	*	1515	26.0	1.3	( 23.4, 28.6)	*
NORTH CAROLINA	934	28.3	1.7	( 25.0, 31.6)	*	1423	23.3	1.4	( 20.5, 26.0)	*
NORTH DAKOTA	834	37.5	1.9	( 33.9, 41.2)	*	1079	33.9	1.6	( 30.7, 37.1)	*
OHIO	621	33.6	2.3	( 29.0, 38.1)	*	963	29.2	1.8	( 25.6, 32.7)	*
OKLAHOMA	1105	29.8	1.7	( 26.5, 33.0)	*	1730	25.8	1.3	( 23.3, 28.3)	*
OREGON	737	38.8	2.0	( 34.9, 42.7)	*	988	27.4	1.6	( 24.2, 30.5)	*
PENNSYLVANIA	1371	28.5	1.4	( 25.7, 31.2)	*	2058	28.5	1.1	( 26.3, 30.7)	*
RHODE ISLAND	1498	27.1	1.4	( 24.4, 29.7)	*	2362	21.8	1.0	( 19.8, 23.8)	*
SOUTH CAROLINA	1346	30.9	1.5	( 27.9, 33.8)	*	1987	24.9	1.1	( 22.7, 27.0)	*
SOUTH DAKOTA	1600	38.4	1.4	( 35.7, 41.1)	*	2322	34.6	1.1	( 32.4, 36.8)	*
TENNESSEE	990	29.7	1.6	( 26.5, 32.8)	*	1898	24.6	1.1	( 22.4, 26.7)	*
TEXAS	1942	34.5	1.4	( 31.7, 37.3)	*	2883	27.3	1.1	( 25.2, 29.4)	*
UTAH	1346	38.4	2.0	( 34.6, 42.3)	*	1758	34.5	1.6	( 31.4, 37.6)	*
VERMONT	1222	31.7	1.5	( 28.8, 34.7)	*	1806	27.6	1.3	( 25.1, 30.1)	*
VIRGINIA	1358	28.5	1.8	( 25.1, 31.9)	*	2002	25.8	1.5	( 22.9, 28.6)	*
WASHINGTON	1539	35.0	1.6	( 31.8, 38.1)	*	1920	28.4	1.3	( 25.9, 30.8)	*
WEST VIRGINIA	1025	33.6	1.7	( 30.3, 36.9)	*	1474	28.7	1.4	( 26.0, 31.4)	*
WISCONSIN	886	34.1	1.8	( 30.6, 37.6)	*	1202	25.9	1.4	( 23.1, 28.7)	*
WYOMING	1033	32.2	1.6	( 29.0, 35.4)	*	1305	30.7	1.5	( 27.8, 33.5)	*
PUERTO RICO	1084	31.6	1.7	( 28.3, 35.0)	*	1874	22.3	1.2	( 20.0, 24.6)	*

SUMMARY STATISTICS:      NUMBER OF PARTICIPANTS = 52      NUMBER OF PARTICIPANTS = 52  
 MEDIAN                              = 33.4      MEDIAN                              = 28.5  
 RANGE                                  = 21.1-44.6      RANGE                                  = 18.6-36.1

\* DID NOT HAVE A CHOLESTEROL SCREENING

June 20, 2000

TABLE 11.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 NO CHOLESTEROL SCREENING IN PAST FIVE YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	29.0	30.4	1.4
ALASKA	37.6	35.1	-2.6
ARIZONA	31.5	24.6	-7.0
ARKANSAS	41.4	35.4	-6.0
CALIFORNIA	34.2	32.5	-1.7
COLORADO	30.3	33.3	3.0
CONNECTICUT	26.7	24.8	-1.9
DELAWARE	30.7	26.5	-4.3
DISTRICT OF CO	20.6	19.7	-0.9
FLORIDA	24.7	24.0	-0.7
GEORGIA	27.4	26.6	-0.8
HAWAII	30.5	32.7	2.2
IDAHO	35.0	36.2	1.2
ILLINOIS	32.8	31.5	-1.3
INDIANA	33.6	29.5	-4.2
IOWA	33.9	32.7	-1.2
KANSAS	45.1	31.1	-14.0
KENTUCKY	34.4	32.8	-1.6
LOUISIANA	33.7	32.9	-0.8
MAINE	28.2	26.8	-1.4
MARYLAND	25.5	23.5	-2.0
MASSACHUSETTS	25.3	22.8	-2.5
MICHIGAN	28.9	28.6	-0.3
MINNESOTA	38.8	40.2	1.4
MISSISSIPPI	37.5	33.5	-4.0
MISSOURI	29.6	34.3	4.7
MONTANA	36.8	35.0	-1.8
NEBRASKA	34.5	35.3	0.9
NEVADA	31.6	32.3	0.8
NEW HAMPSHIRE	27.0	25.7	-1.2
NEW JERSEY	24.7	24.5	-0.1
NEW MEXICO	37.2	37.9	0.6
NEW YORK	27.5	27.6	0.1
NORTH CAROLINA	27.9	25.7	-2.2
NORTH DAKOTA	36.2	35.7	-0.5
OHIO	33.5	31.3	-2.3
OKLAHOMA	25.6	27.7	2.1
OREGON	31.8	32.9	1.1
PENNSYLVANIA	31.5	28.5	-3.0
RHODE ISLAND	25.3	24.3	-1.0
SOUTH CAROLINA	27.8	27.7	-0.1
SOUTH DAKOTA	36.9	36.4	-0.4
TENNESSEE	29.8	27.0	-2.8
TEXAS	32.6	30.8	-1.8
UTAH	34.4	36.4	2.0
VERMONT	31.1	29.6	-1.5
VIRGINIA	26.4	27.1	0.6
WASHINGTON	30.4	31.6	1.2
WEST VIRGINIA	32.8	31.0	-1.8
WISCONSIN	29.8	29.9	0.1
WYOMING	29.8	31.4	1.6
PUERTO RICO	20.7	26.7	6.0

\* DID NOT HAVE A CHOLESTEROL SCREENING

June 20, 2000

TABLE 12.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD CHOLESTEROL IS HIGH\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1528	33.2	1.3	( 30.5, 35.8)
ALASKA	1411	29.0	1.8	( 25.6, 32.5)
ARIZONA	1323	25.0	1.7	( 21.7, 28.3)
ARKANSAS	2099	32.7	1.2	( 30.4, 35.0)
CALIFORNIA	3158	30.0	0.9	( 28.2, 31.8)
COLORADO	2170	25.9	1.3	( 23.4, 28.4)
CONNECTICUT	2745	28.0	1.2	( 25.7, 30.3)
DELAWARE	1941	31.6	1.4	( 28.8, 34.4)
DISTRICT OF CO	1037	22.1	1.5	( 19.2, 24.9)
FLORIDA	4063	33.0	0.9	( 31.3, 34.8)
GEORGIA	1715	29.2	1.3	( 26.8, 31.7)
HAWAII	1466	27.6	1.6	( 24.4, 30.7)
IDAHO	3459	30.1	1.0	( 28.2, 31.9)
ILLINOIS	1048	31.8	1.6	( 28.6, 35.0)
INDIANA	1819	31.6	1.9	( 27.9, 35.2)
IOWA	2630	31.7	1.1	( 29.6, 33.7)
KANSAS	2786	27.1	0.9	( 25.3, 28.9)
KENTUCKY	5331	32.5	0.9	( 30.8, 34.3)
LOUISIANA	1166	26.5	1.4	( 23.7, 29.2)
MAINE	1295	31.2	1.4	( 28.5, 33.9)
MARYLAND	3129	30.6	1.1	( 28.6, 32.7)
MASSACHUSETTS	3925	29.3	0.9	( 27.5, 31.2)
MICHIGAN	1917	32.9	1.2	( 30.5, 35.2)
MINNESOTA	3401	31.3	0.9	( 29.7, 33.0)
MISSISSIPPI	1495	30.1	1.3	( 27.5, 32.7)
MISSOURI	2937	30.2	1.1	( 28.0, 32.4)
MONTANA	1269	30.5	1.4	( 27.6, 33.3)
NEBRASKA	1992	27.4	1.1	( 25.3, 29.5)
NEVADA	1589	35.1	1.8	( 31.5, 38.6)
NEW HAMPSHIRE	971	33.2	1.7	( 29.9, 36.4)
NEW JERSEY	2226	27.1	1.1	( 24.9, 29.3)
NEW MEXICO	2365	27.3	1.0	( 25.2, 29.3)
NEW YORK	2010	28.6	1.2	( 26.4, 30.9)
NORTH CAROLINA	1869	31.2	1.3	( 28.7, 33.7)
NORTH DAKOTA	1407	30.6	1.3	( 27.9, 33.2)
OHIO	1223	32.5	1.6	( 29.4, 35.5)
OKLAHOMA	2205	21.2	0.9	( 19.3, 23.0)
OREGON	1320	29.0	1.4	( 26.3, 31.6)
PENNSYLVANIA	2648	28.2	1.0	( 26.3, 30.2)
RHODE ISLAND	3110	28.6	0.9	( 26.8, 30.4)
SOUTH CAROLINA	2620	27.3	1.0	( 25.3, 29.2)
SOUTH DAKOTA	2886	29.4	0.9	( 27.6, 31.2)
TENNESSEE	2252	29.1	1.1	( 27.0, 31.3)
TEXAS	3637	30.9	1.0	( 29.0, 32.7)
UTAH	2164	28.4	1.3	( 25.8, 31.0)
VERMONT	2408	26.7	1.0	( 24.8, 28.7)
VIRGINIA	2692	31.1	1.4	( 28.4, 33.7)
WASHINGTON	2654	28.3	1.0	( 26.3, 30.3)
WEST VIRGINIA	1886	37.1	1.3	( 34.7, 39.6)
WISCONSIN	1626	31.6	1.3	( 29.0, 34.1)
WYOMING	1754	30.9	1.3	( 28.4, 33.3)
PUERTO RICO	2412	30.0	1.1	( 28.0, 32.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 30.0 RANGE = 21.2-37.1

\* AMONG RESPONDENTS HAVING THEIR CHOLESTEROL CHECKED

June 20, 2000

TABLE 12.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD CHOLESTEROL IS HIGH\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	622	35.7	2.1	( 31.5, 39.9)	*	906	30.9	1.7	( 27.6, 34.1)	*
ALASKA	664	31.1	2.5	( 26.1, 36.0)	*	747	26.9	2.5	( 22.1, 31.7)	*
ARIZONA	551	22.8	2.3	( 18.2, 27.3)	*	772	27.1	2.3	( 22.6, 31.7)	*
ARKANSAS	771	32.3	1.9	( 28.7, 36.0)	*	1328	33.1	1.5	( 30.2, 35.9)	*
CALIFORNIA	1264	31.0	1.5	( 28.1, 33.8)	*	1894	29.2	1.2	( 26.9, 31.5)	*
COLORADO	944	23.1	1.9	( 19.5, 26.7)	*	1226	28.5	1.8	( 25.0, 32.0)	*
CONNECTICUT	1039	29.2	1.8	( 25.6, 32.8)	*	1706	26.9	1.5	( 24.0, 29.8)	*
DELAWARE	768	31.7	2.2	( 27.3, 36.1)	*	1173	31.5	1.8	( 27.9, 35.0)	*
DISTRICT OF CO	373	19.5	2.4	( 14.9, 24.2)	*	664	24.2	1.8	( 20.7, 27.7)	*
FLORIDA	1552	30.2	1.4	( 27.4, 32.9)	*	2511	35.4	1.1	( 33.2, 37.7)	*
GEORGIA	651	28.6	2.0	( 24.7, 32.5)	*	1064	29.8	1.6	( 26.7, 33.0)	*
HAWAII	671	29.8	2.3	( 25.2, 34.3)	*	795	25.4	2.2	( 21.0, 29.8)	*
IDAHO	1419	32.6	1.5	( 29.6, 35.5)	*	2040	27.8	1.2	( 25.5, 30.2)	*
ILLINOIS	401	31.2	2.6	( 26.2, 36.3)	*	647	32.4	2.0	( 28.4, 36.4)	*
INDIANA	741	30.5	2.8	( 25.0, 36.0)	*	1078	32.5	2.4	( 27.8, 37.3)	*
IOWA	1013	32.3	1.7	( 29.0, 35.7)	*	1617	31.1	1.3	( 28.5, 33.8)	*
KANSAS	1100	27.0	1.5	( 24.1, 29.8)	*	1686	27.2	1.2	( 24.9, 29.5)	*
KENTUCKY	1841	29.1	1.4	( 26.4, 31.8)	*	3490	35.4	1.1	( 33.3, 37.5)	*
LOUISIANA	450	25.5	2.2	( 21.1, 29.8)	*	716	27.3	1.8	( 23.8, 30.9)	*
MAINE	552	31.9	2.2	( 27.6, 36.1)	*	743	30.6	1.8	( 27.0, 34.2)	*
MARYLAND	1219	32.2	1.7	( 28.9, 35.5)	*	1910	29.3	1.3	( 26.8, 31.8)	*
MASSACHUSETTS	1512	29.6	1.5	( 26.7, 32.5)	*	2413	29.1	1.2	( 26.8, 31.4)	*
MICHIGAN	762	31.8	1.9	( 28.1, 35.4)	*	1155	33.8	1.6	( 30.8, 36.8)	*
MINNESOTA	1447	31.9	1.3	( 29.3, 34.4)	*	1954	30.9	1.1	( 28.7, 33.1)	*
MISSISSIPPI	571	28.9	2.1	( 24.7, 33.1)	*	924	31.1	1.6	( 27.9, 34.3)	*
MISSOURI	1150	30.2	1.8	( 26.6, 33.8)	*	1787	30.2	1.4	( 27.4, 33.0)	*
MONTANA	522	31.8	2.3	( 27.4, 36.2)	*	747	29.2	1.9	( 25.6, 32.9)	*
NEBRASKA	749	26.6	1.7	( 23.2, 30.0)	*	1243	28.1	1.4	( 25.4, 30.8)	*
NEVADA	786	32.2	2.3	( 27.7, 36.7)	*	803	37.8	2.7	( 32.5, 43.1)	*
NEW HAMPSHIRE	380	33.0	2.6	( 27.9, 38.0)	*	591	33.3	2.2	( 29.1, 37.6)	*
NEW JERSEY	872	27.4	1.8	( 23.9, 30.9)	*	1354	26.9	1.4	( 24.1, 29.6)	*
NEW MEXICO	1007	28.2	1.6	( 25.1, 31.2)	*	1358	26.4	1.4	( 23.8, 29.1)	*
NEW YORK	797	29.3	1.8	( 25.7, 32.9)	*	1213	28.0	1.5	( 25.2, 30.9)	*
NORTH CAROLINA	703	31.7	2.1	( 27.6, 35.7)	*	1166	30.8	1.5	( 27.9, 33.8)	*
NORTH DAKOTA	585	27.9	2.0	( 24.0, 31.7)	*	822	33.1	1.8	( 29.6, 36.5)	*
OHIO	468	32.7	2.5	( 27.8, 37.6)	*	755	32.2	2.0	( 28.3, 36.1)	*
OKLAHOMA	819	19.9	1.5	( 16.9, 22.8)	*	1386	22.3	1.2	( 20.0, 24.7)	*
OREGON	514	27.3	2.1	( 23.1, 31.5)	*	806	30.3	1.8	( 26.9, 33.7)	*
PENNSYLVANIA	1052	27.9	1.5	( 24.9, 30.8)	*	1596	28.6	1.3	( 26.0, 31.1)	*
RHODE ISLAND	1178	28.8	1.5	( 25.9, 31.7)	*	1932	28.4	1.2	( 26.0, 30.7)	*
SOUTH CAROLINA	1031	25.8	1.5	( 22.9, 28.7)	*	1589	28.5	1.3	( 26.0, 31.0)	*
SOUTH DAKOTA	1134	28.5	1.4	( 25.7, 31.3)	*	1752	30.2	1.2	( 27.9, 32.5)	*
TENNESSEE	734	27.8	1.9	( 24.2, 31.5)	*	1518	30.3	1.3	( 27.8, 32.8)	*
TEXAS	1401	29.5	1.4	( 26.8, 32.3)	*	2236	32.0	1.3	( 29.5, 34.5)	*
UTAH	914	29.2	2.1	( 25.1, 33.3)	*	1250	27.6	1.7	( 24.2, 31.0)	*
VERMONT	919	27.4	1.6	( 24.3, 30.5)	*	1489	26.1	1.3	( 23.7, 28.6)	*
VIRGINIA	1055	31.1	2.3	( 26.6, 35.5)	*	1637	31.1	1.6	( 28.0, 34.2)	*
WASHINGTON	1133	28.9	1.6	( 25.8, 32.0)	*	1521	27.7	1.3	( 25.2, 30.2)	*
WEST VIRGINIA	746	34.7	2.0	( 30.8, 38.6)	*	1140	39.1	1.6	( 36.0, 42.2)	*
WISCONSIN	650	32.0	2.0	( 28.0, 35.9)	*	976	31.2	1.7	( 27.9, 34.5)	*
WYOMING	772	32.1	2.0	( 28.3, 36.0)	*	982	29.7	1.6	( 26.6, 32.7)	*
PUERTO RICO	847	31.5	1.8	( 28.0, 35.0)	*	1565	28.8	1.3	( 26.4, 31.3)	*

SUMMARY STATISTICS:      NUMBER OF PARTICIPANTS = 52      NUMBER OF PARTICIPANTS = 52  
 MEDIAN                      = 29.7      MEDIAN                      = 29.8  
 RANGE                        = 19.5-35.7      RANGE                        = 22.3-39.1

\* AMONG RESPONDENTS HAVING THEIR CHOLESTEROL CHECKED

June 20, 2000

TABLE 12.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 EVER TOLD CHOLESTEROL IS HIGH\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	28.8	33.2	4.4
ALASKA	25.5	29.0	3.5
ARIZONA	31.9	25.0	-6.9
ARKANSAS	30.1	32.7	2.6
CALIFORNIA	30.4	30.0	-0.4
COLORADO	27.9	25.9	-2.0
CONNECTICUT	24.3	28.0	3.6
DELAWARE	28.9	31.6	2.7
DISTRICT OF CO	18.2	22.1	3.9
FLORIDA	31.9	33.0	1.1
GEORGIA	24.1	29.2	5.1
HAWAII	31.5	27.6	-4.0
IDAHO	29.8	30.1	0.3
ILLINOIS	34.3	31.8	-2.4
INDIANA	29.0	31.6	2.5
IOWA	28.1	31.7	3.6
KANSAS	28.1	27.1	-0.9
KENTUCKY	30.1	32.5	2.4
LOUISIANA	26.6	26.5	-0.1
MAINE	32.2	31.2	-1.0
MARYLAND	28.6	30.6	2.0
MASSACHUSETTS	25.0	29.3	4.4
MICHIGAN	31.2	32.9	1.7
MINNESOTA	31.2	31.3	0.1
MISSISSIPPI	28.8	30.1	1.3
MISSOURI	30.5	30.2	-0.3
MONTANA	31.0	30.5	-0.5
NEBRASKA	30.0	27.4	-2.6
NEVADA	29.6	35.1	5.5
NEW HAMPSHIRE	30.9	33.2	2.2
NEW JERSEY	27.9	27.1	-0.8
NEW MEXICO	28.0	27.3	-0.7
NEW YORK	28.0	28.6	0.6
NORTH CAROLINA	26.5	31.2	4.8
NORTH DAKOTA	29.8	30.6	0.7
OHIO	27.7	32.5	4.8
OKLAHOMA	22.0	21.2	-0.8
OREGON	32.0	29.0	-3.0
PENNSYLVANIA	25.9	28.2	2.4
RHODE ISLAND	28.1	28.6	0.5
SOUTH CAROLINA	24.5	27.3	2.8
SOUTH DAKOTA	25.7	29.4	3.7
TENNESSEE	29.5	29.1	-0.3
TEXAS	28.7	30.9	2.2
UTAH	26.4	28.4	2.0
VERMONT	25.9	26.7	0.8
VIRGINIA	29.5	31.1	1.6
WASHINGTON	25.6	28.3	2.7
WEST VIRGINIA	32.2	37.1	4.9
WISCONSIN	26.9	31.6	4.6
WYOMING	29.9	30.9	1.0
PUERTO RICO	24.4	30.0	5.6

\* AMONG RESPONDENTS HAVING THEIR CHOLESTEROL CHECKED

TABLE 13.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED\*  
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2089	47.7	1.2	( 45.3, 50.2)
ALASKA	2045	53.0	1.7	( 49.7, 56.2)
ARIZONA	1742	37.9	1.7	( 34.5, 41.3)
ARKANSAS	2989	50.3	1.0	( 48.3, 52.3)
CALIFORNIA	4303	45.1	0.9	( 43.4, 46.8)
COLORADO	3046	48.3	1.3	( 45.7, 50.8)
CONNECTICUT	3507	48.2	1.2	( 45.9, 50.5)
DELAWARE	2555	49.4	1.4	( 46.7, 52.2)
DISTRICT OF CO	1260	36.7	1.5	( 33.7, 39.7)
FLORIDA	5164	47.3	0.9	( 45.7, 49.0)
GEORGIA	2261	46.3	1.2	( 43.9, 48.7)
HAWAII	2152	41.5	1.4	( 38.7, 44.2)
IDAHO	4942	45.8	0.9	( 44.1, 47.5)
ILLINOIS	2906	48.1	1.1	( 45.9, 50.2)
INDIANA	2429	49.0	1.7	( 45.7, 52.3)
IOWA	3613	48.1	1.0	( 46.1, 50.0)
KANSAS	3864	43.6	0.9	( 41.8, 45.3)
KENTUCKY	7524	52.9	0.8	( 51.3, 54.5)
LOUISIANA	1665	44.2	1.4	( 41.4, 46.9)
MAINE	1671	52.8	1.4	( 50.0, 55.6)
MARYLAND	3943	44.5	1.0	( 42.5, 46.5)
MASSACHUSETTS	5003	45.1	0.9	( 43.3, 46.9)
MICHIGAN	2510	48.7	1.1	( 46.5, 51.0)
MINNESOTA	5332	39.5	0.8	( 38.0, 41.0)
MISSISSIPPI	2174	43.2	1.2	( 40.8, 45.6)
MISSOURI	4269	51.8	1.1	( 49.7, 53.9)
MONTANA	1795	47.9	1.3	( 45.4, 50.5)
NEBRASKA	2816	45.3	1.1	( 43.2, 47.4)
NEVADA	2187	60.6	1.6	( 57.4, 63.7)
NEW HAMPSHIRE	1245	51.3	1.6	( 48.1, 54.5)
NEW JERSEY	2806	41.9	1.1	( 39.7, 44.1)
NEW MEXICO	3484	47.1	1.0	( 45.2, 48.9)
NEW YORK	2639	44.9	1.1	( 42.7, 47.0)
NORTH CAROLINA	2435	48.4	1.2	( 46.0, 50.7)
NORTH DAKOTA	1977	46.9	1.3	( 44.5, 49.4)
OHIO	1635	51.6	1.5	( 48.6, 54.5)
OKLAHOMA	2964	43.9	1.1	( 41.8, 46.0)
OREGON	1806	49.6	1.3	( 47.0, 52.1)
PENNSYLVANIA	3553	46.4	1.0	( 44.6, 48.3)
RHODE ISLAND	3986	47.8	0.9	( 46.0, 49.6)
SOUTH CAROLINA	3452	44.0	1.0	( 42.1, 45.9)
SOUTH DAKOTA	4146	47.6	0.9	( 45.9, 49.3)
TENNESSEE	3006	45.6	1.1	( 43.5, 47.6)
TEXAS	4978	45.7	0.9	( 43.9, 47.5)
UTAH	3195	31.2	1.1	( 29.1, 33.4)
VERMONT	3150	48.4	1.0	( 46.4, 50.4)
VIRGINIA	3486	46.7	1.3	( 44.3, 49.2)
WASHINGTON	3595	49.2	1.1	( 47.2, 51.3)
WEST VIRGINIA	2588	51.0	1.1	( 48.8, 53.1)
WISCONSIN	2166	52.3	1.2	( 49.9, 54.7)
WYOMING	2396	48.8	1.2	( 46.5, 51.1)
PUERTO RICO	3044	29.1	1.0	( 27.1, 31.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 47.5 RANGE = 29.1-60.6

\* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

June 20, 2000

TABLE 13.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER SMOKED\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	853	58.2	2.0	( 54.4, 62.0)	*	1236	38.4	1.6	( 35.4, 41.5)	*
ALASKA	999	52.7	2.3	( 48.2, 57.2)	*	1046	53.3	2.4	( 48.6, 57.9)	*
ARIZONA	713	44.7	2.6	( 39.5, 49.9)	*	1029	31.4	2.2	( 27.2, 35.7)	*
ARKANSAS	1160	58.8	1.6	( 55.7, 61.9)	*	1829	42.7	1.3	( 40.2, 45.3)	*
CALIFORNIA	1836	52.0	1.3	( 49.5, 54.6)	*	2467	38.3	1.1	( 36.1, 40.4)	*
COLORADO	1368	52.5	2.0	( 48.7, 56.3)	*	1678	44.3	1.7	( 40.9, 47.6)	*
CONNECTICUT	1367	53.5	1.8	( 50.0, 56.9)	*	2140	43.4	1.6	( 40.3, 46.4)	*
DELAWARE	1026	53.4	2.1	( 49.2, 57.5)	*	1529	45.9	1.8	( 42.3, 49.5)	*
DISTRICT OF CO	458	39.0	2.5	( 34.1, 43.9)	*	802	34.7	1.9	( 31.0, 38.5)	*
FLORIDA	2048	53.5	1.4	( 50.9, 56.2)	*	3116	41.7	1.1	( 39.6, 43.8)	*
GEORGIA	883	55.1	2.0	( 51.2, 59.0)	*	1378	38.2	1.5	( 35.3, 41.1)	*
HAWAII	991	46.6	2.1	( 42.5, 50.7)	*	1161	36.3	1.8	( 32.7, 39.9)	*
IDAHO	2088	51.4	1.4	( 48.8, 54.1)	*	2854	40.4	1.1	( 38.3, 42.5)	*
ILLINOIS	1110	53.4	1.7	( 50.0, 56.8)	*	1796	43.2	1.3	( 40.6, 45.7)	*
INDIANA	1013	57.9	2.6	( 52.8, 63.0)	*	1416	41.0	2.2	( 36.7, 45.2)	*
IOWA	1422	57.0	1.6	( 53.9, 60.1)	*	2191	40.0	1.2	( 37.6, 42.4)	*
KANSAS	1557	52.4	1.4	( 49.6, 55.2)	*	2307	35.3	1.1	( 33.2, 37.5)	*
KENTUCKY	2745	60.7	1.4	( 58.0, 63.4)	*	4779	45.8	1.0	( 43.9, 47.8)	*
LOUISIANA	661	50.4	2.2	( 46.2, 54.7)	*	1004	38.6	1.8	( 35.1, 42.0)	*
MAINE	736	60.3	2.1	( 56.1, 64.5)	*	935	45.9	1.8	( 42.4, 49.4)	*
MARYLAND	1584	48.9	1.6	( 45.8, 52.0)	*	2359	40.5	1.3	( 38.0, 42.9)	*
MASSACHUSETTS	1999	46.3	1.5	( 43.5, 49.2)	*	3004	44.0	1.1	( 41.8, 46.2)	*
MICHIGAN	1034	53.5	1.8	( 50.0, 56.9)	*	1476	44.4	1.4	( 41.7, 47.2)	*
MINNESOTA	2437	42.6	1.1	( 40.4, 44.7)	*	2895	36.6	1.0	( 34.7, 38.6)	*
MISSISSIPPI	849	54.4	2.1	( 50.4, 58.4)	*	1325	33.4	1.4	( 30.6, 36.2)	*
MISSOURI	1719	59.8	1.6	( 56.6, 63.0)	*	2550	44.6	1.4	( 41.9, 47.2)	*
MONTANA	767	50.2	2.0	( 46.2, 54.1)	*	1028	45.9	1.7	( 42.5, 49.2)	*
NEBRASKA	1116	53.8	1.6	( 50.5, 57.0)	*	1700	37.6	1.3	( 35.0, 40.1)	*
NEVADA	1101	63.8	2.3	( 59.3, 68.2)	*	1086	57.3	2.3	( 52.9, 61.7)	*
NEW HAMPSHIRE	503	54.0	2.5	( 49.1, 58.9)	*	742	48.8	2.1	( 44.6, 52.9)	*
NEW JERSEY	1119	45.6	1.8	( 42.1, 49.1)	*	1687	38.6	1.4	( 35.8, 41.4)	*
NEW MEXICO	1531	52.3	1.4	( 49.5, 55.1)	*	1953	42.2	1.3	( 39.7, 44.6)	*
NEW YORK	1064	47.4	1.7	( 44.0, 50.8)	*	1575	42.6	1.4	( 39.8, 45.3)	*
NORTH CAROLINA	961	56.8	1.9	( 53.1, 60.4)	*	1474	40.7	1.5	( 37.8, 43.7)	*
NORTH DAKOTA	859	53.5	1.9	( 49.9, 57.2)	*	1118	40.5	1.6	( 37.2, 43.7)	*
OHIO	633	57.5	2.4	( 52.9, 62.1)	*	1002	46.3	1.9	( 42.6, 50.0)	*
OKLAHOMA	1154	49.4	1.7	( 46.1, 52.8)	*	1810	38.8	1.3	( 36.3, 41.4)	*
OREGON	768	52.5	2.0	( 48.6, 56.4)	*	1038	46.8	1.7	( 43.4, 50.1)	*
PENNSYLVANIA	1417	52.4	1.5	( 49.5, 55.3)	*	2136	41.1	1.2	( 38.8, 43.5)	*
RHODE ISLAND	1553	50.4	1.5	( 47.5, 53.2)	*	2433	45.6	1.2	( 43.3, 47.9)	*
SOUTH CAROLINA	1392	55.2	1.6	( 52.2, 58.3)	*	2060	33.8	1.2	( 31.5, 36.1)	*
SOUTH DAKOTA	1685	53.8	1.4	( 51.1, 56.5)	*	2461	41.7	1.1	( 39.6, 43.9)	*
TENNESSEE	1032	51.4	1.7	( 48.0, 54.8)	*	1974	40.3	1.2	( 38.0, 42.7)	*
TEXAS	1997	54.8	1.4	( 52.1, 57.5)	*	2981	37.2	1.1	( 35.0, 39.4)	*
UTAH	1372	39.0	1.8	( 35.5, 42.5)	*	1823	23.9	1.3	( 21.3, 26.4)	*
VERMONT	1269	52.6	1.6	( 49.5, 55.7)	*	1881	44.5	1.3	( 42.0, 47.1)	*
VIRGINIA	1400	52.4	2.0	( 48.5, 56.3)	*	2086	41.4	1.6	( 38.4, 44.5)	*
WASHINGTON	1593	55.5	1.6	( 52.3, 58.6)	*	2002	43.2	1.3	( 40.6, 45.8)	*
WEST VIRGINIA	1065	61.0	1.7	( 57.7, 64.3)	*	1523	42.0	1.4	( 39.2, 44.8)	*
WISCONSIN	915	55.9	1.9	( 52.3, 59.5)	*	1251	48.9	1.7	( 45.7, 52.2)	*
WYOMING	1060	54.2	1.7	( 50.8, 57.6)	*	1336	43.5	1.6	( 40.5, 46.6)	*
PUERTO RICO	1118	40.9	1.7	( 37.6, 44.1)	*	1926	18.6	1.0	( 16.6, 20.5)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 53.4	MEDIAN	= 41.6
	RANGE	= 39.0-63.8	RANGE	= 18.6-57.3

\* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

June 20, 2000

TABLE 13.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 EVER SMOKED\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	46.2	47.7	1.6
ALASKA	51.6	53.0	1.4
ARIZONA	39.1	37.9	-1.2
ARKANSAS	50.2	50.3	0.1
CALIFORNIA	45.4	45.1	-0.3
COLORADO	47.1	48.3	1.2
CONNECTICUT	49.1	48.2	-0.9
DELAWARE	49.0	49.4	0.5
DISTRICT OF CO	38.7	36.7	-1.9
FLORIDA	48.1	47.3	-0.8
GEORGIA	45.6	46.3	0.7
HAWAII	45.6	41.5	-4.1
IDAHO	45.4	45.8	0.4
ILLINOIS	48.1	48.1	-0.1
INDIANA	52.0	49.0	-3.0
IOWA	47.1	48.1	1.0
KANSAS	42.9	43.6	0.6
KENTUCKY	53.7	52.9	-0.8
LOUISIANA	45.4	44.2	-1.2
MAINE	53.7	52.8	-0.9
MARYLAND	46.6	44.5	-2.1
MASSACHUSETTS	48.9	45.1	-3.8
MICHIGAN	52.8	48.7	-4.1
MINNESOTA	36.6	39.5	2.9
MISSISSIPPI	46.1	43.2	-2.8
MISSOURI	50.0	51.8	1.8
MONTANA	48.2	47.9	-0.2
NEBRASKA	46.5	45.3	-1.2
NEVADA	54.4	60.6	6.2
NEW HAMPSHIRE	51.8	51.3	-0.5
NEW JERSEY	45.5	41.9	-3.6
NEW MEXICO	46.9	47.1	0.2
NEW YORK	48.4	44.9	-3.5
NORTH CAROLINA	48.1	48.4	0.3
NORTH DAKOTA	48.2	46.9	-1.3
OHIO	49.7	51.6	1.9
OKLAHOMA	43.5	43.9	0.4
OREGON	48.4	49.6	1.1
PENNSYLVANIA	46.7	46.4	-0.3
RHODE ISLAND	51.6	47.8	-3.8
SOUTH CAROLINA	45.4	44.0	-1.4
SOUTH DAKOTA	47.6	47.6	0.0
TENNESSEE	47.6	45.6	-2.0
TEXAS	44.5	45.7	1.2
UTAH	32.9	31.2	-1.7
VERMONT	50.9	48.4	-2.5
VIRGINIA	46.1	46.7	0.6
WASHINGTON	48.9	49.2	0.3
WEST VIRGINIA	51.7	51.0	-0.7
WISCONSIN	52.8	52.3	-0.5
WYOMING	48.0	48.8	0.8
PUERTO RICO	32.1	29.1	-3.0

\* SMOKED AT LEAST 100 CIGARETTES IN LIFETIME

June 20, 2000

TABLE 14.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2088	23.5	1.1	( 21.4, 25.5)
ALASKA	2045	27.3	1.5	( 24.4, 30.2)
ARIZONA	1742	20.1	1.4	( 17.3, 22.9)
ARKANSAS	2989	27.2	0.9	( 25.4, 29.0)
CALIFORNIA	4303	18.7	0.7	( 17.4, 20.0)
COLORADO	3046	22.5	1.1	( 20.3, 24.6)
CONNECTICUT	3505	22.8	1.1	( 20.7, 25.0)
DELAWARE	2554	25.5	1.3	( 23.0, 27.9)
DISTRICT OF CO	1260	20.6	1.3	( 17.9, 23.2)
FLORIDA	5164	20.6	0.7	( 19.2, 21.9)
GEORGIA	2261	23.8	1.1	( 21.7, 25.8)
HAWAII	2151	18.5	1.1	( 16.3, 20.7)
IDAHO	4942	21.5	0.7	( 20.1, 22.9)
ILLINOIS	2906	24.2	0.9	( 22.4, 26.1)
INDIANA	2428	27.0	1.5	( 24.0, 30.0)
IOWA	3613	23.5	0.9	( 21.8, 25.1)
KANSAS	3864	21.0	0.8	( 19.5, 22.5)
KENTUCKY	7523	29.7	0.7	( 28.3, 31.2)
LOUISIANA	1665	23.5	1.2	( 21.1, 25.9)
MAINE	1671	23.3	1.3	( 20.7, 25.8)
MARYLAND	3943	20.3	0.8	( 18.7, 21.9)
MASSACHUSETTS	4999	19.3	0.7	( 17.9, 20.7)
MICHIGAN	2510	25.1	1.0	( 23.2, 27.0)
MINNESOTA	5330	19.5	0.6	( 18.3, 20.6)
MISSISSIPPI	2174	22.9	1.0	( 20.9, 25.0)
MISSOURI	4269	27.1	1.0	( 25.2, 28.9)
MONTANA	1795	20.2	1.1	( 18.1, 22.3)
NEBRASKA	2815	23.2	0.9	( 21.4, 25.1)
NEVADA	2186	31.5	1.5	( 28.5, 34.5)
NEW HAMPSHIRE	1244	22.3	1.4	( 19.6, 25.0)
NEW JERSEY	2806	20.6	1.0	( 18.7, 22.5)
NEW MEXICO	3484	22.5	0.8	( 20.9, 24.0)
NEW YORK	2639	21.8	0.9	( 19.9, 23.6)
NORTH CAROLINA	2434	25.1	1.1	( 23.0, 27.2)
NORTH DAKOTA	1976	22.1	1.0	( 20.1, 24.1)
OHIO	1635	27.6	1.3	( 25.0, 30.2)
OKLAHOMA	2963	25.2	1.0	( 23.3, 27.0)
OREGON	1805	21.4	1.1	( 19.3, 23.5)
PENNSYLVANIA	3552	23.1	0.8	( 21.5, 24.7)
RHODE ISLAND	3986	22.3	0.8	( 20.8, 23.7)
SOUTH CAROLINA	3451	23.6	0.9	( 21.9, 25.2)
SOUTH DAKOTA	4146	22.5	0.8	( 21.0, 24.0)
TENNESSEE	3005	24.8	0.9	( 23.0, 26.6)
TEXAS	4978	22.4	0.8	( 20.8, 24.0)
UTAH	3194	14.0	0.8	( 12.4, 15.5)
VERMONT	3150	21.7	0.9	( 20.0, 23.4)
VIRGINIA	3485	21.4	0.9	( 19.5, 23.2)
WASHINGTON	3594	22.4	0.9	( 20.7, 24.1)
WEST VIRGINIA	2588	27.1	1.0	( 25.1, 29.1)
WISCONSIN	2166	23.7	1.0	( 21.7, 25.6)
WYOMING	2396	23.9	1.0	( 21.9, 25.9)
PUERTO RICO	3044	13.7	0.8	( 12.2, 15.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 22.7 RANGE = 13.7-31.5 YEAR 2000 OBJ. 3.4  
 \* SMOKES EVERYDAY OR SOME DAYS June 20, 2000

TABLE 14.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKERS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	852	26.2	1.7	( 22.9, 29.4)	*	1236	21.1	1.3	( 18.5, 23.6)	*
ALASKA	999	25.3	1.8	( 21.8, 28.8)	*	1046	29.5	2.4	( 24.8, 34.1)	*
ARIZONA	713	23.7	2.3	( 19.2, 28.1)	*	1029	16.7	1.8	( 13.2, 20.1)	*
ARKANSAS	1160	29.7	1.5	( 26.7, 32.6)	*	1829	25.0	1.1	( 22.8, 27.3)	*
CALIFORNIA	1836	22.0	1.1	( 19.8, 24.1)	*	2467	15.5	0.8	( 13.9, 17.1)	*
COLORADO	1368	22.7	1.6	( 19.6, 25.9)	*	1678	22.2	1.5	( 19.3, 25.0)	*
CONNECTICUT	1366	25.3	1.6	( 22.2, 28.4)	*	2139	20.6	1.5	( 17.7, 23.5)	*
DELAWARE	1025	27.6	1.9	( 23.9, 31.4)	*	1529	23.5	1.7	( 20.1, 26.8)	*
DISTRICT OF CO	458	21.4	2.1	( 17.3, 25.6)	*	802	19.8	1.7	( 16.5, 23.1)	*
FLORIDA	2048	22.2	1.1	( 20.0, 24.3)	*	3116	19.1	0.8	( 17.5, 20.7)	*
GEORGIA	883	28.3	1.8	( 24.8, 31.8)	*	1378	19.6	1.2	( 17.2, 21.9)	*
HAWAII	990	20.1	1.7	( 16.7, 23.4)	*	1161	16.9	1.5	( 14.0, 19.8)	*
IDAHO	2088	22.5	1.2	( 20.1, 24.9)	*	2854	20.6	0.9	( 18.8, 22.3)	*
ILLINOIS	1110	27.0	1.5	( 24.0, 30.0)	*	1796	21.7	1.1	( 19.6, 23.9)	*
INDIANA	1013	31.0	2.4	( 26.2, 35.7)	*	1415	23.4	1.9	( 19.8, 27.0)	*
IOWA	1422	26.6	1.4	( 23.8, 29.3)	*	2191	20.7	1.0	( 18.7, 22.6)	*
KANSAS	1557	24.3	1.3	( 21.8, 26.8)	*	2307	18.0	0.9	( 16.3, 19.7)	*
KENTUCKY	2745	33.9	1.3	( 31.5, 36.4)	*	4778	25.9	0.9	( 24.2, 27.6)	*
LOUISIANA	661	26.8	2.0	( 23.0, 30.6)	*	1004	20.6	1.5	( 17.6, 23.5)	*
MAINE	736	27.7	2.2	( 23.5, 31.9)	*	935	19.2	1.4	( 16.4, 21.9)	*
MARYLAND	1584	22.1	1.3	( 19.5, 24.7)	*	2359	18.6	1.0	( 16.7, 20.6)	*
MASSACHUSETTS	1996	19.5	1.1	( 17.3, 21.7)	*	3003	19.1	0.9	( 17.3, 20.8)	*
MICHIGAN	1034	26.6	1.5	( 23.6, 29.6)	*	1476	23.8	1.2	( 21.4, 26.2)	*
MINNESOTA	2436	21.7	0.9	( 19.9, 23.5)	*	2894	17.3	0.8	( 15.9, 18.8)	*
MISSISSIPPI	849	27.4	1.8	( 23.9, 30.9)	*	1325	19.0	1.2	( 16.7, 21.3)	*
MISSOURI	1719	30.6	1.5	( 27.6, 33.5)	*	2550	23.9	1.2	( 21.6, 26.2)	*
MONTANA	767	18.5	1.6	( 15.4, 21.6)	*	1028	21.9	1.5	( 19.0, 24.7)	*
NEBRASKA	1116	27.4	1.5	( 24.5, 30.4)	*	1699	19.4	1.1	( 17.1, 21.6)	*
NEVADA	1100	32.8	2.2	( 28.5, 37.0)	*	1086	30.3	2.1	( 26.1, 34.4)	*
NEW HAMPSHIRE	503	21.6	2.1	( 17.4, 25.7)	*	741	23.0	1.8	( 19.5, 26.5)	*
NEW JERSEY	1119	22.0	1.6	( 19.0, 25.1)	*	1687	19.4	1.2	( 17.1, 21.6)	*
NEW MEXICO	1531	24.1	1.2	( 21.7, 26.5)	*	1953	20.9	1.0	( 18.9, 22.9)	*
NEW YORK	1064	22.8	1.5	( 19.9, 25.7)	*	1575	20.8	1.2	( 18.5, 23.2)	*
NORTH CAROLINA	961	27.6	1.7	( 24.3, 30.9)	*	1473	22.9	1.4	( 20.2, 25.6)	*
NORTH DAKOTA	858	23.4	1.6	( 20.3, 26.4)	*	1118	20.9	1.4	( 18.2, 23.6)	*
OHIO	633	29.3	2.2	( 25.0, 33.5)	*	1002	26.1	1.7	( 22.9, 29.3)	*
OKLAHOMA	1154	26.6	1.5	( 23.6, 29.6)	*	1809	23.8	1.2	( 21.5, 26.1)	*
OREGON	767	22.9	1.7	( 19.6, 26.1)	*	1038	20.1	1.4	( 17.4, 22.8)	*
PENNSYLVANIA	1416	24.2	1.3	( 21.7, 26.7)	*	2136	22.1	1.0	( 20.1, 24.1)	*
RHODE ISLAND	1553	23.2	1.2	( 20.8, 25.5)	*	2433	21.5	1.0	( 19.6, 23.4)	*
SOUTH CAROLINA	1392	28.4	1.4	( 25.6, 31.2)	*	2059	19.2	0.9	( 17.3, 21.0)	*
SOUTH DAKOTA	1685	23.1	1.2	( 20.7, 25.4)	*	2461	21.9	0.9	( 20.1, 23.7)	*
TENNESSEE	1032	25.6	1.5	( 22.7, 28.6)	*	1973	24.1	1.0	( 22.1, 26.2)	*
TEXAS	1997	27.4	1.4	( 24.7, 30.1)	*	2981	17.8	0.8	( 16.2, 19.3)	*
UTAH	1371	16.7	1.3	( 14.1, 19.3)	*	1823	11.3	0.9	( 9.5, 13.2)	*
VERMONT	1269	22.8	1.4	( 20.2, 25.5)	*	1881	20.6	1.1	( 18.4, 22.8)	*
VIRGINIA	1399	21.6	1.4	( 18.8, 24.3)	*	2086	21.2	1.2	( 18.8, 23.6)	*
WASHINGTON	1592	24.0	1.3	( 21.4, 26.6)	*	2002	20.9	1.1	( 18.7, 23.0)	*
WEST VIRGINIA	1065	30.2	1.6	( 27.0, 33.4)	*	1523	24.4	1.3	( 21.9, 26.8)	*
WISCONSIN	915	22.9	1.5	( 19.9, 26.0)	*	1251	24.4	1.4	( 21.7, 27.0)	*
WYOMING	1060	25.9	1.5	( 22.9, 28.8)	*	1336	22.0	1.3	( 19.4, 24.6)	*
PUERTO RICO	1118	18.9	1.4	( 16.2, 21.7)	*	1926	9.1	0.8	( 7.6, 10.6)	*

SUMMARY STATISTICS:      NUMBER OF PARTICIPANTS = 52      NUMBER OF PARTICIPANTS = 52  
 MEDIAN                      = 24.2      MEDIAN                      = 20.9  
 RANGE                        = 16.7-33.9      RANGE                        = 9.1-30.3

\* SMOKES EVERYDAY OR SOME DAYS

June 20, 2000

TABLE 14.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 CURRENT SMOKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	24.6	23.5	-1.2
ALASKA	26.1	27.3	1.2
ARIZONA	21.9	20.1	-1.8
ARKANSAS	25.9	27.2	1.3
CALIFORNIA	19.2	18.7	-0.5
COLORADO	22.8	22.5	-0.4
CONNECTICUT	20.9	22.8	1.9
DELAWARE	24.5	25.5	1.0
DISTRICT OF CO	21.6	20.6	-1.1
FLORIDA	22.0	20.6	-1.4
GEORGIA	23.6	23.8	0.1
HAWAII	19.5	18.5	-1.0
IDAHO	20.3	21.5	1.2
ILLINOIS	23.1	24.2	1.1
INDIANA	26.0	27.0	1.0
IOWA	23.4	23.5	0.1
KANSAS	21.1	21.0	-0.1
KENTUCKY	30.8	29.7	-1.1
LOUISIANA	25.5	23.5	-2.0
MAINE	22.4	23.3	0.9
MARYLAND	22.4	20.3	-2.1
MASSACHUSETTS	20.9	19.3	-1.6
MICHIGAN	27.4	25.1	-2.3
MINNESOTA	18.0	19.5	1.5
MISSISSIPPI	24.1	22.9	-1.2
MISSOURI	26.4	27.1	0.7
MONTANA	21.5	20.2	-1.2
NEBRASKA	22.0	23.2	1.2
NEVADA	30.4	31.5	1.2
NEW HAMPSHIRE	23.3	22.3	-1.0
NEW JERSEY	19.1	20.6	1.6
NEW MEXICO	22.6	22.5	-0.1
NEW YORK	24.1	21.8	-2.3
NORTH CAROLINA	24.6	25.1	0.5
NORTH DAKOTA	20.0	22.1	2.1
OHIO	26.1	27.6	1.5
OKLAHOMA	23.9	25.2	1.3
OREGON	21.1	21.4	0.4
PENNSYLVANIA	23.8	23.1	-0.7
RHODE ISLAND	22.6	22.3	-0.3
SOUTH CAROLINA	24.7	23.6	-1.1
SOUTH DAKOTA	27.2	22.5	-4.7
TENNESSEE	26.1	24.8	-1.2
TEXAS	21.9	22.4	0.5
UTAH	14.2	14.0	-0.2
VERMONT	22.3	21.7	-0.6
VIRGINIA	22.9	21.4	-1.5
WASHINGTON	21.4	22.4	1.0
WEST VIRGINIA	27.9	27.1	-0.8
WISCONSIN	23.4	23.7	0.3
WYOMING	22.8	23.9	1.1
PUERTO RICO	15.3	13.7	-1.6

\* SMOKES EVERYDAY OR SOME DAYS

TABLE 15.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT DRINKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2095	39.6	1.2	( 37.2, 42.0)
ALASKA	2046	56.9	1.7	( 53.5, 60.2)
ARIZONA	1741	34.5	1.7	( 31.3, 37.8)
ARKANSAS	2997	38.3	1.0	( 36.3, 40.3)
CALIFORNIA	4154	60.4	0.9	( 58.7, 62.1)
COLORADO	3051	64.2	1.2	( 61.7, 66.6)
CONNECTICUT	3511	59.4	1.2	( 57.1, 61.7)
DELAWARE	2559	60.0	1.3	( 57.5, 62.6)
DISTRICT OF CO	1264	50.3	1.6	( 47.2, 53.4)
FLORIDA	5165	55.1	0.8	( 53.4, 56.7)
GEORGIA	2270	47.7	1.2	( 45.3, 50.1)
HAWAII	2153	48.6	1.4	( 45.8, 51.4)
IDAHO	4953	49.0	0.9	( 47.2, 50.7)
ILLINOIS	1446	57.3	1.5	( 54.4, 60.2)
INDIANA	2438	54.3	1.7	( 50.9, 57.6)
IOWA	3613	55.0	1.0	( 53.1, 56.9)
KANSAS	3869	46.8	0.9	( 45.0, 48.6)
KENTUCKY	7534	34.9	0.8	( 33.3, 36.5)
LOUISIANA	1666	45.4	1.4	( 42.7, 48.2)
MAINE	1670	59.8	1.4	( 57.1, 62.5)
MARYLAND	3956	57.8	1.0	( 55.8, 59.7)
MASSACHUSETTS	5012	61.5	0.9	( 59.8, 63.3)
MICHIGAN	2517	59.2	1.1	( 57.0, 61.3)
MINNESOTA	5337	51.1	0.8	( 49.5, 52.6)
MISSISSIPPI	2175	38.6	1.2	( 36.2, 41.0)
MISSOURI	4270	53.4	1.0	( 51.3, 55.4)
MONTANA	1797	56.9	1.3	( 54.4, 59.5)
NEBRASKA	2813	52.2	1.1	( 50.1, 54.2)
NEVADA	2187	63.1	1.6	( 60.1, 66.1)
NEW HAMPSHIRE	1246	64.2	1.5	( 61.2, 67.2)
NEW JERSEY	2811	54.0	1.2	( 51.8, 56.3)
NEW MEXICO	3485	54.1	1.0	( 52.3, 56.0)
NEW YORK	2644	52.6	1.1	( 50.4, 54.8)
NORTH CAROLINA	2439	41.7	1.2	( 39.4, 44.0)
NORTH DAKOTA	1978	55.4	1.2	( 53.0, 57.9)
OHIO	1636	43.8	1.5	( 40.9, 46.7)
OKLAHOMA	2964	28.6	1.0	( 26.7, 30.5)
OREGON	1804	60.9	1.3	( 58.4, 63.4)
PENNSYLVANIA	3548	51.5	1.0	( 49.6, 53.4)
RHODE ISLAND	3990	58.3	0.9	( 56.5, 60.1)
SOUTH CAROLINA	3457	42.0	1.0	( 40.1, 43.9)
SOUTH DAKOTA	4144	58.6	0.9	( 56.9, 60.2)
TENNESSEE	3010	29.7	1.0	( 27.7, 31.7)
TEXAS	4986	54.8	0.9	( 53.0, 56.5)
UTAH	3194	29.7	1.1	( 27.7, 31.8)
VERMONT	3159	60.5	1.0	( 58.6, 62.5)
VIRGINIA	3498	54.3	1.2	( 51.9, 56.7)
WASHINGTON	3599	60.7	1.0	( 58.7, 62.6)
WEST VIRGINIA	2586	32.6	1.1	( 30.5, 34.7)
WISCONSIN	2174	71.1	1.1	( 68.9, 73.3)
WYOMING	2401	56.1	1.2	( 53.8, 58.4)
PUERTO RICO	3048	31.8	1.1	( 29.7, 33.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 54.2 RANGE = 28.6-71.1

\* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH

June 20, 2000

TABLE 15.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT DRINKERS\* -BY SEX  
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	854	49.8	2.0	( 46.0, 53.6)	*	1241	30.6	1.5	( 27.7, 33.4)	*
ALASKA	998	62.8	2.2	( 58.4, 67.2)	*	1048	50.4	2.6	( 45.4, 55.4)	*
ARIZONA	711	43.4	2.6	( 38.2, 48.6)	*	1030	26.1	2.0	( 22.2, 30.1)	*
ARKANSAS	1166	46.7	1.6	( 43.5, 49.9)	*	1831	30.8	1.2	( 28.4, 33.1)	*
CALIFORNIA	1774	68.4	1.3	( 66.0, 70.9)	*	2380	52.5	1.2	( 50.2, 54.8)	*
COLORADO	1372	72.3	1.7	( 68.9, 75.7)	*	1679	56.3	1.7	( 53.0, 59.6)	*
CONNECTICUT	1369	66.3	1.7	( 63.0, 69.5)	*	2142	53.2	1.6	( 50.1, 56.2)	*
DELAWARE	1026	70.4	1.9	( 66.8, 74.1)	*	1533	50.6	1.8	( 47.1, 54.1)	*
DISTRICT OF CO	459	53.5	2.6	( 48.4, 58.6)	*	805	47.6	2.0	( 43.8, 51.5)	*
FLORIDA	2049	62.6	1.3	( 60.0, 65.2)	*	3116	48.2	1.1	( 46.1, 50.3)	*
GEORGIA	884	59.0	2.0	( 55.1, 62.8)	*	1386	37.4	1.5	( 34.4, 40.3)	*
HAWAII	991	59.3	2.1	( 55.2, 63.4)	*	1162	37.8	1.9	( 34.1, 41.4)	*
IDAHO	2096	55.1	1.3	( 52.5, 57.7)	*	2857	43.1	1.1	( 40.9, 45.2)	*
ILLINOIS	573	67.4	2.3	( 63.0, 71.9)	*	873	47.3	1.9	( 43.7, 50.9)	*
INDIANA	1016	62.5	2.6	( 57.4, 67.5)	*	1422	46.8	2.2	( 42.5, 51.2)	*
IOWA	1423	65.2	1.5	( 62.2, 68.1)	*	2190	45.8	1.2	( 43.4, 48.2)	*
KANSAS	1561	56.8	1.4	( 54.0, 59.5)	*	2308	37.5	1.1	( 35.3, 39.7)	*
KENTUCKY	2747	43.5	1.3	( 40.9, 46.1)	*	4787	27.1	1.0	( 25.3, 29.0)	*
LOUISIANA	660	56.1	2.2	( 51.9, 60.4)	*	1006	35.9	1.7	( 32.5, 39.3)	*
MAINE	736	68.4	1.9	( 64.6, 72.2)	*	934	51.9	1.8	( 48.4, 55.5)	*
MARYLAND	1587	66.1	1.5	( 63.2, 69.0)	*	2369	50.2	1.3	( 47.6, 52.7)	*
MASSACHUSETTS	2001	66.4	1.4	( 63.6, 69.2)	*	3011	57.1	1.1	( 54.9, 59.3)	*
MICHIGAN	1039	68.0	1.6	( 64.8, 71.1)	*	1478	51.1	1.4	( 48.3, 53.9)	*
MINNESOTA	2436	55.5	1.1	( 53.4, 57.7)	*	2901	46.9	1.0	( 44.8, 48.9)	*
MISSISSIPPI	850	50.0	2.0	( 46.1, 54.0)	*	1325	28.6	1.4	( 25.9, 31.2)	*
MISSOURI	1718	62.3	1.6	( 59.3, 65.4)	*	2552	45.3	1.4	( 42.7, 47.9)	*
MONTANA	767	64.3	1.9	( 60.5, 68.1)	*	1030	49.9	1.7	( 46.6, 53.3)	*
NEBRASKA	1113	61.9	1.6	( 58.8, 65.0)	*	1700	43.2	1.4	( 40.6, 45.9)	*
NEVADA	1101	68.9	2.0	( 65.1, 72.7)	*	1086	57.2	2.3	( 52.7, 61.8)	*
NEW HAMPSHIRE	502	71.3	2.2	( 66.9, 75.6)	*	744	57.6	2.1	( 53.5, 61.6)	*
NEW JERSEY	1119	59.3	1.8	( 55.8, 62.9)	*	1692	49.2	1.5	( 46.4, 52.1)	*
NEW MEXICO	1532	62.9	1.4	( 60.2, 65.6)	*	1953	45.8	1.3	( 43.4, 48.3)	*
NEW YORK	1067	60.3	1.7	( 56.9, 63.7)	*	1577	45.7	1.4	( 42.9, 48.5)	*
NORTH CAROLINA	961	52.2	1.9	( 48.5, 55.9)	*	1478	32.2	1.5	( 29.3, 35.0)	*
NORTH DAKOTA	860	65.2	1.8	( 61.7, 68.7)	*	1118	45.9	1.6	( 42.7, 49.1)	*
OHIO	635	52.9	2.3	( 48.3, 57.4)	*	1001	35.7	1.8	( 32.1, 39.3)	*
OKLAHOMA	1154	35.7	1.6	( 32.5, 38.8)	*	1810	22.2	1.1	( 20.0, 24.3)	*
OREGON	767	65.7	1.9	( 62.0, 69.4)	*	1037	56.3	1.7	( 53.0, 59.7)	*
PENNSYLVANIA	1414	60.5	1.5	( 57.7, 63.4)	*	2134	43.5	1.2	( 41.1, 45.8)	*
RHODE ISLAND	1555	65.3	1.4	( 62.6, 68.1)	*	2435	52.1	1.2	( 49.7, 54.4)	*
SOUTH CAROLINA	1391	52.0	1.6	( 49.0, 55.1)	*	2066	32.9	1.2	( 30.6, 35.2)	*
SOUTH DAKOTA	1684	68.1	1.3	( 65.6, 70.6)	*	2460	49.6	1.1	( 47.4, 51.8)	*
TENNESSEE	1033	38.9	1.7	( 35.7, 42.1)	*	1977	21.5	1.0	( 19.4, 23.5)	*
TEXAS	2002	65.8	1.3	( 63.3, 68.2)	*	2984	44.4	1.2	( 42.1, 46.6)	*
UTAH	1371	36.1	1.7	( 32.7, 39.5)	*	1823	23.7	1.3	( 21.1, 26.2)	*
VERMONT	1272	66.8	1.5	( 63.9, 69.6)	*	1887	54.8	1.3	( 52.2, 57.3)	*
VIRGINIA	1408	61.8	1.9	( 58.2, 65.5)	*	2090	47.3	1.6	( 44.2, 50.4)	*
WASHINGTON	1595	66.6	1.5	( 63.6, 69.5)	*	2004	55.0	1.3	( 52.5, 57.6)	*
WEST VIRGINIA	1064	42.1	1.7	( 38.7, 45.4)	*	1522	24.2	1.3	( 21.8, 26.7)	*
WISCONSIN	920	80.6	1.4	( 77.8, 83.4)	*	1254	62.3	1.6	( 59.1, 65.4)	*
WYOMING	1064	66.4	1.6	( 63.4, 69.5)	*	1337	46.0	1.6	( 43.0, 49.1)	*
PUERTO RICO	1117	43.8	1.7	( 40.4, 47.2)	*	1931	21.2	1.1	( 18.9, 23.4)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	52	NUMBER OF PARTICIPANTS =	52
	MEDIAN	= 62.4	MEDIAN	= 46.0
	RANGE	= 35.7-80.6	RANGE	= 21.2-62.3

\* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH

June 20, 2000

TABLE 15.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 CURRENT DRINKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	36.8	39.6	2.8
ALASKA	55.2	56.9	1.7
ARIZONA	41.1	34.5	-6.6
ARKANSAS	30.6	38.3	7.7
CALIFORNIA	61.4	60.4	-1.0
COLORADO	62.0	64.2	2.1
CONNECTICUT	62.1	59.4	-2.7
DELAWARE	54.2	60.0	5.8
DISTRICT OF CO	43.7	50.3	6.7
FLORIDA	54.8	55.1	0.3
GEORGIA	44.6	47.7	3.1
HAWAII	51.1	48.6	-2.5
IDAHO	47.6	49.0	1.4
ILLINOIS	54.7	57.3	2.6
INDIANA	48.5	54.3	5.8
IOWA	53.9	55.0	1.1
KANSAS	41.3	46.8	5.5
KENTUCKY	33.7	34.9	1.2
LOUISIANA	50.6	45.4	-5.1
MAINE	55.4	59.8	4.4
MARYLAND	44.4	57.8	13.4
MASSACHUSETTS	64.8	61.5	-3.3
MICHIGAN	57.4	59.2	1.8
MINNESOTA	53.5	51.1	-2.4
MISSISSIPPI	35.1	38.6	3.5
MISSOURI	49.2	53.4	4.2
MONTANA	57.2	56.9	-0.2
NEBRASKA	58.0	52.2	-5.9
NEVADA	62.9	63.1	0.2
NEW HAMPSHIRE	61.3	64.2	2.9
NEW JERSEY	58.3	54.0	-4.3
NEW MEXICO	53.6	54.1	0.5
NEW YORK	53.5	52.6	-0.9
NORTH CAROLINA	37.6	41.7	4.1
NORTH DAKOTA	53.7	55.4	1.7
OHIO	33.7	43.8	10.1
OKLAHOMA	32.8	28.6	-4.1
OREGON	56.7	60.9	4.2
PENNSYLVANIA	49.3	51.5	2.2
RHODE ISLAND	59.4	58.3	-1.1
SOUTH CAROLINA	39.7	42.0	2.3
SOUTH DAKOTA	55.5	58.6	3.1
TENNESSEE	28.7	29.7	1.0
TEXAS	49.4	54.8	5.4
UTAH	28.3	29.7	1.5
VERMONT	60.2	60.5	0.3
VIRGINIA	54.9	54.3	-0.5
WASHINGTON	60.4	60.7	0.3
WEST VIRGINIA	31.0	32.6	1.6
WISCONSIN	70.2	71.1	0.9
WYOMING	52.6	56.1	3.5
PUERTO RICO	26.7	31.8	5.1

\* HAD ALCOHOLIC BEVERAGES DURING THE PAST MONTH

June 20, 2000

TABLE 16.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR BINGE DRINKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2072	11.7	0.8	( 10.1, 13.3)
ALASKA	2018	18.9	1.3	( 16.4, 21.5)
ARIZONA	1729	8.8	1.0	( 6.8, 10.8)
ARKANSAS	2985	10.3	0.7	( 9.0, 11.6)
CALIFORNIA	4137	15.5	0.7	( 14.2, 16.8)
COLORADO	3015	17.2	1.0	( 15.3, 19.2)
CONNECTICUT	3471	14.0	0.9	( 12.3, 15.7)
DELAWARE	2545	18.9	1.2	( 16.5, 21.3)
DISTRICT OF CO	1251	13.0	1.2	( 10.7, 15.2)
FLORIDA	5120	12.9	0.6	( 11.7, 14.1)
GEORGIA	2251	12.5	0.8	( 10.8, 14.1)
HAWAII	2136	14.0	1.0	( 12.1, 15.9)
IDAHO	4930	14.7	0.7	( 13.4, 16.0)
ILLINOIS	1437	19.7	1.3	( 17.2, 22.2)
INDIANA	2426	19.1	1.4	( 16.4, 21.9)
IOWA	3595	18.3	0.8	( 16.7, 19.9)
KANSAS	3831	11.8	0.6	( 10.5, 13.0)
KENTUCKY	7486	9.8	0.5	( 8.8, 10.9)
LOUISIANA	1645	15.0	1.1	( 12.9, 17.1)
MAINE	1664	14.8	1.2	( 12.5, 17.1)
MARYLAND	3925	15.9	0.8	( 14.3, 17.5)
MASSACHUSETTS	4973	17.4	0.8	( 16.0, 18.9)
MICHIGAN	2462	19.0	1.0	( 17.1, 21.0)
MINNESOTA	5306	16.3	0.6	( 15.2, 17.4)
MISSISSIPPI	2137	12.1	0.9	( 10.3, 13.9)
MISSOURI	4230	16.4	0.8	( 14.7, 18.0)
MONTANA	1779	17.6	1.1	( 15.5, 19.7)
NEBRASKA	2791	16.6	0.9	( 14.9, 18.3)
NEVADA	2170	21.0	1.3	( 18.4, 23.6)
NEW HAMPSHIRE	1231	20.0	1.5	( 17.1, 23.0)
NEW JERSEY	2789	12.3	0.8	( 10.8, 13.8)
NEW MEXICO	3468	15.0	0.7	( 13.6, 16.3)
NEW YORK	2623	13.9	0.8	( 12.3, 15.5)
NORTH CAROLINA	2407	12.0	0.9	( 10.3, 13.7)
NORTH DAKOTA	1967	19.7	1.0	( 17.7, 21.7)
OHIO	1626	12.1	1.1	( 10.0, 14.2)
OKLAHOMA	2939	8.1	0.6	( 6.9, 9.3)
OREGON	1795	14.9	0.9	( 13.0, 16.7)
PENNSYLVANIA	3515	15.9	0.7	( 14.4, 17.3)
RHODE ISLAND	3957	15.6	0.7	( 14.2, 16.9)
SOUTH CAROLINA	3414	12.3	0.7	( 10.9, 13.7)
SOUTH DAKOTA	4103	17.4	0.7	( 16.0, 18.9)
TENNESSEE	2990	7.7	0.6	( 6.5, 8.9)
TEXAS	4944	17.3	0.8	( 15.8, 18.8)
UTAH	3177	10.2	0.7	( 8.8, 11.7)
VERMONT	3127	17.4	0.9	( 15.6, 19.1)
VIRGINIA	3470	12.7	0.8	( 11.2, 14.3)
WASHINGTON	3577	14.4	0.8	( 12.8, 15.9)
WEST VIRGINIA	2578	8.6	0.6	( 7.3, 9.8)
WISCONSIN	2166	27.0	1.2	( 24.6, 29.3)
WYOMING	2388	17.3	0.9	( 15.5, 19.1)
PUERTO RICO	2949	10.6	0.8	( 9.2, 12.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 14.9 RANGE = 7.7-27.0

\* 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH

June 20, 2000



TABLE 16.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 BINGE DRINKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	11.4	11.7	0.3
ALASKA	16.5	18.9	2.4
ARIZONA	8.8	8.8	-0.0
ARKANSAS	9.2	10.3	1.1
CALIFORNIA	15.2	15.5	0.3
COLORADO	15.3	17.2	1.9
CONNECTICUT	15.6	14.0	-1.5
DELAWARE	11.9	18.9	7.0
DISTRICT OF CO	12.1	13.0	0.9
FLORIDA	13.1	12.9	-0.2
GEORGIA	9.4	12.5	3.1
HAWAII	17.1	14.0	-3.1
IDAHO	14.9	14.7	-0.2
ILLINOIS	16.3	19.7	3.4
INDIANA	12.6	19.1	6.5
IOWA	17.9	18.3	0.4
KANSAS	13.3	11.8	-1.6
KENTUCKY	9.4	9.8	0.4
LOUISIANA	15.2	15.0	-0.2
MAINE	13.8	14.8	1.0
MARYLAND	6.3	15.9	9.6
MASSACHUSETTS	17.9	17.4	-0.5
MICHIGAN	18.9	19.0	0.1
MINNESOTA	15.6	16.3	0.7
MISSISSIPPI	9.5	12.1	2.5
MISSOURI	15.1	16.4	1.3
MONTANA	14.0	17.6	3.5
NEBRASKA	16.3	16.6	0.3
NEVADA	19.2	21.0	1.8
NEW HAMPSHIRE	16.1	20.0	3.9
NEW JERSEY	13.1	12.3	-0.8
NEW MEXICO	14.6	15.0	0.4
NEW YORK	9.2	13.9	4.7
NORTH CAROLINA	9.0	12.0	3.0
NORTH DAKOTA	18.4	19.7	1.3
OHIO	8.7	12.1	3.4
OKLAHOMA	8.8	8.1	-0.7
OREGON	14.3	14.9	0.5
PENNSYLVANIA	14.6	15.9	1.3
RHODE ISLAND	14.9	15.6	0.6
SOUTH CAROLINA	9.7	12.3	2.6
SOUTH DAKOTA	20.9	17.4	-3.5
TENNESSEE	7.2	7.7	0.5
TEXAS	17.4	17.3	-0.1
UTAH	7.7	10.2	2.5
VERMONT	16.1	17.4	1.3
VIRGINIA	14.5	12.7	-1.7
WASHINGTON	14.5	14.4	-0.1
WEST VIRGINIA	8.4	8.6	0.2
WISCONSIN	23.3	27.0	3.7
WYOMING	15.4	17.3	1.9
PUERTO RICO	10.9	10.6	-0.3

\* 5 OR MORE DRINKS ON AT LEAST ONE OCCASION DURING THE PAST MONTH

June 20, 2000

TABLE 17.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHRONIC DRINKERS\*  
DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2058	3.1	0.4	( 2.3, 4.0)
ALASKA	2011	4.8	0.7	( 3.5, 6.1)
ARIZONA	1722	4.4	0.7	( 3.0, 5.8)
ARKANSAS	2964	2.5	0.3	( 1.9, 3.1)
COLORADO	3001	4.5	0.6	( 3.3, 5.7)
CONNECTICUT	3426	3.9	0.5	( 2.9, 5.0)
DELAWARE	2524	3.9	0.6	( 2.7, 5.1)
DISTRICT OF CO	1229	3.1	0.6	( 1.9, 4.4)
FLORIDA	5074	3.6	0.3	( 3.0, 4.3)
GEORGIA	2223	3.1	0.5	( 2.2, 4.1)
HAWAII	2130	5.2	0.7	( 3.9, 6.5)
IDAHO	4893	3.5	0.3	( 2.9, 4.2)
ILLINOIS	1426	4.6	0.7	( 3.3, 5.9)
INDIANA	2406	6.2	0.9	( 4.4, 7.9)
IOWA	3580	4.4	0.5	( 3.5, 5.3)
KANSAS	3790	3.3	0.4	( 2.5, 4.2)
KENTUCKY	7449	2.7	0.3	( 2.1, 3.2)
LOUISIANA	1615	4.0	0.6	( 2.9, 5.2)
MAINE	1646	3.1	0.8	( 1.6, 4.6)
MARYLAND	3859	4.1	0.5	( 3.2, 5.0)
MASSACHUSETTS	4899	4.0	0.4	( 3.3, 4.7)
MICHIGAN	2455	5.7	0.7	( 4.3, 7.1)
MINNESOTA	5258	4.7	0.3	( 4.1, 5.4)
MISSISSIPPI	2122	3.6	0.5	( 2.7, 4.6)
MISSOURI	4181	4.1	0.5	( 3.2, 5.0)
MONTANA	1756	3.6	0.6	( 2.5, 4.7)
NEBRASKA	2766	3.0	0.4	( 2.2, 3.7)
NEVADA	2149	7.4	0.9	( 5.8, 9.1)
NEW HAMPSHIRE	1219	4.9	0.9	( 3.2, 6.6)
NEW JERSEY	2752	2.5	0.4	( 1.8, 3.3)
NEW MEXICO	3444	3.8	0.4	( 3.1, 4.6)
NEW YORK	2570	3.7	0.5	( 2.8, 4.6)
NORTH CAROLINA	2380	2.4	0.4	( 1.7, 3.0)
NORTH DAKOTA	1959	3.5	0.5	( 2.5, 4.5)
OHIO	1609	1.7	0.4	( 0.9, 2.5)
OKLAHOMA	2916	2.2	0.3	( 1.5, 2.8)
OREGON	1776	3.5	0.5	( 2.6, 4.5)
PENNSYLVANIA	3468	3.8	0.4	( 3.0, 4.7)
RHODE ISLAND	3897	4.3	0.4	( 3.6, 5.1)
SOUTH CAROLINA	3376	4.0	0.5	( 3.0, 4.9)
SOUTH DAKOTA	4053	3.0	0.3	( 2.3, 3.6)
TENNESSEE	2946	2.1	0.3	( 1.4, 2.7)
TEXAS	4915	4.5	0.4	( 3.8, 5.3)
UTAH	3164	2.9	0.4	( 2.0, 3.7)
VERMONT	3105	4.5	0.5	( 3.4, 5.5)
VIRGINIA	3409	2.9	0.4	( 2.1, 3.6)
WASHINGTON	3525	3.6	0.4	( 2.8, 4.5)
WEST VIRGINIA	2569	2.7	0.4	( 2.0, 3.5)
WISCONSIN	2156	5.0	0.6	( 4.0, 6.1)
WYOMING	2368	3.1	0.4	( 2.3, 3.9)
PUERTO RICO	2896	3.0	0.4	( 2.1, 3.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 3.6 RANGE = 1.7- 7.4

\* 60 OR MORE DRINKS DURING THE PAST MONTH

June 20, 2000

TABLE 17.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHRONIC DRINKERS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	831	5.6	0.9	( 3.9, 7.3)	*	1227	1.0	0.3	( 0.4, 1.5)	*
ALASKA	976	7.8	1.2	( 5.5, 10.1)	*	1035	1.7	0.6	( 0.5, 2.8)	*
ARIZONA	699	8.8	1.5	( 5.9, 11.6)	*	1023	0.4	0.2	( 0.1, 0.7)	*
ARKANSAS	1155	4.7	0.6	( 3.5, 5.9)	*	1809	0.5	0.2	( 0.2, 0.9)	*
COLORADO	1350	7.3	1.1	( 5.1, 9.5)	*	1651	1.9	0.5	( 0.9, 2.8)	*
CONNECTICUT	1317	7.9	1.1	( 5.7, 10.0)	*	2109	0.4	0.1	( 0.1, 0.6)	*
DELAWARE	1007	5.1	0.9	( 3.4, 6.8)	*	1517	2.8	0.9	( 1.0, 4.6)	*
DISTRICT OF CO	445	5.3	1.2	( 2.9, 7.7)	*	784	1.3	0.5	( 0.3, 2.3)	*
FLORIDA	2003	6.3	0.6	( 5.1, 7.5)	*	3071	1.2	0.2	( 0.8, 1.6)	*
GEORGIA	855	6.0	1.0	( 4.1, 7.9)	*	1368	0.6	0.2	( 0.2, 1.0)	*
HAWAII	980	9.1	1.3	( 6.6, 11.5)	*	1150	1.3	0.4	( 0.5, 2.1)	*
IDAHO	2066	6.3	0.7	( 5.0, 7.6)	*	2827	0.9	0.2	( 0.5, 1.2)	*
ILLINOIS	562	6.6	1.2	( 4.2, 8.9)	*	864	2.6	0.7	( 1.4, 3.9)	*
INDIANA	1001	10.8	1.7	( 7.5, 14.1)	*	1405	2.0	0.7	( 0.7, 3.2)	*
IOWA	1405	8.7	0.9	( 6.9, 10.5)	*	2175	0.6	0.2	( 0.2, 0.9)	*
KANSAS	1526	6.2	0.9	( 4.5, 7.9)	*	2264	0.6	0.2	( 0.2, 1.0)	*
KENTUCKY	2688	5.2	0.6	( 4.1, 6.4)	*	4761	0.4	0.1	( 0.2, 0.7)	*
LOUISIANA	638	7.9	1.2	( 5.5, 10.2)	*	977	0.6	0.3	( 0.1, 1.1)	*
MAINE	722	5.9	1.6	( 2.8, 9.0)	*	924	0.5	0.3	( -0.0, 1.1)	*
MARYLAND	1539	6.8	0.8	( 5.1, 8.4)	*	2320	1.7	0.5	( 0.7, 2.6)	*
MASSACHUSETTS	1953	7.4	0.7	( 6.0, 8.7)	*	2946	0.9	0.2	( 0.5, 1.4)	*
MICHIGAN	1008	10.4	1.4	( 7.5, 13.2)	*	1447	1.4	0.3	( 0.8, 2.1)	*
MINNESOTA	2386	7.7	0.6	( 6.5, 8.9)	*	2872	2.0	0.3	( 1.4, 2.5)	*
MISSISSIPPI	826	7.0	1.0	( 5.0, 9.0)	*	1296	0.7	0.2	( 0.2, 1.1)	*
MISSOURI	1669	7.5	0.9	( 5.7, 9.2)	*	2512	1.0	0.3	( 0.4, 1.7)	*
MONTANA	742	7.0	1.1	( 4.8, 9.2)	*	1014	0.4	0.1	( 0.1, 0.6)	*
NEBRASKA	1090	5.3	0.8	( 3.9, 6.8)	*	1676	0.8	0.2	( 0.4, 1.3)	*
NEVADA	1083	11.9	1.5	( 9.0, 14.7)	*	1066	2.9	0.8	( 1.3, 4.5)	*
NEW HAMPSHIRE	487	8.4	1.7	( 5.1, 11.7)	*	732	1.6	0.6	( 0.4, 2.8)	*
NEW JERSEY	1090	4.6	0.8	( 3.0, 6.1)	*	1662	0.7	0.2	( 0.3, 1.2)	*
NEW MEXICO	1505	7.1	0.7	( 5.7, 8.5)	*	1939	0.8	0.2	( 0.4, 1.2)	*
NEW YORK	1034	6.6	0.9	( 4.7, 8.4)	*	1536	1.1	0.3	( 0.5, 1.8)	*
NORTH CAROLINA	924	4.3	0.7	( 2.9, 5.7)	*	1456	0.6	0.2	( 0.2, 1.0)	*
NORTH DAKOTA	846	6.0	1.0	( 4.1, 7.9)	*	1113	1.1	0.4	( 0.3, 2.0)	*
OHIO	623	3.0	0.7	( 1.5, 4.4)	*	986	0.6	0.3	( -0.1, 1.2)	*
OKLAHOMA	1122	3.8	0.7	( 2.4, 5.1)	*	1794	0.7	0.2	( 0.3, 1.2)	*
OREGON	756	5.8	0.9	( 4.1, 7.6)	*	1020	1.4	0.4	( 0.6, 2.2)	*
PENNSYLVANIA	1376	7.1	0.9	( 5.4, 8.8)	*	2092	0.9	0.3	( 0.4, 1.4)	*
RHODE ISLAND	1515	7.4	0.7	( 5.9, 8.8)	*	2382	1.6	0.3	( 1.0, 2.3)	*
SOUTH CAROLINA	1351	7.6	1.0	( 5.8, 9.5)	*	2025	0.6	0.2	( 0.3, 1.0)	*
SOUTH DAKOTA	1644	5.2	0.7	( 3.9, 6.5)	*	2409	0.9	0.2	( 0.4, 1.3)	*
TENNESSEE	999	4.0	0.7	( 2.6, 5.3)	*	1947	0.4	0.2	( 0.1, 0.7)	*
TEXAS	1966	8.1	0.8	( 6.6, 9.6)	*	2949	1.2	0.2	( 0.8, 1.6)	*
UTAH	1355	5.3	0.8	( 3.7, 6.9)	*	1809	0.5	0.2	( 0.0, 1.0)	*
VERMONT	1246	7.9	1.0	( 5.9, 9.9)	*	1859	1.3	0.4	( 0.6, 2.0)	*
VIRGINIA	1365	5.0	0.7	( 3.5, 6.4)	*	2044	0.9	0.3	( 0.3, 1.5)	*
WASHINGTON	1563	6.2	0.9	( 4.5, 7.9)	*	1962	1.2	0.3	( 0.7, 1.7)	*
WEST VIRGINIA	1054	5.6	0.8	( 4.1, 7.1)	*	1515	0.2	0.1	( 0.0, 0.4)	*
WISCONSIN	913	9.1	1.1	( 7.0, 11.2)	*	1243	1.3	0.4	( 0.6, 1.9)	*
WYOMING	1049	6.0	0.8	( 4.4, 7.5)	*	1319	0.3	0.1	( 0.1, 0.6)	*
PUERTO RICO	1017	5.5	0.9	( 3.8, 7.2)	*	1879	1.0	0.3	( 0.3, 1.6)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	51	NUMBER OF PARTICIPANTS =	51
	MEDIAN	= 6.6	MEDIAN	= 0.9
	RANGE	= 3.0-11.9	RANGE	= 0.2- 2.9

\* 60 OR MORE DRINKS DURING THE PAST MONTH

June 20, 2000

TABLE 17.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 CHRONIC DRINKERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	3.0	3.1	0.2
ALASKA	4.0	4.8	0.9
ARIZONA	2.6	4.4	1.8
ARKANSAS	2.3	2.5	0.2
COLORADO	2.7	4.5	1.8
CONNECTICUT	3.2	3.9	0.7
DELAWARE	3.4	3.9	0.5
DISTRICT OF CO	3.0	3.1	0.2
FLORIDA	4.0	3.6	-0.3
GEORGIA	1.8	3.1	1.4
HAWAII	5.1	5.2	0.2
IDAHO	3.0	3.5	0.5
ILLINOIS	3.4	4.6	1.2
INDIANA	3.0	6.2	3.2
IOWA	3.5	4.4	1.0
KANSAS	2.3	3.3	1.0
KENTUCKY	2.7	2.7	-0.1
LOUISIANA	4.6	4.0	-0.6
MAINE	2.7	3.1	0.4
MARYLAND	1.2	4.1	2.9
MASSACHUSETTS	3.6	4.0	0.4
MICHIGAN	3.8	5.7	1.9
MINNESOTA	2.7	4.7	2.0
MISSISSIPPI	2.5	3.6	1.2
MISSOURI	3.2	4.1	0.8
MONTANA	2.6	3.6	1.0
NEBRASKA	2.3	3.0	0.7
NEVADA	4.7	7.4	2.7
NEW HAMPSHIRE	2.5	4.9	2.4
NEW JERSEY	2.0	2.5	0.5
NEW MEXICO	3.6	3.8	0.3
NEW YORK	2.4	3.7	1.3
NORTH CAROLINA	2.7	2.4	-0.3
NORTH DAKOTA	2.5	3.5	1.0
OHIO	2.4	1.7	-0.7
OKLAHOMA	2.2	2.2	-0.0
OREGON	3.5	3.5	-0.0
PENNSYLVANIA	3.1	3.8	0.7
RHODE ISLAND	4.6	4.3	-0.3
SOUTH CAROLINA	3.3	4.0	0.7
SOUTH DAKOTA	3.2	3.0	-0.2
TENNESSEE	1.8	2.1	0.3
TEXAS	4.9	4.5	-0.3
UTAH	1.7	2.9	1.2
VERMONT	4.1	4.5	0.3
VIRGINIA	3.3	2.9	-0.4
WASHINGTON	3.6	3.6	0.0
WEST VIRGINIA	2.1	2.7	0.6
WISCONSIN	5.0	5.0	0.1
WYOMING	2.8	3.1	0.4
PUERTO RICO	3.7	3.0	-0.7

\* 60 OR MORE DRINKS DURING THE PAST MONTH

June 20, 2000

TABLE 18.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DRINKING AND DRIVING\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2089	1.9	0.4	( 1.2, 2.5)
ALASKA	2042	2.1	0.4	( 1.3, 2.9)
ARIZONA	1730	1.8	0.5	( 0.8, 2.8)
ARKANSAS	2994	1.5	0.3	( 0.9, 2.0)
CALIFORNIA	4147	2.3	0.3	( 1.8, 2.9)
COLORADO	3040	3.6	0.5	( 2.5, 4.6)
CONNECTICUT	3503	2.9	0.5	( 1.9, 3.8)
DELAWARE	2553	3.2	0.5	( 2.2, 4.2)
DISTRICT OF CO	1261	1.4	0.4	( 0.5, 2.2)
FLORIDA	5148	2.0	0.2	( 1.5, 2.4)
GEORGIA	2267	1.5	0.3	( 0.9, 2.0)
HAWAII	2146	2.3	0.4	( 1.4, 3.1)
IDAHO	4949	1.8	0.2	( 1.3, 2.2)
ILLINOIS	1443	4.4	0.7	( 3.0, 5.7)
INDIANA	2435	3.2	0.5	( 2.2, 4.3)
IOWA	3609	3.9	0.4	( 3.0, 4.7)
KANSAS	3858	2.8	0.3	( 2.2, 3.5)
KENTUCKY	7500	1.6	0.3	( 1.1, 2.1)
LOUISIANA	1660	3.6	0.6	( 2.5, 4.7)
MAINE	1667	1.1	0.3	( 0.5, 1.7)
MARYLAND	3946	2.4	0.4	( 1.7, 3.2)
MASSACHUSETTS	5002	2.8	0.3	( 2.1, 3.4)
MICHIGAN	2510	3.1	0.4	( 2.3, 3.8)
MINNESOTA	5325	4.1	0.3	( 3.5, 4.7)
MISSISSIPPI	2172	2.7	0.5	( 1.7, 3.7)
MISSOURI	4260	3.0	0.4	( 2.2, 3.8)
MONTANA	1789	3.4	0.5	( 2.4, 4.5)
NEBRASKA	2800	3.7	0.4	( 2.8, 4.5)
NEVADA	2180	5.5	0.7	( 4.0, 6.9)
NEW HAMPSHIRE	1238	3.8	0.7	( 2.5, 5.1)
NEW JERSEY	2807	1.3	0.3	( 0.9, 1.8)
NEW MEXICO	3481	2.3	0.3	( 1.8, 2.9)
NEW YORK	2638	1.6	0.3	( 1.1, 2.1)
NORTH CAROLINA	2431	1.7	0.4	( 0.9, 2.4)
NORTH DAKOTA	1970	4.4	0.6	( 3.3, 5.6)
OHIO	1634	1.2	0.3	( 0.5, 1.8)
OKLAHOMA	2952	2.5	0.3	( 2.0, 3.1)
OREGON	1801	1.9	0.3	( 1.3, 2.6)
PENNSYLVANIA	3537	2.4	0.3	( 1.9, 3.0)
RHODE ISLAND	3980	2.6	0.3	( 2.0, 3.2)
SOUTH CAROLINA	3445	2.1	0.3	( 1.5, 2.7)
SOUTH DAKOTA	4130	4.1	0.4	( 3.4, 4.8)
TENNESSEE	3000	1.5	0.3	( 1.0, 2.0)
TEXAS	4980	2.8	0.3	( 2.3, 3.3)
UTAH	3191	1.2	0.3	( 0.6, 1.7)
VERMONT	3155	2.4	0.4	( 1.7, 3.1)
VIRGINIA	3491	2.4	0.4	( 1.7, 3.1)
WASHINGTON	3589	1.7	0.4	( 1.0, 2.4)
WEST VIRGINIA	2579	1.1	0.2	( 0.6, 1.5)
WISCONSIN	2172	4.9	0.6	( 3.8, 6.0)
WYOMING	2399	2.4	0.3	( 1.8, 3.1)
PUERTO RICO	3008	2.0	0.3	( 1.4, 2.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 2.4 RANGE = 1.1- 5.5  
 \* DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES IN PAST MONTH June 20, 2000



TABLE 18.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 DRINKING AND DRIVING\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	1.5	1.9	0.3
ALASKA	2.2	2.1	-0.1
ARIZONA	1.4	1.8	0.4
ARKANSAS	1.6	1.5	-0.1
CALIFORNIA	2.5	2.3	-0.2
COLORADO	2.5	3.6	1.1
CONNECTICUT	2.1	2.9	0.8
DELAWARE	2.0	3.2	1.2
DISTRICT OF CO	2.5	1.4	-1.2
FLORIDA	1.9	2.0	0.1
GEORGIA	1.1	1.5	0.4
HAWAII	2.4	2.3	-0.1
IDAHO	1.3	1.8	0.5
ILLINOIS	2.8	4.4	1.5
INDIANA	1.9	3.2	1.3
IOWA	3.8	3.9	0.1
KANSAS	2.7	2.8	0.1
KENTUCKY	0.6	1.6	1.0
LOUISIANA	3.2	3.6	0.4
MAINE	0.9	1.1	0.2
MARYLAND	0.9	2.4	1.5
MASSACHUSETTS	1.8	2.8	1.0
MICHIGAN	3.5	3.1	-0.4
MINNESOTA	3.8	4.1	0.3
MISSISSIPPI	1.2	2.7	1.5
MISSOURI	3.0	3.0	0.0
MONTANA	2.6	3.4	0.8
NEBRASKA	3.8	3.7	-0.1
NEVADA	3.1	5.5	2.4
NEW HAMPSHIRE	1.9	3.8	1.9
NEW JERSEY	1.5	1.3	-0.2
NEW MEXICO	1.7	2.3	0.6
NEW YORK	0.9	1.6	0.7
NORTH CAROLINA	1.1	1.7	0.6
NORTH DAKOTA	3.8	4.4	0.7
OHIO	1.0	1.2	0.1
OKLAHOMA	1.5	2.5	1.1
OREGON	1.6	1.9	0.4
PENNSYLVANIA	1.7	2.4	0.7
RHODE ISLAND	1.6	2.6	1.0
SOUTH CAROLINA	0.9	2.1	1.2
SOUTH DAKOTA	3.7	4.1	0.4
TENNESSEE	1.0	1.5	0.5
TEXAS	4.0	2.8	-1.2
UTAH	0.8	1.2	0.3
VERMONT	3.0	2.4	-0.6
VIRGINIA	2.4	2.4	0.1
WASHINGTON	1.7	1.7	0.0
WEST VIRGINIA	0.8	1.1	0.2
WISCONSIN	5.3	4.9	-0.4
WYOMING	2.6	2.4	-0.2
PUERTO RICO	3.2	2.0	-1.2

\* DRIVING AFTER HAVING TOO MUCH TO DRINK, ONE OR MORE TIMES IN PAST MONTH

June 20, 2000

TABLE 19.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO FLU SHOT, AGE 65 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	429	35.4	2.5	( 30.5, 40.2)
ALASKA	160	40.2	5.6	( 29.2, 51.3)
ARIZONA	466	28.7	3.0	( 22.7, 34.6)
ARKANSAS	575	32.7	2.2	( 28.5, 37.0)
CALIFORNIA	616	27.8	2.1	( 23.7, 31.9)
COLORADO	480	25.2	2.8	( 19.7, 30.8)
CONNECTICUT	610	35.2	2.6	( 30.2, 40.2)
DELAWARE	443	32.3	2.8	( 26.8, 37.8)
DISTRICT OF CO	245	44.2	3.4	( 37.4, 50.9)
FLORIDA	1239	36.9	1.8	( 33.3, 40.4)
GEORGIA	319	43.1	3.2	( 36.8, 49.3)
HAWAII	395	25.9	3.1	( 19.8, 32.0)
IDAHO	892	31.0	1.9	( 27.4, 34.7)
ILLINOIS	262	32.5	3.2	( 26.2, 38.7)
INDIANA	401	33.9	3.9	( 26.2, 41.5)
IOWA	880	30.4	1.8	( 26.9, 34.0)
KANSAS	859	33.0	1.8	( 29.5, 36.5)
KENTUCKY	1721	31.6	1.5	( 28.7, 34.6)
LOUISIANA	313	39.7	3.1	( 33.7, 45.7)
MAINE	355	26.3	2.7	( 21.0, 31.6)
MARYLAND	616	37.4	2.5	( 32.5, 42.3)
MASSACHUSETTS	864	30.6	1.9	( 26.9, 34.3)
MICHIGAN	462	30.0	2.3	( 25.5, 34.5)
MINNESOTA	903	36.0	1.7	( 32.6, 39.4)
MISSISSIPPI	380	37.2	2.7	( 31.9, 42.5)
MISSOURI	944	31.6	2.1	( 27.5, 35.7)
MONTANA	384	27.1	2.5	( 22.3, 31.9)
NEBRASKA	714	30.8	1.9	( 27.1, 34.6)
NEVADA	335	37.9	4.2	( 29.6, 46.1)
NEW HAMPSHIRE	220	34.9	3.5	( 28.0, 41.8)
NEW JERSEY	536	34.7	2.3	( 30.1, 39.3)
NEW MEXICO	631	31.2	2.1	( 27.2, 35.2)
NEW YORK	427	36.2	2.6	( 31.2, 41.2)
NORTH CAROLINA	522	35.8	2.3	( 31.3, 40.4)
NORTH DAKOTA	475	32.8	2.3	( 28.2, 37.4)
OHIO	396	31.2	2.7	( 25.9, 36.4)
OKLAHOMA	658	28.2	2.0	( 24.3, 32.0)
OREGON	346	34.8	2.8	( 29.4, 40.3)
PENNSYLVANIA	690	36.9	2.0	( 32.9, 40.9)
RHODE ISLAND	773	24.2	1.8	( 20.8, 27.6)
SOUTH CAROLINA	519	30.1	2.2	( 25.8, 34.4)
SOUTH DAKOTA	950	26.4	1.5	( 23.4, 29.3)
TENNESSEE	574	34.5	2.3	( 30.1, 39.0)
TEXAS	693	30.2	2.0	( 26.2, 34.1)
UTAH	543	24.9	2.4	( 20.1, 29.6)
VERMONT	606	26.6	1.9	( 22.8, 30.3)
VIRGINIA	572	34.3	2.7	( 29.1, 39.6)
WASHINGTON	609	31.1	2.1	( 27.0, 35.2)
WEST VIRGINIA	612	37.1	2.1	( 33.0, 41.3)
WISCONSIN	397	35.1	2.6	( 30.0, 40.2)
WYOMING	398	26.2	2.4	( 21.5, 30.8)
PUERTO RICO	669	59.7	2.1	( 55.6, 63.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 32.6 RANGE = 24.2-59.7 YEAR 2000 OBJ. 20.11  
 \* DID NOT HAVE FLU SHOT WITHIN PAST YEAR June 20, 2000

TABLE 19.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO FLU SHOT, AGE 65 AND OVER\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	156	37.0	4.1	( 29.0, 44.9)	*	273	34.3	3.1	( 28.3, 40.4)	*
ALASKA	72	36.9	7.8	( 21.7, 52.1)	*	88	43.3	8.0	( 27.7, 58.9)	*
ARIZONA	179	29.0	5.0	( 19.2, 38.8)	*	287	28.5	3.8	( 21.1, 35.8)	*
ARKANSAS	172	34.3	3.8	( 26.8, 41.7)	*	403	31.7	2.6	( 26.6, 36.8)	*
CALIFORNIA	223	23.5	3.4	( 16.8, 30.2)	*	393	30.7	2.6	( 25.6, 35.9)	*
COLORADO	178	31.1	5.1	( 21.2, 41.0)	*	302	21.0	3.2	( 14.8, 27.3)	*
CONNECTICUT	221	35.2	4.2	( 26.9, 43.5)	*	389	35.3	3.2	( 29.0, 41.5)	*
DELAWARE	176	35.5	4.6	( 26.5, 44.5)	*	267	30.1	3.5	( 23.2, 36.9)	*
DISTRICT OF CO	86	40.7	6.0	( 29.0, 52.4)	*	159	46.2	4.2	( 37.9, 54.5)	*
FLORIDA	470	36.3	3.0	( 30.4, 42.2)	*	769	37.3	2.1	( 33.2, 41.3)	*
GEORGIA	96	43.7	5.9	( 32.3, 55.2)	*	223	42.6	3.6	( 35.6, 49.6)	*
HAWAII	160	31.0	5.3	( 20.7, 41.3)	*	235	21.7	3.5	( 14.9, 28.5)	*
IDAHO	336	34.5	3.1	( 28.5, 40.5)	*	556	28.4	2.3	( 24.0, 32.9)	*
ILLINOIS	74	30.0	5.6	( 19.1, 40.9)	*	188	34.1	3.8	( 26.7, 41.5)	*
INDIANA	138	29.2	6.4	( 16.6, 41.8)	*	263	37.0	4.9	( 27.4, 46.5)	*
IOWA	283	31.1	3.2	( 24.8, 37.3)	*	597	30.0	2.2	( 25.8, 34.2)	*
KANSAS	296	28.8	2.9	( 23.2, 34.4)	*	563	35.9	2.3	( 31.4, 40.3)	*
KENTUCKY	504	29.4	2.7	( 24.1, 34.7)	*	1217	33.1	1.7	( 29.7, 36.5)	*
LOUISIANA	103	44.6	5.3	( 34.2, 54.9)	*	210	36.5	3.6	( 29.4, 43.6)	*
MAINE	133	32.5	4.9	( 22.9, 42.1)	*	222	22.1	3.0	( 16.2, 28.0)	*
MARYLAND	225	38.0	4.1	( 29.9, 46.0)	*	391	37.1	3.1	( 31.0, 43.1)	*
MASSACHUSETTS	293	29.0	3.1	( 23.0, 35.1)	*	571	31.6	2.4	( 27.0, 36.3)	*
MICHIGAN	167	25.8	3.6	( 18.9, 32.8)	*	295	32.9	3.0	( 27.0, 38.8)	*
MINNESOTA	319	38.7	2.9	( 33.1, 44.3)	*	584	34.1	2.1	( 30.0, 38.2)	*
MISSISSIPPI	114	36.5	4.7	( 27.2, 45.7)	*	266	37.7	3.2	( 31.3, 44.0)	*
MISSOURI	315	33.0	3.6	( 26.0, 40.0)	*	629	30.7	2.6	( 25.6, 35.7)	*
MONTANA	124	27.2	4.3	( 18.8, 35.6)	*	260	27.1	2.9	( 21.4, 32.7)	*
NEBRASKA	227	35.9	3.4	( 29.3, 42.5)	*	487	27.3	2.2	( 23.0, 31.6)	*
NEVADA	153	32.8	5.0	( 23.0, 42.5)	*	182	41.9	6.3	( 29.7, 54.2)	*
NEW HAMPSHIRE	85	35.9	5.6	( 25.0, 46.8)	*	135	34.2	4.6	( 25.2, 43.1)	*
NEW JERSEY	201	33.3	3.9	( 25.6, 40.9)	*	335	35.6	2.9	( 29.9, 41.3)	*
NEW MEXICO	244	33.3	3.2	( 27.1, 39.6)	*	387	29.6	2.7	( 24.4, 34.8)	*
NEW YORK	144	32.4	4.2	( 24.1, 40.6)	*	283	38.7	3.2	( 32.5, 44.9)	*
NORTH CAROLINA	166	35.5	4.1	( 27.5, 43.5)	*	356	36.0	2.8	( 30.6, 41.5)	*
NORTH DAKOTA	157	34.0	4.0	( 26.2, 41.8)	*	318	31.9	2.8	( 26.5, 37.4)	*
OHIO	125	30.0	4.5	( 21.1, 38.8)	*	271	31.9	3.2	( 25.6, 38.3)	*
OKLAHOMA	205	30.1	3.4	( 23.4, 36.8)	*	453	26.9	2.3	( 22.4, 31.4)	*
OREGON	123	35.4	4.6	( 26.5, 44.4)	*	223	34.4	3.4	( 27.7, 41.2)	*
PENNSYLVANIA	250	33.4	3.2	( 27.1, 39.7)	*	440	39.2	2.6	( 34.0, 44.3)	*
RHODE ISLAND	264	23.1	2.9	( 17.4, 28.7)	*	509	24.9	2.2	( 20.6, 29.2)	*
SOUTH CAROLINA	191	28.9	3.6	( 22.0, 35.9)	*	328	30.8	2.8	( 25.4, 36.2)	*
SOUTH DAKOTA	343	25.8	2.5	( 21.0, 30.7)	*	607	26.8	1.9	( 23.0, 30.5)	*
TENNESSEE	161	33.5	4.0	( 25.6, 41.3)	*	413	35.3	2.6	( 30.2, 40.3)	*
TEXAS	239	28.3	3.3	( 21.8, 34.8)	*	454	31.5	2.5	( 26.6, 36.4)	*
UTAH	207	27.2	3.9	( 19.5, 34.8)	*	336	23.1	3.0	( 17.2, 29.0)	*
VERMONT	190	22.9	3.2	( 16.6, 29.2)	*	416	29.1	2.4	( 24.4, 33.7)	*
VIRGINIA	204	34.5	4.3	( 26.0, 42.9)	*	368	34.2	3.4	( 27.6, 40.8)	*
WASHINGTON	223	31.2	3.6	( 24.1, 38.2)	*	386	31.0	2.6	( 26.0, 36.1)	*
WEST VIRGINIA	206	35.5	3.6	( 28.5, 42.5)	*	406	38.2	2.6	( 33.1, 43.3)	*
WISCONSIN	138	35.6	4.4	( 26.9, 44.2)	*	259	34.7	3.2	( 28.5, 41.0)	*
WYOMING	148	30.6	4.0	( 22.8, 38.4)	*	250	22.9	2.9	( 17.2, 28.5)	*
PUERTO RICO	255	59.5	3.3	( 53.0, 66.0)	*	414	59.8	2.7	( 54.5, 65.1)	*

SUMMARY STATISTICS:      NUMBER OF PARTICIPANTS = 52      NUMBER OF PARTICIPANTS = 52  
    MEDIAN = 33.1      MEDIAN = 33.0  
    RANGE = 22.9-59.5      RANGE = 21.0-59.8

\* DID NOT HAVE FLU SHOT WITHIN PAST YEAR

June 20, 2000

TABLE 19.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 NO FLU SHOT, AGE 65 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	37.4	35.4	-2.1
ALASKA	41.7	40.2	-1.5
ARIZONA	27.1	28.7	1.6
ARKANSAS	38.9	32.7	-6.2
CALIFORNIA	34.5	27.8	-6.7
COLORADO	25.6	25.2	-0.3
CONNECTICUT	32.8	35.2	2.5
DELAWARE	31.4	32.3	0.9
DISTRICT OF CO	45.7	44.2	-1.6
FLORIDA	37.7	36.9	-0.8
GEORGIA	41.5	43.1	1.5
HAWAII	28.9	25.9	-3.0
IDAHO	33.6	31.0	-2.5
ILLINOIS	32.2	32.5	0.3
INDIANA	37.5	33.9	-3.6
IOWA	30.3	30.4	0.1
KANSAS	38.5	33.0	-5.6
KENTUCKY	38.8	31.6	-7.1
LOUISIANA	41.6	39.7	-1.9
MAINE	27.9	26.3	-1.6
MARYLAND	36.6	37.4	0.9
MASSACHUSETTS	34.0	30.6	-3.3
MICHIGAN	36.4	30.0	-6.4
MINNESOTA	31.0	36.0	5.0
MISSISSIPPI	38.9	37.2	-1.7
MISSOURI	29.7	31.6	1.9
MONTANA	31.6	27.1	-4.5
NEBRASKA	34.2	30.8	-3.4
NEVADA	43.5	37.9	-5.6
NEW HAMPSHIRE	35.4	34.9	-0.5
NEW JERSEY	39.3	34.7	-4.6
NEW MEXICO	27.2	31.2	4.0
NEW YORK	35.6	36.2	0.6
NORTH CAROLINA	35.4	35.8	0.4
NORTH DAKOTA	35.2	32.8	-2.4
OHIO	34.6	31.2	-3.5
OKLAHOMA	30.7	28.2	-2.5
OREGON	30.2	34.8	4.7
PENNSYLVANIA	34.2	36.9	2.6
RHODE ISLAND	32.3	24.2	-8.1
SOUTH CAROLINA	25.7	30.1	4.4
SOUTH DAKOTA	34.4	26.4	-8.1
TENNESSEE	30.9	34.5	3.6
TEXAS	32.0	30.2	-1.8
UTAH	33.9	24.9	-9.0
VERMONT	30.6	26.6	-4.0
VIRGINIA	32.3	34.3	2.0
WASHINGTON	29.8	31.1	1.3
WEST VIRGINIA	41.8	37.1	-4.7
WISCONSIN	33.9	35.1	1.2
WYOMING	27.6	26.2	-1.4
PUERTO RICO	58.5	59.7	1.2

\* DID NOT HAVE FLU SHOT WITHIN PAST YEAR

TABLE 20.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PNEUMONIA VACCINATION, AGE 65 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	424	46.1	2.6	( 41.1, 51.2)
ALASKA	156	56.2	5.5	( 45.4, 67.0)
ARIZONA	460	46.6	3.4	( 40.0, 53.2)
ARKANSAS	570	49.8	2.3	( 45.3, 54.4)
CALIFORNIA	597	43.0	2.4	( 38.4, 47.6)
COLORADO	472	37.3	3.1	( 31.1, 43.4)
CONNECTICUT	598	51.0	2.7	( 45.8, 56.3)
DELAWARE	440	33.5	2.8	( 27.9, 39.0)
DISTRICT OF CO	240	64.7	3.3	( 58.3, 71.1)
FLORIDA	1207	46.6	1.7	( 43.2, 50.0)
GEORGIA	311	50.3	3.3	( 43.8, 56.7)
HAWAII	388	44.2	3.5	( 37.4, 51.0)
IDAHO	876	44.8	2.0	( 41.0, 48.7)
ILLINOIS	263	52.6	3.4	( 45.9, 59.4)
INDIANA	390	48.4	4.2	( 40.2, 56.5)
IOWA	864	38.8	1.9	( 35.0, 42.5)
KANSAS	845	44.9	1.9	( 41.2, 48.7)
KENTUCKY	1680	48.0	1.7	( 44.7, 51.3)
LOUISIANA	312	59.6	3.1	( 53.6, 65.7)
MAINE	348	42.7	3.0	( 36.9, 48.6)
MARYLAND	598	45.9	2.6	( 40.8, 50.9)
MASSACHUSETTS	837	43.2	2.1	( 39.2, 47.3)
MICHIGAN	453	42.3	2.5	( 37.3, 47.2)
MINNESOTA	892	48.1	1.9	( 44.5, 51.8)
MISSISSIPPI	371	49.6	2.8	( 44.1, 55.2)
MISSOURI	935	47.2	2.3	( 42.8, 51.6)
MONTANA	380	38.8	2.8	( 33.4, 44.2)
NEBRASKA	707	45.2	2.0	( 41.2, 49.1)
NEVADA	332	38.3	4.3	( 29.9, 46.6)
NEW HAMPSHIRE	210	39.7	3.7	( 32.4, 46.9)
NEW JERSEY	518	44.9	2.5	( 40.0, 49.8)
NEW MEXICO	606	46.8	2.3	( 42.2, 51.3)
NEW YORK	418	50.0	2.7	( 44.8, 55.3)
NORTH CAROLINA	513	41.5	2.4	( 36.7, 46.2)
NORTH DAKOTA	464	45.0	2.5	( 40.1, 49.9)
OHIO	389	45.0	2.9	( 39.3, 50.7)
OKLAHOMA	646	46.3	2.2	( 42.1, 50.5)
OREGON	343	43.8	2.9	( 38.1, 49.5)
PENNSYLVANIA	681	47.8	2.1	( 43.6, 51.9)
RHODE ISLAND	764	43.1	2.0	( 39.1, 47.0)
SOUTH CAROLINA	511	43.9	2.4	( 39.2, 48.6)
SOUTH DAKOTA	930	49.6	1.8	( 46.1, 53.1)
TENNESSEE	570	45.7	2.4	( 41.0, 50.4)
TEXAS	680	44.1	2.2	( 39.8, 48.5)
UTAH	540	38.7	2.8	( 33.3, 44.1)
VERMONT	581	43.5	2.3	( 39.1, 48.0)
VIRGINIA	566	44.8	2.8	( 39.2, 50.3)
WASHINGTON	596	44.2	2.3	( 39.7, 48.6)
WEST VIRGINIA	609	45.7	2.2	( 41.4, 50.0)
WISCONSIN	384	46.3	2.8	( 41.0, 51.7)
WYOMING	395	38.5	2.7	( 33.3, 43.7)
PUERTO RICO	629	78.2	1.8	( 74.7, 81.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 45.1 RANGE = 33.5-78.2 YEAR 2000 OBJ. 20.11  
 \* NEVER HAD PNEUMONIA VACCINATION June 20, 2000



TABLE 20.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 NO PNEUMONIA VACCINATION, AGE 65 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	52.6	46.1	-6.4
ALASKA	60.8	56.2	-4.5
ARIZONA	40.6	46.6	6.0
ARKANSAS	61.0	49.8	-11.1
CALIFORNIA	50.2	43.0	-7.2
COLORADO	46.7	37.3	-9.4
CONNECTICUT	57.0	51.0	-5.9
DELAWARE	47.4	33.5	-13.9
DISTRICT OF CO	67.7	64.7	-3.0
FLORIDA	54.5	46.6	-8.0
GEORGIA	51.5	50.3	-1.2
HAWAII	48.3	44.2	-4.0
IDAHO	49.8	44.8	-5.0
ILLINOIS	55.3	52.6	-2.7
INDIANA	62.0	48.4	-13.6
IOWA	48.5	38.8	-9.8
KANSAS	56.3	44.9	-11.4
KENTUCKY	61.4	48.0	-13.4
LOUISIANA	67.8	59.6	-8.1
MAINE	50.0	42.7	-7.3
MARYLAND	59.0	45.9	-13.1
MASSACHUSETTS	47.3	43.2	-4.0
MICHIGAN	54.4	42.3	-12.1
MINNESOTA	51.7	48.1	-3.6
MISSISSIPPI	54.1	49.6	-4.5
MISSOURI	55.7	47.2	-8.5
MONTANA	49.2	38.8	-10.3
NEBRASKA	50.2	45.2	-5.0
NEVADA	46.5	38.3	-8.2
NEW HAMPSHIRE	50.4	39.7	-10.7
NEW JERSEY	66.1	44.9	-21.1
NEW MEXICO	49.9	46.8	-3.1
NEW YORK	61.1	50.0	-11.0
NORTH CAROLINA	49.4	41.5	-7.9
NORTH DAKOTA	59.2	45.0	-14.2
OHIO	61.5	45.0	-16.4
OKLAHOMA	59.6	46.3	-13.3
OREGON	44.1	43.8	-0.3
PENNSYLVANIA	53.0	47.8	-5.2
RHODE ISLAND	57.0	43.1	-13.9
SOUTH CAROLINA	58.4	43.9	-14.5
SOUTH DAKOTA	59.4	49.6	-9.7
TENNESSEE	55.0	45.7	-9.3
TEXAS	55.6	44.1	-11.4
UTAH	51.5	38.7	-12.8
VERMONT	48.4	43.5	-4.8
VIRGINIA	46.4	44.8	-1.6
WASHINGTON	48.4	44.2	-4.3
WEST VIRGINIA	58.7	45.7	-13.1
WISCONSIN	57.4	46.3	-11.1
WYOMING	49.2	38.5	-10.6
PUERTO RICO	66.3	78.2	11.9

\* NEVER HAD PNEUMONIA VACCINATION

June 20, 2000

TABLE 21.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO SIGMOIDSCOPY, AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	928	58.8	1.7	( 55.4, 62.2)
ALASKA	557	49.6	3.3	( 43.1, 56.1)
ARIZONA	822	63.2	2.4	( 58.5, 67.8)
ARKANSAS	1257	57.9	1.6	( 54.9, 61.0)
CALIFORNIA	1438	51.8	1.5	( 48.8, 54.8)
COLORADO	1112	55.8	2.2	( 51.5, 60.2)
CONNECTICUT	1288	51.3	1.9	( 47.6, 54.9)
DELAWARE	987	44.0	2.1	( 39.9, 48.1)
DISTRICT OF CO	502	51.6	2.4	( 46.9, 56.4)
FLORIDA	2365	55.0	1.3	( 52.4, 57.5)
GEORGIA	779	52.6	2.0	( 48.6, 56.6)
HAWAII	849	55.8	2.3	( 51.3, 60.3)
IDAHO	1922	58.7	1.4	( 56.0, 61.3)
ILLINOIS	530	56.1	2.4	( 51.3, 60.9)
INDIANA	871	56.6	2.8	( 51.1, 62.1)
IOWA	1606	53.6	1.5	( 50.7, 56.4)
KANSAS	1594	61.2	1.4	( 58.6, 63.9)
KENTUCKY	3398	65.3	1.2	( 63.0, 67.6)
LOUISIANA	631	63.3	2.1	( 59.2, 67.5)
MAINE	755	57.6	2.0	( 53.7, 61.5)
MARYLAND	1417	49.5	1.7	( 46.2, 52.7)
MASSACHUSETTS	1775	56.1	1.5	( 53.2, 58.9)
MICHIGAN	960	50.5	1.8	( 47.1, 54.0)
MINNESOTA	1911	52.5	1.2	( 50.1, 55.0)
MISSISSIPPI	830	62.6	1.9	( 58.9, 66.2)
MISSOURI	1833	62.0	1.5	( 59.0, 64.9)
MONTANA	798	56.8	2.0	( 53.0, 60.6)
NEBRASKA	1232	66.3	1.5	( 63.4, 69.2)
NEVADA	785	56.9	2.7	( 51.6, 62.3)
NEW HAMPSHIRE	444	53.2	2.6	( 48.0, 58.3)
NEW JERSEY	1108	56.0	1.7	( 52.6, 59.4)
NEW MEXICO	1378	56.9	1.5	( 54.0, 59.8)
NEW YORK	953	57.8	1.8	( 54.2, 61.3)
NORTH CAROLINA	1042	60.9	1.7	( 57.5, 64.3)
NORTH DAKOTA	856	58.0	1.8	( 54.4, 61.5)
OHIO	747	59.3	2.2	( 55.0, 63.5)
OKLAHOMA	1296	62.3	1.5	( 59.4, 65.2)
OREGON	729	52.9	2.0	( 48.9, 56.9)
PENNSYLVANIA	1381	60.3	1.5	( 57.4, 63.2)
RHODE ISLAND	1518	51.3	1.5	( 48.4, 54.2)
SOUTH CAROLINA	1258	58.2	1.5	( 55.2, 61.2)
SOUTH DAKOTA	1726	54.5	1.3	( 52.0, 57.1)
TENNESSEE	1189	60.7	1.6	( 57.5, 63.9)
TEXAS	1708	55.7	1.4	( 52.9, 58.5)
UTAH	1129	55.9	2.0	( 51.9, 59.9)
VERMONT	1270	57.8	1.5	( 54.7, 60.8)
VIRGINIA	1334	55.8	2.0	( 52.0, 59.7)
WASHINGTON	1256	49.2	1.7	( 45.9, 52.6)
WEST VIRGINIA	1210	65.1	1.5	( 62.2, 68.1)
WISCONSIN	814	47.6	1.9	( 43.8, 51.4)
WYOMING	879	53.5	1.8	( 49.9, 57.0)
PUERTO RICO	1381	71.8	1.4	( 69.1, 74.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 56.4 RANGE = 44.0-71.8 YEAR 2000 OBJ. 16.13  
 \* NEVER HAD SIGMOIDSCOPY June 30, 2000

TABLE 21.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO SIGMOIDSCOPY, AGE 50 AND OVER\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	361	58.6	2.8	( 53.2, 64.0)	*	567	59.0	2.2	( 54.6, 63.3)	*
ALASKA	279	45.6	4.6	( 36.6, 54.5)	*	278	53.8	4.6	( 44.8, 62.8)	*
ARIZONA	326	62.1	3.8	( 54.6, 69.6)	*	496	64.0	3.0	( 58.0, 69.9)	*
ARKANSAS	446	57.9	2.6	( 52.9, 62.9)	*	811	58.0	1.9	( 54.2, 61.7)	*
CALIFORNIA	594	47.1	2.4	( 42.5, 51.8)	*	844	55.9	1.9	( 52.1, 59.7)	*
COLORADO	472	52.3	3.6	( 45.4, 59.3)	*	640	58.9	2.7	( 53.5, 64.3)	*
CONNECTICUT	474	46.4	3.0	( 40.5, 52.2)	*	814	54.9	2.3	( 50.4, 59.5)	*
DELAWARE	395	39.6	3.3	( 33.1, 46.0)	*	592	47.3	2.7	( 42.1, 52.6)	*
DISTRICT OF CO	184	44.0	4.1	( 36.0, 51.9)	*	318	57.3	2.9	( 51.6, 63.1)	*
FLORIDA	909	53.3	2.1	( 49.3, 57.3)	*	1456	56.3	1.6	( 53.3, 59.3)	*
GEORGIA	259	48.7	3.5	( 41.9, 55.5)	*	520	55.7	2.4	( 51.1, 60.4)	*
HAWAII	387	53.0	3.4	( 46.4, 59.5)	*	462	58.4	3.2	( 52.1, 64.7)	*
IDAHO	796	58.3	2.1	( 54.1, 62.5)	*	1126	59.0	1.7	( 55.6, 62.4)	*
ILLINOIS	193	60.8	4.0	( 53.0, 68.5)	*	337	52.2	3.0	( 46.4, 58.1)	*
INDIANA	334	54.4	4.5	( 45.5, 63.2)	*	537	58.4	3.5	( 51.6, 65.2)	*
IOWA	591	54.8	2.3	( 50.2, 59.3)	*	1015	52.7	1.8	( 49.1, 56.2)	*
KANSAS	610	60.0	2.2	( 55.7, 64.3)	*	984	62.2	1.7	( 58.9, 65.5)	*
KENTUCKY	1138	66.4	2.0	( 62.4, 70.4)	*	2260	64.4	1.3	( 61.9, 67.0)	*
LOUISIANA	234	60.0	3.5	( 53.1, 66.9)	*	397	65.9	2.6	( 60.9, 70.9)	*
MAINE	323	60.3	3.0	( 54.4, 66.3)	*	432	55.5	2.6	( 50.3, 60.6)	*
MARYLAND	560	44.8	2.6	( 39.6, 50.0)	*	857	53.4	2.1	( 49.3, 57.5)	*
MASSACHUSETTS	646	52.5	2.4	( 47.7, 57.2)	*	1129	58.8	1.8	( 55.3, 62.4)	*
MICHIGAN	372	42.4	2.8	( 37.0, 47.9)	*	588	56.8	2.2	( 52.4, 61.2)	*
MINNESOTA	817	53.3	1.9	( 49.6, 57.0)	*	1094	51.8	1.6	( 48.7, 55.0)	*
MISSISSIPPI	303	61.8	3.1	( 55.8, 67.9)	*	527	63.1	2.3	( 58.6, 67.6)	*
MISSOURI	694	63.2	2.4	( 58.5, 67.9)	*	1139	61.0	1.9	( 57.2, 64.8)	*
MONTANA	319	56.7	3.1	( 50.6, 62.9)	*	479	56.9	2.4	( 52.1, 61.6)	*
NEBRASKA	440	66.2	2.4	( 61.6, 70.9)	*	792	66.4	1.8	( 62.8, 69.9)	*
NEVADA	389	57.9	3.4	( 51.2, 64.5)	*	396	56.1	4.2	( 47.9, 64.3)	*
NEW HAMPSHIRE	176	45.1	4.1	( 37.1, 53.2)	*	268	59.6	3.3	( 53.1, 66.1)	*
NEW JERSEY	422	49.8	2.8	( 44.2, 55.3)	*	686	60.8	2.1	( 56.7, 65.0)	*
NEW MEXICO	575	54.6	2.3	( 50.2, 59.1)	*	803	58.7	2.0	( 54.9, 62.6)	*
NEW YORK	371	54.1	2.9	( 48.4, 59.7)	*	582	60.7	2.3	( 56.2, 65.1)	*
NORTH CAROLINA	390	60.6	2.8	( 55.1, 66.2)	*	652	61.1	2.1	( 57.0, 65.3)	*
NORTH DAKOTA	329	55.0	2.9	( 49.2, 60.7)	*	527	60.4	2.3	( 55.9, 64.9)	*
OHIO	251	57.4	3.8	( 50.0, 64.8)	*	496	60.6	2.5	( 55.7, 65.5)	*
OKLAHOMA	462	59.3	2.5	( 54.4, 64.1)	*	834	64.6	1.8	( 61.1, 68.1)	*
OREGON	276	52.3	3.3	( 45.9, 58.7)	*	453	53.4	2.5	( 48.4, 58.3)	*
PENNSYLVANIA	517	58.0	2.4	( 53.4, 62.6)	*	864	62.1	1.9	( 58.5, 65.8)	*
RHODE ISLAND	555	48.6	2.4	( 44.0, 53.2)	*	963	53.4	1.9	( 49.7, 57.0)	*
SOUTH CAROLINA	494	57.2	2.4	( 52.4, 61.9)	*	764	59.1	2.0	( 55.2, 62.9)	*
SOUTH DAKOTA	659	56.5	2.1	( 52.4, 60.5)	*	1067	52.9	1.7	( 49.7, 56.2)	*
TENNESSEE	367	57.9	2.8	( 52.4, 63.5)	*	822	62.9	1.8	( 59.3, 66.4)	*
TEXAS	622	51.7	2.3	( 47.3, 56.1)	*	1086	58.9	1.8	( 55.4, 62.5)	*
UTAH	475	55.1	3.2	( 49.0, 61.3)	*	654	56.5	2.6	( 51.4, 61.6)	*
VERMONT	465	57.0	2.5	( 52.1, 61.9)	*	805	58.4	1.9	( 54.6, 62.1)	*
VIRGINIA	549	51.4	3.0	( 45.5, 57.3)	*	785	59.6	2.5	( 54.7, 64.5)	*
WASHINGTON	526	46.5	2.7	( 41.1, 51.8)	*	730	51.6	2.1	( 47.4, 55.7)	*
WEST VIRGINIA	455	62.6	2.5	( 57.7, 67.5)	*	755	67.1	1.9	( 63.4, 70.7)	*
WISCONSIN	324	43.8	3.0	( 38.0, 49.6)	*	490	50.7	2.5	( 45.8, 55.5)	*
WYOMING	374	47.7	2.8	( 42.2, 53.2)	*	505	58.6	2.4	( 54.0, 63.2)	*
PUERTO RICO	490	68.7	2.3	( 64.3, 73.2)	*	891	74.1	1.6	( 70.9, 77.3)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 52  
 MEDIAN = 54.9  
 RANGE = 39.6-68.7

NUMBER OF PARTICIPANTS = 52  
 MEDIAN = 58.7  
 RANGE = 47.3-74.1

\* NEVER HAD SIGMOIDSCOPY

June 30, 2000

TABLE 22.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HOME BLOOD STOOL TEST, AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	921	82.4	1.3	( 79.7, 85.0)
ALASKA	552	82.7	2.6	( 77.6, 87.8)
ARIZONA	822	67.1	2.4	( 62.4, 71.8)
ARKANSAS	1256	77.1	1.3	( 74.5, 79.7)
CALIFORNIA	1437	77.0	1.2	( 74.6, 79.4)
COLORADO	1108	70.3	2.0	( 66.5, 74.2)
CONNECTICUT	1269	63.2	1.8	( 59.7, 66.8)
DELAWARE	988	71.0	1.9	( 67.3, 74.8)
DISTRICT OF CO	500	57.0	2.4	( 52.3, 61.7)
FLORIDA	2340	67.6	1.2	( 65.3, 69.8)
GEORGIA	779	78.3	1.6	( 75.1, 81.4)
HAWAII	845	71.3	2.1	( 67.2, 75.3)
IDAHO	1920	76.9	1.2	( 74.6, 79.1)
ILLINOIS	529	74.2	2.1	( 70.0, 78.3)
INDIANA	871	77.6	2.4	( 72.8, 82.3)
IOWA	1585	67.0	1.4	( 64.3, 69.6)
KANSAS	1595	73.9	1.2	( 71.6, 76.2)
KENTUCKY	3374	76.1	1.0	( 74.1, 78.0)
LOUISIANA	627	76.2	1.8	( 72.6, 79.8)
MAINE	758	64.1	1.9	( 60.3, 67.8)
MARYLAND	1401	62.2	1.6	( 59.0, 65.3)
MASSACHUSETTS	1754	64.8	1.4	( 62.0, 67.5)
MICHIGAN	962	66.9	1.6	( 63.7, 70.1)
MINNESOTA	1897	73.8	1.1	( 71.7, 75.9)
MISSISSIPPI	827	82.2	1.5	( 79.3, 85.1)
MISSOURI	1828	75.0	1.4	( 72.3, 77.7)
MONTANA	792	73.6	1.8	( 70.2, 77.1)
NEBRASKA	1217	72.6	1.4	( 69.9, 75.3)
NEVADA	782	80.0	2.0	( 76.0, 83.9)
NEW HAMPSHIRE	440	59.3	2.6	( 54.2, 64.4)
NEW JERSEY	1102	64.6	1.7	( 61.3, 68.0)
NEW MEXICO	1367	76.1	1.3	( 73.6, 78.6)
NEW YORK	936	70.8	1.7	( 67.6, 74.1)
NORTH CAROLINA	1034	61.5	1.7	( 58.1, 64.9)
NORTH DAKOTA	858	75.5	1.6	( 72.4, 78.6)
OHIO	744	69.1	2.0	( 65.3, 72.9)
OKLAHOMA	1294	76.7	1.3	( 74.1, 79.2)
OREGON	721	67.8	1.9	( 64.1, 71.5)
PENNSYLVANIA	1378	69.0	1.4	( 66.3, 71.7)
RHODE ISLAND	1496	68.6	1.4	( 65.9, 71.3)
SOUTH CAROLINA	1237	73.6	1.4	( 70.9, 76.3)
SOUTH DAKOTA	1693	74.8	1.1	( 72.6, 77.1)
TENNESSEE	1189	76.7	1.4	( 74.0, 79.4)
TEXAS	1701	77.0	1.2	( 74.6, 79.4)
UTAH	1123	80.2	1.6	( 77.0, 83.4)
VERMONT	1247	62.6	1.5	( 59.7, 65.6)
VIRGINIA	1338	73.8	1.7	( 70.4, 77.1)
WASHINGTON	1256	64.3	1.7	( 61.0, 67.6)
WEST VIRGINIA	1201	80.9	1.2	( 78.5, 83.3)
WISCONSIN	813	78.7	1.5	( 75.7, 81.6)
WYOMING	882	81.6	1.4	( 78.8, 84.4)
PUERTO RICO	1366	88.2	1.0	( 86.2, 90.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 73.8 RANGE = 57.0-88.2  
 \* DID NOT USE HOME BLOOD STOOL KIT WITHIN 2 YEARS June 30, 2000

TABLE 22.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HOME BLOOD STOOL TEST, AGE 50 AND OVER\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	363	85.2	1.9	( 81.4, 88.9)	*	558	80.1	1.9	( 76.5, 83.8)	*
ALASKA	278	85.9	3.1	( 79.9, 91.8)	*	274	79.4	4.2	( 71.2, 87.6)	*
ARIZONA	327	69.2	3.9	( 61.6, 76.8)	*	495	65.5	3.0	( 59.5, 71.4)	*
ARKANSAS	446	77.6	2.2	( 73.3, 82.0)	*	810	76.7	1.6	( 73.5, 79.8)	*
CALIFORNIA	596	79.3	1.9	( 75.6, 83.0)	*	841	75.0	1.6	( 71.8, 78.2)	*
COLORADO	473	78.8	2.7	( 73.5, 84.2)	*	635	62.8	2.7	( 57.5, 68.1)	*
CONNECTICUT	467	68.7	2.9	( 63.1, 74.3)	*	802	59.2	2.3	( 54.6, 63.8)	*
DELAWARE	399	74.2	2.8	( 68.6, 79.7)	*	589	68.6	2.6	( 63.6, 73.6)	*
DISTRICT OF CO	185	57.3	3.9	( 49.6, 64.9)	*	315	56.8	3.0	( 51.0, 62.6)	*
FLORIDA	898	71.2	1.9	( 67.5, 74.9)	*	1442	64.6	1.5	( 61.6, 67.6)	*
GEORGIA	260	83.7	2.4	( 79.1, 88.4)	*	519	73.8	2.1	( 69.7, 78.0)	*
HAWAII	387	74.5	2.8	( 69.0, 80.0)	*	458	68.3	3.0	( 62.5, 74.2)	*
IDAHO	791	82.1	1.7	( 78.9, 85.4)	*	1129	72.2	1.6	( 69.2, 75.3)	*
ILLINOIS	194	82.2	3.0	( 76.3, 88.2)	*	335	67.4	2.8	( 61.9, 72.9)	*
INDIANA	332	77.9	4.0	( 70.1, 85.7)	*	539	77.3	3.0	( 71.5, 83.1)	*
IOWA	589	73.7	2.0	( 69.7, 77.7)	*	996	61.7	1.8	( 58.2, 65.2)	*
KANSAS	612	79.9	1.7	( 76.6, 83.3)	*	983	69.0	1.6	( 65.8, 72.1)	*
KENTUCKY	1132	77.1	1.8	( 73.6, 80.6)	*	2242	75.2	1.2	( 72.9, 77.5)	*
LOUISIANA	233	77.5	2.9	( 71.7, 83.2)	*	394	75.2	2.4	( 70.6, 79.9)	*
MAINE	325	72.9	2.6	( 67.8, 78.0)	*	433	57.0	2.6	( 51.9, 62.1)	*
MARYLAND	555	63.0	2.6	( 57.9, 68.0)	*	846	61.5	2.1	( 57.5, 65.5)	*
MASSACHUSETTS	644	68.4	2.3	( 64.0, 72.8)	*	1110	62.0	1.8	( 58.5, 65.5)	*
MICHIGAN	377	66.1	2.6	( 60.9, 71.2)	*	585	67.5	2.1	( 63.4, 71.6)	*
MINNESOTA	817	76.8	1.6	( 73.7, 79.9)	*	1080	71.3	1.4	( 68.5, 74.1)	*
MISSISSIPPI	302	84.2	2.4	( 79.5, 88.8)	*	525	80.7	1.9	( 77.0, 84.4)	*
MISSOURI	695	78.3	2.1	( 74.1, 82.4)	*	1133	72.3	1.8	( 68.8, 75.8)	*
MONTANA	316	77.6	2.8	( 72.1, 83.0)	*	476	70.2	2.3	( 65.7, 74.6)	*
NEBRASKA	441	75.8	2.2	( 71.6, 80.0)	*	776	70.1	1.8	( 66.6, 73.5)	*
NEVADA	387	79.7	2.7	( 74.5, 85.0)	*	395	80.2	2.9	( 74.4, 85.9)	*
NEW HAMPSHIRE	178	60.5	4.1	( 52.5, 68.5)	*	262	58.3	3.4	( 51.7, 64.9)	*
NEW JERSEY	423	66.5	2.7	( 61.2, 71.7)	*	679	63.2	2.2	( 58.9, 67.5)	*
NEW MEXICO	573	76.3	1.9	( 72.5, 80.0)	*	794	76.0	1.7	( 72.6, 79.3)	*
NEW YORK	366	72.5	2.6	( 67.4, 77.5)	*	570	69.5	2.2	( 65.3, 73.7)	*
NORTH CAROLINA	388	64.8	2.8	( 59.2, 70.3)	*	646	58.8	2.1	( 54.6, 62.9)	*
NORTH DAKOTA	330	79.8	2.4	( 75.0, 84.5)	*	528	72.0	2.1	( 67.9, 76.1)	*
OHIO	248	72.0	3.2	( 65.7, 78.4)	*	496	67.0	2.4	( 62.2, 71.7)	*
OKLAHOMA	459	75.1	2.2	( 70.9, 79.4)	*	835	77.9	1.6	( 74.8, 80.9)	*
OREGON	275	75.9	2.8	( 70.4, 81.3)	*	446	61.0	2.5	( 56.1, 66.0)	*
PENNSYLVANIA	516	69.6	2.2	( 65.3, 73.9)	*	862	68.5	1.8	( 65.1, 72.0)	*
RHODE ISLAND	552	72.2	2.1	( 68.2, 76.3)	*	944	65.8	1.8	( 62.3, 69.3)	*
SOUTH CAROLINA	490	77.4	2.1	( 73.3, 81.4)	*	747	70.6	1.9	( 67.0, 74.2)	*
SOUTH DAKOTA	653	77.1	1.7	( 73.7, 80.5)	*	1040	73.0	1.5	( 70.0, 75.9)	*
TENNESSEE	367	78.6	2.3	( 74.0, 83.1)	*	822	75.3	1.7	( 71.9, 78.6)	*
TEXAS	621	78.4	1.9	( 74.7, 82.0)	*	1080	75.9	1.6	( 72.7, 79.0)	*
UTAH	473	81.7	2.4	( 77.0, 86.4)	*	650	78.9	2.2	( 74.6, 83.2)	*
VERMONT	461	69.8	2.3	( 65.3, 74.2)	*	786	56.5	1.9	( 52.7, 60.2)	*
VIRGINIA	551	78.2	2.5	( 73.3, 83.2)	*	787	70.0	2.3	( 65.5, 74.4)	*
WASHINGTON	527	67.8	2.6	( 62.6, 73.0)	*	729	61.4	2.1	( 57.2, 65.6)	*
WEST VIRGINIA	454	81.2	1.9	( 77.5, 84.9)	*	747	80.6	1.6	( 77.5, 83.7)	*
WISCONSIN	330	80.7	2.3	( 76.1, 85.3)	*	483	77.0	2.0	( 73.0, 80.9)	*
WYOMING	375	83.2	2.1	( 79.0, 87.3)	*	507	80.2	1.9	( 76.5, 83.9)	*
PUERTO RICO	488	87.7	1.7	( 84.4, 91.0)	*	878	88.5	1.2	( 86.2, 90.8)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 52  
 MEDIAN = 77.1  
 RANGE = 57.3-87.7

NUMBER OF PARTICIPANTS = 52  
 MEDIAN = 70.1  
 RANGE = 56.5-88.5

\* DID NOT USE HOME BLOOD STOOL KIT WITHIN 2 YEARS

June 30, 2000

TABLE 22.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1997 TO 1999 NO HOME BLOOD STOOL TEST, AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1997 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	80.8	82.4	1.6
ALASKA	75.0	82.7	7.7
ARIZONA	70.7	67.1	-3.6
ARKANSAS	78.7	77.1	-1.6
CALIFORNIA	75.2	77.0	1.8
COLORADO	67.8	70.3	2.6
CONNECTICUT	68.9	63.2	-5.7
DELAWARE	69.5	71.0	1.5
DISTRICT OF CO	62.2	57.0	-5.2
FLORIDA	69.9	67.6	-2.4
GEORGIA	72.2	78.3	6.1
HAWAII	67.8	71.3	3.5
IDAHO	74.6	76.9	2.3
ILLINOIS	75.1	74.2	-1.0
INDIANA	77.8	77.6	-0.3
IOWA	72.0	67.0	-5.0
KANSAS	68.5	73.9	5.4
KENTUCKY	73.9	76.1	2.2
LOUISIANA	76.7	76.2	-0.5
MAINE	65.0	64.1	-0.9
MARYLAND	67.9	62.2	-5.8
MASSACHUSETTS	65.8	64.8	-1.0
MICHIGAN	69.0	66.9	-2.1
MINNESOTA	67.7	73.8	6.1
MISSISSIPPI	86.5	82.2	-4.4
MISSOURI	74.9	75.0	0.0
MONTANA	76.0	73.6	-2.3
NEBRASKA	75.6	72.6	-3.0
NEVADA	84.0	80.0	-4.1
NEW HAMPSHIRE	66.1	59.3	-6.8
NEW JERSEY	71.5	64.6	-6.9
NEW MEXICO	78.3	76.1	-2.2
NEW YORK	68.1	70.8	2.7
NORTH CAROLINA	63.5	61.5	-2.0
NORTH DAKOTA	77.3	75.5	-1.8
OHIO	70.6	69.1	-1.5
OKLAHOMA	85.7	76.7	-9.1
OREGON	65.4	67.8	2.4
PENNSYLVANIA	71.2	69.0	-2.1
RHODE ISLAND	73.5	68.6	-4.9
SOUTH CAROLINA	76.0	73.6	-2.4
SOUTH DAKOTA	78.3	74.8	-3.5
TENNESSEE	77.7	76.7	-1.0
TEXAS	75.0	77.0	2.0
UTAH	78.0	80.2	2.2
VERMONT	64.9	62.6	-2.3
VIRGINIA	74.1	73.8	-0.3
WASHINGTON	64.1	64.3	0.2
WEST VIRGINIA	82.0	80.9	-1.1
WISCONSIN	74.6	78.7	4.1
WYOMING	80.7	81.6	0.9
PUERTO RICO	77.1	88.2	11.1

\* DID NOT USE HOME BLOOD STOOL KIT WITHIN 2 YEARS

June 30, 2000

TABLE 23.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY BODY MASS INDEX\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2044	37.5	1.2	( 35.1, 39.9)
ALASKA	2006	37.7	1.6	( 34.6, 40.8)
ARIZONA	1675	22.9	1.4	( 20.1, 25.7)
ARKANSAS	2907	38.1	1.0	( 36.1, 40.0)
CALIFORNIA	4152	33.5	0.9	( 31.9, 35.2)
COLORADO	2986	26.7	1.1	( 24.4, 28.9)
CONNECTICUT	3319	28.6	1.0	( 26.6, 30.7)
DELAWARE	2460	32.0	1.3	( 29.5, 34.5)
DISTRICT OF CO	1242	30.2	1.5	( 27.3, 33.0)
FLORIDA	5043	33.6	0.8	( 32.0, 35.1)
GEORGIA	2202	35.9	1.2	( 33.5, 38.3)
HAWAII	2110	28.8	1.4	( 26.1, 31.4)
IDAHO	4809	33.5	0.8	( 31.9, 35.1)
ILLINOIS	2805	36.0	1.1	( 33.9, 38.1)
INDIANA	2369	34.1	1.6	( 31.0, 37.3)
IOWA	3528	36.6	1.0	( 34.7, 38.4)
KANSAS	3680	33.4	0.9	( 31.6, 35.1)
KENTUCKY	7315	36.7	0.8	( 35.1, 38.3)
LOUISIANA	1617	37.7	1.4	( 35.0, 40.4)
MAINE	1610	32.7	1.3	( 30.1, 35.3)
MARYLAND	3819	32.7	1.0	( 30.8, 34.5)
MASSACHUSETTS	4688	28.3	0.9	( 26.6, 29.9)
MICHIGAN	2445	36.8	1.1	( 34.7, 38.9)
MINNESOTA	5155	29.2	0.7	( 27.9, 30.5)
MISSISSIPPI	2078	41.8	1.3	( 39.3, 44.3)
MISSOURI	4161	35.5	1.0	( 33.5, 37.4)
MONTANA	1743	29.8	1.2	( 27.4, 32.2)
NEBRASKA	2699	35.7	1.0	( 33.7, 37.7)
NEVADA	2131	28.9	1.4	( 26.1, 31.7)
NEW HAMPSHIRE	1191	27.9	1.5	( 24.9, 30.8)
NEW JERSEY	2677	31.4	1.1	( 29.2, 33.5)
NEW MEXICO	3378	33.0	0.9	( 31.2, 34.7)
NEW YORK	2506	31.9	1.1	( 29.8, 33.9)
NORTH CAROLINA	2330	34.9	1.1	( 32.7, 37.1)
NORTH DAKOTA	1927	36.3	1.2	( 33.9, 38.6)
OHIO	1544	35.9	1.4	( 33.1, 38.8)
OKLAHOMA	2810	37.2	1.1	( 35.1, 39.3)
OREGON	1740	32.8	1.2	( 30.3, 35.2)
PENNSYLVANIA	3355	35.5	0.9	( 33.6, 37.3)
RHODE ISLAND	3802	32.3	0.9	( 30.6, 34.0)
SOUTH CAROLINA	3262	37.3	1.0	( 35.4, 39.2)
SOUTH DAKOTA	3999	36.1	0.9	( 34.4, 37.7)
TENNESSEE	2866	34.5	1.0	( 32.5, 36.5)
TEXAS	4733	36.1	0.9	( 34.4, 37.9)
UTAH	3091	30.0	1.1	( 27.7, 32.2)
VERMONT	3013	31.1	1.0	( 29.2, 33.0)
VIRGINIA	3372	35.0	1.2	( 32.7, 37.3)
WASHINGTON	3453	31.8	1.0	( 29.9, 33.7)
WEST VIRGINIA	2504	41.3	1.1	( 39.1, 43.4)
WISCONSIN	2116	33.8	1.1	( 31.6, 36.0)
WYOMING	2343	31.8	1.0	( 29.8, 33.8)
PUERTO RICO	2893	36.5	1.0	( 34.5, 38.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 33.7 RANGE = 22.9-41.8 YEAR 2000 OBJ. 2.3  
 \* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES  
 June 20, 2000



TABLE 23.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 OVERWEIGHT: BY BODY MASS INDEX\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	35.8	37.5	1.7
ALASKA	36.2	37.7	1.4
ARIZONA	22.5	22.9	0.4
ARKANSAS	33.4	38.1	4.6
CALIFORNIA	31.0	33.5	2.6
COLORADO	27.2	26.7	-0.5
CONNECTICUT	27.5	28.6	1.1
DELAWARE	32.8	32.0	-0.8
DISTRICT OF CO	31.8	30.2	-1.6
FLORIDA	32.3	33.6	1.3
GEORGIA	33.3	35.9	2.6
HAWAII	27.8	28.8	0.9
IDAHO	29.9	33.5	3.6
ILLINOIS	33.5	36.0	2.5
INDIANA	34.5	34.1	-0.4
IOWA	34.6	36.6	2.0
KANSAS	31.7	33.4	1.6
KENTUCKY	35.8	36.7	0.9
LOUISIANA	36.1	37.7	1.6
MAINE	31.2	32.7	1.5
MARYLAND	34.5	32.7	-1.8
MASSACHUSETTS	26.9	28.3	1.4
MICHIGAN	34.6	36.8	2.2
MINNESOTA	32.5	29.2	-3.3
MISSISSIPPI	37.5	41.8	4.3
MISSOURI	35.6	35.5	-0.1
MONTANA	29.1	29.8	0.7
NEBRASKA	32.3	35.7	3.4
NEVADA	27.0	28.9	1.9
NEW HAMPSHIRE	28.7	27.9	-0.8
NEW JERSEY	29.7	31.4	1.7
NEW MEXICO	29.9	33.0	3.1
NEW YORK	28.6	31.9	3.2
NORTH CAROLINA	32.7	34.9	2.2
NORTH DAKOTA	33.5	36.3	2.7
OHIO	33.6	35.9	2.3
OKLAHOMA	32.4	37.2	4.8
OREGON	32.8	32.8	-0.0
PENNSYLVANIA	34.8	35.5	0.7
RHODE ISLAND	30.3	32.3	2.0
SOUTH CAROLINA	34.6	37.3	2.8
SOUTH DAKOTA	30.5	36.1	5.6
TENNESSEE	32.4	34.5	2.1
TEXAS	34.9	36.1	1.2
UTAH	28.8	30.0	1.2
VERMONT	27.4	31.1	3.8
VIRGINIA	31.8	35.0	3.1
WASHINGTON	32.3	31.8	-0.5
WEST VIRGINIA	37.8	41.3	3.5
WISCONSIN	34.0	33.8	-0.2
WYOMING	28.1	31.8	3.8
PUERTO RICO	34.7	36.5	1.8

\* BODY MASS INDEX GE 27.8 FOR MALES AND 27.3 FOR FEMALES

TABLE 24.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR OVERWEIGHT: BY PERCENT OF MEDIAN\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2044	43.1	1.3	( 40.6, 45.5)
ALASKA	2004	42.0	1.6	( 38.8, 45.1)
ARIZONA	1672	26.8	1.5	( 23.9, 29.7)
ARKANSAS	2906	42.4	1.0	( 40.3, 44.4)
CALIFORNIA	4146	38.2	0.9	( 36.5, 39.9)
COLORADO	2986	31.4	1.2	( 29.0, 33.8)
CONNECTICUT	3315	33.5	1.1	( 31.4, 35.7)
DELAWARE	2457	36.5	1.3	( 33.9, 39.0)
DISTRICT OF CO	1238	34.3	1.5	( 31.4, 37.3)
FLORIDA	5042	38.5	0.8	( 36.9, 40.1)
GEORGIA	2202	40.3	1.2	( 37.8, 42.7)
HAWAII	2109	33.3	1.4	( 30.6, 36.1)
IDAHO	4799	38.7	0.9	( 37.0, 40.4)
ILLINOIS	2805	41.0	1.1	( 38.8, 43.1)
INDIANA	2363	39.8	1.7	( 36.5, 43.0)
IOWA	3524	41.5	1.0	( 39.6, 43.5)
KANSAS	3673	38.6	0.9	( 36.8, 40.4)
KENTUCKY	7306	42.2	0.8	( 40.5, 43.8)
LOUISIANA	1613	42.1	1.4	( 39.4, 44.9)
MAINE	1613	37.9	1.4	( 35.2, 40.5)
MARYLAND	3808	37.7	1.0	( 35.8, 39.7)
MASSACHUSETTS	4677	32.8	0.9	( 31.0, 34.5)
MICHIGAN	2443	41.8	1.1	( 39.6, 44.0)
MINNESOTA	5147	34.4	0.7	( 33.0, 35.8)
MISSISSIPPI	2077	46.5	1.3	( 43.9, 49.0)
MISSOURI	4158	40.5	1.1	( 38.4, 42.5)
MONTANA	1739	35.1	1.3	( 32.6, 37.5)
NEBRASKA	2697	41.1	1.1	( 39.0, 43.2)
NEVADA	2126	33.6	1.5	( 30.6, 36.5)
NEW HAMPSHIRE	1190	31.8	1.6	( 28.8, 34.9)
NEW JERSEY	2674	36.2	1.1	( 34.0, 38.5)
NEW MEXICO	3375	38.0	0.9	( 36.2, 39.9)
NEW YORK	2500	36.5	1.1	( 34.3, 38.6)
NORTH CAROLINA	2326	40.5	1.2	( 38.2, 42.8)
NORTH DAKOTA	1921	41.5	1.2	( 39.1, 43.9)
OHIO	1546	40.0	1.5	( 37.1, 42.9)
OKLAHOMA	2809	43.0	1.1	( 40.9, 45.1)
OREGON	1737	36.5	1.3	( 34.0, 39.0)
PENNSYLVANIA	3353	39.9	1.0	( 38.0, 41.7)
RHODE ISLAND	3797	37.7	0.9	( 35.8, 39.5)
SOUTH CAROLINA	3256	41.4	1.0	( 39.5, 43.4)
SOUTH DAKOTA	3994	41.7	0.9	( 39.9, 43.4)
TENNESSEE	2864	39.1	1.0	( 37.1, 41.2)
TEXAS	4726	41.2	0.9	( 39.4, 42.9)
UTAH	3089	35.1	1.2	( 32.7, 37.4)
VERMONT	3010	35.7	1.0	( 33.7, 37.6)
VIRGINIA	3368	40.2	1.2	( 37.7, 42.6)
WASHINGTON	3450	36.8	1.0	( 34.8, 38.8)
WEST VIRGINIA	2501	46.2	1.1	( 44.0, 48.5)
WISCONSIN	2114	38.6	1.2	( 36.3, 40.9)
WYOMING	2336	37.4	1.1	( 35.3, 39.6)
PUERTO RICO	2887	41.7	1.1	( 39.6, 43.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 52 MEDIAN = 38.7 RANGE = 26.8-46.5

\* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

June 20, 2000



TABLE 24.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 OVERWEIGHT: BY PERCENT OF MEDIAN\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	41.6	43.1	1.5
ALASKA	40.8	42.0	1.2
ARIZONA	26.6	26.8	0.2
ARKANSAS	38.3	42.4	4.0
CALIFORNIA	35.6	38.2	2.6
COLORADO	32.0	31.4	-0.7
CONNECTICUT	31.7	33.5	1.9
DELAWARE	36.4	36.5	0.1
DISTRICT OF CO	35.7	34.3	-1.4
FLORIDA	37.2	38.5	1.3
GEORGIA	37.1	40.3	3.2
HAWAII	31.2	33.3	2.1
IDAHO	34.9	38.7	3.8
ILLINOIS	38.2	41.0	2.8
INDIANA	38.5	39.8	1.3
IOWA	39.7	41.5	1.8
KANSAS	36.6	38.6	2.0
KENTUCKY	40.6	42.2	1.6
LOUISIANA	41.7	42.1	0.5
MAINE	36.0	37.9	1.8
MARYLAND	38.7	37.7	-1.0
MASSACHUSETTS	32.0	32.8	0.8
MICHIGAN	39.0	41.8	2.8
MINNESOTA	38.8	34.4	-4.4
MISSISSIPPI	42.2	46.5	4.2
MISSOURI	39.6	40.5	0.9
MONTANA	33.1	35.1	2.0
NEBRASKA	37.2	41.1	3.9
NEVADA	31.2	33.6	2.4
NEW HAMPSHIRE	34.0	31.8	-2.2
NEW JERSEY	35.2	36.2	1.0
NEW MEXICO	34.4	38.0	3.7
NEW YORK	33.2	36.5	3.3
NORTH CAROLINA	38.4	40.5	2.1
NORTH DAKOTA	38.9	41.5	2.6
OHIO	38.3	40.0	1.8
OKLAHOMA	36.6	43.0	6.4
OREGON	38.3	36.5	-1.8
PENNSYLVANIA	39.7	39.9	0.2
RHODE ISLAND	35.3	37.7	2.4
SOUTH CAROLINA	39.5	41.4	1.9
SOUTH DAKOTA	34.8	41.7	6.9
TENNESSEE	37.1	39.1	2.0
TEXAS	40.0	41.2	1.2
UTAH	33.6	35.1	1.5
VERMONT	32.2	35.7	3.5
VIRGINIA	36.2	40.2	4.0
WASHINGTON	37.6	36.8	-0.8
WEST VIRGINIA	43.0	46.2	3.2
WISCONSIN	39.2	38.6	-0.5
WYOMING	33.0	37.4	4.5
PUERTO RICO	39.8	41.7	1.9

\* WEIGHT FOR HEIGHT GREATER THAN 120% OF THE MEDIAN

June 20, 2000

TABLE 25.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1612	9.5	0.8	( 8.0, 11.1)
ALASKA	1812	10.3	1.1	( 8.1, 12.5)
ARIZONA	1241	6.3	0.8	( 4.7, 7.9)
ARKANSAS	2367	12.6	0.8	( 11.1, 14.0)
COLORADO	2500	8.0	0.8	( 6.5, 9.5)
CONNECTICUT	2751	8.9	0.8	( 7.4, 10.3)
DELAWARE	2051	7.6	0.7	( 6.2, 9.1)
DISTRICT OF CO	981	7.4	1.0	( 5.6, 9.3)
FLORIDA	3851	10.2	0.7	( 8.9, 11.6)
GEORGIA	1880	14.1	0.9	( 12.3, 15.9)
HAWAII	1686	9.6	0.9	( 7.9, 11.3)
IDAHO	3973	11.8	0.7	( 10.5, 13.1)
ILLINOIS	2380	12.2	0.9	( 10.4, 13.9)
INDIANA	1970	10.4	1.2	( 8.1, 12.8)
IOWA	2609	7.7	0.6	( 6.5, 8.9)
KANSAS	2848	12.4	0.7	( 11.0, 13.7)
KENTUCKY	5595	11.4	0.6	( 10.3, 12.6)
LOUISIANA	1311	13.1	1.0	( 11.0, 15.1)
MAINE	1266	8.0	0.8	( 6.3, 9.6)
MARYLAND	3252	7.9	0.6	( 6.7, 9.0)
MASSACHUSETTS	3894	8.6	0.6	( 7.4, 9.7)
MICHIGAN	1995	10.9	0.8	( 9.4, 12.5)
MINNESOTA	4352	28.9	0.8	( 27.5, 30.4)
MISSISSIPPI	1714	12.4	0.9	( 10.6, 14.1)
MISSOURI	3217	10.3	0.8	( 8.8, 11.8)
MONTANA	1367	15.5	1.1	( 13.5, 17.6)
NEBRASKA	1964	11.2	0.8	( 9.7, 12.6)
NEVADA	1809	10.2	1.0	( 8.3, 12.1)
NEW HAMPSHIRE	985	7.2	0.9	( 5.5, 8.9)
NEW JERSEY	2123	9.6	0.8	( 8.2, 11.1)
NEW MEXICO	2775	10.6	0.7	( 9.3, 11.9)
NEW YORK	2055	8.7	0.7	( 7.3, 10.1)
NORTH CAROLINA	1827	12.4	0.9	( 10.6, 14.2)
NORTH DAKOTA	1456	10.3	0.9	( 8.6, 12.0)
OHIO	1199	8.6	0.9	( 6.8, 10.5)
OKLAHOMA	2245	12.3	0.8	( 10.8, 13.9)
OREGON	1423	10.8	0.9	( 9.0, 12.6)
PENNSYLVANIA	2714	11.7	0.7	( 10.3, 13.1)
RHODE ISLAND	3047	9.9	0.6	( 8.7, 11.2)
SOUTH CAROLINA	2792	12.6	0.7	( 11.1, 14.0)
SOUTH DAKOTA	3065	9.8	0.6	( 8.6, 10.9)
TENNESSEE	2232	17.9	1.0	( 16.0, 19.7)
TEXAS	4167	10.5	0.6	( 9.4, 11.6)
UTAH	2551	11.6	0.9	( 9.9, 13.3)
VERMONT	2464	6.6	0.6	( 5.4, 7.8)
VIRGINIA	2817	10.5	1.0	( 8.6, 12.4)
WASHINGTON	2883	8.4	0.7	( 7.0, 9.7)
WEST VIRGINIA	1943	12.2	0.9	( 10.5, 13.9)
WISCONSIN	1730	8.5	0.8	( 7.0, 9.9)
WYOMING	1954	8.5	0.7	( 7.2, 9.9)
PUERTO RICO	2249	15.1	0.9	( 13.5, 16.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 10.3 RANGE = 6.3-28.9

\* RESPONDENT WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM June 20, 2000

TABLE 25.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64 YEARS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	676	9.5	1.2	( 7.2, 11.8)	*	936	9.6	1.1	( 7.5, 11.6)	*
ALASKA	887	12.2	1.8	( 8.6, 15.7)	*	925	8.2	1.2	( 5.9, 10.6)	*
ARIZONA	520	6.2	1.2	( 3.9, 8.5)	*	721	6.4	1.2	( 4.1, 8.8)	*
ARKANSAS	969	13.8	1.2	( 11.5, 16.2)	*	1398	11.4	0.9	( 9.6, 13.2)	*
COLORADO	1166	8.4	1.2	( 6.1, 10.6)	*	1334	7.6	1.0	( 5.6, 9.5)	*
CONNECTICUT	1100	9.1	1.2	( 6.9, 11.4)	*	1651	8.6	0.9	( 6.7, 10.4)	*
DELAWARE	823	8.7	1.3	( 6.2, 11.1)	*	1228	6.6	0.8	( 5.0, 8.2)	*
DISTRICT OF CO	363	8.7	1.6	( 5.6, 11.8)	*	618	6.3	1.1	( 4.1, 8.4)	*
FLORIDA	1549	10.8	1.1	( 8.6, 13.0)	*	2302	9.7	0.8	( 8.2, 11.2)	*
GEORGIA	760	15.1	1.5	( 12.2, 18.0)	*	1120	13.2	1.2	( 10.9, 15.4)	*
HAWAII	800	8.0	1.2	( 5.8, 10.3)	*	886	11.2	1.3	( 8.6, 13.8)	*
IDAHO	1708	12.5	1.1	( 10.5, 14.6)	*	2265	11.1	0.8	( 9.5, 12.7)	*
ILLINOIS	953	14.2	1.5	( 11.3, 17.1)	*	1427	10.1	0.9	( 8.3, 11.9)	*
INDIANA	854	12.1	1.9	( 8.4, 15.9)	*	1116	8.8	1.4	( 6.0, 11.5)	*
IOWA	1079	7.6	0.9	( 5.8, 9.4)	*	1530	7.8	0.8	( 6.3, 9.3)	*
KANSAS	1203	13.7	1.0	( 11.7, 15.7)	*	1645	11.0	0.8	( 9.3, 12.6)	*
KENTUCKY	2139	12.0	1.0	( 10.1, 13.9)	*	3456	10.9	0.7	( 9.6, 12.3)	*
LOUISIANA	539	14.7	1.7	( 11.4, 17.9)	*	772	11.5	1.3	( 9.0, 14.1)	*
MAINE	578	8.1	1.2	( 5.7, 10.5)	*	688	7.9	1.1	( 5.7, 10.1)	*
MARYLAND	1331	6.4	0.8	( 4.9, 8.0)	*	1921	9.2	0.8	( 7.6, 10.9)	*
MASSACHUSETTS	1605	9.2	1.0	( 7.3, 11.1)	*	2289	7.9	0.7	( 6.5, 9.4)	*
MICHIGAN	837	11.3	1.3	( 8.8, 13.8)	*	1158	10.6	1.0	( 8.6, 12.5)	*
MINNESOTA	2072	31.9	1.1	( 29.7, 34.1)	*	2280	26.0	1.0	( 24.0, 27.9)	*
MISSISSIPPI	702	13.3	1.4	( 10.5, 16.1)	*	1012	11.5	1.1	( 9.4, 13.6)	*
MISSOURI	1350	10.7	1.2	( 8.3, 13.1)	*	1867	9.9	1.0	( 8.0, 11.8)	*
MONTANA	613	17.8	1.7	( 14.5, 21.0)	*	754	13.3	1.3	( 10.8, 15.9)	*
NEBRASKA	831	9.9	1.1	( 7.8, 11.9)	*	1133	12.4	1.1	( 10.4, 14.5)	*
NEVADA	926	9.3	1.3	( 6.8, 11.8)	*	883	11.2	1.4	( 8.4, 13.9)	*
NEW HAMPSHIRE	404	6.5	1.2	( 4.1, 8.9)	*	581	7.8	1.2	( 5.5, 10.2)	*
NEW JERSEY	865	8.8	1.0	( 6.8, 10.8)	*	1258	10.5	1.1	( 8.3, 12.6)	*
NEW MEXICO	1250	12.5	1.1	( 10.4, 14.7)	*	1525	8.8	0.8	( 7.2, 10.3)	*
NEW YORK	858	9.3	1.1	( 7.0, 11.5)	*	1197	8.1	0.9	( 6.4, 9.8)	*
NORTH CAROLINA	766	13.1	1.4	( 10.4, 15.7)	*	1061	11.7	1.3	( 9.2, 14.2)	*
NORTH DAKOTA	678	11.7	1.3	( 9.1, 14.2)	*	778	8.9	1.1	( 6.7, 11.0)	*
OHIO	492	11.1	1.6	( 7.9, 14.2)	*	707	6.4	1.0	( 4.4, 8.3)	*
OKLAHOMA	919	13.4	1.2	( 11.0, 15.8)	*	1326	11.3	1.0	( 9.3, 13.2)	*
OREGON	629	11.8	1.4	( 9.0, 14.5)	*	794	9.8	1.2	( 7.5, 12.1)	*
PENNSYLVANIA	1112	13.1	1.1	( 10.9, 15.4)	*	1602	10.4	0.9	( 8.7, 12.0)	*
RHODE ISLAND	1234	10.9	1.0	( 8.9, 12.9)	*	1813	9.0	0.8	( 7.5, 10.5)	*
SOUTH CAROLINA	1153	12.6	1.1	( 10.5, 14.7)	*	1639	12.5	1.0	( 10.6, 14.4)	*
SOUTH DAKOTA	1275	10.2	0.9	( 8.3, 12.0)	*	1790	9.4	0.8	( 7.8, 10.9)	*
TENNESSEE	795	18.9	1.6	( 15.8, 22.1)	*	1437	16.8	1.1	( 14.7, 19.0)	*
TEXAS	1711	11.8	0.9	( 10.0, 13.5)	*	2456	9.2	0.7	( 7.8, 10.7)	*
UTAH	1121	10.8	1.2	( 8.4, 13.2)	*	1430	12.3	1.3	( 9.8, 14.8)	*
VERMONT	1047	6.1	0.9	( 4.3, 7.9)	*	1417	7.1	0.8	( 5.6, 8.6)	*
VIRGINIA	1161	10.3	1.5	( 7.3, 13.3)	*	1656	10.7	1.2	( 8.3, 13.0)	*
WASHINGTON	1313	8.8	1.1	( 6.6, 11.0)	*	1570	7.9	0.8	( 6.5, 9.4)	*
WEST VIRGINIA	844	14.0	1.3	( 11.5, 16.6)	*	1099	10.4	1.1	( 8.3, 12.6)	*
WISCONSIN	765	9.0	1.1	( 6.8, 11.2)	*	965	8.0	1.0	( 6.1, 9.9)	*
WYOMING	886	8.6	1.0	( 6.6, 10.5)	*	1068	8.5	0.9	( 6.7, 10.3)	*
PUERTO RICO	821	14.4	1.3	( 11.8, 16.9)	*	1428	15.9	1.1	( 13.7, 18.0)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	51	NUMBER OF PARTICIPANTS =	51
	MEDIAN	= 10.9	MEDIAN	= 9.8
	RANGE	= 6.1-31.9	RANGE	= 6.3-26.0

\* RESPONDENT WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

June 20, 2000

TABLE 25.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NOT ENCOURAGE TEEN CONDOM USE, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	9.4	9.5	0.1
ALASKA	10.2	10.3	0.1
ARIZONA	5.4	6.3	0.9
ARKANSAS	21.7	12.6	-9.1
COLORADO	6.2	8.0	1.8
CONNECTICUT	9.8	8.9	-0.9
DELAWARE	7.6	7.6	0.0
DISTRICT OF CO	7.6	7.4	-0.2
FLORIDA	10.5	10.2	-0.2
GEORGIA	12.4	14.1	1.7
HAWAII	8.6	9.6	1.0
IDAHO	12.1	11.8	-0.3
ILLINOIS	10.8	12.2	1.4
INDIANA	10.5	10.4	-0.1
IOWA	8.7	7.7	-1.0
KANSAS	11.6	12.4	0.8
KENTUCKY	13.9	11.4	-2.5
LOUISIANA	11.9	13.1	1.2
MAINE	8.3	8.0	-0.3
MARYLAND	11.3	7.9	-3.4
MASSACHUSETTS	7.5	8.6	1.1
MICHIGAN	8.5	10.9	2.5
MINNESOTA	41.5	28.9	-12.5
MISSISSIPPI	14.8	12.4	-2.4
MISSOURI	11.2	10.3	-0.9
MONTANA	16.8	15.5	-1.3
NEBRASKA	14.0	11.2	-2.8
NEVADA	8.3	10.2	1.9
NEW HAMPSHIRE	7.9	7.2	-0.7
NEW JERSEY	11.4	9.6	-1.7
NEW MEXICO	9.1	10.6	1.5
NEW YORK	8.3	8.7	0.3
NORTH CAROLINA	11.6	12.4	0.8
NORTH DAKOTA	9.8	10.3	0.5
OHIO	9.6	8.6	-1.0
OKLAHOMA	11.1	12.3	1.3
OREGON	10.7	10.8	0.1
PENNSYLVANIA	11.2	11.7	0.6
RHODE ISLAND	9.1	9.9	0.8
SOUTH CAROLINA	13.1	12.6	-0.6
SOUTH DAKOTA	11.1	9.8	-1.3
TENNESSEE	21.1	17.9	-3.2
TEXAS	9.5	10.5	1.0
UTAH	11.2	11.6	0.3
VERMONT	7.0	6.6	-0.4
VIRGINIA	9.1	10.5	1.4
WASHINGTON	8.7	8.4	-0.4
WEST VIRGINIA	12.2	12.2	-0.0
WISCONSIN	8.8	8.5	-0.3
WYOMING	9.5	8.5	-1.0
PUERTO RICO	12.0	15.1	3.1

\* RESPONDENT WOULD NOT ENCOURAGE SEXUALLY ACTIVE TEENAGERS TO USE A CONDOM

June 20, 2000

TABLE 26.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING HIV, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1626	5.0	0.6	( 3.8, 6.1)
ALASKA	1826	6.1	0.8	( 4.5, 7.7)
ARIZONA	1250	2.7	0.6	( 1.5, 3.9)
ARKANSAS	2371	6.4	0.6	( 5.2, 7.6)
COLORADO	2511	5.9	0.7	( 4.5, 7.3)
CONNECTICUT	2770	8.0	0.8	( 6.4, 9.6)
DELAWARE	2061	7.4	0.9	( 5.6, 9.3)
DISTRICT OF CO	971	10.6	1.3	( 8.1, 13.0)
FLORIDA	3857	7.5	0.6	( 6.4, 8.7)
GEORGIA	1879	6.8	0.7	( 5.4, 8.2)
HAWAII	1686	8.9	1.2	( 6.6, 11.2)
IDAHO	4026	5.0	0.5	( 4.1, 6.0)
ILLINOIS	2390	6.7	0.7	( 5.4, 8.1)
INDIANA	1979	7.0	1.1	( 4.9, 9.2)
IOWA	2644	7.6	0.6	( 6.4, 8.8)
KANSAS	2866	7.0	0.6	( 5.8, 8.3)
KENTUCKY	5643	6.6	0.4	( 5.7, 7.4)
LOUISIANA	1312	6.8	0.9	( 5.0, 8.6)
MAINE	1282	4.5	0.7	( 3.2, 5.8)
MARYLAND	3268	7.1	0.6	( 5.9, 8.3)
MASSACHUSETTS	3890	8.4	0.6	( 7.2, 9.6)
MICHIGAN	2016	6.9	0.8	( 5.2, 8.5)
MINNESOTA	4394	8.8	0.5	( 7.8, 9.7)
MISSISSIPPI	1720	7.5	0.7	( 6.1, 8.9)
MISSOURI	3239	6.1	0.6	( 5.0, 7.2)
MONTANA	1384	3.9	0.6	( 2.7, 5.0)
NEBRASKA	1985	6.8	0.7	( 5.4, 8.1)
NEVADA	1821	7.9	1.1	( 5.7, 10.0)
NEW HAMPSHIRE	986	5.8	0.9	( 4.0, 7.5)
NEW JERSEY	2147	8.0	0.8	( 6.5, 9.5)
NEW MEXICO	2776	7.6	0.6	( 6.5, 8.7)
NEW YORK	2051	7.3	0.7	( 5.9, 8.7)
NORTH CAROLINA	1841	6.0	0.6	( 4.8, 7.1)
NORTH DAKOTA	1479	5.6	0.7	( 4.3, 7.0)
OHIO	1206	7.4	1.1	( 5.3, 9.5)
OKLAHOMA	2253	5.0	0.6	( 3.9, 6.1)
OREGON	1432	5.3	0.7	( 3.9, 6.7)
PENNSYLVANIA	2742	5.6	0.5	( 4.6, 6.6)
RHODE ISLAND	3079	5.1	0.5	( 4.2, 6.1)
SOUTH CAROLINA	2810	6.7	0.6	( 5.6, 7.8)
SOUTH DAKOTA	3126	4.9	0.4	( 4.0, 5.7)
TENNESSEE	2309	5.9	0.6	( 4.8, 6.9)
TEXAS	4178	8.0	0.5	( 7.0, 9.0)
UTAH	2610	4.5	0.5	( 3.5, 5.6)
VERMONT	2475	5.0	0.5	( 4.0, 6.0)
VIRGINIA	2820	6.4	0.6	( 5.1, 7.6)
WASHINGTON	2915	3.8	0.4	( 3.0, 4.5)
WEST VIRGINIA	1951	6.9	0.7	( 5.6, 8.2)
WISCONSIN	1746	7.0	0.7	( 5.7, 8.2)
WYOMING	1979	6.4	0.6	( 5.2, 7.6)
PUERTO RICO	2243	11.8	0.9	( 10.1, 13.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 6.7 RANGE = 2.7-11.8

\* CHANCES OF GETTING INFECTED WITH HIV MEDIUM/HIGH

June 20, 2000

TABLE 26.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CHANCE OF GETTING HIV, AGE 18-64 YEARS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	685	5.8	1.0	( 3.9, 7.8)	*	941	4.2	0.7	( 2.8, 5.5)	*
ALASKA	896	5.7	1.1	( 3.5, 7.9)	*	930	6.5	1.2	( 4.1, 8.9)	*
ARIZONA	525	3.5	1.0	( 1.6, 5.5)	*	725	1.9	0.7	( 0.5, 3.3)	*
ARKANSAS	971	6.7	0.9	( 4.9, 8.5)	*	1400	6.1	0.7	( 4.7, 7.6)	*
COLORADO	1165	6.4	1.0	( 4.3, 8.4)	*	1346	5.5	0.9	( 3.7, 7.2)	*
CONNECTICUT	1104	10.1	1.4	( 7.4, 12.8)	*	1666	5.9	0.8	( 4.3, 7.5)	*
DELAWARE	831	9.1	1.4	( 6.2, 11.9)	*	1230	5.8	1.2	( 3.5, 8.1)	*
DISTRICT OF CO	360	13.4	2.2	( 9.0, 17.8)	*	611	7.9	1.2	( 5.6, 10.2)	*
FLORIDA	1551	8.7	1.0	( 6.7, 10.6)	*	2306	6.4	0.6	( 5.2, 7.6)	*
GEORGIA	760	7.9	1.2	( 5.5, 10.2)	*	1119	5.8	0.8	( 4.3, 7.3)	*
HAWAII	796	10.6	1.8	( 7.0, 14.1)	*	890	7.1	1.4	( 4.2, 9.9)	*
IDAHO	1744	5.3	0.8	( 3.7, 6.9)	*	2282	4.7	0.5	( 3.7, 5.7)	*
ILLINOIS	958	7.8	1.2	( 5.5, 10.2)	*	1432	5.6	0.7	( 4.3, 6.9)	*
INDIANA	859	10.1	2.0	( 6.2, 14.0)	*	1120	4.0	0.9	( 2.3, 5.8)	*
IOWA	1100	8.5	1.0	( 6.6, 10.4)	*	1544	6.7	0.7	( 5.3, 8.1)	*
KANSAS	1219	7.1	1.1	( 5.0, 9.1)	*	1647	7.0	0.8	( 5.5, 8.4)	*
KENTUCKY	2161	7.4	0.7	( 6.0, 8.8)	*	3482	5.8	0.6	( 4.7, 6.9)	*
LOUISIANA	544	7.1	1.3	( 4.6, 9.6)	*	768	6.6	1.2	( 4.1, 9.0)	*
MAINE	585	5.8	1.2	( 3.5, 8.1)	*	697	3.2	0.7	( 1.8, 4.6)	*
MARYLAND	1345	7.7	1.0	( 5.6, 9.7)	*	1923	6.6	0.7	( 5.2, 7.9)	*
MASSACHUSETTS	1613	9.9	1.0	( 7.9, 11.9)	*	2277	6.9	0.7	( 5.5, 8.3)	*
MICHIGAN	857	7.9	1.5	( 4.9, 10.8)	*	1159	5.9	0.8	( 4.4, 7.5)	*
MINNESOTA	2101	10.1	0.7	( 8.7, 11.6)	*	2293	7.4	0.6	( 6.2, 8.6)	*
MISSISSIPPI	707	8.0	1.2	( 5.7, 10.3)	*	1013	7.0	0.9	( 5.3, 8.6)	*
MISSOURI	1368	5.9	0.9	( 4.2, 7.6)	*	1871	6.3	0.7	( 4.9, 7.7)	*
MONTANA	628	3.6	0.8	( 2.0, 5.3)	*	756	4.1	0.8	( 2.5, 5.8)	*
NEBRASKA	841	9.0	1.2	( 6.6, 11.3)	*	1144	4.6	0.7	( 3.3, 5.9)	*
NEVADA	932	10.5	1.9	( 6.8, 14.2)	*	889	5.1	1.0	( 3.1, 7.1)	*
NEW HAMPSHIRE	404	5.3	1.2	( 2.9, 7.8)	*	582	6.2	1.3	( 3.8, 8.7)	*
NEW JERSEY	871	8.7	1.2	( 6.4, 11.0)	*	1276	7.4	1.0	( 5.4, 9.4)	*
NEW MEXICO	1258	8.5	0.9	( 6.8, 10.2)	*	1518	6.7	0.7	( 5.3, 8.1)	*
NEW YORK	859	8.2	1.1	( 6.0, 10.4)	*	1192	6.5	0.9	( 4.7, 8.2)	*
NORTH CAROLINA	767	5.9	0.9	( 4.2, 7.6)	*	1074	6.0	0.8	( 4.4, 7.5)	*
NORTH DAKOTA	693	4.6	0.9	( 2.9, 6.3)	*	786	6.7	1.0	( 4.7, 8.7)	*
OHIO	497	7.8	1.8	( 4.4, 11.3)	*	709	6.9	1.2	( 4.5, 9.3)	*
OKLAHOMA	918	5.3	0.9	( 3.5, 7.1)	*	1335	4.7	0.7	( 3.4, 6.1)	*
OREGON	633	5.7	1.1	( 3.6, 7.8)	*	799	4.8	0.9	( 2.9, 6.6)	*
PENNSYLVANIA	1127	6.5	0.9	( 4.7, 8.2)	*	1615	4.8	0.6	( 3.6, 5.9)	*
RHODE ISLAND	1250	5.7	0.8	( 4.1, 7.3)	*	1829	4.6	0.6	( 3.5, 5.7)	*
SOUTH CAROLINA	1155	8.0	1.0	( 6.2, 9.9)	*	1655	5.3	0.6	( 4.1, 6.5)	*
SOUTH DAKOTA	1312	4.2	0.6	( 3.0, 5.4)	*	1814	5.5	0.6	( 4.4, 6.6)	*
TENNESSEE	826	5.9	0.9	( 4.2, 7.6)	*	1483	5.9	0.7	( 4.6, 7.2)	*
TEXAS	1712	8.5	0.8	( 6.9, 10.1)	*	2466	7.5	0.7	( 6.2, 8.9)	*
UTAH	1148	4.4	0.8	( 2.8, 6.0)	*	1462	4.6	0.7	( 3.3, 6.0)	*
VERMONT	1053	5.8	0.8	( 4.2, 7.5)	*	1422	4.2	0.6	( 3.1, 5.4)	*
VIRGINIA	1164	7.2	1.1	( 5.1, 9.3)	*	1656	5.5	0.7	( 4.2, 6.8)	*
WASHINGTON	1336	4.5	0.6	( 3.3, 5.7)	*	1579	3.0	0.5	( 2.1, 3.9)	*
WEST VIRGINIA	850	7.2	1.0	( 5.2, 9.2)	*	1101	6.7	0.8	( 5.0, 8.3)	*
WISCONSIN	775	6.8	1.0	( 4.9, 8.7)	*	971	7.1	0.9	( 5.4, 8.9)	*
WYOMING	905	7.1	0.9	( 5.2, 8.9)	*	1074	5.7	0.8	( 4.2, 7.2)	*
PUERTO RICO	818	14.2	1.5	( 11.3, 17.1)	*	1425	9.6	0.9	( 7.8, 11.4)	*

SUMMARY STATISTICS:           NUMBER OF PARTICIPANTS = 51  
   MEDIAN                         = 7.2  
   RANGE                         = 3.5-14.2

NUMBER OF PARTICIPANTS = 51  
 MEDIAN                         = 5.9  
 RANGE                         = 1.9- 9.6

\* CHANCES OF GETTING INFECTED WITH HIV MEDIUM/HIGH

TABLE 26.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 CHANCE OF GETTING HIV, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	6.1	5.0	-1.1
ALASKA	5.6	6.1	0.5
ARIZONA	1.9	2.7	0.8
ARKANSAS	5.5	6.4	0.9
COLORADO	6.2	5.9	-0.3
CONNECTICUT	6.8	8.0	1.2
DELAWARE	4.7	7.4	2.7
DISTRICT OF CO	8.5	10.6	2.1
FLORIDA	6.1	7.5	1.4
GEORGIA	5.7	6.8	1.1
HAWAII	7.7	8.9	1.1
IDAHO	6.9	5.0	-1.9
ILLINOIS	6.7	6.7	0.0
INDIANA	6.5	7.0	0.5
IOWA	5.4	7.6	2.2
KANSAS	5.0	7.0	2.0
KENTUCKY	5.2	6.6	1.4
LOUISIANA	7.3	6.8	-0.5
MAINE	5.8	4.5	-1.3
MARYLAND	6.0	7.1	1.1
MASSACHUSETTS	7.3	8.4	1.0
MICHIGAN	4.1	6.9	2.8
MINNESOTA	9.3	8.8	-0.5
MISSISSIPPI	7.8	7.5	-0.3
MISSOURI	6.5	6.1	-0.4
MONTANA	3.8	3.9	0.1
NEBRASKA	7.7	6.8	-0.9
NEVADA	6.5	7.9	1.3
NEW HAMPSHIRE	4.2	5.8	1.6
NEW JERSEY	7.4	8.0	0.6
NEW MEXICO	5.5	7.6	2.1
NEW YORK	7.3	7.3	-0.0
NORTH CAROLINA	6.6	6.0	-0.7
NORTH DAKOTA	4.7	5.6	1.0
OHIO	6.0	7.4	1.4
OKLAHOMA	6.2	5.0	-1.2
OREGON	5.9	5.3	-0.7
PENNSYLVANIA	6.6	5.6	-1.0
RHODE ISLAND	6.8	5.1	-1.6
SOUTH CAROLINA	6.8	6.7	-0.1
SOUTH DAKOTA	11.0	4.9	-6.2
TENNESSEE	6.7	5.9	-0.8
TEXAS	7.0	8.0	1.1
UTAH	4.8	4.5	-0.3
VERMONT	5.3	5.0	-0.2
VIRGINIA	5.9	6.4	0.5
WASHINGTON	3.8	3.8	-0.1
WEST VIRGINIA	11.4	6.9	-4.4
WISCONSIN	5.6	7.0	1.4
WYOMING	6.3	6.4	0.1
PUERTO RICO	10.2	11.8	1.6

\* CHANCES OF GETTING INFECTED WITH HIV MEDIUM/HIGH

June 20, 2000

TABLE 27.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO HIV TEST IN PAST 12 MONTHS, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	256	57.2	3.6	( 50.1, 64.3)
ALASKA	245	66.0	4.0	( 58.2, 73.8)
ARIZONA	159	67.6	5.7	( 56.5, 78.8)
ARKANSAS	342	64.5	3.0	( 58.6, 70.4)
COLORADO	321	61.6	3.9	( 54.0, 69.3)
CONNECTICUT	347	64.4	3.4	( 57.6, 71.1)
DELAWARE	340	61.4	3.6	( 54.4, 68.4)
DISTRICT OF CO	145	49.7	4.7	( 40.5, 58.9)
FLORIDA	644	60.5	2.4	( 55.9, 65.2)
GEORGIA	319	57.6	3.4	( 51.0, 64.1)
HAWAII	202	50.1	4.6	( 41.1, 59.0)
IDAHO	452	71.4	2.9	( 65.8, 77.0)
INDIANA	242	57.4	5.3	( 47.1, 67.8)
IOWA	239	65.7	3.7	( 58.5, 72.9)
KANSAS	289	62.0	3.2	( 55.7, 68.2)
KENTUCKY	542	56.7	3.0	( 50.8, 62.7)
LOUISIANA	215	53.3	3.9	( 45.7, 60.9)
MAINE	146	59.7	4.6	( 50.7, 68.7)
MARYLAND	528	63.1	2.8	( 57.7, 68.5)
MASSACHUSETTS	530	72.1	2.4	( 67.3, 76.8)
MICHIGAN	263	64.3	3.2	( 57.9, 70.6)
MINNESOTA	348	67.5	2.7	( 62.2, 72.8)
MISSISSIPPI	278	60.6	3.4	( 54.0, 67.3)
MISSOURI	404	56.7	3.6	( 49.6, 63.8)
MONTANA	122	68.7	4.5	( 59.8, 77.5)
NEBRASKA	192	66.8	3.7	( 59.6, 73.9)
NEVADA	313	65.8	4.1	( 57.8, 73.8)
NEW HAMPSHIRE	123	59.9	5.3	( 49.5, 70.3)
NEW JERSEY	291	64.3	3.3	( 57.7, 70.8)
NEW MEXICO	373	65.2	2.8	( 59.8, 70.6)
NEW YORK	247	52.5	3.6	( 45.6, 59.5)
NORTH CAROLINA	279	53.7	3.5	( 46.8, 60.5)
NORTH DAKOTA	134	60.6	5.0	( 50.8, 70.3)
OHIO	111	73.3	5.0	( 63.4, 83.1)
OKLAHOMA	301	60.0	3.2	( 53.6, 66.3)
OREGON	182	65.6	3.8	( 58.1, 73.1)
PENNSYLVANIA	356	64.1	3.1	( 58.1, 70.2)
RHODE ISLAND	411	59.0	2.9	( 53.3, 64.6)
SOUTH CAROLINA	455	56.7	2.8	( 51.1, 62.3)
SOUTH DAKOTA	324	57.8	3.4	( 51.2, 64.4)
TENNESSEE	320	57.1	3.2	( 50.8, 63.4)
TEXAS	691	61.9	2.2	( 57.6, 66.3)
UTAH	318	69.9	3.7	( 62.6, 77.1)
VERMONT	309	68.9	3.2	( 62.6, 75.2)
VIRGINIA	442	54.7	3.6	( 47.5, 61.8)
WASHINGTON	421	70.2	2.8	( 64.8, 75.7)
WEST VIRGINIA	166	66.0	4.5	( 57.3, 74.7)
WISCONSIN	189	70.5	3.9	( 62.8, 78.2)
WYOMING	240	64.5	3.3	( 58.0, 71.0)
PUERTO RICO	247	62.6	3.5	( 55.8, 69.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 50 MEDIAN = 62.3 RANGE = 49.7-73.3

\* NO HIV TEST OUTSIDE OF BLOOD DONATIONS

June 20, 2000



TABLE 27.3 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO HIV TEST IN PAST 12 MONTHS, AGE 18-64 YEARS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	56.4	57.2	0.9
ALASKA	58.8	66.0	7.1
ARIZONA	69.4	67.6	-1.8
ARKANSAS	61.2	64.5	3.3
COLORADO	68.4	61.6	-6.8
CONNECTICUT	57.8	64.4	6.6
DELAWARE	57.0	61.4	4.4
DISTRICT OF CO	58.8	49.7	-9.1
FLORIDA	56.4	60.5	4.1
GEORGIA	51.2	57.6	6.4
HAWAII	45.4	50.1	4.7
IDAHO	64.5	71.4	6.8
INDIANA	61.6	57.4	-4.2
IOWA	66.1	65.7	-0.4
KANSAS	53.7	62.0	8.2
KENTUCKY	61.8	56.7	-5.0
LOUISIANA	53.3	53.3	-0.1
MAINE	62.2	59.7	-2.4
MARYLAND	57.1	63.1	6.0
MASSACHUSETTS	64.7	72.1	7.4
MICHIGAN	63.2	64.3	1.1
MINNESOTA	61.1	67.5	6.4
MISSISSIPPI	49.8	60.6	10.9
MISSOURI	62.5	56.7	-5.8
MONTANA	59.9	68.7	8.8
NEBRASKA	63.1	66.8	3.6
NEVADA	59.5	65.8	6.3
NEW HAMPSHIRE	61.2	59.9	-1.4
NEW JERSEY	63.2	64.3	1.1
NEW MEXICO	63.0	65.2	2.2
NEW YORK	57.2	52.5	-4.6
NORTH CAROLINA	53.6	53.7	0.0
NORTH DAKOTA	70.7	60.6	-10.1
OHIO	66.4	73.3	6.8
OKLAHOMA	58.4	60.0	1.5
OREGON	61.8	65.6	3.8
PENNSYLVANIA	56.8	64.1	7.4
RHODE ISLAND	62.7	59.0	-3.7
SOUTH CAROLINA	58.7	56.7	-1.9
SOUTH DAKOTA	56.7	57.8	1.1
TENNESSEE	55.7	57.1	1.5
TEXAS	59.4	61.9	2.5
UTAH	69.0	69.9	0.8
VERMONT	63.9	68.9	5.0
VIRGINIA	56.4	54.7	-1.7
WASHINGTON	63.6	70.2	6.7
WEST VIRGINIA	51.9	66.0	14.1
WISCONSIN	63.6	70.5	6.9
WYOMING	67.3	64.5	-2.8
PUERTO RICO	56.3	62.6	6.3

\* NO HIV TEST OUTSIDE OF BLOOD DONATIONS

June 20, 2000

TABLE 28.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET OF DIABETES\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	162	47.5	1.5	(44.7, 50.3)
ALASKA	75	42.3	3.5	(35.5, 49.0)
ARIZONA	88	41.4	1.8	(37.8, 44.9)
ARKANSAS	189	49.6	1.3	(47.0, 52.1)
CALIFORNIA	233	49.0	1.0	(47.1, 50.8)
COLORADO	119	47.5	2.6	(42.4, 52.5)
CONNECTICUT	158	46.7	1.6	(43.6, 49.7)
DISTRICT OF CO	83	48.9	2.0	(45.1, 52.7)
FLORIDA	353	50.6	1.0	(48.7, 52.5)
GEORGIA	125	47.8	1.5	(44.9, 50.7)
HAWAII	93	46.5	2.0	(42.6, 50.4)
IOWA	203	49.9	1.3	(47.4, 52.4)
KENTUCKY	537	49.5	0.9	(47.8, 51.2)
LOUISIANA	102	47.4	1.7	(44.1, 50.7)
MAINE	97	48.5	1.7	(45.2, 51.8)
MASSACHUSETTS	244	50.3	1.3	(47.7, 52.8)
MICHIGAN	135	49.4	1.4	(46.6, 52.2)
MINNESOTA	243	52.7	1.1	(50.6, 54.7)
MONTANA	107	49.7	1.9	(45.9, 53.5)
NEBRASKA	133	50.2	1.7	(46.8, 53.5)
NEVADA	125	54.9	1.5	(52.0, 57.8)
NEW HAMPSHIRE	54	46.0	2.7	(40.7, 51.3)
NEW JERSEY	141	49.3	1.7	(45.9, 52.6)
NEW MEXICO	201	45.5	1.3	(43.1, 48.0)
NORTH CAROLINA	165	46.9	1.4	(44.2, 49.6)
NORTH DAKOTA	105	52.2	1.8	(48.8, 55.7)
OHIO	111	48.7	1.8	(45.2, 52.3)
OKLAHOMA	168	46.4	1.4	(43.7, 49.1)
PENNSYLVANIA	208	50.2	1.3	(47.6, 52.8)
RHODE ISLAND	183	47.5	1.5	(44.7, 50.4)
TENNESSEE	167	44.9	1.5	(41.9, 47.8)
TEXAS	285	45.8	1.1	(43.6, 47.9)
UTAH	143	50.9	1.6	(47.7, 54.0)
VERMONT	144	51.2	1.6	(48.0, 54.4)
VIRGINIA	226	47.2	1.7	(43.8, 50.6)
WEST VIRGINIA	199	50.2	1.1	(47.9, 52.4)
WISCONSIN	105	49.7	1.9	(45.9, 53.4)
WYOMING	106	49.5	1.7	(46.3, 52.8)
PUERTO RICO	349	47.2	0.9	(45.3, 49.0)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 39 MEDIAN = 48.9 RANGE = 41.4-54.9

\* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

June 20, 2000

TABLE 28.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR AGE AT ONSET OF DIABETES\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	68	47.7	2.1	(43.5, 51.9)	*	94	47.3	2.0	(43.4, 51.1)	*
ALASKA	41	39.8	4.8	(30.5, 49.2)	*	34	45.6	4.2	(37.4, 53.8)	*
ARIZONA	37	38.6	2.5	(33.7, 43.4)	*	51	45.3	2.2	(41.1, 49.6)	*
ARKANSAS	68	48.0	2.2	(43.8, 52.2)	*	121	50.8	1.5	(47.8, 53.8)	*
CALIFORNIA	102	49.1	1.4	(46.2, 51.9)	*	131	48.8	1.3	(46.4, 51.3)	*
COLORADO	50	49.9	4.3	(41.5, 58.3)	*	69	45.5	3.0	(39.6, 51.5)	*
CONNECTICUT	62	47.4	1.9	(43.7, 51.1)	*	96	46.0	2.4	(41.2, 50.8)	*
DISTRICT OF CO	29	42.7	2.5	(37.8, 47.5)	*	54	53.5	2.5	(48.6, 58.4)	*
FLORIDA	144	48.6	1.5	(45.6, 51.5)	*	209	52.6	1.2	(50.2, 55.0)	*
GEORGIA	48	47.6	2.3	(43.0, 52.1)	*	77	48.0	1.9	(44.2, 51.7)	*
HAWAII	51	43.7	2.5	(38.9, 48.6)	*	42	51.2	2.8	(45.8, 56.6)	*
IOWA	98	50.0	1.5	(47.0, 53.0)	*	105	49.8	2.1	(45.7, 53.9)	*
KENTUCKY	189	49.2	1.3	(46.6, 51.7)	*	348	49.8	1.2	(47.5, 52.0)	*
LOUISIANA	43	48.9	2.4	(44.1, 53.6)	*	59	46.1	2.4	(41.5, 50.7)	*
MAINE	39	51.3	2.9	(45.5, 57.1)	*	58	46.5	2.0	(42.6, 50.3)	*
MASSACHUSETTS	116	50.7	1.5	(47.6, 53.7)	*	128	49.7	2.4	(45.0, 54.3)	*
MICHIGAN	65	49.2	2.1	(45.1, 53.3)	*	70	49.6	1.9	(45.8, 53.4)	*
MINNESOTA	119	52.9	1.5	(50.0, 55.8)	*	124	52.4	1.5	(49.5, 55.3)	*
MONTANA	49	48.6	2.9	(42.9, 54.3)	*	58	50.9	2.5	(46.1, 55.7)	*
NEBRASKA	56	46.6	2.7	(41.2, 51.9)	*	77	54.0	1.8	(50.6, 57.4)	*
NEVADA	68	54.8	2.0	(51.0, 58.6)	*	57	55.1	2.2	(50.7, 59.5)	*
NEW HAMPSHIRE	23	42.6	3.9	(34.9, 50.3)	*	31	49.7	3.5	(42.9, 56.4)	*
NEW JERSEY	57	52.1	2.0	(48.1, 56.0)	*	84	47.1	2.5	(42.3, 51.9)	*
NEW MEXICO	79	47.0	1.5	(44.1, 49.9)	*	122	44.5	1.8	(41.0, 48.1)	*
NORTH CAROLINA	63	48.0	1.8	(44.5, 51.4)	*	102	46.1	2.1	(42.0, 50.1)	*
NORTH DAKOTA	41	46.6	3.1	(40.6, 52.6)	*	64	56.6	1.8	(53.0, 60.2)	*
OHIO	38	49.9	3.0	(44.0, 55.8)	*	73	48.0	2.3	(43.5, 52.4)	*
OKLAHOMA	74	46.1	1.9	(42.3, 49.9)	*	94	46.8	2.0	(42.8, 50.7)	*
PENNSYLVANIA	88	50.2	1.9	(46.4, 53.9)	*	120	50.2	1.8	(46.6, 53.8)	*
RHODE ISLAND	79	47.6	2.3	(43.0, 52.1)	*	104	47.5	1.8	(44.0, 51.0)	*
TENNESSEE	56	44.6	2.6	(39.4, 49.7)	*	111	45.1	1.8	(41.7, 48.6)	*
TEXAS	120	45.9	1.7	(42.6, 49.3)	*	165	45.6	1.3	(43.0, 48.2)	*
UTAH	64	48.3	2.6	(43.2, 53.4)	*	79	53.4	1.8	(49.9, 56.8)	*
VERMONT	60	49.2	2.5	(44.3, 54.1)	*	84	53.2	2.0	(49.4, 57.1)	*
VIRGINIA	96	46.8	2.7	(41.6, 52.1)	*	130	47.7	1.9	(43.9, 51.5)	*
WEST VIRGINIA	78	48.9	1.9	(45.2, 52.6)	*	121	51.2	1.4	(48.5, 53.9)	*
WISCONSIN	53	49.2	2.7	(43.9, 54.5)	*	52	50.3	2.7	(45.1, 55.5)	*
WYOMING	45	50.6	2.3	(46.0, 55.1)	*	61	48.7	2.4	(44.1, 53.3)	*
PUERTO RICO	109	47.5	1.3	(44.9, 50.1)	*	240	46.9	1.3	(44.4, 49.4)	*

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 39  
 MEDIAN = 48.3  
 RANGE = 38.6-54.8

NUMBER OF PARTICIPANTS = 39  
 MEDIAN = 48.8  
 RANGE = 44.5-56.6

\* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

June 20, 2000

TABLE 29.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETICS TAKING INSULIN\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	170	31.0	3.8	( 23.5, 38.4)
ALASKA	77	29.9	7.7	( 14.8, 45.0)
ARIZONA	90	41.9	7.2	( 27.8, 56.1)
ARKANSAS	195	31.5	3.8	( 24.0, 39.0)
CALIFORNIA	233	26.9	3.4	( 20.2, 33.6)
COLORADO	121	36.1	5.8	( 24.6, 47.5)
CONNECTICUT	166	35.2	4.8	( 25.7, 44.7)
DISTRICT OF CO	92	32.7	5.2	( 22.5, 42.9)
FLORIDA	363	31.9	3.1	( 25.9, 38.0)
GEORGIA	130	35.1	4.7	( 25.8, 44.3)
HAWAII	102	18.8	4.9	( 9.1, 28.4)
IOWA	214	27.1	3.3	( 20.6, 33.7)
KENTUCKY	561	32.0	2.6	( 26.8, 37.1)
LOUISIANA	108	45.4	5.3	( 34.9, 55.8)
MAINE	98	30.1	5.1	( 20.1, 40.1)
MASSACHUSETTS	256	32.1	3.9	( 24.5, 39.7)
MICHIGAN	140	28.9	4.1	( 20.9, 36.9)
MINNESOTA	245	34.2	3.3	( 27.8, 40.6)
MONTANA	107	38.1	5.3	( 27.6, 48.6)
NEBRASKA	136	33.7	4.5	( 24.9, 42.5)
NEVADA	130	40.5	5.9	( 28.9, 52.0)
NEW HAMPSHIRE	57	32.3	7.1	( 18.3, 46.2)
NEW JERSEY	147	23.2	3.8	( 15.8, 30.7)
NEW MEXICO	205	32.6	3.8	( 25.2, 40.0)
NORTH CAROLINA	175	32.2	4.0	( 24.4, 40.0)
NORTH DAKOTA	107	35.3	4.9	( 25.6, 44.9)
OHIO	115	36.2	5.3	( 25.9, 46.5)
OKLAHOMA	182	31.7	3.9	( 24.0, 39.4)
PENNSYLVANIA	214	34.5	3.9	( 26.9, 42.1)
RHODE ISLAND	196	24.9	3.4	( 18.2, 31.6)
TENNESSEE	181	38.0	4.3	( 29.6, 46.4)
TEXAS	296	26.5	3.1	( 20.6, 32.5)
UTAH	148	30.7	5.2	( 20.6, 40.8)
VERMONT	150	31.7	4.3	( 23.3, 40.1)
VIRGINIA	242	25.6	5.1	( 15.7, 35.5)
WEST VIRGINIA	212	26.4	3.3	( 19.9, 32.8)
WISCONSIN	106	30.6	4.8	( 21.1, 40.0)
WYOMING	109	32.9	4.8	( 23.5, 42.3)
PUERTO RICO	373	32.9	2.8	( 27.5, 38.3)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 39 MEDIAN = 32.1 RANGE = 18.8-45.4

\* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

June 20, 2000

TABLE 29.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR DIABETICS TAKING INSULIN\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	70	31.2	5.9	( 19.6, 42.7)	*	100	30.8	5.0	( 21.1, 40.5)	*
ALASKA	42	36.6	11.2	( 14.6, 58.6)	*	35	20.8	7.7	( 5.8, 35.8)	*
ARIZONA	38	48.5	10.8	( 27.3, 69.8)	*	52	32.9	7.5	( 18.2, 47.6)	*
ARKANSAS	69	28.3	5.9	( 16.8, 39.8)	*	126	34.0	5.0	( 24.2, 43.8)	*
CALIFORNIA	101	24.1	5.1	( 14.1, 34.0)	*	132	29.6	4.6	( 20.6, 38.6)	*
COLORADO	51	36.2	9.2	( 18.2, 54.2)	*	70	35.9	7.5	( 21.2, 50.6)	*
CONNECTICUT	68	35.5	7.6	( 20.6, 50.4)	*	98	35.0	6.1	( 23.1, 46.8)	*
DISTRICT OF CO	33	29.6	8.6	( 12.8, 46.4)	*	59	35.2	6.5	( 22.5, 47.8)	*
FLORIDA	145	38.6	5.1	( 28.7, 48.5)	*	218	25.8	3.4	( 19.0, 32.5)	*
GEORGIA	49	36.0	7.6	( 21.2, 50.8)	*	81	34.3	6.0	( 22.6, 46.0)	*
HAWAII	54	15.4	6.4	( 2.9, 28.0)	*	48	24.3	7.7	( 9.3, 39.3)	*
IOWA	101	25.2	4.6	( 16.2, 34.2)	*	113	29.3	4.8	( 19.8, 38.8)	*
KENTUCKY	193	31.3	4.4	( 22.8, 39.8)	*	368	32.5	3.2	( 26.3, 38.7)	*
LOUISIANA	44	37.3	8.1	( 21.5, 53.0)	*	64	52.4	7.1	( 38.6, 66.2)	*
MAINE	39	31.4	7.9	( 15.9, 46.9)	*	59	29.3	6.7	( 16.2, 42.4)	*
MASSACHUSETTS	123	31.7	5.5	( 21.0, 42.4)	*	133	32.7	5.2	( 22.5, 42.8)	*
MICHIGAN	66	26.1	5.6	( 15.2, 37.1)	*	74	32.2	5.9	( 20.6, 43.8)	*
MINNESOTA	119	32.8	4.5	( 24.0, 41.5)	*	126	35.8	4.7	( 26.6, 44.9)	*
MONTANA	49	40.4	8.0	( 24.8, 56.0)	*	58	35.4	6.8	( 22.1, 48.7)	*
NEBRASKA	56	39.7	7.1	( 25.9, 53.5)	*	80	27.7	5.3	( 17.3, 38.0)	*
NEVADA	70	33.9	7.6	( 19.0, 48.8)	*	60	49.5	8.9	( 32.0, 67.0)	*
NEW HAMPSHIRE	23	31.9	11.2	( 10.0, 53.7)	*	34	32.6	8.9	( 15.2, 50.0)	*
NEW JERSEY	58	17.5	5.1	( 7.5, 27.6)	*	89	27.5	5.3	( 17.2, 37.8)	*
NEW MEXICO	79	33.9	5.9	( 22.4, 45.5)	*	126	31.6	4.9	( 22.1, 41.1)	*
NORTH CAROLINA	65	36.1	6.5	( 23.4, 48.8)	*	110	29.3	4.9	( 19.6, 38.9)	*
NORTH DAKOTA	41	44.5	8.2	( 28.5, 60.5)	*	66	28.2	5.9	( 16.7, 39.7)	*
OHIO	38	33.1	8.8	( 15.8, 50.4)	*	77	38.2	6.5	( 25.4, 51.0)	*
OKLAHOMA	81	36.1	5.9	( 24.5, 47.8)	*	101	26.6	4.8	( 17.2, 36.0)	*
PENNSYLVANIA	91	30.4	5.1	( 20.5, 40.3)	*	123	38.4	5.7	( 27.2, 49.6)	*
RHODE ISLAND	83	17.5	4.7	( 8.4, 26.6)	*	113	31.8	5.0	( 22.1, 41.6)	*
TENNESSEE	59	37.7	7.4	( 23.2, 52.2)	*	122	38.2	5.1	( 28.3, 48.1)	*
TEXAS	121	20.2	3.9	( 12.6, 27.8)	*	175	32.6	4.7	( 23.3, 41.8)	*
UTAH	66	27.8	7.3	( 13.6, 42.1)	*	82	33.4	7.3	( 19.2, 47.6)	*
VERMONT	62	30.3	6.5	( 17.4, 43.1)	*	88	33.2	5.4	( 22.5, 43.9)	*
VIRGINIA	102	23.6	8.4	( 7.1, 40.2)	*	140	28.0	4.8	( 18.6, 37.5)	*
WEST VIRGINIA	82	33.7	5.6	( 22.7, 44.8)	*	130	20.5	3.7	( 13.3, 27.8)	*
WISCONSIN	52	23.2	6.3	( 10.9, 35.4)	*	54	39.3	7.2	( 25.1, 53.4)	*
WYOMING	46	32.5	7.4	( 18.1, 46.9)	*	63	33.3	6.4	( 20.8, 45.7)	*
PUERTO RICO	118	31.5	4.7	( 22.3, 40.7)	*	255	33.8	3.4	( 27.3, 40.4)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 39  
 MEDIAN = 31.9  
 RANGE = 15.4-48.5

NUMBER OF PARTICIPANTS = 39  
 MEDIAN = 32.6  
 RANGE = 20.5-52.4

\* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

June 20, 2000

TABLE 30.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR VISITS TO HEALTH PROFESSIONALS IN PAST YEAR\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	157	3.7	0.3	( 3.1, 4.3)
ALASKA	76	4.2	0.5	( 3.2, 5.3)
ARIZONA	85	4.6	0.6	( 3.4, 5.8)
ARKANSAS	186	4.5	0.5	( 3.5, 5.6)
CALIFORNIA	228	5.0	0.4	( 4.2, 5.8)
COLORADO	119	6.3	1.0	( 4.3, 8.3)
CONNECTICUT	158	4.6	0.4	( 3.9, 5.3)
DISTRICT OF CO	83	4.3	0.4	( 3.5, 5.0)
FLORIDA	349	4.2	0.3	( 3.7, 4.8)
GEORGIA	122	3.7	0.3	( 3.1, 4.4)
HAWAII	99	4.4	0.4	( 3.6, 5.2)
IOWA	208	4.4	0.4	( 3.7, 5.1)
KENTUCKY	526	4.8	0.2	( 4.3, 5.2)
LOUISIANA	99	4.8	0.6	( 3.6, 5.9)
MAINE	97	3.8	0.4	( 3.1, 4.6)
MASSACHUSETTS	248	4.0	0.4	( 3.3, 4.8)
MICHIGAN	131	3.9	0.5	( 3.0, 4.9)
MINNESOTA	240	4.0	0.3	( 3.5, 4.5)
MONTANA	101	5.2	0.8	( 3.7, 6.7)
NEBRASKA	136	3.6	0.3	( 3.0, 4.1)
NEVADA	129	5.0	1.1	( 2.9, 7.1)
NEW HAMPSHIRE	55	2.6	0.3	( 2.1, 3.2)
NEW JERSEY	145	4.9	0.4	( 4.1, 5.7)
NEW MEXICO	195	4.5	0.4	( 3.7, 5.2)
NORTH CAROLINA	164	4.8	0.4	( 4.0, 5.6)
NORTH DAKOTA	106	4.5	0.5	( 3.6, 5.3)
OHIO	113	4.3	0.3	( 3.7, 5.0)
OKLAHOMA	175	4.4	0.3	( 3.8, 4.9)
PENNSYLVANIA	204	4.6	0.3	( 4.0, 5.2)
RHODE ISLAND	190	5.2	0.9	( 3.6, 6.9)
TENNESSEE	164	3.7	0.3	( 3.2, 4.2)
TEXAS	283	5.1	0.5	( 4.2, 6.0)
UTAH	148	4.6	0.5	( 3.6, 5.5)
VERMONT	140	3.3	0.3	( 2.7, 3.9)
VIRGINIA	223	4.1	0.4	( 3.4, 4.8)
WEST VIRGINIA	208	4.6	0.4	( 3.9, 5.4)
WISCONSIN	108	3.8	0.4	( 3.1, 4.6)
WYOMING	109	5.2	0.7	( 3.8, 6.6)
PUERTO RICO	340	7.0	0.3	( 6.4, 7.6)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 39 MEDIAN = 4.5 RANGE = 2.6- 7.0

\* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

June 20, 2000

TABLE 30.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED MEAN AND RELEVANT STATISTICS FOR VISITS TO HEALTH PROFESSIONALS IN PAST YEAR\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	MEAN	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	66	4.1	0.5	( 3.2, 5.0)	*	91	3.4	0.4	( 2.6, 4.1)	*
ALASKA	41	3.9	0.6	( 2.7, 5.2)	*	35	4.6	0.9	( 2.8, 6.4)	*
ARIZONA	37	4.7	0.8	( 3.1, 6.4)	*	48	4.4	0.9	( 2.6, 6.1)	*
ARKANSAS	66	3.7	0.6	( 2.6, 4.9)	*	120	5.2	0.8	( 3.5, 6.8)	*
CALIFORNIA	101	4.6	0.5	( 3.7, 5.6)	*	127	5.4	0.6	( 4.2, 6.5)	*
COLORADO	50	7.6	2.1	( 3.6, 11.6)	*	69	5.3	0.9	( 3.6, 6.9)	*
CONNECTICUT	65	4.6	0.5	( 3.6, 5.6)	*	93	4.7	0.5	( 3.6, 5.7)	*
DISTRICT OF CO	30	3.8	0.6	( 2.7, 4.9)	*	53	4.6	0.5	( 3.6, 5.6)	*
FLORIDA	140	4.1	0.5	( 3.2, 5.1)	*	209	4.3	0.3	( 3.7, 4.9)	*
GEORGIA	47	3.7	0.6	( 2.5, 4.8)	*	75	3.8	0.4	( 3.1, 4.6)	*
HAWAII	53	4.1	0.5	( 3.1, 5.1)	*	46	4.9	0.7	( 3.5, 6.2)	*
IOWA	98	4.2	0.6	( 3.1, 5.3)	*	110	4.6	0.5	( 3.7, 5.6)	*
KENTUCKY	183	5.0	0.4	( 4.2, 5.8)	*	343	4.6	0.2	( 4.2, 5.1)	*
LOUISIANA	41	4.7	0.6	( 3.5, 5.8)	*	58	4.8	0.9	( 3.0, 6.7)	*
MAINE	39	2.9	0.4	( 2.1, 3.6)	*	58	4.6	0.6	( 3.4, 5.7)	*
MASSACHUSETTS	121	3.6	0.5	( 2.6, 4.6)	*	127	4.8	0.6	( 3.6, 6.0)	*
MICHIGAN	64	3.3	0.6	( 2.2, 4.3)	*	67	4.8	0.8	( 3.3, 6.3)	*
MINNESOTA	117	4.5	0.4	( 3.7, 5.2)	*	123	3.4	0.3	( 2.9, 4.0)	*
MONTANA	46	5.7	1.0	( 3.8, 7.6)	*	55	4.6	1.2	( 2.2, 7.0)	*
NEBRASKA	56	3.0	0.3	( 2.4, 3.7)	*	80	4.1	0.4	( 3.3, 4.8)	*
NEVADA	70	2.8	0.4	( 2.0, 3.5)	*	59	8.1	2.4	( 3.5, 12.7)	*
NEW HAMPSHIRE	23	2.5	0.4	( 1.7, 3.3)	*	32	2.8	0.4	( 2.0, 3.6)	*
NEW JERSEY	58	4.5	0.6	( 3.3, 5.7)	*	87	5.2	0.6	( 4.1, 6.4)	*
NEW MEXICO	77	3.9	0.4	( 3.1, 4.7)	*	118	4.9	0.6	( 3.7, 6.0)	*
NORTH CAROLINA	63	5.0	0.8	( 3.5, 6.5)	*	101	4.6	0.4	( 3.9, 5.3)	*
NORTH DAKOTA	41	4.5	0.7	( 3.2, 5.8)	*	65	4.4	0.6	( 3.2, 5.6)	*
OHIO	37	4.0	0.4	( 3.1, 4.8)	*	76	4.5	0.5	( 3.6, 5.4)	*
OKLAHOMA	79	3.6	0.3	( 3.0, 4.2)	*	96	5.3	0.5	( 4.2, 6.3)	*
PENNSYLVANIA	87	4.8	0.5	( 3.8, 5.8)	*	117	4.5	0.4	( 3.7, 5.2)	*
RHODE ISLAND	81	6.2	1.7	( 2.8, 9.5)	*	109	4.4	0.4	( 3.7, 5.0)	*
TENNESSEE	55	3.3	0.3	( 2.6, 4.0)	*	109	4.1	0.3	( 3.4, 4.7)	*
TEXAS	120	4.8	0.7	( 3.5, 6.0)	*	163	5.4	0.6	( 4.2, 6.6)	*
UTAH	66	4.1	0.6	( 3.0, 5.2)	*	82	5.0	0.8	( 3.4, 6.5)	*
VERMONT	61	3.4	0.5	( 2.4, 4.3)	*	79	3.3	0.4	( 2.6, 4.0)	*
VIRGINIA	94	3.4	0.4	( 2.7, 4.1)	*	129	4.8	0.6	( 3.7, 6.0)	*
WEST VIRGINIA	81	4.3	0.7	( 3.0, 5.6)	*	127	4.8	0.4	( 4.0, 5.7)	*
WISCONSIN	53	4.0	0.5	( 3.0, 4.9)	*	55	3.7	0.6	( 2.5, 4.8)	*
WYOMING	46	5.2	1.3	( 2.6, 7.8)	*	63	5.2	0.7	( 3.8, 6.6)	*
PUERTO RICO	110	6.3	0.5	( 5.4, 7.2)	*	230	7.5	0.4	( 6.7, 8.3)	*

SUMMARY STATISTICS: NUMBER OF PARTICIPANTS = 39  
 MEDIAN = 4.1  
 RANGE = 2.5- 7.6

NUMBER OF PARTICIPANTS = 39  
 MEDIAN = 4.6  
 RANGE = 2.8- 8.1

\* RESPONDENTS TOLD BY DOCTOR THEY HAVE DIABETES

June 20, 2000

TABLE 31.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	2050	27.7	1.5	( 24.8, 30.6)
ARIZONA	1739	4.7	0.8	( 3.1, 6.3)
COLORADO	2778	21.8	1.1	( 19.5, 24.0)
LOUISIANA	1645	13.8	1.0	( 11.9, 15.7)
MISSISSIPPI	2129	24.5	1.1	( 22.4, 26.7)
MISSOURI	4271	15.7	0.8	( 14.2, 17.3)
MONTANA	1798	27.0	1.2	( 24.6, 29.4)
NEBRASKA	2679	20.7	0.9	( 18.8, 22.5)
NEVADA	2185	14.3	1.1	( 12.1, 16.5)
NEW YORK	2503	7.6	0.6	( 6.4, 8.8)
OHIO	1600	16.4	1.3	( 13.9, 18.9)
OKLAHOMA	2941	16.9	0.9	( 15.3, 18.6)
PENNSYLVANIA	3479	14.1	0.7	( 12.7, 15.5)
TEXAS	4986	14.6	0.6	( 13.4, 15.8)
UTAH	3175	11.1	0.8	( 9.7, 12.6)
VIRGINIA	3495	15.3	0.9	( 13.5, 17.1)
WASHINGTON	3534	20.0	0.9	( 18.2, 21.8)
WEST VIRGINIA	2580	28.5	1.1	( 26.4, 30.5)
WYOMING	2402	32.2	1.1	( 30.0, 34.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 19 MEDIAN = 16.4 RANGE = 4.7-32.2

\* EVER USED OR TRIED PRODUCTS SUCH AS CHEWING TOBACCO OR SNUFF

June 20, 2000

TABLE 31.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER USED SMOKELESS TOBACCO\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALASKA	1001	44.1	2.3	( 39.7, 48.5)	*	1049	9.9	1.5	( 7.0, 12.8)	*
ARIZONA	712	8.2	1.5	( 5.2, 11.1)	*	1027	1.5	0.7	( 0.0, 2.9)	*
COLORADO	1263	38.6	2.0	( 34.7, 42.4)	*	1515	5.3	0.9	( 3.6, 6.9)	*
LOUISIANA	652	26.9	1.9	( 23.1, 30.6)	*	993	2.1	0.4	( 1.2, 3.0)	*
MISSISSIPPI	834	45.3	2.0	( 41.3, 49.2)	*	1295	6.2	0.7	( 4.8, 7.6)	*
MISSOURI	1719	31.6	1.5	( 28.6, 34.6)	*	2552	1.5	0.3	( 0.9, 2.0)	*
MONTANA	768	47.9	2.0	( 44.0, 51.9)	*	1030	7.1	0.9	( 5.3, 9.0)	*
NEBRASKA	1065	38.4	1.6	( 35.1, 41.6)	*	1614	4.4	0.6	( 3.2, 5.5)	*
NEVADA	1099	24.8	1.9	( 21.0, 28.6)	*	1086	3.7	0.8	( 2.2, 5.2)	*
NEW YORK	1012	13.0	1.2	( 10.8, 15.3)	*	1491	2.7	0.5	( 1.7, 3.7)	*
OHIO	623	31.9	2.3	( 27.4, 36.4)	*	977	2.4	0.6	( 1.3, 3.6)	*
OKLAHOMA	1141	32.7	1.6	( 29.5, 35.8)	*	1800	2.5	0.4	( 1.8, 3.3)	*
PENNSYLVANIA	1390	28.2	1.4	( 25.5, 30.9)	*	2089	1.6	0.3	( 1.0, 2.1)	*
TEXAS	2003	27.8	1.2	( 25.5, 30.1)	*	2983	2.1	0.4	( 1.4, 2.9)	*
UTAH	1364	20.7	1.5	( 17.8, 23.6)	*	1811	2.1	0.4	( 1.3, 2.9)	*
VIRGINIA	1405	29.7	1.7	( 26.3, 33.1)	*	2090	1.9	0.3	( 1.3, 2.6)	*
WASHINGTON	1568	35.3	1.6	( 32.2, 38.4)	*	1966	5.3	0.7	( 4.0, 6.7)	*
WEST VIRGINIA	1064	56.0	1.7	( 52.6, 59.3)	*	1516	3.9	0.6	( 2.8, 4.9)	*
WYOMING	1064	54.1	1.7	( 50.7, 57.4)	*	1338	10.9	1.1	( 8.8, 12.9)	*

SUMMARY STATISTICS:           NUMBER OF PARTICIPANTS = 19  
   MEDIAN                         = 31.9  
   RANGE                         = 8.2-56.0

NUMBER OF PARTICIPANTS = 19  
 MEDIAN                         = 2.7  
 RANGE                         = 1.5-10.9

\* EVER USED OR TRIED PRODUCTS SUCH AS CHEWING TOBACCO OR SNUFF

June 20, 2000

TABLE 32.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALASKA	2050	5.4	0.7	( 4.1, 6.6)
ARIZONA	1738	0.8	0.3	( 0.3, 1.3)
COLORADO	2776	3.8	0.5	( 2.7, 4.8)
LOUISIANA	1645	4.1	0.6	( 3.0, 5.2)
MISSISSIPPI	2126	6.1	0.6	( 4.9, 7.3)
MISSOURI	4271	3.9	0.4	( 3.1, 4.7)
MONTANA	1797	6.2	0.7	( 4.9, 7.5)
NEBRASKA	2670	4.5	0.5	( 3.6, 5.4)
NEVADA	2185	3.2	0.7	( 1.8, 4.6)
NEW YORK	2503	0.8	0.2	( 0.4, 1.2)
OHIO	1597	3.0	0.7	( 1.7, 4.3)
OKLAHOMA	2940	5.0	0.6	( 3.9, 6.1)
PENNSYLVANIA	3479	3.4	0.4	( 2.6, 4.2)
TEXAS	4986	3.2	0.3	( 2.6, 3.8)
UTAH	3174	1.8	0.3	( 1.2, 2.4)
VIRGINIA	3495	3.3	0.4	( 2.5, 4.0)
WASHINGTON	3534	2.7	0.3	( 2.1, 3.3)
WEST VIRGINIA	2580	8.6	0.7	( 7.2, 10.0)
WYOMING	2402	8.1	0.6	( 6.9, 9.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 19 MEDIAN = 3.8 RANGE = 0.8- 8.6

\* CURRENT USERS OF SMOKELESS TOBACCO

June 20, 2000

TABLE 32.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENT SMOKELESS TOBACCO USERS\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALASKA	1001	8.7	1.2	( 6.4, 11.0)	*	1049	1.7	0.4	( 0.8, 2.6)	*
ARIZONA	711	1.6	0.5	( 0.6, 2.6)	*	1027	0.0	0.0	( 0.0, 0.0)	*
COLORADO	1263	7.4	1.0	( 5.4, 9.4)	*	1513	0.2	0.2	( -0.2, 0.6)	*
LOUISIANA	652	8.5	1.2	( 6.2, 10.8)	*	993	0.1	0.1	( -0.0, 0.2)	*
MISSISSIPPI	833	11.2	1.2	( 8.8, 13.5)	*	1293	1.6	0.4	( 0.9, 2.3)	*
MISSOURI	1719	8.1	0.8	( 6.4, 9.7)	*	2552	0.2	0.1	( 0.0, 0.4)	*
MONTANA	767	12.5	1.3	( 9.9, 15.0)	*	1030	0.3	0.2	( -0.1, 0.6)	*
NEBRASKA	1062	9.2	1.0	( 7.3, 11.1)	*	1608	0.2	0.1	( -0.0, 0.4)	*
NEVADA	1099	6.4	1.4	( 3.7, 9.0)	*	1086	0.0	0.0	( 0.0, 0.0)	*
NEW YORK	1012	1.2	0.3	( 0.6, 1.9)	*	1491	0.4	0.2	( -0.0, 0.8)	*
OHIO	621	6.3	1.4	( 3.6, 9.0)	*	976	0.0	0.0	( 0.0, 0.0)	*
OKLAHOMA	1141	10.1	1.2	( 7.8, 12.3)	*	1799	0.3	0.2	( 0.0, 0.7)	*
PENNSYLVANIA	1390	7.1	0.8	( 5.5, 8.6)	*	2089	0.1	0.1	( -0.0, 0.2)	*
TEXAS	2003	6.6	0.6	( 5.4, 7.7)	*	2983	0.1	0.1	( -0.0, 0.1)	*
UTAH	1364	3.6	0.6	( 2.3, 4.8)	*	1810	0.1	0.0	( 0.0, 0.2)	*
VIRGINIA	1405	6.5	0.8	( 5.1, 8.0)	*	2090	0.2	0.1	( -0.0, 0.5)	*
WASHINGTON	1568	5.5	0.6	( 4.2, 6.7)	*	1966	0.2	0.1	( -0.0, 0.3)	*
WEST VIRGINIA	1064	17.5	1.4	( 14.8, 20.2)	*	1516	0.6	0.2	( 0.2, 1.1)	*
WYOMING	1064	16.0	1.2	( 13.6, 18.4)	*	1338	0.5	0.2	( 0.1, 0.8)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	19	NUMBER OF PARTICIPANTS =	19
	MEDIAN	= 7.4	MEDIAN	= 0.2
	RANGE	= 1.2-17.5	RANGE	= 0.0- 1.7

\* CURRENT USERS OF SMOKELESS TOBACCO

June 20, 2000

TABLE 33.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENTLY HAS ASTHMA\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ARIZONA	1738	5.5	0.8	( 3.9, 7.0)
ARKANSAS	2991	7.3	0.6	( 6.1, 8.4)
DISTRICT OF CO	1257	6.8	0.8	( 5.2, 8.3)
IDAHO	4937	7.9	0.5	( 6.9, 8.9)
IOWA	3499	7.7	0.6	( 6.6, 8.7)
LOUISIANA	1659	4.9	0.6	( 3.6, 6.1)
MAINE	1662	8.3	0.8	( 6.7, 9.8)
MISSISSIPPI	2137	5.0	0.5	( 3.9, 6.0)
MISSOURI	4220	7.3	0.6	( 6.2, 8.4)
NEBRASKA	2709	6.5	0.5	( 5.5, 7.4)
NEW HAMPSHIRE	1205	8.8	0.9	( 7.0, 10.6)
PENNSYLVANIA	3479	6.8	0.5	( 5.8, 7.8)
SOUTH CAROLINA	3395	5.8	0.5	( 4.9, 6.6)
UTAH	3180	7.5	0.6	( 6.4, 8.7)
VERMONT	3091	8.0	0.6	( 6.9, 9.1)
WASHINGTON	3525	7.0	0.5	( 6.2, 7.9)
WISCONSIN	2166	7.3	0.7	( 6.0, 8.6)
WYOMING	2395	7.1	0.6	( 6.0, 8.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 18 MEDIAN = 7.2 RANGE = 4.9- 8.8

\* RESPONDENTS TOLD BY DOCTOR THEY HAD ASTHMA

June 30, 2000

TABLE 33.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR CURRENTLY HAS ASTHMA\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ARIZONA	710	4.5	1.2	( 2.2,	6.8)	* 1028	6.4	1.1	( 4.2,	8.6)
ARKANSAS	1165	5.3	0.8	( 3.7,	6.9)	* 1826	9.0	0.8	( 7.5,	10.6)
DISTRICT OF CO	456	5.0	1.1	( 2.8,	7.2)	* 801	8.3	1.1	( 6.2,	10.4)
IDAHO	2084	6.8	0.8	( 5.3,	8.4)	* 2853	8.9	0.7	( 7.6,	10.2)
IOWA	1375	6.2	0.8	( 4.7,	7.6)	* 2124	9.1	0.8	( 7.5,	10.6)
LOUISIANA	659	3.9	1.0	( 1.9,	5.8)	* 1000	5.8	0.8	( 4.2,	7.4)
MAINE	732	7.3	1.1	( 5.0,	9.5)	* 930	9.2	1.1	( 7.0,	11.3)
MISSISSIPPI	835	4.3	0.8	( 2.6,	5.9)	* 1302	5.6	0.7	( 4.3,	6.9)
MISSOURI	1696	6.4	0.9	( 4.6,	8.2)	* 2524	8.1	0.7	( 6.7,	9.5)
NEBRASKA	1074	5.2	0.7	( 3.7,	6.6)	* 1635	7.6	0.7	( 6.2,	9.0)
NEW HAMPSHIRE	485	5.2	1.1	( 3.0,	7.4)	* 720	12.1	1.4	( 9.3,	14.9)
PENNSYLVANIA	1388	5.3	0.8	( 3.8,	6.9)	* 2091	8.1	0.7	( 6.8,	9.3)
SOUTH CAROLINA	1365	4.4	0.7	( 3.1,	5.7)	* 2030	7.0	0.6	( 5.8,	8.2)
UTAH	1368	5.9	0.8	( 4.3,	7.5)	* 1812	9.1	0.9	( 7.4,	10.7)
VERMONT	1245	6.2	0.8	( 4.7,	7.7)	* 1846	9.7	0.9	( 8.0,	11.4)
WASHINGTON	1570	4.8	0.6	( 3.7,	5.8)	* 1955	9.2	0.7	( 7.9,	10.6)
WISCONSIN	912	5.8	0.9	( 4.1,	7.5)	* 1254	8.7	1.0	( 6.7,	10.7)
WYOMING	1061	5.2	0.7	( 3.9,	6.6)	* 1334	8.9	0.9	( 7.2,	10.6)

SUMMARY STATISTICS:

NUMBER OF PARTICIPANTS = 18  
 MEDIAN = 5.3  
 RANGE = 3.9- 7.3

NUMBER OF PARTICIPANTS = 18  
 MEDIAN = 8.8  
 RANGE = 5.6-12.1

\* RESPONDENTS TOLD BY DOCTOR THEY HAD ASTHMA

June 30, 2000

TABLE 34.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD A HEART ATTACK OR MYOCARDIAL INFARCTION\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2085	5.1	0.5	( 4.1, 6.0)
ARIZONA	1741	5.1	0.7	( 3.8, 6.5)
DISTRICT OF CO	1255	3.1	0.6	( 2.0, 4.1)
GEORGIA	2233	4.5	0.5	( 3.5, 5.6)
INDIANA	2425	6.2	0.9	( 4.4, 8.0)
IOWA	3504	4.4	0.4	( 3.7, 5.1)
KENTUCKY	7454	5.5	0.4	( 4.7, 6.2)
LOUISIANA	1655	4.4	0.6	( 3.3, 5.5)
MAINE	1663	5.3	0.6	( 4.2, 6.4)
MISSISSIPPI	2130	5.3	0.5	( 4.3, 6.4)
MONTANA	1798	3.8	0.5	( 2.8, 4.8)
NEW MEXICO	3403	3.5	0.3	( 2.8, 4.1)
NEW YORK	2542	3.6	0.4	( 2.8, 4.5)
NORTH CAROLINA	2371	4.8	0.5	( 3.8, 5.8)
NORTH DAKOTA	1966	4.5	0.5	( 3.5, 5.5)
OHIO	1612	5.3	0.6	( 4.1, 6.6)
OKLAHOMA	2943	5.3	0.6	( 4.1, 6.4)
SOUTH CAROLINA	3372	4.4	0.4	( 3.6, 5.2)
VIRGINIA	3423	3.6	0.4	( 2.7, 4.4)
WISCONSIN	2165	4.2	0.5	( 3.3, 5.2)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 20 MEDIAN = 4.5 RANGE = 3.1- 6.2  
 \* TOLD BY A DOCTOR THEY HAD A HEART ATTACK OR MYOCARDIAL INFARCTION June 20, 2000

TABLE 34.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 94  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR HAD A HEART ATTACK OR MYOCARDIAL INFARCTION\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	*
ALABAMA	854	5.9	0.8	( 4.3, 7.4)	*	1231	4.4	0.6	( 3.2, 5.5)	*
ARIZONA	712	5.4	1.0	( 3.4, 7.4)	*	1029	4.9	1.0	( 3.0, 6.7)	*
DISTRICT OF CO	455	3.9	0.9	( 2.1, 5.7)	*	800	2.4	0.6	( 1.1, 3.6)	*
GEORGIA	877	5.8	0.9	( 4.0, 7.5)	*	1356	3.4	0.6	( 2.3, 4.6)	*
INDIANA	1012	8.6	1.7	( 5.3, 11.8)	*	1413	4.1	0.9	( 2.4, 5.8)	*
IOWA	1373	5.8	0.6	( 4.6, 7.1)	*	2131	3.1	0.4	( 2.4, 3.8)	*
KENTUCKY	2712	7.1	0.7	( 5.8, 8.5)	*	4742	4.0	0.3	( 3.3, 4.6)	*
LOUISIANA	657	5.3	0.9	( 3.5, 7.2)	*	998	3.5	0.6	( 2.3, 4.7)	*
MAINE	734	7.1	1.0	( 5.2, 9.0)	*	929	3.7	0.6	( 2.5, 4.9)	*
MISSISSIPPI	835	7.1	0.9	( 5.3, 8.9)	*	1295	3.8	0.6	( 2.7, 4.9)	*
MONTANA	768	5.0	0.8	( 3.4, 6.7)	*	1030	2.7	0.5	( 1.7, 3.6)	*
NEW MEXICO	1496	4.4	0.5	( 3.4, 5.5)	*	1907	2.5	0.4	( 1.8, 3.2)	*
NEW YORK	1025	4.6	0.7	( 3.2, 6.0)	*	1517	2.8	0.5	( 1.8, 3.8)	*
NORTH CAROLINA	941	6.8	1.0	( 4.9, 8.7)	*	1430	3.0	0.5	( 2.1, 3.9)	*
NORTH DAKOTA	855	5.8	0.9	( 4.2, 7.5)	*	1111	3.2	0.5	( 2.2, 4.2)	*
OHIO	625	7.7	1.2	( 5.4, 10.0)	*	987	3.2	0.6	( 2.1, 4.4)	*
OKLAHOMA	1143	7.2	1.2	( 5.0, 9.5)	*	1800	3.5	0.5	( 2.6, 4.3)	*
SOUTH CAROLINA	1352	5.8	0.7	( 4.5, 7.2)	*	2020	3.2	0.4	( 2.3, 4.0)	*
VIRGINIA	1387	4.8	0.8	( 3.3, 6.3)	*	2036	2.4	0.4	( 1.7, 3.1)	*
WISCONSIN	918	5.3	0.8	( 3.7, 6.9)	*	1247	3.2	0.6	( 2.1, 4.3)	*

SUMMARY STATISTICS:	NUMBER OF PARTICIPANTS =	20	NUMBER OF PARTICIPANTS =	20
	MEDIAN	= 5.8	MEDIAN	= 3.2
	RANGE	= 3.9- 8.6	RANGE	= 2.4- 4.9

\* TOLD BY A DOCTOR THEY HAD A HEART ATTACK OR MYOCARDIAL INFARCTION

June 20, 2000

TABLE 35.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD YOU HAD A STROKE\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	2086	2.9	0.4	( 2.2, 3.6)
ARIZONA	1741	2.6	0.5	( 1.7, 3.5)
DISTRICT OF CO	1253	2.0	0.4	( 1.3, 2.7)
GEORGIA	2234	2.6	0.4	( 1.8, 3.4)
INDIANA	2428	2.4	0.6	( 1.3, 3.5)
IOWA	3506	2.3	0.3	( 1.9, 2.8)
KENTUCKY	7456	3.4	0.3	( 2.8, 4.0)
LOUISIANA	1655	1.9	0.3	( 1.3, 2.5)
MAINE	1662	2.3	0.4	( 1.5, 3.1)
MISSISSIPPI	2139	2.8	0.4	( 2.0, 3.5)
MONTANA	1796	2.2	0.4	( 1.6, 2.9)
NEW MEXICO	3406	2.0	0.3	( 1.5, 2.5)
NEW YORK	2538	2.3	0.4	( 1.6, 3.0)
NORTH CAROLINA	2370	3.3	0.4	( 2.5, 4.0)
NORTH DAKOTA	1965	2.3	0.4	( 1.6, 3.0)
OHIO	1614	2.4	0.4	( 1.6, 3.1)
OKLAHOMA	2944	2.4	0.3	( 1.8, 3.0)
SOUTH CAROLINA	3376	2.1	0.3	( 1.6, 2.7)
VIRGINIA	3430	1.7	0.2	( 1.2, 2.2)
WISCONSIN	2165	1.9	0.3	( 1.3, 2.5)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 20 MEDIAN = 2.3 RANGE = 1.7- 3.4

\* RESPONDENTS TOLD BY A DOCTOR THEY HAD A STROKE

June 20, 2000

TABLE 35.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR EVER TOLD YOU HAD A STROKE\* -BY SEX  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	*-----MALES-----*					*-----FEMALES-----*				
	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL		SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL	
ALABAMA	853	2.6	0.5	( 1.6,	3.7)	1233	3.2	0.5	( 2.2,	4.2)
ARIZONA	712	1.8	0.6	( 0.6,	2.9)	1029	3.4	0.7	( 2.0,	4.7)
DISTRICT OF CO	455	1.5	0.5	( 0.5,	2.5)	798	2.4	0.5	( 1.4,	3.4)
GEORGIA	876	2.8	0.7	( 1.5,	4.2)	1358	2.4	0.5	( 1.5,	3.3)
INDIANA	1013	2.5	0.9	( 0.7,	4.4)	1415	2.3	0.7	( 0.9,	3.7)
IOWA	1373	2.2	0.4	( 1.5,	2.9)	2133	2.5	0.3	( 1.8,	3.1)
KENTUCKY	2712	3.5	0.6	( 2.4,	4.6)	4744	3.3	0.3	( 2.7,	3.8)
LOUISIANA	658	1.8	0.5	( 0.9,	2.8)	997	1.9	0.4	( 1.1,	2.7)
MAINE	733	1.8	0.5	( 0.9,	2.7)	929	2.7	0.7	( 1.4,	4.0)
MISSISSIPPI	836	2.8	0.6	( 1.6,	4.0)	1303	2.7	0.5	( 1.8,	3.7)
MONTANA	767	2.0	0.5	( 1.0,	3.0)	1029	2.5	0.5	( 1.5,	3.4)
NEW MEXICO	1497	2.4	0.5	( 1.4,	3.3)	1909	1.7	0.3	( 1.1,	2.3)
NEW YORK	1023	2.6	0.6	( 1.4,	3.7)	1515	2.1	0.4	( 1.3,	2.9)
NORTH CAROLINA	943	3.5	0.6	( 2.3,	4.6)	1427	3.1	0.5	( 2.1,	4.0)
NORTH DAKOTA	855	2.4	0.6	( 1.3,	3.4)	1110	2.3	0.5	( 1.4,	3.2)
OHIO	626	2.8	0.7	( 1.5,	4.1)	988	1.9	0.5	( 1.0,	2.8)
OKLAHOMA	1143	2.2	0.5	( 1.3,	3.1)	1801	2.6	0.4	( 1.9,	3.4)
SOUTH CAROLINA	1352	2.6	0.5	( 1.7,	3.5)	2024	1.7	0.3	( 1.1,	2.3)
VIRGINIA	1389	1.1	0.2	( 0.6,	1.5)	2041	2.3	0.4	( 1.5,	3.1)
WISCONSIN	918	1.8	0.5	( 0.9,	2.7)	1247	2.1	0.4	( 1.2,	2.9)

SUMMARY STATISTICS:           NUMBER OF PARTICIPANTS = 20  
   MEDIAN                               = 2.4  
   RANGE                               = 1.1- 3.5

NUMBER OF PARTICIPANTS = 20  
 MEDIAN                               = 2.4  
 RANGE                                 = 1.7- 3.4

\* RESPONDENTS TOLD BY A DOCTOR THEY HAD A STROKE

June 20, 2000

TABLE 36.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM, WOMEN AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1230	37.6	1.6	( 34.5, 40.8)
ALASKA	1043	44.3	2.2	( 40.1, 48.6)
ARIZONA	1024	34.2	2.2	( 29.8, 38.5)
ARKANSAS	1822	41.8	1.3	( 39.3, 44.3)
COLORADO	1669	42.2	1.7	( 38.8, 45.5)
CONNECTICUT	2097	33.5	1.4	( 30.7, 36.3)
DELAWARE	1515	33.8	1.8	( 30.2, 37.4)
DISTRICT OF CO	796	40.7	1.9	( 37.1, 44.3)
FLORIDA	3104	31.0	1.0	( 29.0, 32.9)
GEORGIA	1369	40.0	1.6	( 36.9, 43.0)
HAWAII	1150	38.2	1.9	( 34.5, 41.9)
IDAHO	2852	43.6	1.1	( 41.5, 45.6)
ILLINOIS	1792	39.1	1.3	( 36.6, 41.6)
INDIANA	1394	39.6	2.2	( 35.2, 43.9)
IOWA	2159	38.6	1.3	( 36.2, 41.1)
KANSAS	2262	38.1	1.2	( 35.8, 40.4)
KENTUCKY	4754	38.6	1.0	( 36.6, 40.6)
LOUISIANA	995	41.7	1.8	( 38.2, 45.2)
MAINE	932	37.6	1.8	( 34.1, 41.1)
MARYLAND	2347	36.5	1.2	( 34.1, 39.0)
MASSACHUSETTS	2917	37.5	1.2	( 35.2, 39.7)
MICHIGAN	1468	36.5	1.4	( 33.7, 39.2)
MINNESOTA	2901	42.4	1.0	( 40.4, 44.4)
MISSISSIPPI	1309	41.2	1.5	( 38.2, 44.2)
MISSOURI	2531	40.0	1.3	( 37.4, 42.6)
MONTANA	1026	39.5	1.7	( 36.0, 42.9)
NEBRASKA	1624	40.0	1.4	( 37.3, 42.6)
NEVADA	1077	33.7	2.0	( 29.7, 37.7)
NEW HAMPSHIRE	725	41.0	2.1	( 36.9, 45.2)
NEW JERSEY	1658	39.5	1.5	( 36.7, 42.4)
NEW MEXICO	1944	42.7	1.3	( 40.2, 45.2)
NEW YORK	1519	39.4	1.5	( 36.5, 42.2)
NORTH CAROLINA	1453	38.8	1.5	( 35.8, 41.8)
NORTH DAKOTA	1112	39.5	1.6	( 36.3, 42.6)
OHIO	994	37.5	1.9	( 33.7, 41.2)
OKLAHOMA	1798	42.7	1.3	( 40.1, 45.3)
OREGON	1030	35.6	1.7	( 32.3, 38.8)
PENNSYLVANIA	2094	36.8	1.2	( 34.4, 39.2)
RHODE ISLAND	2388	36.0	1.2	( 33.7, 38.2)
SOUTH CAROLINA	2031	38.1	1.2	( 35.6, 40.5)
SOUTH DAKOTA	2443	41.0	1.1	( 38.8, 43.2)
TENNESSEE	1946	38.2	1.3	( 35.7, 40.7)
TEXAS	2961	41.2	1.1	( 38.9, 43.4)
UTAH	1812	44.9	1.6	( 41.8, 48.0)
VERMONT	1860	41.3	1.4	( 38.7, 44.0)
VIRGINIA	2075	40.4	1.6	( 37.3, 43.6)
WASHINGTON	1993	39.1	1.3	( 36.6, 41.5)
WEST VIRGINIA	1515	39.6	1.4	( 36.8, 42.4)
WISCONSIN	1238	39.8	1.6	( 36.6, 43.0)
WYOMING	1332	41.6	1.4	( 38.9, 44.4)
PUERTO RICO	1921	45.8	1.3	( 43.2, 48.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 39.5 RANGE = 31.0-45.8

\* NEVER HAD A MAMMOGRAM

June 20, 2000

TABLE 36.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO MAMMOGRAM, WOMEN AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	38.1	37.6	-0.4
ALASKA	44.3	44.3	0.0
ARIZONA	43.8	34.2	-9.6
ARKANSAS	40.7	41.8	1.1
COLORADO	40.0	42.2	2.2
CONNECTICUT	37.3	33.5	-3.8
DELAWARE	35.8	33.8	-2.1
DISTRICT OF CO	36.7	40.7	4.1
FLORIDA	35.0	31.0	-4.0
GEORGIA	42.2	40.0	-2.3
HAWAII	38.0	38.2	0.2
IDAHO	43.7	43.6	-0.1
ILLINOIS	41.5	39.1	-2.4
INDIANA	42.4	39.6	-2.8
IOWA	39.5	38.6	-0.9
KANSAS	41.0	38.1	-2.9
KENTUCKY	40.2	38.6	-1.6
LOUISIANA	42.3	41.7	-0.6
MAINE	40.5	37.6	-2.9
MARYLAND	36.3	36.5	0.2
MASSACHUSETTS	36.9	37.5	0.6
MICHIGAN	38.3	36.5	-1.8
MINNESOTA	51.4	42.4	-9.0
MISSISSIPPI	43.7	41.2	-2.4
MISSOURI	40.4	40.0	-0.4
MONTANA	42.3	39.5	-2.8
NEBRASKA	44.0	40.0	-4.1
NEVADA	40.0	33.7	-6.4
NEW HAMPSHIRE	42.7	41.0	-1.6
NEW JERSEY	40.1	39.5	-0.5
NEW MEXICO	44.4	42.7	-1.7
NEW YORK	38.2	39.4	1.1
NORTH CAROLINA	40.2	38.8	-1.4
NORTH DAKOTA	41.8	39.5	-2.3
OHIO	41.9	37.5	-4.4
OKLAHOMA	47.8	42.7	-5.1
OREGON	40.4	35.6	-4.8
PENNSYLVANIA	37.6	36.8	-0.7
RHODE ISLAND	38.6	36.0	-2.7
SOUTH CAROLINA	39.6	38.1	-1.6
SOUTH DAKOTA	41.9	41.0	-0.9
TENNESSEE	38.7	38.2	-0.5
TEXAS	44.0	41.2	-2.8
UTAH	47.5	44.9	-2.6
VERMONT	43.9	41.3	-2.6
VIRGINIA	41.5	40.4	-1.1
WASHINGTON	39.5	39.1	-0.4
WEST VIRGINIA	40.1	39.6	-0.5
WISCONSIN	41.1	39.8	-1.3
WYOMING	43.5	41.6	-1.9
PUERTO RICO	47.4	45.8	-1.5

\* NEVER HAD A MAMMOGRAM

June 20, 2000

TABLE 37.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM, WOMEN AGE 40 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	780	14.0	1.4	( 11.3, 16.7)
ALASKA	580	17.4	3.8	( 10.0, 24.7)
ARIZONA	679	12.0	2.0	( 8.0, 16.0)
ARKANSAS	1177	18.9	1.4	( 16.2, 21.5)
COLORADO	1019	15.6	1.7	( 12.3, 18.8)
CONNECTICUT	1285	8.1	1.0	( 6.3, 10.0)
DELAWARE	911	8.8	2.1	( 4.8, 12.8)
DISTRICT OF CO	444	10.5	1.6	( 7.3, 13.7)
FLORIDA	2053	9.5	0.7	( 8.1, 11.0)
GEORGIA	837	14.5	1.4	( 11.9, 17.2)
HAWAII	738	11.5	1.5	( 8.6, 14.3)
IDAHO	1718	16.0	1.0	( 14.1, 17.9)
ILLINOIS	1022	13.4	1.1	( 11.2, 15.6)
INDIANA	884	14.6	2.0	( 10.7, 18.5)
IOWA	1431	13.9	1.0	( 11.9, 15.9)
KANSAS	1465	14.1	1.0	( 12.0, 16.1)
KENTUCKY	3161	14.4	0.8	( 12.8, 15.9)
LOUISIANA	584	17.9	1.7	( 14.4, 21.3)
MAINE	623	9.1	1.2	( 6.8, 11.5)
MARYLAND	1368	10.7	1.0	( 8.7, 12.7)
MASSACHUSETTS	1759	9.8	0.9	( 8.1, 11.4)
MICHIGAN	882	10.5	1.2	( 8.2, 12.9)
MINNESOTA	1674	16.0	1.0	( 14.1, 17.9)
MISSISSIPPI	810	19.4	1.5	( 16.5, 22.3)
MISSOURI	1622	16.9	1.3	( 14.4, 19.4)
MONTANA	702	14.2	1.4	( 11.4, 17.0)
NEBRASKA	1067	15.7	1.2	( 13.3, 18.1)
NEVADA	643	11.1	1.7	( 7.8, 14.3)
NEW HAMPSHIRE	455	11.8	1.7	( 8.5, 15.0)
NEW JERSEY	1058	15.4	1.4	( 12.7, 18.0)
NEW MEXICO	1231	16.4	1.2	( 14.1, 18.8)
NEW YORK	914	14.2	1.4	( 11.5, 16.8)
NORTH CAROLINA	947	13.3	1.2	( 10.8, 15.7)
NORTH DAKOTA	722	13.4	1.4	( 10.7, 16.1)
OHIO	661	11.6	1.4	( 8.8, 14.3)
OKLAHOMA	1166	16.9	1.2	( 14.5, 19.3)
OREGON	651	8.5	1.2	( 6.1, 11.0)
PENNSYLVANIA	1349	11.8	1.0	( 9.9, 13.8)
RHODE ISLAND	1467	9.0	0.9	( 7.3, 10.7)
SOUTH CAROLINA	1175	12.6	1.1	( 10.5, 14.8)
SOUTH DAKOTA	1550	13.7	1.0	( 11.8, 15.6)
TENNESSEE	1211	15.7	1.2	( 13.4, 18.1)
TEXAS	1687	13.8	1.0	( 11.9, 15.7)
UTAH	1007	13.9	1.6	( 10.7, 17.1)
VERMONT	1241	12.2	1.0	( 10.2, 14.2)
VIRGINIA	1279	13.7	1.3	( 11.2, 16.2)
WASHINGTON	1143	10.0	1.0	( 8.1, 11.9)
WEST VIRGINIA	1067	17.4	1.3	( 14.9, 19.9)
WISCONSIN	758	11.2	1.2	( 8.8, 13.6)
WYOMING	826	16.7	1.6	( 13.5, 19.9)
PUERTO RICO	1231	19.2	1.3	( 16.7, 21.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 13.8 RANGE = 8.1-19.4

\* NEVER HAD A MAMMOGRAM

June 20, 2000

TABLE 37.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO MAMMOGRAM, WOMEN AGE 40 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.3	14.0	-0.3
ALASKA	10.6	17.4	6.7
ARIZONA	18.0	12.0	-6.0
ARKANSAS	17.5	18.9	1.4
COLORADO	11.0	15.6	4.5
CONNECTICUT	12.9	8.1	-4.8
DELAWARE	9.9	8.8	-1.1
DISTRICT OF CO	8.3	10.5	2.2
FLORIDA	13.0	9.5	-3.5
GEORGIA	15.6	14.5	-1.1
HAWAII	11.1	11.5	0.4
IDAHO	18.2	16.0	-2.2
ILLINOIS	16.5	13.4	-3.1
INDIANA	15.9	14.6	-1.3
IOWA	14.9	13.9	-1.1
KANSAS	16.1	14.1	-2.1
KENTUCKY	17.2	14.4	-2.8
LOUISIANA	19.4	17.9	-1.6
MAINE	13.1	9.1	-4.0
MARYLAND	10.9	10.7	-0.2
MASSACHUSETTS	8.4	9.8	1.4
MICHIGAN	10.7	10.5	-0.2
MINNESOTA	27.5	16.0	-11.5
MISSISSIPPI	19.5	19.4	-0.1
MISSOURI	15.3	16.9	1.6
MONTANA	17.3	14.2	-3.1
NEBRASKA	18.6	15.7	-2.9
NEVADA	13.9	11.1	-2.8
NEW HAMPSHIRE	15.1	11.8	-3.4
NEW JERSEY	16.8	15.4	-1.4
NEW MEXICO	16.0	16.4	0.5
NEW YORK	13.3	14.2	0.8
NORTH CAROLINA	15.2	13.3	-1.9
NORTH DAKOTA	14.6	13.4	-1.2
OHIO	15.4	11.6	-3.9
OKLAHOMA	24.9	16.9	-7.9
OREGON	13.8	8.5	-5.2
PENNSYLVANIA	15.2	11.8	-3.3
RHODE ISLAND	11.7	9.0	-2.7
SOUTH CAROLINA	15.8	12.6	-3.1
SOUTH DAKOTA	15.9	13.7	-2.2
TENNESSEE	15.0	15.7	0.7
TEXAS	18.1	13.8	-4.3
UTAH	15.9	13.9	-2.0
VERMONT	15.8	12.2	-3.6
VIRGINIA	13.6	13.7	0.1
WASHINGTON	11.0	10.0	-1.0
WEST VIRGINIA	17.0	17.4	0.4
WISCONSIN	14.6	11.2	-3.4
WYOMING	17.5	16.7	-0.8
PUERTO RICO	22.7	19.2	-3.6

\* NEVER HAD A MAMMOGRAM

TABLE 38.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM, WOMEN AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	568	23.8	1.9	( 20.0, 27.5)
ALASKA	280	22.9	3.8	( 15.5, 30.3)
ARIZONA	497	15.5	2.6	( 10.4, 20.6)
ARKANSAS	815	29.8	1.9	( 26.2, 33.5)
COLORADO	640	24.8	2.4	( 20.1, 29.6)
CONNECTICUT	814	13.9	1.5	( 10.9, 16.8)
DELAWARE	593	14.0	1.9	( 10.3, 17.7)
DISTRICT OF CO	320	16.7	2.3	( 12.3, 21.2)
FLORIDA	1455	16.6	1.1	( 14.4, 18.8)
GEORGIA	525	24.8	2.1	( 20.8, 28.8)
HAWAII	463	20.7	2.7	( 15.5, 25.9)
IDAHO	1131	31.4	1.6	( 28.3, 34.5)
ILLINOIS	658	23.7	1.8	( 20.2, 27.3)
INDIANA	542	31.3	3.4	( 24.7, 37.8)
IOWA	1016	25.5	1.5	( 22.6, 28.5)
KANSAS	1002	20.1	1.4	( 17.4, 22.7)
KENTUCKY	2269	26.5	1.2	( 24.1, 28.9)
LOUISIANA	395	30.4	2.6	( 25.3, 35.5)
MAINE	435	18.2	1.9	( 14.4, 22.0)
MARYLAND	862	16.9	1.6	( 13.8, 20.0)
MASSACHUSETTS	1145	16.9	1.3	( 14.3, 19.5)
MICHIGAN	587	16.7	1.8	( 13.2, 20.2)
MINNESOTA	1091	26.6	1.4	( 23.9, 29.4)
MISSISSIPPI	530	32.9	2.2	( 28.6, 37.3)
MISSOURI	1144	25.7	1.7	( 22.4, 29.0)
MONTANA	480	28.8	2.2	( 24.5, 33.0)
NEBRASKA	767	25.8	1.7	( 22.5, 29.2)
NEVADA	400	25.0	3.4	( 18.3, 31.7)
NEW HAMPSHIRE	275	15.6	2.3	( 11.2, 20.0)
NEW JERSEY	686	22.2	1.9	( 18.6, 25.9)
NEW MEXICO	812	27.2	2.0	( 23.4, 31.1)
NEW YORK	585	21.5	1.9	( 17.7, 25.2)
NORTH CAROLINA	652	19.8	1.8	( 16.4, 23.3)
NORTH DAKOTA	530	23.3	1.9	( 19.6, 27.1)
OHIO	494	23.2	2.1	( 19.1, 27.4)
OKLAHOMA	832	28.2	1.7	( 24.9, 31.5)
OREGON	454	22.6	2.1	( 18.4, 26.8)
PENNSYLVANIA	866	22.1	1.5	( 19.1, 25.1)
RHODE ISLAND	957	14.9	1.4	( 12.2, 17.5)
SOUTH CAROLINA	764	18.7	1.5	( 15.7, 21.6)
SOUTH DAKOTA	1062	25.1	1.4	( 22.3, 27.8)
TENNESSEE	827	24.9	1.7	( 21.6, 28.2)
TEXAS	1082	25.1	1.5	( 22.2, 28.0)
UTAH	654	27.7	2.4	( 23.0, 32.4)
VERMONT	794	22.3	1.6	( 19.2, 25.4)
VIRGINIA	794	26.0	2.4	( 21.2, 30.7)
WASHINGTON	746	24.5	1.8	( 21.0, 27.9)
WEST VIRGINIA	748	28.0	1.8	( 24.4, 31.6)
WISCONSIN	491	25.2	2.1	( 21.0, 29.4)
WYOMING	511	29.2	2.2	( 24.9, 33.4)
PUERTO RICO	876	29.5	1.7	( 26.2, 32.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 24.5 RANGE = 13.9-32.9

\* DID NOT HAVE A MAMMOGRAM IN PAST TWO YEARS

June 20, 2000

TABLE 38.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO MAMMOGRAM, WOMEN AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	23.9	23.8	-0.1
ALASKA	20.9	22.9	2.0
ARIZONA	17.4	15.5	-1.9
ARKANSAS	34.1	29.8	-4.3
COLORADO	22.6	24.8	2.2
CONNECTICUT	22.0	13.9	-8.2
DELAWARE	18.8	14.0	-4.8
DISTRICT OF CO	10.6	16.7	6.1
FLORIDA	22.0	16.6	-5.3
GEORGIA	27.5	24.8	-2.6
HAWAII	22.9	20.7	-2.2
IDAHO	29.1	31.4	2.3
ILLINOIS	28.6	23.7	-4.9
INDIANA	28.5	31.3	2.8
IOWA	28.6	25.5	-3.0
KANSAS	24.7	20.1	-4.6
KENTUCKY	31.6	26.5	-5.1
LOUISIANA	29.8	30.4	0.6
MAINE	22.9	18.2	-4.7
MARYLAND	19.2	16.9	-2.3
MASSACHUSETTS	15.8	16.9	1.1
MICHIGAN	21.1	16.7	-4.3
MINNESOTA	35.1	26.6	-8.4
MISSISSIPPI	32.1	32.9	0.9
MISSOURI	26.0	25.7	-0.4
MONTANA	27.7	28.8	1.1
NEBRASKA	30.3	25.8	-4.4
NEVADA	28.4	25.0	-3.3
NEW HAMPSHIRE	22.9	15.6	-7.2
NEW JERSEY	24.5	22.2	-2.3
NEW MEXICO	25.0	27.2	2.3
NEW YORK	22.9	21.5	-1.4
NORTH CAROLINA	21.0	19.8	-1.1
NORTH DAKOTA	25.1	23.3	-1.8
OHIO	26.6	23.2	-3.4
OKLAHOMA	29.1	28.2	-0.9
OREGON	21.3	22.6	1.3
PENNSYLVANIA	22.7	22.1	-0.6
RHODE ISLAND	18.8	14.9	-4.0
SOUTH CAROLINA	22.8	18.7	-4.2
SOUTH DAKOTA	25.0	25.1	0.1
TENNESSEE	24.8	24.9	0.1
TEXAS	27.3	25.1	-2.2
UTAH	28.8	27.7	-1.1
VERMONT	23.0	22.3	-0.6
VIRGINIA	22.2	26.0	3.8
WASHINGTON	21.6	24.5	2.9
WEST VIRGINIA	25.0	28.0	3.0
WISCONSIN	26.0	25.2	-0.8
WYOMING	31.3	29.2	-2.2
PUERTO RICO	38.7	29.5	-9.2

\* DID NOT HAVE A MAMMOGRAM IN PAST TWO YEARS

TABLE 39.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CLINICAL BREAST EXAM, WOMEN AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1234	12.3	1.1	( 10.1, 14.5)
ALASKA	1041	9.6	1.4	( 6.9, 12.3)
ARIZONA	1028	18.7	2.0	( 14.7, 22.7)
ARKANSAS	1828	10.5	0.8	( 8.8, 12.1)
COLORADO	1676	9.0	1.0	( 7.0, 10.9)
CONNECTICUT	2118	13.0	1.0	( 11.0, 14.9)
DELAWARE	1525	9.1	1.0	( 7.2, 11.1)
DISTRICT OF CO	802	8.8	1.2	( 6.5, 11.1)
FLORIDA	3106	12.4	0.7	( 11.0, 13.8)
GEORGIA	1371	10.4	1.0	( 8.5, 12.4)
HAWAII	1156	13.4	1.2	( 10.9, 15.8)
IDAHO	2849	9.2	0.6	( 8.0, 10.4)
ILLINOIS	1788	13.5	0.9	( 11.7, 15.3)
INDIANA	1410	10.6	1.3	( 8.1, 13.2)
IOWA	2154	8.6	0.7	( 7.2, 10.0)
KANSAS	2270	14.8	0.9	( 13.1, 16.5)
KENTUCKY	4759	11.4	0.6	( 10.2, 12.6)
LOUISIANA	1003	15.1	1.3	( 12.6, 17.6)
MAINE	929	5.0	0.8	( 3.4, 6.5)
MARYLAND	2352	10.5	0.8	( 8.9, 12.0)
MASSACHUSETTS	2956	12.3	0.8	( 10.7, 13.9)
MICHIGAN	1471	9.0	0.9	( 7.3, 10.8)
MINNESOTA	2896	6.3	0.5	( 5.3, 7.3)
MISSISSIPPI	1309	12.9	1.1	( 10.8, 15.0)
MISSOURI	2537	13.6	0.9	( 11.8, 15.4)
MONTANA	1030	6.2	0.9	( 4.5, 7.9)
NEBRASKA	1629	10.0	0.8	( 8.3, 11.6)
NEVADA	1079	11.1	1.4	( 8.4, 13.8)
NEW HAMPSHIRE	726	6.2	1.1	( 4.0, 8.4)
NEW JERSEY	1676	18.2	1.2	( 15.8, 20.5)
NEW MEXICO	1940	11.4	0.9	( 9.5, 13.2)
NEW YORK	1544	16.3	1.2	( 14.0, 18.6)
NORTH CAROLINA	1459	7.4	1.0	( 5.5, 9.4)
NORTH DAKOTA	1117	7.8	0.9	( 6.1, 9.5)
OHIO	997	8.9	1.2	( 6.6, 11.1)
OKLAHOMA	1796	14.3	1.0	( 12.3, 16.4)
OREGON	1035	9.3	1.1	( 7.1, 11.4)
PENNSYLVANIA	2111	12.8	0.9	( 11.1, 14.5)
RHODE ISLAND	2405	14.8	0.9	( 13.0, 16.6)
SOUTH CAROLINA	2050	13.9	0.9	( 12.2, 15.7)
SOUTH DAKOTA	2448	7.5	0.6	( 6.3, 8.7)
TENNESSEE	1961	14.0	0.9	( 12.1, 15.8)
TEXAS	2979	15.8	0.8	( 14.2, 17.5)
UTAH	1816	8.5	1.0	( 6.5, 10.4)
VERMONT	1875	10.3	0.8	( 8.7, 12.0)
VIRGINIA	2079	11.1	1.1	( 8.9, 13.3)
WASHINGTON	1983	8.7	0.8	( 7.2, 10.1)
WEST VIRGINIA	1519	11.5	1.0	( 9.6, 13.4)
WISCONSIN	1241	8.3	1.0	( 6.2, 10.3)
WYOMING	1336	10.8	1.1	( 8.7, 13.0)
PUERTO RICO	1916	23.7	1.2	( 21.3, 26.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 10.8 RANGE = 5.0-23.7

\* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

June 20, 2000

TABLE 39.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO CLINICAL BREAST EXAM, WOMEN AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	13.3	12.3	-1.0
ALASKA	8.4	9.6	1.2
ARIZONA	18.7	18.7	0.0
ARKANSAS	17.5	10.5	-7.0
COLORADO	8.1	9.0	0.8
CONNECTICUT	13.7	13.0	-0.8
DELAWARE	5.5	9.1	3.6
DISTRICT OF CO	6.1	8.8	2.7
FLORIDA	14.8	12.4	-2.4
GEORGIA	11.2	10.4	-0.8
HAWAII	10.9	13.4	2.5
IDAHO	8.1	9.2	1.1
ILLINOIS	14.3	13.5	-0.9
INDIANA	12.6	10.6	-2.0
IOWA	9.6	8.6	-1.0
KANSAS	13.3	14.8	1.5
KENTUCKY	10.1	11.4	1.3
LOUISIANA	13.2	15.1	1.9
MAINE	9.0	5.0	-4.0
MARYLAND	10.9	10.5	-0.4
MASSACHUSETTS	11.2	12.3	1.1
MICHIGAN	11.2	9.0	-2.2
MINNESOTA	7.1	6.3	-0.8
MISSISSIPPI	14.9	12.9	-2.0
MISSOURI	11.1	13.6	2.5
MONTANA	7.2	6.2	-1.0
NEBRASKA	9.7	10.0	0.3
NEVADA	11.1	11.1	0.0
NEW HAMPSHIRE	8.9	6.2	-2.7
NEW JERSEY	17.3	18.2	0.9
NEW MEXICO	12.8	11.4	-1.5
NEW YORK	15.3	16.3	1.0
NORTH CAROLINA	8.0	7.4	-0.6
NORTH DAKOTA	8.3	7.8	-0.5
OHIO	8.9	8.9	0.0
OKLAHOMA	20.5	14.3	-6.2
OREGON	9.3	9.3	-0.1
PENNSYLVANIA	14.9	12.8	-2.1
RHODE ISLAND	14.1	14.8	0.7
SOUTH CAROLINA	15.7	13.9	-1.7
SOUTH DAKOTA	8.5	7.5	-1.0
TENNESSEE	11.6	14.0	2.4
TEXAS	17.4	15.8	-1.5
UTAH	12.8	8.5	-4.4
VERMONT	12.8	10.3	-2.5
VIRGINIA	11.3	11.1	-0.3
WASHINGTON	7.7	8.7	0.9
WEST VIRGINIA	8.8	11.5	2.8
WISCONSIN	8.7	8.3	-0.4
WYOMING	8.2	10.8	2.6
PUERTO RICO	25.5	23.7	-1.7

\* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

June 20, 2000

TABLE 40.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CLINICAL BREAST EXAM, WOMEN AGE 40 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	776	13.1	1.5	( 10.3, 16.0)
ALASKA	575	6.9	1.6	( 3.7, 10.0)
ARIZONA	677	11.9	2.0	( 8.1, 15.7)
ARKANSAS	1176	11.0	1.0	( 8.9, 13.0)
COLORADO	1015	9.3	1.3	( 6.7, 11.9)
CONNECTICUT	1277	12.0	1.2	( 9.6, 14.4)
DELAWARE	909	8.2	1.2	( 5.8, 10.5)
DISTRICT OF CO	444	7.5	1.3	( 5.0, 10.1)
FLORIDA	2044	10.8	0.8	( 9.2, 12.4)
GEORGIA	832	11.0	1.2	( 8.6, 13.4)
HAWAII	732	11.1	1.5	( 8.2, 14.1)
IDAHO	1712	7.2	0.7	( 5.9, 8.5)
ILLINOIS	1018	10.6	1.0	( 8.5, 12.6)
INDIANA	880	9.7	1.7	( 6.3, 13.1)
IOWA	1422	8.3	0.8	( 6.8, 9.8)
KANSAS	1460	11.8	0.9	( 10.0, 13.6)
KENTUCKY	3157	10.8	0.7	( 9.5, 12.0)
LOUISIANA	585	13.1	1.5	( 10.1, 16.1)
MAINE	622	4.4	0.9	( 2.7, 6.1)
MARYLAND	1358	8.9	0.9	( 7.1, 10.7)
MASSACHUSETTS	1745	9.2	0.8	( 7.6, 10.8)
MICHIGAN	877	5.3	0.9	( 3.6, 7.1)
MINNESOTA	1669	5.5	0.6	( 4.4, 6.6)
MISSISSIPPI	806	13.6	1.5	( 10.6, 16.5)
MISSOURI	1614	13.0	1.1	( 10.8, 15.1)
MONTANA	702	4.9	0.9	( 3.1, 6.6)
NEBRASKA	1062	8.8	0.9	( 7.0, 10.5)
NEVADA	642	9.3	1.6	( 6.2, 12.4)
NEW HAMPSHIRE	453	5.8	1.3	( 3.3, 8.3)
NEW JERSEY	1053	15.5	1.4	( 12.8, 18.2)
NEW MEXICO	1222	10.4	1.2	( 7.9, 12.8)
NEW YORK	911	12.7	1.3	( 10.2, 15.2)
NORTH CAROLINA	942	5.8	0.9	( 4.1, 7.6)
NORTH DAKOTA	723	6.4	0.9	( 4.7, 8.2)
OHIO	658	7.8	1.1	( 5.6, 10.1)
OKLAHOMA	1160	10.9	1.0	( 8.9, 12.8)
OREGON	647	8.6	1.1	( 6.3, 10.8)
PENNSYLVANIA	1346	12.5	1.1	( 10.4, 14.6)
RHODE ISLAND	1458	13.0	1.1	( 10.9, 15.1)
SOUTH CAROLINA	1174	12.6	1.1	( 10.5, 14.6)
SOUTH DAKOTA	1544	7.3	0.7	( 6.0, 8.7)
TENNESSEE	1211	12.2	1.1	( 10.1, 14.3)
TEXAS	1684	12.6	0.9	( 10.7, 14.4)
UTAH	1005	4.6	0.8	( 2.9, 6.2)
VERMONT	1238	8.5	0.8	( 6.9, 10.1)
VIRGINIA	1271	9.5	1.1	( 7.4, 11.5)
WASHINGTON	1136	7.7	0.9	( 5.9, 9.5)
WEST VIRGINIA	1066	11.0	1.0	( 9.0, 13.0)
WISCONSIN	752	6.6	1.0	( 4.6, 8.6)
WYOMING	823	9.1	1.1	( 6.9, 11.2)
PUERTO RICO	1226	13.8	1.1	( 11.6, 15.9)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 9.5 RANGE = 4.4-15.5

\* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

June 20, 2000

TABLE 40.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO CLINICAL BREAST EXAM, WOMEN AGE 40 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	13.2	13.1	-0.1
ALASKA	7.1	6.9	-0.2
ARIZONA	16.8	11.9	-4.9
ARKANSAS	14.6	11.0	-3.7
COLORADO	5.4	9.3	3.8
CONNECTICUT	12.4	12.0	-0.4
DELAWARE	5.1	8.2	3.1
DISTRICT OF CO	4.6	7.5	2.9
FLORIDA	13.5	10.8	-2.8
GEORGIA	11.6	11.0	-0.6
HAWAII	8.5	11.1	2.7
IDAHO	7.1	7.2	0.1
ILLINOIS	13.6	10.6	-3.0
INDIANA	11.7	9.7	-2.0
IOWA	9.3	8.3	-1.0
KANSAS	12.0	11.8	-0.2
KENTUCKY	9.9	10.8	0.8
LOUISIANA	13.8	13.1	-0.7
MAINE	6.8	4.4	-2.4
MARYLAND	11.8	8.9	-2.9
MASSACHUSETTS	10.2	9.2	-1.0
MICHIGAN	11.2	5.3	-5.9
MINNESOTA	7.6	5.5	-2.2
MISSISSIPPI	15.4	13.6	-1.8
MISSOURI	10.8	13.0	2.2
MONTANA	4.8	4.9	0.0
NEBRASKA	10.0	8.8	-1.3
NEVADA	12.7	9.3	-3.4
NEW HAMPSHIRE	9.0	5.8	-3.2
NEW JERSEY	15.5	15.5	0.1
NEW MEXICO	10.9	10.4	-0.5
NEW YORK	13.9	12.7	-1.2
NORTH CAROLINA	7.2	5.8	-1.4
NORTH DAKOTA	7.5	6.4	-1.1
OHIO	7.8	7.8	0.1
OKLAHOMA	15.8	10.9	-4.9
OREGON	7.5	8.6	1.0
PENNSYLVANIA	15.1	12.5	-2.6
RHODE ISLAND	12.4	13.0	0.7
SOUTH CAROLINA	14.8	12.6	-2.2
SOUTH DAKOTA	6.0	7.3	1.4
TENNESSEE	9.4	12.2	2.9
TEXAS	14.5	12.6	-1.9
UTAH	9.1	4.6	-4.6
VERMONT	12.9	8.5	-4.4
VIRGINIA	11.1	9.5	-1.6
WASHINGTON	5.6	7.7	2.1
WEST VIRGINIA	7.5	11.0	3.5
WISCONSIN	7.0	6.6	-0.4
WYOMING	8.5	9.1	0.6
PUERTO RICO	18.7	13.8	-4.9

\* NEVER HAD EXAM BY A DOCTOR, NURSE OR OTHER HEALTH PROFESSIONAL

June 20, 2000

TABLE 41.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO CLINICAL BREAST EXAM, WOMEN AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	555	26.8	2.1	( 22.8, 30.9)
ALASKA	276	20.1	3.4	( 13.4, 26.7)
ARIZONA	494	18.5	2.8	( 13.1, 23.9)
ARKANSAS	812	29.9	1.8	( 26.4, 33.5)
COLORADO	633	25.2	2.4	( 20.4, 29.9)
CONNECTICUT	802	16.8	1.7	( 13.5, 20.0)
DELAWARE	594	17.8	2.1	( 13.7, 21.8)
DISTRICT OF CO	316	15.9	2.2	( 11.7, 20.2)
FLORIDA	1438	19.8	1.2	( 17.4, 22.1)
GEORGIA	513	26.1	2.1	( 22.0, 30.2)
HAWAII	455	23.7	2.8	( 18.3, 29.1)
IDAHO	1125	26.5	1.5	( 23.6, 29.5)
ILLINOIS	650	22.5	1.8	( 19.0, 26.1)
INDIANA	536	31.1	3.4	( 24.4, 37.7)
IOWA	1003	23.8	1.5	( 20.9, 26.7)
KANSAS	997	22.9	1.4	( 20.1, 25.7)
KENTUCKY	2260	24.4	1.2	( 22.1, 26.7)
LOUISIANA	396	27.3	2.5	( 22.4, 32.2)
MAINE	430	11.5	1.6	( 8.3, 14.7)
MARYLAND	852	19.9	1.7	( 16.5, 23.3)
MASSACHUSETTS	1125	16.9	1.4	( 14.3, 19.6)
MICHIGAN	580	17.5	1.8	( 14.0, 21.0)
MINNESOTA	1082	21.9	1.3	( 19.4, 24.5)
MISSISSIPPI	521	29.3	2.2	( 25.1, 33.6)
MISSOURI	1121	26.5	1.7	( 23.1, 29.9)
MONTANA	474	23.2	2.1	( 19.2, 27.3)
NEBRASKA	761	21.2	1.6	( 18.1, 24.3)
NEVADA	396	28.9	4.6	( 19.8, 38.0)
NEW HAMPSHIRE	272	14.4	2.3	( 9.9, 18.9)
NEW JERSEY	680	23.2	1.9	( 19.4, 26.9)
NEW MEXICO	800	26.2	2.0	( 22.4, 30.1)
NEW YORK	585	21.2	1.9	( 17.5, 24.8)
NORTH CAROLINA	648	16.7	1.6	( 13.5, 19.9)
NORTH DAKOTA	526	20.8	1.8	( 17.2, 24.4)
OHIO	492	21.2	2.1	( 17.2, 25.3)
OKLAHOMA	828	25.3	1.6	( 22.1, 28.4)
OREGON	449	23.8	2.2	( 19.6, 28.0)
PENNSYLVANIA	859	23.9	1.6	( 20.7, 27.1)
RHODE ISLAND	953	20.2	1.5	( 17.2, 23.2)
SOUTH CAROLINA	756	21.7	1.6	( 18.5, 24.9)
SOUTH DAKOTA	1053	22.5	1.4	( 19.8, 25.3)
TENNESSEE	817	23.7	1.6	( 20.5, 26.8)
TEXAS	1076	26.6	1.5	( 23.7, 29.6)
UTAH	649	21.7	2.1	( 17.5, 25.9)
VERMONT	793	22.5	1.6	( 19.3, 25.6)
VIRGINIA	780	23.7	2.4	( 19.0, 28.4)
WASHINGTON	735	24.1	1.8	( 20.6, 27.5)
WEST VIRGINIA	747	27.9	1.8	( 24.4, 31.4)
WISCONSIN	482	22.0	2.1	( 17.9, 26.0)
WYOMING	507	26.5	2.1	( 22.4, 30.6)
PUERTO RICO	868	29.6	1.8	( 26.2, 33.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 23.2 RANGE = 11.5-31.1

\* DID NOT HAVE EXAM IN PAST TWO YEARS

June 20, 2000

TABLE 41.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO CLINICAL BREAST EXAM, WOMEN AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	22.0	26.8	4.8
ALASKA	20.2	20.1	-0.2
ARIZONA	17.3	18.5	1.2
ARKANSAS	34.7	29.9	-4.8
COLORADO	22.0	25.2	3.2
CONNECTICUT	22.2	16.8	-5.4
DELAWARE	16.5	17.8	1.2
DISTRICT OF CO	9.8	15.9	6.1
FLORIDA	24.0	19.8	-4.2
GEORGIA	27.1	26.1	-1.0
HAWAII	20.4	23.7	3.3
IDAHO	26.6	26.5	-0.1
ILLINOIS	27.6	22.5	-5.1
INDIANA	26.6	31.1	4.5
IOWA	27.7	23.8	-4.0
KANSAS	24.2	22.9	-1.3
KENTUCKY	27.5	24.4	-3.1
LOUISIANA	28.5	27.3	-1.1
MAINE	18.5	11.5	-7.0
MARYLAND	22.5	19.9	-2.6
MASSACHUSETTS	20.6	16.9	-3.7
MICHIGAN	24.6	17.5	-7.1
MINNESOTA	27.8	21.9	-5.9
MISSISSIPPI	29.9	29.3	-0.6
MISSOURI	26.0	26.5	0.6
MONTANA	22.2	23.2	1.0
NEBRASKA	30.1	21.2	-8.9
NEVADA	29.5	28.9	-0.7
NEW HAMPSHIRE	20.1	14.4	-5.7
NEW JERSEY	28.3	23.2	-5.1
NEW MEXICO	27.1	26.2	-0.9
NEW YORK	23.5	21.2	-2.3
NORTH CAROLINA	18.4	16.7	-1.7
NORTH DAKOTA	23.3	20.8	-2.5
OHIO	24.7	21.2	-3.5
OKLAHOMA	25.5	25.3	-0.3
OREGON	25.2	23.8	-1.4
PENNSYLVANIA	27.0	23.9	-3.1
RHODE ISLAND	22.2	20.2	-2.0
SOUTH CAROLINA	24.8	21.7	-3.1
SOUTH DAKOTA	19.2	22.5	3.4
TENNESSEE	20.5	23.7	3.1
TEXAS	29.3	26.6	-2.6
UTAH	30.1	21.7	-8.5
VERMONT	24.7	22.5	-2.3
VIRGINIA	24.4	23.7	-0.7
WASHINGTON	22.3	24.1	1.8
WEST VIRGINIA	22.8	27.9	5.1
WISCONSIN	25.2	22.0	-3.3
WYOMING	32.2	26.5	-5.8
PUERTO RICO	37.5	29.6	-7.9

\* DID NOT HAVE EXAM IN PAST TWO YEARS

June 20, 2000

TABLE 42.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 40 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	775	22.9	1.8	( 19.4, 26.3)
ALASKA	575	20.0	3.8	( 12.5, 27.5)
ARIZONA	677	17.9	2.3	( 13.5, 22.3)
ARKANSAS	1174	23.5	1.4	( 20.7, 26.3)
COLORADO	1015	20.4	1.8	( 16.9, 24.0)
CONNECTICUT	1276	16.7	1.4	( 14.0, 19.4)
DELAWARE	908	15.0	2.2	( 10.7, 19.4)
DISTRICT OF CO	443	15.6	1.9	( 11.9, 19.2)
FLORIDA	2043	16.5	1.0	( 14.6, 18.4)
GEORGIA	830	21.4	1.6	( 18.3, 24.5)
HAWAII	732	18.8	1.9	( 15.1, 22.4)
IDAHO	1711	20.4	1.1	( 18.3, 22.5)
ILLINOIS	1018	20.8	1.4	( 18.1, 23.5)
INDIANA	879	19.8	2.2	( 15.4, 24.2)
IOWA	1422	18.3	1.1	( 16.0, 20.5)
KANSAS	1457	21.0	1.2	( 18.7, 23.4)
KENTUCKY	3151	19.0	0.9	( 17.2, 20.7)
LOUISIANA	584	24.0	1.9	( 20.2, 27.8)
MAINE	622	11.3	1.3	( 8.7, 14.0)
MARYLAND	1358	15.7	1.2	( 13.3, 18.0)
MASSACHUSETTS	1745	15.3	1.0	( 13.2, 17.3)
MICHIGAN	877	14.4	1.3	( 11.8, 16.9)
MINNESOTA	1669	18.3	1.0	( 16.3, 20.3)
MISSISSIPPI	806	26.7	1.8	( 23.2, 30.3)
MISSOURI	1612	23.2	1.4	( 20.4, 26.0)
MONTANA	702	16.9	1.6	( 13.8, 20.0)
NEBRASKA	1060	19.1	1.3	( 16.5, 21.7)
NEVADA	641	17.5	2.1	( 13.4, 21.6)
NEW HAMPSHIRE	452	14.8	1.9	( 11.1, 18.4)
NEW JERSEY	1052	24.4	1.6	( 21.2, 27.5)
NEW MEXICO	1222	21.3	1.5	( 18.3, 24.2)
NEW YORK	911	21.1	1.5	( 18.1, 24.1)
NORTH CAROLINA	941	15.6	1.3	( 12.9, 18.2)
NORTH DAKOTA	722	16.4	1.5	( 13.5, 19.2)
OHIO	658	15.2	1.6	( 12.1, 18.3)
OKLAHOMA	1160	23.0	1.4	( 20.3, 25.7)
OREGON	647	14.7	1.5	( 11.7, 17.7)
PENNSYLVANIA	1345	19.7	1.2	( 17.3, 22.2)
RHODE ISLAND	1458	18.4	1.2	( 16.1, 20.7)
SOUTH CAROLINA	1172	21.3	1.4	( 18.6, 23.9)
SOUTH DAKOTA	1542	17.8	1.1	( 15.7, 19.9)
TENNESSEE	1208	22.4	1.4	( 19.7, 25.1)
TEXAS	1682	21.6	1.2	( 19.3, 23.8)
UTAH	1004	16.4	1.7	( 13.1, 19.8)
VERMONT	1237	17.1	1.2	( 14.8, 19.3)
VIRGINIA	1271	18.3	1.4	( 15.5, 21.0)
WASHINGTON	1135	15.8	1.2	( 13.5, 18.2)
WEST VIRGINIA	1065	21.6	1.4	( 18.9, 24.3)
WISCONSIN	750	15.3	1.4	( 12.5, 18.1)
WYOMING	823	21.5	1.7	( 18.1, 24.9)
PUERTO RICO	1225	26.0	1.4	( 23.2, 28.7)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 18.8 RANGE = 11.3-26.7

\* NEVER HAD A MAMMOGRAM AND BREAST EXAM

June 20, 2000

TABLE 42.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 40 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	21.5	22.9	1.4
ALASKA	15.0	20.0	5.0
ARIZONA	21.2	17.9	-3.3
ARKANSAS	25.2	23.5	-1.7
COLORADO	15.0	20.4	5.5
CONNECTICUT	20.4	16.7	-3.7
DELAWARE	12.1	15.0	2.9
DISTRICT OF CO	11.6	15.6	3.9
FLORIDA	21.2	16.5	-4.6
GEORGIA	21.9	21.4	-0.5
HAWAII	16.7	18.8	2.1
IDAHO	22.8	20.4	-2.4
ILLINOIS	24.0	20.8	-3.2
INDIANA	23.0	19.8	-3.1
IOWA	19.2	18.3	-0.9
KANSAS	22.4	21.0	-1.4
KENTUCKY	19.8	19.0	-0.9
LOUISIANA	26.3	24.0	-2.2
MAINE	15.4	11.3	-4.1
MARYLAND	19.3	15.7	-3.6
MASSACHUSETTS	15.2	15.3	0.0
MICHIGAN	18.3	14.4	-4.0
MINNESOTA	30.4	18.3	-12.1
MISSISSIPPI	26.9	26.7	-0.2
MISSOURI	21.4	23.2	1.8
MONTANA	18.7	16.9	-1.8
NEBRASKA	24.4	19.1	-5.3
NEVADA	21.8	17.5	-4.3
NEW HAMPSHIRE	19.5	14.8	-4.7
NEW JERSEY	25.4	24.4	-1.0
NEW MEXICO	22.2	21.3	-1.0
NEW YORK	21.4	21.1	-0.3
NORTH CAROLINA	18.4	15.6	-2.8
NORTH DAKOTA	18.7	16.4	-2.3
OHIO	18.3	15.2	-3.1
OKLAHOMA	31.5	23.0	-8.5
OREGON	19.0	14.7	-4.3
PENNSYLVANIA	24.0	19.7	-4.3
RHODE ISLAND	19.5	18.4	-1.1
SOUTH CAROLINA	24.3	21.3	-3.0
SOUTH DAKOTA	18.8	17.8	-0.9
TENNESSEE	19.5	22.4	2.8
TEXAS	25.9	21.6	-4.3
UTAH	22.1	16.4	-5.6
VERMONT	24.0	17.1	-6.9
VIRGINIA	21.3	18.3	-3.0
WASHINGTON	14.9	15.8	0.9
WEST VIRGINIA	19.6	21.6	2.1
WISCONSIN	17.7	15.3	-2.4
WYOMING	22.2	21.5	-0.7
PUERTO RICO	31.4	26.0	-5.4

\* NEVER HAD A MAMMOGRAM AND BREAST EXAM

June 20, 2000

TABLE 43.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	558	34.7	2.2	( 30.3, 39.0)
ALASKA	275	26.9	3.9	( 19.1, 34.6)
ARIZONA	494	23.1	3.0	( 17.3, 28.9)
ARKANSAS	809	38.9	1.9	( 35.1, 42.7)
COLORADO	633	33.9	2.7	( 28.7, 39.1)
CONNECTICUT	803	22.2	1.9	( 18.6, 25.9)
DELAWARE	594	22.8	2.3	( 18.4, 27.3)
DISTRICT OF CO	314	21.8	2.5	( 16.9, 26.7)
FLORIDA	1438	23.9	1.3	( 21.4, 26.5)
GEORGIA	512	33.8	2.3	( 29.4, 38.3)
HAWAII	454	30.5	3.1	( 24.5, 36.5)
IDAHO	1122	36.8	1.7	( 33.6, 40.0)
ILLINOIS	650	32.1	2.0	( 28.2, 36.1)
INDIANA	536	39.3	3.5	( 32.5, 46.2)
IOWA	1002	31.3	1.6	( 28.2, 34.5)
KANSAS	995	28.5	1.6	( 25.5, 31.6)
KENTUCKY	2254	31.7	1.3	( 29.2, 34.2)
LOUISIANA	394	35.0	2.7	( 29.8, 40.2)
MAINE	430	19.9	2.0	( 15.9, 23.9)
MARYLAND	854	24.4	1.9	( 20.7, 28.0)
MASSACHUSETTS	1124	22.3	1.5	( 19.4, 25.3)
MICHIGAN	579	23.5	2.0	( 19.6, 27.4)
MINNESOTA	1082	31.0	1.5	( 28.1, 34.0)
MISSISSIPPI	523	40.7	2.4	( 36.0, 45.3)
MISSOURI	1125	35.1	1.9	( 31.4, 38.8)
MONTANA	475	33.6	2.3	( 29.1, 38.1)
NEBRASKA	762	31.1	1.8	( 27.6, 34.6)
NEVADA	396	35.3	4.6	( 26.2, 44.3)
NEW HAMPSHIRE	274	19.6	2.6	( 14.5, 24.6)
NEW JERSEY	680	32.0	2.1	( 27.9, 36.1)
NEW MEXICO	800	33.7	2.1	( 29.7, 37.7)
NEW YORK	582	29.0	2.1	( 24.9, 33.1)
NORTH CAROLINA	643	23.1	1.9	( 19.5, 26.7)
NORTH DAKOTA	524	28.7	2.1	( 24.6, 32.8)
OHIO	492	29.0	2.3	( 24.4, 33.5)
OKLAHOMA	827	35.3	1.8	( 31.8, 38.8)
OREGON	448	30.7	2.4	( 26.1, 35.4)
PENNSYLVANIA	856	32.1	1.8	( 28.6, 35.6)
RHODE ISLAND	949	25.2	1.6	( 22.0, 28.4)
SOUTH CAROLINA	752	27.4	1.8	( 23.9, 30.8)
SOUTH DAKOTA	1052	31.8	1.5	( 28.9, 34.8)
TENNESSEE	821	32.1	1.8	( 28.6, 35.6)
TEXAS	1072	34.7	1.6	( 31.5, 37.9)
UTAH	647	32.3	2.5	( 27.5, 37.2)
VERMONT	789	30.0	1.8	( 26.6, 33.4)
VIRGINIA	781	32.0	2.5	( 27.1, 36.9)
WASHINGTON	733	32.6	2.0	( 28.7, 36.4)
WEST VIRGINIA	743	34.3	1.9	( 30.6, 38.1)
WISCONSIN	482	31.2	2.3	( 26.7, 35.7)
WYOMING	507	35.9	2.3	( 31.5, 40.4)
PUERTO RICO	866	38.8	1.9	( 35.2, 42.4)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 31.7 RANGE = 19.6-40.7 YEAR 2000 OBJ 16.11  
 \* DID NOT HAVE MAMMOGRAM AND BREAST EXAM IN PAST TWO YEARS June 20, 2000

TABLE 43.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO MAMMOGRAM AND BREAST EXAM, WOMEN AGE 50 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	30.1	34.7	4.5
ALASKA	25.9	26.9	0.9
ARIZONA	19.0	23.1	4.1
ARKANSAS	43.0	38.9	-4.1
COLORADO	29.9	33.9	4.0
CONNECTICUT	30.6	22.2	-8.3
DELAWARE	22.9	22.8	-0.1
DISTRICT OF CO	14.8	21.8	7.0
FLORIDA	31.1	23.9	-7.2
GEORGIA	35.9	33.8	-2.0
HAWAII	30.3	30.5	0.2
IDAHO	36.5	36.8	0.3
ILLINOIS	37.9	32.1	-5.8
INDIANA	36.1	39.3	3.2
IOWA	36.0	31.3	-4.6
KANSAS	33.7	28.5	-5.1
KENTUCKY	36.3	31.7	-4.6
LOUISIANA	36.5	35.0	-1.5
MAINE	26.4	19.9	-6.5
MARYLAND	29.3	24.4	-4.9
MASSACHUSETTS	25.2	22.3	-2.9
MICHIGAN	30.2	23.5	-6.8
MINNESOTA	40.0	31.0	-9.0
MISSISSIPPI	39.9	40.7	0.8
MISSOURI	33.4	35.1	1.7
MONTANA	31.0	33.6	2.6
NEBRASKA	40.5	31.1	-9.4
NEVADA	37.6	35.3	-2.3
NEW HAMPSHIRE	28.4	19.6	-8.8
NEW JERSEY	34.9	32.0	-2.9
NEW MEXICO	33.7	33.7	-0.0
NEW YORK	31.3	29.0	-2.3
NORTH CAROLINA	25.9	23.1	-2.8
NORTH DAKOTA	31.7	28.7	-3.0
OHIO	31.9	29.0	-2.9
OKLAHOMA	37.5	35.3	-2.2
OREGON	32.2	30.7	-1.4
PENNSYLVANIA	33.5	32.1	-1.4
RHODE ISLAND	29.8	25.2	-4.6
SOUTH CAROLINA	33.1	27.4	-5.7
SOUTH DAKOTA	30.5	31.8	1.3
TENNESSEE	29.1	32.1	2.9
TEXAS	37.0	34.7	-2.3
UTAH	38.5	32.3	-6.1
VERMONT	32.9	30.0	-2.9
VIRGINIA	31.9	32.0	0.1
WASHINGTON	29.8	32.6	2.8
WEST VIRGINIA	31.1	34.3	3.3
WISCONSIN	33.2	31.2	-2.0
WYOMING	39.5	35.9	-3.6
PUERTO RICO	47.2	38.8	-8.4

\* DID NOT HAVE MAMMOGRAM AND BREAST EXAM IN PAST TWO YEARS

June 20, 2000

TABLE 44.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PAP SMEAR, WOMEN AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	1228	3.5	0.6	( 2.3, 4.7)
ALASKA	1041	2.9	0.8	( 1.3, 4.6)
ARIZONA	1019	8.1	1.3	( 5.6, 10.6)
ARKANSAS	1820	4.1	0.5	( 3.1, 5.2)
COLORADO	1666	4.3	0.7	( 2.9, 5.7)
CONNECTICUT	2092	5.7	0.7	( 4.3, 7.1)
DELAWARE	1512	3.8	0.8	( 2.3, 5.3)
DISTRICT OF CO	794	5.1	1.0	( 3.1, 7.1)
FLORIDA	3095	5.1	0.5	( 4.0, 6.1)
GEORGIA	1370	3.2	0.6	( 2.0, 4.3)
HAWAII	1148	6.6	1.0	( 4.6, 8.5)
IDAHO	2847	4.8	0.5	( 3.8, 5.9)
ILLINOIS	1787	5.9	0.7	( 4.5, 7.2)
INDIANA	1388	4.7	1.0	( 2.7, 6.7)
IOWA	2155	4.7	0.6	( 3.6, 5.7)
KANSAS	2252	4.9	0.6	( 3.8, 6.0)
KENTUCKY	4749	7.1	0.6	( 6.0, 8.2)
LOUISIANA	996	8.1	1.1	( 6.0, 10.2)
MAINE	929	2.4	0.6	( 1.2, 3.5)
MARYLAND	2338	5.9	0.7	( 4.5, 7.2)
MASSACHUSETTS	2902	7.5	0.7	( 6.1, 8.9)
MICHIGAN	1465	5.3	0.7	( 3.9, 6.7)
MINNESOTA	2898	4.6	0.5	( 3.6, 5.5)
MISSISSIPPI	1306	4.8	0.7	( 3.5, 6.1)
MISSOURI	2528	5.1	0.6	( 4.0, 6.2)
MONTANA	1027	3.0	0.7	( 1.6, 4.4)
NEBRASKA	1614	5.7	0.7	( 4.3, 7.0)
NEVADA	1074	4.9	1.0	( 3.0, 6.8)
NEW HAMPSHIRE	719	4.1	1.0	( 2.2, 6.0)
NEW JERSEY	1646	8.9	1.0	( 7.1, 10.8)
NEW MEXICO	1938	4.9	0.6	( 3.7, 6.1)
NEW YORK	1511	8.9	0.9	( 7.2, 10.6)
NORTH CAROLINA	1450	2.8	0.5	( 1.9, 3.7)
NORTH DAKOTA	1110	3.6	0.6	( 2.4, 4.9)
OHIO	992	5.2	0.9	( 3.3, 7.0)
OKLAHOMA	1798	4.0	0.6	( 3.0, 5.1)
OREGON	1029	4.5	0.8	( 3.0, 6.0)
PENNSYLVANIA	2086	6.0	0.6	( 4.8, 7.3)
RHODE ISLAND	2384	7.3	0.7	( 5.9, 8.7)
SOUTH CAROLINA	2025	4.5	0.6	( 3.4, 5.6)
SOUTH DAKOTA	2430	4.6	0.5	( 3.6, 5.6)
TENNESSEE	1945	7.9	0.8	( 6.4, 9.4)
TEXAS	2958	7.1	0.6	( 5.9, 8.3)
UTAH	1807	6.7	1.0	( 4.8, 8.6)
VERMONT	1855	4.9	0.7	( 3.5, 6.2)
VIRGINIA	2064	5.7	1.0	( 3.8, 7.7)
WASHINGTON	1981	3.5	0.5	( 2.6, 4.5)
WEST VIRGINIA	1510	6.3	0.8	( 4.8, 7.8)
WISCONSIN	1234	4.3	0.7	( 3.0, 5.7)
WYOMING	1331	3.1	0.5	( 2.1, 4.1)
PUERTO RICO	1912	14.6	1.1	( 12.4, 16.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 4.9 RANGE = 2.4-14.6 YEAR 2000 OBJ 16.11  
 \* NEVER HAD PAP SMEAR June 20, 2000

TABLE 44.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999 NO PAP SMEAR, WOMEN AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	4.6	3.5	-1.1
ALASKA	3.1	2.9	-0.1
ARIZONA	13.1	8.1	-5.0
ARKANSAS	7.0	4.1	-2.9
COLORADO	4.2	4.3	0.1
CONNECTICUT	7.7	5.7	-2.0
DELAWARE	4.6	3.8	-0.8
DISTRICT OF CO	3.0	5.1	2.2
FLORIDA	4.6	5.1	0.4
GEORGIA	4.2	3.2	-1.0
HAWAII	4.8	6.6	1.7
IDAHO	5.3	4.8	-0.4
ILLINOIS	5.2	5.9	0.6
INDIANA	5.9	4.7	-1.2
IOWA	5.2	4.7	-0.5
KANSAS	6.2	4.9	-1.3
KENTUCKY	5.4	7.1	1.7
LOUISIANA	7.3	8.1	0.8
MAINE	5.3	2.4	-3.0
MARYLAND	5.0	5.9	0.9
MASSACHUSETTS	6.6	7.5	0.9
MICHIGAN	4.5	5.3	0.8
MINNESOTA	5.5	4.6	-0.9
MISSISSIPPI	6.2	4.8	-1.3
MISSOURI	4.5	5.1	0.6
MONTANA	3.6	3.0	-0.6
NEBRASKA	5.3	5.7	0.4
NEVADA	4.9	4.9	0.0
NEW HAMPSHIRE	5.6	4.1	-1.5
NEW JERSEY	9.9	8.9	-1.0
NEW MEXICO	6.0	4.9	-1.1
NEW YORK	6.8	8.9	2.1
NORTH CAROLINA	4.1	2.8	-1.4
NORTH DAKOTA	4.3	3.6	-0.6
OHIO	5.3	5.2	-0.1
OKLAHOMA	7.4	4.0	-3.4
OREGON	5.6	4.5	-1.1
PENNSYLVANIA	6.7	6.0	-0.6
RHODE ISLAND	9.3	7.3	-1.9
SOUTH CAROLINA	6.0	4.5	-1.5
SOUTH DAKOTA	5.8	4.6	-1.2
TENNESSEE	7.2	7.9	0.7
TEXAS	7.1	7.1	0.0
UTAH	8.6	6.7	-1.9
VERMONT	6.5	4.9	-1.7
VIRGINIA	5.1	5.7	0.7
WASHINGTON	4.4	3.5	-0.9
WEST VIRGINIA	6.3	6.3	0.0
WISCONSIN	5.0	4.3	-0.6
WYOMING	3.0	3.1	0.1
PUERTO RICO	16.4	14.6	-1.8

\* NEVER HAD PAP SMEAR

TABLE 45.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 115  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	801	3.3	0.7	( 1.9, 4.8)
ALASKA	854	2.5	0.7	( 1.0, 3.9)
ARIZONA	720	9.1	1.7	( 5.8, 12.4)
ARKANSAS	1259	4.4	0.7	( 3.1, 5.7)
COLORADO	1268	3.8	0.8	( 2.2, 5.3)
CONNECTICUT	1770	5.7	0.8	( 4.2, 7.3)
DELAWARE	1248	4.4	0.9	( 2.6, 6.2)
DISTRICT OF CO	663	5.0	1.1	( 2.8, 7.1)
FLORIDA	2284	5.2	0.6	( 4.1, 6.4)
GEORGIA	993	3.4	0.8	( 1.9, 4.9)
HAWAII	954	6.4	1.0	( 4.5, 8.4)
IDAHO	2067	5.5	0.7	( 4.2, 6.8)
ILLINOIS	1454	6.0	0.8	( 4.5, 7.5)
INDIANA	1066	4.9	1.2	( 2.6, 7.1)
IOWA	1660	4.6	0.6	( 3.4, 5.9)
KANSAS	1672	5.5	0.7	( 4.2, 6.8)
KENTUCKY	3370	6.8	0.7	( 5.5, 8.1)
LOUISIANA	687	7.2	1.2	( 4.9, 9.6)
MAINE	722	2.1	0.7	( 0.8, 3.5)
MARYLAND	1876	6.0	0.8	( 4.4, 7.6)
MASSACHUSETTS	2508	7.3	0.8	( 5.8, 8.8)
MICHIGAN	1162	5.5	0.8	( 3.9, 7.1)
MINNESOTA	2361	5.1	0.6	( 4.0, 6.1)
MISSISSIPPI	875	5.4	0.9	( 3.6, 7.2)
MISSOURI	1886	4.3	0.6	( 3.1, 5.5)
MONTANA	759	3.0	0.9	( 1.3, 4.7)
NEBRASKA	1214	5.6	0.8	( 4.0, 7.2)
NEVADA	820	5.6	1.2	( 3.3, 7.9)
NEW HAMPSHIRE	590	4.2	1.1	( 2.1, 6.4)
NEW JERSEY	1415	9.3	1.1	( 7.2, 11.4)
NEW MEXICO	1471	5.7	0.7	( 4.2, 7.1)
NEW YORK	1318	9.2	1.0	( 7.3, 11.1)
NORTH CAROLINA	1045	2.8	0.6	( 1.7, 3.9)
NORTH DAKOTA	862	3.8	0.7	( 2.3, 5.2)
OHIO	730	5.6	1.2	( 3.3, 7.8)
OKLAHOMA	1261	4.4	0.7	( 3.0, 5.8)
OREGON	766	4.4	0.9	( 2.6, 6.2)
PENNSYLVANIA	1640	6.4	0.8	( 4.9, 7.9)
RHODE ISLAND	1964	7.4	0.8	( 5.8, 9.0)
SOUTH CAROLINA	1476	5.3	0.7	( 3.9, 6.6)
SOUTH DAKOTA	1891	4.1	0.6	( 3.0, 5.2)
TENNESSEE	1379	7.3	0.9	( 5.5, 9.1)
TEXAS	2182	8.4	0.8	( 6.9, 9.9)
UTAH	1367	8.3	1.2	( 5.9, 10.7)
VERMONT	1542	4.7	0.8	( 3.1, 6.2)
VIRGINIA	1589	6.4	1.3	( 4.0, 8.9)
WASHINGTON	1513	3.7	0.6	( 2.6, 4.8)
WEST VIRGINIA	1055	6.6	1.0	( 4.7, 8.5)
WISCONSIN	998	4.5	0.8	( 3.0, 6.1)
WYOMING	972	3.4	0.6	( 2.1, 4.6)
PUERTO RICO	1499	17.4	1.4	( 14.8, 20.1)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 5.4 RANGE = 2.1-17.4

\* NEVER HAD PAP SMEAR, WOMEN WITH INTACT CERVIX

June 20, 2000

TABLE 45.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999  
 NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	5.7	3.3	-2.4
ALASKA	3.3	2.5	-0.8
ARIZONA	15.1	9.1	-6.0
ARKANSAS	7.9	4.4	-3.5
COLORADO	4.5	3.8	-0.7
CONNECTICUT	7.3	5.7	-1.6
DELAWARE	5.1	4.4	-0.7
DISTRICT OF CO	3.0	5.0	2.0
FLORIDA	5.2	5.2	0.0
GEORGIA	4.3	3.4	-0.9
HAWAII	5.5	6.4	0.9
IDAHO	4.8	5.5	0.7
ILLINOIS	5.6	6.0	0.4
INDIANA	6.3	4.9	-1.5
IOWA	5.2	4.6	-0.5
KANSAS	6.5	5.5	-1.0
KENTUCKY	5.0	6.8	1.8
LOUISIANA	6.4	7.2	0.8
MAINE	4.7	2.1	-2.6
MARYLAND	4.8	6.0	1.1
MASSACHUSETTS	6.2	7.3	1.2
MICHIGAN	4.6	5.5	0.9
MINNESOTA	6.0	5.1	-1.0
MISSISSIPPI	6.7	5.4	-1.3
MISSOURI	4.3	4.3	0.0
MONTANA	3.5	3.0	-0.5
NEBRASKA	5.4	5.6	0.2
NEVADA	4.7	5.6	0.9
NEW HAMPSHIRE	5.6	4.2	-1.4
NEW JERSEY	10.3	9.3	-0.9
NEW MEXICO	6.7	5.7	-1.0
NEW YORK	7.4	9.2	1.8
NORTH CAROLINA	4.6	2.8	-1.8
NORTH DAKOTA	4.5	3.8	-0.7
OHIO	5.4	5.6	0.2
OKLAHOMA	7.3	4.4	-2.9
OREGON	6.3	4.4	-2.0
PENNSYLVANIA	6.7	6.4	-0.3
RHODE ISLAND	8.8	7.4	-1.4
SOUTH CAROLINA	6.9	5.3	-1.7
SOUTH DAKOTA	5.9	4.1	-1.8
TENNESSEE	7.1	7.3	0.2
TEXAS	7.8	8.4	0.6
UTAH	9.7	8.3	-1.4
VERMONT	6.0	4.7	-1.4
VIRGINIA	5.6	6.4	0.9
WASHINGTON	5.2	3.7	-1.5
WEST VIRGINIA	6.7	6.6	-0.1
WISCONSIN	5.4	4.5	-0.8
WYOMING	2.7	3.4	0.7
PUERTO RICO	19.4	17.4	-1.9

\* NEVER HAD PAP SMEAR, WOMEN WITH INTACT CERVIX

TABLE 46.1 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA 117  
 ESTIMATED PREVALENCE AND RELEVANT STATISTICS FOR NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	SAMPLE SIZE	PERCENT	STANDARD ERROR	95% CONFIDENCE INTERVAL
ALABAMA	792	13.4	1.4	( 10.7, 16.0)
ALASKA	850	9.1	1.3	( 6.5, 11.6)
ARIZONA	719	13.9	2.0	( 10.1, 17.8)
ARKANSAS	1253	17.4	1.2	( 15.1, 19.8)
COLORADO	1267	12.6	1.3	( 10.0, 15.1)
CONNECTICUT	1747	11.7	1.1	( 9.6, 13.8)
DELAWARE	1238	12.3	1.3	( 9.7, 14.9)
DISTRICT OF CO	658	9.2	1.3	( 6.5, 11.8)
FLORIDA	2271	14.7	0.9	( 13.0, 16.4)
GEORGIA	988	12.0	1.2	( 9.7, 14.2)
HAWAII	950	12.9	1.4	( 10.3, 15.6)
IDAHO	2059	20.7	1.1	( 18.6, 22.7)
ILLINOIS	1451	15.1	1.1	( 12.9, 17.2)
INDIANA	1060	17.6	2.1	( 13.5, 21.7)
IOWA	1653	14.7	1.0	( 12.8, 16.6)
KANSAS	1656	12.6	0.9	( 10.8, 14.4)
KENTUCKY	3343	16.6	0.9	( 14.9, 18.3)
LOUISIANA	679	14.6	1.6	( 11.5, 17.7)
MAINE	719	10.5	1.2	( 8.1, 12.9)
MARYLAND	1862	10.8	1.0	( 8.9, 12.8)
MASSACHUSETTS	2496	13.8	1.0	( 11.9, 15.7)
MICHIGAN	1152	14.2	1.2	( 11.8, 16.5)
MINNESOTA	2345	12.5	0.8	( 10.9, 14.0)
MISSISSIPPI	863	15.9	1.5	( 13.0, 18.8)
MISSOURI	1868	15.4	1.1	( 13.3, 17.4)
MONTANA	753	14.8	1.5	( 11.9, 17.7)
NEBRASKA	1205	15.2	1.2	( 12.9, 17.5)
NEVADA	820	21.5	2.4	( 16.9, 26.2)
NEW HAMPSHIRE	590	11.6	1.6	( 8.5, 14.7)
NEW JERSEY	1391	15.7	1.2	( 13.3, 18.1)
NEW MEXICO	1462	16.3	1.1	( 14.2, 18.4)
NEW YORK	1311	15.8	1.2	( 13.5, 18.1)
NORTH CAROLINA	1043	9.1	1.0	( 7.3, 11.0)
NORTH DAKOTA	854	13.4	1.2	( 11.1, 15.8)
OHIO	723	14.5	1.6	( 11.4, 17.5)
OKLAHOMA	1256	16.2	1.2	( 13.9, 18.6)
OREGON	761	12.4	1.4	( 9.7, 15.1)
PENNSYLVANIA	1622	16.4	1.1	( 14.3, 18.5)
RHODE ISLAND	1949	14.0	1.0	( 12.0, 16.0)
SOUTH CAROLINA	1463	13.0	1.0	( 10.9, 15.0)
SOUTH DAKOTA	1875	14.6	1.0	( 12.8, 16.5)
TENNESSEE	1367	14.9	1.2	( 12.6, 17.2)
TEXAS	2171	17.3	1.0	( 15.4, 19.3)
UTAH	1363	19.2	1.6	( 16.1, 22.3)
VERMONT	1519	11.6	1.0	( 9.6, 13.5)
VIRGINIA	1579	14.5	1.6	( 11.5, 17.6)
WASHINGTON	1508	14.1	1.1	( 11.9, 16.3)
WEST VIRGINIA	1048	19.3	1.4	( 16.7, 22.0)
WISCONSIN	993	15.3	1.3	( 12.8, 17.8)
WYOMING	967	16.2	1.3	( 13.7, 18.6)
PUERTO RICO	1475	26.9	1.5	( 24.1, 29.8)

SUMMARY STATISTICS: NO. OF PARTICIPANTS = 51 MEDIAN = 14.6 RANGE = 9.1-26.9

\* DID NOT HAVE A PAP SMEAR IN PAST THREE YEARS

June 20, 2000

TABLE 46.2 1999 BEHAVIORAL RISK FACTOR SURVEILLANCE DATA  
 PREVALENCE DIFFERENCES FROM 1998 TO 1999  
 NO PAP SMEAR, WOMEN WITH INTACT CERVIX, AGE 18 AND OVER\*  
 DENOMINATOR EXCLUDES MISSING, DON'T KNOW, AND REFUSED

PARTICIPANT	1998 PERCENT	1999 PERCENT	PERCENT DIFFERENCE
ALABAMA	14.7	13.4	-1.4
ALASKA	10.5	9.1	-1.4
ARIZONA	18.3	13.9	-4.4
ARKANSAS	21.1	17.4	-3.6
COLORADO	13.0	12.6	-0.4
CONNECTICUT	13.8	11.7	-2.1
DELAWARE	12.9	12.3	-0.6
DISTRICT OF CO	6.1	9.2	3.1
FLORIDA	14.1	14.7	0.6
GEORGIA	13.9	12.0	-1.9
HAWAII	13.7	12.9	-0.8
IDAHO	17.5	20.7	3.2
ILLINOIS	15.5	15.1	-0.4
INDIANA	19.1	17.6	-1.4
IOWA	15.5	14.7	-0.8
KANSAS	14.4	12.6	-1.8
KENTUCKY	15.7	16.6	0.9
LOUISIANA	15.2	14.6	-0.6
MAINE	15.0	10.5	-4.5
MARYLAND	10.8	10.8	0.1
MASSACHUSETTS	12.3	13.8	1.6
MICHIGAN	11.9	14.2	2.3
MINNESOTA	15.0	12.5	-2.5
MISSISSIPPI	15.6	15.9	0.3
MISSOURI	14.6	15.4	0.8
MONTANA	19.4	14.8	-4.6
NEBRASKA	17.6	15.2	-2.4
NEVADA	14.9	21.5	6.6
NEW HAMPSHIRE	15.8	11.6	-4.2
NEW JERSEY	18.6	15.7	-2.9
NEW MEXICO	17.4	16.3	-1.1
NEW YORK	16.2	15.8	-0.4
NORTH CAROLINA	12.3	9.1	-3.2
NORTH DAKOTA	14.5	13.4	-1.0
OHIO	15.8	14.5	-1.4
OKLAHOMA	15.2	16.2	1.0
OREGON	16.3	12.4	-3.8
PENNSYLVANIA	16.2	16.4	0.2
RHODE ISLAND	14.8	14.0	-0.8
SOUTH CAROLINA	15.0	13.0	-2.1
SOUTH DAKOTA	13.7	14.6	1.0
TENNESSEE	13.9	14.9	1.0
TEXAS	18.5	17.3	-1.2
UTAH	21.6	19.2	-2.4
VERMONT	14.0	11.6	-2.4
VIRGINIA	15.1	14.5	-0.6
WASHINGTON	14.6	14.1	-0.5
WEST VIRGINIA	19.0	19.3	0.3
WISCONSIN	16.6	15.3	-1.3
WYOMING	18.0	16.2	-1.8
PUERTO RICO	31.0	26.9	-4.0

\* DID NOT HAVE A PAP SMEAR IN PAST THREE YEARS

June 20, 2000