


Guidelines for the Use of Antiretroviral Agents in HIV-Infected Adults and Adolescents

February 4, 2002



**Developed by the panel on Clinical Practices
for Treatment of HIV Infection
convened by the
Department of Health and Human Services
(DHHS) and the
Henry J. Kaiser Family Foundation**

It is emphasized that concepts relevant to HIV management evolve rapidly. The Panel has a mechanism to update recommendations on a regular basis, and the most recent information is available on the **HIV/AIDS Treatment Information Service website** (<http://www.hivatis.org>).

February 4, 2002

Table of Contents

Guidelines Panel Roster	1
Summary	2
Introduction	3
Use of Testing for Plasma HIV RNA Levels and CD4 ⁺ T Cell Count in Guiding Decisions for Therapy	4
Testing for Drug Resistance	5
Background	5
Use of Resistance Assays in Clinical Practice	5
Considerations for Patients with Established HIV Infection	6
Considerations for Initiating Therapy in the Patient with Asymptomatic HIV Infection	7
Considerations for Discontinuing Therapy	9
Adherence to Potent Antiretroviral Therapy	10
Introduction	10
Adherence in HIV Disease	10
Adherence to HAART: Approach to the Patient	11
Patient-related strategies	11
Clinician and health team-related strategies	11
Regimen-related strategies	12
Directly observed therapy	12
Goals of Therapy	13
Tools to Achieve the Goals of Therapy	13
Initiating Therapy in the Patient with Asymptomatic HIV Infection	14
Initiating Therapy in Advanced HIV Disease	15
HAART-Associated Adverse Clinical Events	16
Lactic Acidosis/Hepatic Steatosis	16
Hepatotoxicity	17
Hyperglycemia	18
Fat Maldistribution	18
Hyperlipidemia	19
Increased Bleeding Episodes in Patients with Hemophilia	19
Osteonecrosis, Osteopenia, and Osteoporosis	19
Skin Rash	20
Interruption of Antiretroviral Therapy	21
Considerations for Changing a Failing Regimen	21
Criteria for Changing Therapy	22
Therapeutic Options When Changing Antiretroviral Therapy	23
Acute HIV Infection	24
Treatment Regimen for Primary HIV Infection	25
Patient Follow-up	25
Duration of Therapy for Primary HIV Infection	25
Considerations for Antiretroviral Therapy in the HIV-Infected Adolescent	25
Considerations for Antiretroviral Therapy in the HIV-Infected Pregnant Woman	26

Prevention Counseling in the Patient with HIV Infection	29
Conclusion	31
Tables and Figure	32
References	65

Supplements

Supplement: Considerations for Antiretroviral Therapy in Women	1
Supplement: Hydroxyurea	1
Supplement (new): Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy (5/23/2002)	1

List of Tables and Figure

Table 1. Rating Scheme for Clinical Practice	32
Table 2. Indications for Plasma HIV RNA Testing	33
Table 3. Recommendations for the Use of Drug Resistance Assays	34
Table 4. Risks and Benefits of Delayed Initiation of Therapy and of Early Therapy in the Asymptomatic HIV-Infected Patient	35
Table 5. Risk of Progression to AIDS Defining Illness in a Cohort of Homosexual Men Predicted by Baseline CD4 ⁺ T Cell Count and Viral Load	36
Table 6. Indications for the Initiation of Antiretroviral Therapy in the Chronically HIV-1 Infected Patient	37
Table 7. Strategies to Improve Adherence: Patient and Medication-Related	38
Table 8. Strategies to Improve Adherence: Clinician and Health Team-Related	39
Table 9. Interventions Associated with Improved Adherence	40
Table 10. Goals of HIV Therapy and Tools to Achieve Them	41
Table 11. Advantages and Disadvantages of Class-Sparing Regimens	42
Table 12. Recommended Antiretroviral Agents for Initial Treatment of Established HIV Infection	43
Table 13. Characteristics of Nucleoside Reverse Transcriptase Inhibitors (NRTIs)	44
Table 14. Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)	46
Table 15. Characteristics of Protease Inhibitors (PIs)	48
Table 16. Adverse Drug Reactions Related “Black Box Warnings” in Product Labeling for Antiretroviral Agents	49
Table 17. Drugs That Should Not Be Used With PI Antiretrovirals	51
Table 18. Drug Interactions Between Antiretrovirals and Other Drugs: Protease Inhibitors	53
Table 19. Drug Interactions: Protease Inhibitors Effect of Drug on Levels (AUCs)/Dose	57
Table 20. HIV-Related Drugs with Overlapping Toxicities	59
Table 21. Guidelines for Changing an Antiretroviral Regimen for Suspected Drug Failure	60
Table 22. Acute Retroviral Syndrome: Associated Signs & Symptoms (Expected Frequency)	61
Table 23. Preclinical and Clinical Data Relevant to Use of Antiretrovirals in Pregnancy	62
Table 24. Zidovudine Perinatal Transmission Prophylaxis Regimen	63
<i>Figure 1. Likelihood of Developing AIDS Within 3 Years</i>	64

Guidelines Panel Roster

These Guidelines were developed by the Panel on Clinical Practices for Treatment of HIV Infection convened by the Department of Health and Human Services (DHHS) and the Henry J. Kaiser Family Foundation.

Leadership of the Panel:

Anthony S. Fauci, *National Institutes of Health, Bethesda, MD (co-chair);*
 John G. Bartlett, *Johns Hopkins University, Baltimore, MD (co-chair);*
 Eric P. Goosby, *DHHS, Washington, DC, (co-convenor); and*
 Jennifer Kates, *Henry J. Kaiser Foundation, San Francisco, CA, (co-convenor).*

Current members of the Panel include:

Jean Anderson	Johns Hopkins University, Baltimore, MD
Rodney Armstead	UHP Healthcare, Inc., Inglewood, CA
Cornelius Baker	Whitman-Walker Clinic, Washington, DC
David Barr	Forum for Collaborative HIV Research, Washington, DC
Samuel A. Bozzette	SDVA Medical Center, San Diego, CA
Charles Carpenter	Brown University School of Medicine, Providence, RI
Martin Delaney	Project <i>Inform</i> , San Francisco, CA
Lawrence Deyton	Dept. of Veterans Affairs, Washington, DC
Wafaa El-Sadr	Harlem Hospital, New York, NY
Courtney Fletcher	University of Minnesota, MN
Gregg Gonsalves	Gay Men's Health Crisis, New York, NY
Fred Gordin	Veterans Administration Medical Center, Washington, DC
Mark Harrington	Treatment Action Group, New York, NY
Martin S. Hirsch	Massachusetts General Hospital, Boston, MA
John W. Mellors	University of Pittsburgh, Pittsburgh, PA
James Neaton	University of Minnesota, Minneapolis, MN
Sallie Perryman	New York State Department of Health, New York, NY
John Phair	Northwestern University, Chicago, IL
Robert T. Schooley	University of Colorado, Denver, CO
Renslow Sherer	The CORE Center, Cook County Hospital, Chicago, IL
Stephen A. Spector	University of California San Diego, La Jolla, CA
Sharilyn K. Stanley	Texas Department Of Health, Austin, TX
Paul Volberding	University of California, San Francisco, CA
Suzanne Willard	MCP Hahnemann University, Philadelphia, PA

Participants from the Department of Health and Human Services:

Victoria Cargill	National Institutes of Health
Oren Cohen	National Institutes of Health
Mark Dybul	National Institutes of Health
T. Randolph Graydon	Centers for Medicare and Medicaid Services
Heidi Jolson	Food and Drug Administration
Jonathan Kaplan	Centers for Disease Control and Prevention
Abe Macher	Health Resources and Services Administration
Henry Masur	National Institutes of Health
Lynne Mofenson	National Institutes of Health
Jeffrey Murray	Food and Drug Administration
Joseph O'Neill	Health Resources and Services Administration
Alice Pau	National Institutes of Health
Lucille C. Perez	Substance Abuse and Mental Health Services Administration

Non-voting observers include:

Gerry Bally	Health Canada, Ottawa, Ontario
Douglas Brust	National Institutes of Health
Sophia Chang	Center for Quality HIV Care, V.A., Palo Alto, CA, Washington, DC
Richard Marlink	Harvard AIDS Institute, Cambridge, MA
Celia Maxwell	AIDS Education and Training Center, Washington, DC
Howard Minkoff	Maimonides Medical Center, Brooklyn, NY
James Oleske	UMDNJ, Newark, NJ
Daniel Simpson	Indian Health Service, Rockville, MD

Guidelines for the Use of Antiretroviral Agents in HIV-Infected Adults and Adolescents

SUMMARY

The availability of an increasing number of antiretroviral agents and the rapid evolution of new information has introduced extraordinary complexity into the treatment of HIV-infected persons. In 1996, the Department of Health and Human Services and the Henry J. Kaiser Family Foundation convened the Panel on Clinical Practices for the Treatment of HIV to develop guidelines for the clinical management of HIV-infected adults and adolescents.

This report recommends that care should be supervised by an expert. The report provides guidelines for antiretroviral therapy, including when to start treatment, what drugs to initiate, when to change therapy, and therapeutic options when changing therapy. In addition, the report makes recommendations for laboratory monitoring, including plasma HIV RNA, CD4⁺ T cell counts and HIV drug resistance testing. Special considerations are provided for adolescents and pregnant women. As with treatment of other chronic conditions, therapeutic decisions require a mutual understanding between the patient and the health care provider regarding the benefits and risks of treatment. Antiretroviral regimens are complex, have major side effects, pose adherence difficulties, and carry serious potential consequences from the development of viral resistance due to non-adherence to the drug regimen or suboptimal levels of antiretroviral agents. Patient education and involvement in therapeutic decisions is important for all medical conditions, but it is considered especially critical in cases of HIV infection and its treatment.

Treatment should be offered to all patients with symptoms ascribed to HIV infection. Recommendations for offering antiretroviral therapy in asymptomatic patients require analysis of many real and potential risks and benefits. In general, treatment should be offered to individuals with fewer than 350 CD4⁺ T cells/mm³ or plasma HIV RNA levels exceeding 55,000 copies/mL (by RT-PCR or bDNA assay). The strength of the recommendation to treat asymptomatic patients should be based on the willingness and readiness of the individual to begin therapy; the degree of existing immunodeficiency as determined by the CD4⁺ T cell count; the risk of disease progression as determined by the CD4⁺ T cell

count and level of plasma HIV RNA; the potential benefits and risks of initiating therapy in asymptomatic individuals; and the likelihood, after counseling and education, of adherence to the prescribed treatment regimen. Once the decision has been made to initiate antiretroviral therapy, the goals should be maximal and durable suppression of viral load, restoration and/or preservation of immunologic function, improvement of quality of life, and reduction of HIV-related morbidity and mortality. Results of therapy are evaluated primarily with plasma HIV RNA levels; these are expected to show a one-log₁₀ decrease at eight weeks and no detectable virus (<50 copies/mL) at 4–6 months after initiation of treatment. Failure of therapy at 4–6 months may be ascribed to non-adherence, inadequate potency of drugs, or suboptimal levels of antiretroviral agents, viral resistance, and other factors that are poorly understood. Patients whose therapy fails in spite of a high level of adherence to the regimen should have their regimen changed; this change should be guided by a thorough drug treatment history and the results of drug resistance testing. Optimal changes in therapy may be especially difficult to achieve for patients in whom the primary regimen has failed, due to limitations in the available alternative antiretroviral regimens that have documented efficacy; these decisions are further confounded by problems with adherence, toxicity, and resistance. In some settings, it may be preferable to participate in a clinical trial with or without access to new drugs, or to use a regimen that may not achieve complete suppression of viral replication.

It is emphasized that concepts relevant to HIV management evolve rapidly. The Panel has a mechanism to update recommendations on a regular basis, and the most recent information is available on the **HIV/AIDS Treatment Information Service Web site** (<http://www.hivatis.org>).

Guidelines for the Use of Antiretroviral Agents in HIV-Infected Adults and Adolescents

INTRODUCTION

This document was developed by the Panel on Clinical Practices for Treatment of HIV Infection convened by the Department of Health and Human Services (DHHS) and the Henry J. Kaiser Family Foundation. The document contains recommendations for the clinical use of antiretroviral agents in the treatment of HIV-infected adults and adolescents (defined here as late puberty or Tanner V). Guidance for the use of antiretroviral treatment in pediatric HIV infection is not contained in this document. While the pathogenesis of HIV infection and the general virologic and immunologic principles underlying the use of antiretroviral therapy are similar for all HIV-infected individuals, there are unique therapeutic and management considerations in HIV-infected children. In recognition of these differences, a separate document addresses pediatric-specific issues related to antiretroviral therapy, and is available at (<http://www.hivatis.org>).

These guidelines are intended for use by physicians and other health care providers who use antiretroviral therapy to treat HIV-infected adults and adolescents and serve as a companion to the therapeutic principles formulated by the National Institutes of Health (NIH) Panel to Define Principles of Therapy of HIV Infection [1]. Together the documents should provide the pathogenesis-based rationale for therapeutic strategies as well as practical guidelines for implementing these strategies. While the guidelines represent the current state of knowledge regarding the use of antiretroviral agents, this is a rapidly evolving field of science, and the availability of new agents or new clinical data regarding the use of existing agents will result in changes in therapeutic options and preferences. Thus, in recognition of the need for frequent updates to this document, a subgroup of the Panel, the Antiretroviral Working Group, meets monthly to review new data; recommendations for changes in this document are then submitted to the Panel and incorporated as appropriate. Copies of this document and all updates are available from the

HIV/AIDS Treatment Information Service-ATIS
(1-800-448-0440; TTY 1-888-480-3739;
Fax 301-519-6616) and
on the ATIS Web site (<http://www.hivatis.org>).

They are also available from the National Prevention Information Network (NPIN) Web site (<http://www.cdcpin.org>). These recommendations are not intended to substitute for the judgment of a clinician who is an expert in the care of HIV-infected individuals. It is important to note that the Panel felt that, where possible, the treatment of HIV-infected patients should be directed by a clinician with extensive experience in the care of these patients. When this is not possible, it is important to have access to such expertise through consultations.

Each recommendation is accompanied by a rating that includes a letter and a Roman numeral (**Table 1**); and is similar to the rating schemes used in previous guidelines on the prophylaxis of opportunistic infections (OIs) issued by the U.S. Public Health Service and the Infectious Diseases Society of America [2]. The letter indicates the strength of the recommendation, based on the opinion of the Panel, while the Roman numeral rating reflects the nature of the evidence supporting the recommendation (**Table 1**). Thus, recommendations based on data from clinical trials with clinical endpoints are differentiated from those with laboratory endpoints such as CD4⁺ T lymphocyte count or plasma HIV RNA levels; where no clinical trial data are available, recommendations are based on the opinions of experts familiar with the relevant scientific literature.

This document addresses the following issues: the use of testing for plasma HIV RNA levels (viral load) and CD4⁺ T cell count; the use of testing for antiretroviral drug resistance; considerations for when to initiate therapy in cases of established HIV infection; adherence to antiretroviral therapy; special considerations for therapy in patients with advanced stage disease; therapy-related adverse events; interruption of therapy; considerations for changing therapy and available therapeutic options; the treatment of acute HIV infection; considerations for antiretroviral therapy in adolescents; considerations for antiretroviral therapy in pregnant women; and issues related to the transmission of HIV to others.

USE OF TESTING FOR PLASMA HIV RNA LEVELS AND CD4⁺ T CELL COUNT IN GUIDING DECISIONS FOR THERAPY

Decisions regarding initiation or changes in antiretroviral therapy should be guided by monitoring the laboratory parameters of plasma HIV RNA (viral load) and CD4⁺ T cell count, as well as the clinical condition of the patient. Results of these two laboratory tests give the clinician important information about the virologic and immunologic status of the patient and the risk of disease progression to AIDS [3, 4]. It should be noted that HIV viral load testing has been approved by the FDA for determining prognosis and for monitoring the response to therapy only for the RT-PCR assay (Roche). Multiple analyses in over 5000 patients who participated in approximately 18 trials with viral load monitoring showed a statistically significant dose-response type association between decreases in plasma viremia and improved clinical outcome based on standard endpoints of new AIDS-defining diagnoses and survival. This relationship was observed over a range of patient baseline characteristics including: pretreatment plasma RNA level, CD4⁺ T cell count, and prior drug experience. Thus, it is the consensus of the Panel that viral load testing is an essential parameter in decisions to initiate or change antiretroviral therapies. Measurement of plasma HIV RNA levels (viral load), using quantitative methods, should be performed at the time of diagnosis and every 3–4 months thereafter in the untreated patient (AIII) ([Table 2](#)). CD4⁺ T cell counts should be measured at the time of diagnosis and generally every 3–6 months thereafter (AIII). These intervals between tests are merely recommendations and flexibility should be exercised according to the circumstances of the individual case. Plasma HIV RNA levels should also be measured immediately prior to and again at 2–8 weeks after initiation of antiretroviral therapy (AIII). This second time point allows the clinician to evaluate the initial effectiveness of therapy, since in most patients adherence to a regimen of potent antiretroviral agents should result in a large decrease ($\sim 1.0 \log_{10}$) in viral load by 2–8 weeks. The viral load should continue to decline over the following weeks and in most individuals becomes below detectable levels (currently defined as <50 RNA copies/mL) by 16–20 weeks. The rate of viral load decline towards undetectable is affected by the baseline CD4⁺ T cell count, the initial viral load, potency of the regimen, adherence to the

regimen, prior exposure to antiretroviral agents, and the presence of any OIs.

These individual differences must be considered when monitoring the effect of therapy. However, the absence of a virologic response of the magnitude discussed above should prompt the clinician to reassess patient adherence, rule out malabsorption, consider repeat RNA testing to document lack of response, and/or consider a change in drug regimen. Once the patient is on therapy, HIV RNA testing should be repeated every 3–4 months to evaluate the continuing effectiveness of therapy (AII). With optimal therapy viral levels in plasma at 6 months should be undetectable, that is, below 50 copies of HIV RNA per mL of plasma [5]. Data from clinical trials strongly suggest that lowering plasma HIV RNA to below 50 copies/mL is associated with a more complete and durable viral suppression, compared with reducing HIV RNA to levels between 50–500 copies/mL [6]. If HIV RNA remains detectable in plasma after 16–20 weeks of therapy, the plasma HIV RNA test should be repeated to confirm the result and a change in therapy should be considered, according to the guidelines in the section [“**Considerations for Changing a Failing Regimen**”](#) (see p. 21) (BIII).

When making decisions regarding the initiation of therapy, the CD4⁺ T lymphocyte count and plasma HIV RNA measurement should ideally be performed on two occasions to ensure accuracy and consistency of measurement (BIII). However, in patients who present with advanced HIV disease, antiretroviral therapy should generally be initiated after the first viral load measurement is obtained in order to prevent a potentially deleterious delay in treatment. It is recognized that the requirement for two measurements of viral load may place a significant financial burden on patients or payers. Nonetheless, the Panel feels that two measurements of viral load will provide the clinician with the best information for subsequent follow-up of the patient. Plasma HIV RNA levels should not be measured during or within four weeks after successful treatment of any intercurrent infection, resolution of symptomatic illness, or immunization. Because there are differences among commercially available tests, confirmatory plasma HIV RNA levels should be measured by the same laboratory using the same technique in order to ensure consistent results.

A minimally significant change in plasma viremia is considered to be a 3-fold or $0.5 \log_{10}$ increase or decrease. A significant decrease in CD4⁺ T lymphocyte

count is a decrease of >30% from baseline for absolute cell numbers and a decrease of >3% from baseline in percentages of cells [7]. Discordance between trends in CD4⁺ T cell numbers and plasma HIV RNA levels can occur and was found in 20% of patients in one cohort studied [8]. Such discordance can complicate decisions regarding antiretroviral therapy and may be due to a number of factors that affect plasma HIV RNA testing. In general, viral load and trends in viral load are felt to be more informative for guiding decisions regarding antiretroviral therapy than are CD4⁺ T cell counts; exceptions to this rule do occur, however. For further discussion refer to “[Considerations for Changing a Failing Regimen](#),” p. 21. In many such cases, expert consultation should be considered.

TESTING FOR DRUG RESISTANCE

Background

Testing for HIV resistance to antiretroviral drugs is a rational adjunct to guide antiretroviral therapy. When combined with a detailed drug history and efforts aimed at maximizing drug adherence, these assays may help to maximize the benefits of antiretroviral therapy. Many studies in treatment experienced patients have shown strong associations between the presence of drug resistance (identified by either genotyping or phenotyping resistance assays) and failure of the antiretroviral treatment regimen to suppress HIV replication. Genotyping assays detect drug resistance mutations that are present in the relevant viral genes (i.e., RT and protease). Some genotyping assays involve sequencing of the entire RT and protease genes, while others utilize probes to detect selected mutations that are known to confer drug resistance. Genotyping assays can be performed relatively rapidly, such that results can be reported within 1–2 weeks of sample collection. Interpretation of test results requires an appreciation of the range of mutations that are selected for by various antiretroviral drugs, as well as the potential for cross-resistance to other drugs conferred by some of these mutations (see the <http://hiv-web.lanl.gov> Web site). Consultation with an expert in HIV drug resistance is encouraged to facilitate interpretation of genotypic test results.

Phenotyping assays measure the ability of viruses to grow in various concentrations of antiretroviral drugs. Automated, recombinant phenotyping assays are commercially available with turn-around times of 2–3 weeks; however, phenotyping assays are generally more costly to perform compared with genotypic

assays. Recombinant phenotyping assays involve insertion of the RT and protease gene sequences derived from patient plasma HIV RNA into the backbone of a laboratory clone of HIV either by cloning or *in vitro* recombination. Replication of the recombinant virus at various drug concentrations is monitored by expression of a reporter gene and is compared with replication of a reference strain of HIV. The concentrations of drugs that inhibit 50% and 90% of viral replication (i.e., the IC₅₀ and IC₉₀) are calculated, and the ratio of the IC₅₀s of the test and reference viruses is reported as the fold increase in IC₅₀, or fold resistance. Interpretation of phenotyping assay results is complicated by the paucity of data on the specific level of resistance (fold increase in IC₅₀) that is associated with failure of different drugs; again, consultation with an expert may be helpful for interpretation of test results.

Further limitations of both genotyping and phenotyping assays include the lack of uniform quality assurance for all assays that are currently available, relatively high cost, and insensitivity for minor viral species; if drug-resistant viruses are present but constitute less than 10–20% of the circulating virus population, they will likely not be detected by current assays. This limitation is of particular importance when interpreting data about susceptibility to drugs that the patient has taken in the past but are not part of the current antiretroviral regimen. If drug resistance had developed to a drug that was subsequently discontinued, the drug-resistant virus can become a minor species because its growth advantage is lost [9]. Consequently, resistance assays should be performed while the patient is taking his/her antiretroviral regimen, and data suggesting the absence of resistance should be interpreted carefully in relation to the prior treatment history.

Use of Resistance Assays in Clinical Practice

Resistance assays may be useful in the setting of virologic failure on antiretroviral therapy and in acute HIV infection ([Table 3](#)). Recent prospective data supporting the use of resistance testing in clinical practice come from trials in which the utility of resistance tests were assessed in the setting of virologic failure. The VIRADAPT [10] and GART [11] studies compared virologic responses to antiretroviral treatment regimens when genotyping resistance tests were available to help guide therapy with those observed when changes in therapy were guided solely by clinical judgment. The results of both studies indicated that the short-term virologic response to

therapy was significantly greater when results of resistance testing were available. Similarly, a recent prospective, randomized, multicenter trial has shown that therapy selected on the basis of phenotypic resistance testing significantly improves the virological response to antiretroviral therapy, compared with therapy selected without the aid of phenotypic testing [12]. Thus, resistance testing appears to be a useful tool in selecting active drugs when changing antiretroviral regimens in the setting of virologic failure (BII). Similar rationale applies to the potential use of resistance testing in the setting of suboptimal viral load reduction, as detailed in “[Criteria for Changing Therapy](#),” p. 22 (BIII). It should be noted that virologic failure in the setting of HAART (Highly Active Antiretroviral Therapy) is in some instances associated with resistance only to one component of the regimen [13]; in this situation, it may be possible to substitute individual drugs in a failing regimen, although this concept requires clinical validation (“[Considerations for Changing a Failing Regimen](#),” p. 21). There are currently no prospective data to support the use of one type of resistance assay over the other (i.e., genotyping vs. phenotyping) in different clinical situations. Therefore, one type of assay is generally recommended per sample; however, in the setting of a complex prior treatment history, both assays may provide important and complementary information.

Transmission of drug-resistant strains of HIV has been documented, and may be associated with a suboptimal virologic response to initial antiretroviral therapy [14-17]. If the decision is made to initiate therapy in an individual with acute HIV infection, optimization of the initial antiretroviral regimen through the use of resistance testing is a reasonable, albeit untested, strategy (CIII). Because of its more rapid turnaround time, the use of a genotypic assay may be preferred in this setting. The use of resistance testing prior to initiation of antiretroviral therapy in chronic HIV infection is not generally recommended (DIII) because of uncertainty about the prevalence of resistance in treatment-naïve individuals and the fact that currently available resistance assays may fail to detect drug resistant species that were transmitted at the time of primary infection but became a minor species in the absence of selective drug pressure. The currently favored approach would be to reserve resistance testing for cases in which viral load suppression was suboptimal after initiation of therapy (see above), although this may change as more

information becomes available on the prevalence of resistant virus in antiretroviral-naïve individuals.

In general, recommendations for resistance testing in pregnancy should be the same as for non-pregnant patients: acute HIV infection, virologic failure on an antiretroviral regimen, or suboptimal viral load suppression after initiation of antiretroviral therapy are all appropriate indications for resistance testing. If an HIV⁺ pregnant woman is taking an antiretroviral regimen that does not include zidovudine, or if zidovudine was discontinued because of maternal drug resistance, intrapartum and neonatal zidovudine prophylaxis should still be administered to prevent mother-to-infant HIV transmission (“[Considerations for Antiretroviral Therapy in the HIV-Infected Pregnant Woman](#),” see p. 26 and [Table 24](#)). It is important to note that not all of zidovudine’s activity in preventing mother-to-infant transmission of HIV can be accounted for by its effect on maternal viral load [20]; furthermore, preliminary data indicate that the rate of perinatal transmission following zidovudine prophylaxis may not differ between those with and without zidovudine resistance mutations [21, 22]. Further studies are needed to determine the best strategy to prevent mother-to-infant HIV transmission in the presence of zidovudine resistance.

CONSIDERATIONS FOR PATIENTS WITH ESTABLISHED HIV INFECTION

Patients with established HIV infection are discussed in two arbitrarily defined clinical categories:

1. Asymptomatic infection or
2. Symptomatic disease (wasting, thrush or unexplained fever for >2 weeks) including AIDS, defined according to the 1993 CDC classification system [23].

All patients in the second category should be offered antiretroviral therapy. Considerations for initiating antiretroviral therapy in the first category of patients are complex and are discussed separately below. Before initiating therapy in any patient, however, the following evaluation should be performed:

- Complete history and physical (AII)

- Complete blood count, chemistry profile (including serum transaminases and lipid profile (AII))
- CD4⁺ T lymphocyte count (AI)
- Plasma HIV RNA Measurement (AI)

Additional evaluation should include routine tests pertinent to the prevention of OIs, if not already performed (RPR or VDRL, tuberculin skin test, toxoplasma IgG serology, and gynecologic exam with Pap smear), and other tests as clinically indicated (e.g., chest X-ray, hepatitis C virus (HCV) serology, ophthalmologic exam) (AII). Hepatitis B virus (HBV) serology is indicated in a patient who is a candidate for the hepatitis B vaccine or has abnormal liver function tests (AII), and CMV serology may be useful in certain individuals, as discussed in the “USPHS/IDSA Guidelines for the Prevention of Opportunistic Infections in Persons Infected with the Human Immunodeficiency Virus” [2] (BIII).

CONSIDERATIONS FOR INITIATING THERAPY IN THE PATIENT WITH ASYMPTOMATIC HIV INFECTION

While randomized clinical trials provide strong evidence for treating patients with <200 CD4⁺ T cells/mm³ [24–26], the optimal time to initiate antiretroviral therapy among asymptomatic patients with CD4⁺ T cell counts >200 cells/mm³ is not known. For individuals with >200 CD4⁺ T cells/mm³, the strength of the recommendation for therapy must balance the readiness of the patient for treatment, consideration of the prognosis for disease-free survival as determined by baseline CD4⁺ T cell count and viral load levels, and assessment of the risks and potential benefits associated with initiating antiretroviral therapy.

In regard to the prognosis based on the patient’s CD4⁺ T cell count and viral load, there is an absence of data on clinical endpoints from randomized, controlled trials for individuals with >200 CD4⁺ T cells/mm³ to guide the decision on when to initiate therapy. However, despite their limitations, observational cohorts of HIV-infected individuals either treated or untreated with antiretroviral therapy provide important data to assist in the assessment of the risk of disease progression.

Observational cohorts have provided critical data on the prognostic importance of viral load and CD4⁺ T cell count in the absence of treatment. These data indicate a strong relationship between plasma HIV

RNA levels and CD4⁺ T cell counts in terms of the risk of progression to AIDS for untreated individuals and provide potent support for the conclusion that therapy should be initiated before the CD4⁺ T cell count declines to <200 cells/mm³ (Figure 1 and Table 5). In addition, these studies are useful for the identification of high-risk, asymptomatic individuals with CD4⁺ T cell counts >200 cells/mm³ who may be candidates for antiretroviral therapy or more frequent CD4⁺ T cell count monitoring. In this regard, the Multicenter AIDS Cohort Study (MACS) demonstrated that the 3-year risk of progression to AIDS was 38.5% among patients with 201 to 350 CD4⁺ T cells/mm³ compared to 14.3% for patients with CD4⁺ T cell counts >350 cells/mm³. However, the short-term risk of progression also was related to the level of plasma HIV RNA, and the risk is relatively low in those with <20,000 copies/mL. An evaluation of 231 individuals with CD4⁺ T cell counts of 201–350 cells/mm³ demonstrated that the 3-year risk of progression to AIDS was 4.1% for the 74 patients with HIV RNA ≤20,000; 36.4% for those 53 patients with HIV RNA 20,001–55,000 copies/mL; and 64.4% for those 104 patients with HIV RNA >55,000 copies/mL. Similar risk gradations by viral load are evident for patients with CD4⁺ T cell counts >350 cells/mm³ (Figure 1 and Table 5) (A. Munoz, unpublished observation). These data indicate that for many patients with CD4⁺ T cell counts >200 cells/mm³, the 3-year risk of disease progression to AIDS in the absence of treatment is very high. Thus, although observational studies of untreated individuals cannot assess the effects of therapy, and therefore cannot determine the optimal time to initiate therapy, these studies do provide important guidance on the risks of progression in the absence of therapy based on a patient’s CD4⁺ T cell count and viral load.

Data from observational studies of HAART-treated cohorts also provide important information to guide the use of antiretroviral therapy in asymptomatic patients [27–30]. A collaborative analysis of data from 13 cohort studies from Europe and North America indicates that in drug-naïve patients without AIDS-defining illness and a viral load <100,000 copies/mL, the 3-year probability of progression to AIDS or death was 15.8% among those who initiated therapy with CD4⁺ T cell counts 0–49 cells/mm³; 12.5% among those with CD4⁺ T cell counts 50–99 cells/mm³; 9.3% among those with CD4⁺ T cell counts 100–199 cells/mm³; 4.7% among those with CD4⁺ T cell counts 200–349 cells/mm³; and 3.4% among those with CD4⁺ T cell counts of 350 cells/mm³ or higher [30]. These data suggest that the prognosis may be better for patients who initiate therapy above 200 cells/mm³; but

risk following the initiation of therapy does not vary considerably above 200 cells/mm³. The risk of progression also was related to plasma HIV RNA levels in this study. There was a substantial increase in risk to progression among patients with a viral load >100,000 copies/mL among all patients. In other cohort studies, an apparent benefit in terms of disease progression was reported among individuals who began antiretroviral therapy when CD4⁺ T cell counts were >350 cells/mm³ compared to those who deferred therapy [31, 32]. For example, in the Swiss cohort study, there was approximately a 7-fold decrease in disease progression to AIDS in individuals who initiated therapy with a CD4⁺ T cell count >350 cells/mm³ compared to those who were monitored without therapy over a 2 year period [32]. However, there was a substantial incidence of adverse treatment effects among patients who initiated therapy: 40% of patients had at least one treatment switch due to adverse effects, and 20% were no longer on treatment after 2 years [32]. Unfortunately, observational studies of individuals treated with HAART also have limitations regarding the ability to determine an optimal time to initiate therapy. The relative risks of disease progression for individuals with CD4⁺ T cell counts 200–349 and >350 cells/mm³ cannot be precisely compared due to the low level of disease progression of these patients during the follow-up period. In addition, the groups may differ in important known and unknown prognostic factors that bias the comparison.

In addition to the risks of disease progression, the decision to initiate antiretroviral therapy also is influenced by an assessment of other potential risks and benefits associated with treatment. [Table 4](#) summarizes the potential benefits and risks of early and of delayed initiation of therapy in the asymptomatic patient that the clinician and the patient should consider in deciding when to initiate therapy.

Potential benefits of early therapy include

1. earlier suppression of viral replication
2. preservation of immune function
3. prolongation of disease-free survival, and
4. decrease in the risk of viral transmission.

Risks of early therapy include

1. the adverse effects of the drugs on quality of life
2. the inconvenience of most of the suppressive regimens currently available, leading to reduced adherence;
3. development of drug resistance over time because of sub-optimal suppression of viral replication;

4. limitation of future treatment options due to premature cycling of the patient through the available drugs;
5. the risk of transmission of virus resistant to antiretroviral drugs;
6. serious toxicities associated with some antiretroviral drugs (e.g., elevations in serum levels of cholesterol and triglycerides, alterations in the distribution of body fat, insulin resistance, and even frank diabetes mellitus); and
7. the unknown durability of effect of the currently available therapies.

The potential benefits of delayed therapy include

1. minimization of treatment-related negative effects on quality of life and drug-related toxicities;
2. preservation of treatment options; and
3. delay in the development of drug resistance.

Potential risks of delayed therapy include

1. the possibility that some damage to the immune system that might otherwise be salvaged by earlier therapy is irreversible;
2. the possibility that suppression of viral replication may be more difficult at a later stage of disease; and
3. the increased risk of HIV transmission to others during a longer untreated period.

Finally, for certain individuals, it may be difficult to ascertain the precise time at which the CD4⁺ T cell count will decrease to a level where the risk of disease is high and there may be a period of time required to identify an effective, tolerable regimen; this may be better accomplished prior to reaching a CD4⁺ T cell count of 200 cells/mm³.

When considering the available data in terms of the relative risk of progression to AIDS at certain CD4⁺ T cell counts and viral loads, and the potential risks and benefits associated with initiating therapy, many experts believe that the weight of the evidence supports a general guideline that consideration be given to initiating therapy in asymptomatic HIV-infected individuals with a CD4⁺ T cell count <350 cells/mm³ or a viral load >55,000 copies/mL. For asymptomatic patients with CD4⁺ T cell counts >350 cells/mm³, rationale exists for both conservative and aggressive approaches to therapy. The conservative approach is based on the recognition that robust immune reconstitution still occurs in most patients who initiate therapy with CD4⁺ T cell counts in the 200–350 cells/mm³ range, and that toxicities and adherence challenges may outweigh benefits of initiating therapy at CD4⁺ T cell counts >350 cells/mm³. In the

conservative approach, high levels of plasma HIV RNA (i.e., >55,000 by bDNA or RT-PCR) are an indication for more frequent monitoring of CD4⁺ T cell counts and plasma HIV RNA levels, but not necessarily for initiation of therapy. In the aggressive approach, asymptomatic patients with CD4⁺ T cell counts >350 cells/mm³ and levels of plasma HIV RNA >55,000 copies/ml would be treated because of the risk of immunologic deterioration and disease progression. The aggressive approach is supported by the observation in many studies that suppression of plasma HIV RNA by antiretroviral therapy is easier to achieve and maintain at higher CD4⁺ T cell counts and lower levels of plasma viral load [6, 33–36]. Long-term clinical outcomes data, however, are not available to fully endorse this approach.

Data conflict regarding sex-specific differences in viral load and CD4⁺ T cell counts (see “[Considerations for Antiretroviral Therapy in Women](#),” p. 75). Several studies [37–43], though not others [44–47], have concluded that after adjustment for CD4⁺ T cell counts, levels of HIV RNA are lower in women compared with men. In those studies that have indicated a possible gender difference in HIV RNA levels, women have had RNA levels that ranged between 0.13 to 0.28 log₁₀ lower than levels observed in men. In two studies of HIV seroconverters, HIV RNA copy numbers were significantly lower in women than in men at seroconversion, but these differences decreased over time, and median viral load in women and men became similar within 5–6 years after seroconversion [38, 39, 43]. Some data suggest that CD4⁺ T cell counts may be higher in women than in men [48]. However, importantly, rates of disease progression do not differ in a sex-dependent manner [41, 43, 49, 50]. Taken together, these data suggest that sex-based differences in viral load occur predominantly during a window of time when the CD4⁺ T cell count is relatively preserved, when treatment is recommended only in the setting of high levels of plasma HIV RNA. Clinicians may wish to consider lower plasma HIV RNA thresholds for initiating therapy in women with CD4⁺ T cell counts >350 cells/mm³, although there are insufficient data to determine an appropriate threshold. In patients with CD4⁺ T cell counts <350 cells/mm³, very small sex-based differences in viral load have been observed; therefore, no changes in treatment guidelines for women are recommended for this group.

In summary, the decision to begin therapy in the asymptomatic patient with >200 CD4⁺ T cells/mm³ is complex and must be made in the setting of careful patient counseling and education. The factors that must be considered in this decision are:

1. the willingness, ability and readiness of the individual to begin therapy;
2. the degree of existing immunodeficiency as determined by the CD4⁺ T cell count;
3. the risk of disease progression as determined by the CD4⁺ T cell count and level of plasma HIV RNA ([Tables 5 and 6](#) and [Figure 1](#); see also reference [1]);
4. the potential benefits and risks of initiating therapy in asymptomatic individuals, including short-term and long-term adverse drug effects; and
5. the likelihood, after counseling and education, of adherence to the prescribed treatment regimen. In this regard, no individual patient should automatically be excluded from consideration for antiretroviral therapy simply because he or she exhibits a behavior or other characteristic judged by some to lend itself to nonadherence. Rather, the likelihood of patient adherence to a long-term, complex drug regimen should be discussed and determined by the individual patient and clinician before therapy is initiated. To achieve the level of adherence necessary for effective therapy, providers are encouraged to utilize strategies for assessing and assisting adherence; in this regard, intensive patient education and support regarding the critical need for adherence should be provided, specific goals of therapy should be established and mutually agreed upon, and a long-term treatment plan should be developed with the patient. Intensive follow up should take place to assess adherence to treatment and to continue patient counseling for the prevention of sexual and drug injection-related transmission (see below “[Adherence to Potent Antiretroviral Therapy](#)”).

CONSIDERATIONS FOR DISCONTINUING THERAPY

As recommendations evolve, patients who have begun highly active antiretroviral therapy at CD4⁺ T cell counts >350/mm³ may wish to discontinue treatment. There are no clinical data addressing whether or not this should be done or can be accomplished safely. Potential benefits include reduction of toxicities, drug-drug interactions, and selection of drug resistant variants, and improvement in the quality of life. Potential risks include rebound in viral replication and renewed immunologic deterioration. If the patient and clinician agree to discontinue therapy the patient should be closely monitored.

ADHERENCE TO POTENT ANTIRETROVIRAL THERAPY

Introduction

These Guidelines call for many people living with HIV, many of whom are asymptomatic, to be treated with highly active antiretroviral therapy (HAART) for the rest of their lives. The ability of the patient to adhere to the regimen is essential for successful treatment. Excellent adherence has been shown to increase the likelihood of sustained virologic control, which is important for reducing HIV-related morbidity and mortality. Conversely, poor adherence has been shown to increase the likelihood of virologic failure and has been associated with increased morbidity and mortality [51, 52]. Poor adherence leads to the development of drug resistance, limiting the effectiveness of therapy [53]. The determinants, measurements, and interventions to improve adherence to HAART are poorly characterized and understood, and more research on this critical topic is needed.

Adherence in HIV Disease

Adherence is an important determinant of both the degree and duration of virologic suppression. In addition, numerous studies have found an association between poor adherence and virologic failure. In several studies, non-adherence in patients on HAART was the strongest predictor of failure to achieve viral suppression below the level of detection [52, 53]. A high degree of adherence is necessary for optimal virologic suppression with HAART; several studies have shown that 90–95% of doses must be taken for optimal suppression, and lesser degrees of adherence are more often associated with virologic failure [51, 54]. To date, there is no conclusive evidence that the degree of adherence required varies with different classes of agents or different specific medications in the HAART regimen.

Imperfect adherence is common. Surveys have shown that one-third of patients missed doses within 3 days of the survey [55]. The reasons for missed doses were predictable and included forgetting, being too busy, being out of town, being asleep, being depressed, having adverse side effects, and being too ill [56]. One fifth of HIV infected patients in one urban center never filled their prescriptions. The instability of homelessness may lead to poor adherence, but not without exception; one recent program achieved a 70% adherence rate among the homeless utilizing flexible

clinic hours, accessible clinical staff, and incentives [57].

Many predictors of poor adherence to HIV medications have been identified. These include

1. poor clinician-patient relationship
2. active drug and alcohol use
3. active mental illness, in particular depression
4. lack of patient education and inability of patients to identify their medications (see ref. [56]), and
5. lack of reliable access to primary medical care or medication [58].

Other sources of instability that may influence adherence include domestic violence and discrimination [58]. Medication side effects may also cause poor adherence [59]. More recently, fear of or the experience of metabolic and morphologic side effects of HAART has been associated with poor adherence [59].

Predictors of good adherence to HIV medications have also been identified. These include

1. availability of emotional and practical life supports;
2. the ability of patients to fit the medications into their daily routine;
3. the understanding that poor adherence leads to resistance;
4. the recognition that taking all medication doses is important; and
5. feeling comfortable taking medications in front of people [60].
6. Importantly, optimal viral suppression is associated with keeping clinic appointments [34].

The measurement of adherence is imperfect and lacking a gold standard. Patient self-report is weakly predictive of the likelihood of adherence; however, an estimate of poor adherence by a patient has a strong predictive value and should be regarded seriously [60, 61]. Clinician estimation of a patient's likelihood of adherence is a poor predictor [62]. Each of several aids to measure adherence, such as pill counts, pharmacy records, smart pill bottles with computer chips recording each opening (i.e. Medication Event Monitoring Systems or “MEMSCaps”), and other devices may be of use, though each requires comparison to patient self-reports [61, 63]. In some studies, clinician and patient estimates of the degree of adherence have been found to exceed measures based on MEMSCaps. Due to its complexity and cost, MEMSCaps technology is best used as an adjunct to adherence research, but is not useful in most clinical settings.

Self-report should include a short term assessment of each dose that was taken over the recent past (e.g., the past 3 days), and a general inquiry regarding adherence since the last visit, with explicit attention to the circumstances of missed doses and possible measures to prevent further missed doses. It may also be helpful for patients to bring their medications and medication diary to clinic visits.

Adherence to HAART: Approach to the Patient

Patient-related strategies

Suggestions for strategies to improve adherence are noted in [Tables 7–9](#). The first principle is to negotiate a treatment plan that the patient understands and to which he/she commits [64, 65]. Before the first prescription is written, patient “readiness” to take medication should be clearly established. Such negotiation takes time, commonly two or three office visits, and patience. Specific education should include the goals of therapy, including a review of expected outcomes based on baseline viral load and CD4⁺ T cell counts (i.e., the Multicenter AIDS Cohort Study (MACS) data from the Guidelines), the reason for the need for adherence, and the plan for and mechanics of adherence. Patients must understand that the first HAART regimen has the best chance of long-term success [1]. The clinician and health team should develop a concrete plan for the specific regimen in question, including the timing of doses of medications around meals and other daily activities. Some centers offer “dry runs” with jelly beans in order to familiarize the patient with the rigors of HAART; however, there are no data to indicate whether or not this exercise improves adherence. Daily or weekly pillboxes, timers with alarms, pagers and other devices may be useful. The development of side effects can affect the ability to adhere to treatment. Clinicians should inform patients in advance about possible side effects and when they are likely to occur; treatment for likely side effects should be included with the first prescription along with instructions on the appropriate response and the possible need to contact the clinician. In some studies, low literacy has been associated with poor adherence; clinicians should take care to assess a patient’s literacy level before relying on written information, and to tailor the adherence intervention to the individual patient. Visual aids and audio or video sources of information may be useful in these patients [66].

Education of family and friends regarding the importance of adherence, as well as recruitment of family and friends to become participants in the plan for medication adherence can be invaluable. Community interventions can be of assistance, including adherence support groups, or the addition of adherence issues to regular support group interactions. Community-based case managers and peer educators can greatly assist adherence education and adherence strategies in individual patients.

Temporary postponement of HAART initiation has been proposed for patients with identified risks for poor adherence [67, 68]. For example, a patient with active substance abuse or mental illness may benefit from immediate psychiatric treatment or treatment for chemical dependency. Appropriate therapy during the 1–2 months needed for treatment of these conditions may be limited to opportunistic infection prophylaxis, if indicated, and therapy directed towards the symptoms of drug withdrawal and detoxification or the underlying mental illness. In addition, readiness for HAART can be assessed and adherence education can be instituted during this time. Other sources of patient instability, such as homelessness, may also be addressed during this interval. Patients should be informed and in agreement with such a plan for future treatment and time-limited treatment deferral.

Clinicians are reminded that such factors as gender, race, socio-economic status, educational level, and a past history of drug use do not reliably predict poor adherence. Conversely, a higher socio-economic status and educational levels and a lack of a history of drug abuse do not predict adequate adherence [68]. No individual patient should automatically be excluded from consideration from antiretroviral therapy simply because he or she exhibits a behavior or other characteristics judged by some to lend itself to nonadherence.

Clinician and health team-related strategies

Clinician and health team-related strategies to enhance adherence are noted in [Table 8](#). A trusting relationship is essential. The clinician should commit to a feasible mechanism for communication between clinic visits, to ongoing monitoring of adherence, and to timely and appropriate responses to adverse events or interim illness. Interim management during clinician vacations or other absences must be clarified.

Adherence requires full deployment of the available health care team. Regular reinforcement by two or more team members will assist the goals of adherence. Provider attitudes and behaviors that are supportive and non-judgmental will encourage patients to be honest about their adherence and about problems they have experienced with adherence. Interventions that have been associated with improved adherence include a pharmacist-based adherence clinic [69], a street-level drop-in center with medication storage and flexible hours for the homeless [70], an adolescent-specific adherence training program [71], and medication counseling and behavioral intervention [72]; these and others are noted in [Table 9](#). For all health care team members, specific training on HAART and adherence should be offered and updated regularly.

Monitoring may identify periods of poor adherence. There is evidence that adherence wanes over time, even in highly adherent patients, a phenomenon described as “pill fatigue” or “treatment fatigue” [67, 73]. Thus monitoring adherence at every clinical encounter is essential. Reasonable responses to decreasing adherence include increasing the intensity of clinical follow up, shortening the follow up interval, and recruiting additional health team members, depending on the nature of the problem [68]. Intermittent drug use or emotional crisis might lead to referral for mental health or chemical dependency assessment or further recruitment and intervention with family or friends. Some patients may require ongoing assistance from support team members from the outset, including chemically dependent patients, mentally retarded patients in the care of another, children and adolescents, or patients in crisis.

New diagnoses or symptoms may influence adherence. For example, depression may require referral, management, and consideration of the short and long-term impact on adherence. Cessation of all medications at the same time may be more desirable than uncertain adherence during a 2-month exacerbation of chronic depression.

The response to the problem of adherence in special populations has not been well studied. There is evidence that programs designed specifically for adolescents, for women and families, for injection drug users, and for homeless persons increase the likelihood of medication adherence [69, 71, 74, 75]. In particular, the incorporation of adherence interventions into convenient primary care settings, the training and deployment of peer educators, pharmacists, nurses, and other health care personnel in adherence interventions, and the monitoring of clinician and patient performance

regarding adherence are beneficial [70, 76, 77]. In the absence of data, a reasonable response is to address and monitor adherence in all HIV primary care encounters and incorporate adherence goals in all patient treatment plans and interventions. This may require the full use of a support team including bilingual providers and peer educators for non-English speaking populations, incorporation of adherence into support group agendas and community forums, and inclusion of adherence goals and interventions into the work of chemical dependency counselors and programs.

Regimen-related strategies

To the extent possible, regimens should be simplified by reducing the number of pills and the frequency of therapy, and by minimizing drug interactions and side effects. This is particularly true for patients with strong biases against many pills and frequent dosing; for some patients, these issues are of lesser importance. There is evidence that simplified regimens with reduced pill numbers and dose frequencies improve adherence [78, 79]. With the numerous effective options for initial therapy noted in these Guidelines and the observed benefit of less frequent dosing on adherence, twice daily dosing of HAART regimens is feasible in most circumstances. Regimens should be chosen with review and discussion of specific food requirements in mind and patient understanding and agreement to such restrictions. Regimens requiring an empty stomach numerous times daily may be difficult for patients with wasting, just as regimens requiring high fat intake may be difficult for patients with lactose intolerance or fat aversion. Fortunately, an increasing number of effective regimens have no specific food requirements.

Directly observed therapy

Directly observed therapy (DOT), in which a health care provider observes the ingestion of medication, has been shown to be successful in the management of tuberculosis, specifically in patients who are poorly adherent to medications. However, it is labor-intensive, expensive, intrusive, and programmatically complex to initiate and complete and, unlike tuberculosis, HIV requires lifelong therapy.

Several pilot programs have studied DOT in HIV patients with some preliminary success [80–83]. Programs have studied once daily regimens in prisons, in methadone programs, and in other cohorts of patients with a record of repeated poor adherence. Modified DOT programs have also been studied, in

which the morning dose is observed and evening and weekend doses are self-administered. The goal of these programs is to improve patient education and medication self-administration over a time-limited (i.e., 3–6 months) period. It is too early to judge the outcomes of these programs, particularly with regard to long term adherence following completion of DOT.

GOALS OF THERAPY

Eradication of HIV infection cannot be achieved with currently available antiretroviral regimens; in large measure, this is due to the establishment of a pool of latently infected CD4⁺ T cells during the very earliest stages of acute HIV infection [84] that persists with an extremely long half-life, even with prolonged suppression of plasma viremia to <50 copies/mL [85–88]. The primary goals of antiretroviral therapy are maximal and durable suppression of viral load, restoration and/or preservation of immunologic function, improvement of quality of life, and reduction of HIV-related morbidity and mortality (Table 10). In fact, adoption of treatment strategies articulated in these guidelines has resulted in substantial reductions in HIV-related morbidity and mortality [89–91].

Plasma viremia is a strong prognostic indicator in HIV infection [3]. Furthermore, reductions in plasma viremia achieved with antiretroviral therapy account for much of the clinical benefit associated with therapy [92]. Therefore, suppression of plasma viremia as much as possible for as long as possible is an important goal of antiretroviral therapy. However, this goal must be balanced against the need to preserve effective treatment options. Switching antiretroviral regimens for any detectable level of plasma viremia may rapidly exhaust treatment options; reasonable parameters that may prompt a change in therapy are discussed below (see “[Criteria for Changing Therapy](#),” p. 22).

HAART often leads to increases in the CD4⁺ T cell count of 100–200 cells/μl or more, although individual responses are quite variable. CD4⁺ T cell responses are generally related to the degree of viral load suppression [93]. In turn, continued viral load suppression is more likely among those who achieve higher CD4⁺ T cell counts during therapy [94]. A favorable CD4⁺ T cell response can occur with incomplete viral load suppression and may not necessarily indicate a poor prognosis [95]. The durability of these immunologic responses that occur with suboptimal suppression of viremia is unknown. Therefore, while viral load is the

strongest single predictor of long-term clinical outcomes, strong consideration should also be given to sustained rises in CD4⁺ T cell counts and partial immune restoration. The urgency of the need to change therapy in the presence of low level viremia is clearly tempered by this observation. The expectation that continuing the existing therapy in this situation will inevitably lead to rapid accumulation of drug resistant virus may not always be realized. One reasonable strategy is maintenance of the regimen, but with redoubled efforts at optimizing adherence, and more frequent monitoring.

Partial reconstitution of immune function induced by HAART may allow for elimination of unnecessary therapies, such as some of those used for prevention and maintenance therapy against opportunistic infections. The appearance of naïve T cells [96, 97], partial normalization of perturbed T cell receptor VB repertoires [98], and evidence of residual thymic function in patients receiving HAART [99, 100] suggest that partial immune reconstitution frequently occurs in these patients. Further evidence of functional immune restoration can be found in the return during HAART of *in vitro* responses to microbial antigens associated with opportunistic infections [101], and the lack of cases of *Pneumocystis carinii* pneumonia (PCP) among patients who discontinued primary PCP prophylaxis when their CD4⁺ T cell counts rose to >200 cells/mm³ during HAART [102–104]. Current guidelines include some recommendations regarding the discontinuation of prophylaxis and maintenance therapy for certain opportunistic infections in the setting of HAART-induced increases in CD4⁺ T cell counts [2].

Tools to Achieve the Goals of Therapy

Although as many as 70–90% of antiretroviral drug-naïve patients achieve maximal viral load suppression 6–12 months after initiation of therapy, only about 50% of patients in a city clinic setting achieve similar results [33, 34]. Predictors of virologic success include low baseline viremia and high baseline CD4⁺ T cell count [33–35], rapid decline of viremia [6], decline of viremia to <50 HIV RNA copies/mL [6], adequate serum levels of antiretroviral drugs [6, 105], and adherence to the drug regimen [34, 51, 106]. While optimal strategies for achieving the goals of antiretroviral therapy have not yet been fully delineated, efforts to improve patient adherence to therapy are likely important (see “[Adherence to Potent Antiretroviral Therapy](#)” p. 9).

Another tool to maximize the benefits of antiretroviral therapy is the rational sequencing of drugs and the preservation of future treatment options for as long as possible. [Table 11](#) shows the possible advantages and disadvantages of three alternative regimens, including a PI with 2 NRTIs, an NNRTI with 2 NRTIs, or a 3 NRTI regimen. The goal of a class-sparing regimen is to preserve or “spare” one or more than one class of drugs for later use. By sequencing drugs in this fashion, it may be possible to extend the overall long-term effectiveness of the available therapy options. Moreover, this strategy makes it possible to selectively delay the risk of certain side effects uniquely associated with a single class of drugs. The efficacy of PI-containing HAART regimens has been demonstrated to include durable viral load suppression, partial immunologic restoration, and decreased incidence of AIDS and death [24–26]. Viral load suppression and CD4⁺ T cell responses that are similar to those observed with PI-containing regimens have been achieved with selected PI-sparing regimens, such as efavirenz + 2 NRTIs [107] or abacavir + 2 NRTIs [108]; however, it is not yet known whether such PI-sparing regimens will provide comparable efficacy with regard to clinical endpoints.

The presence of drug resistant HIV in treatment-experienced patients is a strong predictor of virologic failure and disease progression [109–111]. The results of several prospective studies indicate that the virologic response to a new antiretroviral regimen after virologic failure on a previous regimen can be significantly improved when results of resistance testing were available to guide the choice of drugs in the new regimen [10, 11]. Thus, resistance testing appears to be a useful tool in selecting active drugs when changing antiretroviral regimens in the setting of virologic failure (see [“Testing for Drug Resistance,”](#) p. 5).

INITIATING THERAPY IN THE PATIENT WITH ASYMPTOMATIC HIV INFECTION

When initiating therapy in the patient naïve to antiretroviral therapy, one should begin with a regimen that is expected to achieve sustained suppression of plasma HIV RNA, a sustained increase in CD4⁺ T cell count, and a favorable clinical outcome (i.e., delayed progression to AIDS and death). Additional consideration should be given to the regimen’s pill

burden, dosing frequency, food requirements, convenience, toxicity, and drug interaction profile compared with other regimens. Strongly recommended regimens include either indinavir, nelfinavir, ritonavir + saquinavir, ritonavir + indinavir, ritonavir + lopinavir or efavirenz in combination with one of several 2 NRTI combinations ([Table 12](#)). Clinical outcome data support the use of a PI in combination with 2 NRTIs [24–26] (BI). It should be noted that ritonavir as the sole PI is considered as an alternative agent because of the difficulty many patients have tolerating standard doses of ritonavir [34], and because of the drug’s many interactions. A similar rationale applies to saquinavir-SGC, because of the difficulty many patients have tolerating standard doses and because of the large pill burden associated with its use; however, then there is no reason to switch a patient off of a ritonavir or saquinavir-based regimen if they are tolerating it and if the regimen is effective.

The use of ritonavir to increase plasma concentrations of other protease inhibitors (PIs) has rapidly evolved from an investigational concept to widespread practice. Standard doses of individual PIs result in trough drug levels that are often only slightly higher than the effective antiviral concentration; this may afford an opportunity for viral replication. In contrast, protease “boosting” or “enhancement” by ritonavir increases the trough levels of other protease inhibitors well above the IC₅₀ or IC₉₅, minimizing opportunities for viral replication, and potentially allowing for drug activity even against moderately resistant strains of virus. In addition, these dual PI combinations often lead to more convenient regimens in terms of pill burden, scheduling, and elimination of food restrictions. They also may prevent efavirenz or nevirapine-induced drug interactions.

Ritonavir increases plasma concentrations of other PIs by at least two mechanisms, including inhibition of gastrointestinal CYP450 during absorption, and metabolic inhibition of hepatic CYP450. The 20-fold increase in saquinavir plasma concentrations with ritonavir co-administration is likely caused by inhibition of CYP450 at both sites and leads to a marked increase primarily in the saquinavir C_{max} [112]. For lopinavir, the addition of ritonavir increases both the peak concentration and the half-life (which subsequently results in a higher trough concentration). The result is a lopinavir AUC that is 100-fold higher compared to lopinavir alone [113]. For other PIs, metabolism in the gastrointestinal tract plays a relatively minor role and the enhancement is primarily due to CYP450 inhibition in the liver. The addition of

ritonavir to amprenavir, nelfinavir, or indinavir results in marked increases in half-life and trough levels, with a more moderate or minimal increase in the peak concentration [114, 115].

The dose of ritonavir that is used for PI “boosting” also appears to be important for some protease inhibitors but not others. With saquinavir and amprenavir, increases in the ritonavir dose above 100 mg BID do not significantly increase the PI levels further [114, 116]. However, increasing ritonavir doses above 100 mg BID do appear to provide additional enhancement for indinavir and nelfinavir [115, 117]. While a considerable body of pharmacokinetic data support the use of most of these ritonavir + PI combinations, few efficacy data are available yet for combinations other than ritonavir + saquinavir [118] or ritonavir + lopinavir [119]. In addition, the long term risks and toxicities of dual PI combinations remain unknown.

Disappointing results with antiretroviral regimens prescribed in the setting of virologic failure with a previous regimen suggest that the first regimen affords the best opportunity for long-term control of viral replication. Because the genetic barrier to resistance is greatest with PIs, many would consider a PI + 2 NRTIs to be the preferred initial regimen. However, efavirenz + 2NRTIs appears to be at least as effective as a PI + 2 NRTIs in suppressing plasma viremia and increasing CD4⁺ T cell counts [107], and many would argue that such a regimen is the preferred initial regimen because it may spare the toxicities of PIs for a considerable time (BII). Although no direct comparative trials exist that would allow a ranking of the relative efficacy of the NNRTIs, the demonstrated ability of efavirenz in combination with 2 NRTIs to suppress viral replication and increase CD4⁺ T cell counts to a similar degree as a PI with 2 NRTIs support a preference for efavirenz over the other available NNRTIs at this time. Abacavir + 2 NRTIs, a triple NRTI regimen, has been used with some success as well [108] (CII). Such a regimen, however, may have short-lived efficacy when the baseline viral load is >100,000 copies/mL. Using 2 NRTIs alone does not achieve the goal of suppressing viremia to below detectable levels as consistently as does a regimen in the “strongly recommended” or “alternative” categories and should be used only if more potent treatment is not possible (DI). Use of antiretroviral agents as monotherapy is contraindicated (DI), except when there are no other options, or in pregnancy to reduce perinatal transmission as noted below. When initiating antiretroviral therapy, all drugs should be started simultaneously at full dose with the

following three exceptions: dose escalation regimens are recommended for ritonavir, nevirapine, and, in some cases, ritonavir plus saquinavir.

Hydroxyurea has been used investigationaly in combination with antiretroviral agents for treatment of HIV infection, however its utility in this setting has not been established. Clinicians considering use of hydroxyurea in a treatment regimen for HIV should be aware of the limited and conflicting nature of data in support of its efficacy, and the importance of monitoring patients closely for potentially serious toxicity (See “[Hydroxyurea](#)”, p. 77).

Detailed information comparing the different nucleoside RT inhibitors, non-nucleoside RT inhibitors, the protease inhibitors, and drug interactions between the protease inhibitors and other agents can be found in [Tables 13–20](#). Particular attention should be paid to [Tables 17–19](#) regarding drug interactions between the protease inhibitors and other agents, as these are extensive and often require dose modification or substitution of various drugs. Toxicity assessment is an ongoing process; assessment at least twice during the first month of therapy and every 3 months thereafter is a reasonable management approach.

INITIATING THERAPY IN ADVANCED HIV DISEASE

All patients diagnosed with advanced HIV disease, which is defined as any condition meeting the 1993 CDC definition of AIDS [23] should be treated with antiretroviral agents regardless of plasma viral levels (AI). All patients with symptomatic HIV infection without AIDS, defined as the presence of thrush or unexplained fever, should also be treated. When the patient is acutely ill with an OI or other complication of HIV infection, the clinician should consider clinical issues such as drug toxicity, ability to adhere to treatment regimens, drug interactions, and laboratory abnormalities when determining the timing of initiation of antiretroviral therapy. Once therapy is initiated, a maximally suppressive regimen should be used, as indicated in [Table 12](#). Advanced stage patients being maintained on an antiretroviral regimen should not have the therapy discontinued during an acute opportunistic infection or malignancy, unless there are concerns regarding drug toxicity, intolerance, or drug interactions.

Patients who have progressed to AIDS are often treated with complicated combinations of drugs and the potential for multiple drug interactions must be appreciated by clinician and patient. Thus, the choice of which antiretroviral agents to use must be made with consideration given to potential drug interactions and overlapping drug toxicities, as outlined in [Tables 13–20](#). For instance, the use of rifampin to treat active tuberculosis is problematic in a patient receiving a protease inhibitor, which adversely affects the metabolism of rifampin but is frequently needed to effectively suppress viral replication in these advanced patients. Conversely, rifampin lowers the blood level of protease inhibitors, which may result in suboptimal antiretroviral therapy. While rifampin is contraindicated or not recommended for use with all of the protease inhibitors, one might consider using rifabutin at a reduced dose, as indicated in [Table 18](#); this topic is discussed in greater detail elsewhere [120]. Other factors complicating advanced disease are wasting and anorexia, which may prevent patients from adhering to the dietary requirements for efficient absorption of certain protease inhibitors. Bone marrow suppression associated with ZDV and the neuropathic effects of ddC, d4T and ddI may combine with the direct effects of HIV to render the drugs intolerable. Hepatotoxicity associated with certain protease inhibitors may limit the use of these drugs, especially in patients with underlying liver dysfunction. The absorption and half-life of certain drugs may be altered by antiretroviral agents, particularly the protease inhibitors and NNRTIs whose metabolism involves the hepatic cytochrome p450 (CYP450) enzymatic pathway. PIs inhibit the CYP450 pathway, whereas NNRTIs have variable effects; nevirapine is an inducer, delavirdine is an inhibitor, and efavirenz is a mixed inducer/inhibitor. CYP450 inhibitors have the potential to increase blood levels of drugs metabolized by this pathway. At times, adding a CYP450 inhibitor can improve the pharmacokinetic profile of selected agents (such as adding ritonavir therapy to saquinavir) as well as contribute an additive antiviral effect; however, these interactions can also result in life threatening drug toxicity, as indicated in [Tables 17–20](#). Thus, health care providers should inform their patients of the need to discuss any new drugs, including over the counter agents and alternative medications, that they may consider taking, and careful attention should be given to the relative risks versus benefits of specific combinations of agents.

Initiation of potent antiretroviral therapy is often associated with some degree of recovery of immune function. In this setting, patients with advanced HIV

disease and subclinical opportunistic infections such as MAI or CMV may develop a new immunologic response to the pathogen and thus new symptoms may develop in association with the heightened immunologic and/or inflammatory response. This should not be interpreted as a failure of antiretroviral therapy and these newly presenting opportunistic infections should be treated appropriately while maintaining the patient on the antiretroviral regimen. Viral load measurement is helpful in clarifying this situation.

HAART-ASSOCIATED ADVERSE CLINICAL EVENTS

Potential adverse events associated with antiretroviral agents are outlined in [Tables 13–16](#). A summary of Box warnings is provided in [Table 16](#). Drug interactions of concern are listed in [Tables 17–18](#).

Lactic Acidosis/ Hepatic Steatosis

Chronic compensated hyperlactatemia can occur during treatment with nucleoside/nucleotide analogue reverse transcriptase inhibitors (NRTIs) [121, 122]. Although cases of severe decompensated lactic acidosis with hepatomegaly and steatosis are rare (estimated incidence of 1.3 cases per 1000 person-years of NRTI exposure), this syndrome is associated with a high mortality rate [123–126]. Severe lactic acidosis with or without pancreatitis, including three fatal cases, were reported in later-stage pregnant or postpartum women and women whose antiretroviral therapy during pregnancy included stavudine and didanosine in combination with other antiretroviral agents [125, 127, 128]. Other risk factors for the development of this toxicity include female gender, obesity, and prolonged use of NRTIs, although some cases have been reported to occur without known risk factors [125].

The mitochondrial basis of NRTI-induced lactic acidosis and hepatic steatosis is one possible mechanism of cellular injury as NRTIs also inhibit DNA polymerase gamma, which is the enzyme responsible for mitochondrial DNA synthesis. The ensuing mitochondrial dysfunction may also result in several other adverse effects such as pancreatitis, peripheral neuropathy, myopathy and cardiomyopathy [129]. It has also been hypothesized that some features of the lipodystrophy syndrome are also tissue-specific mitochondrial toxicities caused by NRTI treatment [130–132].

The initial clinical presentations of patients with the lactic acidosis syndrome are variable and may include nonspecific gastrointestinal symptoms without dramatic elevation of hepatic enzymes [133]. The clinical “prodrome” may include otherwise unexplained onset and persistence of abdominal distention, nausea, abdominal pain, vomiting, diarrhea, anorexia, dyspnea, generalized weakness, ascending neuromuscular weakness, myalgias, paresthesias, weight loss and hepatomegaly [134]. In addition to hyperlactatemia, laboratory evaluation may reveal an increased anion gap ($\text{Na} - [\text{Cl} + \text{CO}_2] > 16$), elevated aminotransferases, CPK, LDH, lipase and amylase [124, 133, 135].

Echotomography and CT scans may demonstrate an enlarged fatty liver; histopathological examination of the liver reveals microvesicular steatosis [133]. Since there are significant technical problems associated with lactate testing, routine monitoring of lactate level is not generally recommended. One must first rely on other laboratory abnormalities PLUS symptoms when lactic acidosis is suspected. Measurement of lactate requires a standardized mode of sample handling including pre-chilled fluoride-oxalate tubes which should be transported immediately on ice to the laboratory and processed within four hours of collection; blood should be collected without the use of a tourniquet, without fist-clenching, and, if possible, without stasis [136, 137]. When interpreting serum lactate, levels from 2 to 5 mmol/dL are considered elevated and need to be correlated with symptoms, levels above 5 mmol/dL are clearly abnormal, and levels above 10 mmol/dL indicate serious and possibly life-threatening situation. Some experts also suggest monitoring of serum bicarbonate and electrolytes for the early identification of an increased anion gap every three months.

In some cases the adverse event has resolved after discontinuation of NRTIs [133, 138], and some patients have tolerated rechallenge with a new NRTI-containing regimen [133, 139]; however, at present there are insufficient data to recommend this strategy versus treatment with an NRTI-sparing regimen. If NRTI treatment is not discontinued, in some patients progressive mitochondrial toxicity may produce severe lactic acidosis manifested clinically by tachypnea and dyspnea; respiratory failure may follow, requiring mechanical ventilation. In addition to discontinuation of antiretroviral treatment and intensive therapeutic strategies that include bicarbonate infusions and hemodialysis [140] (AD), some clinicians have administered thiamine [141] and riboflavin [127], based upon the pathophysiological hypothesis that sustained cellular dysfunctions of the mitochondrial respiratory chain cause this fulminant clinical

syndrome; the efficacy of these latter interventions requires clinical validation.

In conclusion, antiretroviral treatment should be suspended if clinical and laboratory manifestations of the lactic acidosis syndrome occur (BIII).

Hepatotoxicity

Hepatotoxicity, defined as at least three to five times increase in serum transaminases (AST, ALT, GGT) with or without clinical hepatitis, has been reported in patients receiving HAART. All the currently marketed NNRTIs and PIs have been associated with serum transaminase elevation. Most of the cases are asymptomatic, some of which resolve spontaneously without therapy interruption or modification [142]. Hepatic steatosis in the presence of lactic acidosis is a rare but serious adverse effect associated with the nucleoside analogs. A more detailed discussion of this syndrome can be seen in the “[Lactic Acidosis/Hepatic Steatosis](#),” section of this document (on p. 16).

Among the NNRTIs, nevirapine has the greatest potential of causing clinical hepatitis. Martinez and colleagues reported an incidence of 12.5% of hepatotoxicity in patients initiating nevirapine, with clinical hepatitis seen in 1.1% of these patients [143]. In a randomized trial in Africa in which stavudine was the backbone NRTI, and either nevirapine or efavirenz was added to emtricitabine or lamivudine, 9.4% of the nevirapine-treated patients developed grade 4 liver enzyme elevation as compared to none of the efavirenz-treated patients. Two of these patients died of liver failure. The incidence in female patients was twice that seen in male patients (12% versus 6%, $p=0.05$) [144]. Nevirapine-associated hepatitis may also be present as part of a hypersensitivity syndrome, with a constellation of other symptoms such as skin rash, fever, and eosinophilia. Approximately two-thirds of the cases of nevirapine-associated clinical hepatitis occur within the first 12 weeks of therapy. Fulminant and even fatal cases of hepatic necrosis have been reported. The patients may present with non-specific gastrointestinal and flu-like symptoms with or without liver enzyme abnormalities. The syndrome can progress rapidly to hepatomegaly, jaundice, and hepatic failure within a few days [145]. A two-week lead-in dosing of nevirapine with 200mg once daily before dose escalation to twice daily may reduce the incidence of hepatotoxicity. In light of the potential severity of clinical hepatitis, some experts advise close monitoring of liver enzymes and clinical symptoms

after nevirapine initiation, e.g. every two weeks for the first month, then monthly for 12 weeks, and every one to three months thereafter. Patients who experience severe clinical hepatotoxicity while on nevirapine should not be rechallenged.

Unlike the early-onset hepatotoxicity seen with nevirapine, PI-associated liver enzyme abnormalities can occur any time during the course of treatment. In a retrospective review, severe hepatotoxicity (defined as >5x increase over baseline ALT or AST) was seen more often in patients receiving ritonavir- or ritonavir/saquinavir-containing regimens than those receiving other PIs (indinavir, nelfinavir, or saquinavir) [146]. Co-infection with hepatitis C virus is reported to be a major risk factor for development of hepatotoxicity after PI initiation [147, 148]. Other potential risk factors for hepatotoxicity include hepatitis B infection [142, 147, 149], alcohol abuse [148], baseline elevated liver enzymes [150], stavudine use [149], and concomitant use of other hepatotoxic agents. HAART-induced immune reconstitution rather than direct liver toxic effects of the PIs have been suggested as the cause of liver decompensation in hepatitis C or hepatitis B coinfecting patients.

Hyperglycemia

Hyperglycemia, new onset diabetes mellitus, diabetic ketoacidosis, and exacerbation of pre-existing diabetes mellitus have been reported in patients receiving HAART [151–153]. These metabolic derangements are strongly associated with PI use [154], though they may occur independently of PI use as well [155]. The incidence of new onset hyperglycemia was reported as 5% in a 5-year historical cohort analysis in a population of 221 HIV-infected patients. PIs were independently associated with hyperglycemia and it did not vary significantly by PIs [156]. It was also noted that viral load suppression and increase in body weight did not reduce the magnitude of the association with PIs. The pathogenesis of these abnormalities has not been fully elucidated; however, it appears that hyperglycemia may result from peripheral and hepatic insulin resistance, relative insulin deficiency, an impaired ability of the liver to extract insulin, and a longer exposure to antiretroviral medications [157, 158]. Hyperglycemia with or without diabetes has been reported in 3–17% of patients in various retrospective studies. Among these reports, symptoms of hyperglycemia were reported at a median of approximately 60 days, ranging from 2–390 days following initiation of PI therapy. Hyperglycemia resolved in some patients who discontinued PI therapy;

however, the reversibility of these events is currently unknown due to limited data. Some patients continued PI therapy and initiated treatment with oral hypoglycemic agents or insulin. Clinicians are advised to monitor HIV-infected patients with pre-existing diabetes closely when PIs are prescribed, and to be aware of the risk for drug-related new-onset diabetes in patients without a history of diabetes (BIII). Patients should be advised about the warning signs of hyperglycemia (i.e., polydipsia, polyphagia, and polyuria) and the need to maintain ideal body weight when these medications are prescribed. Some experts recommend routine fasting blood glucose measurements at 3–4 month intervals during the first year of PI treatment in patients with no prior history of diabetes (CIII). Routine use of glucose tolerance tests to detect this complication is not recommended (DIII). Because pregnancy is an independent risk factor for impaired glucose tolerance, closer monitoring of blood glucose levels should be done in pregnant women on PI-containing regimens. There are no data to aid in the decision to continue or discontinue drug therapy in cases of new-onset or worsening diabetes; however, most experts would recommend continuation of HAART in the absence of severe diabetes (BIII). Several studies have attempted to examine the potential of reversing insulin resistance after switching from PI containing HAART regimens to NNRTI-based regimens; the results have been inconclusive.

Fat Maldistribution

Changes in body fat distribution, sometimes referred to as “lipodystrophy syndrome” or “pseudo-Cushing's syndrome” have been observed in 6 to 80 percent of patients receiving HAART; the wide range of estimates of the incidence of this syndrome reflects the lack of a uniform case definition and other variables that are poorly understood. The morphologic changes occur gradually, and are generally not apparent until months after the initiation of HAART. Clinical findings include central obesity, peripheral fat wasting, and lipomas; pathologic changes may include visceral fat accumulation, dorsocervical fat accumulation (“buffalo hump”), extremity wasting with venous prominence, facial thinning, and breast enlargement [159–162]. Some patients may have a cushingoid appearance despite the absence of confounding medications (i.e., corticosteroids) or adrenal function laboratory abnormalities [160]. Hyperlipidemia and insulin resistance are frequently but not always associated with lipodystrophy [163]; it is unclear whether these various clinical manifestations represent distinct entities with

different etiologies, or whether they occur as a result of a single pathologic process. Lipodystrophy has been associated with the use of PIs [159, 163], but may occur with NRTI therapy [160, 164, 165] or in the absence of therapy [166]. Compared with PI-associated lipodystrophy, the NRTI-associated syndrome(s) may be associated with recent onset fatigue and nausea; weight loss; higher levels of lactate and alanine aminotransferase; and lower levels of albumin, cholesterol, triglycerides, glucose, and insulin [165]. Therapeutic strategies aimed at reversing or halting the progression of lipodystrophy include switching classes of antiretroviral agents [167, 168] exercise training [169]; however, insufficient data are currently available to guide the management of lipodystrophy.

Hyperlipidemia

Changes in triglycerides and/or cholesterol have occurred with or without the clinical findings of fat maldistribution, and may occur during the first month of HAART [154, 155, 159, 163]. In clinical studies all PIs have been implicated; however, increases in cholesterol and triglyceride levels may be more dramatic during treatment with ritonavir [170]. The mechanism of these effects has not yet been defined, but may be due in part to interference by protease inhibitors with normal cellular proteins involved in lipid metabolism [171]. Although the long-term consequences of dysregulated lipid metabolism are unknown, substantial increases in triglycerides or cholesterol are of concern because of the possible association with cardiovascular events and pancreatitis. In this regard, case reports have appeared describing premature coronary artery disease, cerebrovascular disease, pancreatitis and cholelithiasis in patients receiving PI therapy. Controlled studies have not yet demonstrated an increased risk of cardiovascular events associated with PI therapy; however, longer follow-up time will be needed to accurately assess this issue [172, 173]. Some experts recommend monitoring serum levels of cholesterol and triglycerides (preferably fasting) at 3–4 month intervals during PI therapy (CIII). For individuals with elevated triglyceride levels at baseline and who may be at increased risk of pancreatitis, it is preferable to repeat a lipid profile sooner (e.g., within 1–2 months of initiating HAART). Assessment should include evaluation for independent risks for cardiovascular disease (i.e., family history, medical history, smoking, diet, weight, etc.) and the magnitude of lipid changes. Intervention is often recommended for triglyceride levels > 750–1000 mg/dL and/or LDL cholesterol levels > 130 mg/dL (in individuals without known

coronary disease and with 2 or more coronary risk factors) or >160 mg/dL (in individuals without known coronary disease and with fewer than 2 coronary risk factors) [174]. The effectiveness of lifestyle modifications (i.e., dietary changes, exercise, and smoking cessation) and lipid lowering drugs such as gemfibrozil, niacin, and the HMG coenzyme A reductase inhibitors (i.e., “statins”) is not clear. Concurrent use of PIs and statins should be undertaken with caution due to the potential for enhanced statin-related toxicity in this setting (Tables 13 and 14). Some patients have had resolution of serum lipid abnormalities following discontinuation of PIs and substitution of PI-sparing antiretroviral regimens; however, this decision requires a risk-benefit analysis.

Increased Bleeding Episodes in Patients with Hemophilia

Increased spontaneous bleeding episodes in patients with hemophilia A and B have been observed with the use of protease inhibitors [175]. Most of the reported episodes involved joints and soft tissues; however, more serious bleeding episodes including intracranial and gastrointestinal bleeding have been reported. The bleeding episodes occurred a median of 22 days after initiation of protease inhibitor therapy. Some patients received additional coagulation factor while continuing protease inhibitor therapy.

Osteonecrosis, Osteopenia, and Osteoporosis

Avascular necrosis and decreased bone density are now recognized as emerging metabolic complications of HIV infection that may be linked to highly active antiretroviral regimens. Both of these bone abnormalities have been reported in adults and children with HIV infection who are now surviving longer with their disease in part because of HAART [176–178].

Avascular necrosis involving the hips was first described in HIV-infected adults and more recently in HIV-infected children (known as Legg Calve Parthese Disease). The diagnosis of osteonecrosis is usually made by CT scan or MRI, when these studies are performed in response to patient complaints of pain in an affected hip or spine. However, asymptomatic disease with MRI findings can occur in 5% of HIV patients [179]. It does not appear that avascular necrosis is clearly associated with a specific antiretroviral regimen in HIV-infected adults, but it has been linked to corticosteroid use in some patients [180]. Factors

associated with osteonecrosis include alcohol abuse, hemoglobinopathies, corticosteroid treatment, hyperlipidemia, and hypercoagulability states. The occurrence of hyperlipidemia suggests an indirect link between antiretroviral therapy and the occurrence of osteonecrosis in HIV-infected patients; however, prospective clinical studies will be required to establish this association. There is no generally accepted medical therapy for avascular necrosis, and surgery sometimes becomes necessary for disabling symptoms.

The decrease in bone mineral density (BMD)—both moderate (osteopenia) and severe (osteoporosis)—is a reflection of the competing effects of bone reabsorption by osteoclast and bone deposition by osteoblast and is measured by bone densitometry. Prior to HAART, there were reports of marginal decreases in BMD in HIV-infected individuals [181]. Decreased bone formation and turnover has been demonstrated with more potent antiretroviral therapy, in particular protease inhibitors [182]. Studies of bone demineralization in a limited number of patients receiving HAART have shown that up to 50% of patients receiving a PI-based regimen developed evidence of osteopenia compared to 20% of patients who are untreated or receiving a non-PI-containing regimen [183]. Other studies have shown that patients with lipodystrophy with extensive prior PI therapy had associated findings of osteopenia (28%) or osteoporosis (9%) [184]. The preliminary observations that there are increased serum and urinary markers of bone turnover in patients on protease containing HAART who have osteopenia support the possible link of bone abnormalities to other metabolic abnormalities observed in HIV-infected patients [185, 186]. There is no recommendation at the present time for routine measurement of bone density in asymptomatic patients by dual energy X-ray absorptiometry (DEXA) or by newer measurements such as quantitative ultrasound (QUS). Specific prophylaxis or treatment recommendations to prevent more significant osteoporosis have not been developed for HIV-infected patients with osteopenia.

Based on experience in the treatment of primary osteoporosis, it would be reasonable to recommend adequate intake of calcium and vitamin D and appropriate weight bearing exercise. When fractures occur or osteoporosis is documented, more specific and aggressive therapies with bisphosphonates, raloxifene or calcitonin may be indicated [187]. Hormone replacement therapy including estrogen may be considered in the setting of significant decreased bone density in postmenopausal women on HAART.

Skin Rash

Skin rash occurs most commonly with the NNRTI class of drugs. Most cases are mild to moderate in nature, occurring within the first few weeks of therapy. Some experts suggest managing skin rash with antihistamines for symptomatic relief without drug discontinuation, although the wisdom of treating through such rashes has been questioned [188]. More serious cutaneous manifestations such as Stevens-Johnson syndrome (SJS) and toxic epidermal necrosis (TEN) should result in the prompt and permanent discontinuation of NNRTIs or other offending agents.

Most reactions resulting in skin rash are confined to cutaneous reactions. However, a severe or even life-threatening syndrome of drug rash with eosinophilia and systemic symptoms (DRESS) has also been described [189, 190]. The systemic symptoms may include fever, hematological abnormalities, and multiple organ involvement.

Among the NNRTIs, skin rash occurs more frequently and in greater severity with nevirapine. Using a two-week lead-in dose escalation schedule when initiating nevirapine therapy may reduce the incidence of rash. In a case-controlled multi-national study, SJS and/or TEN were reported in 18 HIV-infected patients. Fifteen of the 18 patients were receiving nevirapine. The median time from initiation of nevirapine to onset of cutaneous eruption was 11 days, with two-thirds of the cases occurring during the initial dosing period [188]. Female patients appear to have as much as a seven-fold higher risk for developing grade 3 or 4 skin rashes than male patients [191, 192]. The use of systemic corticosteroid or antihistamine therapy at the time of the initiation of nevirapine to prevent development of skin rash has not proven effective [192, 193]. In fact, a higher incidence of skin rash has been reported in the steroid- or antihistamine-treated patients. At present, prophylactic use of corticosteroids should be discouraged.

Skin rash appears to be a “class adverse reaction” of the NNRTIs. The incidence of cross hypersensitivity reactions between these agents is not known. In a small number of reports, patients with prior histories of nevirapine-associated skin rash had been able to tolerate efavirenz without increased rates of cutaneous reactions [194, 195]. Most experts would not recommend the use of another NNRTI in those patients who experienced SJS or TEN with one NNRTI. Initiating NNRTI in a patient with a history of mild to moderate skin rash with another NNRTI should be done with caution and close follow-up.

Among the NRTIs, skin rash occurs most frequently with abacavir. Skin rash may be one of the presenting symptoms of abacavir-associated systemic hypersensitivity reaction, in which case therapy should be discontinued without rechallenge.

Of the PIs, skin rash occurs most frequently with amprenavir, with an incidence of up to 27% in clinical trials. Although amprenavir is a sulfonamide, the potential of cross reactivity between amprenavir and other sulfa drugs is not known. As a result, amprenavir should be used with caution in patients with a history of sulfa allergies.

INTERRUPTION OF ANTIRETROVIRAL THERAPY

There are multiple reasons for temporary or permanent discontinuation of antiretroviral therapy. If there is a need to discontinue any antiretroviral medication, clinicians and patients should be aware of the theoretical advantage of stopping all antiretroviral agents simultaneously, rather than continuing one or two agents, to minimize the emergence of resistant viral strains. If a decision is made to interrupt therapy the patient should be closely monitored including clinical and laboratory evaluations. Prophylaxis should be initiated as needed based on the CD4⁺ T cell count.

There has been interest in what is commonly called structured or supervised treatment interruptions (STI). The concepts underlying STI vary depending on patient populations and encompass at least three major strategies:

1. STI as part of *salvage* therapy,
2. STI for “*auto-immunization*” and better immune control of HIV, and
3. STI for the sole purpose of allowing *less total time on antiretroviral therapy*.

As a consequence of limited data, currently, none of these approaches can be recommended.

Salvage STI is directed at patients whose virus has developed significant antiretroviral drug resistance and who have persistent plasma viremia and relatively low CD4⁺ T cell counts despite receiving therapy. The theoretical goal of STI in this patient population is to allow for the re-emergence of HIV that is susceptible to antiretroviral therapy. Although HIV that was sensitive to antiretroviral agents was detected in the plasma of many individuals following several weeks to months of interrupting treatment, the emergence of drug-sensitive HIV was associated with a significant

decline in CD4⁺ T cells and a significant increase in plasma viremia, suggesting improved replicative fitness and pathogenicity of wild type virus [196]. In addition, drug resistant HIV persisted in CD4⁺ T cells. The observed decrease in CD4⁺ T cells is of particular concern in this patient population, and STI cannot currently be recommended for this group of patients.

Auto-immunization STI and STI for the reduction of total time on drugs are directed at individuals who have maintained suppression of plasma viremia below the limit of detection for prolonged periods of time and who have relatively high CD4⁺ T cell counts. The theoretical goal of auto-immunization STI is to allow several short bursts of viral replication to augment HIV-specific immune responses. This strategy is being studied in individuals who began HAART during either the very early or chronic stages of HIV infection [197–199]. STI for the purpose of less time on therapy utilizes predetermined periods of long or short cycle intermittent antiretroviral therapy. The numbers of patients and duration of follow-up are not sufficient for adequate evaluation of these approaches at this time. Potential risks include a decline in CD4⁺ T cell counts, an increase in transmission and the development of drug resistance.

Due to insufficient data in the situations discussed, STI cannot be recommended for use in general clinical practice at this time. Further research is necessary in each of these areas.

CONSIDERATIONS FOR CHANGING A FAILING REGIMEN

As with the initiation of antiretroviral therapy, the decision to change regimens should be approached with careful consideration of several complex factors. These factors include: recent clinical history and physical examination; plasma HIV RNA levels measured on two separate occasions; absolute CD4⁺ T lymphocyte count and changes in these counts; assessment of adherence to medications; remaining treatment options; potential resistance patterns from prior antiretroviral therapies; and preparation of the patient for the implications of the new regimen which include side effects, drug interactions, dietary requirements and possible need to alter concomitant medications. Failure of a regimen may occur for many reasons, including initial viral resistance to one or more agents, altered absorption or metabolism of the drug, multi-drug pharmacokinetics that adversely affect therapeutic drug levels, and poor patient adherence to a

regimen. In this regard, it is important to carefully assess patient adherence prior to changing antiretroviral therapy; health care workers involved in the care of the patient, such as the case manager or social worker, may be of assistance in this evaluation. Clinicians should be aware of the prevalence of mental health disorders and psychoactive substance use disorders in certain HIV-infected persons; inadequate mental health treatment services may jeopardize the ability of such individuals to adhere to their medical treatment. Proper identification of and intervention in these mental health disorders can greatly enhance adherence to medical HIV treatment.

It is important to distinguish between the need to change therapy due to drug failure versus drug toxicity. In the latter case, it is appropriate to substitute one or more alternative drugs of the same potency and from the same class of agents as the agent suspected to be causing the toxicity. In the case of drug failure where more than one drug had been used, a detailed history of current and past antiretroviral medications, as well as other HIV-related medications, should be obtained. Testing for antiretroviral drug resistance may also be very helpful in maximizing the number of active drugs in a regimen (see "[Testing for Drug Resistance](#)," p. 5). Viral resistance to antiretroviral drugs is an important, but not the only, reason for treatment failure. Genetically distinct viral variants emerge in each HIV-infected individual over time after initial infection. Viruses with single drug resistant mutations exist even prior to therapy, but are selected for replication by antiviral regimens that are only partially suppressive. The more potent a regimen is in durably suppressing HIV replication, the less likely the emergence of resistant variants. Thus the goal of therapy should be to reduce plasma HIV RNA to below detectable limits using the most sensitive assay available (<50 copies/mL), thereby providing the strongest genetic barrier possible to the emergence of resistance.

Three different populations of patients should be considered with regard to a change in therapy:

1. Individuals who are receiving incompletely suppressive antiretroviral therapy, such as single or double nucleoside therapy, with detectable or undetectable plasma viral load (discussed further below);
2. Individuals who have been on potent combination therapy and whose viremia was initially suppressed to undetectable levels but has again become detectable; and

3. Individuals who have been on potent combination therapy and whose viremia was never suppressed to below detectable limits.

CRITERIA FOR CHANGING THERAPY

The goal of antiretroviral therapy, to improve the length and quality of the patient's life, is likely best accomplished by maximal suppression of viral replication to below detectable levels (currently defined as <50 copies/mL) sufficiently early to preserve immune function. However, this is not always achievable with a given therapeutic regimen and frequently regimens must be modified. In general, the plasma HIV RNA level is the most important parameter to evaluate response to therapy, and increases in levels of viremia that are significant, confirmed and not attributable to intercurrent infection or vaccination indicate failure of the drug regimen regardless of changes in the CD4⁺ T cell counts. Clinical complications and sequential changes in CD4⁺ T cell count may complement the viral load test in evaluating a response to treatment. Specific criteria that should prompt consideration for changing therapy include:

- *Less than a 0.5–0.75 log₁₀ reduction in plasma HIV RNA by 4 weeks following initiation of therapy, or less than a 1 log₁₀ reduction by 8 weeks (CIII);*
- *Failure to suppress plasma HIV RNA to undetectable levels within 4–6 months of initiating therapy (BIII).* In this regard, the degree of initial decrease in plasma HIV RNA and the overall trend in decreasing viremia should be considered. For instance, a patient with 10⁶ viral copies/mL prior to therapy who stabilizes after 6 months of therapy at an HIV RNA level that is detectable but <10,000 copies/mL may not warrant an immediate change in therapy.
- *Repeated detection of virus in plasma after initial suppression to undetectable levels, suggesting the development of resistance (BIII).* However the degree of plasma HIV RNA increase should be considered; the clinician may consider short-term further observation in a patient whose plasma HIV RNA increases from undetectable to low-level detectability (e.g., 50–5000 copies/mL) at 4 months. In this situation the patient should be followed very closely. It should be noted, however, that most patients who fall into this category will subsequently show progressive increases in plasma viremia that will likely require a change in the antiretroviral regimen.

- *Any reproducible significant increase, defined as 3-fold or greater, from the nadir of plasma HIV RNA not attributable to intercurrent infection, vaccination, or test methodology except as noted above (BIII);*
- *Undetectable viremia in the patient receiving double nucleoside therapy (BIII).* Patients currently receiving 2 NRTIs who have achieved the goal of no detectable virus have the option of continuing this regimen or may have modification to conform to regimens in the strongly recommended category ([Table 12](#)). Prior experience indicates that most of these patients on double nucleoside therapy will eventually have virologic failure with a frequency that is substantially greater compared to patients treated with the strongly recommended regimens.
- *Persistently declining CD4⁺ T cell numbers, as measured on at least two separate occasions (CIII);*
- *Clinical deterioration (DIII).* In this regard, a new AIDS-defining diagnosis that was acquired after the time treatment was initiated suggests clinical deterioration but may or may not suggest failure of antiretroviral therapy. If the antiretroviral effect of therapy was poor (e.g., <10-fold reduction in viral RNA), then a judgment of therapeutic failure could be made. However, if the antiretroviral effect was good but the patient was already severely immunocompromised, the appearance of a new opportunistic disease may not necessarily reflect a failure of antiretroviral therapy, but rather a persistence of severe immunocompromise that did not improve despite adequate suppression of virus replication. Similarly, an accelerated decline in CD4⁺ T cell counts suggests progressive immune deficiency providing there are sufficient measurements to assure quality control of CD4⁺ T cell measurements.

A final consideration in the decision to change therapy is the recognition of the still limited choice of available agents and the knowledge that a decision to change may reduce future treatment options for the patient. This may influence the clinician to be somewhat more conservative when deciding to change therapy. Consideration of alternative options should include potency of the substituted regimen and probability of tolerance of or adherence to the alternative regimen. Clinical trials have shown that partial suppression of virus is superior to no suppression of virus. On the other hand, some clinicians and patients may prefer to suspend treatment in order to preserve future options or because a sustained antiviral effect cannot be achieved. Referral to or consultation with an experienced HIV clinician is appropriate when one is considering a

change in therapy. When possible, patients requiring a change in an antiretroviral regimen but without treatment options using currently approved drugs should be referred for consideration for inclusion in an appropriate clinical trial.

THERAPEUTIC OPTIONS WHEN CHANGING ANTIRETROVIRAL THERAPY

Recommendations for changes in treatment differ according to the indication for the change. If the desired virologic objectives have been achieved in patients who have intolerance or toxicity, there should be substitution for the offending drug, preferably using an agent in the same class with a different toxicity or tolerance profile. If virologic objectives have been achieved, but the patient is receiving a regimen not in the preferred category (such as two NRTIs or monotherapy), there is the option to continue treatment with careful monitoring of viral load or to add drugs to the current regimen to comply with strongly recommended treatment regimens. As discussed above, most authorities feel that treatment with regimens not in the strongly recommended or alternative categories is associated with eventual failure and recommend the latter tactic.

At present there are very few clinical data to support specific strategies for changing therapy in patients who have failed the strongly recommended regimens; however, a number of theoretical considerations should guide decisions. Because of the relatively rapid mutability of HIV, viral strains with resistance to one or more agents often emerge during therapy, particularly when viral replication has not been maximally suppressed. Of major concern is the possibility of broad cross-resistance among drugs within a class. Evidence indicates that viral strains that become resistant to one PI or NNRTI often have reduced susceptibility to most or all other PIs or NNRTIs.

[Table 21](#) summarizes some of the most important guidelines to follow when changing a patient's antiretroviral therapy. As stated above, a change in regimen because of treatment failure should ideally be guided by results of resistance testing. Dose modifications may be required to account for drug interactions when using combinations of PIs or a PI and NNRTI ([Table 19](#)). In some individuals, options may be limited because of prior antiretroviral use,

toxicity or intolerance. In the clinically stable patient with detectable viremia for whom an optimal change in therapy is not possible, it may be prudent to delay changing therapy in anticipation of the availability of newer and more potent agents. It is recommended that the decision to change therapy and design a new regimen should be made with assistance from a clinician experienced in the treatment of HIV infected patients through consultation or referral.

ACUTE HIV INFECTION

It has been estimated that at least 50% and as many as 90% of patients acutely infected with HIV will experience at least some symptoms of the acute retroviral syndrome ([Table 22](#)) and can thus be identified as candidates for early therapy [200–203]. However, acute HIV infection is often not recognized in the primary care setting because of the similarity of the symptom complex with those of the “flu” or other common illnesses. Additionally, acute primary infection may occur without symptoms. **Health care providers** should maintain a high level of suspicion for HIV infection in all patients presenting with a compatible clinical syndrome ([Table 22](#)) and should obtain appropriate laboratory testing. **Evidence includes detectable HIV RNA in plasma using sensitive PCR or bDNA assays together with a negative or indeterminate HIV antibody test.** While measurement of plasma HIV RNA is the preferred method of diagnosis, a test for p24 antigen may be useful when RNA testing is not readily available. It should be noted, however, that a negative p24 antigen test does not rule out acute infection, and there may be low titer (<10,000 copies/mL), false-positive tests with HIV RNA levels. When suspicion for acute infection is high, such as in a patient with a report of recent risk behavior in association with symptoms and signs listed in [Table 22](#), a test for HIV RNA should be performed (BII). Patients diagnosed with HIV infection by HIV RNA testing should have confirmatory testing performed ([Table 2](#)).

Information regarding treatment of acute HIV infection from clinical trials is very limited. Preliminary data suggest that treatment of primary HIV infection with combination therapy has a beneficial effect on laboratory markers of disease progression [18, 19, 204–206]. **However, the potential disadvantages of initiating therapy include additional exposure to antiretroviral therapy without a known clinical benefit that may result in significant toxicities, development of antiretroviral drug resistance, and adverse effect on**

quality of life. Ongoing clinical trials are addressing the question of the long term clinical benefit of potent treatment regimens.

The theoretical rationale for early intervention is as follows:

- to decrease the severity of acute disease;
- to potentially alter the initial viral “set point,” which may ultimately affect the rate of disease progression;
- to possibly reduce the rate of viral mutation due to the suppression of viral replication;
- to preserve immune function;
- **to possibly reduce the risk of viral transmission.**

The **potential** risks of therapy for acute HIV infection include

- adverse effects on quality of life resulting from drug toxicities and dosing constraints;
- the potential, if therapy fails to effectively suppress viral replication, for the development of drug resistance which may limit future treatment options;
- the potential need for continuing therapy indefinitely.

These considerations are similar to those for initiating therapy in the asymptomatic patient and were discussed in greater detail in the section **“[Considerations for Initiating Therapy in the Patient with Asymptomatic HIV-Infection](#),”** p. 7.

The health care provider and the patient should be fully aware that therapy for primary HIV infection is based on theoretical considerations, and the potential benefits should be weighed against the potential risks. Many authorities endorse treatment of acute HIV infection based on the theoretical rationale and limited but supportive clinical trial data.

Apart from patients with acute primary HIV infection, many experts would also consider therapy for patients in whom seroconversion has been documented to have occurred within the previous six months (CIII). Although the initial burst of viremia in infected adults has usually resolved by two months, treatment during the 2–6 month period after infection is based on the likelihood that virus replication in lymphoid tissue is still not maximally contained by the immune system during this time [207]. Decisions regarding therapy for patients who test antibody positive and who believe the infection is recent but for whom the time of infection cannot be documented should be made using the algorithm mentioned previously; see **“[Considerations for Patients with Established HIV Infection](#),”** p. 6 (CIII). Except in the setting of post-exposure prophylaxis with antiretroviral agents [208], no patient

should be treated for HIV infection until the infection is documented. In this regard, all patients presenting without a formal medical record of a positive HIV test, such as those who have tested positive by available home testing kits, should undergo ELISA and an established confirmatory test such as the Western Blot (AI) to document HIV infection.

Treatment Regimen for Primary HIV Infection

Once the clinician and patient have made the decision to use antiretroviral therapy for primary HIV infection, treatment should be implemented with the goal of suppressing plasma HIV RNA levels to below detectable levels (AIII). There are insufficient data to make firm conclusions regarding specific drug recommendations; potential combinations of agents available are much the same as those used in established infection, listed in [Table 12](#). It is recognized that these aggressive regimens may be associated with several disadvantages, including drug toxicity, large pill burden, cost of drugs, and the possibility of developing drug resistance that may limit future options; the latter is likely if virus replication is not adequately suppressed or if the patient has been infected with a viral strain that is already resistant to one or more agents. The patient should be carefully counseled regarding these potential limitations and individual decisions made only after weighing the risks and sequelae of therapy against the theoretical benefit of treatment (see above). Since 1) the ultimate goal of therapy is suppression of viral replication to below the level of detection, and 2) the benefits of therapy are based primarily on theoretical considerations and 3) long term clinical outcome benefit has not been documented, any regimen that is not expected to maximally suppress viral replication is not considered appropriate for treating the acutely HIV-infected individual (EIII). Additional clinical studies are needed to delineate further the role of antiretroviral therapy in the primary infection period.

Patient Follow-up

Testing for plasma HIV RNA levels and CD4⁺ T cell count and toxicity monitoring should be performed as described above in [“Use of Testing for Plasma HIV RNA Levels and CD4+ T Cell Count in Guiding Decisions for Therapy,”](#) p. 4 (i.e., on initiation of therapy, after 4 weeks, and every 3–4 months thereafter) (AII). Some experts feel that testing for

plasma HIV RNA levels at 4 weeks is not helpful in evaluating the effect of therapy for acute infection as viral loads may be decreasing from peak viremia levels even in the absence of therapy.

Duration of Therapy for Primary HIV Infection

Once therapy is initiated many experts would continue to treat the patient with antiretroviral agents indefinitely because viremia has been documented to reappear or increase after discontinuation of therapy (CII). The optimal duration and composition of therapy are unknown and ongoing clinical trials are expected to provide data relevant to these issues. The difficulties inherent in determining the optimal duration and composition of therapy initiated for acute infection should be considered when first counseling the patient regarding therapy.

CONSIDERATIONS FOR ANTIRETROVIRAL THERAPY IN THE HIV-INFECTED ADOLESCENT

HIV-infected adolescents who were infected sexually or via injection drug use during adolescence appear to follow a clinical course that is more similar to HIV disease in adults than in children. In contrast, adolescents who were infected perinatally or via blood products as young children have a unique clinical course that may differ from other adolescents and long-term surviving adults. Currently, most HIV-infected adolescents were infected sexually during the adolescent period and are in a relatively early stage of infection.

Puberty is a time of somatic growth and hormonally-mediated changes, with females developing more body fat and males more muscle mass. Although theoretically these physiologic changes could affect drug pharmacology, particularly in the case of drugs with a narrow therapeutic index that are used in combination with protein-bound medicines or hepatic enzyme inducers or inhibitors, no clinically significant impact of puberty has been noted to date with the use of NRTIs. Clinical experience with PIs and NNRTIs has been limited. Thus, it is currently recommended that medications used to treat HIV and opportunistic infections in adolescents should be dosed based on Tanner staging of puberty and not specific age. Adolescents in early puberty (Tanner I–II) should be

dosed under pediatric guidelines, while those in late puberty (Tanner V) should be dosed by adult guidelines. Youth who are in the midst of their growth spurt (Tanner III females and Tanner IV males) should be closely monitored for medication efficacy and toxicity when choosing adult or pediatric dosing guidelines.

CONSIDERATIONS FOR ANTIRETROVIRAL THERAPY IN THE HIV-INFECTED PREGNANT WOMAN

Antiretroviral treatment recommendations for HIV-infected pregnant women are based on the belief that therapies of known benefit to women should not be withheld during pregnancy unless the risk of adverse effects outweighs the expected benefit to the woman. Combination antiretroviral therapy is the recommended standard treatment for HIV-infected non-pregnant adults. Additionally, a three-part regimen of ZDV, given orally starting at 14 weeks gestation and continued throughout pregnancy, intravenously during labor and to the newborn for the first six weeks of life, reduced the risk of perinatal transmission by 66% in a randomized, double-blind clinical trial (PACTG protocol 076) [20] and is recommended for all pregnant women [209]. Pregnancy should not preclude the use of optimal therapeutic regimens. However, recommendations regarding the choice of antiretroviral drugs for treatment of infected women are subject to unique considerations including:

- a. potential changes in dosing requirement resulting from physiologic changes associated with pregnancy,
- b. potential effects of antiretroviral drugs on the pregnant woman,
- c. effect on the risk of perinatal HIV transmission,
- d. the potential short- and long-term effects of the antiretroviral drug on the fetus and newborn, which may not be known for many antiretroviral drugs [209] (See [Public Health Service Task Force Recommendations for the Use of Antiretroviral Drugs in Pregnant HIV-1 Infected Women for Maternal Health and Interventions to Reduce Perinatal HIV-1 Transmission in the United States](#)).

As discussed further below, the decision to use any antiretroviral drug during pregnancy should be made by the woman following discussion with her health care provider regarding the known and unknown

benefits and risks to her and her fetus. Long-term follow-up is recommended for all infants born to women who have received antiretroviral drugs during pregnancy.

Women who are in the first trimester of pregnancy and who are not receiving antiretroviral therapy may wish to consider delaying initiation of therapy until after 10 to 12 weeks gestation, since this is the period of organogenesis when the embryo is most susceptible to potential teratogenic effects of drugs; the risks of antiretroviral therapy to the fetus during that period are unknown. However, this decision should be carefully considered and discussed between the health care provider and the patient and should include an assessment of the woman's health status and the potential benefits and risks of delaying initiation of therapy for several weeks. If clinical, virologic or immunologic parameters were such that therapy would be recommended for nonpregnant individuals, many of the Panel members would recommend initiating therapy regardless of gestational age. Nausea and vomiting in early pregnancy affecting the ability to adequately take and absorb oral medications may be a factor in the decision regarding treatment during the first trimester.

Standard combination antiretroviral therapy is recommended as initial therapy for HIV-infected pregnant women whose clinical, immunologic or virologic status would suggest the need for treatment if non-pregnant. When initiation of antiretroviral therapy would be considered optional based on current guidelines for treatment of non-pregnant individuals but HIV-1 RNA levels are $\geq 1,000$ copies/mL, infected pregnant women should be counseled regarding the potential benefits of standard combination therapy and should be offered such therapy including the three-part ZDV chemoprophylaxis regimen ([Table 24](#)). Although such women are at low risk for clinical disease progression if combination therapy is delayed, antiretroviral therapy that successfully reduces HIV-1 RNA levels to below 1,000 copies/mL substantially lowers the risk of perinatal transmission [210–212] and limits the need to consider elective cesarean delivery as an intervention to reduce transmission risk [209] (See [Public Health Service Task Force Recommendations for the Use of Antiretroviral Drugs in Pregnant HIV-1 Infected Women for Maternal Health and Interventions to Reduce Perinatal HIV-1 Transmission in the United States](#)).

Use of antiretroviral prophylaxis has been shown to provide benefit in preventing perinatal transmission

even for infected pregnant women with HIV-1 RNA levels <1,000 copies/mL. In a meta-analysis of factors associated with perinatal transmission among women who had infected infants despite having HIV-1 RNA <1,000 copies/mL at or near delivery, transmission was only 1.0% among women receiving ZDV prophylaxis compared to 9.8% among those receiving no antiretroviral treatment [210]. The time-limited use of ZDV alone during pregnancy for chemoprophylaxis of perinatal transmission is controversial. The potential benefits of standard combination antiretroviral regimens for treatment of HIV infection should be discussed with and offered to all pregnant HIV-infected women regardless of viral load, and is recommended for all pregnant women with HIV-1 RNA levels \geq 1,000 copies/mL. However, some women may wish to restrict exposure of their fetus to antiretroviral drugs during pregnancy but still wish to reduce the risk of transmitting HIV to their infant. Additionally, for women with HIV-1 RNA levels <1,000 copies/mL, time-limited use of ZDV during the second and third trimesters of pregnancy is less likely to induce the development of resistance due to the limited viral replication existing in the patient and the time-limited exposure to the antiretroviral drug. For example, the development of ZDV resistance was unusual among the healthy population of women who participated in PACTG 076 [21]. The use of ZDV chemoprophylaxis alone during pregnancy might be an appropriate option for these women.

When combination therapy is given principally to reduce perinatal transmission and would have been considered optional for treatment if non-pregnant, consideration may be given to discontinuing therapy postnatally, with the decision to reinstitute treatment based on standard criteria for non-pregnant individuals. If drugs are discontinued postnatally, all drugs should be stopped simultaneously. Discussion regarding the decision to continue or stop combination therapy postpartum should occur prior to initiation of therapy during pregnancy.

Some women already receiving antiretroviral therapy may recognize their pregnancy early enough in gestation that concern for potential teratogenicity may lead them to consider temporarily stopping antiretroviral therapy until after the first trimester. There are insufficient data to support or refute teratogenic risk of antiretroviral drugs in humans when administered during the first 10–12 weeks of gestation. However, treatment with EFV should be avoided during the first trimester because significant teratogenic effects in rhesus macaques were seen at

drug exposures similar to those representing human exposure. Hydroxyurea is a potent teratogen in a variety of animal species and should also be avoided during the first trimester.

There is concern that temporary discontinuation of antiretroviral therapy could result in a rebound in viral levels that theoretically could be associated with increased risk of early *in utero* HIV transmission or could potentiate disease progression in the woman [213]. Although the effects of all antiretroviral drugs on the developing fetus during the first trimester are uncertain, many experts recommend continuation of a maximally suppressive regimen even during the first trimester. If antiretroviral therapy is discontinued during the first trimester for any reason, all agents should be stopped simultaneously to avoid development of resistance. Once the drugs are reinstated, they should be introduced simultaneously for the same reason.

There are currently limited data available on the pharmacokinetics and safety of antiretroviral agents during pregnancy for drugs other than ZDV (see “[Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy](#),” p. 78). In the absence of data, drug choices need to be individualized based on discussion with the patient and available data from preclinical and clinical testing of the individual drugs. The FDA pregnancy classification for all currently approved antiretroviral agents and selected other information relevant to the use of antiretroviral drugs in pregnancy is shown in [Table 23](#). It is important to recognize that the predictive value of *in vitro* and animal screening tests for adverse effects in humans is unknown. Many drugs commonly used to treat HIV infection or its consequences may have positive findings on one or more of these screening tests. For example, acyclovir is positive on some *in vitro* assays for chromosomal breakage and carcinogenicity and is associated with some fetal abnormalities in rats; however, data on human experience from the Acyclovir in Pregnancy Registry indicate no increased risk of birth defects to date in infants with *in utero* exposure to acyclovir [214].

When combination antiretroviral therapy is given during pregnancy, ZDV should be included as a component of antenatal therapy whenever possible. There may be circumstances where this is not feasible, such as the occurrence of significant ZDV-related toxicity. In addition, women receiving an antiretroviral regimen that does not contain ZDV but who have HIV-1 RNA levels that are consistently very low or

undetectable have a very low risk of perinatal transmission and there may be concerns that the addition of ZDV to the current regimen could compromise adherence to the regimen. Regardless of the antepartum antiretroviral regimen, intravenous intrapartum ZDV and the standard six week course of ZDV for the infant is recommended. If the woman has not received ZDV as a component of her antenatal therapeutic antiretroviral regimen, intravenous ZDV should still be administered to the pregnant woman during the intrapartum period when feasible. Additionally, in women receiving combination antiretroviral treatment, the maternal antenatal antiretroviral treatment regimen should be continued on schedule as much as possible during labor to provide maximal virologic effect and to minimize the chance of development of drug resistance. Because ZDV and d4T should not be administered together due to potential pharmacologic antagonism, options for women receiving oral d4T as part of their antenatal therapy include continuation of oral d4T during labor without intravenous ZDV, or withholding the oral d4T during the period of intravenous administration during labor.

Toxicity related to mitochondrial dysfunction has been reported in infected patients receiving long-term treatment with nucleoside analogues, and may be of particular concern for pregnant women. Symptomatic lactic acidosis and hepatic steatosis may have a female preponderance [125]. Additionally, these syndromes have similarities to the rare but life-threatening syndromes of acute fatty liver of pregnancy and hemolysis, elevated liver enzymes and low platelets (the HELLP syndrome) that occur during the third trimester of pregnancy. Some data suggest that a disorder of mitochondrial fatty acid oxidation in the mother or her fetus during late pregnancy may play a role in the etiology of acute fatty liver of pregnancy and HELLP syndrome [215, 216], and possibly contribute to susceptibility to antiretroviral-associated mitochondrial toxicity.

It is unclear if pregnancy augments the incidence of the lactic acidosis/hepatic steatosis syndrome reported in non-pregnant individuals receiving nucleoside analogue treatment. Bristol-Myers Squibb has reported three maternal deaths due to lactic acidosis, two with and one without accompanying pancreatitis, in women who were either pregnant or postpartum and whose antepartum therapy during pregnancy included d4T and ddI in combination with other antiretroviral agents (either a PI or NVP) [217]. All cases were in women who were receiving treatment with these agents at the time of conception and continued for the duration of

pregnancy; all presented late in gestation with symptomatic disease that progressed to death in the immediate postpartum period. Two cases were also associated with fetal demise. Non-fatal cases of lactic acidosis in pregnant women have also been reported.

Because pregnancy itself can mimic some of the early symptoms of the lactic acidosis/hepatic steatosis syndrome or be associated with other significant disorders of liver metabolism, these cases emphasize the need for clinicians caring for HIV-infected pregnant women receiving nucleoside analogue drugs to be alert for early diagnosis of this syndrome. Pregnant women receiving nucleoside analogue drugs should have hepatic enzymes and electrolytes assessed more frequently during the last trimester of pregnancy and any new symptoms should be evaluated thoroughly. Additionally, because of the reports of several cases of maternal mortality secondary to lactic acidosis with prolonged use of the combination of d4T and ddI by HIV-infected pregnant women, clinicians should prescribe this antiretroviral combination during pregnancy with caution and generally only when other nucleoside analogue drug combinations have failed or caused unacceptable toxicity or side effects [217].

The antenatal ZDV dosing regimen used in the perinatal transmission prophylaxis trial PACTG 076 was ZDV 100 mg administered five times daily, and was selected based on the standard ZDV dosage for adults at the time the study was designed in 1989 (Table 24). However, data indicate that administration of ZDV three times daily will maintain intracellular ZDV triphosphate at levels comparable with those observed with more frequent dosing [218, 219]. Comparable clinical response also has been observed in clinical trials among persons receiving ZDV twice daily [220–222]. Thus, the current standard ZDV dosing regimen for adults is 200 mg three times daily, or 300 mg twice daily. A less frequent dosing regimen would be expected to enhance maternal adherence to the ZDV perinatal prophylaxis regimen, and therefore is an acceptable alternative antenatal dosing regimen for ZDV prophylaxis.

In a short-course antenatal/intrapartum ZDV perinatal transmission prophylaxis trial in Thailand, administration of ZDV 300 mg twice daily for 4 weeks antenatally and 300 mg every 3 hours orally during labor was shown to reduce perinatal transmission by approximately 50% compared to placebo [223]. The lower efficacy of the short-course 2-part ZDV prophylaxis regimen studied in Thailand compared to the 3-part ZDV prophylaxis regimen used in PACTG 076 and recommended for use in the U.S. could result

from the shorter antenatal duration of ZDV, oral rather than intravenous administration during labor, lack of treatment for the infant, or a combination of these factors. In the United States, identification of HIV-infected pregnant women before or as early as possible during the course of pregnancy and use of the full 3-part PACTG 076 ZDV regimen is recommended for prevention of perinatal HIV transmission.

Monitoring and use of HIV-1 RNA for therapeutic decision-making during pregnancy should be performed as recommended for non-pregnant individuals. Data from untreated as well as ZDV-treated infected pregnant women indicate that HIV-1 RNA levels correlate with risk of transmission [20, 211, 212]. However, although the risk of perinatal transmission in women with HIV-1 RNA below the level of assay quantitation appears to be very low, transmission from mother to infant has been reported in women with all levels of maternal HIV-1 RNA. Additionally, antiretroviral prophylaxis is effective in reducing transmission even among women with low RNA levels [20, 210]. The mechanism by which antiretroviral prophylaxis reduces transmission is likely multifactorial. Reduction in maternal antenatal viral load is likely an important component of prophylaxis. However, in addition, pre- and post-exposure prophylaxis of the infant is provided by passage of antiretroviral drugs across the placenta, resulting in inhibitory drug levels in the fetus during and immediately following the birth process [224]; the extent of transplacental passage varies between antiretroviral drugs (Table 23). Additionally, while there is general correlation between plasma and genital tract viral load, discordance has also been reported [225–227]; in addition, differential evolution of viral sequence diversity occurs between the peripheral blood and genital tract [227, 228]. Studies are needed to define the relationship between viral load suppression by antiretroviral therapy in plasma and levels of HIV in the genital tract, and the relationship between these compartment-specific effects and the risk of perinatal HIV transmission. The full ZDV chemoprophylaxis regimen, including intravenous ZDV during delivery and the administration of ZDV to the infant for the first six weeks of life, in combination with other antiretrovirals or alone in selected cases, should be discussed with and offered to all infected pregnant women regardless of their HIV-1 RNA level.

Health care providers who are treating HIV-infected pregnant women are strongly encouraged to report cases of prenatal exposure to antiretroviral drugs (either administered alone or in combinations) to the Antiretroviral Pregnancy Registry. The registry collects

observational, nonexperimental data regarding anti-retroviral exposure during pregnancy for the purpose of assessing potential teratogenicity. Registry data will be used to supplement animal toxicology studies and assist clinicians in weighing the potential risks and benefits of treatment for individual patients. The registry is a collaborative project with an advisory committee of obstetric and pediatric practitioners, staff from CDC and NIH, and staff from pharmaceutical manufacturers. The registry allows the anonymity of patients, and birth outcome follow-up is obtained by registry staff from the reporting clinician.

Referrals should be directed to
Antiretroviral Pregnancy Registry
Research Park
1011 Ashes Drive
Wilmington, NC 28405
Telephone 1–800–258–4263
Fax 1–800–800–1052
Internet access <http://www.apregistry.com/>

PREVENTION COUNSELING IN THE PATIENT WITH HIV INFECTION

Ongoing prevention counseling is an essential component of the management of persons with HIV infection [229]. Each patient encounter provides an opportunity to reinforce HIV prevention messages. Each encounter should include assessment and documentation of

1. the patient's knowledge and understanding of the means of HIV transmission; and
2. the patient's HIV transmission behaviors since the last encounter with a member of the health care team.

This should be followed by a discussion of strategies to prevent transmission that may be useful to the individual patient. The physician, nurse, or other health care team member should routinely provide this counseling. Partner notification is an important component of HIV detection and prevention and should be pursued by the provider or by referral services.

Although the core elements of HIV prevention messages have been unchanged since the introduction of HAART, important observations regarding the biology of HIV transmission, the impact of HAART upon transmission, and individual risk behaviors have

been noted. For example, sustained low plasma viremia that results from successful HIV therapy substantially reduces the likelihood of HIV transmission. In one study, for each one-log reduction in plasma viral load, the likelihood of transmission between discordant couples was reduced 2.5-fold [230]. Similarly, mother to child HIV transmission was observed to decline in a linear fashion with each log reduction in maternal delivery viral load [212]. While in general this relationship is linear, important exceptions should be noted. For example, mother to child transmission have been reported even in women with very low or “undetectable” viral loads [210].

Similarly, the relationship between viral load in the plasma, and the levels in the genital fluid of women and the seminal fluid of men is complex. In general, studies have shown a rough correlation between plasma HIV levels and genital HIV levels, but important exceptions have been observed [231]. Viral evolution may occur in the genital compartment that is distinct from the viral evolution in the plasma, and transmissions have been documented in the presence of an undetectable plasma viral load [20, 210, 232]. Thus, while durably effective HAART substantially reduces the likelihood of HIV transmission, the degree of protection is incomplete.

Many biologic factors other than plasma viral load have also been shown to influence sexual transmission of HIV. These include ulcerative and nonulcerative sexually transmitted infections [233], vaginitis (including bacterial vaginosis and candida vaginitis) [234], genital irritation associated with frequent use of products containing nonoxynol-9 (N-9), menstruation, lack of circumcision in men [235], oral contraceptive use [236], estrogen deficiency [237], progesterone excess [238], and deficiencies of vitamin A [239] and selenium [240].

Behavioral changes in HIV-infected individuals that impact prevention have been observed during the HAART era. Unfortunately, there is evidence that awareness of the potential benefits of highly active antiretroviral therapy is leading some individuals to relapse into high-risk activities. For example, reports from urban gay communities in the U.S. suggest rising HIV seroprevalence rates, as well as rising rates of unsafe sexual practices, corroborated by the rising rates of other sexually transmitted infections. Recently an association between knowledge of the benefits of HAART among gay men and relapse to high-risk activity was observed [241, 242].

Women may have unprotected sex because they wish to become pregnant. For women of childbearing potential, the desire for pregnancy should be assessed at each encounter; women wishing to pursue pregnancy should be referred for preconception counseling to reduce risks of perinatal transmission and transmission to uninfected sexual partners. In women of childbearing age who wish to avoid pregnancy, condoms should be encouraged in addition to other forms of contraception for prevention of HIV and STD transmission (dual method use) or used as a single method for prevention of pregnancy as well (dual protection). In a randomized placebo-controlled clinical trial conducted among commercial sex workers with high rates of sexual activity, use of N-9 did not protect against HIV infection, resulted in more vaginal lesions, and may have caused more transmission. Although these adverse effects might not occur with less frequent use, given current evidence, spermicides containing N-9 should not be recommended as an effective means of HIV prevention.

Good adherence to antiretroviral regimens has been directly associated with a lower risk of morbidity and mortality, and indirectly associated with a reduction in the risk of HIV transmission due to its association with lower viral loads [243]. Poor adherence to HIV medication recently has been shown to be a predictor of poor adherence to HIV prevention strategies [244]. More intensive adherence and prevention counseling may be appropriate for individuals who demonstrate repeated deficiencies in either area.

In summary, despite the strong association between a reduced risk of HIV transmission and sustained low viral load, the message of HIV prevention for patients should remain simple: Once infected, an HIV-infected individual may transmit the virus at any time, and there is no substitute for latex or polyurethane male or female condoms, other safer sexual behaviors (e.g., partner reduction, abstinence), and cessation of any sharing of drug paraphernalia. Prevention counseling for patients known to have HIV infection, including easy access to condoms and other means of prevention, remains a critical component of HIV primary care. Clinicians may wish to directly address with their patients the risks of using viral load outcomes as a factor in considering high-risk behavior. HIV-infected persons who use injection drugs should be advised to enroll in drug rehabilitation programs. If this advice is not followed, or if these services are not available, the patient should receive counseling on risks associated with sharing needles and paraphernalia.

Finally, the most successful and effective prevention messages are those tailored to the individual. These messages are culturally appropriate, practical, and relevant to the individual's knowledge, beliefs, and behaviors [229]. The message, the manner of delivery, and the cultural context vary greatly depending upon the patient. For more information about these strategies, as well as recommendations on prevention, please see "HIV Transmission and Prevention in Specific Populations" on the HIV InSite web page (<http://hivinsite.ucsf.edu/InSite.jsp?page=kb-07>).

CONCLUSION

The panel has attempted to use the advances in our understanding of the pathogenesis of HIV in the infected person to translate scientific principles and data obtained from clinical experience into recommendations that can be used by the clinician and patient to make therapeutic decisions. The recommendations are offered in the context of an ongoing dialogue between the patient and the clinician after having defined specific therapeutic goals with an acknowledgment of uncertainties. It is necessary for the patient to be entered into a continuum of medical care and services, including social, psychosocial, and nutritional services, with the availability of expert referral and consultation. In order to achieve the maximal flexibility in tailoring therapy to each patient over the duration of his or her infection, it is imperative that drug formularies allow for all FDA-approved NRTI, NNRTI, and PI as treatment options. The Panel strongly urges industry and the public/private sectors to conduct further studies to allow refinement of these guidelines. Specifically, studies are needed to optimize recommendations for first line therapy; to define second line therapy; and to more clearly delineate the reason(s) for treatment failure. The Panel remains committed to revising their recommendations as such new data become available.

- Information included in these guidelines may not represent FDA approval or approved labeling for the particular products or indications in question. Specifically, the terms "safe" and "effective" may not be synonymous with the FDA-defined legal standards for product approval.


Table 1. Rating Scheme for Clinical Practice

Table 2. Indications for Plasma HIV RNA Testing*

Clinical Indication	Information	Use
Syndrome consistent with acute HIV infection	Establishes diagnosis when HIV antibody test is negative or indeterminate	Diagnosis [†]
Initial evaluation of newly diagnosed HIV infection	Baseline viral load “set point”	Decision to start or defer therapy
Every 3-4 months in patients not on therapy	Changes in viral load	Decision to start therapy
2-8 weeks after initiation of antiretroviral therapy	Initial assessment of drug efficacy	Decision to continue or change therapy
3-4 months after start of therapy	Maximal effect of therapy	Decision to continue or change therapy
Every 3-4 months in patients on therapy	Durability of antiretroviral effect	Decision to continue or change therapy
Clinical event or significant decline in CD4 ⁺ T cells	Association with changing or stable viral load	Decision to continue, initiate, or change therapy

* Acute illness (e.g., bacterial pneumonia, tuberculosis, HSV, PCP, etc.) and immunizations can cause increase in plasma HIV RNA for 2-4 weeks; viral load testing should not be performed during this time. Plasma HIV RNA results should usually be verified with a repeat determination before starting or making changes in therapy.

† Diagnosis of HIV infection made by HIV RNA testing should be confirmed by standard methods such as Western blot serology performed 2-4 months after the initial indeterminate or negative

Table 3. Recommendations for the Use of Drug Resistance Assays

Clinical Setting/Recommendation	Rationale
<p>Recommended</p> <p>Virologic failure during HAART (see page 21)</p> <p>Suboptimal suppression of viral load after initiation of antiretroviral therapy (see page 13)</p>	<p>Determine the role of resistance in drug failure and maximize the number of active drugs in the new regimen if indicated.</p> <p>Determine the role of resistance and maximize the number of active drugs in the new regimen if indicated.</p>
<p>Consider</p> <p>Acute HIV infection</p>	<p>Determine if drug resistant virus was transmitted and change regimen accordingly.</p>
<p>Not generally recommended</p> <p>Chronic HIV infection prior to initiation of therapy</p> <p>After discontinuation of drugs</p> <p>Plasma viral load <1000 HIV RNA copies/mL</p>	<p>Uncertain prevalence of resistant virus. Current assays may not detect minor drug resistant species.</p> <p>Drug resistance mutations may become minor species in the absence of selective drug pressure. Current assays may not detect minor drug resistant species.</p> <p>Resistance assays cannot be reliably performed because of low copy number of HIV RNA.</p>

Table 4. Risks and Benefits of Delayed Initiation of Therapy and of Early Therapy in the Asymptomatic HIV-Infected Patient

Risks and benefits of delayed therapy*
<p>Benefits of delayed therapy</p> <ul style="list-style-type: none"> • Avoid negative effects on quality of life (i.e., inconvenience) • Avoid drug-related adverse events • Delay in development of drug resistance • Preserve maximum number of available and future drug options when HIV disease risk is highest <p>Risks of delayed therapy</p> <ul style="list-style-type: none"> • Possible risk of irreversible immune system depletion • Possible greater difficulty in suppressing viral replication • Possible increased risk of HIV transmission
Risks and benefits of early therapy*
<p>Benefits of early therapy</p> <ul style="list-style-type: none"> • Control of viral replication easier to achieve and maintain • Delay or prevention of immune system compromise • Lower risk of resistance with complete viral suppression • Possible decreased risk of HIV transmission[†] <p>Risks of early therapy</p> <ul style="list-style-type: none"> • Drug-related reduction in quality of life • Greater cumulative drug-related adverse events • Earlier development of drug resistance, if viral suppression is suboptimal • Limitation of future antiretroviral treatment options

* See **Table 6** for consensus recommendations regarding when to initiate therapy.

[†] The risk of viral transmission still exists; antiretroviral therapy cannot substitute for primary HIV prevention measures (e.g., use of condoms and safer sex practices).

Table 5. Risk of Progression to AIDS Defining Illness in a Cohort of Homosexual Men Predicted by Baseline CD4⁺ T Cell Count and Viral Load*

CD4 < 200		% AIDS			
Plasma Viral Load (copies/ml) †		(AIDS-defining complication) ‡			
bDNA	RT-PCR	n	3 years	6 years	9 years
≤ 500	≤ 1,500	0 §	–	–	–
501 – 3,000	1,501 – 7,000	3 §	–	–	–
3,001 – 10,000	7,001 – 20,000	7	14.3	28.6	64.3
10,001 – 30,000	20,001 – 55,000	20	50.0	75.0	90.0
> 30,000	> 55,000	70	85.5	97.9	100.0
CD4 201 – 350 [∞]		% AIDS			
Plasma Viral Load (copies/ml)		(AIDS-defining complication)			
bDNA	RT-PCR	n	3 years	6 years	9 years
≤ 500	≤ 1,500	3 §	–	–	–
501 – 3,000	1,501 – 7,000	27	0	20.0	32.2
3,001 – 10,000	7,001 – 20,000	44	6.9	44.4	66.2
10,001 – 30,000	20,001 – 55,000	53	36.4	72.2	84.5
> 30,000	> 55,000	104	64.4	89.3	92.9
CD4 > 350		% AIDS			
Plasma Viral Load (copies/ml)		(AIDS-defining complication)			
bDNA	RT-PCR	n	3 years	6 years	9 years
≤ 500	≤ 1,500	119	1.7	5.5	12.7
501 – 3,000	1,501 – 7,000	227	2.2	16.4	30.0
3,001 – 10,000	7,001 – 20,000	342	6.8	30.1	53.5
10,001 – 30,000	20,001 – 55,000	323	14.8	51.2	73.5
> 30,000	> 55,000	262	39.6	71.8	85.0

* Data from the Multi-Center AIDS Cohort Study (MACS) adapted from reference 3 by Alvaro Muñoz, Ph.D. (personal communication.)

† MACS numbers reflect plasma HIV RNA values obtained by version 2.0 bDNA testing. RT-PCR values are consistently 2–2.5 fold higher than first generation bDNA values, as indicated. It should be noted that the current generation bDNA assay (3.0) gives similar HIV-1 RNA values as RT-PCR except at the lower end of the linear range (<1,500 copies/ml).

‡ In this study AIDS was defined according to the 1987 CDC definition and does not include asymptomatic individuals with CD4⁺ T cells <200 mm³.

∞ A recent evaluation of data from the MACS cohort by of 231 individuals with CD4⁺ T cell counts >200 and <350 cells/mm³ demonstrated that of 40 (17%) individuals with plasma HIV RNA <10,000 copies/ml, none progressed to AIDS by 3 years (Alvaro Munoz, personal communication). Of 28 individuals (29%) with plasma viremia of 10,000–20,000 copies/ml, 4% and 11% progressed to AIDS at 2 and 3 years, respectively. Plasma HIV RNA was calculated as RT-PCR values from measured bDNA values.

§ Too few subjects were in the category to provide a reliable estimate of AIDS risk.

Table 6. Indications for the Initiation of Antiretroviral Therapy in the Chronically HIV-1 Infected Patient

The optimal time to initiate therapy in asymptomatic individuals with >200 CD4⁺ T cells is not known. This table provides general guidance rather than absolute recommendations for an individual patient. All decisions to initiate therapy should be based on prognosis as determined by the CD4⁺ T cell count and level of plasma HIV RNA shown in [Table 5](#), the potential benefits and risks of therapy shown in [Table 4](#), and the willingness of the patient to accept therapy.

Clinical Category	CD4 ⁺ T Cell Count	Plasma HIV RNA	Recommendation
Symptomatic (AIDS, severe symptoms)	Any value	Any value	Treat
Asymptomatic, AIDS	CD4 ⁺ T cells $<200/\text{mm}^3$	Any value	Treat
Asymptomatic	CD4 ⁺ T cells $>200/\text{mm}^3$ but $<350/\text{mm}^3$	Any value	Treatment should generally be offered, though controversy exists.*
Asymptomatic	CD4 ⁺ T cells $>350/\text{mm}^3$	$>55,000$ (by bDNA or RT-PCR) ^φ	Some experts would recommend initiating therapy, recognizing that the 3-year risk of developing AIDS in untreated patients is $>30\%$ and some would defer therapy and monitor CD4 ⁺ T cell counts more frequently.
Asymptomatic	CD4 ⁺ T cells $>350/\text{mm}^3$	$<55,000$ (by bDNA or RT-PCR) ^φ	Many experts would defer therapy and observe, recognizing that the 3-year risk of developing AIDS in untreated patients is $<15\%$.

* Clinical benefit has been demonstrated in controlled trials only for patients with CD4⁺ T cells $<200/\text{mm}^3$. However, most experts would offer therapy at a CD4⁺ T cell threshold $<350/\text{mm}^3$. A recent evaluation of data from the MACS cohort of 231 individuals with CD4⁺ T cell counts >200 and <350 cells/ mm^3 demonstrated that of 40 (17%) individuals with plasma HIV RNA $<10,000$ copies/ml, none progressed to AIDS by 3 years (Alvaro Munoz, personal communication). Of 28 individuals (29%) with plasma viremia of 10,000–20,000 copies/mL, 4% and 11% progressed to AIDS at 2 and 3 years respectively. Plasma HIV RNA was calculated as RT-PCR values from measured bDNA values. For further information, see [“Considerations for Initiating Therapy in the Patient with Asymptomatic HIV Infection,”](#) p. 7.

φ Although there was a 2-2.5 fold difference between RT-PCR and the first bDNA assay (version 2.0), with the current bDNA assay (version 3.0), values obtained by bDNA and RT-PCR are similar except at the lower end of the linear range ($<1,500$ copies/mL).

Table 7. Strategies to Improve Adherence: Patient and Medication-Related

- Inform patient, anticipate, and treat side effects.
- Simplify food requirements.
- Avoid adverse drug interactions.
- If possible, reduce dose frequency and number of pills.
- Negotiate a treatment plan, which the patient understands and to which he/she commits.
- Take time, multiple encounters to educate and explain goals of therapy and need for adherence.
- Establish readiness to take medication *before* first prescription is written.
- Recruit family and friends to support the treatment plan.
- Develop concrete plan for specific regimen, relation to meals, daily schedule, side effects.
- Provide written schedule and pictures of medications, daily or weekly pill boxes, alarm clocks, pagers, other mechanical aids to adherence.
- Develop adherence support groups, or add adherence issues to regular agenda of support groups.
- Develop linkages with local community-based organizations around adherence with educational sessions and practical strategies.
- Consider “pill trials” with jelly beans.

Table 8. Strategies to Improve Adherence: Clinician and Health Team Related

<ul style="list-style-type: none"> • Establish trust.
<ul style="list-style-type: none"> • Serve as educator, source of information, ongoing support and monitoring.
<ul style="list-style-type: none"> • Provide access between visits for questions/problems via page number, including vacation/conference coverage.
<ul style="list-style-type: none"> • Monitor ongoing adherence; intensify management in periods of low adherence (i.e., more frequent visits, recruitment of family/friends, deployment of other team members, referral for mental health or chemical dependency services).
<ul style="list-style-type: none"> • Utilize health team for all patients, for difficult patients, for special needs (e.g., peer educators for adolescents or for injection drug users).
<ul style="list-style-type: none"> • Consider impact of new diagnoses on adherence (e.g., depression, liver disease, wasting, recurrent chemical dependency), and include adherence intervention in management.
<ul style="list-style-type: none"> • Utilize nurses, pharmacists, peer educators, volunteers, case managers, drug counselors, physician's assistants, nurse practitioners, research nurses to reinforce message of adherence.
<ul style="list-style-type: none"> • Provide training to support team related to antiretroviral therapy and adherence.
<ul style="list-style-type: none"> • Add adherence interventions to job descriptions of HIV support team members; add continuity-of-care role to improve patient access.

Table 9. Interventions Associated with Improved Adherence

- ❖ **Pharmacist-based adherence encounters/clinics**
- ❖ **Adherence encounters at each visit, often multi-disciplinary**
- ❖ **Reminders, alarms, pagers, timers on pillboxes**
- ❖ **Patient education aids**, including regimen pictures, calendars, stickers
- ❖ **Clinician education aids** (e.g., medication guides, pictures, calendars)

Table 10. Goals of HIV Therapy and Tools to Achieve Them

Goals of Therapy
<ul style="list-style-type: none">• Maximal and durable suppression of viral load• Restoration and/or preservation of immunologic function• Improvement of quality of life• Reduction of HIV-related morbidity and mortality
Tools to Achieve Goals of Therapy
<ul style="list-style-type: none">• Maximize adherence to the antiretroviral regimen• Rational sequencing of drugs• Preservation of future treatment options• Use of resistance testing in selected clinical settings

Table 11. Advantages and Disadvantages of Class-Sparing Regimens

Regimen	Possible Advantages	Possible Disadvantages	Drug Interaction Complications	Impact on Future Options
PI-based HAART regimen (NNRTI-sparing)	<ul style="list-style-type: none"> • Clinical, virologic, and immunologic efficacy well-documented • Continued benefits sometimes seen despite viral breakthrough • Resistance requires multiple mutations • Targets HIV at two steps of viral replication (RT and PI) 	<ul style="list-style-type: none"> • May be difficult to use and adhere to • Long-term side effects may include lipodystrophy*, hyperlipidemia, and insulin resistance 	<ul style="list-style-type: none"> • Mild to severe inhibition of cytochrome P450 pathway; ritonavir is most potent inhibitor, but this effect can be exploited to boost levels of other PIs 	<ul style="list-style-type: none"> • Preserves NNRTIs for use in treatment failure • Resistance primes for cross-resistance with other PIs
NNRTI-based HAART regimen (PI-sparing)	<ul style="list-style-type: none"> • Sparing of PI-related side effects • Generally easier to use and adhere to compared with PIs 	<ul style="list-style-type: none"> • Comparability to PI-containing regimens with regard to clinical endpoints unknown • Resistance conferred by a single or few mutations 	<ul style="list-style-type: none"> • Fewer drug-drug interactions compared with PIs 	<ul style="list-style-type: none"> • Preserves PIs for later use • Resistance usually leads to cross-resistance across entire NNRTI class
Triple NRTI regimen (NNRTI- and PI-sparing)	<ul style="list-style-type: none"> • Generally easier to use and adhere to compared with PIs • Sparing of PI and NNRTI side effects • Resistance to 1 NRTI does not confer cross-resistance to entire class 	<ul style="list-style-type: none"> • Comparability to PI-containing regimens with regard to clinical endpoints unknown • Long-term virologic efficacy with high baseline plasma viral load (i.e., >100,000 copies/ml) may be suboptimal 	<ul style="list-style-type: none"> • Generally manageable drug interaction problems 	<ul style="list-style-type: none"> • Preserves both PI and NNRTI classes for later use • Limited cross-resistance within the NRTI class

* Some side effects being attributed to protease inhibitor therapy, such as lipodystrophy, have not been proven to be strictly associated with the use of protease inhibitor-containing regimens. Lipodystrophy has also been described uncommonly in patients on NRTIs alone and in patients on no antiretroviral therapy.

Table 12. Recommended Antiretroviral Agents for Initial Treatment of Established HIV Infection

This table provides a guide to the use of available treatment regimens for individuals with no prior or limited experience on HIV therapy. In accordance with the established goals of HIV therapy, priority is given to regimens in which clinical trials data suggest the following: sustained suppression of HIV plasma RNA (particularly in patients with high baseline viral load) and sustained increase in CD4⁺ T cell count (in most cases over 48 weeks), and favorable clinical outcome (i.e., delayed progression to AIDS and death). Particular emphasis is given to regimens that have been compared directly with other regimens that perform sufficiently well with regard to these parameters to be included in the “Strongly Recommended” category. Additional consideration is given to the regimen’s pill burden, dosing frequency, food requirements, convenience, toxicity, and drug interaction profile compared with other regimens.

It is important to note that all antiretroviral agents, including those in the “Strongly Recommended” category, have potentially serious toxic and adverse events associated with their use. The reader is strongly encouraged to consult tables 13–19 while formulating an antiretroviral regimen.

Antiretroviral drug regimens are comprised of one choice each from columns A and B. Drugs are listed in alphabetical, not priority, order.

Strongly Recommended	Column A Efavirenz Indinavir Nelfinavir Ritonavir + Indinavir* [†] Ritonavir + Lopinavir* [‡] Ritonavir + Saquinavir* (SGC [§] or HGC [§])	Column B Didanosine + Lamivudine Stavudine + Didanosine [¶] Stavudine + Lamivudine Zidovudine + Didanosine Zidovudine + Lamivudine
Recommended as Alternatives	Column A Abacavir Amprenavir Delavirdine Nelfinavir + Saquinavir-SGC Nevirapine Ritonavir Saquinavir-SGC	Column B Zidovudine + Zalcitabine
No Recommendation: Insufficient Data[#]	Hydroxyurea in combination with antiretroviral drugs Ritonavir + Amprenavir* Ritonavir + Nelfinavir* Tenofovir ••	
Not Recommended: Should Not Be Offered	All monotherapies, whether from column A or B** Column A Saquinavir-HGC †† Column B Stavudine + Zidovudine Zalcitabine + Didanosine Zalcitabine + Lamivudine Zalcitabine + Stavudine	

* See page 14 for more information on optimizing protease inhibitor exposure with ritonavir.

† Based on expert opinion.

‡ Co-formulated as Kaletra.

§ Saquinavir-SGC, soft-gel capsule (Fortovase); Saquinavir-HGC, hard-gel capsule (Invirase).

¶ Pregnant women may be at increased risk for lactic acidosis and liver damage when treated with the combination of stavudine and didanosine. This combination should be used in pregnant women only when the potential benefit clearly outweighs the potential risk.

This category includes drugs or combinations for which information is too limited to allow a recommendation for or against use.

** Zidovudine monotherapy may be considered for prophylactic use in pregnant women with low viral load and high CD4⁺ T cell counts to prevent perinatal transmission, as discussed under “Considerations for Antiretroviral Therapy in the HIV-Infected Pregnant Woman.”

†† Use of Saquinavir-HGC (Invirase) is not recommended, except in combination with ritonavir.

•• Data from clinical trials are limited to use in salvage. Data from trials of Tenofovir as initial therapy may be available in the near future.

Table 13. Characteristics of Nucleoside Reverse Transcriptase Inhibitors (NRTIs)

Generic Name	Zidovudine (AZT, ZDV)	Didanosine (ddI)	Zalcitabine (ddC)	Tenofovir Disoproxil Fumarate Viread
Trade Name	Retrovir	Videx	HIVID	
Form	100 mg capsules 300 mg tablets 10 mg/mL IV solution 10 mg/mL oral solution	25, 50, 100, 150, 200* mg chewable/dispersible buffered tablets 100, 167, 250 mg buffered powder for oral solution 400 mg enteric coated capsules (EC)	0.375, 0.75 mg tablets	300 mg tablet
Dosing Recommendations	200 mg tid or 300 mg bid or with 3TC as Combivir [†] , 1 bid or with abacavir and 3TC as Trizivir [‡] , 1 bid	>60kg: 200 mg bid (buff. tablets), 250 mg bid (buff. powder) or 400 mg qd [§] (buff. tablets or EC capsules) <60kg: 125 mg bid (buff. tablets), 167 mg bid (buff. powder) or 250 mg qd [§] (buff. tablets or EC capsules)	0.75 mg tid	300 mg QD - for pts with creatinine clearance ≥ 60 mL/min Not recommended for patient with creatinine clearance <60 mL/min
Food Effect	Take without regard to meals	Levels ↓ 55% Take 1/2 hour before or 2 hours after meal	Take without regard to meals	Increased bioavailability when taken with food
Oral bioavailability	60%	30-40%	85%	25% in fasting state 39% with high fat meal
Serum half-life	1.1 hours	1.6 hours	1.2 hours	17 hours
Intracellular half-life	3 hours	25-40 hours	3 hours	10 to 50 hours
Elimination	Metabolized to AZT glucuronide (GAZT) Renal excretion of GAZT	Renal excretion 50%	Renal excretion 70%	Primarily renally excreted by glomerular filtration and active tubular secretion
Adverse Events	Bone marrow suppression: Anemia and/or neutropenia Subjective complaints: GI intolerance, headache, insomnia, asthenia Lactic acidosis with hepatic steatosis is a rare but potentially life-threatening toxicity with the use of NRTIs.	Pancreatitis [¶] Peripheral neuropathy Nausea Diarrhea Lactic acidosis with hepatic steatosis is a rare but potentially life-threatening toxicity with the use of NRTIs. [#]	Peripheral neuropathy Stomatitis Lactic acidosis with hepatic steatosis is a rare but potentially life- threatening toxicity with the use of NRTIs.	Asthenia Headache Diarrhea Nausea Vomiting Flatulence

* For once daily dosing only. Twice daily dosing is preferred; however, once daily dosing may be appropriate for patients who require a simplified dosing schedule.

† Each Combivir tablet contains 300 mg ZDV and 150 mg 3TC.

‡ Each Trizivir tablet contains 300 mg ZDV, 150 mg 3TC, and 300 mg abacavir.

§ Twice daily dosing is preferred; however, once daily dosing may be appropriate for patients who require a simplified dosing schedule.

¶ Cases of fatal and nonfatal pancreatitis have occurred in treatment-naïve and treatment-experienced patients during therapy with ddI or in combination with other drugs, particularly d4T or d4T + hydroxyurea.

Pregnant women may be at increased risk for lactic acidosis and liver damage when treated with the combination of stavudine and didanosine. This combination should be used in pregnant women only when the potential benefit clearly outweighs the potential risk.

Table 13. Characteristics of Nucleoside Reverse Transcriptase Inhibitors (NRTIs)

Generic Name Trade Name	Stavudine (d4T) Zerit	Lamivudine (3TC) Epivir	Abacavir (ABC) Ziagen
Form	15, 20, 30, 40 mg capsules 1mg/mL for oral solution	150 mg tablets 10 mg/mL oral solution	300 mg tablets 20 mg/mL oral solution
Dosing Recommendations	>60kg: 40 mg bid <60kg: 30 mg bid	150 mg bid <50kg: 2 mg/kg bid or with ZDV as Combivir, 1 bid, or with ZDV and abacavir as Trizivir‡, 1 bid	300 mg bid or with ZDV and 3TC as Trizivir‡, 1 bid
Food Effect	Take without regard to meals	Take without regard to meals	Take without regard to meals Alcohol ↑ ABC levels 41%; no effect on alcohol
Oral bioavailability	86%	86%	83%
Serum half-life	1.0 hour	3-6 hours	1.5 hours
Intracellular half-life	3.5 hours	12 hours	3.3 hours
Elimination	Renal excretion 50%	Renal excretion unchanged	Metabolized by alcohol dehydrogenase and glucuronyl transferase. Renal excretion of metabolites 82%
Adverse Events	Pancreatitis [¶] Peripheral neuropathy Lactic acidosis with hepatic steatosis is a rare but potentially life-threatening toxicity with the use of NRTIs. [#]	(Minimal toxicity) Lactic acidosis with hepatic steatosis is a rare but potentially life-threatening toxicity with the use of NRTIs.	Hypersensitivity reaction (can be fatal) ^{**} ; fever, rash, nausea, vomiting, malaise or fatigue, and loss of appetite. Respiratory symptoms may also be component (sore throat, cough, SOB). Lactic acidosis with hepatic steatosis is a rare but potentially life-threatening toxicity with the use of NRTIs.

‡ Each Trizivir tablet contains 300 mg ZDV, 150 mg 3TC, and 300 mg abacavir.

¶ Cases of fatal and nonfatal pancreatitis have occurred in treatment-naïve and treatment-experienced patients during therapy with ddI or in combination with other drugs, particularly d4T or d4T + hydroxyurea.

Pregnant women may be at increased risk for lactic acidosis and liver damage when treated with the combination of stavudine and didanosine. This combination should be used in pregnant women only when the potential benefit clearly outweighs the potential risk.

**Patients who develop signs or symptoms of hypersensitivity (which may include fever, rash, fatigue, nausea, vomiting, diarrhea, and abdominal pain) should discontinue abacavir as soon as a hypersensitivity reaction is suspected. Abacavir should not be re-started, because more severe symptoms will recur within hours and may include life-threatening hypotension and death. Cases of abacavir hypersensitivity syndrome should be reported to the Abacavir Hypersensitivity Registry at 1-800-270-0425.

Table 14. Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)

Generic Name Trade Name	Nevirapine Viramune	Delavirdine Rescriptor	Efavirenz Sustiva
Form	200 mg tablets 50 mg/5 mL oral suspension	100 mg tablets 200 mg tablets	50, 100, 200 mg capsules
Dosing Recommendation	200 mg po qd x 14 days, then 200 mg po bid	400 mg po tid; 100 mg tablets can be dispersed in ≥ 3 oz water to produce slurry Separate dosing with ddI or antacids by 1 hour	600 mg po qHS
Food Effect	Take without regard to meals	Take without regard to meals	Avoid taking after high fat meals, Levels \uparrow 50%
Oral bioavailability	> 90%	85%	Data not available
Serum half-life	25-30 hours	5.8 hours	40-55 hours
Elimination	Metabolized by cytochrome P450 (3A inducer); 80% excreted in urine (glucuronidated metabolites, < 5% unchanged), 10% in feces	Metabolized by cytochrome P450 (3A inhibitor) 51% excreted in urine (<5% unchanged), 44% in feces	Metabolized by cytochrome P450 (3A mixed inducer/inhibitor); 14-34% excreted in urine (glucuronidated metabolites, < 1% unchanged), 16-61% in feces.
Adverse Events	Rash* Increased transaminase levels Hepatitis	Rash* Increased transaminase levels Headaches	Rash* Central nervous system symptoms [†] Increase transaminase levels False positive cannabinoid test Teratogenic in monkeys [‡]
Drug Interactions	For information on Drug Interactions please see Table 18.		

- In clinical trials, the NNRTI was discontinued because of rash in 7% of patients taking nevirapine, 4.3% of patients taking delavirdine, and 1.7% of patients taking efavirenz. Rare cases of Stevens-Johnson Syndrome have been reported with the use of all three NNRTIs.

[†] May include dizziness, somnolence, insomnia, abnormal dreams, confusion, abnormal thinking, impaired concentration, amnesia, agitation, depersonalization, hallucinations, and euphoria. The overall frequency of any of these symptoms associated with use of efavirenz was 52% compared with 26% in controls; 2.6% of those on efavirenz discontinued the drug due to these symptoms; symptoms usually subside spontaneously over 2-4 weeks.

[‡] No data are available regarding teratogenicity of other NNRTIs in non-human primates.

Table 15. Characteristics of Protease Inhibitors (PIs)

Generic Name Trade Name	Indinavir Crixivan	Ritonavir Norvir	Nelfinavir Viracept
Form	200, 333, 400 mg capsules	100 mg capsules 600 mg/7.5 mL po solution	250 mg tablets 50 mg/g oral powder
Dosing Recommendations	800 mg q8h Separate dosing with ddI by 1 hour	600 mg q12h* Separate dosing with ddI by 2 hours	750 mg tid or 1250mg bid
Food Effect	Levels decrease 77% Take 1 hour before or 2 hours after meals; may take with skim milk or low fat meal	Levels increase 15% Take with food if possible; this may improve tolerability	Levels increase 2-3 fold Take with meal or snack
Oral bioavailability	65%	(not determined)	20-80%
Serum half-life	1.5-2 hours	3-5 hours	3.5-5 hours
Route of Metabolism	P450 cytochrome 3A4 inhibitor (less than ritonavir)	P450 cytochrome 3A4 > 2D6 Potent 3A4 inhibitor	P450 cytochrome 3A4 inhibitor (less than ritonavir)
Storage	Room temperature	Refrigerate capsules Oral solution should NOT be refrigerated	Room temperature
Adverse Effects	<ul style="list-style-type: none"> • Nephrolithiasis • GI intolerance, nausea • Lab: Increased indirect bilirubinemia (inconsequential) • Misc.: Headache, asthenia, blurred vision, dizziness, rash, metallic taste, thrombocytopenia, alopecia • Hyperglycemia[†] • Fat redistribution and lipid abnormalities[‡] • Possible increased bleeding episodes in patients with hemophilia 	<ul style="list-style-type: none"> • GI intolerance, nausea, vomiting, diarrhea • Paresthesias – circumoral and extremities • Hepatitis • Pancreatitis • Asthenia • Taste perversion • Lab.: Triglycerides increase > 200%, transaminase elevation, elevated CPK and uric acid • Hyperglycemia[†] • Fat redistribution and lipid abnormalities[‡] • Possible increased bleeding episodes in patients with hemophilia 	<ul style="list-style-type: none"> • Diarrhea • Hyperglycemia[†] • Fat redistribution and lipid abnormalities[‡] • Possible increased bleeding episodes in patients with hemophilia
Drug Interactions	For more information on Drug Interactions please see Table 18.		

* Dose escalation for Ritonavir: Day 1-2: 300 mg bid; day 3-5: 400 mg bid; day 6-13: 500 mg bid; day 14: 600 mg bid
Combination treatment regimen with Saquinavir (400 mg po bid) plus Ritonavir (400 mg po bid).

[†] Cases of worsening glycemic control in patients with pre-existing diabetes, and cases of new-onset diabetes, including diabetic ketoacidosis, have been reported with the use of all protease inhibitors.

[‡] Fat redistribution and lipid abnormalities have been increasingly recognized with the use of protease inhibitors. Patients with hypertriglyceridemia or hypercholesterolemia should be evaluated for risk for cardiovascular events and pancreatitis. Possible interventions include dietary modification, lipid lowering agents, or discontinuation of PIs.

Table 15. Characteristics of Protease Inhibitors (PIs)

Generic Name Trade Name	Saquinavir		Amprenavir	Lopinavir + Ritonavir
	Invirase	Fortovase	Agenerase	Kaletra
Form	200 mg capsules	200 mg capsules	50 mg, 150 mg capsules 15 mg/mL oral solution (capsules and solution NOT interchangeable on mg per mg basis)	133.3 mg lopinavir + 33.3 mg ritonavir capsules 80 mg lopinavir + 20 mg ritonavir per ml oral solution
Dosing Recommendations	400 mg bid with ritonavir; Invirase not recommended otherwise	1,200 mg tid §	>50 kg: 1200 mg bid (capsules) 1400 mg bid (oral solution) < 50 kg: 20mg/kg bid (capsules) maximum 2400 mg daily total. <50 kg: 1.5mL/kg bid (oral solution) maximum 2800 mg daily total	400 mg lopinavir + 100 mg ritonavir bid
Food Effect	No food effect when taken with ritonavir	Levels increase 6-fold. Take with large meal	High fat meal decreases AUC 21%; can be taken with or without food, but high fat meal should be avoided.	Moderate fat meal increases AUC of capsules and solution by 48% and 80%, respectively. Take with food.
Oral bioavailability	Hard gel capsule: 4% erratic	Soft gel capsule (not determined)	Not determined in humans	Not determined in humans
Serum half-life	1-2 hours	1-2 hours	7.1-10.6 hours	5-6 hours
Route of Metabolism	P450 cytochrome 3A4 inhibitor (less than ritonavir)	P450 cytochrome 3A4 inhibitor (less than ritonavir)	P450 cytochrome 3A4 inhibitor (less than ritonavir; similar to indinavir, nelfinavir)	P450 cytochrome 3A4 inhibitor
Storage	Room temperature	Refrigerate or store at room temperature (up to 3 months)	Room temperature	Refrigerated capsules stable until date on label. If stored at room temperature, stable for 2 months
Adverse Effects	<ul style="list-style-type: none"> GI intolerance, nausea and diarrhea Headache Elevated transaminase enzymes Hyperglycemia † Fat redistribution and lipid abnormalities ‡ Possible increased bleeding episodes in patients with hemophilia 	<ul style="list-style-type: none"> GI intolerance, nausea, diarrhea, abdominal pain and dyspepsia Headache Elevated transaminase enzymes Hyperglycemia † Fat redistribution and lipid abnormalities ‡ Possible increased bleeding episodes in patients with hemophilia 	<ul style="list-style-type: none"> GI intolerance, nausea, vomiting, diarrhea Rash Oral paresthesias Lab: Increase in liver function tests Hyperglycemia † Fat redistribution and lipid abnormalities ‡ Possible increased bleeding episodes in patients with hemophilia Oral solution contains propylene glycol; contraindicated in pregnant women and children <4 years old, patients with hepatic or renal failure, and patients treated with disulfiram or metronidazole 	<ul style="list-style-type: none"> GI intolerance, nausea, vomiting, diarrhea Asthenia Elevated transaminase enzymes Hyperglycemia † Fat redistribution and lipid abnormalities ‡ Possible increased bleeding episodes in patients with hemophilia Oral solution contains 42% alcohol
Drug Interactions	For more information on Drug Interactions please see Table 18.			

§ Saquinavir soft gel capsule given as 1600 bid produced lower daily exposure and trough serum concentrations compared with the standard 1200 mg tid regimen. Trends in immunologic and virologic responses favored the standard tid regimen. The clinical significance of the inferior trends observed in the bid dosing group are not known; however, until the availability of the results from longer follow-up studies, bid dosing of saquinavir soft gel capsules is not recommended.

† Cases of worsening glycemic control in patients with pre-existing diabetes, and cases of new-onset diabetes, including diabetic ketoacidosis, have been reported with the use of all protease inhibitors.

‡ Fat redistribution and lipid abnormalities have been increasingly recognized with the use of protease inhibitors. Discontinuation of PIs may be required to reverse fat redistribution. Patients with hypertriglyceridemia or hypercholesterolemia should be evaluated for risk for cardiovascular events and pancreatitis. Possible interventions include dietary modification, lipid lowering agents or discontinuation of PIs.

Table 16: Adverse Drug Reactions and Related “Black Box Warnings” in Product Labeling for Antiretroviral Agents

Special problems associated with a prescription drug, particularly those that may lead to death or serious injury, may be required by the Food and Drug Administration to be placed in a prominently displayed box, more commonly known as a “black box.” Please note that there are other serious toxicities associated with some antiretroviral agents not listed in this table. For more extensive lists of adverse effects associated with individual antiretroviral drugs, or for drug interactions, please refer to [Tables 13, 14, 15, 17](#) and [18](#).

Anti-Retroviral Drug	Pertinent Black Box Warning Information
Abacavir (Ziagen™), or as combination product with zidovudine and lamivudine as Trizivir™)	<ul style="list-style-type: none"> • Fatal hypersensitivity reactions reported: <ul style="list-style-type: none"> ✓ Signs or symptoms include: fever, skin rash, fatigue, gastrointestinal symptoms (nausea, vomiting, diarrhea, or abdominal pain), and respiratory symptoms (pharyngitis, dyspnea, or cough) ✓ Abacavir should be discontinued as soon as hypersensitivity reaction is suspected ✓ Abacavir SHOULD NOT be restarted ✓ If restarted, more severe symptoms will recur within hours and may include life-threatening hypotension and death • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside analogues alone or in combination
Amprenavir (Agenerase™) Oral Solution	<ul style="list-style-type: none"> • Because of the potential risk of toxicity from large amounts of the excipient propylene glycol in Agenerase Oral Solution, it is contraindicated in the following patient populations: <ul style="list-style-type: none"> ✓ children age < 4 years ✓ pregnant women ✓ patients with renal or hepatic failure ✓ patients treated with disulfiram or metronidazole • Oral solution should be used only when Agenerase capsules or other protease inhibitors cannot be used
Delavirdine (Rescriptor™)	None
Didanosine (Videx™, Videx-EC™)	<ul style="list-style-type: none"> • Fatal and nonfatal pancreatitis have occurred with didanosine alone or in combination with other antiretroviral agents <ul style="list-style-type: none"> ✓ Didanosine should be held if pancreatitis is suspected ✓ Didanosine should be discontinued if pancreatitis is confirmed • Fatal lactic acidosis has been reported in pregnant women who received a combination of didanosine and stavudine along with other antiretroviral combinations <ul style="list-style-type: none"> ✓ Didanosine and stavudine combination should only be used during pregnancy if the potential benefit clearly outweighs the potential risks • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside analogues alone or in combination
Efavirenz (Sustiva™)	None
Indinavir (Crixivan™)	None
Lamivudine (Epivir™), or as combination product in Combivir™ and Trizivir™)	<ul style="list-style-type: none"> • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside analogues alone or in combination
Lopinavir/ritonavir (Kaletra™)	None

Anti-Retroviral Drug	Pertinent Black Box Warning Information
Nelfinavir (Viracept™)	None
Nevirapine (Viramune™)	<ul style="list-style-type: none"> • Severe, life-threatening hepatotoxicity, including fulminant and cholestatic hepatitis, hepatic necrosis, and hepatic failure. Patients should be advised to seek medical evaluation immediately should signs and symptoms of hepatitis occur. • Severe, life-threatening, and even fatal skin reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, and hypersensitivity reactions characterized by rash, constitutional findings, and organ dysfunction have occurred with nevirapine treatment • Patients should be monitored intensively during the first 12 weeks of nevirapine therapy to detect potentially life-threatening hepatotoxicity or skin reactions. • A 14-day lead-in period with nevirapine 200mg daily must be strictly followed. • Nevirapine should not be restarted after severe hepatic, skin, or hypersensitivity reactions
Ritonavir (Norvir™)	<ul style="list-style-type: none"> • Co-administration of ritonavir with certain medications may result in potentially serious and/or life-threatening adverse events due to effects of ritonavir on hepatic metabolism of certain drugs
Saquinavir (Fortovase™, Invirase™)	None
Stavudine (Zerit™)	<ul style="list-style-type: none"> • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside analogues alone or in combination • Fatal lactic acidosis has been reported in pregnant women who received a combination of stavudine and didanosine along with other antiretroviral combinations <ul style="list-style-type: none"> ✓ Stavudine and didanosine combination should only be used during pregnancy if the potential benefit clearly outweighs the potential risks • Fatal and non-fatal pancreatitis have occurred when stavudine was part of a combination regimen with didanosine with or without hydroxyurea
Tenofovir (Viread™)	<ul style="list-style-type: none"> • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogs alone or in combination with other antiretrovirals
Zalcitabine (Hivid™)	<ul style="list-style-type: none"> • Zalcitabine can cause severe peripheral neuropathy, use with caution in patients with pre-existing neuropathy • It may rarely cause pancreatitis, therapy should be held until pancreatitis is excluded • Rare cases of hepatic failure and death have been reported in patients with underlying hepatitis B infection • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside analogues alone or in combination
Zidovudine (Retrovir™), or as combination products in Combivir™ and Trizivir™	<ul style="list-style-type: none"> • Zidovudine may be associated with hematologic toxicities, including granulocytopenia and severe anemia, particularly in advanced HIV patients • Prolonged zidovudine use has been associated with symptomatic myopathy • Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of antiretroviral nucleoside analogues alone or in combination

Table 17. Drugs That Should Not Be Used With PI Antiretrovirals

Drug Category	Indinavir	Ritonavir*	Saquinavir	Nelfinavir	Amprenavir	Lopinavir + Ritonavir
Ca++ channel blocker	(none)	bepidil	(none)	(none)	bepidil	(none)
Cardiac	(none)	amiodarone flecainide propafenone quinidine	(none)	(none)	(none)	flecainide propafenone
Lipid Lowering Agents	simvastatin lovastatin	simvastatin lovastatin	simvastatin lovastatin	simvastatin lovastatin	simvastatin lovastatin	simvastatin lovastatin
Anti-Mycobacterial	rifampin	(none)	rifampin rifabutin	rifampin	rifampin	rifampin
Antihistamine	astemizole terfenadine	astemizole terfenadine	astemizole terfenadine	astemizole terfenadine	astemizole terfenadine	astemizole terfenadine
Gastrointestinal Drugs	cisapride	cisapride	cisapride	cisapride	cisapride	cisapride
Neuroleptic	(none)	clozapine pimozide	(none)	(none)	(none)	pimozide
Psychotropic	midazolam triazolam	midazolam triazolam	midazolam triazolam	midazolam triazolam	midazolam triazolam	midazolam triazolam
Ergot Alkaloids (vasoconstrictor)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)
Herbs	St. John's wort	St. John's wort	St. John's wort	St. John's wort	St. John's wort	St. John's wort

* Some of the contraindicated drugs listed are based on theoretical considerations. Thus, drugs with low therapeutic indices yet with suspected major metabolic contribution from cytochrome P450 3A, CYP2D6, or unknown pathways, are included in this table. Actual interactions may or may not occur in patients.

† This is likely a class effect.

Suggested Alternatives

Simvastatin, lovastatin: atorvastatin, pravastatin, fluvastatin, cerivastatin (alternatives should be used with caution)

Rifabutin: clarithromycin, azithromycin (MAI prophylaxis); clarithromycin, azithromycin, ethambutol (MAI treatment)

Astemizole, terfenadine: loratadine, fexofenadine, cetirizine

Midazolam, triazolam: temazepam, lorazepam

Table 17. Drugs That Should Not Be Used With NNRTI Antiretrovirals

Drug Category	Nevirapine	Delavirdine	Efavirenz
Ca⁺⁺ channel blocker	(none)	(none)	(none)
Cardiac	(none)	(none)	(none)
Lipid Lowering Agents	(none)	simvastatin lovastatin	(none)
Anti-Mycobacterial	Insufficient data	rifampin rifabutin	(none)
Antihistamine	(none)	astemizole terfenadine	astemizole terfenadine
Gastrointestinal Drugs	(none)	cisapride H-2 blockers Proton pump inhibitors	cisapride
Neuroleptic	(none)	(none)	(none)
Psychotropic	(none)	midazolam triazolam	midazolam triazolam
Ergot Alkaloids (vasoconstrictor)	(none)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)	dihydroergotamine (D.H.E. 45) ergotamine [†] (various forms)

† This is likely a class effect.

Suggested Alternatives

Simvastatin, lovastatin: atorvastatin, pravastatin, fluvastatin, cerivastatin (alternatives should be used with caution)

Rifabutin: clarithromycin, azithromycin (MAI prophylaxis); clarithromycin, ethambutol (MAI treatment)

Astemizole, terfenadine: loratadine, fexofenadine, cetirizine

Midazolam, triazolam: temazepam, lorazepam

Table 18. Drug Interactions Between Antiretrovirals and Other Drugs: Protease Inhibitors (PIs)

Drug Interactions Requiring Dose Modifications or Cautious Use			
Drugs Affected	Indinavir (IDV)	Ritonavir* (RTV)	Saquinavir† (SQV)
ANTIFUNGALS			
Ketoconazole	Levels: IDV ↑ 68% Dose: IDV 600 mg tid	Levels: ketoconazole ↑ 3X Dose: Use with caution; do not exceed 200 mg ketoconazole daily	Levels: SQV ↑ 3X Dose: Standard
ANTI-MYCOBACTERIALS			
Rifampin	Levels: IDV ↓ 89% Contraindicated	Levels: RTV ↓ 35% Dose: No Data Increased liver toxicity possible	Levels: SQV ↓ 84% Contraindicated, unless using RTV+SQV, then use rifampin 600 mg qd or 2-3x/week
Rifabutin	Levels: IDV ↓ 32% Rifabutin ↑ 2X Dose: ↓ rifabutin to 150 mg qd or 300 mg 2-3x/week IDV 1000 mg tid	Levels: Rifabutin ↑ 4X Dose: ↓ rifabutin to 150 mg qod Or dose 3x per week RTV: Standard	Levels: SQV ↓ 40% No dose adjustment unless using RTV+SQV, then use rifabutin 150 mg 2-3x/week
Clarithromycin	Levels: Clarithromycin ↑ 53% No dose adjustment	Levels: Clarithromycin ↑ 77% Dose adjust for renal insufficiency	Levels: Clarithromycin ↑ 45% SQV ↑ 177% No dose adjustment
ORAL CONTRACEPTIVES	Levels: Norethindrone ↑ 26% Ethinylestradiol ↑ 24% No dose adjustment	Levels: Ethinyl estradiol ↓ 40% Use alternative or additional method	No data
LIPID LOWERING AGENTS			
Simvastatin Lovastatin	Levels: Potential for large increase in statin levels. Avoid concomitant use.	Levels: Potential for large increase in statin levels. Avoid concomitant use.	Levels: Potential for large increase in statin levels. Avoid concomitant use.
ANTICONVULSANTS			
Phenobarbitol Phenytoin Carbamazepine	Carbamazepine markedly ↓ IDV AUC. Consider alternative agent	Unknown Use with caution Monitor anticonvulsant levels.	Unknown but may decrease SQV levels substantially Monitor anticonvulsant levels.
METHADONE	No change in methadone levels	Methadone ↓ 37%. Monitor and titrate dose if needed. May require ↑ methadone dose.	No data
MISCELLANEOUS	Grapefruit juice ↓ IDV levels by 26% Sildenafil AUC ↑ 2-11 fold. Do not exceed 25 mg in a 48 hr. period	Desipramine ↑ 145%, reduce dose Theophylline ↓ 47%, monitor theo. levels. Many possible interactions Sildenafil AUC ↑ 2-11 fold. Do not exceed 25 mg in a 48 hr. period	Grapefruit juice increases SQV levels Dexamethasone decreases SQV levels Sildenafil AUC ↑ 2-11 fold. Use a 25 mg starting dose of sildenafil.

* Drugs for which plasma concentrations may be decreased by coadministration with Ritonavir: anticoagulants (warfarin), anticonvulsants (phenytoin, divaproex, lamotrigine), antiparasitics (atovaquone).

† Some drug interaction studies were conducted with Invirase. May not necessarily apply to use with Fortovase.

Table 18. Drug Interactions Between Antiretrovirals and Other Drugs: Protease Inhibitors (PIs)

Drug Interactions Requiring Dose Modifications or Cautious Use			
Drugs Affected	Nelfinavir (NFV)	Amprenavir (APV)	Lopinavir (LPV)
ANTIFUNGALS			
Ketoconazole	No dose adjustment necessary	Levels: APV ↑ 31% Keto ↑ 44%. Combination under investigation	Levels: LPV AUC ↓ 13%. Keto ↑ 3-fold
ANTI-MYCOBACTERIALS			
Rifampin	Levels ↓ 82% Contraindicated	Levels: APV AUC ↓ 82% No change in rifampin AUC. Avoid concomitant use.	Levels: LPV AUC ↓ 75% Avoid concomitant use
Rifabutin	Levels: NFV ↓ 32% Rifabutin ↑ 2X Dose: ↓rifabutin to 150mg qd Or 300 mg 2-3x/week ↑ NFV dose to 1000 mg tid.	Levels: APV AUC ↓ 15% Rifabutin ↑ 193% Dose: No change in APV dose; Decrease rifabutin to 150 mg qd or 300 mg 2-3x/week.	Levels: Rifabutin AUC ↑ 3-fold 25-O-desacetyl metabolite ↑ 47.5-fold. Decrease rifabutin dose to 150 mg qod LPV/r: Standard
Clarithromycin	No data	Levels: APV AUC ↑ 18%. No change in clarithromycin AUC. No dose adjustment	No data
ORAL CONTRACEPTIVES	Levels: Norethindrone ↓ 18% Ethinyl estradiol ↓ 47% Use alternative or additional method	Levels: Potential for metabolic interactions; use alternative or additional method.	Levels: ethinyl estradiol ↓ 42% Use alternative or additional method
LIPID LOWERING AGENTS			
Simvastatin Lovastatin Atorvastatin Pravastatin	Levels: Potential for large increase in statin levels. Avoid concomitant use. Atorvastatin AUC ↑ 74%—use with caution. Simvastatin AUC ↑ 505%—not recommended. Potential for large increase in Lovastatin AUC—not recommended.	Levels: Potential for large increase in statin levels. Avoid concomitant use with lovastatin and simvastatin.	Levels: Potential for large increase in statin levels. Avoid concomitant use. Atorvastatin AUC ↑ 5.88-fold. Use with caution and monitoring. Pravastatin AUC ↑ 33%; no dosage adjustment necessary
ANTICONVULSANTS			
Phenobarbitol Phenytoin Carbamazepine	Unknown, but may decrease NFV levels substantially Monitor anticonvulsant levels.	Unknown, but may decrease APV levels substantially Monitor anticonvulsant levels.	Unknown, but may decrease LPV levels substantially. Monitor anticonvulsant levels.
METHADONE	NFV may decrease methadone levels, but minimal effect on maintenance dose. Monitor and titrate dose if needed. May require ↑ methadone dose.	No data	Methadone AUC ↓ 53% Monitor and titrate dose if needed. May require ↑ methadone dose.
SILDENAFIL	Sildenafil AUC ↑ 2-11 fold. Do not exceed 25 mg in a 48 hr period.	Sildenafil AUC ↑ 2-11 fold. Do not exceed 25 mg in a 48 hr period.	Probable substantial ↑ in sildenafil AUC. Do not exceed 25 mg in a 48 hr period.

**Table 18. Drug Interactions Between Antiretrovirals and Other Drugs:
Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)**

Drug Interactions Requiring Dose Modifications or Cautious Use			
Drugs Affected	Nevirapine (NVP)	Delavirdine (DLV)	Efavirenz (EFV)
ANTIFUNGALS			
Ketoconazole	Levels: Keto. ↓ 63% NVP ↑ 15-30% Dose: Not recommended	No data	No data
ANTI-MYCOBACTERIALS			
Rifampin	Levels: NVP ↓ 37% Not recommended	Levels: DLV ↓ 96% Contraindicated	Levels: EFV ↓ 25% No dose adjustment
Rifabutin	Levels: NVP ↓ 16% No dose adjustment*	Levels: DLV ↓ 80% Rifabutin ↑ 100% Not Recommended	Levels: EFV unchanged; Rifabutin ↓ 35% Dose: ↑ rifabutin dose to 450-600 mg qd or 600 mg 2-3x/week.* EFV: Standard
Clarithromycin	Levels: NVP ↑ 26%, clarithromycin ↓ 30%. No dose adjustment.	Levels: Clarithromycin ↑ 100%, DLV ↑ 44% Dose adjust for renal failure	Levels: Clarithromycin ↓ 39% Alternative recommended
ORAL CONTRACEPTIVES	Levels: ethinyl estradiol ↓ approx 20%. Use alternative or additional methods.	No data	Levels: Ethinyl estradiol ↑ 37%. No data on other component. Use alternative or additional methods
LIPID LOWERING AGENTS			
Simvastatin Lovastatin	No data	Levels: Potential for large increase in statin levels. Avoid concomitant use.	No data
ANTICONSULSANTS			
Phenobarbital Phenytoin Carbamazepine	Unknown Use with caution Monitor anticonvulsant levels.	Unknown but may decrease DLV levels substantially Monitor anticonvulsant levels.	Unknown. Use with caution Monitor anticonvulsant levels
METHADONE	Levels: NVP unchanged, methadone ↓ significantly. Titrate methadone dose to effect.	No data	Levels: methadone ↓ significantly. Titrate methadone dose to effect.
MISCELLANEOUS	No data	May increase levels of dapsone, warfarin and quinidine Sildenafil: potential for increased concentrations and adverse effects. Do not exceed 25 mg in a 48 hr period	Monitor warfarin when used concomitantly

* These recommendations apply to regimens that do not include PIs, which can substantially increase rifabutin levels.

**Table 18. Drug Interactions Between Antiretrovirals and Other Drugs:
Nucleoside/Nucleotide Reverse Transcriptase Inhibitors (NRTIs)**

Drug Interactions Requiring Dose Modifications or Cautious Use				
Drugs Affected	Zidovudine (ZDV)	Stavudine (d4T)	Didanosine (ddI)	Tenofovir
METHADONE	No data	Levels: d4T ↓ 27%, methadone unchanged. No dose adjustment.	Levels: ddI ↓ 41%, methadone unchanged. Consider ddI dose increase.	No data
MISCELLANEOUS				
Ribavirin	Ribavirin inhibits phosphorylation of ZDV; this combination should be avoided if possible.	No data	No data	No data
Didanosine buffered tablets	No data	No data	No data	<ul style="list-style-type: none"> • ddI AUC increase by 44%, Cmax increased by 28% • Monitor for ddI-associated toxicities
Cidofovir, Ganciclovir, Valganciclovir	No data	No data	No data	<ul style="list-style-type: none"> • Possibly competes for active tubular secretion with tenofovir, may increase serum concentration of these drugs and/or tenofovir • Monitor for dose-related toxicities

Table 19. Drug Interactions Involving Protease Inhibitors: Effect of Drug on Levels (AUCs)/Dose

Drug Affected	Ritonavir	Saquinavir*	Nelfinavir	Amprenavir	Lopinavir/Ritonavir
Indinavir (IDV)	Levels: IDV ↑ 2-5X Dose: IDV 400 mg bid + RTV 400 mg bid, or IDV 800 mg bid + RTV 100 or 200 mg bid	Levels: IDV no effect SQV ↑ 4-7x † Dose: Insufficient data	Levels: IDV ↑ 50% NFV ↑ 80% Dose: Limited data for IDV 1200 mg bid + NFV 1250 mg bid	Levels: APV AUC ↑ 33%. Dose: no change	Levels: IDV AUC and Cmin increased. Dose: IDV 600 mg bid
Ritonavir (RTV)	•	Levels: RTV no effect SQV ↑ 20x †‡ Dose: Inivrase or Fortovase 400 mg bid + RTV 400 mg bid	Levels: RTV no effect NFV ↑ 1.5x Dose: RTV 400 mg bid + NFV 500-750 mg bid	Levels: APV AUC ↑ 2.5-fold. Dose: Limited data for APV 600-1200 mg bid + RTV 100-200 mg bid	Lopinavir is co-formulated with ritonavir as Kaletra.
Saquinavir (SQV)	•	•	Levels: SQV ↑ 3-5x NFV ↑ 20% † Dose: Standard NFV Fortovase 800 mg tid or 1200 mg bid	Levels: APV AUC ↓ 32% Dose: insufficient data	Levels: SQV † AUC and Cmin increased Dose: SQV 800 mg bid, LPV/r standard
Nelfinavir (NFV)	•	•	•	Levels: APV AUC ↑ 1.5-fold. Dose: insufficient data	No data
Amprenavir (APV)	•	•	•	•	Levels: APV AUC and Cmin increased Dose: APV 600-750 mg bid, LPV/r standard

* Several drug interaction studies have been completed with saquinavir given as Inivrase or Fortovase. Results from studies conducted with Inivrase may not be applicable to Fortovase.

† Conducted with Fortovase

‡ Conducted with Inivrase

Table 19. Drug Interactions Involving Protease Inhibitors and Non-Nucleoside Reverse Transcriptase Inhibitors: Effect of Drug on Levels (AUCs)/Dose

Drug Affected	Nevirapine	Delavirdine	Efavirenz
Indinavir (IDV)	Levels: IDV ↓ 28% NVP no effect Dose: IDV 1000 mg q8h NVP standard	Levels: IDV ↑ >40% DLV no effect Dose: IDV 600 mg q 8h DLV: standard	Levels: IDV ↓ 31% Dose: IDV 1000mg q 8h EFV standard
Ritonavir (RTV)	Levels: RTV ↓ 11% NVP no effect Dose: Standard	Levels: RTV ↑ 70% DLV: no effect Dose: DLV: standard RTV: no data	Levels: RTV ↑ 18% EFV ↑ 21% Dose: RTV 600 mg bid (500 mg bid for intolerance) EFV standard
Saquinavir (SQV)	Levels: SQV ↓ 25% NVP no effect Dose: No data	Levels: SQV ↑ 5x‡ DLV no effect Dose: Fortovase 800 mg tid, DLV standard (monitor transaminase levels)	Levels: SQV ↓ 62%‡ EFV ↓ 12% Co-administration not recommended
Nelfinavir (NFV)	Levels: NFV ↑ 10% NVP no effect Dose: Standard	Levels: NFV ↑ 2x DLV ↓ 50% Dose: No data (monitor for neutropenic complications)	Levels: NFV ↑ 20% Dose: Standard
Amprenavir (APV)	No data	No data	Levels: APV AUC ↓ 36% Dose: APV 1200 mg tid as single PI, or 1200 mg bid + RTV 200 mg bid EFV standard
Lopinavir/Ritonavir (LPV/RTV)	Levels: LPV Cmin ↓ 55%. Dose: Consider LPV/r 533/133 mg bid in PI-experienced patients NVP standard	Levels: LPV levels expected to increase. Dose: Insufficient data	Levels: LPV AUC ↓ 40% EFV no change Dose: Consider LPV/r 533/133 mg bid in PI-experienced patients EFV standard
Nevirapine (NVP)	•	No data	Levels: NVP: no effect EFV: AUC ↓ 22%
Delavirdine (DLV)	No data	•	No data

‡ Conducted with Inivrase

Table 20. HIV-Related Drugs with Overlapping Toxicities

Bone Marrow Suppression	Peripheral Neuropathy	Pancreatitis	Nephrotoxicity	Hepato-toxicity	Rash	Diarrhea	Ocular Effects
cidofovir	didanosine	cotrimoxazole	adefovir	delavirdine	abacavir	didanosine	didanosine
cotrimoxazole	isoniazid	didanosine	aminoglycosides	efavirenz	amprenavir	clindamycin	ethambutol
cytotoxic chemotherapy	stavudine	lamivudine (children)	amphotericin B	fluconazole	cotrimoxazole	nelfinavir	rifabutin
dapsone	zalcitabine	pentamidine	cidofovir	isoniazid	dapsone	ritonavir	cidofovir
flucytosine		ritonavir	foscarnet	itraconazole	NNRTIs	lopinavir/ritonavir	
ganciclovir		stavudine	indinavir	ketoconazole	sulfadiazine	tenofovir	
hydroxyurea			pentamidine	nevirapine			
interferon- α				NRTIs			
primaquine				protease inhibitors			
pyrimethamine				rifabutin			
ribavirin				rifampin			
rifabutin							
sulfadiazine							
trimetrexate							
zidovudine							

Table 21.

Guidelines for Changing an Antiretroviral Regimen for Suspected Drug Failure

- Criteria for changing therapy include a suboptimal reduction in plasma viremia after initiation of therapy, re-appearance of viremia after suppression to undetectable, significant increases in plasma viremia from the nadir of suppression, and declining CD4⁺ T cell numbers. Please refer to the more extensive discussion of these on page 19.
- When the decision to change therapy is based on viral load determination, it is preferable to confirm with a second viral load test.
- Distinguish between the need to change a regimen due to drug intolerance or inability to comply with the regimen versus failure to achieve the goal of sustained viral suppression; single agents can be changed in the event of drug intolerance.
- In general, do not change a single drug or add a single drug to a failing regimen; it is important to use at least two new drugs and preferably to use an entirely new regimen with at least three new drugs. If susceptibility testing indicates resistance to only one agent in a combination regimen, it may be possible to replace only that drug; however, this approach requires clinical validation.
- Many patients have limited options for new regimens of desired potency; in some of these cases, it is rational to continue the prior regimen if partial viral suppression was achieved.
- In some cases, regimens identified as suboptimal for initial therapy are rational due to limitations imposed by toxicity, intolerance or nonadherence. This especially applies in late stage disease. For patients with no rational options who have virologic failure with return of viral load to baseline (pretreatment levels) and declining CD4⁺ T cell count, there should be consideration for discontinuation of antiretroviral therapy.
- Experience is limited with regimens using combinations of two protease inhibitors or combinations of protease inhibitors with NNRTIs; for patients with limited options due to drug intolerance or suspected resistance these regimens provide possible alternative treatment options.

There is limited information about the value of restarting a drug that the patient has previously received. Susceptibility testing may be useful in this situation if clinical evidence suggestive of the emergence of resistance is observed. However, testing for phenotypic or genotypic resistance in peripheral blood virus may fail to detect minor resistant variants. Thus, the presence of resistance is more useful information in altering treatment strategies than the absence of detectable resistance.

- Avoid changing from ritonavir to indinavir or vice versa for drug failure, since high level cross resistance is likely.
- Avoid changing among NNRTIs for drug failure, since high level cross resistance is likely.
- The decision to change therapy and the choice of a new regimen requires that the clinician have considerable expertise in the care of people living with HIV. Physicians who are less experienced in the care of persons with HIV infection are strongly encouraged to obtain assistance through consultation with or referral to a clinician with considerable expertise in the care of HIV-infected patients.

Table 22. Acute Retroviral Syndrome: Associated Signs and Symptoms (Expected Frequency) [see reference 201]

- ◆ Fever (96%)
- ◆ Lymphadenopathy (74%)
- ◆ Pharyngitis (70%)
- ◆ Rash (70%)
 - ✓ Erythematous maculopapular with lesions on face and trunk and sometimes extremities, including palms and soles.
 - ✓ Mucocutaneous ulceration involving mouth, esophagus or genitals.
- ◆ Myalgia or arthralgia (54%)
- ◆ Diarrhea (32%)
- ◆ Headache (32%)
- ◆ Nausea and vomiting (27%)
- ◆ Hepatosplenomegaly (14%)
- ◆ Weight Loss (13%)
- ◆ Thrush (12%)
- ◆ Neurologic symptoms (12%)
 - ✓ Meningoencephalitis or aseptic meningitis
 - ✓ Peripheral neuropathy or radiculopathy
 - ✓ Facial palsy
 - ✓ Guillain-Barré syndrome
 - ✓ Brachial neuritis
 - ✓ Cognitive impairment or psychosis

Table 23. Preclinical and Clinical Data Relevant to Use of Antiretrovirals in Pregnancy

Antiretroviral Drug	FDA Pregnancy Category*	Placental Passage [Newborn:Maternal Drug Ratio]	Long-Term Animal Carcinogenicity Studies	Rodent Teratogen
zidovudine [†]	C	Yes (human) [0.85]	Positive (rodent, vaginal tumors)	Positive (near lethal dose)
zalcitabine	C	Yes (rhesus) [0.30 – 0.50]	Positive (rodent, thymic lymphomas)	Positive (hydrocephalus at high dose)
didanosine	B	Yes (human) [0.5]	Negative (no tumors, lifetime rodent study)	Negative
stavudine	C	Yes (rhesus) [0.76]	Positive (rodent, liver and bladder tumors)	Negative (but sternal bone calcium decreases)
lamiduvine	C	Yes (human) [~1.0]	Negative (no tumors, lifetime rodent study)	Negative
abacavir	C	Yes (rats)	Not completed	Positive (anasarca and skeletal malformations at 1000 mg/kg [35x human exposure] during organogenesis)
saquinavir	B	Unknown	Not completed	Negative
indinavir	C	Yes (rats) (“Significant” in rats, low in rabbits)	Not completed	Negative (but extra ribs in rats)
ritonavir	B	Yes (rats) [mid-term fetus, 1.15; late-term fetus, 0.15 – 0.64]	Positive (rodent, liver tumors)	Negative (but cryptorchidism in rats) [‡]
nelfinavir	B	Unknown	Not completed	Negative
amprenavir	C	Unknown	Not completed	Positive (thymic elongation; incomplete ossification of bones; low body weight)
lopinavir/ ritonavir	C	Lopinavir – yes (rats) [0.08 at 6 hrs post-dose]	Lopinavir – not completed. Ritonavir – see above	Negative (but delayed skeletal ossification and increase in skeletal variations in rats at maternally toxic doses)
nevirapine	C	Yes (human) [~1.0]	Not completed	Negative
delavirdine	C	Yes (rats) [late-term fetus, blood, 0.15; late-term fetus, liver 0.04]	Positive (rodent, liver and bladder tumors)	Ventricular septal defect
efavirenz	C	Yes (cynomolgus monkeys, rats, rabbits) [~1.0]	Not completed	Anencephaly; anophthalmia; microphthalmia (cynomolgus monkeys)

* FDA Pregnancy Categories are:

- A – Adequate and well-controlled studies of pregnant women fail to demonstrate a risk to the fetus during the first trimester of pregnancy (and there is no evidence of risk during later trimesters);
- B – Animal reproduction studies fail to demonstrate a risk to the fetus and adequate but well-controlled studies of pregnant women have not been conducted;
- C – Safety in human pregnancy has not been determined, animal studies are either positive for fetal risk or have not been conducted, and the drug should not be used unless the potential benefit outweighs the potential risk to the fetus;
- D – Positive evidence of human fetal risk based on adverse reaction data from investigational or marketing experiences, but the potential benefits from the use of the drug in pregnant women may be acceptable despite its potential risks;
- X – Studies in animals or reports of adverse reactions have indicated that the risk associated with the use of the drug for pregnant women clearly outweighs any possible benefit.


[†] Despite certain animal data showing potential teratogenicity of ZDV when near-lethal doses are given to pregnant rodents, considerable human data are available to date indicating that the risk to the fetus, if any, is extremely small when given to the pregnant mother beyond 14 weeks gestation. Follow-up for up to 6 years of age for 734 infants born to HIV-infected women who had in utero exposure to ZDV has not demonstrated any tumor development (228). However, no data are available on longer follow-up for late effects.

[‡] These effects seen only at maternally toxic doses.

Table 24. Zidovudine Perinatal Transmission Prophylaxis Regimen

ANTEPARTUM	<p>Initiation at 14-34 weeks gestation and continued throughout pregnancy</p> <p>A. PACTG 076 Regimen: ZDV 100 mg 5 times daily</p> <p>B. Acceptable Alternative Regimen:</p> <p style="text-align: center;">ZDV 200 mg 3 times daily</p> <p style="text-align: center;">or</p> <p style="text-align: center;">ZDV 300 mg 2 times daily</p>
INTRAPARTUM	<p>During labor, ZDV 2 mg/kg intravenously over 1 hour, followed by a continuous infusion of 1 mg/kg/hr intravenously until delivery.</p>
POSTPARTUM	<p>Oral administration of ZDV to the newborn (ZDV syrup, 2 mg/kg every 6 hours) for the first 6 weeks of life, beginning at 8-12 hours after birth.</p>

Figure 1. Likelihood of Developing AIDS Within Three Years



MACS bDNA:	>30K	10K-30K	3K-10K	501-3K	<500
RT -PCR:	>55K	20K-55K	7K-20K	1.5K-7K	<1500

Plasma Viral Load (copies/ml)

Likelihood of developing an AIDS-related illness in three years. Viral load represents the actual data obtained on the specimens from the MACS cohort as well as the values showing the equivalent expected RT-PCR values. Values shown in this figure differ slightly from those in Table 5 because better discrimination of outcome was achieved by re-analysis of the data using viral load as the initial parameter for categorization followed by CD4⁺ T lymphocyte stratification of the patients. (Adapted from reference 4)

References:

1. Report of the NIH panel to define principles of therapy of HIV infection. *MMWR*, 1998. 47(RR-5): p. 1-41.
2. 1999 USPHS/IDSA guidelines for the prevention of opportunistic infections in persons infected with human immunodeficiency virus. *MMWR*, 1999. 48(RR-10): p. 1-67.
3. Mellors JW, Rinaldo CR, Gupta P, et al. Prognosis in HIV-1 infection predicted by the quantity of virus in plasma. *Science*, 1996. 272(5265): p. 1167-1170.
4. Mellors JW, Munoz A, Giorgi JV, et al. Plasma viral load and CD4+ lymphocytes as prognostic markers of HIV-1 infection. *Ann Intern Med*, 1997. 126(12): p. 946-954.
5. Perelson AS, Essunger P, Cao Y, et al. Decay characteristics of HIV-1 infected compartments during combination therapy. *Nature*, 1997. 387(6629): p. 188-191.
6. Powderly WG, Saag MS, Chapman S, et al. Predictors of optimal virological response to potent antiretroviral therapy. *AIDS*, 1999. 13(14): p. 1873-1880.
7. Stein DS, Korvick JA, Vermund SH. CD4+ lymphocyte cell enumeration for prediction of clinical course of human immunodeficiency virus disease: A review. *J Infect Dis*, 1992. 165(2): p. 352-363.
8. Raboud JM, Montaner JS, Conway B, et al. Variation in plasma RNA levels, CD4 cell counts, and p24 antigen levels in clinically stable men with human immunodeficiency virus infection. *J Infect Dis*, 1996. 174(1): p. 191-194.
9. Devereux HL, Youle M, Johnson MA, Loveday C. Rapid decline in detectability of HIV-1 drug resistance mutations after stopping therapy. *AIDS*, 1999. 13(18): p. F123-F127.
10. Durant J, Clevenbergh P, Halfon P, et al. Drug-resistance genotyping in HIV-1 therapy: The VIRADAPT randomised controlled trial. *Lancet*, 1999. 353(9171): p. 2195-2199.
11. Baxter JD, Mayers DL, Wentworth DN, et al. CPCRA 046 Study Team. A pilot study of the short-term effects of antiretroviral management based on plasma genotypic antiretroviral resistance testing (GART) in patients failing antiretroviral therapy. 6th Conference on Retroviruses and Opportunistic Infections. Chicago, IL, 1999. (Abstract LB8).
12. Cohen C, Hunt S, Sension M, et al. Phenotypic resistance testing significantly improves response to therapy: A randomized trial (VIRA3001). 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 237).
13. Havlir DV, Hellmann NS, Petropoulos CJ, et al. Drug susceptibility in HIV infection after viral rebound in patients receiving indinavir-containing regimens. *JAMA*, 2000. 283(2): p. 229-234.
14. Hecht FM, Grant RM, Petropoulos CJ, et al. Sexual transmission of an HIV-1 variant resistant to multiple reverse-transcriptase and protease inhibitors. *N Engl J Med*, 1998. 339(5): p. 307-311.
15. Yerly S, Kaiser L, Race E, et al. Transmission of antiretroviral-drug-resistant HIV-1 variants. *Lancet*, 1999. 354(9180): p. 729-733.
16. Boden D, Hurley A, Zhang L, et al. HIV-1 drug resistance in newly infected individuals. *J Am Med Assoc*, 1999. 282(12): p. 1135-1141.
17. Little SJ, Daar ES, D'Aquila RT, et al. Reduced antiretroviral drug susceptibility among patients with primary HIV infection. *J Am Med Assoc*, 1999. 282(12): p. 1142-1149.
18. Kinloch-De Loes S, Hirschel BJ, Hoen B, et al. A controlled trial of zidovudine in primary human immunodeficiency virus infection. *N Engl J Med*, 1995. 333(7): p. 408-413.
19. Hoen B, Dumon B, Harzic M, et al. Highly active antiretroviral treatment initiated early in the course of symptomatic primary HIV-1 infection: Results of the ANRS 053 trial. *J Infect Dis*, 1999. 180(4): p. 1342-1346.
20. Sperling RS, Shapiro DE, Coombs RW, et al. Maternal viral load, zidovudine treatment, and the risk of transmission of human immunodeficiency virus type 1 from mother to infant. Pediatric AIDS Clinical Trials Group Protocol 076 Study Group. *N Engl J Med*, 1996. 335(22): p. 1621-1629.
21. Eastman PS, Shapiro DE, Coombs RW, et al. Maternal viral genotypic zidovudine resistance and infrequent failure of zidovudine therapy to prevent perinatal transmission of human immunodeficiency virus type 1 in pediatric AIDS Clinical Trials Group Protocol 076. *J Infect Dis*, 1998. 177(3): p. 557-564.
22. Palumbo R, Dobbs T, Holland B. Antiretroviral (ARV) resistance mutations among pregnancy, HIV-infected women: Frequency and clinical correlates. Global Strategies for the Prevention of HIV Transmission from Mothers to Infants. Montreal, Canada, 1999. (Abstract 256).
23. Centers for Disease Control and Prevention. 1993 revised classification system for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. *MMWR*, 1992. 41(51): p. 1-19.
24. Gulick RM, Mellors JW, Havlir D, et al. Treatment with indinavir, zidovudine, and lamivudine in adults with human immunodeficiency virus infection and prior antiretroviral therapy. *N Engl J Med*, 1997. 337(11): p. 734-739.
25. Hammer SM, Squires KE, Hughes MD, et al. A controlled trial of two nucleoside analogues plus indinavir in persons with human immunodeficiency virus infection and CD4 cell counts of 200 per cubic

- millimeter or less. *N Engl J Med*, 1997. 337(11): p. 725-733.
26. Cameron DW, Heath-Chiozzi M, Danner S, et al. Randomised placebo-controlled trial of ritonavir in advanced HIV-1 disease. The Advanced HIV Disease Ritonavir Study Group. *Lancet*, 1998. 351(9102): p. 543-549.
 27. Hogg RS, Yip B, Chan KJ, et al. Rates of disease progression by baseline CD4 cell count and viral load after initiating triple-drug therapy. *JAMA*, 2001. 286(20): p. 2568-2577.
 28. Chen R, Westfall A, Cloud G, et al. Long-term survival after initiation of antiretroviral therapy. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 341).
 29. Sterling TR, C.R., Bartlett JG, Moore RD. CD4+ lymphocyte level is better than HIV-1 plasma viral load in determining when to initiate HAART. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 519).
 30. Egger M. Art Cohort Collaboration. Progression of HIV-1 infected drug-naïve patients starting potent antiretroviral therapy: multicohort analysis of 12,040 patients. 41st ICAAC. Chicago, IL. 2001. (Abstract LB-18).
 31. Kaplan J, Hanson DL, Karon J, et al. Early initiation of combination antiretroviral therapy (ART): Does it affect clinical outcome? 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract LbPeB7051).
 32. Opravil M, Ledergerber, Furrer H, et al. and the Swiss HIV Cohort Study. Clinical benefit of early initiation of HAART in patients with asymptomatic HIV and CD4 counts >350/mm³. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract LB-6).
 33. Deeks SG, Hecht FM, Swanson M, et al. HIV RNA and CD4 cell count response to protease inhibitor therapy in an urban AIDS clinic: Response to both initial and salvage therapy. *AIDS*, 1999. 13(6): p. F35-43.
 34. Lucas GM, Chaisson RE, Moore RD. Highly active antiretroviral therapy in a large urban clinic: Risk factors for virologic failure and adverse drug reactions. *Ann Intern Med*, 1999. 131(2): p. 81-87.
 35. Ledergerber B, Egger M, Opravil M, et al. Clinical progression and virological failure on highly active antiretroviral therapy in HIV-1 patients: A prospective cohort study. Swiss HIV Cohort Study. *Lancet*, 1999. 353(9156): p. 863-868.
 36. Chaisson RE, Keruly JC, Moore RD. Association of initial CD4 cell count and viral load with response to highly active antiretroviral therapy. *J Am Med Assoc*, 2000. 284(24): p. 3128-3129.
 37. Farzadegan H, Hoover DR, Astemborski J, et al. Sex differences in HIV-1 viral load and progression to AIDS. *Lancet*, 1998. 352(9139): p. 1510-1514.
 38. Sterling TR, Lyles CM, Vlahov D, et al. Sex differences in longitudinal human immunodeficiency virus type 1 RNA levels among seroconverters. *J Infect Dis*, 1999. 180(3): p. 666-672.
 39. Evans JS, Nims T, Cooley J, et al. Serum levels of virus burden in early-stage human immunodeficiency virus type 1 disease in women. *J Infect Dis*, 1997. 175(4): p. 795-800.
 40. Katzenstein DA, Hammer SM, Hughes MD, et al. The relation of virologic and immunologic markers to clinical outcomes after nucleoside therapy in HIV-infected adults with 200 to 500 CD4 cells per cubic millimeter. AIDS Clinical Trials Group Study 175 Virology Study Team. *N Engl J Med*, 1996. 335(15): p. 1091-1098.
 41. Junghans C, Ledergerber B, Chan P, et al. Sex differences in HIV-1 viral load and progression to AIDS. Swiss HIV Cohort Study. *Lancet*, 1999. 353(9152): p. 589.
 42. Moroni M. Sex differences in HIV-1 viral load and progression to AIDS. ICONA Study Group. Italian cohort of HIV-1 positive individuals. *Lancet*, 1999. 353(9152): p. 589-590.
 43. Sterling TR, Vlahov D, Astemborski J, et al. Initial Plasma HIV-1 RNA levels and progression to AIDS in women and men. *N Engl J Med*, 2001. 344(10): p. 720-725.
 44. Moore RD, Cheever L, Keruly JC, Chaisson RE. Lack of sex difference in CD4 to HIV-1 RNA viral load ratio [letter]. *Lancet*, 1999. 353(9151): p. 463-464.
 45. Bush CE, Donovan RM, Markowitz N, et al. Gender is not a factor in serum human immunodeficiency virus type 1 RNA levels in patients with viremia. *J Clin Microbiol*, 1996. 34(4): p. 970-972.
 46. Anastos K, Gange SJ, Lau B, et al. Gender specific differences in quantitative HIV-1 RNA levels. 6th Conference on Retroviruses and Opportunistic Infections. Chicago, IL, 1999. (Abstract 274).
 47. Kalish LA, Collier AC, Flanigan TP, Kumar PN. Plasma HIV RNA viral load in women and men with advanced HIV disease. 39th Interscience Conference on Antimicrobial Agents and Chemotherapy. San Francisco, CA, 1999. (Abstract 2190).
 48. Prins M, Robertson JR, Brettle RP, et al. Do gender differences in CD4 cell counts matter? *AIDS*, 1999. 13(17): p. 2361-2364.
 49. Webber MP, Schoenbaum EE, Gourevitch MN, et al. A prospective study of HIV disease progression in female and male drug users. *AIDS*, 1999. 13(2): p. 257-262.
 50. Blair J, Hanson D, Jones J, et al. Do gender differences in viral load predict differences in HIV disease progression? 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 195).
 51. Paterson DL, Swindells S, Mohr J, et al. Adherence to protease inhibitor therapy and outcomes in

- patients with HIV infection. *Ann Intern Med*, 2000. 133(1): p. 21-30.
52. Carmona A, Knobel H, Guelar A, et al. Factors influencing survival in HIV infected patients treated with HAART. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract TuOrB417).
 53. Walsh JC, Hertogs K, Gazzard B. Viral drug resistance, adherence and pharmacokinetic indices in HIV-1 infected patients on successful and failing protease inhibitor based HAART. 40th Interscience Conference of Antimicrobial Agents and Chemotherapy. Toronto, Canada. 2000. (Abstract 699).
 54. Arnsten J, Demas P, Gourevitch M, et al. Adherence and viral load in HIV-infected drug users: Comparison of self-report and medication events monitors. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 69).
 55. Ickovics JR, Meisler AW. Adherence in AIDS clinical trials: A framework for clinical research and clinical care. *J Clin Epidemiol*, 1997. 50(4): p. 385-391.
 56. Chesney MA. Factors affecting adherence to antiretroviral therapy. *Clin Infect Dis*, 2000. 30 Suppl 2: p. S171-176.
 57. Bangsberg D, Tulsy JP, Hecht FM, Moss AR. Protease inhibitors in the homeless. *J Am Med Assoc*, 1997. 278(1): p. 63-65.
 58. Shapiro MF, Morton SC, McCaffrey DF, et al. Variations in the care of HIV-infected adults in the United States: Results from the HIV Cost and Services Utilization Study. *J Am Med Assoc*, 1999. 281(24): p. 2305-2315.
 59. Max B, Sherer R. Management of the adverse effects of antiretroviral therapy and medication adherence. *Clin Infect Dis*, 2000. 30 Suppl 2: p. S96-S116.
 60. Cheever L. Forum for Collaborative HIV Research. What do we know about adherence levels in different populations? Adherence to HIV therapy: Building a bridge to success. Available at <http://www.gwu.edu/~chsrp/>. Washington, D.C. 1999: p. 10.
 61. Greenberg RN. Overview of patient compliance with medication dosing: a literature review. *Clin Ther*, 1984. 6(5): p. 592-599.
 62. Crespo-Fierro M. Compliance/adherence and care management in HIV disease. *J Assoc Nurses AIDS Care*, 1997. 8(4): p. 43-54.
 63. Singh N, Squier C, Sivek C, et al. Determinants of compliance with antiretroviral therapy in patients with human immunodeficiency virus: Prospective assessment with implications for enhancing compliance. *AIDS Care*, 1996. 8(3): p. 261-269.
 64. Williams A, Friedland G. Adherence, compliance, and HAART. *AIDS Clin Care*, 1997. 9(7): p. 51-54, 58.
 65. Fowler ME. Recognizing the phenomenon of readiness: Concept analysis and case study. *J Assoc Nurses AIDS Care*, 1998. 9(3): p. 72-76.
 66. Kalichman SC, Ramachandran B, Catz S. Adherence to combination antiretroviral therapies in HIV patients of low health literacy. *J Gen Intern Med*, 1999. 14(5): p. 267-273.
 67. Mannherheimer S, Friedland G, Matts J, et al. Self-reported antiretroviral adherence correlates with HIV viral load and declines over time. 13th International AIDS Conference. Durban, South Africa, 2000. (Abstract TuOrB421).
 68. Sherer R. Adherence and antiretroviral therapy in injection drug users. *J Am Med Assoc*, 1998. 280(6): p. 567-568.
 69. Luber AD, Sherman M, Gotterer H, et al. Community collaborations between physicians and pharmacists improved adherence with HIV Consensus Panel Guidelines and enhances the care of HIV infected individuals. 40th Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto, Canada, 2000. (Abstract 800).
 70. Bangsberg DR, Hecht FM, Clague H, et al. Provider estimate and structured patient report of adherence compared with unannounced pill count. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 70).
 71. Rogers A. Forum for Collaborative HIV Research. TREAT - a clinical management tool for adolescents. Adherence to HIV therapy: Building a bridge to success. Available at <http://www.gwu.edu/~chsrp/>. Washington, D.C. 1999: p. 16-17.
 72. McPherson-Baker S, Malow RM, Penedo F, et al. Enhancing adherence to combination antiretroviral therapy in non-adherent HIV-positive men. *AIDS Care*, 2000. 12(4): p. 399-404.
 73. Nieuwkerk P, Burger D, Hugen P, et al. Patient adherence to highly active antiretroviral therapy for HIV-1 infection in a nationwide cohort study in the Netherlands. 13th International AIDS Conference. Durban, South Africa, 2000. (Abstract MoPpD1055).
 74. Ahdieh L, Silverberg M, Palacio H, et al. Factors associated with discontinuation of HAART in a large cohort of HIV+ women. 13th International AIDS Conference. Durban, South Africa, 2000. (Abstract WePeB4293).
 75. Tyndall M, Yip B, Hogg R, et al. Coverage, adherence, and sustainability of antiretroviral therapy among injection drug users in Vancouver, Canada. 13th International AIDS Conference. Durban, South Africa, 2000. (Abstract ThPeB4990).
 76. Parienti JJ, Massari V, Verdon R, et al. Adherence to antiretroviral therapy is associated with results of pills identification test. 40th Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto, Canada, 2000. (Abstract 793).

77. Knobel H, Carmona A, Grau S, et al. Strategies to optimize adherence to highly active antiretroviral treatment. 12th International AIDS Conference. Geneva, Switzerland, 1998. (Abstract 32322).
78. Bartlett J, DeMasi R, Quinn J, et al. Correlation between antiretroviral pill burden and durability of virologic response. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract ThPeB4998).
79. Cahn P. Potential advantages of a compact triple nucleoside regimen: Efficacy and adherence with combivir/abacavir versus combivir/indinavir in an open-label randomised comparative study (CNAB3014). 40th Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto, Canada. 2000. (Abstract 695).
80. Flanigan T. Forum for Collaborative HIV Research. Directly observed therapy. Adherence to HIV therapy: Building a bridge to success. Available at <http://www.gwu.edu/~chsrp/>. Washington, D.C. 1999: p. 21-22.
81. Rappaport E. Forum for Collaborative HIV Research. Correctional Institutions. Adherence to HIV therapy: Building a bridge to success. Available at <http://www.gwu.edu/~chsrp/>. Washington, D.C. 1999: p. 31-32.
82. Fischl M, Rodriguez A, Scerpella E, et al. Impact of directly observed therapy on outcomes in HIV clinical trials. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 71).
83. Stenzel MS, McKenzie M, Adelson-Mitty J, Flanigan T. Modified directly observed therapy to enhance highly active therapy: 12 month follow-up. 13th International AIDS Conference. Durban, South Africa, 2000. (Abstract ThPeB4992).
84. Chun TW, Engel D, Berrey MM, et al. Early establishment of a pool of latently infected, resting CD4(+) T cells during primary HIV-1 infection. *Proc Natl Acad Sci USA*, 1998. 95(15): p. 8869-8873.
85. Chun TW, Stuyver L, Mizell SB, et al. Presence of an inducible HIV-1 latent reservoir during highly active antiretroviral therapy. *Proc Natl Acad Sci USA*, 1997. 94(24): p. 13193-13197.
86. Finzi D, Hermankova M, Pierson T, et al. Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science*, 1997. 278(5341): p. 1295-1300.
87. Wong JK, Hezareh M, Gunthard HF, et al. Recovery of replication-competent HIV despite prolonged suppression of plasma viremia. *Science*, 1997. 278(5341): p. 1291-1295.
88. Finzi D, Blankson J, Siliciano JD, et al. Latent infection of CD4⁺ T cells provides a mechanism for lifelong persistence of HIV-1, even in patients on effective combination therapy. *Nat Med*, 1999. 5(5): p. 512-517.
89. Mocroft A, Vella S, Benfield TL, et al. Changing patterns of mortality across Europe in patients infected with HIV-1. EuroSIDA Study Group. *Lancet*, 1998. 352(9142): p. 1725-1730.
90. Palella FJ Jr, Delaney KM, Moorman AC, et al. Declining morbidity and mortality among patients with advanced human immunodeficiency virus infection. HIV Outpatient Study Investigators. *N Engl J Med*, 1998. 338(13): p. 853-860.
91. Vittinghoff E, Scheer S, O'Malley P, et al. Combination antiretroviral therapy and recent declines in AIDS incidence and mortality. *J Infect Dis*, 1999. 179(3): p. 717-720.
92. O'Brien WA, Hartigan PM, Martin D, et al. Changes in plasma HIV-1 RNA and CD4⁺ lymphocyte counts and the risk of progression to AIDS. *N Engl J Med*, 1996. 334(7): p. 426-431.
93. Staszewski S, Miller V, Sabin C, et al. Determinants of sustainable CD4 lymphocyte count increases in response to antiretroviral therapy. *AIDS*, 1999. 13(8): p. 951-956.
94. Miller V, Staszewski S, Sabin C, et al. CD4 lymphocyte count as a predictor of the duration of highly active antiretroviral therapy-induced suppression of human immunodeficiency virus load. *J Infect Dis*, 1999. 180(2): p. 530-533.
95. Kaufmann D, Pantaleo G, Sudre P, Telenti A. CD4-cell count in HIV-1-infected individuals remaining viraemic with highly active antiretroviral therapy (HAART). Swiss HIV Cohort Study [letter]. *Lancet*, 1998. 351(9104): p. 723-724.
96. Autran B, Carcelain G, Li TS, et al. Positive effects of combined antiretroviral therapy on CD4⁺ T cell homeostasis and function in advanced HIV disease. *Science*, 1997. 277(5322): p. 112-116.
97. Pakker NG, Notermans DW, de Boer RJ, et al. Biphasic kinetics of peripheral blood T cells after triple combination therapy in HIV-1 infection: A composite of redistribution and proliferation. *Nat Med*, 1998. 4(2): p. 208-214.
98. Gorochoy G, Neumann AU, Kereveur A, et al. Perturbation of CD4⁺ and CD8⁺ T-cell repertoires during progression to AIDS and regulation of the CD4⁺ repertoire during antiviral therapy. *Nat Med*, 1998. 4(2): p. 215-221.
99. McCune JM, Loftus R, Schmidt DK, et al. High prevalence of thymic tissue in adults with human immunodeficiency virus-1 infection. *J Clin Invest*, 1998. 101(11): p. 2301-2308.
100. Douek DC, McFarland RD, Keiser PH, et al. Changes in thymic function with age and during the treatment of HIV infection. *Nature*, 1998. 396(6712): p. 690-695.
101. Komanduri KV, Viswanathan MN, Wieder ED, et al. Restoration of cytomegalovirus-specific CD4⁺ T-lymphocyte responses after ganciclovir and highly active antiretroviral therapy in individuals infected with HIV-1. *Nat Med*, 1998. 4(8): p. 953-956.

102. Schneider MM, Borleffs JC, Stolk RP, et al. Discontinuation of prophylaxis for *Pneumocystis carinii* pneumonia in HIV-1-infected patients treated with highly active antiretroviral therapy. *Lancet*, 1999. 353(9148): p. 201-203.
103. Weverling GJ, Mocroft A, Ledergerber B, et al. Discontinuation of *Pneumocystis carinii* pneumonia prophylaxis after start of highly active antiretroviral therapy in HIV-1 infection. EuroSIDA Study Group. *Lancet*, 1999. 353(9161): p. 1293-1298.
104. Furrer H, Egger M, Opravil M, et al. Discontinuation of primary prophylaxis against *Pneumocystis carinii* pneumonia in HIV-1-infected adults treated with combination antiretroviral therapy. Swiss HIV Cohort Study. *N Engl J Med*, 1999. 340(17): p. 1301-1306.
105. Acosta EP, Henry K, Baken L, et al. Indinavir concentrations and antiviral effect. *Pharmacotherapy*, 1999. 19(6): p. 708-712.
106. Montaner JS, Reiss P, Cooper D, et al. A randomized, double-blind trial comparing combinations of nevirapine, didanosine, and zidovudine for HIV-infected patients: The INCAS Trial. Italy, The Netherlands, Canada and Australia Study. *J Am Med Assoc*, 1998. 279(12): p. 930-937.
107. Staszewski S, Morales-Ramirez J, Tashima KT, et al. Efavirenz plus zidovudine and lamivudine, efavirenz plus indinavir, and indinavir plus zidovudine and lamivudine in the treatment of HIV-1 infection in adults. *N Engl J Med*, 1999. 341(25): p. 1865-1873.
108. Staszewski S, Keiser P, Gathe J, et al. Comparison of antiviral response with abacavir/combivir to indinavir/combivir in therapy-naive adults at 48 weeks (CNA3005). 39th Interscience Conference on Antimicrobial Agents and Chemotherapy. San Francisco, CA, 1999. (Abstract 505).
109. D'Aquila RT, Johnson VA, Welles SL, et al. Zidovudine resistance and HIV-1 disease progression during antiretroviral therapy. AIDS Clinical Trials Group Protocol 116B/117 Team and the Virology Committee Resistance Working Group. *Ann Intern Med*, 1995. 122(6): p. 401-408.
110. Deeks SG, Hellmann NS, Grant RM, et al. Novel four-drug salvage treatment regimens after failure of a human immunodeficiency virus type 1 protease inhibitor-containing regimen: Antiviral activity and correlation of baseline phenotypic drug susceptibility with virologic outcome. *J Infect Dis*, 1999. 179(6): p. 1375-1381.
111. Parkin NT, Lie YS, Hellmann N, et al. Phenotypic changes in drug susceptibility associated with failure of human immunodeficiency virus type 1 (HIV-1) triple combination therapy. *J Infect Dis*, 1999. 180(3): p. 865-870.
112. Hsu A, Granneman GR, Cao G, et al. Pharmacokinetic interactions between two human immunodeficiency virus protease inhibitors, ritonavir and saquinavir. *Clin Pharmacol Ther*, 1998. 63(4): p. 453-464.
113. Sham HL, Kempf DJ, Molla A, et al. ABT-378, a highly potent inhibitor of the human immunodeficiency virus protease. *Antimicrob Agents Chemother*, 1998. 42(12): p. 3218-3224.
114. Piscitelli S, Bechtel C, Sadler B, Falloon J. The addition of a second protease inhibitor eliminated amprenavir-efavirenz interactions and increased amprenavir concentrations. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 90).
115. Saah AJ, Winchell G, Seniuk M, Deutsch P. Multiple-dose pharmacokinetics and tolerability of indinavir-ritonavir combinations in healthy volunteers. 6th Conference on Retroviruses and Opportunistic Infections. Chicago, IL, 1999. (Abstract 136).
116. Kilby M, Hill AM, Buss N. The effect of ritonavir on increases in saquinavir plasma concentration is independent of ritonavir dosage: Combined analysis of 120 subjects. 40th Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto, Canada, 2000. (Abstract 336).
117. Kurowski M, Kaeser B, Mroziekiewicz A, et al. The influence of low doses of ritonavir on the pharmacokinetics of nelfinavir 1250 mg bid. 40th Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto, Canada, 2000. (Abstract 333).
118. Cameron DW, Japour AJ, Xu Y, et al. Ritonavir and saquinavir combination therapy for the treatment of HIV infection. *AIDS*, 1999. 13(2): p. 213-224.
119. Murphy RL, Brun S, Hicks C, et al. ABT-378/ritonavir plus stavudine and lamivudine for the treatment of antiretroviral-naive adults with HIV-1 infection: 48-week results. *AIDS*, 2001. 15(1): p. F1-F9.
120. Clinical update: impact of HIV protease inhibitors on the treatment of HIV-infected tuberculosis patients with rifampin. *MMWR*, 1996. 45(42): p. 921-925.
121. John M, Moore C, James I, et al. Chronic hyperlactatemia in HIV-infected patients taking antiretroviral therapy. *AIDS*, 2001. 15(6): p. 717-723.
122. Harris M, Tesiorowski A, Chan K, et al. Lactic acidosis complicating antiretroviral therapy: Frequency and correlates. *Antiviral Therapy*, 2000. 5(Suppl 2):31.
123. Fortgang IS, Belitsos PC, Chaisson RE, Moore RD. Hepatomegaly and steatosis in HIV-infected patients receiving nucleoside analog antiretroviral therapy. *Am J Gastroenterol*, 1995. 90(9): p. 1433-1436.
124. Boubaker K, Sudre P, Flepp M, et al. Hyperlactatemia and antiretroviral therapy in the Swiss HIV Cohort Study. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 57).

125. Boxwell DE, Styrts BA. Lactic acidosis in patients receiving nucleoside reverse transcriptase inhibitors. 39th Interscience Conference on Antimicrobial Agents and Chemotherapy. San Francisco, CA, 1999. (Abstract 1284).
126. ter Hofstede HJ, de Marie S, Foudraine NA, et al. Clinical features and risk factors of lactic acidosis following long-term antiretroviral therapy: 4 fatal cases. *Int J STD AIDS*, 2000. 11(9): p. 611-616.
127. Luzzati R, Del Bravo P, Di Perri G, et al. Riboflavin and severe lactic acidosis [letter]. *Lancet*, 1999. 353(9156): p. 901-902.
128. Bristol Myers Squibb Company. Healthcare Provider Important Drug Warning Letter. 5 January 2001.
129. Kakuda TN. Pharmacology of nucleoside and nucleotide reverse transcriptase inhibitor-induced mitochondrial toxicity. *Clin Ther*, 2000. 22(6): p. 685-708.
130. Brinkman K, Smeitink J, Romijn J, Reiss P. Mitochondrial toxicity induced by nucleoside-analogue reverse-transcriptase inhibitors is a key factor in the pathogenesis of antiretroviral therapy related lipodystrophy. *Lancet*, 1999. 354(9184): p. 1112-1115.
131. Kakuda T, Brundage R, Anderson P, et al. NRTI-induced mitochondrial toxicity as an aetiology for fat redistribution syndrome. *AIDS*, 1999. 13: p. 2311-2332.
132. Mallal S, John M, Moore C, et al. Contribution of nucleoside analogue reverse transcriptase inhibitors to subcutaneous fat wasting in patients with HIV infection. *AIDS*, 2000. 14(10): p. 1309-1316.
133. Lonergan JT, Behling C, Pfander H, et al. Hyperlactatemia and hepatic abnormalities in 10 human immunodeficiency virus-infected patients receiving nucleoside analogue combination regimens. *Clin Infect Dis*, 2000. 31(1): p. 162-166.
134. Bristol Myers Squibb Company. Healthcare Provider Important Drug Warning Letter. 28 September 2001.
135. Moore R, Keruly J, Chaisson R. Differences in anion gap with different nucleoside RTI combinations. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 55).
136. Brinkman K. Editorial response: Hyperlactatemia and hepatic steatosis as features of mitochondrial toxicity of nucleoside analogue reverse transcriptase inhibitors. *Clin Infect Dis*, 2000. 31(1): p. 167-169.
137. Toffaletti JG. Blood lactate: Biochemistry, laboratory methods, and clinical interpretation. *Crit Rev Clin Lab Sci*, 1991. 28(4): p. 253-268.
138. Miller KD, Cameron M, Wood LV, et al. Lactic acidosis and hepatic steatosis associated with use of stavudine: Report of four cases. *Ann Intern Med*, 2000. 133(3): p. 192-196.
139. Mokrzycki MH, Harris C, May H, et al. Lactic acidosis associated with stavudine administration: a report of five cases. *Clin Infect Dis*, 2000. 30(1): p. 198-200.
140. Chodock R, Mylonakis E, Shemin D, et al. Survival of a human immunodeficiency patient with nucleoside-induced lactic acidosis--role of haemodialysis treatment. *Nephrol Dial Transplant*, 1999. 14(10): p. 2484-2486.
141. Schramm C, Wanitschke R, Galle PR. Thiamine for the treatment of nucleoside analogue-induced severe lactic acidosis. *Eur J Anaesthesiol*, 1999. 16(10): p. 733-735.
142. denBrinker M, Wit FW, Wertheim-van Dillen PM, et al. Hepatitis B and C virus co-infection and the risk for hepatotoxicity of highly active antiretroviral therapy in HIV-1 infection. *AIDS*, 2000. 14(18): p. 2895-2902.
143. Martinez E, Bianco JL, Arnaiz JA, et al. Hepatotoxicity in HIV-1-infected patients receiving nevirapine-containing antiretroviral therapy. *AIDS*, 2001. 15(10): p. 1261-1268.
144. Barlett J. Severe liver toxicity in patients receiving two nucleoside analogues and a non-nucleoside reverse transcriptase inhibitor. In Program & Abstracts: 8th Conference on Retroviruses & Opportunistic Infections. Chicago, IL. Feb. 2001. (Abstract 19).
145. Boehringer-Ingelheim/Roxane Laboratory "Dear Health Care Professional Letter - Re: Severe, life-threatening and fatal cases of hepatotoxicity with VIRAMUNE". November 2000.
146. Sulkowski MS, Thomas DL, Chaisson RE, Moore RD. Hepatotoxicity associated with antiretroviral therapy in adults infected with human immunodeficiency virus and the role of hepatitis C or B virus infection. *JAMA*, 2000. 283(1): p. 74-80.
147. Saves M, Raffi F, Clevenbergh P, et al. and the APROCO Study Group. Hepatitis B or hepatitis C virus infection is a risk factor for severe hepatic cytolysis after initiation of a protease inhibitor-containing antiretroviral regimen in human immunodeficiency virus-infected patients. *Antimicrob Agents & Chemother*, 2000. 44(12): p. 3451-3455.
148. Nunez M, Lana R, Mendoza JL, et al. Risk factors for severe hepatic injury after introduction of highly active antiretroviral therapy. *J Acquir Immune Defic Syndr*, 2001. 27(5): p. 426-431.
149. Gisolf EH, Dreezen C, Danner SA, et al. for the Prometheus Study Group. Risk factors for hepatotoxicity in HIV-1-infected patients receiving ritonavir and saquinavir with or without stavudine. *Clin Infect Dis*, 2000. 31(5): p. 1234-1239.
150. Bonfanti P, Landonio S, Ricci E, et al. for the CISAI study group. Risk factors for hepatotoxicity in patients treated with highly active antiretroviral therapy. *J Acquir Immune Defic Syndr*, 2001. 27(3): p. 316-318.
151. Dube MP. Metabolic complications of antiretroviral therapies. *AIDS Clinical Care*, 1998. 10: p. 41.

152. Visnegarwala F, Krause KL, Musher DM. Severe diabetes associated with protease inhibitor therapy. *Ann Intern Med*, 1997. 127(10): p. 947.
153. Eastone JA, Decker CF. New-onset diabetes mellitus associated with use of protease inhibitor. *Ann Intern Med*, 1997. 127(10): p. 948.
154. Mulligan K, Grunfeld C, Tai VW, et al. Hyperlipidemia and insulin resistance are induced by protease inhibitors independent of changes in body composition in patients with HIV infection. *J Acquir Immune Defic Syndr*, 2000. 23(1): p. 35-43.
155. Behrens G, Dejam A, Schmidt H, et al. Impaired glucose tolerance, beta cell function and lipid metabolism in HIV patients under treatment with protease inhibitors. *AIDS*, 1999. 13(10): p. F63-70.
156. Tsiodras S, Mantzoros C, Hammer S, Samore M. Effects of protease inhibitors on hyperglycemia, hyperlipidemia, and lipodystrophy: a 5-year cohort study. *Arch Intern Med*, 2000. 160(13): p. 2050-2056.
157. Yarasheski K. 3rd International Workshop on Adverse Drug Reactions and Lipodystrophy in HIV Athens, Greece. Oct 23-26, 2001. (Abstract 6).
158. Murata H, Hruz P. 3rd International Workshop on Adverse Drug Reactions and Lipodystrophy in HIV. Athens, Greece. Oct 23-26, 2001. (Abstracts 1 and 2).
159. Miller KD, Jones E, Yanovski JA, et al. Visceral abdominal-fat accumulation associated with use of indinavir. *Lancet*, 1998. 351(9106): p. 871-875.
160. Lo JC, Mulligan K, Tai VW, et al. "Buffalo hump" in men with HIV-1 infection. *Lancet*, 1998. 351(9106): p. 867-870.
161. Herry I, Bernard L, de Truchis P, Perronne C. Hypertrophy of the breasts in a patient treated with indinavir. *Clin Infect Dis*, 1997. 25(4): p. 937-938.
162. Hengel RL, Watts NB, Lennox JL. Benign symmetric lipomatosis associated with protease inhibitors [letter]. *Lancet*, 1997. 350(9091): p. 1596.
163. Carr A, Samaras K, Thorisdottir A, et al. Diagnosis, prediction, and natural course of HIV-1 protease-inhibitor associated lipodystrophy, hyperlipidaemia, and diabetes mellitus: a cohort study. *Lancet*, 1999. 353(9170): p. 2093-2099.
164. Saint-Marc T, Partisani M, Poizot-Martin I, et al. A syndrome of peripheral fat wasting (lipodystrophy) in patients receiving long-term nucleoside analogue therapy. *AIDS*, 1999. 13(13): p. 1659-1667.
165. Carr A, Miller J, Law M, Cooper DA. A syndrome of lipoatrophy, lactic acidemia and liver dysfunction associated with HIV nucleoside analogue therapy: Contribution to protease inhibitor-related lipodystrophy syndrome. *AIDS*, 2000. 14(3): p. F25-32.
166. Kotler DP, Rosenbaum K, Wang J, Pierson RN. Studies of body composition and fat distribution in HIV-infected and control subjects. *J Acquir Immune Defic Syndr Hum Retrovirol*, 1999. 20(3): p. 228-237.
167. Martinez E, Conget I, Lozano L, et al. Reversion of metabolic abnormalities after switching from HIV-1 protease inhibitors to nevirapine. *AIDS*, 1999. 13(7): p. 805-810.
168. Barreiro P, Soriano V, Blanco F, et al. Risks and benefits of replacing protease inhibitors by nevirapine in HIV-infected subjects under long-term successful triple combination therapy. *AIDS*, 2000. 14(7): p. 807-812.
169. Roubenoff R, Weiss L, McDermott A, et al. A pilot study of exercise training to reduce trunk fat in adults with HIV-associated fat redistribution. *AIDS*, 1999. 13(11): p. 1373-1375.
170. Periard D, Telenti A, Sudre P, et al. Atherogenic dyslipidemia in HIV-infected individuals treated with protease inhibitors. The Swiss HIV Cohort Study. *Circulation*, 1999. 100(7): p. 700-705.
171. Carr A, Samaras K, Chisholm DJ, Cooper DA. Pathogenesis of HIV-1-protease inhibitor-associated peripheral lipodystrophy, hyperlipidaemia, and insulin resistance. *Lancet*, 1998. 351(9119): p. 1881-1883.
172. Klein D, Hurley L, Sidney S. Do protease inhibitors increase the risk for coronary heart disease among HIV-positive patients? Additional follow-up. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 33).
173. Coplan P, Cormier K, Japour A, et al. Myocardial infarction incidence in clinical trials of 4 protease inhibitors. 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 34).
174. Summary of the second report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel II). *J Am Med Assoc*, 1993. 269(23): p. 3015-3023.
175. Racoosin JA, Kessler CM. Bleeding episodes in HIV-positive patients taking HIV protease inhibitors: A case series. *Haemophilia*, 1999. 5(4): p. 266-269.
176. Scribner AN, T.-C.P., Cox BA, et al. Osteonecrosis in HIV: A case-control study. *J Acquir Immune Defic Syndr Hum Retrovirol*, 2000. 25: p. p. 19-25.
177. Gaughan DM, M.L., Hughes MD, et al. Avascular necrosis of the hip (Legg-Calve-Perthes Disease [LCPD]) in HIV-infected children in long-term follow up PACTG Protocol 219. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 686).
178. Mora S, Sala N, Bricalli D, et al. Bone mineral loss through increased bone turnover in HIV-infected children treated with highly active antiretroviral therapy. *AIDS*, 2001. 15(14): p. 1823-1829.
179. Masur H, Miller KD, Jones EC, et al. High prevalence of avascular necrosis of the hip in HIV

- infection. Program and abstracts of the 38th Annual Meeting of the Infectious Diseases Society of America. New Orleans, Louisiana. September 7-10, 2000. (Abstract 15).
180. Glesby M, Caamonde C. Case-control study of avascular necrosis in HIV-infected patients. *Antiviral Therapy*, 2000. 5(Suppl 5): p. 21.
 181. Paton NI, Macallan DC, Griffin GE, Pazianas M. Bone mineral density in patients with human immunodeficiency virus infection. *Calcif Tissue Int*, 1997. 61(1): p. 30-32.
 182. Aukrust P, Haug CJ, Ueland T, et al. Decreased bone formative and enhanced resorptive markers in human immunodeficiency virus infection: Indication of normalization of the bone-remodeling process during highly active antiretroviral therapy. *J Clin Endocrinol Metab*, 1999. 84(1): p. 145-150.
 183. Tebas P, Powderly WG, Claxton S, et al. Accelerated bone mineral loss in HIV-infected patients receiving potent antiretroviral therapy. *AIDS*, 2000. 14(4): p. F63-67.
 184. Hoy J, Hudson J, Law M, Cooper DA. Osteopenia in a randomized, multicenter study of protease inhibitor (PI) substitution in patients with the lipodystrophy syndrome and well-controlled HIV viremia. Program and abstracts of the 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA. Jan 30-Feb 2, 2000. (Abstract 208).
 185. Cunney RJ, Calligan D, Skehan S, et al. Reduced bone mineral density of HIV-positive men: Association with protease-inhibitor therapy and low serum testosterone. Program and abstracts of the 37th Annual Meeting of the Infectious Disease Society of America. Philadelphia, PA. Nov. 18-20, 1999. (Abstract 331).
 186. Tebas P, Yarasheski K, Whyte M, et al. Serum and urine markers of bone mineral metabolism in HIV-infected patients taking protease inhibitor containing potent antiretroviral therapy. *Antiviral Therapy*, 2000. 5 (Suppl 5):21: p. 19-19. (Abstract 029).
 187. Dunlop MB, Lane NE. Osteoporosis: Diagnosis, prevention and treatment of established disease. *Bull Rheum Di*, 1999. 48(6): p. 1-4.
 188. Fagot JP, Mockenhaupt M, Bouwes-Bavinck JN, et al. for the EuroSCAR study group. Nevirapine and the risk of Stevens-Johnson syndrome or toxic epidermal necrolysis. *AIDS*, 2001. 15(14): p. 1843-1848.
 189. Bourezane Y, Salard D, Hoen B, et al. DRESS (drug rash with eosinophilia and systemic symptoms) syndrome associated with nevirapine therapy. *Clin Infect Dis*, 1998. 27(5): p. 1321-1322.
 190. Bossi P, Colin D, Bricaire F, Caumes E. Hypersensitivity syndrome associated with efavirenz therapy. *Clin Infect Dis*, 2000. 30(1): p. 227-228.
 191. Bersoff-Matcha SJ, Miller WC, Aberg JA, et al. Sex difference in nevirapine rash. *Clin Infect Dis*, 2001. 32(1): p. 124-129.
 192. Antinori A, Baldini F, Girardi E, et al. Female sex and the use of anti-allergic agents increase the risk of developing cutaneous rash associated with nevirapine therapy. *AIDS*, 2001. 15(12): p. 1579-1581.
 193. Knobel H, Miro JM, De-Miguel V, et al. Efficacy of a short term prednisone regimen in nevirapine associated rash prevention: A double-blind, placebo-controlled clinical trial. Results of the GESIDA 09/99 study. In Program & Abstracts of 40th Interscience Conference on Antimicrobial Agents and Chemotherapy. Toronto. Sept. 2000. (Abstract L-15).
 194. Soriano V, Dona C, Barreiro P, Gonzalez-Lahoz J. Is there cross-toxicity between nevirapine and efavirenz in subjects developing rash? *AIDS*, 2000. 14(11): p. 1672-1673.
 195. Clarke S, Harrington P, Barry M, Mulcahy F. The tolerability of efavirenz after nevirapine-related adverse events. *Clin Infect Dis*, 2000. 31(3): p. 506-507.
 196. Deeks SG, Wrin T, Liegler T, et al. Virologic and immunologic consequences of discontinuing combination antiretroviral-drug therapy in HIV-infected patients with detectable viremia. *N Engl J Med*, 2001. 344(7): p. 472-480.
 197. Rosenberg ES, Altfeld M, Poon SH, et al. Immune control of HIV-1 after early treatment of acute infection. *Nature*, 2000. 407(6803): p. 523-526.
 198. Fagard C, Lebraz M, Gunthard H, et al. SSITT: A prospective trial of strategic treatment interruptions in 128 patients. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 357).
 199. Dybul M, Yoder C, Hidalgo B, et al. A randomized controlled trial on intermittent versus continuous HAART. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 357).
 200. Tindall B, Cooper DA. Primary HIV infection: Host responses and intervention strategies. *AIDS*, 1991. 5(1): p. 1-14.
 201. Niu MT, Stein DS, Schnittman SM. Primary human immunodeficiency virus type 1 infection: review of pathogenesis and early treatment intervention in humans and animal retrovirus infections. *J Infect Dis*, 1993. 168(6): p. 1490-1501.
 202. Kinloch-De Loes S, de Saussure P, Saurat JH, et al. Symptomatic primary infection due to human immunodeficiency virus type 1: Review of 31 cases. *Clin Infect Dis*, 1993. 17(1): p. 59-65.
 203. Schacker T, Collier AC, Hughes J, et al. Clinical and epidemiologic features of primary HIV infection. *Ann Intern Med*, 1996. 125(4): p. 257-264.
 204. Lafeuillade A, Poggi C, Tamalet C, et al. Effects of a combination of zidovudine, didanosine, and lamivudine on primary human immunodeficiency

- virus type 1 infection. *J Infect Dis*, 1997. 175(5): p. 1051-1055.
205. Lillo FB, Ciuffreda D, Veglia F, et al. Viral load and burden modification following early antiretroviral therapy of primary HIV-1 infection. *AIDS*, 1999. 13(7): p. 791-796.
 206. Malhotra U, Berrey MM, Huang Y, et al. Effect of combination antiretroviral therapy on T-cell immunity in acute human immunodeficiency virus type 1 infection. *J Infect Dis*, 2000. 181(1): p. 121-131.
 207. Pantaleo G, Cohen OJ, Schacker T, et al. Evolutionary pattern of human immunodeficiency virus (HIV) replication and distribution in lymph nodes following primary infection: Implications for antiviral therapy. *Nat Med*, 1998. 4(3): p. 341-345.
 208. Update: provisional Public Health Service recommendations for chemoprophylaxis after occupational exposure to HIV. *MMWR*, 1996. 45(22): p. 468-480.
 209. Centers for Disease Control and Prevention, Perinatal HIV Guidelines Working Group., Public Health Service Task Force recommendations for the use of antiretroviral drugs in pregnant women infected with HIV-1 for maternal health and for reducing perinatal HIV-1 transmission in the United States. *MMWR*, 1998. 47(1-30). For updated guidelines, See HIV/ATIS Treatment Information Service website <http://www.hivatis.org>.
 210. Ioannidis JPA, Abrams EJ, Ammann A, et al. Perinatal transmission of human immunodeficiency virus type 1 by pregnant women with RNA virus loads <1000 copies/ml. *J Infect Dis*, 2001. 183(4): p. 539-545.
 211. Mofenson LM, Lambert JS, Stiehm ER, et al. Risk factors for perinatal transmission of human immunodeficiency virus type 1 in women treated with zidovudine. Pediatric AIDS Clinical Trials Group Study 185 Team. *N Engl J Med*, 1999. 341(6): p. 385-393.
 212. Garcia PM, Kalish LA, Pitt J, et al. Maternal levels of plasma human immunodeficiency virus type 1 RNA and the risk of perinatal transmission. Women and Infants Transmission Study Group. *N Engl J Med*, 1999. 341(6): p. 394-402.
 213. Minkoff H, Augenbraun M. Antiretroviral therapy for pregnant women. *Am J Obstet Gynecol*, 1997. 176(2): p. 478-489.
 214. Pregnancy outcomes following systemic prenatal acyclovir exposure--June 1, 1984-June 30, 1993. *MMWR*, 1993. 42(41): p. 806-809.
 215. Ibdah JA, Yang Z, Bennett MJ. Liver disease in pregnancy and fetal fatty acid oxidation defects. *Mol Genet Metab*, 2000. 71(1-2): p. 182-189.
 216. Ibdah JA, Bennett MJ, Rinaldo P, et al. A fetal fatty-acid oxidation disorder as a cause of liver disease in pregnant women. *N Engl J Med*, 1999. 340(22): p. 1723-1731.
 217. Bristol-Myers Squibb Company. Healthcare provider important drug warning letter. January 5, 2001.
 218. Rodman JH, Robbins B, Flynn PM, Fridland A. A systemic and cellular model for zidovudine plasma concentrations and intracellular phosphorylation in patients. *J Infect Dis*, 1996. 174(3): p. 490-499.
 219. Barry MG, Khoo SH, Veal GJ, et al. The effect of zidovudine dose on the formation of intracellular phosphorylated metabolites. *AIDS*, 1996. 10(12): p. 1361-1367.
 220. Mulder JW, Cooper DA, Mathiesen L, et al. Zidovudine twice daily in asymptomatic subjects with HIV infection and a high risk of progression to AIDS: A randomized, double-blind, placebo-controlled study. The European-Australian Collaborative Group (Study 017). *AIDS*, 1994. 8(3): p. 313-321.
 221. Mannucci PM, Gringeri A, Savidge G, et al. Randomized double-blind, placebo-controlled trial of twice-daily zidovudine in asymptomatic haemophiliacs infected with the human immunodeficiency virus type 1. European-Australian Haemophilia Collaborative Study Group. *Br J Haematol*, 1994. 86(1): p. 174-179.
 222. Cooper DA, Gatell JM, Kroon S, et al. Zidovudine in persons with asymptomatic HIV infection and CD4+ cell counts greater than 400 per cubic millimeter. The European-Australian Collaborative Group. *N Engl J Med*, 1993. 329(5): p. 297-303.
 223. Administration of zidovudine during late pregnancy and delivery to prevent perinatal HIV transmission--Thailand, 1996-1998. *MMWR*, 1998. 47(8): p. 151-154.
 224. Kourtis AP, Bulterys M, Nesheim SR, Lee FK. Understanding the Timing of HIV Transmission From Mother to Infant. *JAMA*, 2001. 285(6): p. 709-711. <http://jama.ama-assn.org/issues/v285n6/full>.
 225. Iversen AK, Larsen AR, Jensen T, et al. Distinct determinants of human immunodeficiency virus type 1 RNA and DNA loads in vaginal and cervical secretions. *J Infect Dis*, 1998. 177(5): p. 1214-1220.
 226. Hart CE, Lennox JL, Pratt-Palmore M, et al. Correlation of human immunodeficiency virus type 1 RNA levels in blood and the female genital tract. *J Infect Dis*, 1999. 179(4): p. 871-882.
 227. Shaheen F, Sison AV, McIntosh L, et al. Analysis of HIV-1 in the cervicovaginal secretions and blood of pregnant and nonpregnant women. *J Hum Virol*, 1999. 2(3): p. 154-166.
 228. Overbaugh J, Anderson RJ, Ndinya-Achola JO, Kreiss JK. Distinct but related human immunodeficiency virus type 1 variant populations in genital secretions and blood. *AIDS Res Hum Retroviruses*, 1996. 12(2): p. 107-115.
 229. Libman H, Makadon HJ, Ed. *HIV*. Philadelphia, PA: American College of Physicians. 2000.
 230. Quinn TC, Wawer M, Sewankambo N, et al. Viral load and heterosexual transmission of human

- immunodeficiency virus Type 1. *N Engl J Med*, 2000. 342(13): p. 921-929.
231. Kovacs A, Wasserman SS, Burns D, et al. Determinants of HIV-1 shedding in the genital tract of women. *Lancet*, 2001. 358(9293): p. 1593-1601.
232. The European Collaborative Study. Maternal viral load and vertical transmission of HIV-1: an important factor but not the only one. *AIDS*, 1999. 13(11): p. 1377-1385.
233. Wabwire-Mangen F, Gray RH, Kigozi G, et al. Randomized trial of STD control for maternal-infant health: Rakai, Uganda. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract MoOrC242).
234. Hoffman I, Taha TE, Martinson F, et al. Adverse health events occurring during an N-9 gel pilot study: Malawi. 13th International AIDS Conference, Durban, South Africa. 2000. (Abstract TuPpC1171).
235. Bailey R. Male circumcision as an effective HIV prevention strategy: current evidence. 8th Conference on Retroviruses and Opportunistic Infections, Chicago, IL. 2001. (Abstract S22).
236. Marx P, Smith S, Baskin G. Estrogen protects against vaginal transmission of simian immunodeficiency virus. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract MoOrA227).
237. Gray R, Wawer MJ, Sewakambo NK, et al. HIV incidence associated with male circumcision and viral load in discordant couples: Rakai, Uganda. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract MoOrC193).
238. Bukusi E, C.C., Karanja J, et al. Bacterial vaginosis: evidence for sexual transmission and association with HIV-1 infection. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract MoOrA229).
239. Baeten J, Mostad S, Hughes M, et al. Selenium deficiency is associated with shedding of HIV-1 infected cells in the female genital tract. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract MoOrA226).
240. Buve A, Auvert B, Lagarde E, et al. Male circumcision and HIV spread in sub-Saharan Africa. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract MoOrC192).
241. Valleroy L, MacKellar D, Karon J, et al. HIV prevalence and associated risks in young men who have sex with men. *JAMA*, 2000. 284(2): p. 198-204.
242. Bingham T, Johnson D, Harawa N. Confidence in HAART and recent unprotected anal sex among men who have sex with men (MSM). 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 213).
243. Hogg R, Yip B, Chan K, et al. Non-adherence to triple combination therapy is predictive of AIDS progression and death in HIV-positive men and women. 13th International AIDS Conference. Durban, South Africa. 2000. (Abstract TuOrB419).
244. Flaks R, Burman P, Gourley C, et al. Unsafe sex among persons in HIV care is associated with decreased adherence with antiretroviral therapy. 8th Conference on Retroviruses and Opportunistic Infections. Chicago, IL. 2001. (Abstract 214).

Considerations for Antiretroviral Therapy in Women

Several studies have suggested that plasma HIV RNA levels are significantly lower in adult women compared to men. Several analyses have been reported from the ALIVE cohort of intravenous drug users in Baltimore. In a cross-sectional study from this cohort, there was a consistent trend toward lower viral load (quantitative microculture as well as HIV RNA measured by branched chain DNA and RT-PCR) in women compared to men after adjustment for CD4⁺ lymphocyte count, race and drug use within the prior 6 months; the difference in RNA levels was approximately 0.25 log [1]. When women and men were matched for CD4⁺ T cell count there was no difference in the risk for progression to AIDS. However, when matched for RNA copy number, the risk of AIDS was 1.6-fold higher for women. In a further longitudinal case-control evaluation of seroconverters from this cohort, the sex-specific difference in viral load was present at seroconversion, but viral load tended to increase more rapidly in women and median viral loads in women and men became similar within 5-6 years of seroconversion [2]. The relationship between initial HIV RNA level at seroconversion and progression to AIDS was examined in a longitudinal study of 202 seroconverters (156 men and 46 women) from this cohort [3]. HIV RNA levels following seroconversion were significantly lower in women than men (by approximately 0.5 log), but these differences became attenuated over time. There was no significant sex-specific difference in rates of progression to AIDS. In another longitudinal study of 14 women and 28 men in the armed forces, median RNA levels were lower in women, but these differences were less than 0.5 log and diminished over time; no differences in HIV DNA load were observed [4]. In a virology substudy of ACTG 175, cross-sectional HIV RNA levels were 0.28 log lower in 71 women at baseline compared with men after adjustment for CD4⁺ T cell count [5].

Other large cohort studies have had less convincing results. In 647 women from the Swiss HIV Cohort Study, there was a slightly lower viral load among female injection drug users (0.13 log) but not among heterosexually infected women [6]. Additionally, there was no difference in disease progression between women and men matched for HIV RNA level and CD4⁺ T cell count. In 712 women in the ICONA study, viral load was only 0.13 log lower in women after adjustment for CD4⁺ T cell count; however, in contrast to the Swiss HIV Cohort Study, the sex-specific difference was larger in women with heterosexually

acquired HIV infection compared with injection drug use-acquired HIV infection [7]. Data reported from Johns Hopkins showed little evidence of lower viral load after stratification by CD4⁺ T cell count [8], and in a comparison of 1262 women from the Women's Interagency HIV Study and men from the Multicenter AIDS Cohort Study, a small viral load difference of ~0.10-0.14 log was present only at higher CD4 count levels [9]. Finally, in an analysis of adults with advanced transfusion-acquired HIV infection, no significant differences in HIV RNA levels between women and men were observed [10] and no difference in viral load by sex was observed for age and CD4⁺ T cell-matched antiretroviral naïve men and women either before or after antiretroviral therapy [11].

Limited studies in HIV-infected adults have indicated that women may have higher CD4⁺ T cell count than men. In a French study, this difference was observed only for CD4 percentage and was of borderline significance for CD4 absolute number once women and men were matched for age [12]. In a second European study, while absolute CD4⁺ T cell count was higher in women than men, these differences were only statistically significant at AIDS onset and not at seroconversion or death [13]. Neither study evaluated the relationship of sex and CD4⁺ T count to disease progression. However, other studies have shown similar rates of disease progression between men and women matched for CD4⁺ T cell count and/or HIV RNA level [6, 14, 15].

Taken together, these data suggest that gender-based differences in viral load occur predominantly during a window of time when the CD4⁺ T cell count is relatively preserved and treatment is recommended only in the setting of high levels of plasma HIV RNA. Clinicians may wish to consider lower plasma HIV RNA thresholds for initiating therapy in women with CD4⁺ T cell counts >350 cells/mm³, although there are insufficient data to determine an appropriate threshold. In patients with CD4⁺ T cell counts <350 cells/mm³, very small sex-based differences in viral load are apparent; therefore, no changes in treatment guidelines for women are recommended for this group.

Further study is warranted regarding sex differences in viral and immunologic parameters. It is likely that any such differences would be hormonally related; estrogen-related effects have been described on immune function [16]. Consistent with this hypothesis are some preliminary studies of variation in viral load

according to menstrual cycle. One study has suggested that the ovulatory cycle influences circulating HIV-1 RNA levels [17]. Additionally, another study suggests that pharmacokinetic parameters may vary over the ovulatory cycle; considerable variations in indinavir pharmacokinetics were found during the menstrual cycle, with a trend to more drug exposure during the follicular phase [18].

References

1. Farzadegan H, Hoover DR, Astemborski J, et al. Sex differences in HIV-1 viral load and progression to AIDS. *Lancet*, 1998. 352(9139): p. 1510-4.
2. Sterling TR, Lyles CM, Vlahov D, et al. Sex differences in longitudinal human immunodeficiency virus type 1 RNA levels among seroconverters. *J Infect Dis*, 1999. 180(3): p. 666-72.
3. Sterling TR, Vlahov D, Astemborski J, et al. Initial Plasma HIV-1 RNA levels and progression to AIDS in women and men. *N Engl J Med*, 2001. 344(10): p. 720-5.
4. Evans JS, Nims T, Cooley J, et al. Serum levels of virus burden in early-stage human immunodeficiency virus type 1 disease in women. *J Infect Dis*, 1997. 175(4): p. 795-800.
5. Katzenstein DA, Hammer SM, Hughes MD, et al. The relation of virologic and immunologic markers to clinical outcomes after nucleoside therapy in HIV-infected adults with 200 to 500 CD4 cells per cubic millimeter. AIDS Clinical Trials Group Study 175 Virology Study Team. *N Engl J Med*, 1996. 335(15): p. 1091-8.
6. Junghans C, Ledergerber B, Chan P, et al. Sex differences in HIV-1 viral load and progression to AIDS. Swiss HIV Cohort Study. *Lancet*, 1999. 353(9152): p. 589.
7. Moroni M. Sex differences in HIV-1 viral load and progression to AIDS. ICONA Study Group. Italian cohort of HIV-1 positive individuals. *Lancet*, 1999. 353(9152): p. 589-90.
8. Moore RD, Cheever L, Keruly JC, Chaisson RE. Lack of sex difference in CD4 to HIV-1 RNA viral load ratio [letter]. *Lancet*, 1999. 353(9151): p. 463-4.
9. Anastos K, Gange SJ, Lau B, et al. Gender specific differences in quantitative HIV-1 RNA levels. 6th Conference on Retroviruses and Opportunistic Infections. Chicago, IL, 1999. (Abstract 274).
10. Kalish LA, Collier AC, Flanigan TP, Kumar PN. Plasma HIV RNA viral load in women and men with advanced HIV disease. 39th Interscience Conference on Antimicrobial Agents and Chemotherapy. San Francisco, CA, 1999. (Abstract 2190).
11. Bush CE, Donovan RM, Markowitz N, et al. Gender is not a factor in serum human immunodeficiency virus type 1 RNA levels in patients with viremia. *J Clin Microbiol*, 1996. 34(4): p. 970-2.
12. Delmas MC, Jadand C, De Vincenzi I, et al. Gender difference in CD4+ cell counts persist after HIV-1 infection. SEROCO Study Group [letter]. *AIDS*, 1997. 11(8): p. 1071-3.
13. Prins M, Robertson JR, Brettelle RP, et al. Do gender differences in CD4 cell counts matter? *AIDS*, 1999. 13(17): p. 2361-4.
14. Webber MP, Schoenbaum EE, Gourevitch MN, et al. A prospective study of HIV disease progression in female and male drug users. *AIDS*, 1999. 13(2): p. 257-62.
15. Blair J, Hanson D, Jones J, et al. Do gender differences in viral load predict differences in HIV disease progression? 7th Conference on Retroviruses and Opportunistic Infections. San Francisco, CA, 2000. (Abstract 195).
16. Styrt B, Sugarman B. Estrogens and infection. *Rev Infect Dis*, 1991. 13(6): p. 1139-50.
17. Greenblatt RM, Ameli N, Grant RM, et al. Impact of the ovulatory cycle on virologic and immunologic markers in HIV-infected women. *J Infect Dis*, 2000. 181(1): p. 82-90.
18. Adams J, Frost C, Shelton M. et al. Indinavir pharmacokinetics and menstrual cycle physiology. 5th Conference on Retroviruses and Opportunistic Infections. Chicago, IL, 1998. (Abstract 356).

Hydroxyurea

Hydroxyurea is indicated for use in the treatment of certain malignancies and in sickle cell anemia, and has been used investigationally for the treatment of HIV. Its potential safety and effectiveness for treatment of HIV have not been established, and clinicians should be aware of important safety precautions regarding its use. Hydroxyurea does not have direct antiretroviral activity; rather, it inhibits the cellular enzyme ribonucleotide reductase, resulting in reduced intracellular levels of deoxynucleoside triphosphates (dNTPs) that are necessary for DNA synthesis.

Depletion of the dNTP pool results in arrest of the cell cycle in the G1 phase prior to DNA synthesis; in an HIV-infected cell, incomplete reverse transcription of the viral genome also results from depletion of the dNTP pool [1]. Hydroxyurea preferentially depletes intracellular dATP; therefore, it has been hypothesized that the antiretroviral activity of ddI and d4T may be enhanced in combination with hydroxyurea.

Hydroxyurea also induces the activity of cellular kinases that phosphorylate nucleoside analogue reverse transcriptase inhibitors, potentially further enhancing their antiretroviral activity.

Few data are available from controlled clinical trials that provide support for the clinical utility of hydroxyurea as an adjunct in the treatment of HIV infection. In limited studies, the addition of hydroxyurea to a regimen of ddI +d4T or ddI alone appeared to result in moderately enhanced antiretroviral activity [2-4], although the optimal dosage and dosing schedule were not determined. In contrast, in ACTG 5025, a randomized, controlled clinical trial conducted in subjects on potent antiretroviral therapy with levels of plasma viremia <200 copies/mL [5], no statistically significant differences in viral load suppression were observed in patients receiving hydroxyurea 600 mg twice daily in combination with ddI+d4T+indinavir compared to those receiving the combination regimen without hydroxyurea. Importantly, this trial was prematurely closed due to higher rates of drug toxicity in patients randomized to the hydroxyurea-containing arm. Among 68 patients randomized to hydroxyurea, three deaths related to complications of pancreatitis were reported, and a substantial decrease in median CD4⁺T cell count was observed in the hydroxyurea treatment group. The increased frequency of fatal pancreatitis in the hydroxyurea-containing arm was not statistically

significant and had not been reported previously.

These cases of fatal pancreatitis do, however, raise the question of whether hydroxyurea in combination with ddI+d4T may increase the risk of ddI-associated pancreatitis. Additional concerns regarding the use of hydroxyurea in HIV infection have been raised in this trial and other studies, and include an increased risk of persistent cytopenias [6] and hepatotoxicity [7], the drug's teratogenic properties, and the possibility of an increased risk of neuropathy. Given these concerns, more data on the potential safety and efficacy of lower doses of hydroxyurea are necessary to determine if hydroxyurea in combination with antiretroviral agents has a therapeutic role for HIV infection. Clinicians considering the use of hydroxyurea in a treatment regimen for HIV should be aware of the limited and conflicting nature of data in support of its efficacy, and the importance of monitoring patients closely for potentially serious toxicity (DII).

References

1. Rutschmann OT, Opravil M, Iten A, et al. A placebo-controlled trial of didanosine plus stavudine, with and without hydroxyurea, for HIV infection. The Swiss HIV Cohort Study. *AIDS*, 1998. 12(8): p. F71-7.
2. Montaner JS, Zala C, Conway B, et al. A pilot study of hydroxyurea among patients with advanced human immunodeficiency virus (HIV) disease receiving chronic didanosine therapy: Canadian HIV trials network protocol 080. *J Infect Dis*, 1997. 175(4): p. 801-6.
3. Gao WY, Cara A, Gallo RC, Lori F. Low levels of deoxynucleotides in peripheral blood lymphocytes: A strategy to inhibit human immunodeficiency virus type 1 replication. *Proc Natl Acad Sci U S A*, 1993. 90(19): p. 8925-8.
4. Federici ME, Lupo S, Cahn P, et al. Hydroxyurea in combination regimens for the treatment of antiretroviral-naive HIV-infected adults. 12th World AIDS Conference. Geneva, 1998. (Abstract 12235).
5. ACTG 5025 Team, *Closure of ACTG 5025*. 1999.
6. Goodrich J, Khardori N. Hydroxyurea toxicity in human immunodeficiency virus-positive patients. *Clin Infect Dis*, 1999. 29(3): p. 692-3.
7. Weissman SB, Sinclair GI, Green CL, Fissell WH. Hydroxyurea-induced hepatitis in human immunodeficiency virus-positive patients. *Clin Infect Dis*, 1999. 29(1): p. 223-4.

Safety and Toxicity of Individual Antiretroviral Agents in Pregnancy

NUCLEOSIDE & NUCLEOTIDE ANALOGUE REVERSE TRANSCRIPTASE INHIBITORS

There are currently six approved nucleoside analogue reverse transcriptase inhibitors. Data are available from clinical trials in human pregnancy for zidovudine and lamivudine, while didanosine and stavudine are under study. Zalcitabine and abacavir have not been studied in pregnant women. Tenofovir disoproxil fumarate is the first acyclic nucleotide analogue reverse transcriptase inhibitor. The nucleoside analogue drugs require three intracellular phosphorylation steps to form the triphosphate nucleoside, which is the active drug moiety; tenofovir, an acyclic nucleotide analogue drug, contains a monophosphate component attached to the adenine base, and hence only requires two phosphorylation steps to form the active moiety.

Abacavir (ZIAGEN[®], ABC) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies
Long-term animal carcinogenicity studies of abacavir in rodents are not completed; however, some *in vitro* mutagenesis and clastogenesis screening tests are positive.
- Reproduction/fertility animal studies
No effect of abacavir on reproduction or fertility in male and female rodents has been seen at doses of up to 500 mg/kg/day (about 8 times that of human therapeutic exposure).
- Teratogenicity/developmental toxicity animal studies
Abacavir is associated with developmental toxicity (decreased fetal body weight and reduced crown-rump length) and increased incidence of fetal anasarca and skeletal malformations in rats treated with abacavir during organogenesis at doses of 1000 mg/kg (about 35 times that of human therapeutic exposure based on area under the curve (AUC)). Toxicity to the developing embryo and fetus (increased resorptions and decreased fetal body weight) occurred with abacavir administration to pregnant rodents at 500 mg/kg/day. The offspring of female rats treated with 500 mg/kg of abacavir beginning at embryo implantation and ending at weaning had an increased incidence of stillbirth and lower body weight throughout life.

However, in the rabbit, no evidence of drug-related developmental toxicity was observed and no increase in fetal malformations was observed at doses up to 700 mg/kg (about 8.5 times that of human therapeutic exposure).

- Placental and breast milk passage in animal studies
Abacavir crosses the placenta and is excreted into the breast milk of lactating rats.
- Human studies in pregnancy
No studies have been conducted with abacavir in pregnant women or neonates. Serious hypersensitivity reactions have been associated with abacavir therapy in non-pregnant adults and have rarely been fatal; symptoms include fever, skin rash, fatigue, and gastrointestinal symptoms such as nausea, vomiting, diarrhea, or abdominal pain. Abacavir should not be restarted following a hypersensitivity reaction because more severe symptoms will recur within hours and may include life-threatening hypotension and death.

Didanosine (Videx[®], ddl) is classified as FDA pregnancy category B.

- Animal carcinogenicity studies
Long-term animal carcinogenicity screening studies in rodents given didanosine have been negative.
- Reproduction/fertility animal studies
There has been no effect of didanosine on reproduction or fertility in rodents or on preimplantation mouse embryos [1].
- Teratogenicity/developmental toxicity animal studies
No evidence of teratogenicity or toxicity was observed with administration of high doses of didanosine to pregnant rats, mice, or rabbits.
- Placental and breast milk passage in humans
Placental transfer of didanosine was limited in a phase I/II safety and pharmacokinetic study (cord-to-maternal blood ratio, 0.35–0.11) [2]. Didanosine is excreted in the milk of lactating rats; it is not known if didanosine is excreted in human breast milk.

- Human studies in pregnancy

A phase I study (PACTG 249) of didanosine was conducted in 14 HIV-infected pregnant women enrolled at gestational age 26 to 36 weeks and treated through 6 weeks postpartum [2]. The drug was well-tolerated during pregnancy by the women and the fetuses. Preliminary analyses indicate that pharmacokinetic parameters after oral administration were not significantly affected by pregnancy, and that dose modification from the usual adult dosage is not needed.

Lamivudine (EPIVIR[®], 3TC) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies

Long-term animal carcinogenicity screening studies in rodents administered lamivudine have been negative.

- Reproduction/fertility animal studies

There appears to be no effect of lamivudine on reproduction or fertility in rodents.

- Teratogenicity/developmental toxicity animal studies

There is no evidence of lamivudine-induced teratogenicity. Early embryoletality was seen in rabbits but not rats at doses similar to human therapeutic exposure.

- Placental and breast milk passage in humans

Lamivudine readily crosses the placenta in humans, achieving comparable cord blood and maternal concentrations [3]. Lamivudine is excreted into human breast milk.

- Human studies in pregnancy

A small phase I study in South Africa evaluated the safety and pharmacokinetics of lamivudine alone or in combination with zidovudine in 20 HIV-infected pregnant women; therapy was started at 38 weeks gestation, continued through labor, and given for 1 week following birth to the infants [3]. The drug was well-tolerated in the women at the recommended adult dose of 150 mg orally twice daily; pharmacokinetics were similar to those observed in nonpregnant adults, and no pharmacokinetic interaction with zidovudine was observed.

Zidovudine and lamivudine, given in combination orally intrapartum, were well-tolerated. Lamivudine was well-tolerated in the neonates, but clearance was about 50% that of older children, requiring a reduced dosing regimen (4 mg/kg/day in neonates compared to

8 mg/kg/day for infants older than 3 months). There are currently no data on the pharmacokinetics of lamivudine between 2 to 6 weeks of age, and the exact age at which lamivudine clearance begins to approximate that in older children is not known.

Stavudine (Zerit[®], d4T) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies

Long-term animal carcinogenicity studies of stavudine in rodents are not completed; some *in vitro* and *in vivo* mutagenesis and clastogenicity tests are positive.

- Reproduction/fertility animal studies

No effect of stavudine on reproduction or fertility in rodents has been seen. A dose-related cytotoxic effect on preimplantation mouse embryos, with inhibition of blastocyst formation at a concentration of stavudine of 100 μ M and of postblastocyst development at 10 μ M [1].

- Teratogenicity/developmental toxicity animal studies

No evidence of teratogenicity of stavudine has been observed in pregnant rats and rabbits. Developmental toxicity, consisting of a small increase in neonatal mortality and minor skeletal ossification delay, occurred at the highest dose in rats.

- Placental and breast milk passage in animals

Stavudine crosses the rat placenta *in vivo* and the human placenta *ex vivo*, resulting in a fetal/maternal concentration of approximately 0.50. In primates (pigtailed macaques), fetal/maternal plasma concentrations were approximately 0.80 [4]. Stavudine is excreted into the breast milk of lactating rats.

- Human studies in pregnancy

A phase I/II safety and pharmacokinetic study of combination stavudine and lamivudine in pregnant HIV-infected women and their infants (PACTG 332) is being conducted, but data are not yet available. In primate studies, pregnancy did not affect the pharmacokinetics of stavudine [5].

Tenofovir disoproxil fumerate [DF] (Viread™) is classified as FDA pregnancy category B.

- Animal carcinogenicity studies
Long-term animal carcinogenicity studies of tenofovir DF in rodents are not completed; however, some *in vitro* mutagenesis and clastogenesis screening tests are positive.
- Reproduction/fertility animal studies
Reproductive toxicity has been evaluated in rats and rabbits. Tenofovir had no adverse effects on fertility or general reproductive performance in rats at doses up to 600 mg/kg/day (exposure equivalent to approximately 10 times the human dose based on body surface area comparisons). However, there was an alteration of the estrous cycle in female rats administered 600 mg/kg/day of tenofovir.
- Teratogenicity/developmental toxicity animal studies
No adverse effects on embryo/fetal development were seen when tenofovir was given in doses up to 450 mg/kg/day to pregnant rats and 300 mg/kg/day to pregnant rabbits. When tenofovir was administered to pregnant rats in doses of 450–600 mg/kg/day, which are maternally toxic doses, peri- and post-natal development studies of their offspring showed reduced survival and slight delay in sexual maturation. However, there were no adverse effects on growth, development, behavior, or reproductive parameters when tenofovir was administered to pregnant rodents at doses that were not associated with maternal toxicity (150 mg/kg/day). Chronic exposure of fetal monkeys to tenofovir at a high dose of 30 mg/kg (exposure equivalent to 25 times the AUC achieved with therapeutic dosing in humans) from days 20–150 of gestation did not result in gross structural abnormalities.[6] However, significantly lower fetal circulating insulin-like growth factor (IGF)-1 (a primary regulator of linear growth) and higher IGF binding protein (IGFBP)-3 levels were shown and were associated with overall body weights approximately 13% lower than untreated controls. A slight reduction in fetal bone porosity was also observed. Effects on these parameters were observed within 2 months of maternal treatment. Significant changes in maternal monkey bone biomarkers were noted but were primarily limited to the treatment period and were reversible.

Continued administration of tenofovir at 30 mg/kg/day to the infant monkey postnatally resulted in significant growth restriction and severe bone toxicity in 25% of eight infants and effects on bone biomarkers and

defective bone mineralization in all animals. Chronic administration of tenofovir to immature animals of multiple species has resulted in reversible bone abnormalities; these effects were dose-, exposure-, age-, and species-specific. Abnormalities ranged from minimal decrease in bone mineral density and content (with oral dosing in rats and dogs that achieved drug exposures 6 to 10 times that achieved with therapeutic dosing in humans) to severe, pathologic osteomalacia (with subcutaneous dosing given to monkeys). Juvenile monkeys given chronic subcutaneous tenofovir at 30 mg/kg/day (exposure equivalent to 25 times the AUC achieved with therapeutic dosing in humans) developed osteomalacia, bone fractures, and marked hypophosphatemia. However, no clinical or radiologic bone toxicity was seen when juvenile monkeys received subcutaneous dosing of 10 mg/kg/day (exposure equivalent to 8 times the AUC achieved with therapeutic dosing in humans). Evidence of nephrotoxicity was observed in newborn and juvenile monkeys given tenofovir in doses resulting in exposures 12 to 50 times higher than the human dose based on body surface area comparisons.

- Placental and breast milk passage in humans
Studies in rats have demonstrated that tenofovir is secreted in milk. Intravenous administration of tenofovir to pregnant cynomolgus monkeys resulted in a fetal/maternal concentration of 17%, demonstrating that tenofovir does cross the placenta [7]. There are no data on whether tenofovir crosses the placenta or is excreted in breast milk in humans.
- Human studies in pregnancy
No studies of tenofovir have been conducted in pregnant women or neonates.

Zalcitabine (HIVID®, ddC) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies
High doses of zalcitabine (over 1,000 times that of human therapeutic exposure) have been associated with the development of thymic lymphomas in rodents.
- Reproduction/fertility animal studies
No effect of zalcitabine on reproduction or fertility in rodents has been seen. However, there is a dose-related cytotoxic effect on preimplantation mouse embryos, with inhibition at a zalcitabine concentration of 100 µM; no inhibition of postblastocyst development was observed [1].

- Teratogenicity/developmental toxicity animal studies
Teratogenicity (hydrocephalus) occurred in rats given very high doses (over 1,000 times the maximally recommended human exposure) of zalcitabine. Developmental toxicity, consisting of decreased fetal weight and skeletal defects, has been seen in rodents at moderate to high zalcitabine doses. Cytotoxic effects were observed on rat fetal thymocytes at zalcitabine concentrations as low as 10 μ M (approximately 100 times human therapeutic exposure).
- Placental and breast milk passage in animal studies
In primate and placental perfusion studies, zalcitabine crosses the placenta (fetal-to-maternal drug ratio approximately 0.50 to 0.60) [8]. In rodents, zalcitabine concentrates in the fetal kidney and a relatively small proportion (approximately 20%) reaches the fetal brain. It is unknown if ddC is excreted in breast milk.
- Human studies in pregnancy
No studies of zalcitabine have been conducted in pregnant women or neonates.

Zidovudine (Retrovir[®]) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies
Prolonged, continuous, high-dose zidovudine administration to adult rodents is associated with the development of nonmetastasizing vaginal squamous tumors in 13% of female rodents (at estimated drug concentrations 3 and 24 times that of human therapeutic exposure in mice and rats, respectively) [9]. In rodents, unmetabolized zidovudine is concentrated in urine with reflux into the vaginal vault. Therefore, vaginal tumors could be a topical effect of chronic zidovudine exposure on the vaginal mucosa. The observation that vaginal squamous cell carcinomas were observed in rodents exposed to 20 mg/mL zidovudine intravaginally is consistent with this hypothesis [9]. In humans, only metabolized zidovudine is excreted in the urine. No increase in tumors in other organ sites has been seen in adult rodent studies.

Two transplacental carcinogenicity studies of zidovudine were conducted in mice, with differing results. In one study, two very high daily doses of zidovudine were administered during the last third of gestation in mice [10]. These doses were near the maximum dose beyond which lethal fetal toxicity would be observed and approximately 25 and 50 times

greater than the daily dose given to humans (although the cumulative dose was similar to the cumulative dose received by a pregnant woman taking 6 months of zidovudine). In the offspring of zidovudine-exposed pregnant mice at the highest dose level followed for 12 months, a statistically significant increase in lung, liver, and female reproductive organ tumors was observed; the investigators also documented incorporation of zidovudine into the DNA of a variety of newborn mouse tissues, although this did not clearly correlate with the presence of tumors. In the second study, pregnant mice were given one of several regimens of zidovudine, at doses intended to achieve blood levels approximately threefold higher than human therapeutic exposure [11]. The daily doses received by the mice during gestation ranged from one-twelfth to one-fiftieth the daily doses received in the previous study. Some of the offspring also received zidovudine for varying periods of time over their lifespan. No increase in the incidence of tumors was observed in the offspring of these mice, except among those that received additional lifetime zidovudine exposure, in which vaginal tumors were again noted.

Transplacental carcinogenicity studies have not been performed for any of the other available antiretroviral drugs or combinations of drugs. In January 1997, the National Institutes of Health convened an expert panel to review these animal data [12]. The panel concluded that the known benefit of zidovudine in reducing vertical transmission of HIV by nearly 70% (7.2 versus 21.9% with placebo) [13] far outweighs the theoretical risks of transplacental carcinogenicity. The panel also concluded that infants with *in utero* exposure to zidovudine (or any other antiretroviral) should have long-term followup for potential adverse effects. No tumors have been observed in 727 children with *in utero* ZDV exposure followed for over 1,100 person-years [14]. While these data are reassuring, followup is still limited and needs to be continued into adulthood before it can be concluded that there is no carcinogenic risk.

- Reproduction/fertility animal studies
No effect of zidovudine on reproduction or fertility in rodents has been seen. A dose-related cytotoxic effect on preimplantation mouse embryos can occur, with inhibition of blastocyst and postblastocyst development at a zidovudine concentrations similar to levels achieved with human therapeutic doses [15].
- Teratogenicity/developmental toxicity animal studies
No evidence of teratogenicity or toxicity was observed with administration of doses up to 500 to 600

mg/kg/day of zidovudine to pregnant rats, mice or rabbits. However, marked maternal toxicity and an increase in fetal malformations were noted in rats given a zidovudine dose of 3000 mg/kg/day (near the lethal dose, and 350 times the peak human plasma concentration).

In humans, data from PACTG 076 study and the Antiretroviral Pregnancy Registry do not demonstrate an increased incidence of congenital abnormalities in infants born to women with antepartum ZDV exposure [13, 16-18]. In the PACTG 076 study, the incidence of minor and major congenital abnormalities were similar between zidovudine and placebo groups, and no specific pattern of defects was seen [13, 17]. However, definitive conclusions regarding teratogenic risk cannot be made due to the limited numbers of children that have been evaluated.

- **Placental and breast milk passage in humans**
Zidovudine rapidly crosses the human placenta, achieving cord-to-maternal blood ratios of about 0.80. ZDV is excreted into human breast milk.
- **Human studies in pregnancy**
Zidovudine is well-tolerated in pregnancy at recommended adult doses and in the full-term neonate at 2 mg/per/kg body weight orally every 6 hours [13, 19]. Long-term data on the safety of *in utero* drug exposure in humans are not available for any antiretroviral drug; however, short-term data on the safety of zidovudine are reassuring. No difference in disease progression between women in PACTG 076 who received zidovudine and those who received placebo has been seen in followup through 4 years postpartum [20]. Infants with *in utero* zidovudine exposure followed for nearly 6 years have shown no significant differences from those who received placebo in immunologic, neurologic and growth parameters [17, 21]; followup of these infants is continuing.

Issues Related to Use of Nucleoside Analogue Drugs and Mitochondrial Toxicity

Nucleoside analogue drugs are known to induce mitochondrial dysfunction, as the drugs have varying affinity for mitochondrial gamma DNA polymerase. This affinity can result in interference with mitochondrial replication, resulting in mitochondrial DNA depletion and dysfunction [22]. The relative potency of the nucleosides in inhibiting mitochondrial gamma DNA polymerase *in vitro* is highest for zalcitabine (ddC), followed by didanosine (ddI), stavudine (d4T), lamivudine (3TC), ZDV, and abacavir (ABC). Toxicity related to mitochondrial dysfunction has been reported in infected patients receiving long-term treatment with nucleoside analogues, and generally has resolved with discontinuation of the drug or drugs; a possible genetic susceptibility to these toxicities has been suggested [23]. These toxicities may be of particular concern for pregnant women and for infants with *in utero* exposure to nucleoside analogue drugs.

Issues in Pregnancy: Clinical disorders linked to mitochondrial toxicity include neuropathy, myopathy, cardiomyopathy, pancreatitis, hepatic steatosis, and lactic acidosis. Among these disorders, symptomatic lactic acidosis and hepatic steatosis may have a female preponderance [24].

These syndromes have similarities to the rare but life-threatening syndromes of acute fatty liver of pregnancy and hemolysis, elevated liver enzymes and low platelets (the HELLP syndrome) that occur during the third trimester of pregnancy. A number of investigators have correlated these pregnancy-related disorders with a recessively-inherited mitochondrial abnormality in the fetus/infant that results in an inability to oxidize fatty acids [25-27]. Since the mother would be a heterozygotic carrier of the abnormal gene, there may be an increased risk of liver toxicity due to an inability to properly oxidize both maternal and accumulating fetal fatty acids [28]. Additionally, animal studies show that in late gestation pregnant mice have significant reductions (25%–50%) in mitochondrial fatty acid oxidation, and that exogenously administered estradiol and progesterone can reproduce these effects [29, 30]; whether this can be translated to humans is unknown. However, these data suggest that a disorder of mitochondrial fatty acid oxidation in the mother or her fetus during late pregnancy may play a role in the etiology of acute fatty liver of pregnancy and HELLP

syndrome, and possibly contribute to susceptibility to antiretroviral-associated mitochondrial toxicity.

Lactic acidosis with microvacuolar hepatic steatosis is a toxicity related to nucleoside analogue drugs that is thought to be related to mitochondrial toxicity; it has been reported in infected individuals treated with nucleoside analogue drugs for long periods of time (>6 months). Initially, most cases were associated with AZT, but subsequently other nucleoside analogue drugs have been associated with the syndrome, particularly d4T. In a report from the FDA Spontaneous Adverse Event Program of 106 individuals with this syndrome (60 in patients receiving combination and 46 receiving single nucleoside analogue therapy), typical initial symptoms included 1 to 6 weeks of nausea, vomiting, abdominal pain, dyspnea, and weakness [24]. Metabolic acidosis with elevated serum lactate and elevated hepatic enzymes was common. Patients in this report were predominantly female gender and high body weight. The incidence of this syndrome may be increasing, possibly due to increased use of combination nucleoside analogue therapy or increased recognition of the syndrome. In a cohort of infected patients receiving nucleoside analogue therapy followed at Johns Hopkins University between 1989 and 1994, the incidence of the hepatic steatosis syndrome was 0.13% per year [31]. However, in a report from a cohort of 964 HIV-infected individuals followed in France between 1997 and 1999, the incidence of symptomatic hyperlactatemia was 0.8% per year for all patients and 1.2% for patients receiving a regimen including d4T [32].

The frequency of this syndrome in pregnant HIV-infected women receiving nucleoside analogue treatment is unknown. In 1999, Italian researchers reported a case of severe lactic acidosis in an infected pregnant woman who was receiving d4T/3TC at the time of conception and throughout pregnancy who presented with symptoms and fetal demise at 38 weeks gestation [33]. Bristol-Myers Squibb has reported three maternal deaths due to lactic acidosis, two with and one without accompanying pancreatitis, in women who were either pregnant or postpartum and whose antepartum therapy during pregnancy included d4T and ddI in combination with other antiretroviral agents (either a protease inhibitor or nevirapine) [34, 35]. All cases were in women who were receiving treatment with these agents at the time of conception and continued for the duration of pregnancy; all presented late in gestation with symptomatic disease that progressed to death in the immediate postpartum period. Two cases were also associated with fetal demise.

It is unclear if pregnancy augments the incidence of the lactic acidosis/hepatic steatosis syndrome reported in non-pregnant individuals receiving nucleoside analogue treatment. However, because pregnancy itself can mimic some of the early symptoms of the lactic acidosis/hepatic steatosis syndrome or be associated with other significant disorders of liver metabolism, these cases emphasize the need for physicians caring for HIV-infected pregnant women receiving nucleoside analogue drugs to be alert for early diagnosis of this syndrome. Pregnant women receiving nucleoside analogue drugs should have hepatic enzymes and electrolytes assessed more frequently during the last trimester of pregnancy, and any new symptoms should be evaluated thoroughly. Additionally, because of the reports of several cases of maternal mortality secondary to lactic acidosis with prolonged use of the combination of d4T and ddI by HIV-infected pregnant women, clinicians should prescribe this antiretroviral combination during pregnancy with caution and generally only when other nucleoside analogue drug combinations have failed or caused unacceptable toxicity or side effects.

Issues with *In Utero* Exposure: A French group reported eight cases of uninfected infants with *in utero* and/or neonatal exposure to either ZDV/3TC (four infants) or ZDV alone (four infants) who developed indications of mitochondrial dysfunction after the first few months of life [32]. Two of these infants developed severe neurologic disease and died (both of whom had been exposed to ZDV/3TC), three had mild to moderate symptoms, and three had no symptoms but had transient laboratory abnormalities. It is important to note that an association between these findings and *in utero* exposure to antiretroviral drugs has not been established.

In infants followed through age 18 months in PACTG 076, the occurrence of neurologic events was rare — seizures occurred in one child exposed to ZDV and two exposed to placebo, and one child in each group had reported spasticity; mortality at 18 months was 1.4% in ZDV-exposed compared to 3.5% in placebo infants [17]. In a large database that included 223 deaths in over 20,000 children with and without antiretroviral drug exposure who were born to HIV-infected women followed prospectively in several large cohorts in the United States, no deaths similar to those reported from France were identified [36]. However, most of the infants with antiretroviral exposure had been exposed to ZDV alone, and only a relatively small proportion (approximately 6%) had been exposed to ZDV/3TC. Evaluation is ongoing to determine if there is any evidence of mitochondrial dysfunction among any of the

living children in these cohorts. Data have been reviewed relating to neurologic adverse events in 1,798 children that participated in PETRA, an African perinatal trial that compared three regimens of ZDV/3TC (before, during, and one week postpartum; during labor and postpartum; and during labor only) to placebo for prevention of transmission. No increased risk of neurologic events was observed among children treated with ZDV/3TC compared to placebo, regardless of the intensity of treatment [37]. Echocardiograms were prospectively performed every 4 to 6 months during the first 5 years of life in 382 uninfected infants born to HIV-infected women; 9% of infants had been exposed to ZDV prenatally [38]. No significant differences in ventricular function were observed between infants exposed and unexposed to ZDV.

If the association of mitochondrial dysfunction and *in utero* antiretroviral exposure proves to be real, the development of severe or fatal mitochondrial disease in these infants appears to be extremely rare, and should be compared to the clear benefit of ZDV in reducing transmission of a fatal infection by nearly 70% [39]. These data emphasize the importance of the existing Public Health Service recommendation for long-term followup for any child with *in utero* exposure to antiretroviral drugs.

NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS

Delavirdine (RESCRIPTOR[®]) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies
Long-term animal carcinogenicity studies with delavirdine in rodents are not completed; *in vitro* screening tests have been negative.
- Reproduction/fertility animal studies
Delavirdine does not impair fertility in rodents. Teratogenicity/developmental toxicity animal studies: Delavirdine is teratogenic in rats; doses of 50 to 200 mg/kg/day during organogenesis caused ventricular septal defects. Exposure of rats to doses approximately 5 times human therapeutic exposure resulted in marked maternal toxicity, embryotoxicity, fetal developmental delay, and reduced pup survival.

Abortions, embryotoxicity, and maternal toxicity were observed in rabbits at doses approximately 6 times human therapeutic exposure.

- Placental and breast milk passage in animal studies
Whether delavirdine crosses the placenta is unknown. Delavirdine is excreted in the milk of lactating rats; however, it is unknown if the drug is excreted in human breast milk.
- Human studies in pregnancy
Delavirdine has not been evaluated in HIV-infected pregnant women. In premarketing clinical studies, the outcomes of seven unplanned pregnancies were reported: three resulted in ectopic pregnancies, three resulted in healthy live births, and one infant was born prematurely with a small muscular ventricular septal defect to a patient who received approximately 6 weeks of treatment with delavirdine and zidovudine early in the course of pregnancy.

Efavirenz (SUSTIVA[®]) is FDA pregnancy category C.

- Animal carcinogenicity studies
Long-term animal carcinogenicity studies with efavirenz in rats and mice are not completed; *in vitro* screening tests have been negative.
- Reproduction/fertility animal studies
No effect of efavirenz on reproduction or fertility in rodents has been seen. An increase in fetal resorptions has been observed in rats at doses comparable to or lower than those used to achieve human therapeutic exposure.
- Teratogenicity/developmental toxicity animal studies
Significant central nervous system malformations were observed in 3 of 20 infants born to pregnant cynomolgus monkeys receiving efavirenz from gestational days 20 to 150 at a dose of 30 mg/kg twice daily (resulting in plasma concentrations comparable to systemic human therapeutic exposure) [40]. The malformations included anencephaly and unilateral anophthalmia in one; microphthalmia in another; and cleft palate in the third. Primate teratogenicity studies have not been conducted for delavirdine or nevirapine.
- Placental and breast milk passage in animal studies
Efavirenz crosses the placenta in rats, rabbits, and primates, producing cord blood concentrations similar to concentrations in maternal plasma. It is unknown whether efavirenz is excreted in human breast milk.
- Human studies in pregnancy
No **clinical trials** with efavirenz in pregnant humans are **planned**. **There has been a case report of**

myelomeningocele in a human infant born to a woman who was receiving efavirenz at the time of conception and during the first trimester [41, 42]. Because of the potential for teratogenicity, pregnancy should be avoided in women receiving efavirenz. It should be noted that non-nucleoside reverse transcriptase inhibitors like nevirapine and efavirenz as well as the protease inhibitors may affect estrogen and/or norethindrone blood concentrations in women receiving oral contraceptives; additional or alternative contraception should be used by women using oral contraceptives who are receiving these antiretroviral agents. There are insufficient data on drug interactions with injectable hormones (DEPO-PROVERA) to make recommendations regarding the need for additional contraception. Theoretically, since hormone levels are much higher with injectable than oral contraceptives, interactions with antiretroviral drugs may be less significant.

Nevirapine (Viramune®) is FDA pregnancy category C.

▪ Animal carcinogenicity studies

Long-term animal carcinogenicity studies with nevirapine in rats and mice are not completed; *in vitro* screening tests have been negative.

▪ Reproduction/fertility animal studies

Evidence of impaired fertility was seen in female rats at nevirapine doses providing systemic exposure comparable to human therapeutic exposure.

▪ Teratogenicity/developmental toxicity animal studies

Teratogenic effects of nevirapine have not been observed in reproductive studies with rats and rabbits. In rats, however, a significant decrease in fetal weight occurred at doses producing systemic concentrations approximately 50% higher than human therapeutic exposure.

▪ Placental and breast milk passage in humans

Nevirapine crosses the placenta and achieves neonatal blood concentrations equivalent to that in the mother (cord-to-maternal blood ratio approximately 0.90) [43]. Nevirapine is excreted into human breast milk; the median concentration in four breast milk samples obtained from three women during the first week after delivery was approximately 76% (range 54 to 104%) of serum levels [43].

▪ Human studies in pregnancy

A phase I study (PACTG 250) evaluated the safety and pharmacokinetics of nevirapine, administered to infected pregnant women as a single 200 mg dose at the onset of labor and as a single 2 mg/kg dose to the infant at age 48 to 72 hours [43]. No adverse effects were seen in the women or the infants.

Pharmacokinetic parameters in pregnant women receiving intrapartum nevirapine were similar though somewhat more variable than in nonpregnant adults, possibly due to incomplete drug absorption associated with impaired gastrointestinal function during labor. Pharmacokinetic data on chronic antenatal nevirapine dosing in pregnant women are under study but not yet available. Nevirapine elimination was prolonged in the infants. The regimen maintained serum concentrations associated with antiviral activity in the infants for the first week of life. The HIVNET 012 study in Uganda compared nevirapine (200 mg orally to the mother at the onset of labor and 2 mg/kg to the neonate within 72 hours of birth) with zidovudine (600 mg orally to the mother at the onset of delivery and 300 mg every 3 hours until delivery, and 4 mg/kg orally twice daily for the first 7 days of life to the neonate). In this study, nevirapine lowered the risk of HIV transmission by nearly 50% during the first 14–16 weeks of life compared with zidovudine [44]. However, the women in this African trial were not receiving any other antiretroviral therapy. In the U.S., most infected women who know their HIV status during pregnancy receive standard ZDV prophylaxis combined with whatever antiretroviral therapy is needed for treatment of their HIV disease; it is unknown whether adding the HIVNET 012 nevirapine regimen to standard antiretroviral prophylaxis and treatment offers any additional benefit in terms of reducing perinatal transmission. A phase III perinatal trial (PACTG 316) being conducted in the United States, Europe, the Bahamas and Brazil is evaluating this regimen in combination with standard maternal antiretroviral therapy and ZDV antiretroviral therapy and ZDV prophylaxis for the prevention of perinatal HIV transmission. Selection of nevirapine-resistant virus was found at 6 weeks postpartum in both the untreated and antiretroviral-treated pregnant women who received a single dose of nevirapine in labor in HIVNET 012 and PACTG 316. In HIVNET 012, 7 of 31 women (23%) evaluated developed genotypic resistance mutations at 6 weeks postpartum; these mutations were no longer present in 4 women studied at 13–18 months postpartum [37, 45]. In the antiretroviral-treated women in PACTG 316, 4 of 32 women (13%, 95% CI 4–25%) with HIV-1 RNA above 3,000 copies/mL at delivery who received

nevirapine developed genotypic nevirapine resistance mutations compared to none of 38 women in the placebo arm [39].

Severe, life-threatening, and in some cases, fatal hepatotoxicity, including fulminant and cholestatic hepatitis, hepatic necrosis, and hepatic failure, has been reported in HIV-infected patients receiving nevirapine in combination with other drugs for treatment of HIV disease and in a small number of individuals receiving nevirapine as part of a combination regimen for post-exposure prophylaxis of nosocomial or sexual HIV exposure [46]. These events have generally occurred during the first 12 weeks of therapy, and may present with non-specific prodromal signs or symptoms of hepatitis. This has not been reported in women or infants receiving two-dose nevirapine (the HIVNET 012 regimen) for prevention of perinatal transmission. Severe, life-threatening hypersensitivity skin reactions, including Stevens-Johnson syndrome, have been reported in HIV-infected individuals receiving nevirapine for treatment, usually during the first 12 weeks of therapy. This has not been reported with use of the HIVNET 012 two-dose nevirapine regimen.

PROTEASE INHIBITORS

Issues Related to the Use of Protease Inhibitors

Hyperglycemia and diabetes mellitus:

Hyperglycemia, new onset diabetes mellitus, exacerbation of existing diabetes mellitus, and diabetic ketoacidosis have been reported with administration of protease inhibitor antiretroviral drugs in HIV-infected patients [47-50]. In addition, pregnancy is itself a risk factor for hyperglycemia; it is unknown if the use of protease inhibitors will exacerbate the risk for pregnancy-associated hyperglycemia. Clinicians caring for HIV-infected pregnant women who are receiving protease inhibitor therapy should be aware of the risk of this complication, and closely monitor glucose levels. Symptoms of hyperglycemia should be discussed with pregnant women who are receiving protease inhibitors.

Combination Therapy: There are limited data concerning combination antiretroviral therapy in pregnancy. A retrospective Swiss report evaluated the pregnancy outcome in 37 HIV-infected pregnant women treated with combination therapy; all received two reverse transcriptase inhibitors and 16 received one or two protease inhibitors [51]. Almost 80% of women developed one or more typical adverse effects of the drugs such as anemia, nausea/vomiting, aminotransferase elevation, or hyperglycemia. A possible association of combination antiretroviral therapy with preterm births was noted, as 10 of 30 babies were born prematurely. The preterm birth rate did not differ between women receiving combination therapy with or without protease inhibitors. The contribution of maternal HIV disease stage and other covariates that might be associated with a risk for prematurity were not assessed. Furthermore, some studies have shown elevated preterm birth rates in HIV-infected women who have not received any antiretroviral therapy [52-54]. To evaluate the baseline rates of adverse pregnancy outcome and risk factors for such outcomes in HIV-infected pregnant women, a meta-analysis of multiple PACTG perinatal trials and cohort studies is in progress. Preliminary analyses do not indicate an elevated risk of preterm delivery among infants born to women receiving combination antiretroviral therapy with or without protease inhibitors compared to those receiving single drug or no antiretroviral therapy. Until more information is known, it is recommended that HIV-infected pregnant women who are receiving combination therapy for treatment of their HIV infection should continue their provider-recommended regimen. They should receive careful, regular monitoring for pregnancy complications and for potential toxicities.

Individual Agents: Protease Inhibitors

Phase I studies of four of the approved protease inhibitors (indinavir, ritonavir, nelfinavir and saquinavir soft gel capsule in combination with ZDV and 3TC) in pregnant HIV-infected women and their infants are ongoing in the United States. However, complete data are not yet available regarding drug dosage, safety, and tolerance of the protease inhibitors in pregnancy or in neonates. Amprenavir and lopinavir/ritonavir (Kaletra™), two more recently approved protease inhibitors, have not yet been studied in pregnant women or neonates.

Amprenavir (AGENERASE[®]) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies
Long-term animal carcinogenicity studies of amprenavir in rats and mice are not completed; *in vitro* screening tests have been negative.
- Reproduction/fertility animal studies
No effect has been seen on reproductive performance, fertility, or embryo survival in rats at exposures about twice those of human therapeutic exposure.
- Teratogenicity/developmental toxicity animal studies
In pregnant rabbits, administration of amprenavir resulting in systemic exposures about one-twentieth of that observed with human therapeutic exposure was associated with abortions and an increased incidence of minor skeletal variations resulting from deficient ossification of the femur, humerus trochlea and humerus. In rat fetuses, thymic elongation and incomplete ossification of bones were also attributed to amprenavir at systemic exposures about one-half that associated with the recommended human dose. Reduced body weights of approximately 10–20% were observed in offspring of rodents administered amprenavir from day 7 of gestation to day 22 of lactation (exposures approximately twice that observed with the human therapeutic dose). However, the subsequent development of the offspring, including fertility and reproductive performance, was not affected by maternal administration of amprenavir.
- Placental and breast milk passage in animals
Whether amprenavir crosses the placenta is unknown. Amprenavir is excreted in the milk of lactating rats; it is not known if it is excreted in human milk.
- Human studies in pregnancy
There have been no studies of amprenavir in pregnant women or neonates.

Indinavir (CRIXIVAN[®]) is classified as FDA pregnancy category C.

- Animal carcinogenicity studies
Long-term animal carcinogenicity studies with indinavir in rats and mice are not completed; *in vitro* screening tests have been negative.
- Reproduction/fertility animal studies
No effect of indinavir has been seen on reproductive performance, fertility, or embryo survival in rats.

- Teratogenicity/developmental toxicity animal studies
There has been no evidence of teratogenicity of indinavir in rats, rabbits or dogs. In rats, developmental toxicity manifested by an increase in supernumerary and cervical ribs was observed at doses comparable to those administered to humans. No treatment-related external, visceral or skeletal changes were seen in rabbits (fetal exposure limited, approximately 2% of maternal levels) or dogs (fetal exposure approximately 50% of maternal levels). Indinavir was administered to Rhesus monkeys during the third trimester of pregnancy (at doses up to 160 mg/kg twice daily) and to neonatal Rhesus monkeys (at doses up to 160 mg/kg twice daily). When administered to neonates, indinavir caused an exacerbation of the transient physiologic hyperbilirubinemia seen in this species after birth; serum bilirubin values were approximately fourfold above controls at 160 mg/kg twice daily. A similar exacerbation did not occur in neonates after *in utero* exposure to indinavir during the third trimester of pregnancy. In Rhesus monkeys, fetal plasma drug levels were approximately 1–2% of maternal plasma drug levels approximately 1 hour after maternal dosing at 40, 80, or 160 mg/kg twice daily.
- Placental and breast milk passage in animals
Significant placental passage of indinavir occurs in rats and dogs, but only limited placental transfer occurs in rabbits. Indinavir is excreted in the milk of lactating rats at concentrations slightly above maternal levels (milk-to-plasma ratio 1.26 to 1.45); it is not known if indinavir is excreted in human milk.
- Human studies in pregnancy:
A phase I/II safety and pharmacokinetic study (PACTG 358) of indinavir (800 mg tid) in combination with ZDV and lamivudine in pregnant HIV-infected women and their infants is being conducted (the infants do not receive indinavir in this study). Preliminary data are available from five women and infants [55]. One woman discontinued indinavir due to nausea and vomiting; adverse effects in the women included one case of moderately severe hyperbilirubinemia and one case of flank pain without renal stones, both of which resolved spontaneously and did not require drug discontinuation. Pharmacokinetic data from three women indicate that the plasma area under the curve (AUC) indinavir level was lower during pregnancy than postpartum or than observed in non-pregnant HIV-infected individuals. However, HIV RNA levels in the four women who completed the study decreased to undetectable levels (<400 copies/mL) prior to delivery and CD4 cell number and percentage significantly increased. The median

gestational age of the five infants was 39 weeks (range 36–39 weeks). In a pharmacokinetic study of two pregnant HIV-infected women receiving combination therapy including indinavir (800 mg tid), a marked difference was noted between the AUC indinavir exposure between the third trimester and postpartum evaluations [56]. The AUC during the third trimester was reduced by 63% in one and 86% in the other woman when compared to 9–12 week postpartum evaluations in the same women. Similar reductions in maximum plasma indinavir concentrations were observed.

Lopinavir + Ritonavir (KALETRA™) is classified as FDA pregnancy category C.

▪ Animal carcinogenicity studies

Long-term animal carcinogenicity screening studies of lopinavir + ritonavir in animal systems are not completed. *In vitro* mutagenicity and clastogenicity screening tests are negative for both lopinavir and ritonavir.

Carcinogenicity studies in mice and rats have been carried out for ritonavir. In male mice, at levels of 50, 100 or 200 mg/kg/day, a dose-dependent increase in liver adenomas and combined adenomas and carcinomas was observed; based on AUC, exposure in male mice at the highest dose was approximately fourfold that in male humans at the recommended therapeutic dose (400 mg lopinavir/100 mg ritonavir bid). No carcinogenic effects were observed in female mice with exposures ninefold that of female humans at the recommended therapeutic dose. No carcinogenic effects were observed in rats at exposures up to 0.7-fold that of humans at the recommended therapeutic dose.

▪ Reproduction/fertility animal studies

Lopinavir in combination with ritonavir at a 2:1 ratio produced no effects on fertility in male and female rats with exposures approximately 0.7-fold for lopinavir and 1.8-fold for ritonavir of the exposures in humans at the recommended therapeutic dose.

▪ Teratogenicity/developmental toxicity animal studies

There has been no evidence of teratogenicity with administration of lopinavir + ritonavir to pregnant in rats or rabbits. In rats treated with maternally toxic dosage (100 mg lopinavir/50 mg ritonavir/kg/day), embryonic and fetal developmental toxicities (early resorption, decreased fetal viability, decreased fetal body weight, increased incidence of skeletal variations and skeletal ossification delays) were observed; drug

exposure in the pregnant rats was 0.7-fold for lopinavir and 1.8-fold for ritonavir of the exposures in humans at the recommended therapeutic dose. In a peri- and postnatal study in rats, a decrease in survival of pups between birth and postnatal day 21 occurred with exposures of 40 mg lopinavir/20 mg ritonavir/kg/day or greater. In rabbits, no embryonic or fetal developmental toxicities were observed with maternally toxic dosage, where drug exposure was 0.6-fold for lopinavir and 1.0-fold for ritonavir of the exposures in humans at recommended therapeutic dose.

▪ Placental and breast milk passage in animals

Data on placental passage of lopinavir in animals are not available. For ritonavir, transplacental passage has been observed in rat fetuses at mid- and late-gestation. Lopinavir and ritonavir are secreted in the breast milk of lactating rats; it is not known if either drug is excreted in human milk.

▪ Human studies in pregnancy

No studies of lopinavir in human pregnancies have been conducted. A phase I/II safety and pharmacokinetic study of ritonavir given at therapeutic doses (600 mg bid) in combination with ZDV and lamivudine in pregnant HIV-infected women and their infants (PACTG 354) is being conducted but complete data are not yet available; preliminary data indicate that there is minimal, if any, placental passage of ritonavir in humans.

Nelfinavir (VIRACEPT®) is classified as FDA pregnancy category B.

▪ Animal carcinogenicity studies

Long-term animal carcinogenicity studies of nelfinavir in rats and mice are not completed; *in vitro* screening tests have been negative.

▪ Reproduction/fertility animal studies

No effect of nelfinavir has been seen on reproductive performance, fertility, or embryo survival in rats at exposures comparable to human therapeutic exposure.

▪ Teratogenicity/developmental toxicity animal studies

No teratogenicity or effect on fetal development by nelfinavir has been demonstrated in rodent or rabbit studies at exposures comparable to human therapeutic exposure.

▪ Placental and breast milk passage in animals

Whether nelfinavir crosses the placenta is unknown. Nelfinavir is excreted in the milk of lactating rats; it is not known if it is excreted in human milk.

- Human studies in pregnancy

A phase I/II safety and pharmacokinetic study (PACTG 353) of nelfinavir in combination with ZDV and lamivudine in pregnant HIV-infected women and their infants is being conducted, but complete data are not yet available. In preliminary data from this study, the standard adult dose of nelfinavir (750 mg tid) produced drug exposures in the first nine pregnant HIV-infected women enrolled in the study that were variable and generally lower than those reported in non-pregnant adults for both tid and bid dosing. Therefore, the study has been modified to evaluate an increased dose of nelfinavir (1250 mg) administered bid. In infants, nelfinavir was not detectable in cord blood from four infants born to mothers receiving 750 mg nelfinavir tid; in one additional infant, the cord blood nelfinavir concentration was 11.7% that detected in maternal blood at delivery.

Ritonavir (Norvir[®]) is classified as FDA pregnancy category B.

- Animal carcinogenicity studies

In vitro mutagenicity and clastogenicity screening tests are negative for ritonavir. Carcinogenicity studies in mice and rats have been completed. In male mice, at levels of 50, 100 or 200 mg/kg/day, a dose-dependent increase in liver adenomas and combined adenomas and carcinomas was observed; based on AUC, exposure in male mice at the highest dose was approximately fourfold that in male humans at the recommended therapeutic dose (400 mg lopinavir/100 mg ritonavir bid). No carcinogenic effects were observed in female mice with exposures ninefold that of female humans at the recommended therapeutic dose. No carcinogenic effects were observed in rats at exposures up to 0.7-fold that of humans at the recommended therapeutic dose.

- Reproduction/fertility animal studies

No effect of ritonavir has been seen on reproductive performance or fertility in rats at drug exposures 40% (male) and 60% (female) of that achieved with human therapeutic dosing; higher doses were not feasible due to hepatic toxicity in the rodents.

- Teratogenicity/developmental toxicity animal studies

No ritonavir-related teratogenicity has been observed in rats or rabbits. Developmental toxicity was observed in rats, including early resorptions, decreased body weight, ossification delays, and developmental variations such as wavy ribs and enlarged fontanelles; however, these effects occurred only at maternally toxic dosages (exposure equivalent to 30% of human

therapeutic exposure). In addition, a slight increase in cryptorchidism was also noted in rats at exposures equivalent to 22% of the human therapeutic dose. In rabbits, developmental toxicity (resorptions, decreased litter size, and decreased fetal weight) was observed only at maternally toxic doses (1.8 times human therapeutic exposure).

- Placental and breast milk passage in animals

Transplacental passage of ritonavir has been observed in rats with fetal tissue to maternal serum ratios >1.0 at 24 hours post-dose in mid- and late-gestation fetuses. In a human placental perfusion model, the clearance index of ritonavir was very low, with little accumulation in the fetal compartment and no accumulation in placental tissue [57]. Ritonavir is excreted in the milk of lactating rats; it is unknown if it is excreted in human milk.

- Human studies in pregnancy

A phase I/II safety and pharmacokinetic study (PACTG 354) of ritonavir in combination with zidovudine and lamivudine in pregnant HIV-infected women and their infants is being conducted, but complete data are not yet available. Preliminary data indicate minimal, if any, placental passage of ritonavir.

Saquinavir (INVIRASE[®] [Hard Gel Capsule]/FORTOVASE[®] [Soft Gel Capsule]) is classified as FDA pregnancy category B.

- Animal carcinogenicity studies

Long-term animal carcinogenicity studies of saquinavir in rats and mice are not completed; *in vitro* screening tests have been negative.

- Reproduction/fertility animal studies

No effect of saquinavir has been seen on reproductive performance, fertility, or embryo survival in rats. Administration of low doses of saquinavir to newborn rats was associated with gastrointestinal toxicity, including inflammation at the rectoanal junction and red anal fluid; mortality was seen at very high doses (1200 mg/kg/day).

- Teratogenicity/developmental toxicity animal studies

No evidence for embryotoxicity or teratogenicity of saquinavir has been found in animal studies.

- Placental and breast milk transfer in animal studies

Placental transfer of saquinavir in the rat and rabbit was minimal. Saquinavir is excreted in the milk of lactating rats; it is not known if it is excreted in human milk.

- Human studies in pregnancy
A phase I/II safety and pharmacokinetic study (PACTG 386) of saquinavir in combination with ZDV and lamivudine in pregnant HIV-infected women and their infants is being conducted, but complete data are not yet available. In preliminary data from this study, the standard adult dose of saquinavir (1200 mg tid) was not sufficient to produce adequate drug exposure in the first four pregnant HIV-infected women enrolled in the study compared to those obtained with standard dosing in non-pregnant adults. Therefore, the study has been modified to evaluate a dose of saquinavir 800 mg combined with zidovudine 100 mg both administered bid.
- Reproduction/fertility animal studies
Hydroxyurea administered to male rats at doses of 60 mg/kg/day (about 0.3 times the maximum recommended human daily dose on a mg/m² basis) produced testicular atrophy, decreased spermatogenesis, and significantly reduced their ability to impregnate females.
- Teratogenicity/developmental toxicity animal studies
Potent teratogenic effects have been observed in all animal species tested, with defects reported in multiple organ systems [58-64]. Administration of hydroxyurea to pregnant rats at doses as low as 180 mg/kg/day (about 0.8 times the maximum recommended human daily dose on a mg/m² basis) and pregnant rabbits at 30 mg/kg/day (about 0.3 times the maximum recommended human daily dose on a mg/m² basis) was associated with embryotoxicity and fetal malformations. In pregnant rats administered doses ranging from 185 to 1000 mg/kg body weight, fetal defects that have been observed include central nervous system, cardiovascular, ocular, craniofacial, and skeletal anomalies, limb deformities, and diaphragmatic hernia, with the pattern of defects dependent on gestational day of exposure [58, 61, 62]. Exposure early in gestation was associated with embryo death in a large percentage of cases. In pregnant rats, single doses of 375 mg/kg body weight or more (about 1.7 times the maximum recommended human daily dose on a mg/m² basis), were associated with growth retardation and impaired learning ability in their offspring. In hamsters, neural tube defects and cardiovascular abnormalities were produced after a 50 mg dose of hydroxyurea was given intravenously [59]. In pregnant rhesus monkeys administered a cumulative dose greater than 500 mg/kg body weight, multiple skeletal, genitourinary, cardiac, and ocular anomalies were found in their offspring [61]. Teratogenicity was also demonstrated in pregnant cats given a single oral dose of 50 or 100 mg/kg body weight [60].

MISCELLANEOUS AGENTS

Hydroxyurea is classified as FDA pregnancy category D.

Hydroxyurea is a cytotoxic and antimetabolic agent that inhibits DNA synthesis and has been used for treatment of myeloproliferative disorders and sickle cell anemia. It has recently been studied for treatment of HIV disease in combination with nucleoside analogue antiretroviral agents. By inhibiting ribonucleotide reductase, it depletes the pool of deoxynucleoside triphosphates, particularly dATP, thereby potentiating the incorporation of the nucleoside analogue drugs into viral DNA and increasing their antiretroviral effect. However, the drug has significant toxicities and its role in HIV therapy is not well defined.

- Animal carcinogenicity studies and human data
Hydroxyurea is genotoxic in a wide range of *in vitro* and *in vivo* animal test systems, causes cellular transformation to a tumorigenic phenotype, and is a transspecies carcinogen, which implies a potential carcinogenic risk to humans. Conventional long-term animal carcinogenicity studies have not been performed. However, intraperitoneal administration of 125 to 250 mg/kg of hydroxyurea (approximately 0.6 to 1.2 times the maximum recommended human oral dose on a mg/m² basis) three times weekly for 6 months to female rats increased the incidence of mammary tumors in rats surviving to 18 months compared to controls.

In humans receiving long-term hydroxyurea for myeloproliferative disorders such as polycythemia vera, secondary leukemias have been reported. It is unknown whether this leukemogenic effect is secondary to hydroxyurea or is associated with the patients' underlying disease. Skin cancer has also been reported in patients receiving long-term therapy.

- Placental and breast milk passage in animal studies
Hydroxyurea has been shown to cross the placenta in animals.
- Placental and breast milk passage in humans
Hydroxyurea is excreted in human milk [65].
- Human studies in pregnancy
Published reports of hydroxyurea during human pregnancy include 16 women, all treated for primary hematologic illnesses (e.g., chronic myeloid leukemia, sickle cell anemia, primary thrombocytopenia) [66]. Doses ranged from 0.5 to 3 g/day and 13 women had first trimester exposure. No fetal anomalies were seen

and normal pregnancy outcomes were reported, except for one stillbirth with eclampsia at 26 weeks gestation and four elective pregnancy terminations.

Because of concerns raised by the significant anomalies seen in multiple animal species exposed to hydroxyurea and limited human information, as well as the uncertain role of hydroxyurea in HIV therapy, hydroxyurea use as an antiretroviral regimen component should be avoided during pregnancy. Clinicians should counsel women of childbearing potential about potential risks of teratogenicity if they are treated with hydroxyurea and become pregnant, and encouraged to use effective contraception and avoid becoming pregnant while being treated with hydroxyurea.

ANTIRETROVIRAL PREGNANCY REGISTRY

The Antiretroviral Pregnancy Registry is an epidemiologic project to collect observational, nonexperimental data on antiretroviral exposure during pregnancy for the purpose of assessing the potential teratogenicity of these drugs. Registry data will be used to supplement animal toxicology studies and assist clinicians in weighing the potential risks and benefits of treatment for individual patients. The registry is a collaborative project of the pharmaceutical manufacturers with an advisory committee of obstetric and pediatric practitioners.

It is strongly recommended that health care providers who are treating HIV-1-infected pregnant women and their newborns report cases of prenatal exposure to antiretroviral drugs (either alone or in combination) to the Antiretroviral Pregnancy Registry. The registry does not use patient names, and birth outcome followup is obtained by registry staff from the reporting physician.

Referrals should be directed to
Antiretroviral Pregnancy Registry
Research Park
1011 Ashes Drive
Wilmington, NC 28405
Telephone 1-800-258-4263
Fax 1-800-800-1052
Internet access www.apregistry.com

References

1. Toltzis P, Mourton T, Magnuson T. Comparative embryonic cytotoxicity of antiretroviral nucleosides. *J Infect Dis*, 1994. 169(5): p. 1100-2.
2. Wang Y, Livingston E, Patil S, et al. Pharmacokinetics of didanosine in antepartum and postpartum human immunodeficiency virus-infected pregnant women and their neonates: an AIDS clinical trials group study. *J Infect Dis*, 1999. 180(5): p. 1536-41.
3. Moodley J, Moodley D, Pillay K, et al. Pharmacokinetics and antiretroviral activity of lamivudine alone or when coadministered with zidovudine in human immunodeficiency virus type 1-infected pregnant women and their offspring. *J Infect Dis*, 1998. 178(5): p. 1327-33.
4. Odinecs A, Nosbisch C, Keller RD, et al. In vivo maternal-fetal pharmacokinetics of stavudine (2',3'-didehydro- 3'-deoxythymidine) in pigtailed macaques (*Macaca nemestrina*). *Antimicrob Agents Chemother*, 1996. 40(1): p. 196-202.
5. Odinecs A, Pereira C, Nosbisch C, Unadkat JD. Prenatal and postpartum pharmacokinetics of stavudine (2',3'-didehydro- 3'-deoxythymidine) and didanosine (dideoxyinosine) in pigtailed macaques (*Macaca nemestrina*). *Antimicrob Agents Chemother*, 1996. 40(10): p. 2423-5.
6. Tarantal AF, Castillo A, Ekert JE, et al. Fetal and maternal outcome after administration of tenofovir to gravid rhesus monkeys (*Macaca mulatta*). *J Acquir Immune Defic Syndr*, 2002. 29(3): p. 207-20.
7. Tarantal AF, Marthas ML, Shaw JP, et al. Administration of 9-[2-(R)-(phosphonomethoxy)propyl]adenine (PMPA) to gravid and infant rhesus macaques (*Macaca mulatta*): safety and efficacy studies. *J Acquir Immune Defic Syndr Hum Retrovirol*, 1999. 20(4): p. 323-33.
8. Sandberg JA, Binienda Z, Lipe G, et al. Placental transfer and fetal disposition of 2',3'-dideoxycytidine and 2',3'-dideoxyinosine in the rhesus monkey. *Drug Metab Dispos*, 1995. 23(8): p. 881-4.
9. Ayers KM, Clive D, Tucker WE, Jr., et al. Nonclinical toxicology studies with zidovudine: genetic toxicity tests and carcinogenicity bioassays in mice and rats. *Fundam Appl Toxicol*, 1996. 32(2): p. 148-58.
10. Olivero OA, Anderson LM, Diwan BA, et al. Transplacental effects of 3'-azido-2',3'-dideoxythymidine (AZT): tumorigenicity in mice and genotoxicity in mice and monkeys. *J Natl Cancer Inst*, 1997. 89(21): p. 1602-8.
11. Ayers KM, Torrey CE, Reynolds DJ. A transplacental carcinogenicity bioassay in CD-1 mice with zidovudine. *Fundam Appl Toxicol*, 1997. 38(2): p. 195-8.
12. Reggy AA, Rogers MF, Simonds RJ. Using 3'-azido-2',3'-dideoxythymidine (AZT) to prevent perinatal human immunodeficiency virus transmission and risk

- of transplacental carcinogenesis. *J Natl Cancer Inst*, 1997. 89(21): p. 1566-7.
13. Connor EM, Sperling RS, Gelber R, et al. Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. Pediatric AIDS Clinical Trials Group Protocol 076 Study Group. *N Engl J Med*, 1994. 331(18): p. 1173-80.
 14. Hanson IC, Antonelli TA, Sperling RS, et al. Lack of tumors in infants with perinatal HIV-1 exposure and fetal/neonatal exposure to zidovudine. *J Acquir Immune Defic Syndr Hum Retrovirol*, 1999. 20(5): p. 463-7.
 15. Toltzis P, Marx CM, Kleinman N, et al. Zidovudine-associated embryonic toxicity in mice. *J Infect Dis*, 1991. 163(6): p. 1212-8.
 16. Antiretroviral pregnancy registry. PharmaResearch Corp., Wilmington, NC. January 1989 - July 1997.
 17. Sperling RS, Shapiro DE, McSherry GD, et al. Safety of the maternal-infant zidovudine regimen utilized in the Pediatric AIDS Clinical Trial Group 076 Study. *AIDS*, 1998. 12(14): p. 1805-13.
 18. White A, Eldridge R, Andrews E. Birth outcomes following zidovudine exposure in pregnant women: the Antiretroviral Pregnancy Registry. *Acta Paediatr Suppl*, 1997. 421: p. 86-8.
 19. O'Sullivan MJ, Boyer PJ, Scott GB, et al. The pharmacokinetics and safety of zidovudine in the third trimester of pregnancy for women infected with human immunodeficiency virus and their infants: phase I acquired immunodeficiency syndrome clinical trials group study (protocol 082). Zidovudine Collaborative Working Group. *Am J Obstet Gynecol*, 1993. 168(5): p. 1510-6.
 20. Bardeguez AD, Mofenson LM, Spino C, et al. Lack of clinical and immunologic disease progression with transient use of zidovudine (ZDV) to reduce perinatal HIV-1 transmission in PACTG 076. *12th World Conference on AIDS*, 1998. (Abstract 12233).
 21. Culnane M, Fowler M, Lee SS, et al. Lack of long-term effects of *in utero* exposure to zidovudine among uninfected children born to HIV-infected women. Pediatric AIDS Clinical Trials Group Protocol 219/076 Teams. *J Am Med Assoc*, 1999. 281(2): p. 151-7.
 22. Brinkman K, ter Hofstede HJ, Burger DM, et al. Adverse effects of reverse transcriptase inhibitors: mitochondrial toxicity as common pathway. *AIDS*, 1998. 12(14): p. 1735-44.
 23. Mofenson LM. Perinatal exposure to zidovudine--benefits and risks. *N Engl J Med*, 2000. 343(11): p. 803-5.
 24. Boxwell DE, Styr BA. Lactic acidosis in patients receiving nucleoside reverse transcriptase inhibitors. 39th Interscience Conference on Antimicrobial Agents and Chemotherapy. San Francisco, CA, 1999. (Abstract 1284).
 25. Ibdah JA, Bennett MJ, Rinaldo P, et al. A fetal fatty-acid oxidation disorder as a cause of liver disease in pregnant women. *N Engl J Med*, 1999. 340(22): p. 1723-31.
 26. Strauss AW, Bennett MJ, Rinaldo P, et al. Inherited long-chain 3-hydroxyacyl-CoA dehydrogenase deficiency and a fetal-maternal interaction cause maternal liver disease and other pregnancy complications. *Semin Perinatol*, 1999. 23(2): p. 100-12.
 27. Sims HF, Brackett JC, Powell CK, et al. The molecular basis of pediatric long chain 3-hydroxyacyl-CoA dehydrogenase deficiency associated with maternal acute fatty liver of pregnancy. *Proc Natl Acad Sci U S A*, 1995. 92(3): p. 841-5.
 28. Ibdah JA, Yang Z, Bennett MJ. Liver disease in pregnancy and fetal fatty acid oxidation defects. *Mol Genet Metab*, 2000. 71(1-2): p. 182-9.
 29. Grimbert S, Fromenty B, Fisch C, et al. Decreased mitochondrial oxidation of fatty acids in pregnant mice: possible relevance to development of acute fatty liver of pregnancy. *Hepatology*, 1993. 17(4): p. 628-37.
 30. Grimbert S, Fisch C, Deschamps D, et al. Effects of female sex hormones on mitochondria: possible role in acute fatty liver of pregnancy. *Am J Physiol*, 1995. 268(1 Pt 1): p. G107-15.
 31. Fortgang IS, Belitsos PC, Chaisson RE, Moore RD. Hepatomegaly and steatosis in HIV-infected patients receiving nucleoside analog antiretroviral therapy. *Am J Gastroenterol*, 1995. 90(9): p. 1433-6.
 32. Gerard Y, Maulin L, Yazdanpanah Y, et al. Symptomatic hyperlactataemia: an emerging complication of antiretroviral therapy. *AIDS*, 2000. 14(17): p. 2723-30.
 33. Luzzati R, Del Bravo P, Di Perri G, et al. Riboflavine and severe lactic acidosis [letter]. *Lancet*, 1999. 353(9156): p. 901-2.
 34. Bristol-Myers Squibb Company. Healthcare provider important drug warning letter. January 5, 2001.
 35. Sarner L, Fakoya A. Acute onset lactic acidosis and pancreatitis in the third trimester of pregnancy in HIV-1 positive women taking antiretroviral medication. *Sex Transm Inf*, 2002. 78(1): p. 58-9.
 36. Chun TW, Stuyver L, Mizell SB, et al. Presence of an inducible HIV-1 latent reservoir during highly active antiretroviral therapy. *Proc Natl Acad Sci USA*, 1997. 94(24): p. 13193-7.
 37. Jackson JB, Becker-Pergola G, Guay LA, et al. Identification of the K103N resistance mutation in Ugandan women receiving nevirapine to prevent HIV-1 vertical transmission. *AIDS*, 2000. 14(11): p. F111-5.
 38. Lipshultz SE, Easley KA, Orav EJ, et al. Absence of cardiac toxicity of zidovudine in infants. Pediatric Pulmonary and Cardiac Complications of Vertically Transmitted HIV Infection Study Group. *N Engl J Med*, 2000. 343(11): p. 759-66.
 39. Cunningham CK, Chaix M-L, Rekacewica C, et al., for the PACTG 316 Team. Development of resistance

- mutations in women on standard antiretroviral therapy who received intrapartum nevirapine to prevent perinatal HIV-1 transmission: a substudy of Pediatric AIDS Clinical Trials Group Protocol 316. *J Infect Dis*, 2002 (in press).
40. Nightingale SL. From the Food and Drug Administration. *JAMA*, 1998. 280(17): p. 1472.
 41. De Santis M, Carducci B, De Santis L, et al. Periconceptional exposure to efavirenz and neural tube defects. *Arch Intern Med*, 2002. 162(3): p. 355.
 42. Fundaro C, Genovese O, Rendeli C, et al. Myelomeningocele in a child with intrauterine exposure to efavirenz. *AIDS*, 2002. 16(2): p. 299-300.
 43. Mirochnick M, Fenton T, Gagnier P, et al. Pharmacokinetics of nevirapine in human immunodeficiency virus type 1-infected pregnant women and their neonates. Pediatric AIDS Clinical Trials Group Protocol 250 Team. *J Infect Dis*, 1998. 178(2): p. 368-74.
 44. Guay LA, Musoke P, Fleming T, et al. Intrapartum and neonatal single-dose nevirapine compared with zidovudine for prevention of mother-to-child transmission of HIV-1 in Kampala, Uganda: HIVNET 012 randomised trial. *Lancet*, 1999. 354(9181): p. 795-802.
 45. Jackson JB, Mrcanz M, Guay LA, et al. Selection of nevirapine (NVP) resistance mutations in Ugandan women and infants receiving NVP prophylaxis to prevent HIV-1 vertical transmission (HIVNET 012). XIII International AIDS Conference. Durban, South Africa. July 9-14, 2000. (Abstract LbOr13).
 46. Centers for Disease Control and Prevention. Serious adverse events attributed to nevirapine regimens for postexposure prophylaxis after HIV exposures worldwide, 1997-2000. *MMWR*, 2001. 49(51-52): p. 1153-6.
 47. FDA. Food and Drug Administration Advisory. Reports of diabetes and hyperglycemia in patients receiving protease inhibitors for treatment of human immunodeficiency virus (HIV). 1997.
 48. Visnegarwala F, Krause KL, Musher DM. Severe diabetes associated with protease inhibitor therapy [letter]. *Ann Intern Med*, 1997. 127(10): p. 947.
 49. Eastone JA, Decker CF. New-onset diabetes mellitus associated with use of protease inhibitor. *Ann Intern Med*, 1997. 127(10): p. 948.
 50. Dube MP. Metabolic complications of antiretroviral therapies. *AIDS Clinical Care*, 1998. 10(6): p. 41-4.
 51. Lorenzi P, Spicher VM, Laubereau B, et al. Antiretroviral therapies in pregnancy: maternal, fetal and neonatal effects. Swiss HIV Cohort Study, the Swiss Collaborative HIV and Pregnancy Study, and the Swiss Neonatal HIV Study. *AIDS*, 1998. 12(18): p. F241-7.
 52. Martin R, Boyer P, Hammill H, et al. Incidence of premature birth and neonatal respiratory disease in infants of HIV-positive mothers. The Pediatric Pulmonary and Cardiovascular Complications of Vertically Transmitted Human Immunodeficiency Virus Infection Study Group. *J Pediatr*, 1997. 131(6): p. 851-6.
 53. Leroy V, Ladner J, Nyiraziraje M, et al. Effect of HIV-1 infection on pregnancy outcome in women in Kigali, Rwanda, 1992-1994. Pregnancy and HIV Study Group. *AIDS*, 1998. 12(6): p. 643-50.
 54. Brocklehurst P, French R. The association between maternal HIV infection and perinatal outcome: a systematic review of the literature and meta-analysis. *Br J Obstet Gynaecol*, 1998. 105(8): p. 836-48.
 55. Wara D, Tuomala R, Bryson Y. PACTG 358 - safety, pharmacokinetics and antiretroviral activity of indinavir, zidovudine (ZDV), and lamivudine (3TC) in HIV-1 seropositive pregnant women and infants. 2nd Conference on Global Strategies for the Prevention of HIV Transmission from Mothers to Infants. Montreal, Canada. 1999. (Abstract #447).
 56. Hayashi S, Beckerman K, Homma M, et al. Pharmacokinetics of indinavir in HIV-positive pregnant women [letter]. *AIDS*, 2000. 14(8): p. 1061-2.
 57. Casey BM, Bawdon RE. Placental transfer of ritonavir with zidovudine in the ex vivo placental perfusion model. *Am J Obstet Gynecol*, 1998. 179(3 Pt 1): p. 758-61.
 58. Chaube S, Murphy ML. The effects of hydroxyurea and related compounds on the rat fetus. *Cancer Res*, 1966. 26(7): p. 1448-57.
 59. Ferm VH. Severe developmental malformations. *Arch Pathol*, 1966. 81: p. 174-7.
 60. Khara KS. A teratogenicity study on hydroxyurea and diphenylhydantoin in cats. *Teratology*, 1979. 20(3): p. 447-52.
 61. Wilson JG, Scott WJ, Ritter EJ, Fradkin R. Comparative distribution and embryotoxicity of hydroxyurea in pregnant rats and rhesus monkeys. *Teratology*, 1975. 11(2): p. 169-78.
 62. Aliverti V, Bonanomi L, Giavini E. Hydroxyurea as a reference standard in teratological screening. Comparison of the embryotoxic and teratogenic effects following single intraperitoneal or repeated oral administrations to pregnant rats. *Arch Toxicol Suppl*, 1980. 4: p. 239-47.
 63. DeSesso JM, Jordan RL. Drug-induced limb dysplasias in fetal rabbits. *Teratology*, 1977. 15(2): p. 199-211.
 64. Fritz H, Hess R. Effects of hydroxyurea on postnatal growth and behaviour of rats. *Agents Actions*, 1980. 10(4): p. 389-93.
 65. Sylvester RK, Lobell M, Teresi ME, et al. Excretion of hydroxyurea into milk. *Cancer*, 1987. 60(9): p. 2177-8.
 66. Diav-Citrin O, Hunnisett L, Sher GD, Koren G. Hydroxyurea use during pregnancy: a case report in sickle cell disease and review of the literature. *Am J Hematol*, 1999. 60(2): p. 148-50.