

Table X-1

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1975-2000

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000
All Areas ^b	59	55	55	153	588	1241	1876	2234	2221	1947	1405	456	371
All Areas but SF-Oakland	45	41	39	56	158	349	586	848	910	925	738	270	232
San Francisco-Oakland SMSA	14	14	16	97	430	892	1290	1386	1311	1022	667	186	139
San Francisco City/County	7	7	8	75	366	755	1063	1067	1001	749	485	118	82

AGE-ADJUSTED CANCER INCIDENCE RATES^a

By SEER Location and Year of Diagnosis, 1975-2000

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000
All Areas ^b	0.5	0.4	0.4	0.9	2.9	5.7	8.3	9.5	9.0	7.6	5.4	1.8	1.4
All Areas but SF-Oakland	0.5	0.4	0.4	0.5	1.0	2.0	3.1	4.3	4.4	4.3	3.4	1.3	1.1
San Francisco-Oakland SMSA	0.7	0.7	0.8	3.0	12.1	23.7	33.0	34.2	31.7	24.1	15.8	4.4	3.2
San Francisco City/County	1.3	1.2	1.3	10.0	44.7	89.2	122.8	123.8	114.5	83.8	54.3	12.8	8.9

^a SEER 9 areas. Rates are per 100,000 males and are age-adjusted to the 2000 U.S. standard population. Each rate has been age-adjusted by 5-year age groups.

^b All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table X-2

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1975-2000

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000
All Areas ^b	6	6	7	95	508	1160	1752	2106	2073	1835	1286	387	307
All Areas but SF-Oakland	6	5	<4	13	105	302	517	774	843	867	665	224	188
San Francisco-Oakland SMSA	0	<4	6	82	403	858	1235	1332	1230	968	621	163	119
San Francisco City/County	0	<4	5	72	353	729	1026	1034	948	714	454	110	74

AGE-ADJUSTED CANCER INCIDENCE RATES^a

By SEER Location and Year of Diagnosis, 1975-2000

All Races, Males, Ages 20-54

Year of Diagnosis

SEER Location	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000
All Areas ^b	0.1	0.1	0.1	0.9	4.7	10.5	15.1	17.5	16.5	14.0	9.6	2.8	2.2
All Areas but SF-Oakland	0.1	0.1	-	0.1	1.2	3.2	5.2	7.6	7.9	7.8	5.8	2.0	1.6
San Francisco-Oakland SMSA	0.0	-	0.4	4.8	21.9	45.3	62.6	64.8	58.1	44.9	28.7	7.4	5.2
San Francisco City/County	0.0	-	1.5	19.3	85.7	171.2	235.6	238.0	213.0	157.5	99.4	23.4	15.5

^a SEER 9 areas. Rates are per 100,000 males aged 20-54 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.

Table X-3

KAPOSI'S SARCOMANUMBER OF CASES

By SEER Location and Year of Diagnosis, 1975-2000

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000
All Areas ^b	53	49	48	58	78	80	122	127	147	112	119	69	64
All Areas but SF-Oakland	39	36	38	43	52	47	69	74	66	58	73	46	44
San Francisco-Oakland SMSA	14	13	10	15	26	33	53	53	81	54	46	23	20
San Francisco City/County	7	6	<4	<4	13	26	37	32	53	35	31	8	8

AGE-ADJUSTED CANCER INCIDENCE RATES^a

By SEER Location and Year of Diagnosis, 1975-2000

All Races, Males, Ages 55+

Year of Diagnosis

SEER Location	1975-1976	1977-1978	1979-1980	1981-1982	1983-1984	1985-1986	1987-1988	1989-1990	1991-1992	1993-1994	1995-1996	1997-1998	1999-2000
All Areas ^b	2.2	1.9	1.8	2.0	2.5	2.3	3.3	3.4	3.9	2.7	2.9	1.7	1.5
All Areas but SF-Oakland	2.0	1.6	1.7	1.8	1.9	1.8	2.4	2.5	2.2	1.7	2.2	1.4	1.3
San Francisco-Oakland SMSA	3.4	3.2	2.6	2.8	5.4	4.9	7.7	8.1	12.8	8.0	6.7	3.5	2.7
San Francisco City/County	5.9	4.6	-	-	9.1	17.1	23.8	21.9	37.8	24.0	21.7	5.3	5.3


^a SEER 9 areas. Rates are per 100,000 males aged 55+ and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b All Areas refers to all SEER data collection areas. The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi's Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.


Figure X-1

All Cancer Sites vs Kaposi's Sarcoma SEER Incidence Rates, 1975-2000 Males, Ages 20-54




Source: SEER 9 areas. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups.
Regression lines are calculated using the Joinpoint Regression Program.

All SEER excluding
San Francisco/Oakland
Males, Ages 20-54
1975-2000



San Francisco/Oakland
Males, Ages 20-54
1975-2000



San Francisco County/City
Males, Ages 20-54
1975-2000




Figure X-2

Source: SEER 9 areas. Rates are age-adjusted to the 2000 U.S. standard million population by 5-year age groups.

* Kaposi's sarcoma excluded from Colorectal, Lung, and NHL[^].

[^] Non-Hodgkin Lymphoma.