

Table XXIII-1
STOMACH CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2000
 WITH UP TO THREE JOINPOINTS BY RACE AND SEX

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	Male and Female	1975-2000	-1.6 ^c						
All Races	Male	1975-1988	-1.1 ^c	1988-2000	-2.1 ^c				
All Races	Female	1975-2000	-1.7 ^c						
White	Male and Female	1975-2000	-1.9 ^c						
White	Male	1975-2000	-1.9 ^c						
White	Female	1975-2000	-2.2 ^c						
Black	Male and Female	1975-2000	-1.1 ^c						
Black	Male	1975-2000	-1.2 ^c						
Black	Female	1975-2000	-0.8 ^c						
<u>SEER Delay-Adjusted Incidence^a</u>									
White	Male	1975-2000	-1.8 ^c						
White	Female	1975-2000	-2.1 ^c						
Black	Male	1975-2000	-1.1 ^c						
Black	Female	1975-2000	-0.8 ^c						
<u>U.S. Cancer Mortality^b</u>									
All Races	Male and Female	1975-1987	-2.5 ^c	1987-1990	-0.4	1990-2000	-2.9 ^c		
All Races	Male	1975-1987	-2.4 ^c	1987-1990	-0.4	1990-2000	-3.3 ^c		
All Races	Female	1975-1987	-2.8 ^c	1987-1990	-0.6	1990-2000	-2.5 ^c		
White	Male and Female	1975-1987	-2.8 ^c	1987-1990	-0.8	1990-2000	-3.2 ^c		
White	Male	1975-1994	-2.3 ^c	1994-2000	-3.9 ^c				
White	Female	1975-1977	-4.7	1977-2000	-2.6 ^c				
Black	Male and Female	1975-1987	-1.6 ^c	1987-1990	0.9	1990-2000	-2.5 ^c		
Black	Male	1975-1991	-1.1 ^c	1991-2000	-2.6 ^c				
Black	Female	1975-2000	-1.2 ^c						

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups using the Joinpoint Regression Program.

^a Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file.

^c The APC is significantly different from zero (p<.05).

Table XXIII-2
STOMACH CANCER (Invasive)

SEER INCIDENCE RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES^a, BY RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DIAGNOSIS:</u>									
1975	11.7	17.1	7.8	10.7	16.0	7.0	17.0	24.8	11.5
1976	12.2	17.5	8.3	11.2	15.9	7.7	15.8	23.5	10.1
1977	11.5	16.6	7.9	10.3	15.0	7.0	17.3	24.6	11.7
1978	11.5	16.5	7.9	10.3	15.0	7.0	17.6	22.9	13.5
1979	12.0	17.4	8.2	10.7	15.6	7.5	18.5	28.2	11.7
1980	11.3	17.1	7.2	10.3	15.7	6.6	15.1	26.3	7.3
1981	11.1	16.6	7.4	10.1	15.1	6.7	17.6	27.9	10.3
1982	10.9	16.3	7.1	9.7	14.6	6.3	19.0	31.2	10.3
1983	10.9	15.8	7.4	9.5	13.7	6.6	18.0	27.8	11.2
1984	10.6	15.3	7.3	9.4	13.9	6.4	16.4	21.7	12.3
1985	10.2	15.1	6.9	8.9	13.3	6.0	16.4	23.5	11.4
1986	10.2	15.0	6.8	9.0	13.5	5.9	16.5	23.4	11.9
1987	10.2	15.1	6.8	9.0	13.3	6.0	17.0	26.5	10.7
1988	10.2	15.6	6.5	8.8	13.7	5.5	16.3	24.9	10.1
1989	10.0	15.2	6.3	8.8	13.6	5.3	15.8	23.2	10.7
1990	9.3	13.8	6.2	8.0	12.0	5.2	14.5	21.9	9.4
1991	9.7	14.2	6.6	8.3	12.4	5.5	16.5	25.7	10.5
1992	9.2	13.4	6.1	8.0	11.7	5.2	14.0	20.0	9.7
1993	9.0	13.7	5.8	7.7	11.6	4.9	14.4	23.8	8.3
1994	9.1	13.7	5.7	7.8	11.9	4.8	15.1	24.9	8.8
1995	8.4	12.6	5.2	7.3	11.3	4.4	13.4	18.8	9.6
1996	8.5	12.6	5.5	7.3	11.0	4.6	14.7	22.9	9.0
1997	8.6	12.7	5.6	7.2	10.8	4.5	15.6	21.9	11.1
1998	8.5	12.1	5.9	7.1	10.5	4.7	14.8	20.0	11.4
1999	8.5	12.0	5.9	7.3	10.5	5.0	12.7	17.7	9.5
2000	8.0	11.6	5.3	6.7	9.7	4.3	12.8	19.9	8.1
<u>AGE-ADJUSTED RATES, 1996-2000</u>									
<u>AGE AT DIAGNOSIS:</u>									
All ages	8.4	12.2	5.7	7.1	10.5	4.6	14.1	20.5	9.8
Under 65	2.9	4.0	1.9	2.4	3.4	1.5	4.8	6.8	3.2
65 and over	46.6	69.0	31.9	39.6	59.6	26.4	78.2	114.7	55.5
All ages (world std.) ^b	5.1	7.3	3.3	4.3	6.2	2.7	8.4	12.4	5.5
<u>AGE-SPECIFIC RATES, 1996-2000</u>									
<u>AGE AT DIAGNOSIS:</u>									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-
20-24	-	-	-	-	-	-	-	-	-
25-29	0.4	0.5	-	-	-	-	-	-	-
30-34	1.0	1.0	0.9	0.7	0.7	0.7	-	-	-
35-39	1.4	1.7	1.1	0.9	1.2	0.7	3.0	-	-
40-44	2.6	3.1	2.0	2.2	2.6	1.7	4.3	5.6	-
45-49	4.3	5.7	2.9	3.3	4.4	2.2	7.8	11.1	4.9
50-54	7.0	9.7	4.4	6.1	8.8	3.4	12.4	16.1	9.2
55-59	12.1	17.4	7.1	10.6	15.3	6.0	16.2	23.3	10.2
60-64	19.6	28.7	11.3	16.4	24.2	9.2	33.4	51.9	18.6
65-69	27.9	40.8	17.0	23.7	35.1	13.6	42.3	69.1	22.1
70-74	41.5	60.8	26.5	34.6	51.2	21.5	68.2	104.4	43.8
75-79	51.1	75.7	33.9	43.8	66.2	28.2	79.8	127.0	50.2
80-84	62.0	90.4	45.2	52.9	78.0	38.1	110.4	155.4	87.5
85+	72.9	111.8	56.9	62.2	98.3	47.9	137.5	168.0	126.0

^a SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^b SEER 9 areas. Rates are per 100,000 and are age-adjusted to the world standard population.

- Statistic not shown. Rate based on less than 25 cases for the time interval.

Table XXIII-3
STOMACH CANCER (Invasive)

U.S. DEATH RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES^a, BY RACE AND SEX

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>AGE-ADJUSTED RATES</u>									
<u>YEAR OF DEATH:</u>									
1975	8.5	12.3	5.9	7.9	11.4	5.6	13.9	21.0	8.6
1976	8.3	12.0	5.7	7.7	11.2	5.4	13.5	20.2	8.6
1977	7.9	11.5	5.4	7.3	10.7	5.0	13.4	19.5	9.0
1978	7.7	11.0	5.5	7.2	10.2	5.1	12.6	18.7	8.3
1979	7.6	11.0	5.2	7.0	10.2	4.8	12.9	18.7	8.7
1980	7.4	10.6	5.1	6.8	9.8	4.8	12.2	18.3	8.0
1981	7.3	10.6	5.0	6.7	9.8	4.6	12.5	18.1	8.6
1982	7.0	10.3	4.9	6.5	9.4	4.5	12.2	19.2	7.4
1983	6.8	9.9	4.8	6.2	9.0	4.4	12.3	18.4	8.1
1984	6.8	9.9	4.7	6.2	9.1	4.3	11.9	17.9	7.9
1985	6.5	9.5	4.4	6.0	8.8	4.1	11.5	17.0	7.8
1986	6.4	9.4	4.2	5.8	8.6	3.8	11.4	17.4	7.4
1987	6.2	9.1	4.2	5.6	8.3	3.8	11.3	17.0	7.5
1988	6.1	8.8	4.2	5.5	7.9	3.8	11.0	17.0	7.1
1989	6.2	9.2	4.2	5.6	8.3	3.8	11.5	17.7	7.3
1990	6.1	8.9	4.2	5.4	8.0	3.7	11.5	17.2	7.7
1991	6.0	8.9	4.0	5.4	8.0	3.6	11.5	17.6	7.5
1992	5.6	8.1	3.9	5.0	7.3	3.4	10.7	15.6	7.5
1993	5.6	8.2	3.8	5.0	7.3	3.4	10.7	16.3	7.1
1994	5.4	7.9	3.7	4.8	7.0	3.3	10.3	15.4	7.0
1995	5.3	7.7	3.7	4.7	6.9	3.2	10.4	14.9	7.4
1996	5.1	7.4	3.5	4.5	6.6	3.1	10.0	15.1	6.8
1997	4.9	7.2	3.4	4.3	6.3	2.9	9.7	14.6	6.5
1998	4.8	6.8	3.4	4.3	6.1	3.0	9.3	13.7	6.4
1999	4.6	6.6	3.3	4.0	5.8	2.8	9.2	13.7	6.5
2000	4.6	6.4	3.2	4.0	5.7	2.8	9.0	12.8	6.5
<u>AGE-ADJUSTED RATES, 1996-2000</u>									
<u>AGE AT DEATH:</u>									
All ages	4.8	6.9	3.4	4.2	6.1	2.9	9.4	14.0	6.5
Under 65	1.5	2.1	1.0	1.3	1.8	0.8	3.0	4.3	1.9
65 and over	27.5	40.1	19.5	24.6	35.9	17.3	54.0	80.7	38.8
All ages (world std.) ^b	2.8	3.9	1.9	2.4	3.4	1.6	5.4	8.0	3.6
<u>AGE-SPECIFIC RATES, 1996-2000</u>									
<u>AGE AT DEATH:</u>									
<1	-	-	-	-	-	-	-	-	-
1-4	-	-	-	-	-	-	-	-	-
5-9	-	-	-	-	-	-	-	-	-
10-14	-	-	-	-	-	-	-	-	-
15-19	-	-	-	-	-	-	-	-	-
20-24	0.1	0.1	0.1	0.0	-	-	-	-	-
25-29	0.2	0.2	0.2	0.2	0.2	0.1	0.5	0.4	0.6
30-34	0.4	0.5	0.4	0.3	0.4	0.3	0.8	0.9	0.7
35-39	0.8	0.9	0.7	0.6	0.7	0.5	1.6	2.0	1.3
40-44	1.4	1.7	1.1	1.1	1.4	0.9	2.7	3.7	1.9
45-49	2.3	3.0	1.6	1.9	2.5	1.4	4.6	6.7	2.7
50-54	3.8	5.3	2.5	3.3	4.6	2.0	7.7	10.9	5.0
55-59	6.3	9.0	3.8	5.5	7.8	3.2	12.3	18.4	7.4
60-64	10.0	14.4	6.1	8.6	12.3	5.2	19.1	29.8	10.8
65-69	14.8	21.3	9.3	13.1	19.0	8.0	27.4	41.4	17.1
70-74	21.1	30.6	13.8	18.7	27.2	12.0	41.8	64.3	27.0
75-79	29.1	42.7	19.7	25.7	37.9	17.3	59.3	89.6	41.0
80-84	38.6	56.7	28.1	34.8	51.1	25.3	74.5	117.1	53.6
85+	53.3	77.1	43.8	48.0	70.4	39.2	104.8	143.5	90.2

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^b NCHS public use data file. Rates are per 100,000 and are age-adjusted to the world standard population.

- Statistic not shown. Rate based on less than 25 cases for the time interval.

Table XXIII-4
STOMACH CANCER (Invasive)

SURVIVAL RATES, BY RACE, SEX, DIAGNOSIS YEAR, STAGE AND AGE

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-1963 ^a	-	-	-	11	10	13	8	5	14
1970-1973 ^a	-	-	-	13	12	14	13	15	10
1974-1976 ^b	15.4	14.2	17.3	14.7	13.5	16.5	16.5	15.9	17.6
1977-1979 ^b	16.4	15.0	18.6	15.7	14.1	18.2	15.0	14.9	15.2
1980-1982 ^b	17.7	16.6	19.5	16.5	15.6	17.9	19.1	18.2	21.2
1983-1985 ^b	17.4	15.9	19.7	16.3	14.6	18.9	18.9	18.4	19.6
1986-1988 ^b	20.2	17.2	25.2	19.1	16.2	24.2	19.3	15.3	25.7
1989-1991 ^b	21.1	17.6	26.8	18.4	15.1	24.1	24.8	22.3	28.8
1992-1999 ^b	22.5 ^c	21.3 ^c	24.6 ^c	21.4 ^c	20.0 ^c	23.8 ^c	20.5	19.4	22.3
1992-1999 ^b									
ALL STAGES	22.5	21.3	24.6	21.4	20.0	23.8	20.5	19.4	22.3
LOCALIZED	59.0	58.5	59.8	55.4	54.3	57.1	56.1	54.4 ^d	58.8 ^d
REGIONAL	21.7	20.8	23.5	21.1	20.1	23.1	19.8	20.9	18.2
DISTANT	2.5	2.6	2.5	2.5	2.5	2.7	2.0	1.9	2.2
UNSTAGED	13.2	11.8	14.4	12.9	11.7	13.8	15.3	12.6	17.3 ^d
<u>STAGE DISTRIBUTION (%) 1992-1999^b</u>									
ALL STAGES									
Number of cases	13,540	8,402	5,138	9,616	6,069	3,547	1,713	1,018	695
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	23	21	24	23	21	25	22	20	23
REGIONAL	32	34	29	31	33	28	30	31	29
DISTANT	32	34	30	32	34	30	34	37	30
UNSTAGED	14	12	17	14	12	18	15	12	18
<u>5-YR RELATIVE SURVIVAL RATES, 1992-1999^b</u>									
AGE AT DIAGNOSIS:									
<45	27.4	24.3	32.1	28.8	24.0	37.6	27.4	25.6	30.0 ^d
45-54	22.5	19.8	29.5	21.8	18.9	30.0	21.5	16.9	32.9 ^d
55-64	23.1	22.3	24.9	21.4	20.4	23.9	23.0	23.6	21.8
65-74	23.5	22.3	25.9	21.9	21.2	23.4	21.7	19.5	25.5
75+	20.0	19.1	21.0	19.5	18.0	20.9	12.6	10.7	14.1
Under 65	23.8	21.9	27.8	22.7	20.5	28.2	24.1	22.2	28.1
65 and over	21.8	20.9	23.0	20.7	19.8	22.0	17.5	16.3	18.9

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.
^b Rates are from the SEER 9 areas. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients into 2000.
^c The difference in rates between 1974-1976 and 1992-1999 is statistically significant (p<.05).
^d The standard error of the survival rate is between 5 and 10 percentage points.
^e The standard error of the survival rate is greater than 10 percentage points.
- Statistic could not be calculated.

Table XXIII-5
STOMACH CANCER (Invasive)

INCIDENCE, DEATH, AND SURVIVAL RATES^c

By Year of Diagnosis/Death

All Races, Males and Females

Year of Diagnosis/Death

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
Incidence																		
SEER ^a	11.8	10.9	10.2	10.2	10.2	10.2	10.0	9.3	9.7	9.2	9.0	9.1	8.4	8.5	8.6	8.5	8.5	8.0
Mortality																		
U.S. ^b	8.0	7.0	6.5	6.4	6.2	6.1	6.2	6.1	6.0	5.6	5.6	5.4	5.3	5.1	4.9	4.8	4.6	4.6
SEER ^a	8.7	7.7	7.2	6.8	6.6	6.5	6.4	6.5	6.1	6.1	5.8	5.8	5.5	5.1	5.2	5.0	5.1	5.0
Relative Survival Rates (SEER ^a)																		
1-year	39.8	42.9	43.6	45.1	43.6	47.0	46.4	46.4	44.6	47.0	45.3	46.3	47.2	47.1	46.5	49.4	47.5	
2-year	25.6	27.5	28.6	29.1	27.4	32.6	31.7	32.4	30.6	32.4	30.8	33.6	32.2	32.7	34.0	34.4		
3-year	20.3	21.8	22.3	23.5	21.8	26.4	25.5	25.3	25.2	26.2	24.4	27.0	27.2	27.6	27.3			
4-year	17.9	19.0	19.6	21.4	19.4	22.6	22.5	22.8	22.7	23.8	21.4	24.5	24.0	24.3				
5-year	16.3	17.4	18.0	20.2	18.6	21.7	21.3	20.6	21.4	22.6	20.0	22.7	22.7					
6-year	15.3	16.7	17.1	19.3	17.8	20.5	20.3	18.9	20.0	21.6	19.2	20.9						
7-year	14.9	16.0	16.9	19.2	16.2	19.8	19.0	18.3	19.6	20.2	18.1							
8-year	14.3	15.4	16.1	18.8	15.7	19.5	18.6	17.9	19.1	19.2								
9-year	14.0	15.1	15.9	18.2	14.9	19.4	18.1	17.0	18.4									
10-year	13.7	15.0	15.2	17.7	14.1	18.7	17.4	16.4										
11-year	13.2	14.5	15.1	17.1	13.5	18.6	17.4											
12-year	13.1	14.1	15.1	17.1	13.2	18.3												
13-year	12.9	14.1	15.1	16.7	12.9													
14-year	12.5	14.1	15.1	16.4														
15-year	12.2	13.8	15.1															
16-year	11.9	13.5																
17-year	11.8																	
18-year	11.4																	
19-year	11.4																	
20-year	11.1																	

^a Rates are from the SEER 9 areas.

^b Rates are from the NCHS public use data file.

^c Incidence and death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups. Survival rates are relative rates expressed as percents.

Table XXIII-6
STOMACH CANCER (Invasive)

INCIDENCE, DEATH, AND SURVIVAL RATES^c

By Year of Diagnosis/Death

All Races, Males

Year of Diagnosis/Death

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	
Incidence																			
SEER ^a	17.0	16.2	15.1	15.0	15.1	15.6	15.2	13.8	14.2	13.4	13.7	13.7	12.6	12.6	12.7	12.1	12.0	11.6	
Mortality																			
U.S. ^b	11.5	10.2	9.5	9.4	9.1	8.8	9.2	8.9	8.9	8.1	8.2	7.9	7.7	7.4	7.2	6.8	6.6	6.4	
SEER ^a	12.7	11.2	10.9	10.3	9.8	9.8	9.7	9.9	9.2	8.9	8.7	8.5	8.1	7.7	7.8	7.0	6.8	6.9	
Relative Survival Rates (SEER ^a)																			
1-year	39.3	42.5	42.9	42.4	42.4	46.0	47.0	46.2	42.5	45.4	44.2	46.3	48.0	45.1	47.0	50.7	47.2		
2-year	24.7	26.7	27.7	26.8	24.8	30.4	30.6	31.3	26.9	30.2	28.6	32.7	32.2	29.4	32.8	35.8			
3-year	19.2	21.0	21.0	21.2	19.4	23.9	23.3	24.0	21.3	24.3	23.1	26.1	26.4	24.6	25.8				
4-year	16.7	17.9	18.4	18.8	16.7	19.7	19.4	20.9	18.4	22.0	20.5	23.8	23.2	21.8					
5-year	15.2	16.3	16.1	17.6	15.6	18.4	18.1	18.3	16.4	20.8	18.9	21.8	21.6						
6-year	14.2	15.4	15.1	16.7	15.3	17.6	17.1	17.2	14.6	19.4	17.8	19.8							
7-year	13.7	14.7	14.5	16.4	13.6	16.8	15.7	16.5	13.7	18.4	16.2								
8-year	13.3	14.2	13.8	16.3	12.7	16.6	15.2	16.0	13.0	17.4									
9-year	13.2	14.0	13.6	15.8	12.4	16.2	14.4	15.2	12.5										
10-year	12.8	14.0	12.6	15.6	11.9	16.1	13.9	14.4											
11-year	12.5	13.5	12.2	15.2	11.0	16.0	13.8												
12-year	12.4	12.9	12.2	15.2	11.0	15.9													
13-year	12.4	12.9	12.2	15.2	10.9														
14-year	11.7	12.9	12.2	15.2															
15-year	11.3	12.8	12.1																
16-year	11.2	12.3																	
17-year	11.0																		
18-year	10.3																		
19-year	10.3																		
20-year	10.3																		

^a Rates are from the SEER 9 areas.

^b Rates are from the NCHS public use data file.

^c Incidence and death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups. Survival rates are relative rates expressed as percents.

Table XXIII-7
STOMACH CANCER (Invasive)

INCIDENCE, DEATH, AND SURVIVAL RATES^c

By Year of Diagnosis/Death

All Races, Females

Year of Diagnosis/Death

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	
Incidence																			
SEER ^a	8.0	7.3	6.9	6.8	6.8	6.5	6.3	6.2	6.6	6.1	5.8	5.7	5.2	5.5	5.6	5.9	5.9	5.3	
Mortality																			
U.S. ^b	5.6	4.9	4.4	4.2	4.2	4.2	4.2	4.2	4.0	3.9	3.8	3.7	3.7	3.5	3.4	3.4	3.3	3.2	
SEER ^a	6.0	5.3	4.7	4.5	4.4	4.2	4.2	4.2	4.1	4.2	3.8	3.8	3.7	3.4	3.4	3.5	3.8	3.7	
Relative Survival Rates (SEER ^a)																			
1-year	40.6	43.6	44.7	49.5	45.7	48.7	45.4	46.7	48.0	49.7	47.2	46.4	45.8	50.3	45.7	47.3	47.8		
2-year	27.0	28.9	30.0	32.8	31.6	36.5	33.6	34.1	36.3	36.1	34.6	35.1	32.1	38.1	36.0	32.2			
3-year	22.0	23.2	24.3	27.3	25.7	30.6	29.3	27.3	31.3	29.3	26.7	28.7	28.7	32.5	29.8				
4-year	19.6	20.8	21.5	25.7	23.8	27.7	28.0	25.8	29.4	26.8	22.8	25.8	25.3	28.4					
5-year	17.9	19.3	21.0	24.7	23.5	27.3	26.9	24.2	29.2	25.5	21.9	24.4	24.7						
6-year	17.1	18.7	20.3	23.7	21.9	25.7	25.8	21.6	28.2	25.2	21.7	22.9							
7-year	16.7	18.1	20.3	23.7	20.4	25.1	24.8	21.1	28.2	23.2	21.4								
8-year	15.8	17.3	19.8	23.0	20.4	24.4	24.6	21.1	28.2	22.3									
9-year	15.1	16.9	19.5	22.2	18.9	24.4	24.5	19.9	27.4										
10-year	15.0	16.7	19.3	21.2	17.7	23.4	23.6	19.4											
11-year	14.4	16.2	19.3	20.1	17.3	23.1	23.6												
12-year	14.1	16.1	19.3	19.9	16.6	22.4													
13-year	13.7	16.0	19.3	18.7	16.1														
14-year	13.6	16.0	19.3	18.2															
15-year	13.4	15.5	19.3																
16-year	13.1	15.3																	
17-year	12.9																		
18-year	12.9																		
19-year	12.7																		
20-year	11.9																		

^a Rates are from the SEER 9 areas.

^b Rates are from the NCHS public use data file.

^c Incidence and death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups. Survival rates are relative rates expressed as percents.

Table XXIII-8
STOMACH CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
 Given Cancer Free At Current Age By Race and Sex

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
 Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1998-2000

All Races

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.25	0.00	0.00	0.00	0.79
10	0.00	0.00	0.02	1.27	0.00	0.00	0.01	0.81
20	0.00	0.02	0.06	1.28	0.00	0.01	0.04	0.81
30	0.02	0.06	0.19	1.29	0.01	0.04	0.10	0.81
40	0.04	0.17	0.47	1.30	0.03	0.09	0.23	0.80
50	0.13	0.44	0.88	1.30	0.06	0.21	0.45	0.79
60	0.33	0.81	1.16	1.25	0.15	0.40	0.66	0.76

Lifetime Risk of Being Diagnosed = 1.25%
 Lifetime Risk of Dying = 0.61%

Lifetime Risk of Being Diagnosed = 0.79%
 Lifetime Risk of Dying = 0.41%

Whites

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.00	1.09	0.00	0.00	0.00	0.66
10	0.00	0.00	0.02	1.10	0.00	0.00	0.01	0.66
20	0.00	0.02	0.05	1.11	0.00	0.01	0.04	0.66
30	0.01	0.05	0.16	1.12	0.01	0.03	0.08	0.67
40	0.04	0.15	0.41	1.12	0.03	0.08	0.19	0.66
50	0.12	0.39	0.76	1.12	0.05	0.17	0.36	0.65
60	0.29	0.69	1.00	1.07	0.12	0.33	0.54	0.62

Lifetime Risk of Being Diagnosed = 1.09%
 Lifetime Risk of Dying = 0.56%

Lifetime Risk of Being Diagnosed = 0.66%
 Lifetime Risk of Dying = 0.37%

Blacks

Current Age	Males				Females			
	+10 yrs	+20 yrs	+30 yrs	Eventually	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.32	0.00	0.00	0.00	1.03
10	0.00	0.01	0.03	1.37	0.00	0.00	0.02	1.06
20	0.01	0.03	0.09	1.38	0.00	0.02	0.05	1.07
30	0.02	0.09	0.24	1.41	0.01	0.05	0.13	1.07
40	0.07	0.23	0.56	1.43	0.04	0.12	0.30	1.08
50	0.17	0.53	1.05	1.47	0.09	0.27	0.58	1.08
60	0.42	1.03	1.38	1.51	0.20	0.53	0.89	1.08

Lifetime Risk of Being Diagnosed = 1.32%
 Lifetime Risk of Dying = 0.92%

Lifetime Risk of Being Diagnosed = 1.03%
 Lifetime Risk of Dying = 0.69%

Table XXIII-9
STOMACH CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1996-2000			Trend 1992-2000		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	9.1	13.0	6.2	-1.3 ^d	-1.9 ^d	-0.5
White	7.7	11.2	5.1	-1.5 ^d	-2.1 ^d	-0.8
White Hispanic ^b	13.8	18.7	10.3	-1.7 ^d	-2.2	-1.1
White Non-Hispanic ^b	6.8	10.0	4.3	-1.9 ^d	-2.4 ^d	-1.4
Black	14.0	19.9	9.9	-1.0	-2.0	0.5
Asian/Pacific Islander	17.3	23.0	12.8	-2.8 ^d	-2.9 ^d	-2.6 ^d
Amer Ind/Alaska Nat	11.0	14.4	8.3	3.4	-	-
Hispanic ^b	13.3	18.1	10.0	-1.6 ^d	-2.1	-1.0

Mortality^c

RACE/ETHNICITY	Rate 1996-2000			Trend 1992-2000		
	Rate per 100,000 persons			APC (%)		
	Total	Males	Females	Total	Males	Females
All Races	4.8	6.9	3.4	-2.8 ^d	-3.2 ^d	-2.5 ^d
White	4.2	6.1	2.9	-3.1 ^d	-3.5 ^d	-2.8 ^d
White Hispanic ^b	7.4	10.1	5.5	-2.0 ^d	-2.1 ^d	-1.8
White Non-Hispanic ^b	3.9	5.7	2.7	-3.2 ^d	-3.4 ^d	-3.1 ^d
Black	9.4	14.0	6.5	-2.3 ^d	-2.5 ^d	-1.9 ^d
Asian/Pacific Islander	9.6	12.5	7.4	-3.2 ^d	-3.2 ^d	-3.0 ^d
Amer Ind/Alaska Nat	5.4	7.0	4.2	-1.6	-2.3	-0.6
Hispanic ^b	7.2	9.9	5.3	-1.9 ^d	-2.0 ^d	-1.8

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Oklahoma, New York, and New Hampshire.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table XXIII-10
STOMACH CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES^a AND SEER DEATH RATES^b
 By Registry, Race and Sex

	All Races			Whites			Blacks		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
<u>SEER INCIDENCE RATES^a, 1996-2000</u>									
REGISTRY									
12 registries	9.1	13.0	6.2	7.7	11.2	5.1	14.0	19.9	9.9
11 registries	9.1	13.0	6.2	7.7	11.2	5.1	14.0	19.9	9.9
9 registries	8.4	12.2	5.7	7.1	10.5	4.6	14.1	20.5	9.8
San Francisco-Oakland	9.4	13.2	6.7	7.8	11.1	5.3	14.7	22.6	9.5
Connecticut	9.0	13.2	5.9	8.6	12.6	5.7	15.4	24.1	10.5
Metropolitan Detroit	9.9	14.5	6.7	8.5	12.7	5.7	15.0	21.4	10.5
Hawaii	15.6	21.2	10.8	8.7	11.1	6.5	-	-	-
Iowa	5.7	9.0	3.2	5.5	8.7	3.1	-	-	-
New Mexico	7.8	10.6	5.7	7.4	10.3	5.3	-	-	-
Seattle-Puget Sound	7.4	11.0	4.6	6.5	10.0	3.8	14.6	19.0	-
Utah	5.6	7.5	4.2	5.5	7.3	4.1	-	-	-
Metropolitan Atlanta	7.3	10.5	5.2	5.3	8.0	3.6	12.1	17.0	9.0
San Jose-Monterey	8.8	12.4	6.0	7.6	10.8	5.0	-	-	-
Los Angeles	11.3	15.8	8.0	9.7	13.8	6.8	13.8	19.0	10.2
<u>SEER DEATH RATES^b, 1996-2000</u>									
REGISTRY									
12 registries	5.5	7.8	3.8	4.6	6.6	3.2	8.9	12.8	6.2
11 registries	5.5	7.8	3.8	4.6	6.6	3.2	8.9	12.8	6.2
9 registries	5.1	7.2	3.5	4.3	6.2	2.9	8.8	12.7	6.2
San Francisco-Oakland	6.0	8.4	4.3	4.9	6.9	3.4	11.5	18.3	6.9
Connecticut	5.4	7.7	3.7	5.2	7.3	3.6	10.0	15.8	6.6
Metropolitan Detroit	5.5	7.7	4.0	4.8	6.9	3.3	8.2	10.9	6.3
Hawaii	9.6	12.7	7.0	5.3	6.3	4.2	-	-	-
Iowa	3.1	4.9	1.9	3.0	4.7	1.8	-	-	-
New Mexico	5.4	7.3	4.0	5.0	6.8	3.6	-	-	-
Seattle-Puget Sound	4.3	6.2	2.8	3.8	5.9	2.3	8.3	-	-
Utah	3.8	4.8	3.0	3.6	4.5	2.9	-	-	-
Metropolitan Atlanta	4.6	6.9	3.2	3.4	5.6	2.2	7.6	10.7	5.7
San Jose-Monterey	5.6	8.2	3.8	4.5	6.5	3.1	-	-	-
Los Angeles	6.8	9.5	4.8	5.8	8.2	4.2	8.9	13.0	6.3

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.

^a SEER 9 areas. Incidence rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b NCHS public use data file. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

- Statistic could not be calculated or populations are inadequate.

Table XXIII-11
STOMACH CANCER (Invasive)AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, 1996-2000

All Races, Males and Females

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
TOTAL U.S.	4.8	0.02							
High Five States									
Hawaii	9.6 ^{bc}	0.40	(01)	100.0					
District of Columbia	8.5 ^{bc}	0.55	(02)	77.1					
Alaska	6.6 ^b	0.68	(03)	37.5					
Rhode Island	6.2 ^{bc}	0.33	(04)	29.2					
New Jersey	6.1 ^{bc}	0.12	(05)	27.1					
Low Five States									
Nebraska	3.4 ^{bc}	0.20	(47)	-29.2					
Montana	3.4 ^{bc}	0.27	(48)	-29.2					
Idaho	3.3 ^{bc}	0.24	(49)	-31.3					
Vermont	3.3 ^{bc}	0.33	(50)	-31.3					
Iowa	3.1 ^{bc}	0.14	(51)	-35.4					
Alabama	4.7	0.15	(19)	-2.1	Montana	3.4 ^{bc}	0.27	(48)	-29.2
Alaska	6.6 ^b	0.68	(03)	37.5	Nebraska	3.4 ^{bc}	0.20	(47)	-29.2
Arizona	3.9 ^{bc}	0.13	(37)	-18.8	Nevada	4.2	0.24	(31)	-12.5
Arkansas	4.5	0.18	(24)	-6.3	New Hampshire	4.1	0.27	(34)	-14.6
California	5.5 ^c	0.06	(08)	14.6	New Jersey	6.1 ^{bc}	0.12	(05)	27.1
Colorado	4.0 ^{bc}	0.16	(36)	-16.7	New Mexico	5.4	0.26	(11)	12.5
Connecticut	5.4	0.17	(12)	12.5	New York	5.9 ^{bc}	0.08	(07)	22.9
Delaware	5.2	0.37	(13)	8.3	North Carolina	4.4	0.11	(28)	-8.3
District of Columbia	8.5 ^{bc}	0.55	(02)	77.1	North Dakota	4.6	0.36	(22)	-4.2
Florida	4.6 ^c	0.07	(23)	-4.2	Ohio	4.5 ^c	0.09	(25)	-6.3
Georgia	4.6	0.12	(21)	-4.2	Oklahoma	3.6 ^{bc}	0.14	(44)	-25.0
Hawaii	9.6 ^{bc}	0.40	(01)	100.0	Oregon	3.8 ^{bc}	0.15	(39)	-20.8
Idaho	3.3 ^{bc}	0.24	(49)	-31.3	Pennsylvania	4.8	0.08	(18)	0.0
Illinois	5.1	0.09	(16)	6.3	Rhode Island	6.2 ^{bc}	0.33	(04)	29.2
Indiana	3.5 ^{bc}	0.11	(45)	-27.1	South Carolina	5.5 ^c	0.17	(10)	14.6
Iowa	3.1 ^{bc}	0.14	(51)	-35.4	South Dakota	3.9 ^b	0.31	(38)	-18.8
Kansas	3.5 ^{bc}	0.16	(46)	-27.1	Tennessee	4.4	0.13	(30)	-8.3
Kentucky	4.4	0.15	(29)	-8.3	Texas	5.1	0.08	(14)	6.3
Louisiana	6.0 ^{bc}	0.17	(06)	25.0	Utah	3.8 ^{bc}	0.23	(40)	-20.8
Maine	4.5	0.25	(26)	-6.3	Vermont	3.3 ^{bc}	0.33	(50)	-31.3
Maryland	5.1	0.15	(15)	6.3	Virginia	4.7	0.12	(20)	-2.1
Massachusetts	5.5 ^c	0.13	(09)	14.6	Washington	4.2 ^c	0.13	(33)	-12.5
Michigan	4.4 ^c	0.10	(27)	-8.3	West Virginia	4.2	0.20	(32)	-12.5
Minnesota	3.7 ^{bc}	0.13	(42)	-22.9	Wisconsin	4.0 ^{bc}	0.12	(35)	-16.7
Mississippi	5.0	0.19	(17)	4.2	Wyoming	3.7 ^b	0.41	(43)	-22.9
Missouri	3.8 ^{bc}	0.11	(41)	-20.8					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table XXIII-12
STOMACH CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, 1996-2000

All Races, Males

State	Rate	SE	Rank	PD
TOTAL U.S.	6.9	0.04		
High Five States				
Hawaii	12.7 ^{bc}	0.69	(01)	84.1
District of Columbia	11.2 ^{bc}	1.04	(02)	62.3
Alaska	9.8 ^b	1.31	(03)	42.0
Rhode Island	9.6 ^{bc}	0.66	(04)	39.1
Louisiana	8.8 ^{bc}	0.34	(05)	27.5
Low Five States				
Utah	4.8 ^{bc}	0.40	(47)	-30.4
Indiana	4.8 ^{bc}	0.20	(48)	-30.4
Nebraska	4.7 ^{bc}	0.36	(49)	-31.9
Idaho	4.7 ^{bc}	0.45	(50)	-31.9
Vermont	4.3 ^{bc}	0.57	(51)	-37.7

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	6.6	0.28	(23)	-4.3	Montana	5.2 ^b	0.53	(42)	-24.6
Alaska	9.8 ^b	1.31	(03)	42.0	Nebraska	4.7 ^{bc}	0.36	(49)	-31.9
Arizona	5.3 ^{bc}	0.23	(40)	-23.2	Nevada	5.5 ^b	0.41	(37)	-20.3
Arkansas	6.1	0.33	(31)	-11.6	New Hampshire	6.2	0.52	(29)	-10.1
California	7.7 ^c	0.12	(10)	11.6	New Jersey	8.6 ^{bc}	0.23	(06)	24.6
Colorado	5.5 ^{bc}	0.28	(39)	-20.3	New Mexico	7.3	0.47	(14)	5.8
Connecticut	7.7	0.33	(11)	11.6	New York	8.4 ^{bc}	0.15	(07)	21.7
Delaware	6.8	0.68	(18)	-1.4	North Carolina	6.3	0.22	(28)	-8.7
District of Columbia	11.2 ^{bc}	1.04	(02)	62.3	North Dakota	6.7	0.66	(21)	-2.9
Florida	6.4	0.12	(26)	-7.2	Ohio	6.7	0.17	(22)	-2.9
Georgia	6.7	0.24	(20)	-2.9	Oklahoma	5.2 ^{bc}	0.27	(43)	-24.6
Hawaii	12.7 ^{bc}	0.69	(01)	84.1	Oregon	5.7 ^{bc}	0.28	(36)	-17.4
Idaho	4.7 ^{bc}	0.45	(50)	-31.9	Pennsylvania	7.1	0.16	(16)	2.9
Illinois	7.5	0.18	(12)	8.7	Rhode Island	9.6 ^{bc}	0.66	(04)	39.1
Indiana	4.8 ^{bc}	0.20	(48)	-30.4	South Carolina	8.2 ^{bc}	0.35	(09)	18.8
Iowa	4.9 ^{bc}	0.27	(45)	-29.0	South Dakota	5.0 ^b	0.54	(44)	-27.5
Kansas	5.3 ^{bc}	0.31	(41)	-23.2	Tennessee	6.5	0.25	(25)	-5.8
Kentucky	6.0	0.28	(34)	-13.0	Texas	6.9	0.15	(17)	0.0
Louisiana	8.8 ^{bc}	0.34	(05)	27.5	Utah	4.8 ^{bc}	0.40	(47)	-30.4
Maine	6.8	0.50	(19)	-1.4	Vermont	4.3 ^{bc}	0.57	(51)	-37.7
Maryland	7.2	0.29	(15)	4.3	Virginia	6.6	0.24	(24)	-4.3
Massachusetts	8.2 ^{bc}	0.26	(08)	18.8	Washington	6.0 ^c	0.24	(33)	-13.0
Michigan	6.4	0.19	(27)	-7.2	West Virginia	6.2	0.39	(30)	-10.1
Minnesota	5.5 ^{bc}	0.24	(38)	-20.3	Wisconsin	6.1	0.24	(32)	-11.6
Mississippi	7.4	0.38	(13)	7.2	Wyoming	4.9 ^b	0.72	(46)	-29.0
Missouri	5.8 ^{bc}	0.23	(35)	-15.9					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table XXIII-13
STOMACH CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, 1996-2000

All Races, Females

State	Rate	SE	Rank	PD
TOTAL U.S.	3.4	0.02		
High Five States				
Hawaii	7.0 ^{bc}	0.47	(01)	105.9
District of Columbia	6.5 ^{bc}	0.61	(02)	91.2
New York	4.3 ^{bc}	0.09	(03)	26.5
New Jersey	4.3 ^{bc}	0.13	(04)	26.5
Rhode Island	4.1 ^b	0.34	(05)	20.6
Low Five States				
Missouri	2.3 ^{bc}	0.12	(47)	-32.4
Nebraska	2.3 ^{bc}	0.21	(48)	-32.4
Kansas	2.3 ^{bc}	0.17	(49)	-32.4
Montana	2.1 ^{bc}	0.28	(50)	-38.2
Iowa	1.9 ^{bc}	0.14	(51)	-44.1

State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.4	0.16	(18)	0.0	Montana	2.1 ^{bc}	0.28	(50)	-38.2
Alaska	3.9	0.68	(08)	14.7	Nebraska	2.3 ^{bc}	0.21	(48)	-32.4
Arizona	2.8 ^{bc}	0.15	(36)	-17.6	Nevada	3.1	0.27	(27)	-8.8
Arkansas	3.4	0.20	(19)	0.0	New Hampshire	2.6 ^b	0.27	(40)	-23.5
California	3.9 ^c	0.07	(09)	14.7	New Jersey	4.3 ^{bc}	0.13	(04)	26.5
Colorado	2.8 ^b	0.17	(35)	-17.6	New Mexico	4.0 ^b	0.30	(07)	17.6
Connecticut	3.7	0.18	(12)	8.8	New York	4.3 ^{bc}	0.09	(03)	26.5
Delaware	3.9	0.42	(10)	14.7	North Carolina	3.2	0.12	(24)	-5.9
District of Columbia	6.5 ^{bc}	0.61	(02)	91.2	North Dakota	3.0	0.38	(30)	-11.8
Florida	3.1 ^c	0.07	(25)	-8.8	Ohio	2.9 ^c	0.09	(32)	-14.7
Georgia	3.3	0.13	(22)	-2.9	Oklahoma	2.5 ^{bc}	0.16	(41)	-26.5
Hawaii	7.0 ^{bc}	0.47	(01)	105.9	Oregon	2.5 ^{bc}	0.16	(44)	-26.5
Idaho	2.4 ^{bc}	0.27	(46)	-29.4	Pennsylvania	3.2	0.09	(23)	-5.9
Illinois	3.4	0.10	(16)	0.0	Rhode Island	4.1 ^b	0.34	(05)	20.6
Indiana	2.6 ^{bc}	0.12	(39)	-23.5	South Carolina	3.5	0.18	(15)	2.9
Iowa	1.9 ^{bc}	0.14	(51)	-44.1	South Dakota	3.1	0.37	(28)	-8.8
Kansas	2.3 ^{bc}	0.17	(49)	-32.4	Tennessee	3.0	0.14	(29)	-11.8
Kentucky	3.4	0.17	(20)	0.0	Texas	3.8 ^c	0.09	(11)	11.8
Louisiana	4.0 ^b	0.18	(06)	17.6	Utah	3.0	0.27	(31)	-11.8
Maine	2.8 ^b	0.26	(34)	-17.6	Vermont	2.4 ^b	0.37	(45)	-29.4
Maryland	3.7	0.16	(13)	8.8	Virginia	3.4	0.14	(17)	0.0
Massachusetts	3.7	0.13	(14)	8.8	Washington	2.8 ^{bc}	0.14	(37)	-17.6
Michigan	3.1	0.11	(26)	-8.8	West Virginia	2.9	0.21	(33)	-14.7
Minnesota	2.5 ^{bc}	0.14	(42)	-26.5	Wisconsin	2.5 ^{bc}	0.13	(43)	-26.5
Mississippi	3.4	0.21	(21)	0.0	Wyoming	2.7 ^b	0.46	(38)	-20.6
Missouri	2.3 ^{bc}	0.12	(47)	-32.4					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b Absolute percent difference between state rate and total U.S. rate is 15% or more.

^c Difference between state rate and total U.S. rate is statistically significant (p<=.0002).

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table XXIII-14
STOMACH CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2000
By Race/Ethnicity, Sex and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <10	0 to <25	>=25 ^f	Complete ^g
<u>Race/Sex</u>									
All Races ^b	29,179	11,579	7,614	4,212	2,495	40,758	55,079	4,448	59,527
Males	17,369	6,351	4,031	2,349	1,379	23,720	31,479	2,426	33,905
Females	11,810	5,228	3,583	1,863	1,116	17,038	23,600	2,022	25,622
White ^b	22,513	8,768	5,968	3,213	2,067	31,281	42,529	3,457	45,986
Males	13,688	4,884	3,213	1,851	1,126	18,572	24,762	1,939	26,701
Females	8,825	3,884	2,755	1,362	941	12,709	17,767	1,518	19,285
Black ^b	4,191	1,675	1,039	649	-	5,866	7,815	431	8,246
Males	2,235	911	515	330	-	3,146	4,154	213	4,367
Females	1,956	764	524	319	-	2,720	3,661	218	3,879
Asian/Pacific Islander ^c	2,211	972	+	+	+	3,183	+	+	+
Males	1,294	539	+	+	+	1,833	+	+	+
Females	917	433	+	+	+	1,350	+	+	+
Hispanic ^d	2,933	1,056	+	+	+	3,989	+	+	+
Males	1,625	585	+	+	+	2,210	+	+	+
Females	1,308	471	+	+	+	1,779	+	+	+

Estimated prevalence percent^a on January 1, 2000, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence, Race/Ethnicity and Sex

Age at Prevalence	Age Specific (Crude)										Age-Adjusted ^e
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race/Sex</u>											
All Races ^c	0.0159%	-	-	0.0005%	0.0029%	0.0088%	0.0205%	0.0519%	0.0981%	0.1196%	0.0177%
Males	0.0186%	-	-	-	0.0028%	0.0104%	0.0266%	0.0709%	0.1352%	0.1643%	0.0238%
Females	0.0133%	-	-	-	0.0030%	0.0071%	0.0146%	0.0351%	0.0704%	0.0964%	0.0131%
White ^c	0.0133%	-	-	-	0.0024%	0.0064%	0.0167%	0.0426%	0.0747%	0.0947%	0.0140%
Males	0.0157%	-	-	-	0.0026%	0.0076%	0.0224%	0.0596%	0.1035%	0.1300%	0.0190%
Females	0.0109%	-	-	-	0.0023%	0.0052%	0.0112%	0.0272%	0.0530%	0.0766%	0.0101%
Black ^c	0.0157%	-	-	-	-	0.0153%	0.0273%	0.0650%	0.1280%	0.1645%	0.0240%
Males	0.0173%	-	-	-	-	0.0189%	0.0334%	0.0903%	0.1717%	0.1744%	0.0301%
Females	0.0143%	-	-	-	-	0.0121%	0.0222%	0.0450%	0.0992%	0.1601%	0.0194%
Asian/Pacific Islander ^c	0.0344%	-	-	-	0.0051%	0.0190%	0.0407%	0.1049%	0.2611%	0.3506%	0.0432%
Males	0.0410%	-	-	-	-	0.0227%	0.0519%	0.1406%	0.3628%	0.4566%	0.0570%
Females	0.0282%	-	-	-	-	0.0157%	0.0308%	0.0759%	0.1840%	0.2763%	0.0326%
Hispanic ^d	0.0111%	-	-	-	0.0050%	0.0116%	0.0274%	0.0645%	0.1397%	0.1971%	0.0254%
Males	0.0118%	-	-	-	0.0051%	0.0111%	0.0317%	0.0932%	0.1823%	0.2582%	0.0325%
Females	0.0103%	-	-	-	0.0049%	0.0121%	0.0234%	0.0411%	0.1094%	0.1645%	0.0204%

^a US 2000 cancer prevalence counts are based on 2000 cancer prevalence proportions from the SEER registries and 1/1/2000 US population estimates based on the average of 1999 and 2000 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.
^b ^c ^d Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.
^e Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.
^f ^g (f) Cases diagnosed 25 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (g) Complete prevalence is obtained by summing 0 to <25 with >=25.
 - Statistics not shown. Rate based on less than 25 cases estimated alive for the time interval.
 + Not available.

SEER Incidence, Delay Adjusted Incidence and US Death Rates+ Stomach Cancer, by Race and Sex




Figure XXIII-1

+ Source: SEER 9 areas and NCHS public use data file. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines and APCs are calculated using the Joinpoint Regression Program. The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).