

Table 27. Suggested Minimum Target Trough Concentrations for Persons with Wild-Type HIV-1

Drug	Concentration (ng/mL)
Amprenavir (Agenerase)	400
Indinavir (Crixivan)	100
Lopinavir/ritonavir (Kaletra)	1000
Nelfinavir (Viracept) ^a	800
Ritonavir (Norvir) ^b	2100
Saquinavir (Fortovase, Invirase)	100-250
Efavirenz (Sustiva)	1000
Nevirapine (Viramune)	3400

a. Measurable active (M8) metabolite.

b. Ritonavir given as a single PI.

Sources:

- Acosta EP, and Gerber JG. Position paper on therapeutic drug monitoring of antiretroviral agents. *AIDS Research Human Retroviruses* 2002; 18(12):825-34.
- Back D, Gatti G, Fletcher CV, et al. Therapeutic drug monitoring in HIV infection: current status and future directions. *AIDS* 2002; 16 (suppl 1) S5-S37.
- Burger DM, Aarnoutse RE, Hugen PWH. Pros and cons of therapeutic drug monitoring of antiretroviral agents. *Curr Opin Infect Dis* 2002;15(1):17-22.
- Optimizing TDM in HIV clinical care. (May 20, 2003. <http://www.hivpharmacology.com>)