References and Included Studies (appearing in text and Evidence Tables)

1. Acheson RM, Collart AB. New Haven survey of joint diseases. XVII. Relationship between some systemic characteristics and osteoarthrosis in a general population. Ann Rheum Dis 1975;34(5):379-87.

2. Peyron JG. Osteoarthritis. The epidemiologic viewpoint. Clin Orthop 1986;213:13-9.

3. CDC. Public health and aging: Projected prevalence of self-reported arthritis or chronic joint symptoms among persons aged 65 years--United States, 2005-2030. Morbidity and Mortality Weekly Report 2003;52(21, May 30, 2003):489.

4. Elders MJ. The increasing impact of arthritis on public health. J Rheumatol Suppl 2000;60:6-8.

5. Callahan CM, Drake BG, Heck DA, et al. Patient outcomes following tricompartmental total knee replacement. A meta-analysis postoperative alignment of total knee replacement. Its effect on survival. JAMA 1994;271(17):1349-57.

6. Alicea J. Scoring systems and their validation for the arthritic knee. In: Insall JN, Churchill SN, eds. Surgery of the Knee. 3rd ed. New York: Livingston; 2001:1507-15.

7. Insall JN, Dorr LD, Scott RD, et al. Rationale of the Knee Society clinical rating system. Orthopaedics & Related Research 1989;248:13-4.

8. Bellamy N, Buchanan WW, Goldsmith CH, et al. Validation study of WOMAC: A health status instrument for measuring clinically important patient relevant outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. The Journal of Rheumatology 1988;15(12):1833-40.

9. Orthopedic Network News. 2002 hip and knee implant review. CMS MedPar (www.OrthopedicNetworkNews.com). Accessed 9/8/03.

10. National Center for Health Statistics. American Academy and American Association of Orthopaedic Surgeons Bulletin 1999;47(3):14.

11. Tierney WM, Fitzgerald JF, Heck D, et al. Tricompartmental knee replacement: A comparison of orthopaedic surgeons' self reported performance rates with surgical indications, intraindications, and expected outcomes. Clinical Orthopaedics 1994;305:209-17.

12. Wright JG, Coyte P, Hawker G, et al. Variation in orthopedic surgeons' perceptions for and outcomes of knee replacement. Can Med Assoc J 1995;152:687-97.

13. Mancuso CA, Ranawat CS, Esdaile JM, et al. Indications for total hip and total knee arthroplasties. Results of orthopaedic surveys. J Arthroplasty 1996 Jan;11(1):34-46.

14. Coyte PC, Hawker G, Croxford R, et al. Variation in rheumatologists' and family physicians' perceptions of the indications for and outcomes of knee replacement surgery. J Rheumatol 1996 Apr;23(4):730-8.

15. Wright JG, Hawker GA, Bombardier C, et al. Physician enthusiasm as an explanation for area variation in the utilization of knee replacement surgery. Med Care 1999 Sep;37(9):946-56.

16. Malmlin LA, Melfi CA, Parchman ML, et al. Management of osteoarthritis of the knee by primary care physicians. Arch Fam Med 1998;7:563-7.

17. Naylor CD, Williams JI. Primary hip and knee replacement surgery: Ontario criteria for case selection and surgical priority. Qual Health Care 1996;5:20-30.

18. Dieppe P, Basler HD, Chard J, et al. Knee replacement surgery for osteoarthritis: Effectiveness, practice variations, indications and possible determinants of utilization. Rheumatology (Oxford) 1999;38(1):73-83.

19. Hadorn DC, Holmes AC. The New Zealand priority criteria project. Part 1: Overview. BMJ 1997a;314(7074):131-4.

20. Hadorn DC, Holmes AC. The New Zealand priority criteria project. Part 2: Coronary artery bypass graft surgery. BMJ 1997b;314(7074):135-8.

21. Saleh KJ, Dykes DC, Tweedie RL, et al. Functional outcome after total knee arthroplasty revision: a meta-analysis. J Arthroplasty 2002 Dec;17(8):967-77.

22. Kane RL, ed. Understanding Health Care Outcomes Research. Gaithersburg, MD: Aspen Publishers, Inc; 1997.

23. Borenstein M, Rothstein H. Comprehensive Meta Analysis: A computer program for research synthesis. Englewood, NJ: Biostat, Inc.; 1999.

24. Leclerc JR, Gent M, Hirsh J, et al. The incidence of symptomatic venous thromboembolism during and after prophylaxis with enoxaparin: a multi-institutional cohort study of patients who underwent hip or knee arthroplasty. Canadian Collaborative Group. Arch Intern Med 1998; 158(8):873-8.

25. Mauerhan DR, Nelson CL, Smith DL, et al Prophylaxis against infection in total joint arthroplasty. One day of cefuroxime compared with three days of cefazolin. J Bone Joint Surg Am 1994 Jan;76(1):39-45.

26. Jones RE, Skedros JG, Chan AJ, et al. Total knee arthroplasty using the S-ROM mobile-bearing hinge prosthesis. J Arthroplasty 2001 Apr;16(3):279-87.

27. Deshmukh R, Hayes J, Pinder I. Does body weight influence outcome after total knee arthroplasty? A 1-year analysis. J Arthroplasty 2002 Apr;17(3):315-9.

28. Fortin P, Clarke A, Joseph L, et al. Outcomes of total hip and knee replacement: preoperative functional status predicts outcomes at six months after surgery. Arthritis Rheum 1999 Aug;42(8):1722-8.

29. Hawker G, Wright J, Coyte P, et al. Health-related quality of life after knee replacement. J Bone Joint Surg Am 1998 Feb;80(2):163-73.

30. Konig A, Kirschner S, Walther M, et al. Hybrid total knee arthroplasty. Arch Orthop Trauma Surg 1998;118(1-2):66-9.

31. Hawker GA, Wright JG, Coyte PC, et al. Differences between men and women in the rate of use of hip and knee arthroplasty.[comment]. New England Journal of Medicine 2000 Apr 6; 342(14):1016-22.

32. Dunlop DD, Song J, Manheim LM, et al. Racial disparities in joint replacement use among older adults. Med Care 2003 Feb;41(2):288-98.

33. Wilson MG, May DS, Kelly JJ. Racial differences in the use of total knee arthroplasty for osteoarthritis among older Americans. Ethn Dis 1994 Winter;4(1):57-67.

34. Insall JN, Dethmers DA. Revision of total knee arthroplasty. Clinical Orthopaedics & Related Research 1982;170:123-30.

35. Mendenhall S. Get the lowdown on orthopaedic implants. Mater Manag Health Care 1996;5(4):30-2.

36. Haas SB, Insall JN, Montgomery W III, et al. Revision total knee arthroplasty with use of modular components with stems inserted without cement. Journal of Bone & Joint Surgery—American Volume 1995;77(11):1700-7.

37. Whiteside LA, Bicalho PS. Radiologic and hisologic analysis of morselized allegraft in revision total knee replacement. Clinical Orthopaedics & Related Research 1998;357:149-56.

38. Callahan CM, Drake BG, Heck DA, et al. Patient outcomes following unicompartmental or bicompartmental arthroplasty. A meta-analysis. J Arthroplasty 1995;10(2):141-50.

39. Brooks DH, Fehring TK, Griffin WL, et al. Polyethylene exchange only for prosthetic knee instability. Clin Orthop 2002 Dec;(405):182-8.

40. Babis GC, Trousdale RT, Morrey BF. The effectiveness of isolated tibial insert exchange in revision total knee arthroplasty. J Bone Joint Surg Am 2002 Jan;84-A(1):64-8.

41. Miller M, Benjamin JB, Marson B, et al. The effect of implant constraint on results of conversion of unicompartmental knee arthroplasty to total knee arthroplasty. Orthopedics 2002 Dec;25(12):1353-7; discussion 7.

42. Christensen CP, Crawford JJ, Olin MD, et al. Revision of the stiff total knee arthroplasty. J Arthroplasty 2002 Jun;17(4):409-15.

43. Gofton WT, Tsigaras H, Butler RA, et al. Revision total knee arthroplasty: fixation with modular stems. Clin Orthop 2002 Nov;(404):158-68.

44. Nazarian DG, Mehta S, Booth Jr RE. A comparison of stemmed and unstemmed components in revision knee arthroplasty. Clin Orthop 2002 Nov;(404):256-62.

45. Lonner JH, Lotke PA, Kim J, et al. Impaction grafting and wire mesh for uncontained defects in revision knee arthroplasty. Clin Orthop 2002 Nov;(404):145-51.

46. Benjamin J, Engh G, Parsley B, et al. Morselized bone grafting of defects in revision total knee arthroplasty. Clin Orthop 2001 Nov;(392):62-7.

47. Hanssen AD. Bone-grafting for severe patellar bone loss during revision knee arthroplasty. J Bone Joint Surg Am 2001 Feb;83-A(2):171-6.

48. Leopold SS, Silverton CD, Barden RM, et al. Isolated revision of the patellar component in total knee arthroplasty. J Bone Joint Surg Am 2003 Jan;85-A(1):41-7.

49 Parvizi J, Seel MJ, Hanssen AD, et al. Patellar component resection arthroplasty for the severely compromised patella. Clin Orthop 2002 Apr;(397):356-61.

50. Werle JR, Goodman SB, Imrie SN. Revision total knee arthroplasty using large distal femoral augments for severe metaphyseal bone deficiency: a preliminary study. Orthopedics 2002 Mar;25(3):325-7.

51. Springer BD, Hanssen AD, Sim FH, et al. The kinematic rotating hinge prosthesis for complex knee arthroplasty. Clin Orthop 2001 Nov;(392):283-91.

52. Ververeli P, Sutton D, Hearn S, et al. Continuous passive motion after total knee arthroplasty. Analysis of cost and benefits. Clin Orthop 1995 Dec;(321):208-15.

53. Worland R, Arredondo J, Angles F, et al. Home continuous passive motion machine versus professional physical therapy following total knee replacement. J Arthroplasty 1998 Oct;13(7):784-7.

54. Hasegawa M, Ohashi T, Uchida A. Heterotopic ossification around distal femur after total knee arthroplasty. Arch Orthop Trauma Surg 2002;122(5):274-8.

55. Hsu R, Tsai Y, Huang T, et al. Hybrid total knee arthroplasty: a 3- to 6-year outcome analysis. J Formos Med Assoc 1998 Jun;97(6):410-5.

56. Larson C, McDowell C, Lachiewicz P. One-peg versus three-peg patella component fixation in total knee arthroplasty. Clin Orthop 2001 Nov;(392):94-100.

57. Liu T, Chen S. Simultaneous bilateral total knee arthroplasty in a single procedure. Int Orthop 1998;22(6):390-3.

58. Moskal J, Diduch D. Postoperative radiographs after total knee arthroplasty: a cost-containment strategy. Am J Knee Surg 1998 Spring;11(2):89-93.

59. Pereira D, Jaffe F, Ortiguera C. Posterior cruciate ligament-sparing versus posterior cruciate ligament-sacrificing arthroplasty. Functional results using the same prosthesis. J Arthroplasty 1998 Feb;13(2):138-44.

60. Rand J, Gustilo R. Comparison of inset and resurfacing patellar prostheses in total knee arthroplasty. J Orthop Surg 1996;4(1):13-22.

61. Diduch D, Insall J, Scott W, et al. Total knee replacement in young, active patients. Long-term follow-up and functional outcome. J Bone Joint Surg Am 1997 Apr;79(4):575-82.

62. Evanich C, Tkach T, von Glinski S, et al. 6- to 10-year experience using countersunk metal-backed patellas. J Arthroplasty 1997 Feb;12(2):149-54.

63. Healy W, Iorio R, Ko J, et al. Impact of cost reduction programs on short-term patient outcome and hospital cost of total knee arthroplasty. J Bone Joint Surg Am 2002 Mar;84-A(3):348-53.

64. Ikejiani C, Leighton R, Petrie D. Comparison of patellar resurfacing versus nonresurfacing in total knee arthroplasty. Can J Surg 2000 Feb;43(1):35-8.

65. Malkani A, Rand J, Bryan R, et al. Total knee arthroplasty with the kinematic condylar prosthesis. A ten-year follow-up study. J Bone Jt Surg Am 1995 Mar;77(3):423-31.

66. O'Rourke M, Callaghan J, Goetz D, et al. Osteolysis associated with a cemented modular posterior-cruciate-substituting total knee design : five to eight-year follow-up. J Bone Jt Surg Am 2002 Aug;84-A(8):1362-71.

67. Regner L, Carlsson L, Karrholm J, et al. Clinical and radiologic survivorship of cementless tibial components fixed with finned polyethylene pegs. J Arthroplasty 1997 Oct;12(7):751-8.

68. Schroder H, Berthelsen A, Hassani G, et al. Cementless porous-coated total knee arthroplasty: 10-year results in a consecutive series. J Arthroplasty 2001 Aug;16(5):559-67.

69. Bert J, Gross M, Kline C. Patient demand matching in total knee arthroplasty: is it necessary? Am J Knee Surg 2001 Winter;14(1):39-42.

70. Bourne R, Rorabeck C, Vaz M, et al. Resurfacing versus not resurfacing the patella during total knee replacement. Clin Orthop 1995 Dec;(321):156-61.

71. Cohen R, Forrest C, Benjamin J. Safety and efficacy of bilateral total knee arthroplasty. J Arthroplasty 1997 Aug;12(5):497-502.

72. Heck D, Robinson R, Partridge C, et al. Patient outcomes after knee replacement. Clin Orthop 1998 Nov;(356):93-110.

73. Lin Y, Su J, Lin G, et al. Impact of a clinical pathway for total knee arthroplasty. Kaohsiung J Med Sci 2002 Mar;18(3):134-40.

74. Matsueda M, Gustilo R. Subvastus and medial parapatellar approaches in total knee arthroplasty. Clin Orthop 2000 Feb;(371):161-8.

75. Bullens P, van Loon C, de Waal Malefijt M, et al. Patient satisfaction after total knee arthroplasty: a comparison between subjective and objective outcome assessments. J Arthroplasty 2001 Sep;16(6):740-7.

76. Elke R, Meier G, Warnke K, et al. Outcome analysis of total knee-replacements in patients with rheumatoid arthritis versus osteoarthritis. Arch Orthop Trauma Surg 1995;114(6):330-4.

77. Jenny J, Jenny G. Preservation of anterior cruciate ligament in total knee arthroplasty. Arch Orthop Trauma Surg 1998;118(3):145-8.

78. Meding J, Ritter M, Faris P, et al. Does the preoperative radiographic degree of osteoarthritis correlate to results in primary total knee arthroplasty? J Arthroplasty 2001 Jan;16(1):13-6.

79. Ranawat C, Luessenhop C, Rodriguez J. The press-fit condylar modular total knee system. Four-to-six-year results with a posterior-cruciate-substituting design. J Bone Jt Surg Am 1997 Mar;79(3):342-8.

80. Rodriguez J, Saddler S, Edelman S, et al. Long-term results of total knee arthroplasty in class 3 and 4 rheumatoid arthritis. J Arthroplasty 1996 Feb;11(2):141-5.

81. Yang K, Yeo S, Lee B, et al. Total knee arthroplasty in diabetic patients: a study of 109 consecutive cases. J Arthroplasty 2001 Jan;16(1):102-6.

82. Brown T, Diduch D, Moskal J. Component size asymmetry in bilateral total knee arthroplasty. Am J Knee Surg 2001 Spring;14(2):81-4.

83. Cloutier J, Sabouret P, Deghrar A. Total knee arthroplasty with retention of both cruciate ligaments. A 9 to 11 year follow-up study. Eur J Orthop Surg Traumatol 2001;11(1):41-6.

84. Duffy G, Berry D, Rand J. Cement versus cementless fixation in total knee arthroplasty. Clin Orthop 1998 Nov;(356):66-72.

85. Ewald F, Wright R, Poss R, et al. Kinematic total knee arthroplasty: a 10- to 14-year prospective follow-up review. J Arthroplasty 1999 Jun;14(4):473-80.

86. Gill G, Joshi A. Long-term results of cemented, posterior cruciate ligament-retaining total knee arthroplasty in osteoarthritis. Am J Knee Surg 2001 Fall;14(4):209-14.

87. Indelli P, Aglietti P, Buzzi R, et al. The Insall-Burstein II prosthesis: A 5- to 9-year follow-up study in osteoarthritic knees. J Arthroplasty 2002;17(5):544-9.

88. Martin S, McManus J, Scott R, et al. Press-fit condylar total knee arthroplasty. 5- to 9-year follow-up evaluation. J Arthroplasty 1997 Sep;12(6):603-14.

89. Miyasaka K, Ranawat C, Mullaji A. 10- to 20-year followup of total knee arthroplasty for valgus deformities. Clin Orthop 1997 Dec;(345):29-37.

90. Mokris J, Smith S, Anderson S. Primary total knee arthroplasty using the Genesis Total Knee Arthroplasty System: 3- to 6-year follow-up study of 105 knees. J Arthroplasty 1997 Jan;12(1):91-8.

91. Mont M, Yoon T, Krackow K, et al. Eliminating patellofemoral complications in total knee arthroplasty: clinical and radiographic results of 121 consecutive cases using the Duracon system. J Arthroplasty 1999 Jun;14(4):446-55.

92. Rinta-Kiikka I, Savilahti S, Pajamaki J, et al. A five to seven years follow-up of 102 cementless Synatomic knee arthroplasties. Ann Chir Gynaecol 1996;85(1):77-85.

93. Sextro G, Berry D, Rand J. Total knee arthroplasty using cruciate-retaining kinematic condylar prosthesis. Clin Orthop 2001 Jul;388:33-40.

94. Bachmeier C, March L, Cross M, et al. A comparison of outcomes in osteoarthritis patients undergoing total hip and knee replacement surgery. Osteoarthritis Cartilage 2001 Feb;9(2):137-46.

95. Beaupre L, Davies D, Jones C, et al. Exercise combined with continuous passive motion or slider board therapy compared with exercise only: a randomized controlled trial of patients following total knee arthroplasty. Phys Ther 2001 Apr;81(4):1029-37.

96. Jones CA, Voaklander DC, Johnston DW, et al. The effect of age on pain, function, and quality of life after total hip and knee arthroplasty. Arch Intern Med 2001 Feb;161(3):454-60.

97. Stickles B, Phillips L, Brox W, et al. Defining the relationship between obesity and total joint arthroplasty. Obes Res 2001 Mar;9(3):219-23.

98. Clark C, Rorabeck C, MacDonald S, et al. Posterior-stabilized and cruciate-retaining total knee replacement: a randomized study. Clin Orthop 2001 Nov;(392):208-12.

99. Bert J, Gross M, Kline C. Outcome results after total knee arthroplasty: does the patient's physical and mental health improve? Am J Knee Surg 2000 Fall;13(4):223-7.

100. Kiebzak G, Campbell M, Mauerhan D. The SF-36 general health status survey documents the burden of osteoarthritis and the benefits of total joint arthroplasty: But why should we use it? Am J Managed Care 2002;8(5):463-74.

101. Baldwin J, Rubinstein RJ. The effect of bone quality on the outcome of ingrowth total knee arthroplasty. Am J Knee Surg 1996 Spring;9(2):45-9; discussion 9-50.

102. Harwin S. Patellofemoral complications in symmetrical total knee arthroplasty. J Arthroplasty 1998 Oct;13(7):753-62.

103. Gioe T, Bowman K. A randomized comparison of all-polyethylene and metal-backed tibial components. Clin Orthop 2000 Nov;(380):108-15.

104. Hube R, Sotereanos N, Reichel H. The midvastus approach for total knee arthroplasty. Orthop Traumatol 2002;10(3):235-44.

105. Jordan L, Olivo J, Voorhorst P. Survivorship analysis of cementless meniscal bearing total knee arthroplasty. Clin Orthop 1997 May;(338):119-23.

106. Konig A, Scheidler M, Rader C, et al. The need for a dual rating system in total knee arthroplasty. Clin Orthop 1997 Dec;(345):161-7.

107. Title C, Rodriguez J, Ranawat C. Posterior cruciate-sacrificing versus posterior cruciate-substituting total knee arthroplasty: a study of clinical and functional outcomes in matched patients. J Arthroplasty 2001 Jun;16(4):409-14.

108. Gill G, Joshi A, Mills D. Total condylar knee arthroplasty. 16- to 21-year results. Clin Orthop 1999 Oct;(367):210-5.

109. Ritter M, Worland R, Saliski J, et al. Flat-on-flat, nonconstrained, compression molded polyethylene total knee replacement. Clin Orthop 1995 Dec;(321):79-85.

110. Griffin F, Scuderi G, Insall J, et al. Total knee arthroplasty in patients who were obese with 10 years followup. Clin Orthop 1998 Nov;(356):28-33.

111. Konig A, Walther M, Kirschner S, et al. Balance sheets of knee and functional scores 5 years after total knee arthroplasty for osteoarthritis: a source for patient information. J Arthroplasty 2000 Apr;15(3):289-94.

112. Lombardi Jr A, Mallory T, Fada R, et al. An algorithm for the posterior cruciate ligament in total knee arthroplasty. Clin Orthop 2001 Nov;(392):75-87.

113. Blanchard J, Meuwly JY, Leyvraz PF, et al. Prevention of deep-vein thrombosis after total knee replacement. Randomised comparison between a low-molecular-weight heparin (nadroparin) and mechanical prophylaxis with a foot-pump system. J Bone Joint Surg Br 1999; 81(4):654-9.

114. Colwell CW Jr, Spiro TE, Trowbridge AA, et al. Efficacy and safety of enoxaparin versus unfractionated heparin for prevention of deep venous thrombosis after elective knee arthroplasty. Enoxaparin Clinical Trial Group. Clin Orthop 1995; (321):19-27.

115. Francis CW, Pellegrini VD Jr, Leibert KM, et al. Comparison of two warfarin regimens in the prevention of venous thrombosis following total knee replacement. Thromb Haemost 1996; 75(5):706-11.

116. Heit JA, Elliott CG, Trowbridge AA, et al. Ardeparin sodium for extended out-of-hospital prophylaxis against venous thromboembolism after total hip or knee replacement. A randomized, double-blind, placebo-controlled trial. Ann Intern Med 2000; 132(11):853-61.

117. Leclerc JR, Geerts WH, Desjardins L, et al. Prevention of venous thromboembolism after knee arthroplasty. A randomized, double-blind trial comparing enoxaparin with warfarin. Ann Intern Med 1996; 124(7):619-26.

118. Perhoniemi V, Vuorinen J, Myllynen P, et al. The effect of enoxaparin in prevention of deep venous thrombosis in hip and knee surgery--a comparison with the dihydroergotamine-heparin combination. Ann Chir Gynaecol 1996; 85(4):359-63.

119. Robinson KS, Anderson DR, Gross M, et al. Ultrasonographic screening before hospital discharge for deep venous thrombosis after arthroplasty: the post-arthroplasty screening study. A randomized, controlled trial. Ann Intern Med 1997; 127(6):439-45.

120. Westrich GH, Sculco TP. Prophylaxis against deep venous thrombosis after total knee arthroplasty: Pneumatic plantar compression and aspirin compared with aspirin alone. J Bone Jt Surg Am 1996; 78(6):826-34.

121. Chiu FY, Chen CM, Lin CF, et al. Cefuroxime-impregnated cement in primary total knee arthroplasty: a prospective, randomized study of three hundred and forty knees. J Bone Joint Surg Am 2002; 84-A(5):759-62.

122. Periti P, Stringa G, Mini E. Comparative multicenter trial of teicoplanin versus cefazolin for antimicrobial prophylaxis in prosthetic joint implant surgery. Italian Study Group for Antimicrobial Prophylaxis in Orthopedic Surgery. Eur J Clin Microbiol Infect Dis 1999; 18(2):113-9.

123. Abdel-Salam A, Eyres KS. Effects of tourniquet during total knee arthroplasty. A prospective randomised study. J Bone Joint Surg Br 1995; 77(2):250-3.

124. Wakankar HM, Nicholl JE, Koka R, et al. The tourniquet in total knee arthroplasty. A prospective, randomised study. J Bone Jt Surg Br 1999; 81(1):30-3.

125. Katz B, Freund D, Heck D, et al. Demographic variation in the rate of knee replacement: a multi-year analysis. Health Serv Res 1996 Jun;31(2):125-40.

126. McBean AM, Gornick M. Differences by race in the rates of procedures performed in hospitals for Medicare beneficiaries. Health Care Financ Rev 1994 Summer;15(4):77-90.

127. Escarce JJ, Epstein KR, Colby DC, et al. Racial differences in the elderly's use of medical procedures and diagnostic tests. Am J Public Health. Jul 1993;83(7):948-54.

128. Norton E, Garfinkel S, McQuay L, et al. The effect of hospital volume on the in-hospital complication rate in knee replacement patients. Health Serv Res 1998 Dec 1998;33(5 Pt 1):1191-210. Comment in: Health Serv Res 1998 Dec;33(5 Pt 1):85-90. PMID: 9865216.

129. Brighton B, Tornetta P. Part I. Methodological issues in the design of orthopaedic studies: Hierarchy of evidence. From case reports to randomized controlled trials. Clinical Orthopaedics & Related Research 2003;413:19-24.

130. Moseley JB, O'Malley K, Petersen NJ, et al. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. New Engl J Med 2002;347:81-8.

131. Cho MK, Bero LA. Instruments for assessing the quality of drug studies published in the medical literature. JAMA 1994;272(2):101-4.

132. Harbour R, Miller J. A new system for grading recommendations in evidence based guidelines. BMJ 2001;323(7308):334-6.

133. Reisch JS, Tyson JE, Mize SG. Aid to the evaluation of therapeutic studies. Pediatrics 1989;84(5):815-27.

134. Gross M. A critique of the methodologies used in clinical studies of hip-joint arthroplasty published in the English-language orthopaedic literature. Journal of Bone & Joint Surgery—American Volume 1988;70(9):1364-71.

135. Gartland JJ. Orthopaedic clinical research. Deficiencies in experimental design and determinations of outcomes. Journal of Bone & Joint Surgery—American Volume 1988;70(9):1357-64.

136. Smith JS, Watts HG. Methods for locating missing patients for the purpose of long-term clinical studies. Journal of Bone & Joint Surgery—American Volume 1998;80(3):431-8.

137. Saleh KJ, Radosevich DM, Kassim RA, et al. Comparison of commonly used orthopaedic outcome measures using palm-top computers and paper surveys. J Orthop Res 2002;20(6):1146-51.

138. Larson C, Lachiewicz P. Patellofemoral complications with the Insall-Burstein II posterior-stabilized total knee arthroplasty. J Arthroplasty 1999 Apr;14(3):288-92.

139. Ritter M, Berend M, Meding J, et al. Long-term followup of anatomic graduated components posterior cruciate-retaining total knee replacement. Clin Orthop 2001 Jul;388:51-7.

140. Baldwin J, El-Saied M, Rubinstein RJ. Uncemented total knee arthroplasty: report of 109 titanium knees with cancellous-structured porous coating. Orthopedics 1996 Feb;19(2):123-30.