References and Bibliography

Miller, G. and Ramer, J. C. (ed.). Static encephalopathies of infancy and childhood. New York: Raven Press; 1992:372

Vermont-Oxford Neonatal Network: Vermont Neonatal Network 2000 Annual Report. 2000. Burlington, Vermont.

Postnatal corticosteroids and sensorineural outcome at 5 years of age. J Paediatr Child Health 2000 Jun;36(3):256-61.

Agustines LA, Lin YG, Rumney PJ et al. Outcomes of extremely low-birth-weight infants between 500 and 750 g. Am J Obstet Gynecol 2000;182(5):1113-6.

Akerman BA, Thomassen PA. The fate of "small twins": a four-year follow-up study of low birthweight and prematurely born twins. Acta Genet Med Gemellol (Roma) 1992;41(2-3):97-104.

Algawi K, Goggin M, O'Keefe M. Refractive outcome following diode laser versus cryotherapy for eyes with retinopathy of prematurity. Br J Ophthalmol 1994;78(8):612-4.

Allan WC, Vohr B, Makuch RW et al. Antecedents of cerebral palsy in a multicenter trial of indomethacin for intraventricular hemorrhage. Arch Pediatr Adolesc Med 1997;151(6):580-5.

Als H, Lawhon G, Duffy FH et al. Individualized developmental care for the very low-birth-weight preterm infant. Medical and neurofunctional effects. JAMA 1994 Sep 21;272(11):853-8.

Ambalavanan N, Nelson KG, Alexander G et al. Prediction of neurologic morbidity in extremely low birth weight infants. J Perinatol 2000;20(8 Pt 1):496-503.

American Association on Mental Retardation. Mental Retardation: Definition, Classification, and Systems of Supports. 9thed. Washington, DC: American Association on Mental Retardation; 1992

Anderson AE, Wildin SR, Woodside M et al. Severity of medical and neurologic complications as a determinant of neurodevelopmental outcome at 6 and 12 months in very low birth weight infants. J Child Neurol 1996;11(3):215-9.

Avila-Diaz M, Flores-Huerta S, Martinez-Muniz I et al. Increments in whole body bone mineral content associated with weight and length in pre-term and full-term infants during the first 6 months of life. Arch Med Res 2001;32(4):288-92.

Bader D, Ramos AD, Lew CD et al. Childhood sequelae of infant lung disease: exercise and pulmonary function abnormalities after bronchopulmonary dysplasia. J Pediatr 1987 May;110(5):693-9.

Bagnato SJ, Campbell TF. Comprehensive neurodevelopmental evaluation of children with brain insults. In: Miller G and Ramer JC, editors. Static encephalopathies of infancy and childhood. New York: Raven Press Ltd; 1992. p. 27-44.

Baraldi E, Monciotti C, Filippone M et al. Gas exchange during exercise in diabetic children. Pediatr Pulmonol 1992 Jul;13(3):155-60.

Baraldi E, Filippone M, Trevisanuto D et al. Pulmonary function until two years of life in infants with bronchopulmonary dysplasia. Am J Respir Crit Care Med 1997 Jan;155(1):149-55.

Barrington KJ. The adverse neuro-developmental effects of postnatal steroids in the preterm infant: a systematic review of RCTs. BMC Pediatr 2001;1(1):1.

Battin M, Ling EW, Whitfield MF et al. Has the outcome for extremely low gestational age (ELGA) infants improved following recent advances in neonatal intensive care? Am J Perinatol 1998;15(8):469-77.

Bayley N. Bayley Scales of Development. 2ed. New York, NY: Psychology Corporatoin, Harcourt Brace & Co.; 1993

Blayney M, Kerem E, Whyte H et al. Bronchopulmonary dysplasia: improvement in lung function between 7 and 10 years of age. J Pediatr 1991 Feb;118(2):201-6.

Blitz RK, Wachtel RC, Blackmon L et al. Neurodevelopmental outcome of extremely low birth weight infants in Maryland. Md Med J 1997;46(1):18-24.

Bos AF, Martijn A, Okken A et al. Quality of general movements in preterm infants with transient periventricular echodensities. Acta Paediatr 1998;87(3):328-35.

Botting N, Powls A, Cooke RW et al. Attention deficit hyperactivity disorders and other psychiatric outcomes in very low birthweight children at 12 years. J Child Psychol Psychiatry 1997 Nov;38(8):931-41.

Brazy JE, Eckerman CO, Oehler JM et al. Nursery Neurobiologic Risk Score: important factor in predicting outcome in very low birth weight infants. J Pediatr 1991;118(5):783-92.

Brazy JE, Goldstein RF, Oehler JM et al. Nursery neurobiologic risk score: levels of risk and relationships with nonmedical factors. J Dev Behav Pediatr 1993;14(6):375-80.

Breslau N, Brown GG, DelDotto JE et al. Psychiatric sequelae of low birth weight at 6 years of age. J Abnorm Child Psychol 1996 Jun;24(3):385-400.

Breslau N, Chilcoat HD. Psychiatric sequelae of low birth weight at 11 years of age. Biol Psychiatry 2000;47(11):1005-11.

Briscoe J, Gathercole SE, Marlow N. Short-term memory and language outcomes after extreme prematurity at birth. J Speech Lang Hear Res 1998;41(3):654-66.

Brooks AM, Byrd RS, Weitzman M et al. Impact of low birth weight on early childhood asthma in the United States. Arch Pediatr Adolesc Med 2001 Mar;155(3):401-6.

Buhrer C, Grimmer I, Metze B et al. The CRIB (Clinical Risk Index for Babies) score and neurodevelopmental impairment at one year corrected age in very low birth weight infants. Intensive Care Med 2000;26(3):325-9.

Burguet A, Monnet E, Pauchard JY et al. Some risk factors for cerebral palsy in very premature infants: importance of premature rupture of membranes and monochorionic twin placentation. Biol Neonate 1999;75(3):177-86.

Casiro O, Bingham W, MacMurray B et al. One-year follow-up of 89 infants with birth weights of 500 to 749 grams and respiratory distress syndrome randomized to two rescue doses of synthetic surfactant or air placebo. J Pediatr 1995;126(5 Pt 2):53-60.

Chen CH, Shen WC, Wang TM et al. Cerebral magnetic resonance imaging of preterm infants after corrected age of one year. Zhonghua Min Guo Xiao Er Ke Yi Xue Hui Za Zhi 1995;36(4):261-5.

Cherkes-Julkowski M. Learning disability, attention-deficit disorder, and language impairment as outcomes of prematurity: a longitudinal descriptive study. J Learn Disabil 1998;31(3):294-306.

Cheung PY, Prasertsom W, Finer NN et al. Rescue high frequency oscillatory ventilation for preterm infants: neurodevelopmental outcome and its prediction. Biol Neonate 1997;71(5):282-91.

Cheung PY, Peliowski A, Robertson CM. The outcome of very low birth weight neonates (</=1500 g) rescued by inhaled nitric oxide: neurodevelopment in early childhood. J Pediatr 1998;133(6):735-9.

Cheung PY, Barrington KJ, Finer NN et al. Early childhood neurodevelopment in very low birth weight infants with predischarge apnea. Pediatr Pulmonol 1999;27(1):14-20.

Chye JK, Gray PH. Rehospitalization and growth of infants with bronchopulmonary dysplasia: a matched control study. J Paediatr Child Health 1995;31(2):105-11.

Cioni G, Bertuccelli B, Boldrini A et al. Correlation between visual function, neurodevelopmental outcome, and magnetic resonance imaging findings in infants with periventricular leucomalacia. Arch Dis Child Fetal Neonatal Ed 2000;82(2):134-40.

Cioni G, Bos AF, Einspieler C et al. Early neurological signs in preterm infants with unilateral intraparenchymal echodensity. Neuropediatrics 2000;31(5):240-51.

Connolly BP, McNamara JA, Sharma S et al. A comparison of laser photocoagulation with trans-scleral cryotherapy in the treatment of threshold retinopathy of prematurity. Ophthalmology 1998;105(9):1628-31.

Connors JM, O'Callaghan MJ, Burns YR et al. The influence of growth on development outcome in extremely low birthweight infants at 2 years of age. J Paediatr Child Health 1999;35(1):37-41.

Contractor CP, Leslie GI, Bowen JR et al. The Neonatal Neurobiologic Risk Score: does it predict outcome in very premature infants? Indian Pediatr 1996;33(2):95-101.

Cooke RW, Abernethy LJ. Cranial magnetic resonance imaging and school performance in very low birth weight infants in adolescence. Arch Dis Child Fetal Neonatal Ed 1999;81(2):116-21.

Cooke RW. Trends in incidence of cranial ultrasound lesions and cerebral palsy in very low birthweight infants 1982-93. Arch Dis Child Fetal Neonatal Ed 1999;80(2):115-7.

Corbet A, Long W, Schumacher R et al. Double-blind developmental evaluation at 1-year corrected age of 597 premature infants with birth weights from 500 to 1350 grams enrolled in three placebo-controlled trials of prophylactic synthetic surfactant. J Pediatr 1995;126(5 Pt 2):5-12.

Crocker AC. Causes of mental retardation. Pediatr Ann 1989;18:623-35.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. Multicenter trial of cryotherapy for retinopathy of prematurity. One-year outcome--structure and function. Arch Ophthalmol 1990;108(10):1408-16.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. Multicenter trial of cryotherapy for retinopathy of prematurity. 3 1/2-year outcome--structure and function. Arch Ophthalmol 1993 Mar;111(3):339-44.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. The natural ocular outcome of premature birth and retinopathy. Status at 1 year. Arch Ophthalmol 1994;112(7):903-12.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. Multicenter trial of cryotherapy for retinopathy of prematurity. Snellen visual acuity and structural outcome at 5 1/2 years after randomization. Arch Ophthalmol 1996;114(4):417-24.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. Effect of retinal ablative therapy for threshold retinopathy of prematurity: results of Goldmann perimetry at the age of 10 years. Arch Ophthalmol 2001;119(8):1120-5.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. Multicenter Trial of Cryotherapy for Retinopathy of Prematurity: ophthalmological outcomes at 10 years. Arch Ophthalmol 2001;119(8):1110-8.

Cryotherapy for Retinopathy of Prematurity Cooperative Group. Contrast sensitivity at age 10 years in children who had threshold retinopathy of prematurity. Arch Ophthalmol 2001;119(8):1129-33.

Dammann O, Dammann CE, Allred EN et al. Fetal growth restriction is not associated with a reduced risk for bilateral spastic cerebral palsy in very-low-birthweight infants. Early Hum Dev 2001;64(2):79-89.

de Vonderweid U, Spagnolo A, Corchia C et al. Italian multicentre study on very low-birth-weight babies. Neonatal mortality and two-year outcome. Acta Paediatr 1994;83(4):391-6.

de Vries LS, Eken P, Groenendaal F et al. Correlation between the degree of periventricular leukomalacia diagnosed using cranial ultrasound and MRI later in infancy in children with cerebral palsy. Neuropediatrics 1993 Oct;24(5):263-8.

deRegnier RA, Roberts D, Ramsey D et al. Association between the severity of chronic lung disease and first-year outcomes of very low birth weight infants. J Perinatol 1997;17(5):375-82.

Dezoete JA, MacArthur BA, Aftimos S. Developmental outcome at 18 months of children less than 1000 grams. N Z Med J 1997;110(1045):205-7.

Dogru M, Shirabe H, Nakamura M et al. Development of grating acuity in infants with Retinopathy of Prematurity. Acta Ophthalmol Scand 1999;77(1):72-5.

Doyle LW, Casalaz D, The Victorian Infant Collaborative Study Group. Outcome at 14 years of extremely low birthweight infants: a regional study. Arch Dis Child Fetal Neonatal Ed 2001;85(3):159-64.

Doyle LW, Cheung MM, Ford GW et al. Birth weight <1501 g and respiratory health at age 14. Arch Dis Child 2001 Jan;84(1):40-4.

Doyle LW. Outcome at 5 years of age of children 23 to 27 weeks' gestation: refining the prognosis. Pediatrics 2001 Jul;108(1):134-41.

Duvanel CB, Fawer CL, Cotting J et al. Long-term effects of neonatal hypoglycemia on brain growth and psychomotor development in small-for-gestational-age preterm infants. J Pediatr 1999;134(4):492-8.

Ehrenkranz RA, Walsh-Sukys M. New consensus definition of bronchopulmonary dysplasia predicts pulmonary and neurodevelopmental outcomes in early infancy. Pediatr Res 2001;49(4):276A.

Ekert PG, Keenan NK, Whyte HE et al. Visual evoked potentials for prediction of neurodevelopmental outcome in preterm infants. Biol Neonate 1997;71(3):148-55.

Emsley HC, Wardle SP, Sims DG et al. Increased survival and deteriorating developmental outcome in 23 to 25 week old gestation infants, 1990-4 compared with 1984-9. Arch Dis Child Fetal Neonatal Ed 1998;78(2):99-104.

Finer NN, Tarin T, Vaucher YE et al. Intact survival in extremely low birth weight infants after delivery room resuscitation. Pediatrics 1999;104(4):e40.

Finnstrom O, Otterblad Olausson P, Sedin G et al. Neurosensory outcome and growth at three years in extremely low birthweight infants: follow-up results from the Swedish national prospective study. Acta Paediatr 1998;87(10):1055-60.

Ford GW, Doyle LW, Davis NM et al. Very low birth weight and growth into adolescence. Arch Pediatr Adolesc Med 2000;154(8):778-84.

Foreman N, Fielder A, Minshell C et al. Visual search, perception, and visual-motor skill in "healthy" children born at 27-32 weeks' gestation. J Exp Child Psychol 1997;64(1):27-41.

Forslund M. Growth and motor performance in preterm children at 8 years of age. Acta Paediatr 1992;81(10):840-2.

French NP, Hagan R, Evans SF et al. Repeated antenatal corticosteroids: size at birth and subsequent development. Am J Obstet Gynecol 1999;180(1 Pt 1):114-21.

Futagi Y, Suzuki Y, Goto M et al. Neurodevelopmental outcomes of infants with birth weights of less than 1000 g: comparison between periods before and after the introduction of surfactant. Brain Dev 1999;21(7):453-7.

Gaillard EA, Cooke RW, Shaw NJ. Improved survival and neurodevelopmental outcome after prolonged ventilation in preterm neonates who have received antenatal steroids and surfactant. Arch Dis Child Fetal Neonatal Ed 2001;84(3):194-6.

Gaudet LM, Smith GN. Cerebral palsy and chorioamnionitis: the inflammatory cytokine link. Obstet Gynecol Surv 2001 Jul;56(7):433-6.

Gerdes J, Gerdes M, Beaumont E et al. Health and neurodevelopmental outcome at 1-year adjusted age in 508 infants weighing 700 to 1100 grams who received prophylaxis with one versus three doses of synthetic surfactant. J Pediatr 1995;126(5 Pt 2):26-32.

Gerhardt T, Hehre D, Feller R et al. Serial determination of pulmonary function in infants with chronic lung disease. J Pediatr 1987 Mar;110(3):448-56.

Gerner EM, Katz-Salamon M, Hesser U et al. Psychomotor development at 10 months as related to neonatal health status: the Stockholm Neonatal Project. Acta Paediatr Suppl 1997 Apr;419:37-43.

Goetz MC, Gretebeck RJ, Oh KS et al. Incidence, timing, and follow-up of periventricular leukomalacia. Am J Perinatol 1995 Sep;12(5):325-7.

Goldstein RF, Thompson RJ, Jr., Oehler JM et al. Influence of acidosis, hypoxemia, and hypotension on neurodevelopmental outcome in very low birth weight infants. Pediatrics 1995 Feb;95(2):238-43.

Gosch A, Brambring M, Gennat H et al. Longitudinal study of neuropsychological outcome in blind extremely-low-birthweight children. Dev Med Child Neurol 1997 May;39(5):297-304.

Gregoire MC, Lefebvre F, Glorieux J. Health and developmental outcomes at 18 months in very preterm infants with bronchopulmonary dysplasia. Pediatrics 1998 May;101(5):856-60.

Grether JK, Hoogstrate J, Walsh-Greene E et al. Magnesium sulfate for tocolysis and risk of spastic cerebral palsy in premature children born to women

without preeclampsia. Am J Obstet Gynecol 2000;183(3):717-25.

Gross SJ, Iannuzzi DM, Kveselis DA et al. Effect of preterm birth on pulmonary function at school age: a prospective controlled study. J Pediatr 1998 Aug;133(2):188-92.

Guyer B, Hoyert DL, Martin JA et al. Annual summary of vital statistics--1998. Pediatrics 1999 Dec;104(6):1229-46.

Hack M, Breslau N, Aram D et al. The effect of very low birth weight and social risk on neurocognitive abilities at school age. J Dev Behav Pediatr 1992 Dec;13(6):412-20.

Hack M, Weissman B, Borawski-Clark E. Catch-up growth during childhood among very low-birth-weight children. Arch Pediatr Adolesc Med 1996 Nov;150(11):1122-9.

Hack M, Friedman H, Fanaroff AA. Outcomes of extremely low birth weight infants. Pediatrics 1996;98(5):931-7.

Hack M, Wilson-Costello D, Friedman H et al. Neurodevelopment and predictors of outcomes of children with birth weights of less than 1000 g: 1992-1995. Arch Pediatr Adolesc Med 2000;154(7):725-31.

Hack M, Flannery DJ, Schluchter M et al. Outcomes in young adulthood for very-low-birth-weight infants. N Engl J Med 2002 Jan 17;346(3):149-57.

Harvey EM, Dobson V, Luna B. Long-term grating acuity and visual-field development in preterm children who experienced bronchopulmonary dysplasia. Dev Med Child Neurol 1997 Mar;39(3):167-73.

Hille ET, den Ouden AL, Bauer L et al. School performance at nine years of age in very premature and very low birth weight infants: perinatal risk factors and predictors at five years of age. Collaborative Project on Preterm and Small for Gestational Age (POPS) Infants in The Netherlands. J Pediatr 1994 Sep;125(3):426-34.

Hille ET, den Ouden AL, Saigal S et al. Behavioural problems in children who weigh 1000 g or less at birth in four countries. Lancet 2001 May 26;357(9269):1641-3.

Holling EE, Leviton A. Characteristics of cranial ultrasound white-matter echolucencies that predict disability: a review. Dev Med Child Neurol 1999 Feb;41(2):136-9.

Hollingshead A. Four-Factir Index of Social Status. New Haven: Yale University; 1975

Horwood A, Williams B. Does neonatal ocular misalignment predict later abnormality? Eye 2001;15(Pt 4):485-91.

Horwood LJ, Darlow BA, Mogridge N. Breast milk feeding and cognitive ability at 7-8 years. Arch Dis Child Fetal Neonatal Ed 2001;84(1):23-7.

Hoyt CS, Good WV. The many challenges of childhood blindness. Arch Dis Child 2001;85(6):452-3.

Iai M, Takashima S. Thalamocortical development of parvalbumin neurons in normal and periventricular leukomalacia brains. Neuropediatrics 1999;30(1):14-8.

Iles R, Edmunds AT. Assessment of pulmonary function in resolving chronic lung disease of prematurity. Arch Dis Child Fetal Neonatal Ed 1997 Mar;76(2):F113-F117.

Jacob SV, Lands LC, Coates AL et al. Exercise ability in survivors of severe bronchopulmonary dysplasia. Am J Respir Crit Care Med 1997 Jun;155(6):1925-9.

Jankov RP, Asztalos EV, Skidmore MB. Favourable neurological outcomes following delivery room cardiopulmonary resuscitation of infants < or = 750 g at birth. J Paediatr Child Health 2000;36(1):19-22.

Kato EH, Yamada H, Matsumoto Y et al. Relation between perinatal factors and outcome of very low birth weight infants. J Perinat Med 1996;24(6):677-86.

Katz-Salamon M, Gerner EM, Jonsson B et al. Early motor and mental development in very preterm infants with chronic lung disease. Arch Dis Child Fetal Neonatal Ed 2000 Jul;83(1):F1-F6.

Katz KS, Dubowitz LM, Henderson S et al. Effect of cerebral lesions on continuous performance test responses of school age children born prematurely. J Pediatr Psychol 1996 Dec;21(6):841-55.

Keith CG, Doyle LW. Retinopathy of prematurity in extremely low birth weight infants. Pediatrics 1995;95(1):42-5.

Kiely M. The prevalence of mental retardation. Epidemiol Rev 1987;9:194-218.

Kim JN, Namgung R, Chang W et al. Prospective evaluation of perinatal risk factors for cerebral palsy and delayed development in high risk infants. Yonsei Med J 1999 Aug;40(4):363-70.

Koller H, Lawson K, Rose SA et al. Patterns of cognitive development in very low birth weight children during the first six years of life. Pediatrics 1997 Mar;99(3):383-9.

Korkman M, Liikanen A, Fellman V. Neuropsychological consequences of very low birth weight and asphyxia at term: follow-up until school-age. J Clin Exp Neuropsychol 1996 Apr;18(2):220-33.

Krageloh-Mann I, Toft P, Lunding J et al. Brain lesions in preterms: origin, consequences and compensation. Acta Paediatr 1999 Aug;88(8):897-908.

Kraybill EN, Bose CL, Corbet AJ et al. Double-blind evaluation of developmental and health status to age 2 years of infants weighing 700 to 1350 grams treated prophylactically at birth with a single dose of synthetic surfactant or air placebo. J Pediatr 1995 May;126(5 Pt 2):S33-S42.

Kuban KC, Leviton A. Cerebral palsy. N Engl J Med 1994 Jan 20;330(3):188-95.

Kurkinen-Raty M, Koivisto M, Jouppila P. Perinatal and neonatal outcome and late pulmonary sequelae in infants born after preterm premature rupture of membranes. Obstet Gynecol 1998 Sep;92(3):408-15.

Kurkinen-Raty M, Koivisto M, Jouppila P. Preterm delivery for maternal or fetal indications: maternal morbidity, neonatal outcome and late sequelae in infants. BJOG 2000 May;107(5):648-55.

Ladd AP, Rescorla FJ, West KW et al. Long-term follow-up after bowel resection for necrotizing enterocolitis: factors affecting outcome. J Pediatr Surg 1998 Jul;33(7):967-72.

Lee BE, Cheung PY, Robinson JL et al. Comparative study of mortality and morbidity in premature infants (birth weight, < 1,250 g) with candidemia or candidal meningitis. Clin Infect Dis 1998 Sep;27(3):559-65.

Lefebvre F, Gregoire MC, Dubois J et al. Nursery Neurobiologic Risk Score and outcome at 18 months. Acta Paediatr 1998 Jul;87(7):751-7.

Leonard CH, Clyman RI, Piecuch RE et al. Effect of medical and social risk factors on outcome of prematurity and very low birth weight. J Pediatr 1990 Apr;116(4):620-6.

Levene MI. Cerebral ultrasound and neurological impairment: telling the future. Arch Dis Child 1990 May;65(5):469-71.

Ling CS, Fleck BW, Wright E et al. Diode laser treatment for retinopathy of prematurity: structural and functional outcome. Br J Ophthalmol 1995;79(7):637-41.

Lucas A, Morley R, Cole TJ et al. Breast milk and subsequent intelligence quotient in children born preterm. Lancet 1992;339(8788):261-4.

Lucas A, Fewtrell MS, Morley R et al. Randomized outcome trial of human milk fortification and developmental outcome in preterm infants. Am J Clin Nutr 1996;64(2):142-51.

Mandich M, Simons CJ, Ritchie S et al. Motor development, infantile reactions and postural responses of preterm, at-risk infants. Dev Med Child Neurol 1994;36(5):397-405.

Marlow ES, Hunt LP, Marlow N. Sensorineural hearing loss and prematurity. Arch Dis Child Fetal Neonatal Ed 2000 Mar;82(2):F141-F144.

Marlow N, Roberts L, Cooke R. Outcome at 8 years for children with birth weights of 1250 g or less. Arch Dis Child 1993 Mar;68(3 Spec No):286-90.

Mattia FR, deRegnier RA. Chronic physiologic instability is associated with neurodevelopmental morbidity at one and two years in extremely premature infants. Pediatrics 1998;102(3):E35.

McCormick MC, Gortmaker SL, Sobol AM. Very low birth weight children: behavior problems and school difficulty in a national sample. J Pediatr 1990 Nov;117(5):687-93.

McCormick MC, Brooks-Gunn J, Workman-Daniels K et al. The health and developmental status of very low-birth-weight children at school age. JAMA 1992 Apr 22;267(16):2204-8.

Ment LR, Duncan CC, Ehrenkranz RA et al. Randomized low-dose indomethacin trial for prevention of intraventricular hemorrhage in very low birth weight neonates. J Pediatr 1988;112(6):948-55.

Ment LR, Oh W, Ehrenkranz RA et al. Low-dose indomethacin and prevention of intraventricular hemorrhage: a multicenter randomized trial. Pediatrics 1994 Apr;93(4):543-50.

Ment LR, Oh W, Ehrenkranz RA et al. Low-dose indomethacin therapy and extension of intraventricular hemorrhage: a multicenter randomized trial. J Pediatr 1994;124(6):951-5.

Ment LR, Vohr B, Oh W et al. Neurodevelopmental outcome at 36 months' corrected age of preterm infants in the Multicenter Indomethacin Intraventricular Hemorrhage Prevention Trial. Pediatrics 1996 Oct;98(4 Pt 1):714-8.

Ment LR, Westerveld M, Makuch R et al. Cognitive outcome at 4 1/2 years of very low birth weight infants enrolled in the multicenter indomethacin intraventricular hemorrhage prevention trial. Pediatrics 1998;102(1 Pt 1):159-60.

Ment LR, Vohr B, Allan W et al. The etiology and outcome of cerebral ventriculomegaly at term in very low birth weight preterm infants. Pediatrics 1999 Aug;104(2 Pt 1):243-8.

Ment LR, Vohr B, Allan W et al. Outcome of children in the indomethacin intraventricular hemorrhage prevention trial. Pediatrics 2000 Mar;105(3 Pt 1):485-91.

Millet V, Bartoli JM, Lacroze V et al. Predictive significance of magnetic resonance imaging at 4 months of adjusted age in infants after a perinatal neurologic insult. Biol Neonate 1998;73(4):207-19.

Mitchell SH, Teague WG. Reduced gas transfer at rest and during exercise in school-age survivors of bronchopulmonary dysplasia. Am J Respir Crit Care Med 1998 May;157(5 Pt 1):1406-12.

Morley R, Lucas A. Randomized diet in the neonatal period and growth performance until 7.5-8 y of age in preterm children. Am J Clin Nutr 2000 Mar;71(3):822-8.

Murphy DJ, Hope PL, Johnson A. Neonatal risk factors for cerebral palsy in very preterm babies: case-control study. BMJ 1997 Feb 8;314(7078):404-8.

Mutch L, Alberman E, Hagberg B et al. Cerebral palsy epidemiology: where are we now and where are we going? Dev Med Child Neurol 1992 Jun;34(6):547-51.

Myers VS, Gidlewski N, Quinn GE et al. Distance and near visual acuity, contrast sensitivity, and visual fields of 10-year-old children. Arch Ophthalmol 1999 Jan;117(1):94-9.

Nadeau L, Boivin M, Tessier R et al. Mediators of behavioral problems in 7-year-old children born after 24 to 28 weeks of gestation. J Dev Behav Pediatr 2001;22(1):1-10.

Northern Neonatal Nursing Initiative Trial Group. Randomised trial of prophylactic early fresh-frozen plasma or gelatin or glucose in preterm babies: outcome at 2 years. Lancet 1996 Jul 27;348(9022):229-32.

O'Connor AR, Stephenson T, Johnson A et al. Long-term ophthalmic outcome of low birth weight children with and without retinopathy of prematurity. Pediatrics 2002 Jan;109(1):12-8.

O'Connor DL, Hall R, Adamkin D et al. Growth and development in preterm infants fed long-chain polyunsaturated fatty acids: a prospective, randomized controlled trial. Pediatrics 2001;108(2):359-71.

O'Shea TM, Klinepeter KL, Goldstein DJ et al. Survival and developmental disability in infants with birth weights of 501 to 800 grams, born between 1979 and 1994. Pediatrics 1997;100(6):982-6.

O'Shea TM, Klinepeter KL, Dillard RG. Prenatal events and the risk of cerebral palsy in very low birth weight infants. Am J Epidemiol 1998;147(4):362-9.

O'Shea TM, Preisser JS, Klinepeter KL et al. Trends in mortality and cerebral palsy in a geographically based cohort of very low birth weight neonates born between 1982 to 1994. Pediatrics 1998 Apr;101(4 Pt 1):642-7.

O'Shea TM, Kothadia JM, Klinepeter KL et al. Randomized placebo-controlled trial of a 42-day tapering course of dexamethasone to reduce the duration of ventilator dependency in very low birth weight infants: outcome of study participants at 1-year adjusted age. Pediatrics 1999 Jul;104(1 Pt 1):15-21.

Oehler JM, Goldstein RF, Catlett A et al. How to target infants at highest risk for developmental delay. MCN Am J Matern Child Nurs 1993;18(1):20-3.

Oehler JM, Thompson RJ, Jr., Goldstein RF et al. Behavioral characteristics of very-low-birth-weight infants of varying biologic risk at 6, 15, and 24 months of age. J Obstet Gynecol Neonatal Nurs 1996;25(3):233-9.

Page JM, Schneeweiss S, Whyte HE et al. Ocular sequelae in premature infants. Pediatrics 1993;92(6):787-90.

Palfrey JS, Singer JD, Walker DK et al. Early identification of children's special needs: a study in five metropolitan communities. J Pediatr 1987 Nov;111(5):651-9.

Palmer EA, Flynn JT, Hardy RJ et al. Incidence and early course of retinopathy of prematurity. The Cryotherapy for Retinopathy of Prematurity Cooperative Group. Ophthalmology 1991;98(11):1628-40.

Palta M, Sadek-Badawi M, Evans M et al. Functional assessment of a multicenter very low-birth-weight cohort at age 5 years. Arch Pediatr Adolesc Med 2000;154(1):23-30.

Pasman JW, Rotteveel JJ, Maassen B et al. Neonatal risk factors and risk scores including auditory evoked responses. Eur J Pediatr 1998;157(3):230-5.

Pelkonen AS, Hakulinen AL, Turpeinen M. Bronchial lability and responsiveness in school children born very preterm. Am J Respir Crit Care Med 1997 Oct;156(4 Pt 1):1178-84.

Pennefather PM, Clarke MP, Strong NP et al. Risk factors for strabismus in children born before 32 weeks' gestation. Br J Ophthalmol 1999 May;83(5):514-8.

Pennefather PM, Tin W. Ocular abnormalities associated with cerebral palsy after preterm birth. Eye 2000 Feb;14 (Pt 1):78-81.

Piecuch RE, Leonard CH, Cooper BA et al. Outcome of extremely low birth weight infants (500 to 999 grams) over a 12-year period. Pediatrics 1997 Oct;100(4):633-9.

Piecuch RE, Leonard CH, Cooper BA et al. Outcome of infants born at 24-26 weeks' gestation: II. Neurodevelopmental outcome. Obstet Gynecol 1997 Nov;90(5):809-14.

Piecuch RE, Leonard CH, Cooper BA. Infants with birth weight 1,000-1,499 grams born in three time periods: has outcome changed over time? [see comments]. Clin Pediatr (Phila) 1998;37(9):537-45.

Pierrat V, Duquennoy C, van Haastert IC et al. Ultrasound diagnosis and neurodevelopmental outcome of localised and extensive cystic periventricular leucomalacia. Arch Dis Child Fetal Neonatal Ed 2001 May;84(3):F151-F156.

Rademaker KJ, Groenendaal F, Jansen GH et al. Unilateral haemorrhagic parenchymal lesions in the preterm infant: shape, site and prognosis. Acta Paediatr 1994 Jun;83(6):602-8.

Ramer JC, Miller G. Overview of mental retardation. In: Miller G and Ramer JC, editors. Static encephalopathies of infancy and childhood. New York: Raven Press Ltd; 1992. p. 1-10.

Redline RW, Wilson-Costello D, Borawski E et al. Placental lesions associated with neurologic impairment and cerebral palsy in very low-birth-weight infants. Arch Pathol Lab Med 1998 Dec;122(12):1091-8.

Repka MX, Summers CG, Palmer EA et al. The incidence of ophthalmologic interventions in children with birth weights less than 1251 grams. Results through 5 1/2 years. Ophthalmology 1998;105(9):1621-7.

Robertson C, Sauve RS, Christianson HE. Province-based study of neurologic disability among survivors weighing 500 through 1249 grams at birth. Pediatrics 1994;93(4):636-40.

Robson AL, Pederson DR. Predictors of individual differences in attention among low birth weight children. J Dev Behav Pediatr 1997 Feb;18(1):13-21.

Rogers B, Andrus J, Msall ME et al. Growth of preterm infants with cystic periventricular leukomalacia. Dev Med Child Neurol 1998 Sep;40(9):580-6.

Roth S, Amess P, Kirkbride V et al. A comparision of linear-array and mechanical-sector cranial ultrasound scanning techniques to predict neurodevelopmental outcome at 8 years in preterm newborn infants. Ultrasound Med Biol 2001 Apr;27(4):467-72.

Saarela T, Vaarala A, Lanning P et al. Incidence, ultrasonic patterns and resolution of nephrocalcinosis in very low birthweight infants. Acta Paediatr 1999;88(6):655-60.

Saarela T, Lanning P, Koivisto M. Prematurity-associated nephrocalcinosis and kidney function in early childhood. Pediatr Nephrol 1999;13(9):886-90.

Saarela T, Lanning P, Koivisto M et al. Nephrocalcinosis in full-term infants receiving furosemide treatment for congestive heart failure: a study of the incidence and 2-year follow up. Eur J Pediatr 1999;158(8):668-72.

Saigal S, Stoskopf BL, Streiner DL et al. Physical growth and current health status of infants who were of extremely low birth weight and controls at adolescence. Pediatrics 2001 Aug;108(2):407-15.

Sajaniemi N, Salokorpi T, von Wendt L. Temperament profiles and their role in neurodevelopmental assessed preterm children at two years of age. Eur Child Adolesc Psychiatry 1998 Sep;7(3):145-52.

Sajaniemi N, Hakamies-Blomqvist L, Makela J et al. Cognitive development, temperament and behavior at 2 years as indicative of language development at 4 years in pre-term infants. Child Psychiatry Hum Dev 2001;31(4):329-46.

Salokorpi T, Rajantie I, Viitala J et al. Does perinatal hypocarbia play a role in the pathogenesis of cerebral palsy? Acta Paediatr 1999 May;88(5):571-5.

Santuz P, Baraldi E, Zaramella P et al. Factors limiting exercise performance in long-term survivors of bronchopulmonary dysplasia. Am J Respir Crit Care Med 1995 Oct;152(4 Pt 1):1284-9.

Schaffer DB, Palmer EA, Plotsky DF et al. Prognostic factors in the natural course of retinopathy of prematurity. The Cryotherapy for Retinopathy of Prematurity Cooperative Group. Ophthalmology 1993;100(2):230-7.

Schendel DE, Stockbauer JW, Hoffman HJ et al. Relation between very low birth weight and developmental delay among preschool children without disabilities. Am J Epidemiol 1997 Nov 1;146(9):740-9.

Scherjon SA, Oosting H, Smolders-DeHaas H et al. Neurodevelopmental outcome at three years of age after fetal 'brain-sparing'. Early Hum Dev 1998;52(1):67-79.

Schmidt B, Davis P, Moddemann D et al. Long-term effects of indomethacin prophylaxis in extremely-low-birth-weight infants. N Engl J Med 2001 Jun 28;344(26):1966-72.

Scrutton D. The classification of the cerebral palsies. Dev Med Child Neurol 1992 Sep;34(9):833.

Sethi VD, Macfarlane PI. Neurodevelopmental outcome at age two years amongst very low birth weight infants: results from a district general hospital. Public Health 1996 Jul;110(4):211-4.

Shalev B, Farr AK, Repka MX. Randomized comparison of diode laser photocoagulation versus cryotherapy for threshold retinopathy of prematurity: seven-year outcome. Am J Ophthalmol 2001;132(1):76-80.

Shennan AT, Dunn MS, Ohlsson A et al. Abnormal pulmonary outcomes in premature infants: prediction from oxygen requirement in the neonatal period. Pediatrics 1988 Oct;82(4):527-32.

Shepherd AJ, Saunders KJ, McCulloch DL et al. Prognostic value of flash visual evoked potentials in preterm infants. Dev Med Child Neurol 1999;41(1):9-15.

Shinwell ES, Karplus M, Reich D et al. Early postnatal dexamethasone treatment and increased incidence of cerebral palsy. Arch Dis Child Fetal Neonatal Ed 2000 Nov;83(3):F177-F181.

Singer L, Yamashita T, Lilien L et al. A longitudinal study of developmental outcome of infants with

bronchopulmonary dysplasia and very low birth weight. Pediatrics 1997 Dec;100(6):987-93.

Singer LT, Siegel AC, Lewis B et al. Preschool language outcomes of children with history of bronchopulmonary dysplasia and very low birth weight. J Dev Behav Pediatr 2001 Feb;22(1):19-26.

Slonim AD, Patel KM, Ruttimann UE et al. The impact of prematurity: a perspective of pediatric intensive care units. Crit Care Med 2000;28(3):848-53.

Smith KE, Landry SH, Swank PR et al. The relation of medical risk and maternal stimulation with preterm infants' development of cognitive, language and daily living skills. J Child Psychol Psychiatry 1996 Oct;37(7):855-64.

Speechley KN, Avison WR. Admission to a neonatal intensive care unit as a predictor of long-term health: a 12-year follow-up. J Dev Behav Pediatr 1995;16(6):397-405.

Spinillo A, Fazzi E, Stronati M et al. Early morbidity and neurodevelopmental outcome in low-birthweight infants born after third trimester bleeding. Am J Perinatol 1994 Mar;11(2):85-90.

Spinillo A, Fazzi E, Capuzzo E et al. Meconium-stained amniotic fluid and risk for cerebral palsy in preterm infants. Obstet Gynecol 1997;90(4 Pt 1):519-23.

Spinillo A, Capuzzo E, Orcesi S et al. Antenatal and delivery risk factors simultaneously associated with neonatal death and cerebral palsy in preterm infants. Early Hum Dev 1997 Apr 25;48(1-2):81-91.

Spinillo A, Capuzzo E, Cavallini A et al. Preeclampsia, preterm delivery and infant cerebral palsy. Eur J Obstet Gynecol Reprod Biol 1998;77(2):151-5.

Stathis SL, O'Callaghan M, Harvey J et al. Head circumference in ELBW babies is associated with learning difficulties and cognition but not ADHD in the school-aged child. Dev Med Child Neurol 1999 Jun;41(6):375-80.

Subhedar NV, Shaw NJ. Changes in pulmonary arterial pressure in preterm infants with chronic lung disease. Arch Dis Child Fetal Neonatal Ed 2000 May;82(3):F243-F247.

Task Force for the Handbook of Psychiatric Measures. Handbook of Psychiatric Measures. American Psychiatric Association; 2000

The Victorian Infant Collaborative Study Group. Improvement of outcome for infants of birth weight under 1000 g. Arch Dis Child 1991;66(7 Spec No):765-9.

The Victorian Infant Collaborative Study Group. Changing outcome for infants of birth-weight 500-999 g born outside level 3 centres in Victoria. Aust N Z J Obstet Gynaecol 1997;37(3):253-7.

The Victorian Infant Collaborative Study Group. Outcome at 2 years of children 23-27 weeks' gestation born in Victoria in 1991-92. J Paediatr Child Health 1997 Apr;33(2):161-5.

The Victorian Infant Collaborative Study Group. Improved outcome into the 1990s for infants weighing 500-999 g at birth. Arch Dis Child Fetal Neonatal Ed 1997 Sep;77(2):F91-F94.

Thompson Jr. RJ, Gustafson KE, Oehler JM et al. Developmental outcome of very low birth weight infants at four years of age as a function of biological risk and psychosocial risk. J Dev Behav Pediatr 1997;18(2):91-6.

Thompson CM, Buccimazza SS, Webster J et al. Infants of less than 1250 grams birth weight at Groote Schuur Hospital: outcome at 1 and 2 years of age. Pediatrics 1993 May;91(5):961-8.

Thompson RJ, Jr., Goldstein RF, Oehler JM et al. Developmental outcome of very low birth weight infants as a function of biological risk and psychosocial risk. J Dev Behav Pediatr 1994;15(4):232-8.

Thompson RJ, Jr., Gustafson KE, Oehler JM et al. Developmental outcome of very low birth weight infants at four years of age as a function of biological risk and psychosocial risk. J Dev Behav Pediatr 1997 Apr;18(2):91-6.

Thorndike RL, Hagen EP, Sattler JM. The Stanford-Binet Intelligence Scale: Guide for Administration and Scoring. 4thed. Chicago, IL: Riverside Publishing Company; 1986

Tin W, Fritz S, Wariyar U et al. Outcome of very preterm birth: children reviewed with ease at 2 years

differ from those followed up with difficulty. Arch Dis Child Fetal Neonatal Ed 1998;79(2):83-7.

Tin W, Milligan DW, Pennefather P et al. Pulse oximetry, severe retinopathy, and outcome at one year in babies of less than 28 weeks gestation. Arch Dis Child Fetal Neonatal Ed 2001 Mar;84(2):F106-F110.

Torrioli MG, Frisone MF, Bonvini L et al. Perceptual-motor, visual and cognitive ability in very low birthweight preschool children without neonatal ultrasound abnormalities. Brain Dev 2000;22(3):163-8.

Valkama AM, Paakko EL, Vainionpaa LK et al. Magnetic resonance imaging at term and neuromotor outcome in preterm infants. Acta Paediatr 2000 Mar;89(3):348-55.

van Wassenaer AG, Kok JH, Briet JM et al. Thyroid function in preterm newborns; is T4 treatment required in infants < 27 weeks' gestational age? Exp Clin Endocrinol Diabetes 1997;105 Suppl 4:12-8.

van Wassenaer AG, Kok JH, de Vijlder JJ et al. Effects of thyroxine supplementation on neurologic development in infants born at less than 30 weeks' gestation. N Engl J Med 1997 Jan 2;336(1):21-6.

Vohr B, Allan WC, Scott DT et al. Early-onset intraventricular hemorrhage in preterm neonates: incidence of neurodevelopmental handicap. Semin Perinatol 1999 Jun;23(3):212-7.

Vohr BR, Wright LL, Dusick AM et al. Neurodevelopmental and functional outcomes of extremely low birth weight infants in the National Institute of Child Health and Human Development Neonatal Research Network, 1993-1994. Pediatrics 2000 Jun;105(6):1216-26.

Volpe JJ. Neurobiology of periventricular leukomalacia in the premature infant. Pediatr Res 2001 Nov;50(5):553-62.

Wang JS, Cherng WS, Chen SN. Nursery neurobiologic risk score as a predictor of mortality in premature infants. Zhonghua Min Guo Xiao Er Ke Yi Xue Hui Za Zhi 1994;35(4):288-93.

Wang Z, Sauve RS. Assessment of postneonatal growth in VLBW infants: selection of growth references and age adjustment for prematurity. Can J Public Health 1998 Mar;89(2):109-14.

Wauben IP, Atkinson SA, Shah JK et al. Growth and body composition of preterm infants: influence of nutrient fortification of mother's milk in hospital and breastfeeding post-hospital discharge. Acta Paediatr 1998 Jul;87(7):780-5.

Whitaker AH, Feldman JF, Van Rossem R et al. Neonatal cranial ultrasound abnormalities in low birth weight infants: relation to cognitive outcomes at six years of age. Pediatrics 1996 Oct;98(4 Pt 1):719-29.

Wildin SR, Anderson A, Woodside M et al. Prediction of 12-month neurodevelopmental outcome from a 6-month neurologic examination in premature infants. Clin Pediatr (Phila) 1995 Jun;34(6):290-9.

Wildin SR, Smith K, Anderson A et al. Prediction of developmental patterns through 40 months from 6- and 12-month neurologic examinations in very low birth weight infants. J Dev Behav Pediatr 1997 Aug;18(4):215-21.

Wilkinson I, Bear J, Smith J et al. Neurological outcome of severe cystic periventricular leukomalacia. J Paediatr Child Health 1996 Oct;32(5):445-9.

Wodrich D. Children's Psychological Testing: A Guide for Nonpsychologists. Baltimore: Paul H. Brookes Publishing Co.; 1984

Wolke D, Meyer R. Cognitive status, language attainment, and prereading skills of 6-year-old very preterm children and their peers: the Bavarian Longitudinal Study. Dev Med Child Neurol 1999 Feb;41(2):94-109.

Wood NS, Marlow N, Costeloe K et al. Neurologic and developmental disability after extremely preterm birth. EPICure Study Group. N Engl J Med 2000 Aug 10;343(6):378-84.