Bibliography of Included Studies

Agren JJ, Hanninen O, Laitinen M, Seppanen K, Bernhardt I, Fogelholm L et al. Boreal freshwater fish diet modifies the plasma lipids and prostanoids and membrane fatty acids in man. Lipids 1988; 23(10):924-929.

Agren JJ, Pekkarinen H, Litmanen H, Hanninen O. Fish diet and physical fitness in relation to membrane and serum lipids, prostanoid metabolism and platelet aggregation in female students. European Journal of Applied Physiology & Occupational Physiology 1991; 63(5):393-398.

Agren JJ, Hanninen O, Julkunen A, Fogelholm L, Vidgren H, Schwab U et al. Fish diet, fish oil and docosahexaenoic acid rich oil lower fasting and postprandial plasma lipid levels. European Journal of Clinical Nutrition 1996; 50(11):765-771.

Agren JJ, Vaisanen S, Hanninen O, Muller AD, Hornstra G. Hemostatic factors and platelet aggregation after a fish-enriched diet or fish oil or docosahexaenoic acid supplementation. Prostaglandins Leukotrienes & Essential Fatty Acids 1997; 57(4-5):419-421.

Alaswad K, Pogson GW, Harris WS, Sherwani K, Bell HH. Effects of phenytoin, niacin, gemfibrozil, and omega-3 fatty acids on high density lipoproteins in patients with hypoalphalipoproteinemia. Preventive Cardiology 1999; 2(4):144-150.

Allman-Farinelli MA, Hall D, Kingham K, Pang D, Petocz P, Favaloro EJ. Comparison of the effects of two low fat diets with different alpha-linolenic:linoleic acid ratios on coagulation and fibrinolysis. Atherosclerosis 1999; 142(1):159-168.

Angerer P, Kothny W, Stork S, von Schacky C. Effect of dietary supplementation with omega-3 fatty acids on progression of atherosclerosis in carotid arteries. Cardiovascular Research 2002; 54(1):183-190.

Bairati I, Roy L, Meyer F. Double-blind, randomized, controlled trial of fish oil supplements in prevention of recurrence of stenosis after coronary angioplasty. Circulation **1992a**; 85(3):950-956.

Bairati I, Roy L, Meyer F. Effects of a fish oil supplement on blood pressure and serum lipids in patients treated for coronary artery disease. Canadian Journal of Cardiology **1992b**; 8(1):41-46.

Balestrieri GP, Maffi V, Sleiman I, Spandrio S, Di Stefano O, Salvi A et al. Fish oil supplementation in patients with heterozygous familial hypercholesterolemia. Recenti Progressi in Medicina 1996; 87(3):102-105.

Bellamy CM, Schofield PM, Faragher EB, Ramsdale DR. Can supplementation of diet with omega-3 polyunsaturated fatty acids reduce coronary angioplasty restenosis rate? European Heart Journal 1992; 13(12):1626-1631.

Bemelmans WJE, Lefrandt JD, Feskens EJM, Broer J, Tervaert JWC, May JF et al. Change in saturated fat intake is associated with progression of carotid and femoral intima-media thickness, and with levels of soluble intercellular adhesion molecule-1. Atherosclerosis 2002; 163(1):113-120.

Berrettini M, Parise P, Ricotta S, Iorio A, Peirone C, Nenci GG. Increased plasma levels of tissue factor pathway inhibitor (TFPI) after n-3 polyunsaturated fatty acids supplementation in patients with chronic atherosclerotic disease. Thrombosis & Haemostasis 1996; 75(3):395-400.

Bonaa KH, Bjerve KS, Nordoy A. Docosahexaenoic and eicosapentaenoic acids in plasma phospholipids are divergently associated with high density lipoprotein in humans. Arteriosclerosis & Thrombosis 1992; 12(6):675-681.

Bonnema SJ, Jespersen LT, Marving J, Gregersen G. Supplementation with olive oil rather than fish oil increases small arterial compliance in diabetic patients. Diabetes Nutr Metab 1995; 8(2):81-87.

Borchgrevink CF, Skaga E, Berg KJ, Skjaeggestad O. Absence of prophylactic effect of linolenic acid in patients with coronary heart-disease. Lancet 1966; 2(7456):187-189.

Brox J, Olaussen K, Osterud B, Elvevoll EO, Bjornstad E, Brattebog G et al. A long-term seal- and cod-liver-oil supplementation in hypercholesterolemic subjects. Lipids 2001; 36(1):7-13.

Cairns JA, Gill J, Morton B, Roberts R, Gent M, Hirsh J et al. Fish oils and low-molecular-weight heparin for the reduction of restenosis after percutaneous transluminal coronary angioplasty. The EMPAR Study. Circulation 1996; 94(7):1553-1560.

Chan DC, Watts GF, Barrett PH, Beilin LJ, Mori TA. Effect of atorvastatin and fish oil on plasma high-sensitivity C-reactive protein concentrations in individuals with visceral obesity. Clinical Chemistry 2002; 48(6:Pt 1):877-883.

Chan DC, Watts GF, Mori TA, Barrett PH, Redgrave TG, Beilin LJ. Randomized controlled trial of the effect of n-3 fatty acid supplementation on the metabolism of apolipoprotein B-100 and chylomicron remnants in men with visceral obesity. Am J Clin Nutr 2003; 77(2):300-307.

Christensen JH, Gustenhoff P, Korup E, Aaroe J, Toft E, Moller J et al. Effect of fish oil on heart rate variability in survivors of myocardial infarction: a double blind randomised controlled trial. BMJ 1996; 312(7032):677-678.

Christensen JH, Korup E, Aaroe J, Toft E, Moller J, Rasmussen K et al. Fish consumption, n-3 fatty acids in cell membranes, and heart rate variability in survivors of myocardial infarction with left ventricular dysfunction. American Journal of Cardiology 1997; 79(12):1670-1673.

Christensen JH, Christensen MS, Dyerberg J, Schmidt EB. Heart rate variability and fatty acid content of blood cell membranes: a dose-response study with n-3 fatty acids. Am J Clin Nutr 1999; 70(3):331-337.

Cobiac L, Clifton PM, Abbey M, Belling GB, Nestel PJ. Lipid, lipoprotein, and hemostatic effects of fish vs fish-oil n-3 fatty acids in mildly hyperlipidemic males. American Journal of Clinical Nutrition 1991; 53(5):1210-1216.

Conquer JA, Cheryk LA, Chan E, Gentry PA, Holub BJ. Effect of supplementation with dietary seal oil on selected cardiovascular risk factors and hemostatic variables in healthy male subjects. Thromb Res 1999; 96(3):239-250.

de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I et al. Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. Lancet 1994; 343(8911):1454-1459.

Deck C, Radack K. Effects of modest doses of omega-3 fatty acids on lipids and lipoproteins in hypertriglyceridemic subjects. A randomized controlled trial. Archives of Internal Medicine 1989; 149(8):1857-1862.

Dehmer GJ, Popma JJ, van den Berg EK, Eichhorn EJ, Prewitt JB, Campbell WB et al. Reduction in the rate of early restenosis after coronary angioplasty by a diet supplemented with n-3 fatty acids. New England Journal of Medicine 1988; 319(12):733-740.

DeLany JP, Vivian VM, Snook JT, Anderson PA. Effects of fish oil on serum lipids in men during a controlled feeding trial. American Journal of Clinical Nutrition 1990; 52(3):477-485.

Deslypere JP. Influence of supplementation with N-3 fatty acids on different coronary risk factors in men--a placebo controlled study. Verhandelingen - Koninklijke Academie voor Geneeskunde van Belgie 1992; 54(3):189-216.

Djousse L, Folsom AR, Province MA, Hunt SC, Ellison RC. Dietary linolenic acid and carotid atherosclerosis: the National Heart, Lung, and Blood Institute Family Heart Study. Am J Clin Nutr 2003; 77(4):819-825.

Dunstan DW, Mori TA, Puddey IB, Beilin LJ, Burke V, Morton AR et al. The independent and combined effects of aerobic exercise and dietary fish intake on serum lipids and glycemic control in NIDDM. A randomized controlled study. Diabetes Care 1997; 20(6):913-921.

Dunstan DW, Mori TA, Puddey IB, Beilin LJ, Burke V, Morton AR et al. Exercise and fish intake: Effects on serum lipids and glycemic control for type 2 diabetics. Cardiology Review 1998; 15(8):34-37.

Dunstan DW, Mori TA, Puddey IB, Beilin LJ, Burke V, Morton AR et al. A randomised, controlled study of the effects of aerobic exercise and dietary fish on coagulation and fibrinolytic factors in type 2 diabetics. Thrombosis & Haemostasis 1999; 81(3):367-372.

Durrington PN, Bhatnagar D, Mackness MI, Morgan J, Julier K, Khan MA et al. An omega-3 polyunsaturated fatty acid concentrate administered for one year decreased triglycerides in simvastatin treated patients with coronary heart disease and persisting hypertriglyceridaemia. Heart (British Cardiac Society) 2001; 85(5):544-548.

Eritsland J, Arnesen H, Berg K, Seljeflot I, Abdelnoor M. Serum Lp(a) lipoprotein levels in patients with coronary artery disease and the influence of long-term n-3 fatty acid supplementation. Scandinavian Journal of Clinical & Laboratory Investigation **1995a**; 55(4):295-300.

Eritsland J, Arnesen H, Seljeflot I, Hostmark AT. Long-term metabolic effects of n-3 polyunsaturated fatty acids in patients with coronary artery disease. Am J Clin Nutr **1995b**; 61(4):831-836.

Eritsland J, Arnesen H, Seljeflot I, Kierulf P. Long-term effects of n-3 polyunsaturated fatty acids on haemostatic variables and bleeding episodes in patients with coronary artery disease. Blood Coagulation & Fibrinolysis **1995c**; 6(1):17-22.

Finnegan YE, Minihane AM, Leigh-Firbank EC, Kew S, Meijer GW, Muggli R et al. Plant- and marine-derived n-3 polyunsaturated fatty acids have differential effects on fasting and postprandial blood lipid concentrations and on the susceptibility of LDL to oxidative modification in moderately hyperlipidemic subjects. Am J Clin Nutr 2003; 77(4):783-795.

Franzen D, Schannwell M, Oette K, Hopp HW. A prospective, randomized, and double-blind trial on the effect of fish oil on the incidence of restenosis following PTCA. Catheterization & Cardiovascular Diagnosis 1993; 28(4):301-310.

Freese R, Mutanen M, Valsta LM, Salminen I. Comparison of the effects of two diets rich in monounsaturated fatty acids differing in their linoleic/alpha-linolenic acid ratio on platelet aggregation. Thrombosis & Haemostasis 1994; 71(1):73-77.

Freese R, Mutanen M. Small effects of linseed oil or fish oil supplementation on postprandial changes in hemostatic factors. Thrombosis Research **1997a**; 85(2):147-152.

Freese R, Mutanen M. Alpha-linolenic acid and marine long-chain n-3 fatty acids differ only slightly in their effects on hemostatic factors in healthy subjects. Am J Clin Nutr **1997b**; 66(3):591-598.

Gans RO, Bilo HJ, Weersink EG, Rauwerda JA, Fonk T, Popp-Snijders C et al. Fish oil supplementation in patients with stable claudication. American Journal of Surgery 1990; 160(5):490-495.

GISSI-Prevenzione Investigators. Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. Lancet 1999; 354(9177):447-455.

Green P, Fuchs J, Schoenfeld N, Leibovici L, Lurie Y, Beigel Y et al. Effects of fish-oil ingestion on cardiovascular risk factors in hyperlipidemic subjects in Israel: a randomized, double-blind crossover study. Am J Clin Nutr 1990; 52(6):1118-1124.

Grigg LE, Kay TW, Valentine PA, et al. Determinants of restenosis and lack of effect of dietary supplementation with eicosapentaenoic acid on the incidence of coronary artery restenosis after angioplasty. Journal of the American College of Cardiology 1989 Mar 1;13(3):665-72.

Grimsgaard S, Bonaa KH, Hansen JB, Nordoy A. Highly purified eicosapentaenoic acid and docosahexaenoic acid in humans have similar triacylglycerol-lowering effects but divergent effects on serum fatty acids. Am J Clin Nutr 1997; 66(3):649-659.

Grundt H, Nilsen DW, Hetland O, et al. Improvement of serum lipids and blood pressure during intervention with n-3 fatty acids was not associated with changes in insulin levels in subjects with combined hyperlipidaemia. Journal of Internal Medicine 1995 Mar;237(3):249-59.

Grundt H, Nilsen DW, Hetland O, et al. Atherothrombogenic risk modulation by n-3 fatty acids was not associated with changes in homocysteine in subjects with combined hyperlipidaemia. Thrombosis & Haemostasis 1999 Apr;81(4):561-5.

Haines AP, Sanders TA, Imeson JD, et al. Effects of a fish oil supplement on platelet function, haemostatic variables and albuminuria in insulin-dependent diabetics. Thrombosis Research 1986 Sep 15;43(6):643-55.

Hamazaki T, Sawazaki S, Asaoka E, et al. Docosahexaenoic acid-rich fish oil does not affect serum lipid concentrations of normolipidemic young adults. Journal of Nutrition 1996 Nov;126(11):2784-9.

Hanninen OO, Agren JJ, Laitinen MV, et al. Dose-response relationships in blood lipids during moderate freshwater fish diet. Annals of Medicine 1989 Jun;21(3):203-7.

Hansen JB, Olsen JO, Wilsgard L, et al. Effects of dietary supplementation with cod liver oil on monocyte thromboplastin synthesis, coagulation and fibrinolysis. Journal of Internal Medicine Supplement 1989;225(731):133-9.

Hansen JB, Olsen JO, Wilsgard L, et al. Comparative effects of prolonged intake of highly purified fish oils as ethyl ester or triglyceride on lipids, haemostasis and platelet function in normolipaemic men. European Journal of Clinical Nutrition **1993a** Jul;47(7):497-507.

Hansen JB, Berge LN, Svensson B, et al. Effects of cod liver oil on lipids and platelets in males and females. European Journal of Clinical Nutrition **1993b** Feb;47(2):123-31.

Harris WS, Ginsberg HN, Arunakul N, et al. Safety and efficacy of Omacor in severe hypertriglyceridemia. Journal of Cardiovascular Risk 1997 Oct;4(5-6):385-91.

Hendra TJ, Britton ME, Roper DR, et al. Effects of fish oil supplements in NIDDM subjects. Controlled study. Diabetes Care 1990 Aug;13(8):821-9.

Jain S, Gaiha M, Bhattacharjee J, et al. Effects of low-dose omega-3 fatty acid substitution in type-2 diabetes mellitus with special reference to oxidative stress--a prospective preliminary study. J Assoc Physicians India 2002 Aug;50:1028-33.

Jensen T, Stender S, Goldstein K, et al. Partial normalization by dietary cod-liver oil of increased microvascular albumin leakage in patients with insulin-dependent diabetes and albuminuria. New England Journal of Medicine 1989 Dec 7;321(23):1572-7.

Johansen O, Brekke M, Seljeflot I, et al. N-3 fatty acids do not prevent restenosis after coronary angioplasty: results from the CART study. Coronary Angioplasty Restenosis Trial. Journal of the American College of Cardiology 1999 May;33(6):1619-26.

Junker R, Kratz M, Neufeld M, et al. Effects of diets containing olive oil, sunflower oil, or rapeseed oil on the hemostatic system. Thrombosis & Haemostasis 2001 Feb;85(2):280-6.

Kaul U, Sanghvi S, Bahl VK, et al. Fish oil supplements for prevention of restenosis after coronary angioplasty. International Journal of Cardiology 1992 Apr;35(1):87-93.

Kwon JS, Snook JT, Wardlaw GM, et al. Effects of diets high in saturated fatty acids, canola oil, or safflower oil on platelet function, thromboxane B2 formation, and fatty acid composition of platelet phospholipids. American Journal of Clinical Nutrition 1991 Aug;54(2):351-8.

Leigh-Firbank EC, Minihane AM, Leake DS, et al. Eicosapentaenoic acid and docosahexaenoic acid from fish oils: differential associations with lipid responses. Br J Nutr 2002 May;87(5):435-45.

Leng GC, Lee AJ, Fowkes FG, et al. Randomized controlled trial of gamma-linolenic acid and eicosapentaenoic acid in peripheral arterial disease. Clinical Nutrition 1998 Dec;17(6):265-71.

Lungershausen YK, Abbey M, Nestel PJ, et al. Reduction of blood pressure and plasma triglycerides by omega-3 fatty acids in treated hypertensives. Journal of Hypertension 1994 Sep;12(9):1041-5.

Lungershausen YK, Howe PRC, Clifton PM, et al. Evaluation of an omega-3 fatty acid supplement in diabetics with microalbuminuria. Annals of the New York Academy of Sciences, Vol 827 (pp 369-381), 1997 1997.

Luo J, Rizkalla SW, Vidal H, et al. Moderate intake of n-3 fatty acids for 2 months has no detrimental effect on glucose metabolism and could ameliorate the lipid profile in type 2 diabetic men. Results of a controlled study. Diabetes Care 1998 May;21(5):717-24.

Mackness MI, Bhatnagar D, Durrington PN, et al. Effects of a new fish oil concentrate on plasma lipids and lipoproteins in patients with hypertriglyceridaemia. Eur J Clin Nutr 1994 Dec;48(12):859-65.

Madsen T, Christensen JH, Blom M, Schmidt EB. The effect of dietary n-3 fatty acids on serum concentrations of C-reactive protein: a dose-response study. British Journal of Nutrition 2003; 89(4):517-522.

Madsen T, Skou HA, Hansen VE, et al. C-reactive protein, dietary n-3 fatty acids, and the extent of coronary artery disease. American Journal of Cardiology 2001;88(10):1139-42.

Marckmann P, Bladbjerg EM, Jespersen J. Dietary fish oil (4 g daily) and cardiovascular risk markers in healthy men. Arteriosclerosis Thrombosis & Vascular Biology 1997 Dec;17(12):3384-91.

Maresta A, Balduccelli M, Varani E, et al. Prevention of postcoronary angioplasty restenosis by omega-3 fatty acids: main results of the Esapent for Prevention of Restenosis ITalian Study (ESPRIT). American Heart Journal 2002 Jun;143(6):E5.

McGrath LT;Brennan GM;Donnelly JP;Johnston GD;Hayes JR;McVeigh GE; Effect of dietary fish oil supplementation on peroxidation of serum lipids in patients with non-insulin dependent diabetes mellitus. Atherosclerosis 1996; 121(2):275-83.

McVeigh GE, Brennan GM, Johnston GD, et al. Dietary fish oil augments nitric oxide production or release in patients with type 2 (non-insulin-dependent) diabetes mellitus. Diabetologia 1993 Jan;36(1):33-8.

Mezzano D, Leighton F, Martinez C, et al. Complementary effects of Mediterranean diet and moderate red wine intake on haemostatic cardiovascular risk factors. European Journal of Clinical Nutrition 2001 Jun;55(6):444-51.

Milner MR, Gallino RA, Leffingwell A, et al. Usefulness of fish oil supplements in preventing clinical evidence of restenosis after percutaneous transluminal coronary angioplasty. American Journal of Cardiology 1989 Aug 1;64(5):294-9.

Misso NLA, Thompson PJ. Fish oil supplementation inhibits platelet aggregation and ATP release induced by platelet-activating factor and other agonists. Platelets 1995;6(5):275-82.

Mori TA, Vandongen R, Beilin LJ, et al. Effects of varying dietary fat, fish, and fish oils on blood lipids in a randomized controlled trial in men at risk of heart disease. Am J Clin Nutr 1994 May;59(5):1060-8.

Mori TA, Bao DQ, Burke V, et al. Dietary fish as a major component of a weight-loss diet: effect on serum lipids, glucose, and insulin metabolism in overweight hypertensive subjects. Am J Clin Nutr 1999 Nov;70(5):817-25.

Mori TA, Burke V, Puddey IB, et al. Purified eicosapentaenoic and docosahexaenoic acids have differential effects on serum lipids and lipoproteins, LDL particle size, glucose, and insulin in mildly hyperlipidemic men. Am J Clin Nutr 2000 May;71(5):1085-94.

Muller AD, van Houwelingen AC, Dam-Mieras MC, et al. Effect of a moderate fish intake on haemostatic parameters in healthy males. Thrombosis & Haemostasis 1989 Jun 30;61(3):468-73.

Natvig H, Borchgrevink CF, Dedichen J, et al. A controlled trial of the effect of linolenic acid on incidence of coronary heart disease. The Norwegian vegetable oil experiment of 1965-66. Scandinavian Journal of Clinical & Laboratory Investigation - Supplement 1968;105:1-20.

Nenseter MS, Osterud B, Larsen T, et al. Effect of Norwegian fish powder on risk factors for coronary heart disease among hypercholesterolemic individuals. Nutrition Metabolism & Cardiovascular Diseases 2000 Dec;10(6):323-30.

Nikkila M. Influence of fish oil on blood lipids in coronary artery disease. European Journal of Clinical Nutrition 1991 Apr;45(4):209-13.

Nilsen DW, Albrektsen G, Landmark K, et al. Effects of a high-dose concentrate of n-3 fatty acids or corn oil introduced early after an acute myocardial infarction on serum triacylglycerol and HDL cholesterol. Am J Clin Nutr 2001 Jul;74(1):50-6.

Nordoy A, Bonaa KH, Nilsen H, et al. Effects of Simvastatin and omega-3 fatty acids on plasma lipoproteins and lipid peroxidation in patients with combined hyperlipidaemia. Journal of Internal Medicine 1998 Feb;243(2):163-70.

Nordoy A, Bonaa KH, Sandset PM, et al. Effect of omega-3 fatty acids and simvastatin on hemostatic risk factors and postprandial hyperlipemia in patients with combined hyperlipemia. Arteriosclerosis Thrombosis & Vascular Biology 2000 Jan;20(1):259-65.

Nye ER, Ablett MB, Robertson MC, et al. Effect of eicosapentaenoic acid on restenosis rate, clinical course and blood lipids in patients after percutaneous transluminal coronary angioplasty. Australian & New Zealand Journal of Medicine 1990 Aug;20(4):549-52.

Osterud B, Elvevoll E, Barstad H, et al. Effect of marine oils supplementation on coagulation and cellular activation in whole blood. Lipids 1995 Dec;30(12):1111-8.

Pedersen H, Petersen M, Major-Pedersen A, Jensen T, Nielsen NS, Lauridsen ST, Marckmann P. Influence of fish oil supplementation on *in vivo* and *in vitro* oxidation resistance of low-density lipoprotein in type 2 diabetes. European Journal of Clinical Nutrition 2003; 57:716-20.

Prisco D, Paniccia R, Filippini M, et al. No changes in PAI-1 levels after four-month n-3 PUFA ethyl ester supplementation in healthy subjects. Thrombosis Research 1994 Nov 1;76(3):237-44.

Radack K, Deck C, Huster G. Dietary supplementation with low-dose fish oils lowers fibrinogen levels: a randomized, double-blind controlled study. Ann Intern Med 1989 Nov 1;111(9):757-8.

Radack K, Deck C, Huster G. The effects of low doses of n-3 fatty acid supplementation on blood pressure in hypertensive subjects. A randomized controlled trial. Archives of Internal Medicine 1991 Jun;151(6):1173-80.

Radack KL, Deck CC, Huster GA. n-3 fatty acid effects on lipids, lipoproteins, and apolipoproteins at very low doses: results of a randomized controlled trial in hypertriglyceridemic subjects. Am J Clin Nutr 1990 Apr;51(4):599-605.

Reis GJ, Boucher TM, Sipperly ME, et al. Randomised trial of fish oil for prevention of restenosis after coronary angioplasty. Lancet 1989 Jul 22;2(8656):177-81.

Rivellese AA, Maffettone A, Iovine C, et al. Long-term effects of fish oil on insulin resistance and plasma lipoproteins in NIDDM patients with hypertriglyceridemia. Diabetes Care 1996 Nov;19(11):1207-13.

Rossing P, Hansen BV, Nielsen FS, et al. Fish oil in diabetic nephropathy. Diabetes Care 1996 Nov;19(11):1214-9.

Sacks FM, Hebert P, Appel LJ, et al. Short report: the effect of fish oil on blood pressure and high-density lipoprotein-cholesterol levels in phase I of the Trials of Hypertension Prevention. Journal of Hypertension 1994 Feb;12(2):209-13.

Salachas A, Papadopoulos C, Sakadamis G, et al. Effects of a low-dose fish oil concentrate on angina, exercise tolerance time, serum triglycerides, and platelet function. Angiology 1994 Dec;45(12):1023-31.

Salonen R, Nikkari T, Seppanen K, et al. Effect of omega-3 fatty acid supplementation on platelet aggregability and platelet produced thromboxane. Thrombosis & Haemostasis 1987 Jun 3;57(3):269-72.

Schaefer EJ, Lichtenstein AH, Lamon-Fava S, et al. Effects of National Cholesterol Education Program Step 2 diets relatively high or relatively low in fish-derived fatty acids on plasma lipoproteins in middle-aged and elderly subjects. Am J Clin Nutr 1996 Feb;63(2):234-41.

Schectman G, Kaul S, Kissebah AH. Effect of fish oil concentrate on lipoprotein composition in NIDDM. Diabetes 1988 Nov;37(11):1567-73.

Schectman G, Kaul S, Kissebah AH. Heterogeneity of low density lipoprotein responses to fish-oil supplementation in hypertriglyceridemic subjects. Arteriosclerosis 1989; 9(3):345-354.

Seljeflot I, Arnesen H, Brude IR, et al. Effects of omega-3 fatty acids and/or antioxidants on endothelial cell markers. European Journal of Clinical Investigation 1998 Aug;28(8):629-35.

Silva JM, Souza I, Silva R, et al. The triglyceride lowering effect of fish oils is affected by fish consumption. International Journal of Cardiology 1996 May;57(1):75-80.

Singh RB, Dubnov G, Niaz MA, et al. Effect of an Indo-Mediterranean diet on progression of coronary artery disease in high risk patients (Indo-Mediterranean Diet Heart Study): A randomised single-blind trial. Lancet 2002;360(9344):1455-61.

Sirtori CR, Gatti E, Tremoli E, et al. Olive oil, corn oil, and n-3 fatty acids differently affect lipids, lipoproteins, platelets, and superoxide formation in type II hypercholesterolemia. Am J Clin Nutr 1992 Jul;56(1):113-22.

Sirtori CR, Crepaldi G, Manzato E, et al. One-year treatment with ethyl esters of n-3 fatty acids in patients with hypertriglyceridemia and glucose intolerance: reduced triglyceridemia, total cholesterol and increased HDL-C without glycemic alterations. Atherosclerosis 1998 Apr;137(2):419-27.

Solomon SA, Cartwright I, Pockley G, et al. A placebo-controlled, double-blind study of eicosapentaenoic acid-rich fish oil in patients with stable angina pectoris. Current Medical Research & Opinion 1990;12(1):1-11.

Swahn E, von Schenck H, Olsson AG. Omega-3 ethyl ester concentrate decreases total apolipoprotein CIII and increases antithrombin III in postmyocardial infarction patients. Clinical Drug Investigation 1998;15(6):473-82.

Toft I, Bonaa KH, Ingebretsen OC, et al. Effects of n-3 polyunsaturated fatty acids on glucose homeostasis and blood pressure in essential hypertension. A randomized, controlled trial. Ann Intern Med 1995 Dec 15;123(12):911-8.

Toft I, Bonaa KH, Ingebretsen OC, et al. Fibrinolytic function after dietary supplementation with omega3 polyunsaturated fatty acids. Arteriosclerosis Thrombosis & Vascular Biology 1997 May;17(5):814-9.

Toth K, Ernst E, Habon T, et al. Hemorheological and hemodynamical effects of fish oil (Ameu) in patients with ischemic heart disease and hyperlipoproteinemia. Clinical Hemorheology 1995;15(6):867-75.

Verheugt FWA, Schouten JA, Eeltink JC, et al. Omega-3 polyunsaturated fatty acids in the treatment of angina pectoris: Effect on objective signs of exercise-induced myocardial ischemia. Current Therapeutic Research, Clinical & Experimental 1986;39(2):208-13.

von Schacky C, Angerer P, Kothny W, et al. The effect of dietary omega-3 fatty acids on coronary atherosclerosis. A randomized, double-blind, placebo-controlled trial. Ann Intern Med 1999 Apr 6;130(7):554-62.

Warren SE, Siguel EN, Gervino E, et al. Effects of cod liver oil on plasma lipids, eicosanoids, platelet aggregation, and exercise in stable angina pectoris. Journal of Applied Cardiology 1988;3(4):227-36.

Wensing AG, Mensink RP, Hornstra G. Effects of dietary n-3 polyunsaturated fatty acids from plant and marine origin on platelet aggregation in healthy elderly subjects. Br J Nutr 1999 Sep;82(3):183-91.

Westerveld HT, de Graaf JC, van Breugel HH, et al. Effects of low-dose EPA-E on glycemic control, lipid profile, lipoprotein(a), platelet aggregation, viscosity, and platelet and vessel wall interaction in NIDDM. Diabetes Care 1993 May;16(5):683-8.

Wilt TJ. Lofgren RP. Nichol KL. Schorer AE. Crespin L. Downes D. Eckfeldt J. Fish oil supplementation does not lower plasma cholesterol in men with hypercholesterolemia. Results of a randomized, placebo-controlled crossover study. Annals of Internal Medicine 1989; 111(11):900-5

Woodman RJ, Mori TA, Burke V, et al. Effects of purified eicosapentaenoic and docosahexaenoic acids on glycemic control, blood pressure, and serum lipids in type 2 diabetic patients with treated hypertension. Am J Clin Nutr 2002;76(5):1007-15.

Yamada T, Malcom GT, Strong JP, et al. Difference in atherosclerosis between the populations of a fishing and a farming village in Japan. Ann N Y Acad Sci 1997;811:412-9.