Bibliography

Abotalib ZM, Soltlan MH, Chowdhury N, et al. Obstetric outcome in uncomplicated prolonged pregnancy. Int J Gynecol Obstet 1996;55(3):225-30.

Acker DB, Sachs BP, Friedman EA. Risk factors for shoulder dystocia. Obstet Gynecol 1985;66(6):762-8.

ACOG. Ultrasonography in pregnancy. Technical Bulletin No. 187. 1993.

ACOG. Induction of labor. Technical Bulletin No. 217. 1995.

ACOG. Routine ultrasound in low-risk pregnancy. 1997.

ACOG. Fetal macrosomia. ACOG Practice Bulletin No. 22. 2000.

Adair CD, Weeks JW, Barrilleaux PS, et al. Labor induction with oral versus vaginal misoprostol: A randomized, double-blind trial. Am J Obstet Gynecol 1998;178:93.

Adetoro OO, Agah A. The implications of childbearing in postpubertal girls in Sokoto, Nigeria. Int J Gynaecol Obstet 1988;27(1):73-7.

Agarwal N, Mathur T, Gupta N, et al. Role of intra-amniotic instillation of betamethasone in prolonged pregnancy. Indian J Med Res 1981;73(3):335-41.

Ahlden S, Andersch B, Stigsson L, et al. Prediction of sepsis neonatorum following a full-term pregnancy. Gynecol Obstet Invest 1988;25(3):181-5.

Ahmed AI, Versi E. Prolonged pregnancy. [Review]. Curr Opin Obstet Gynecol 1993;5(5):669-74.

Al-Najashi SS, Al-Mulhim AA. Prolongation of pregnancy in multiple pregnancy. Int J Gynecol Obstet 1996;54(2):131-5.

Alberico S, Fadalti M, Grimaldi E, et al. Eligibility criteria for labor induction with prostaglandins. Clin Exp Obstet Gynecol 1997;24(2):61-6.

Alexander JM, McIntire DD, Leveno KJ. Forty weeks and beyond: pregnancy outcomes by week of gestation. Obstet Gynecol 2000;96(2):291-4.

Alexander JM, Mcintire DD, Leveno KJ. Prolonged pregnancy: induction of labor and cesarean births. Obstet Gynecol 2001;97(6911-5).

Alfirevic Z, Howarth G, Gaussmann A. Oral misoprostol for induction of labor (Cochrane Review). In: The Cochrane Library, Issue 2, 2000. Oxford: Update Software.

Alfirevic Z, Luckas M, Walkinshaw SA, et al. A randomised comparison between amniotic fluid index and maximum pool depth in the monitoring of post-term pregnancy. Br J Obstet Gynaecol 1997;104(2):207-11.

Alfirevic Z, Neilson JP. Biophysical profile for fetal assessment in high risk pregnancies (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Alfirevic Z, Walkinshaw SA. Management of post-term pregnancy: to induce or not? [Review] . Br J Hosp Med 1994;52(5):218-21.

Alfirevic Z, Walkinshaw SA. A randomised controlled trial of simple compared with complex antenatal fetal monitoring after 42 weeks of gestation. Br J Obstet Gynaecol 1995;102(8):638-43.

Allott HA, Palmer CR. Sweeping the membranes: A valid procedure in stimulating the onset of labour? Br J Obstet Gynaecol 1993;100(10):898-903.

Almstrom H, Granstrom L, Ekman G. Serial antenatal monitoring compared with labor induction in post-term pregnancies. Acta Obstet Gynecol Scand 1995;74(8):599-603.

Amano K, Saito K, Shoda T, et al. Elective induction of labor at 39 weeks of gestation: A prospective randomized trial. J Obstet Gynaecol Res 1999;25(1):33-7.

Amiel-Tison C. Cerebral handicap in full-term newborns related to late pregnancy and/or labor. Eur J Obstet Gynecol Reprod Biol 1988;28(2):157-63.

Anderson T. Cochrane made simple. Routine ultrasound in early pregnancy. Pract Midwife 1998;1(12):13-4.

Anderson T. Cochrane made simple. Post-term pregnancy. Pract Midwife 1999;2(11):10-2.

Anderson T. Post-term pregnancy. [Review]. Pract Midwife 1999;2(11):10-2.

Anonymous. Task force on cerebral palsy and neonatal asphyxia. In: The Society of Obstetricians and Gynaecologists of Canada: Policy Statement, Committee Opinion, Clinical Practice Guidelines. Ottawa, ON : SOGC, 1996. 72 P.

Anonymous. Prolonged pregnancy: the management debate. BMJ 1988;297(6650):715-7.

Anonymous. Induction and augmentation of labor: ACOG Technical bulletin number 157 (replaces No. 110, November 1987). Int J Gynecol Obstet 1992;39(2):139-42.

Anonymous. The management of postpartum haemorrhage. Drug Ther Bull 1992;30(23):89-92.

Anonymous. Post-term induction: don't kill the messenger. Lancet 1992;340(8823):824.

Anonymous. Postterm pregnancy. Int J Gynecol Obstet 1996;53(1):89-90.

Anonymous. ACOG practice patterns. Management of postterm pregnancy. Number 6, October 1997. American College of Obstetricians and Gynecologists. Int J Gynaecol Obstet 1997;60(1):86-91.

Anonymous. Managing post-term pregnancy. [Review]. Drug Ther Bull 1997;35(3):17-8.

Anonymous. Electronic fetal heart rate monitoring: research guidelines for interpretation. The National Institute of Child Health and Human Development Research Planning Workshop. J Obstet Gynecol Neonat Nurs 1997;26(6):635-40.

Apgar J, Everett GA, Fitzgerald JA. Dietary zinc deprivation affects parturition and outcome of pregnancy in the ewe. Nutr Res 1993;13(3):319-30.

Arabin B, Becker R, Mohnhaupt A, et al. Prediction of fetal distress and poor outcome in prolonged pregnancy using Doppler ultrasound and fetal heart rate monitoring combined with stress tests (II). Fetal Diagn Ther 1994;9(1):1-6.

Arabin B, Snyjders R, Mohnhaupt A, et al. Evaluation of the fetal assessment score in pregnancies at risk for intrauterine hypoxia. Am J Obstet Gynecol 1993;169(3):549-54.

Arias F. Predictability of complications associated with prolongation of pregnancy. Obstet Gynecol 1987;70(1):101-6.

Ariceta JM, Matorras R, Diez J, et al. Sequential vacuum extractor and forceps delivery. J Obstet Gynaecol 1994;14(5):312-16.

Arntzen KJ, Brekke O-L, Vatten L, et al. Reduced production of PGEinf 2 and PGF(2alpha) from decidual cell cultures supplemented with N-3 polyunsaturated fatty acids. Prostaglandins Other Lipid Mediat 1998;56(2-3):183-95.

Arrowsmith S, Hamlin EC, Wall LL. Obstructed labor injury complex: Obstetric fistula formation and the multifaceted morbidity of maternal birth trauma in the developing world. Obstet Gynecol Surv 1996;51(9):568-74.

Atad J, Hallak M, Auslender R, et al. A randomized comparison of prostaglandin E2, oxytocin, and the double-balloon device in inducing labor. Obstet Gynecol 1996;87(2):223-7.

Augensen K, Bergsjø P, Eikeland T, et al. Randomised comparison of early versus late induction of labour in post-term pregnancy. BMJ 1987;294(6581):1192-5.

Ayres-de-Campos D, Bernardes J, Costa-Pereira A, et al. Inconsistencies in classification by experts of cardiotocograms and subsequent clinical decision. Br J Obstet Gynaecol 1999;106(12):1307-10.

Babinszki A, Kerenyi T, Torok O, et al. Perinatal outcome in grand and great-grand multiparity: effects of parity on obstetric risk factors. Am J Obstet Gynecol 1999;181(3):669-74.

Bailey SM, Sarmandal P, Grant JM. A comparison of three methods of assessing inter-observer variation applied to measurement of the symphysis-fundal height. Br J Obstet Gynaecol 1989;96(11):1266-71.

Baines DL, Folkesson HG, Norlin A, et al. The influence of mode of delivery, hormonal status and postnatal Oinf 2 environment on epithelial sodium channel (ENaC) expression in perinatal guinea-pig lung. J Physiol 2000;522(1):147-57.

Bakimer R, Blank M, Kosashvilli D, et al. Antiphospholipid syndrome and the idiotypic network. Lupus 1995;4(3):204-8.

Bakos O, Backstrom T. Induction of labor: a prospective, randomized study into amniotomy and oxytocin as induction methods in a total unselected population. Acta Obstet Gynecol Scand 1987;66(6):537-41.

Barrett JF, Tyrrell S, Lilford RJ. Prolonged pregnancy. BMJ 1987;294(6563):56-7.

Barss VA, Frigoletto FD, Diamond F. Stillbirth after nonstress testing. Obstet Gynecol 1985;65(4):541-4.

Bartnicki J, Ratanasiri T, Meyenburg M, et al. Postterm pregnancy: computer analysis of the antepartum fetal heart rate patterns. Int J Gynaecol Obstet 1992;37(4):243-6.

Barton DPJ, Robson MS, Turner MJ, et al. Prolonged spontaneous labour in primigravidae whose labour was actively managed: Results of an audit. J Obstet Gynaecol 1992;12(5):304-8.

Baskett TF. Gestational age and fetal biophysical assessment. Am J Obstet Gynecol 1988;158(2):332-4.

Baskett TF, Allen AC. Perinatal implications of shoulder dystocia. Obstet Gynecol 1995;86(1):14-7.

Bastian H, Keirse MJ, Lancaster PA. Perinatal death associated with planned home birth in Australia: population based study. BMJ 1998;317(7155):384-8.

Battaglia C, Artini PG, Ballestri M, et al. Hemodynamic, hematological and hemorrheological evaluation of post-term pregnancy. Acta Obstet Gynecol Scand 1995;74(5):336-40.

Battaglia C, Larocca E, Lanzani A, et al. Doppler velocimetry in prolonged pregnancy. Obstet Gynecol 1991;77(2):213-6.

Beck I, Clayton JK. Hazards of prostaglandin pessaries in postmaturity. Lancet 1982;2(8290):161.

Belizan JM, Villar J, Nardin JC, et al. Diagnosis of intrauterine growth retardation by a simple clinical method: measurement of uterine height. Am J Obstet Gynecol 1978;131(6):643-6.

Bell RJ, Permezel M, MacLennan A, et al. A randomized, double-blind, placebo-controlled trial of the safety of vaginal recombinant human relaxin for cervical ripening. Obstet Gynecol 1993;82(3):328-33.

Benacerraf BR, Frigoletto FD. Fetal respiratory movements: only part of the biophysical profile. Obstet Gynecol 1986;67(4):556-7.

Benedetti TJ, Easterling T. Antepartum testing in postterm pregnancy. J Reprod Med 1988;33(3):252-8.

Benifla JL, Goffinet F, Darai E, et al. Emergency cervical cerclage after 20 weeks' gestation: A retrospective study of 6 years' practice in 34 cases. Fetal Diagn Ther 1997;12(5):274-8.

Bennett BB. Uterine rupture during induction of labor at term with intravaginal misoprostol. Obstet Gynecol 1997;89(5 Pt 2):832-3.

Berghella V, Rogers RA, Lescale K. Stripping of membranes as a safe method to reduce prolonged pregnancies. Obstet Gynecol 1996;87(6):927-31.

Bergsjo P, Bakketeig LS, Eikhom SN. Case-control analysis of post-term induction of labor. Acta Obstet Gynecol Scand 1982;61(4):317-24.

Bergsjø P, Huang GD, Yu SQ, et al. Comparison of induced versus non-induced labor in post-term pregnancy. A randomized prospective study. Acta Obstet Gynecol Scand 1989;68(8):683-7.

Bernardes J, Costa-Pereira A, Ayres-de-Campos D, et al. Evaluation of interobserver agreement of cardiotocograms. Int J Gynaecol Obstet 1997;57(1):33-7.

Bernstein IM, Watson M, Simmons GM, et al. Maternal brain death and prolonged fetal survival. Obstet Gynecol 1989;74(3 II SUPPL.):434-7.

Berzosa J, Martinez-Guisasola J, Guerrero M, et al. Application of prostaglandin Einf 2 gel in post-term pregnancy (> 287 days). Clin Invest Ginecol Obstet 1996;23(3):101-5.

Bishop EH. Pelvic scoring for elective induction. Obstet Gynecol 1964;24:266-8.

Blakemore KJ, Qin NG, Petrie RH, et al. A prospective comparison of hourly and quarter-hourly oxytocin dose increase intervals for the induction of labor at term. Obstet Gynecol 1990;75(5):757-61.

Blanchette HA, Nayak S, Erasmus S. Comparison of the safety and efficacy of intravaginal misoprostol (prostaglandin E1) with those of dinoprostone

(prostaglandin E2) for cervical ripening and induction of labor in a community hospital. Am J Obstet Gynecol 1999;180(6 Pt 1):1551-9.

Blanco JD, Collins M, Willis D, et al. Prostaglandin E2 gel induction of patients with a prior low transverse cesarean section. Am J Perinatol 1992;9(2):80-3.

Bobby PD, Divon MY. Fetal testing in postdates. Curr Opin Obstet Gynecol 1997;9(2):79-82.

Bobrowski RA, Bottoms SF. Underappreciated risks of the elderly multipara. Am J Obstet Gynecol 1995;172(6):1764-7.

Bochner CJ, Medearis AL, Davis J, et al. Antepartum predictors of fetal distress in postterm pregnancy. Am J Obstet Gynecol 1987;157(2):353-8.

Bochner CJ, Medearis AL, Ross MG, et al. The role of antepartum testing in the management of postterm pregnancies with heavy meconium in early labor. Obstet Gynecol 1987;69(6):903-7.

Bochner CJ, Williams J 3d, Castro L, et al. The efficacy of starting postterm antenatal testing at 41 weeks as compared with 42 weeks of gestational age. Am J Obstet Gynecol 1988;159(3):550-4.

Bors-Koefoed R, Zylstra S, Resseguie LJ, et al. A cost analysis of ambulatory antenatal testing in a tertiary care center. J Matern Fetal Invest 1993;2(1):5-9.

Boulvain M, Fraser WD, Marcoux S, et al. Does sweeping of the membranes reduce the need for formal induction of labour? A randomised controlled trial. Br J Obstet Gynaecol 1998;105(1):34-40.

Boulvain M, Irion O. Stripping/sweeping the membranes for inducing labour or preventing post-term pregnancy (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Boulvain M, Irion O, Lohse C, et al. Mechanical methods for inducing labour (Cochrane Protocol). In: The Cochrane Library, Issue 4, 2000. Oxford: Update Software. (withdrawn).

Boulvain M, Irion O, Marcoux S, et al. Sweeping of the membranes to prevent post-term pregnancy and to induce labour: a systematic review. [Review]. Br J Obstet Gynaecol 1999;106(5):481-5.

Boulvain M, Stan C, Irion O. Membrane sweeping for induction of labour (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Bourgeois FJ, Harbert GM, Andersen WA, et al. Early versus late tocolytic treatment of preterm premature membrane rupture. Am J Obstet Gynecol 1988;159(3):742-8.

Boyd ME, Usher RH, McLean FH. Fetal macrosomia: prediction, risks, proposed management. Obstet Gynecol 1983;61(6):715-22.

Boyd ME, Usher RH, McLean FH, et al. Obstetric consequences of postmaturity. Am J Obstet Gynecol 1988;158(2):334-8.

Boylan P, McParland P. Fetal assessment in postterm pregnancy. [Review]. Curr Opin Obstet Gynecol 1991;3(1):41-4.

Brace RA, Wolf EJ. Normal amniotic fluid volume changes throughout pregnancy. Am J Obstet Gynecol 1989;161(2):382-8.

Bracero LA, Roshanfekr D, Byrne DW. Analysis of antepartum fetal heart rate tracing by physician and computer. J Matern Fetal Med 2000;9(3):181-5.

Brar HS, Horenstein J, Medearis AL, et al. Cerebral, umbilical, and uterine resistance using Doppler velocimetry in postterm pregnancy. J Ultrasound Med 1989;8(4):187-91.

Brennand JE, Calder AA, Leitch CR, et al. Recombinant human relaxin as a cervical ripening agent. Br J Obstet Gynaecol 1997;104(7):775-80.

Bresadola M, Lo Mastro F, Arena V, et al. Prognostic value of biophysical profile score in post-date pregnancy. Clin Exp Obstet Gynecol 1995;22(4):330-8.

Bricker L, Luckas M. Amniotomy alone for induction of labor (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Bricker L, Neilson JP. Routine Doppler ultrasound in pregnancy (Cochrane Review). In: The Cochrane Library, Issue 3, 2000. Oxford: Update Software.

Bricker L, Neilson JP. Routine ultrasound in late pregnancy (after 24 weeks gestation) (Cochrane Review). In: The Cochrane Library, Issue 3, 2000.

Oxford: Update Software.

Brooks GG, Lewis DF, Gallaspy JW, et al. Labor in the gravida with 10 or more years between pregnancies. J Reprod Med Obstet Gynecol 1992;37(4):336-8.

Brost BC, Newman RB, Hendricks SK, et al. Effect of hemodialysis on serum progesterone level in pregnant women. Am J Kidney Dis 1999;33(5):917-9.

Brown VA, Sawers RS, Parsons RJ, et al. The value of antenatal cardiotocography in the management of high-risk pregnancy: a randomized controlled trial. Br J Obstet Gynaecol 1982;89(9):716-22.

Buser D, Mora G, Arias F. A randomized comparison between misoprostol and dinoprostone for cervical ripening and labor induction in patients with unfavorable cervices. Obstet Gynecol 1997;89(4):581-5.

Buttino LT, Garite TJ. Intracervical prostaglandin in postdate pregnancy. A randomized trial. J Reprod Med 1990;35(2):155-8.

Cabbad MF, Minkoff H, Faustin D. Fetal heart rate decelerations after oxytocin infusion in an abdominal pregnancy. Obstet Gynecol 1985;66(3 Suppl):2S-4S.

Cammu H, Haitsma V. Sweeping of the membranes at 39 weeks in nulliparous women: a randomised controlled trial. Br J Obstet Gynaecol 1998;105(1):41-4.

Campbell MK. Factors affecting outcome in post-term birth. Curr Opin Obstet Gynecol 1997;9(6):356-60.

Campbell MK, Ostbye T, Irgens LM. Post-term birth: Risk factors and outcomes in a 10-year cohort of Norwegian births. Obstet Gynecol 1997;89(4):543-8.

Cardozo L. Is routine induction of labour at term ever justified? BMJ 1993;306(6881):840-1.

Cardozo L, Fysh J, Pearce JM. Prolonged pregnancy: the management debate. BMJ 1986;293(6554):1059-63.

Cario GM. Conservative management of prolonged pregnancy using fetal heart rate monitoring only: a prospective study. Br J Obstet Gynaecol 1984;91(1):23-30.

Carlan SJ, Angel JL, Knuppel RA. Shoulder dystocia. Am Fam Physician 1991;43(4):1307-11.

Carlomagno G, Candussi G, Zavino S, et al. Postmaturity: How far is it a clinical entity in its own right? Clin Exp Obstet Gynecol 1996;23(1):41-7.

Carr MH, Towers CV, Eastenson AR, et al. Prolonged bedrest during pregnancy: Does the risk of deep vein thrombosis warrant the use of routine heparin prophylaxis? J Matern Fetal Med 1997;6(5):264-7.

Castro LC, Arora CP, Roll KE, et al. Perinatal factors influencing atrial natriuretic peptide levels in umbilical arterial plasma at the time of delivery [published erratum appears in Am J Obstet Gynecol 1989 Nov;161(5):1238]. Am J Obstet Gynecol 1989;161(3):623-7.

Center for Reviews and Dissemination Reviewers. Antenatal ultrasound scanning. Database of Abstracts of Reviews of Effectiveness Issue 1, 2000 .

Center for Reviews and Dissemination Reviewers. Misoprostol for cervical ripening and labor induction: a meta-analysis. Database of Abstracts of Reviews of Effectiveness Issue 1, 2000 .

Center for Reviews and Dissemination Reviewers. Umbilical Doppler velocimetry: results of controlled trials in high risk and low risk populations. Database of Abstracts of Reviews of Effectiveness Issue 1, 2000 .

Chang CH, Chang FM. Randomized comparison of misoprostol and dinoprostone for preinduction cervical ripening and labor induction. J Formos Med Assoc 1997;96(5):366-9.

Chang TC, Tan KT, Neow P, et al. Computerised analysis of foetal heart rate variation: Prediction of adverse perinatal outcome in patients undergoing prostaglandin induction of labour at term. Ann Acad Med Singapore 1997;26(6):772-5.

Chari RS, Friedman SA, O'Brien JM, et al. Daily antenatal testing in women with severe preeclampsia. Am J Obstet Gynecol 1995;173(4):1207-10.

Chatterjee MS, Ramchandran K, Ferlita J, et al. Prostaglandin E2 (PGE2) vaginal gel for cervical ripening. Eur J Obstet Gynecol Reprod Biol 1991;38(3):197-202.

Chauhan SP, Hendrix NW, Magann EF, et al. Limitations of clinical and sonographic estimates of birth weight: experience with 1034 parturients. Obstet Gynecol 1998;91(1):72-7.

Chauhan SP, Lutton PM, Bailey KJ, et al. Intrapartum clinical, sonographic, and parous patients' estimates of newborn birth weight. Obstet Gynecol 1992;79(6):956-8.

Chauhan SP, Magann EF, Sullivan CA, et al. Amniotic fluid index as an admission test may increase the incidence of cesarean section in a community hospital. J Matern Fetal Invest 1994;4(4):233-6.

Chauhan SP, Sullivan CA, Lutton TC, et al. Parous patients' estimate of birth weight in postterm pregnancy. J Perinatol 1995;15(3):192-4.

Chauhan SP, Sullivan CA, Magann EF, et al. Estimate of birthweight among post-term pregnancy: Clinical versus sonographic. J Matern Fetal Med 1994;3(5):208-11.

Chayen B, Tejani N, Verma U. Induction of labor with an electric breast pump. J Reprod Med 1986;31(2):116-8.

Chervenak JL. Macrosomia in the postdates pregnancy. [Review] . Clin Obstet Gynecol 1992;35(1):151-5.

Chervenak JL, Divon MY, Hirsch J, et al. Macrosomia in the postdate pregnancy: is routine ultrasonographic screening indicated? Am J Obstet Gynecol 1989;161(3):753-6.

Chez RA, Jonas WB. Complementary and alternative medicine. Part I: Clinical studies in obstetrics. Obstet Gynecol Surv 1997;52(11):704-8.

Chi IC, Agoestina T, Harbin J. Maternal mortality at twelve teaching hospitals in Indonesia-an epidemiologic analysis. Int J Gynaecol Obstet 1981;19(4):259-66.

Choe JK, Baggish MS. Hysteroscopic treatment of septate uterus with Neodymium-YAG laser. Fertil Steril 1992;57(1):81-4.

Christensen KK, Ingemarsson I, Leideman T, et al. Effect of ritodrine on labor after premature rupture of the membranes. Obstet Gynecol 1980;55(2):187-90.

Chuck FJ, Huffaker BJ. Labor induction with intravaginal misoprostol versus intracervical prostaglandin E2 gel (Prepidil gel): randomized comparison. Am J Obstet Gynecol 1995;173(4):1137-42.

Cibils LA. On prolonged pregnancy. Am J Obstet Gynecol 1995;172(4 Pt 1):1321-2.

Cibils LA, Votta R. Clinical significance of fetal heart rate patterns during labor. IX: Prolonged pregnancy. J Perinat Med 1993;21(2):107-16.

Clark A, Cook V, Hill P, et al. Cervical ripening and labor induction: misoprostol vs dinoprostone. Am J Obstet Gynecol 1998;178:S30.

Clark MJ. Use of the oxytocin challenge test in the management of postdate pregnancy. J Am Osteopath Assoc 1980;79(10):632-5.

Clark SL. Intrapartum management of the postdate patient. [Review]. Clin Obstet Gynecol 1989;32(2):278-84.

Clarkson CP, Magann EF, Siddique SA, et al. Hematological complications of Gaucher's disease in pregnancy. Mil Med 1998;163(7):499-501.

Clausson B, Cnattingius S, Axelsson O. Outcomes of post-term births: the role of fetal growth restriction and malformations. Obstet Gynecol 1999;94(5 Pt 1):758-62.

Clement D, Schifrin BS, Kates RB. Acute oligohydramnios in postdate pregnancy. Am J Obstet Gynecol 1987;157(4 Pt 1):884-6.

Clifford SH. Postmaturity--with placental dysfunction. J Pediatr 1954;44(1):1-13.

Cnattingius S, Taube A. Stillbirths and rate of neonatal deaths in 76,761 postterm pregnancies in Sweden, 1982-1991; a register study. Acta Obstet Gynecol Scand 1998;77(5):582-3.

Collins E. Maternal and fetal effects of acetaminophen and salicylates in pregnancy. [Review]. Obstet Gynecol 1981;58(5 Suppl):57S-62S.

Combs CA, Singh NB, Khoury JC. Elective induction versus spontaneous labor after sonographic diagnosis of fetal macrosomia. Obstet Gynecol 1993;81(4):492-6.

Cosner KR, Dougherty MC, Bishop KR. Dynamic characteristics of the circumvaginal muscles during pregnancy and the postpartum. J Nurse Midwifery 1991;36(4):221-5.

Cotzias CS, Paterson-Brown S, Fisk NM. Prospective risk of unexplained stillbirth in singleton pregnancies at term: population based analysis. BMJ 1999;319(7205):287-8.

Cousins ME. The reproductive risk potential of prolonged gestation. [Review]. Issues Health Care Women 1981;3(3):139-50.

Crane J, Bennett K, Young D, et al. The effectiveness of sweeping membranes at term: a randomized trial. Obstet Gynecol 1997;89(4):586-90.

Crowley P. Interventions for preventing or improving the outcome of delivery at or beyond term (Cochrane Review). In: The Cochrane Library, Issue 2, 2000. Oxford: Update Software.

Crowley P, O'Herlihy C, Boylan P. The value of ultrasound measurement of amniotic fluid volume in the management of prolonged pregnancies. Br J Obstet Gynaecol 1984;91(5):444-8.

Crowther CA. Prevention of preterm birth in multiple pregnancy. Bailliere's Clin Obstet Gynaecol 1998;12(1):67-75.

Cruikshank DP. Amniocentesis for determination of fetal maturity. [Review]. Clin Obstet Gynecol 1982;25(4):773-85.

Cruikshank DP, Linyear AS. Term stillbirth: causes and potential for prevention in Virginia. Obstet Gynecol 1987;69(6):841-4.

Crump WJ. Postdate pregnancy in a network of community hospitals: management and outcome. J Fam Pract 1988;26(1):41-4.

Crump WJ, Smith CW. The postdate pregnancy. When to wait, when to induce labor. Postgrad Med 1986;80(5):291-3, 296-7.

Csapo AI, Eskola J, Ruttner Z. The biological meaning of progesterone levels. Prostaglandins 1980;19(2):203-11.

Cucco C, Osborne MA, Cibils LA. Maternal-fetal outcomes in prolonged pregnancy. Am J Obstet Gynecol 1989;161(4):916-20.

Cundy T, Gamble G, Townend K, et al. Perinatal mortality in Type 2 diabetes mellitus. Diabet Med 2000;17(1):33-9.

Curtis PD, Matthews TG, Clarke TA, et al. Neonatal seizures: the Dublin Collaborative Study. Arch Dis Child 1988;63(9):1065-8.

Davis DL, Stewart M, Harmon RJ. Postponing pregnancy after perinatal death: perspectives on doctor advice. J Am Acad Child Adolesc Psychiatry 1989;28(4):481-7.

Davis L. The use of castor oil to stimulate labor in patients with premature rupture of membranes. J Nurse Midwifery 1984;29(6):366-70.

Deayton JM, Young IR, Hollingworth SA, et al. Effect of late hypothalamo-pituitary disconnection on the development of the HPA axis in the ovine fetus and the initiation of parturition. J Neuroendocrinol 1994;6(1):25-31.

Delpapa EH, Mueller-Heubach E. Pregnancy outcome following ultrasound diagnosis of macrosomia. Obstet Gynecol 1991;78(3 Pt 1):340-3.

Dennis KJ. Prolonged pregnancy: the management debate. BMJ 1986;293(6559):1434-5.

Devoe L, Golde S, Kilman Y, et al. A comparison of visual analyses of intrapartum fetal heart rate tracings according to the new national institute of child health and human development guidelines with computer analyses by an automated fetal heart rate monitoring system. Am J Obstet Gynecol 2000;183(2):361-6.

Devoe LD, Sholl JS. Postdates pregnancy. Assessment of fetal risk and obstetric management. J Reprod Med 1983;28(9):576-80.

Divon MY, Haglund B, Nisell H, et al. Fetal and neonatal mortality in the postterm pregnancy: the impact of gestational age and fetal growth restriction. Am J Obstet Gynecol 1998;178(4):726-31.

Dizon-Townson D, Ward K. The genetics of labor. [Review]. Clin Obstet Gynecol 1997;40(3):479-84.

Doany W, McCarty J. Outpatient management of the uncomplicated postdate pregnancy with intravaginal prostaglandin E2 gel and membrane stripping. J Matern Fetal Med 1997;6(2):71-8.

Donker DK, van Geijn HP, Hasman A. Interobserver

variation in the assessment of fetal heart rate recordings. Eur J Obstet Gynecol Reprod Biol 1993;52(1):21-8.

Dove D, Johnson P. Oral evening primrose oil: its effect on length of pregnancy and selected intrapartum outcomes in low-risk nulliparous women. J Nurse Midwifery 1999;44(3):320-4.

Druzin ML, Karver ML, Wagner W, et al. Prospective evaluation of the contraction stress and nonstress tests in the management of post-term pregnancy. Surg Gynecol Obstet 1992;174(6):507-12.

Duff C, Sinclair M. Exploring the risks associated with induction of labour: a retrospective study using the NIMATS database. Northern Ireland Maternity System. J Adv Nurs 2000;31(2):410-7.

Dunn PA, Rogers D, Halford K. Transcutaneous electrical nerve stimulation at acupuncture points in the induction of uterine contractions. Obstet Gynecol 1989;73(2):286-90.

Dyson DC. Fetal surveillance vs. labor induction at 42 weeks in postterm gestation. J Reprod Med 1988;33(3):262-70.

Dyson DC, Miller PD, Armstrong MA. Management of prolonged pregnancy: induction of labor versus antepartum fetal testing. Am J Obstet Gynecol 1987;156(4):928-34.

Ecker JL, Greenberg JA, Norwitz ER, et al. Birth weight as a predictor of brachial plexus injury. Obstet Gynecol 1997;89(5 Pt 1):643-7.

Eden RD. Postdate pregnancy: antenatal assessment of fetal well-being. [Review]. Clin Obstet Gynecol 1989;32(2):235-44.

Eden RD, Gergely RZ, Schifrin BS, et al. Comparison of antepartum testing schemes for the management of the postdate pregnancy. Am J Obstet Gynecol 1982;144(6):683-92.

Eden RD, Seifert LS, Kodack LD, et al. A modified biophysical profile for antenatal fetal surveillance. Obstet Gynecol 1988;71(3 Pt 1):365-9.

Eden RD, Seifert LS, Winegar A, et al. Perinatal characteristics of uncomplicated postdate pregnancies. Obstet Gynecol 1987;69(3 Pt 1):296-9.

Eden RD, Seifert LS, Winegar A, et al. Postdate pregnancies: a review of 46 perinatal deaths. Am J Perinatol 1987;4(4):284-7.

Edozien LC. What do maternity statistics tell us about induction of labour? J Obstet Gynaecol 1999;19(4):343-44.

Eganhouse DJ. Monitoring the postdate pregnancy, part I: antepartum issues. Mother Baby J 2000;5(1):7-10, 26-9.

Eganhouse DJ. Monitoring the postdate pregnancy, part II: intrapartum issues. Mother Baby J 2000;5(2):5-20.41-4.

Egarter C, Kofler E, Fitz R, et al. Is induction of labor indicated in prolonged pregnancy? Results of a prospective randomised trial. Gynecol Obstet Invest 1989;27(1):6-9.

Egarter CH, Husslein PW, Rayburn WF. Uterine hyperstimulation after low-dose prostaglandin E2 therapy: tocolytic treatment in 181 cases. Am J Obstet Gynecol 1990;163(3):794-6.

Ekman G, Perssen PH, Ulmsten U, et al. The impact on labor induction of intracervically applied PGEinf 2-gel, related to gestational age in patients with an unripe cervix. Acta Obstet Gynecol Scand 1983;62(Suppl. 113):173-5.

Ekman G, Persson PH, Ulmsten U. Induction of labor in postterm pregnant women. Int J Gynaecol Obstet 1986;24(1):47-52.

El-Azeem S, Samuels P, Welch G, et al. Term labor induction with PGE1 isoprostol versus PGE2 dinoprostone. Am J Obstet Gynecol 1997;176:S113.

el-Damarawy H, el-Sibaie F, Tawfik TA. Antepartum fetal surveillance in post-date pregnancy. Int J Gynaecol Obstet 1993;43(2):145-50.

El-Sherbiny MT, El-Gharieb IH, Gewely HA. Vaginal misoprostol for induction of labor: 25 vs. 50 µg dose regimen. Int J Gynecol Obstet 2001;72:25-30.

El-Torkey M, Grant JM. Sweeping of the membranes is an effective method of induction of labour in prolonged pregnancy: a report of a randomized trial. Br J Obstet Gynaecol 1992;99(6):455-8.

Elliott CL, Brennand JE, Calder AA. The effects of

mifepristone on cervical ripening and labor induction in primigravidae. Obstet Gynecol 1998;92(5):804-9.

Elliott JP, Flaherty JF. The use of breast stimulation to prevent postdate pregnancy. Am J Obstet Gynecol 1984;149(6):628-32.

Elpek G, Uner M, Trak B, et al. Intravaginal naproxen in preterm labour: A preliminary study. J Obstet Gynaecol 1998;18(3):236-7.

Engstrom JL, Sittler CP, Swift KE. Fundal height measurement. Part 5—The effect of clinician bias on fundal height measurements. J Nurse Midwifery 1994;39(3):130-41.

Escudero F, Contreras H. A comparative trial of labor induction with misoprostol versus oxytocin. Int J Gynaecol Obstet 1997;57(2):139-43.

Essel JK, Opai-Tetteh ET. Macrosomia - Maternal and fetal risk factors. S Afr Med J 1995;85(1):43-6.

Evans MI, Dougan MB, Moawad AH, et al. Ripening of the human cervix with porcine ovarian relaxin. Am J Obstet Gynecol 1983;147(4):410-4.

Ewigman BG, Crane JP, Frigoletto FD, et al. Effect of prenatal ultrasound screening on perinatal outcome. N Engl J Med 1993;329(12):821-7.

Fabre E, Gonzalez de Aguero R, de Agustin JL, et al. Perinatal mortality in term and post-term births. J Perinat Med 1996;24(2):163-9.

Facchinetti F, Neri I, Genazzani AR. Factors predicting labour onset in patients treated with prostaglandin Einf 2 for cervical ripening. Eur J Obstet Gynecol Reprod Biol 1995;60(2):129-32.

Fairlie FM, Lang GD, Greer IA, et al. Umbilical artery Doppler flow velocity waveforms and maternal prostaglandin E2 and F2 alpha metabolite concentrations during cervical ripening with prostaglandin E2. Eur J Obstet Gynecol Reprod Biol 1990;37(1):7-13.

Fait G, Grisaru D, Shenhav M, et al. Balloon catheter with extra-amniotic saline instillation: a method of induction in pregnancies at 41 or more gestational weeks. Aust N Z J Obstet Gynaecol 1997;37(2):174-6.

Farah LA, Sanchez-Ramos L, Rosa C, et al. Randomized trial of two doses of the prostaglandin E1 analog misoprostol for labor induction. Am J Obstet Gynecol 1997;177(2):364-9.

Farmakides G, Schulman H, Ducey J, et al. Uterine and umbilical artery Doppler velocimetry in postterm pregnancy. J Reprod Med 1988;33(3):259-61.

Farmakides G, Schulman H, Winter D, et al. Prenatal surveillance using nonstress testing and Doppler velocimetry. Obstet Gynecol 1988;71(2):184-7.

Farooqi A, Holmgren PA, Engberg S, et al. Survival and 2-year outcome with expectant management of second- trimester rupture of membranes. Obstet Gynecol 1998;92(6):895-901.

Feldman GB. Prospective risk of stillbirth. Obstet Gynecol 1992;79(4):547-53.

Flamm BL. Vaginal birth after cesarean section: controversies old and new. Clin Obstet Gynecol 1985;28(4):735-44.

Flamm BL. Was the death of this baby avoidable? Birth 1990;17(4):226-7.

Fleischer A, Schulman H, Farmakides G, et al. Antepartum nonstress test and the postmature pregnancy. Obstet Gynecol 1985;66(1):80-3.

Fletcher H, Mitchell S, Frederick J, et al. Intravaginal misoprostol versus dinoprostone as cervical ripening and labor-inducing agents. Obstet Gynecol 1994;83(2):244-7.

Fletcher HM, Mitchell S, Simeon D, et al. Intravaginal misoprostol as a cervical ripening agent. Br J Obstet Gynaecol 1993;100(7):641-4.

Flynn AM, Kelly J, Mansfield H, et al. A randomized controlled trial of non-stress antepartum cardiotocography. Br J Obstet Gynaecol 1982;89(6):427-33.

Forouzan I, Cohen AW. Can umbilical and arcuate artery Doppler velocimetry predict fetal distress among prolonged pregnancies? J Ultrasound Med 1991;10(1):15-7.

Francome C, Savage W. Caesarean section in Britain and the United States 12% or 24%: is either the right rate? Soc Sci Med 1993;37(10):1199-218.

Franz WB 3d. Fetal assessment. Prim Care 1983;10(2):173-203.

Freeman RK, Garite TJ, Mondanlou H, et al. Postdate pregnancy: utilization of contraction stress testing for primary fetal surveillance. Am J Obstet Gynecol 1981;140(2):128-35.

Friedrichs PE. An active management approach to the postdate fetus with a reactive nonstress test and fetal heart rate decelerations. Obstet Gynecol 1988;71(4):657-8.

Friesen CD, Miller AM, Rayburn WF. Influence of spontaneous or induced labor on delivering the macrosomic fetus. Am J Perinatol 1995;12(1):63-6.

Froen JF, Arnestad M, Frey K, et al. Risk factors for sudden intrauterine unexplained death: epidemiologic characteristics of singleton cases in Oslo, Norway, 1986-1995. American Journal of Obstetrics & Gynecology 2001;184(4):694-702.

Frydman R, Lelaidier C, Baton-Saint-Mleux C, et al. Labor induction in women at term with mifepristone (RU 486): a double-blind, randomized, placebo-controlled study. Obstet Gynecol 1992;80(6):972-5.

Gabbe SG. Effects of identifying a high risk population. Diabetes Care 1980;3(3):486-8.

Gaffney G, Flavell V, Johnson A, et al. Cerebral palsy and neonatal encephalopathy. Arch Dis Child Fetal & Neonatal Edition 1994;70(3):F195-200.

Gardosi J, Vanner T, Francis A. Gestational age and induction of labour for prolonged pregnancy. Br J Obstet Gynaecol 1997;104(7):792-7.

Garry D, Figueroa R, Guillaume J, et al. Use of castor oil in pregnancies at term. Altern Ther Health Med 2000;6(1):77-9.

Gauthier RJ, Griego BD, Goebelsmann U. Estriol in pregnancy. VII. Unconjugated plasma estriol in prolonged gestation. Am J Obstet Gynecol 1981;139(4):382-9.

Gebhart F. Induced labor, along with use of labor drugs, on the rise. Drug Topics 2000;144(12):54.

Gegor CL, Paine LL, Johnson TR. Antepartum fetal assessment. A nurse-midwifery perspective. [Review]. J Nurse Midwifery 1991;36(3):153-67.

Gerhard I, Postneek F. Auricular acupuncture in the treatment of female infertility. Gynecol Endocrinol 1992;6(3):171-81.

Ghidini A, Doria V, Kirn V, et al. Successful outcome after antibiotic treatment of postamniocentesis membrane rupture and chorioamnionitis in multiple pregnancy. Am J Perinatol 1999;16(8):403-6.

Giacalone P-L, Daures J-P, Faure J-M, et al. The effects of mifepristone on uterine sensitivity to oxytocin and on fetal heart rate patterns. Eur J Obstet Gynecol Reprod Biol 2001;97:30-4.

Giacalone PL, Targosz V, Laffargue F, et al. Cervical ripening with mifepristone before labor induction: A randomized study. Obstet Gynecol 1998;92(4 I):487-92.

Gibb DM, Cardozo LD, Studd JW, et al. Prolonged pregnancy: is induction of labour indicated? A prospective study. Br J Obstet Gynaecol 1982;89(4):292-5.

Gibbs RS, Castillo MS, Rodgers PJ. Management of acute chorioamnionitis. Am J Obstet Gynecol 1980;136(6):709-13.

Gibbs RS, Eschenbach DA. Use of antibiotics to prevent preterm birth. Am J Obstet Gynecol 1997;177(2):375-80.

Gichangi PB, Ndinya-Achola JO, Ombete J, et al. Antimicrobial prophylaxis in pregnancy: A randomized, placebo-controlled trial with cefetamet-pivoxil in pregnant women with a poor obstetric history. Am J Obstet Gynecol 1997;177(3):680-4.

Gilbert L, Porter W, Brown VA. Postpartum haemorrhage—a continuing problem. Br J Obstet Gynaecol 1987;94(1):67-71.

Gilbert WM, Nesbitt TS, Danielsen B. Associated factors in 1611 cases of brachial plexus injury. Obstet Gynecol 1999;93(4):536-40.

Gilby JR, Williams MC, Spellacy WN. Fetal abdominal circumference measurements of 35 and 38 cm as predictors of macrosomia. J Reprod Med 2000;45:936-8.

Giles WB. Vascular Doppler techniques. Obstet Gynecol Clin North Am 1999;26(4):595-606.

Gilson GJ, O'Brien ME, Vera RW, et al. Prolonged pregnancy and the biophysical profile. A birthing center perspective. J Nurse Midwifery 1988;33(4):171-7.

Gimovsky ML, Bruce SL. Aspects of FHR tracings as warning signals. Clin Obstet Gynecol 1986;29(1):51-63.

Gjessing HK, Skjaerven R, Wilcox AJ. Errors in gestational age: evidence of bleeding early in pregnancy. Am J Public Health 1999;89(2):213-8.

Goer H. Postterm pregnancy: putting the merits of a policy of induction of labor into perspective. Birth 1996;23(3):180-1.

Goeree R, Hannah M, Hewson S. Cost-effectiveness of induction of labour versus serial antenatal monitoring in the Canadian Multicentre Postterm Pregnancy Trial. CMAJ 1995;152(9):1445-50.

Gohar J, Mazor M, Leiberman JR. GnRH in pregnancy. Arch Gynecol Obstet 1996;259(1):1-6.

Goharkhay N, Stanczyk FZ, Gentzschein E, et al. Plasma prostaglandin E2 metabolite levels during labor induction with a sustained-release prostaglandin E2 vaginal insert. J Soc Gynecol Investig 2000;7:338-42.

Goldberg J, Newman RB, Rust PF. Interobserver reliability of digital and endovaginal ultrasonographic cervical length measurements. Am J Obstet Gynecol 1997;177(4):853-8.

Gonen O, Rosen DJ, Dolfin Z, et al. Induction of labor versus expectant management in macrosomia: a randomized study. Obstet Gynecol 1997;89(6):913-7.

Gonen R, Spiegel D, Abend M. Is macrosomia predictable, and are shoulder dystocia and birth trauma preventable? Obstet Gynecol 1996;88(4 Pt 1):526-9.

Goni S, Sawhney H, Gopalan S. Oxytocin induction of labor: a comparison of 20- and 60-min dose increment levels. Int J Gynaecol Obstet 1995;48(1):31-6.

Goodlin RC. Prolongation of pregnancy and survival of second twin. Am J Obstet Gynecol 1990;163(1 Pt 1):270-1.

Gottschall D, Borgida AF, Feldman DM, et al. Preinduction cervical ripening comparing 50 and 100 mcg of misoprostol. Am J Obstet Gynecol 1998;178:S93.

Gottschall DS, Borgida AF, Mihalek JJ, et al. A randomized clinical trial comparing misoprostol with prostaglandin E2 gel for preinduction cervical ripening. Am J Obstet Gynecol 1997;177(5):1067-70.

Granados JL. Survey of the management of postterm pregnancy. Obstet Gynecol 1984;63(5):651-3.

Grant A, Hepburn M. Merits of an individualized approach to fetal movement counting compared with fixed-time and fixed-number methods. Br J Obstet Gynaecol 1984;91(11):1087-90.

Grant JM. The fetal heart rate trace is normal, isn't it? Observer agreement of categorical assessments. Lancet 1991;337(8735):215-8.

Grant JM. Induction of labour confers benefits in prolonged pregnancy. [Review]. Br J Obstet Gynaecol 1994;101(2):99-102.

Grant JM. Induction of labour confers benefits in prolonged pregnancy. Br J Obstet Gynaecol 1995;102:80-1.

Graves GR, Baskett TF, Gray JH, et al. The effect of vaginal administration of various doses of prostaglandin E2 gel on cervical ripening and induction of labor. Am J Obstet Gynecol 1985;151(2):178-81.

Gregory KD, Henry OA, Ramicone E, et al. Maternal and infant complications in high and normal weight infants by method of delivery. Obstet Gynecol 1998;92(4 Pt 1):507-13.

Griffiths M. Stillbirths and rate of neonatal deaths in 76,761 postterm pregnancies in Sweden, 1982-1991; a register study. Acta Obstet Gynecol Scand 1998;77(5):583-4.

Grubb DK, Rabello YA, Paul RH. Post-term pregnancy: fetal death rate with antepartum surveillance. [Review]. Obstet Gynecol 1992;79(6):1024-6.

Grünberger W, Spona J. The effect of pericervical PGEinf 2 instillation on levels of maternal serum 13,14-dihydro-15-keto-PGF(2alpha) and progesterone. Arch Gynecol 1986;239(2):93-9.

Guidetti DA, Divon MY, Cavalieri RL, et al. Fetal umbilical artery flow velocimetry in postdate pregnancies. Am J Obstet Gynecol 1987;157(6):1521-3.

Guidetti DA, Divon MY, Langer O. Postdate fetal surveillance: is 41 weeks too early? Am J Obstet Gynecol 1989;161(1):91-3.

Gunderson E. Post-term pregnancy: fetal death rate with antepartum surveillance. Obstet Gynecol 1992;80(4):729-30.

Gupta R, Vasishta K, Sawhney H, et al. Safety and efficacy of stripping of membranes at term. Int J Gynaecol Obstet 1998;60(2):115-21.

Gyetvai K, Hannah ME, Hodnett ED, et al. Tocolytics for preterm labor: A systematic review. Obstet Gynecol 1999;94(5 SUPPL. 1):869-77.

Hagglund L, Christensen KK, Christensen P, et al. Risk factors in cesarean section infection. Obstet Gynecol 1983;62(2):145-50.

Halta VE. Calendars, clocks and choices: prolonged pregnancy [discussion appears in Midwifery Today Childbirth Educ 1996;38:38]. Midwifery Today Childbirth Educ 1996;38:36-7.

Halta VE. Calendars, clocks and choices. Midwifery Today Childbirth Educ 1996;(38):36-8.

Hanley ML, Vintzileos AM. Biophysical testing in premature rupture of the membranes. Semin Perinatol 1996;20(5):418-25.

Hann L, McArdle C, Sachs B. Sonographic biophysical profile in the postdate pregnancy. J Ultrasound Med 1987;6(4):191-5.

Hannah M. Management of post-term pregnancy. In: The Society of Obstetricians and Gynaecologists of Canada: Policy Statement, Committee Opinion, Clinical Practice Guidelines. Ottawa, ON : SOGC, 1996. 10 P.

Hannah M. Post-term pregnancy. In: The Society of Obstetricians and Gynaecologists of Canada: Policy Statement, Committee Opinion, Clinical Practice Guidelines. Ottawa, ON : SOGC, 1997. 7 P.

Hannah ME. Which tests of postterm fetal well-being work? Birth 1990;17(4):228-9.

Hannah ME, Hannah WJ, Caritis SN, et al. Comment on the effectiveness of induction of labor for postterm pregnancy (3). Am J Obstet Gynecol 1995;172(1):240-1.

Hannah ME, Hannah WJ, Hellmann J. Efficacy of induced labor vs serial antenatal monitoring in postterm pregnancy. J Am Osteopath Assoc 1992;92(8):982, 989.

Hannah ME, Hannah WJ, Hellmann J, et al. Induction of labor as compared with serial antenatal monitoring in post-term pregnancy. A randomized controlled trial. The Canadian Multicenter Post-term Pregnancy Trial Group [published erratum appears in N Engl J Med 1992 Jul 30;327(5):368]. N Engl J Med 1992;326(24):1587-92.

Hannah ME, Hannah WJ, Willan A. Comment on the effectiveness of induction of labor for postterm pregnancy. Am J Obstet Gynecol 1995;172(1 Pt 1):240-1.

Hannah ME, Huh C, Hewson SA, et al. Postterm pregnancy: putting the merits of a policy of induction of labor into perspective. Birth 1996;23(1):13-9.

Haque MN. Prolonged pregnancy: Is induction of labour still necessary? J Obstet Gynaecol 1989;9(3):196-8.

Harding K, Evans S, Newnham J. Screening for the small fetus: a study of the relative efficacies of ultrasound biometry and symphysiofundal height. Aust N Z J Obstet Gynaecol 1995;35(2):160-4.

Harman JH, Kim A. Current trends in cervical ripening and labor induction [published erratum appears in Am Fam Physician 1999 Nov 15;60(8):2238]. [Review]. Am Fam Physician 1999;60(2):477-84.

Harris B, Eden RD, Garite TJ, et al. Prolonged pregnancy. Part I: Identifying the patient at risk. Female Patient Pract Obstet Gynecol Med 1991;16(2):43-6, 48, 50.

Harris B, Eden RD, Garite TJ, et al. Prolonged Pregnancy: Part II: Monitoring and intervention. Female Patient Total Health Care Women 1991;16(3):47-9, 53, 56-57, 60.

Harris BA, Huddleston JF, Sutliff G, et al. The unfavorable cervix in prolonged pregnancy. Obstet Gynecol 1983;62(2):171-4.

Harrison JM. Clinical. The initiation of labour: physiological mechanisms. Br J Midwifery 2000;8(5):281-6.

Hart G. Far from settled... the issue of relative risks of "postdates pregnancy" is far from settled. Midwifery Today Childbirth Educ 1996;38:38.

Hauth JC, Goodman MT, Gilstrap LC 3d, et al. Post-term pregnancy. I. Obstet Gynecol 1980;56(4):467-70.

Hauth JC, Hankins GD, Gilstrap LC 3d, et al. Uterine contraction pressures with oxytocin induction/augmentation. Obstet Gynecol 1986;68(3):305-9.

Hedén L, Ingemarsson I, Ahlström H, et al. Induction of labor versus conservative management in prolonged pregnancy: Controlled study. Int J Feto-Matern Med 1991;4(4):231-6.

Hendricksen A. Prolonged pregnancy. A literature review. [Review]. J Nurse Midwifery 1985;30(1):33-42.

Henriksen TB, Wilcox AJ, Hedegaard M, et al. Bias in studies of preterm and postterm delivery due to ultrasound assessment of gestational age. Epidemiology 1995;6(5):533-7.

Herabutya Y, Prasertsawat P, Pokpirom J. A comparison of intravaginal misoprostol and intracervical prostaglandin E2 gel for ripening of unfavorable cervix and labor induction. J Obstet Gynaecol Res 1997;23(4):369-74.

Herabutya Y, Prasertsawat PO, Tongyai T, et al. Prolonged pregnancy: the management dilemma. Int J Gynaecol Obstet 1992;37(4):253-8.

Herbst A, Wolner-Hanssen P, Ingemarsson I. Risk factors for acidemia at birth. Obstet Gynecol 1997;90(1):125-30.

Hilder L, Costeloe K, Thilaganathan B. Prolonged pregnancy: evaluating gestation-specific risks of fetal and infant mortality. Br J Obstet Gynaecol 1998;105(2):169-73.

Hill WC, Fleming AD, Martin RW, et al. Home uterine activity monitoring is associated with a reduction in preterm birth. Obstet Gynecol 1990;76(1 SUPPL.):13S-8S.

Hobart JM. Post-term pregnancy. [Review]. Clin Perinatol 1989;16(4):909-15.

Hofmeyr GJ, Alfirevic Z, Kelly T, et al. Methods for cervical ripening and labour induction in late pregnancy (Cochrane Protocol). In: The Cochrane Library, Issue 3, 2000. Oxford: Update Software.

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and labour induction in late pregnancy (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Holcomb WL, Smeltzer JS. Cervical effacement: variation in belief among clinicians. Obstet Gynecol 1991;78(1):43-5.

Holst N, Jenssen TG, Burhol PG, et al. Gastrointestinal regulatory peptides during oxytocin infusion in post-term pregnancies. Acta Physiol Scand 1988;132(1):23-7.

Hourvitz A, Alcalay M, Korach J, et al. A prospective study of high- versus low-dose oxytocin for induction of labor. Acta Obstet Gynecol Scand 1996;75(7):636-41.

Howarth GR, Funk M, Steytler P, et al. A randomised controlled trial comparing vaginally administered misoprostol to vaginal dinoprostone gel in labour induction. J Obstet Gynaecol 1996;16:474-8.

Hsieh TT, Hung TH, Chen KC, et al. Perinatal outcome of oligohydramnios without associated premature rupture of membranes and fetal anomalies. Gynecol Obstet Invest 1998;45(4):232-6.

Huang DY, Usher RH, Kramer MS, et al. Determinants of unexplained antepartum fetal deaths. Obstet Gynecol 2000;95(2):215-21.

Hughes EG, Kelly AJ, Kavanagh J. Dinoprostone vaginal insert for cervical ripening and labor induction: a meta-analysis. Obstet Gynecol 2001;97(5):847-55.

Hulet B, Platt LD. Sinusoidal heart rate pattern and face presentation in a fetus from a postterm pregnancy. A case report. J Reprod Med 1987;32(3):211-2.

Hunter IW. Advanced extra-uterine pregnancy—two cases and a review of the literature. Ir Med J 1983;76(2):82-3.

Hutchon DJ. Gestational age and induction of labour for prolonged pregnancy. Br J Obstet Gynaecol 1998;105(2):247-8.

Hutson JM, Petrie RH. Possible limitations of fetal monitoring. Clin Obstet Gynecol 1986;29(1):104-13.

Idrisa A, Obisesan KA, Adeleye JA. Fetal membrane sweeping for stimulation of labour in prolonged pregnancy: A controlled study. J Obstet Gynaecol 1993;13(4):235-7.

Iffy L, Apuzzio JJ, Mitra S, et al. Rates of cesarean section and perinatal outcome. Perinatal mortality. Acta Obstet Gynecol Scand 1994;73(3):225-30.

Imai M, Tani A, Saito M, et al. Significance of fetal fibronectin and cytokine measurement in the cervicovaginal secretions of women at term in predicting term labor and post-term pregnancy. Eur J Obstet Gynecol Reprod Biol 2001;97:53-8.

Ingardia CJ. Antepartum testing in postterm pregnancy. Am J Obstet Gynecol 1982;143(1):113-4.

Ingemarsson I, Kallen K. Stillbirths and rate of neonatal deaths in 76,761 postterm pregnancies in Sweden, 1982-1991: a register study. [Review]. Acta Obstet Gynecol Scand 1997;76(7):658-62.

Irion O, Boulvain M. Induction of labour for suspected fetal macrosomia (Cochrane Review). In: The Cochrane Library, Issue 4, 2000. Oxford: Update Software.

Ismail AA, Khowesah MM, Shaala SA, et al. Induction of labor by oral prostaglandin E2 in protracted pregnancy. Int J Gynaecol Obstet 1989;29(4):325-8.

Jacobsen G. Prediction of fetal growth deviations by use of symphysis-fundus height measurements. Internat J Tech Assess Health Care 1992;8(Suppl 1):152-9.

Jarvelin MR, Hartikainen-Sorri AL, Rantakallio P. Labour induction policy in hospitals of different levels of specialisation. Br J Obstet Gynaecol 1993;100(4):310-5.

Jazayeri A, Heffron JA, Phillips R, et al. Macrosomia prediction using ultrasound fetal abdominal circumference of 35 centimeters or more. Obstet Gynecol 1999;93(4):523-6.

Jazayeri A, Tsibris JCM, Spellacy WN. Elevated umbilical cord plasma erythropoietin levels in prolonged pregnancies. Obstet Gynecol 1998;92(1):61-3.

Jocums SB, Berg CJ, Entman SS, et al. Postdelivery mortality in Tennessee, 1989-1991. Obstet Gynecol 1998;91(5 I):766-70.

Johanson RB, Menon BKV. Vacuum extraction vs forcepts for assisted vaginal delivery (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Johnson JM, Harman CR, Lange IR, et al. Biophysical profile scoring in the management of the postterm pregnancy: an analysis of 307 patients. Am J Obstet Gynecol 1986;154(2):269-73.

Johnson JW, Longmate JA, Frentzen B. Excessive maternal weight and pregnancy outcome. Am J Obstet Gynecol 1992;167(2):353-70.

Johnson TR, Paine LL, Strobino DM, et al. Population differences affect the interpretation of fetal nonstress test results. Am J Obstet Gynecol 1998;179(3 Pt 1):779-83.

Johnstone MJ. Prolonged pregnancy: is induction of labour indicated? A prospective study. Br J Obstet Gynaecol 1982;89(9):778-9.

Johnstone MJ, Downie G, Gaskell A. Induction of labor for postmaturity. Lancet 1982;2(8296):496.

Jones DC, Hayslett JP. Outcome of pregnancy in women with moderate or severe renal insufficiency. N Engl J Med 1996;335(4):226-32.

Jongsma HW, Nijhuis JG. Critical analysis of the validity of electronic fetal monitoring. J Perinat Med 1991;19(1-2):33-7.

Jorge CS, Artal R, Paul RH, et al. Antepartum fetal surveillance in diabetic pregnant patients. Am J Obstet Gynecol 1981;141(6):641-5.

Jüni P, Altman DG, Egger M. Assessing the quality of controlled clinical trials. BMJ 2001;323:42-6.

Jüni P, Witschi A, Bloch R, et al. The hazards of scoring the quality of clinical trials for meta-analysis. JAMA 1999;282(11):1054-60.

Kadanali S, Küçüközkan T, Zor N, et al. Comparison of labor induction with misoprostol vs. oxytocin/prostaglandin E2 in term pregnancy. Int J Gynaecol Obstet 1996;55(2):99-104.

Kadar N, Tapp A, Wong A. The influence of nipple

stimulation at term on the duration of pregnancy. J Perinatol 1990;10(2):164-6.

Kanakura Y, Kometani K, Nagata T, et al. Uterine reconstruction for complete septate uterus. Keio J Med 1999;48(4):201-3.

Kaplan B, Goldman GA, Peled Y, et al. The outcome of post-term pregnancy. A comparative study. J Perinat Med 1995;23(3):183-9.

Karande VC, Deshmukh MA, Virkud A. Management of post-term pregnancy. J Postgrad Med 1985;31(2):98-101.

Kassis A, Mazor M, Leiberman JR, et al. Management of post-date pregnancy: a case control study. Isr J Med Sci 1991;27(2):82-6.

Katz VL, Farmer RM, Dean CA, et al. Use of misoprostol for cervical ripening. Southern Med J 2000;93(9):881-4.

Katz Z, Yemini M, Lancet M, et al. Non-aggressive management of post-date pregnancies. Eur J Obstet Gynecol Reprod Biol 1983;15(2):71-9.

Kaufmann RC, McBride P, Amankwah KS, et al. The effect of minor degrees of glucose intolerance on the incidence of neonatal macrosomia. Obstet Gynecol 1992;80(1):97-101.

Keirse MJ. Amniotomy or oxytocin for induction of labor. Re-analysis of a randomized controlled trial. Acta Obstet Gynecol Scand 1988;67(8):731-5.

Keirse MJ. Roundtable: postterm pregnancy. Part II. In the final analysis. Birth 1991;18(2):114-5.

Keirse MJ. Prostaglandins in preinduction cervical ripening. Meta-analysis of worldwide clinical experience. J Reprod Med 1993;38(1 Suppl):89-100.

Keirse MJN. Postterm pregnancy: new lessons from an unresolved debate. Birth 1993;20(2):102-5.

Keirse MJNC. New perspectives for the effective treatment of preterm labor. Am J Obstet Gynecol 1995;173(2):618-28.

Keith RD, Beckley S, Garibaldi JM, et al. A multicentre comparative study of 17 experts and an intelligent computer system for managing labour using the cardiotocogram. Br J Obstet Gynaecol 1995;102(9):688-700.

Kelly AJ, Kavanagh J, Thomas J. Castor oil, bath and/or enema for cervical priming and induction of labour (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Kelly AJ, Kavanagh J, Thomas J. Relaxin for cervical ripening and induction of labour (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Kelly AJ, Kavanagh J, Thomas J. Vaginal prostaglandin (PGE2 and PGF2a) for induction of labour at term (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Kemp B, Winkler M, Rath W. Induction of labor by prostaglandin E2 in relation to the Bishop score. Int J Gynecol Obstet 2000;71:13-7.

Keng MTY, Eng BTS. Oral contraceptives postdatism as a complication. Med J Malaysia 1982;37(4):338-43.

Kesmodel U, Olsen SF. Smoking habits among pregnant Danish women: reliability of information recorded after delivery. J Epidemiol Community Health 1999;53(4):239-42.

Khan KK, Dinnes J, Kleijnen J. Systematic reviews to evaluate diagnostic tests. Eur J Obstet Gynecol Reprod Biol 2001;95:6-11.

Khouzami VA, Johnson JW, Hernandez E, et al. Urinary estrogens in postterm pregnancy. Am J Obstet Gynecol 1981;141(2):205-11.

Kidd LC, Patel NB, Smith R. Non-stress antenatal cardiotocography—a prospective randomized clinical trial. Br J Obstet Gynaecol 1985;92(11):1156-9.

Kilpatrick SJ, Safford KL. Maternal hydration increases amniotic fluid index in women with normal amniotic fluid. Obstet Gynecol 1993;81(1):49-52.

Knox GE, Huddleston JF, Flowers CE. Management of prolonged pregnancy: results of a prospective randomized trial. Am J Obstet Gynecol 1979;134(4):376-84.

Kochenour NK. Estrogen assay during pregnancy. [Review]. Clin Obstet Gynecol 1982;25(4):659-72.

Kochenour NK. Other causes of fetal death. [Review]. Clin Obstet Gynecol 1987;30(2):312-21.

Kolderup LB, Laros RK, Musci TJ. Incidence of persistent birth injury in macrosomic infants: association with mode of delivery. Am J Obstet Gynecol 1997;177(1):37-41.

Konje JC, Obisesan KA, Ladipo OA. Obstructed labor in Ibadan. Int J Gynecol Obstet 1992;39(1):17-21.

Kramer M, et al. The validity of gestational age estimation by menstrual dating in term, preterm, and postterm estations. JAMA 1988;260(22):3306-8.

Kramer RL, Gilson GJ, Morrison DS, et al. A randomized trial of misoprostol and oxytocin for induction of labor: safety and efficacy. Obstet Gynecol 1997;89(3):387-91.

Krammer J, O'Brien WF. Mechanical methods of cervical ripening. [Review]. Clin Obstet Gynecol 1995;38(2):280-6.

Krammer J, Williams MC, Sawai SK, et al. Pre-induction cervical ripening: a randomized comparison of two methods. Obstet Gynecol 1995;85(4):614-8.

Kulkarni SK, Matadial L. Prolonged pregnancy—a rational approach to management. West Indian Med J 1986;35(4):314-7.

Kumari S, Jain S, Pruthi PK, et al. Perinatal risks in postdated pregnancy. Indian Pediatr 1984;21(1):21-7.

Lagrew DC, Freeman RK. Management of postdate pregnancy. Am J Obstet Gynecol 1986;154(1):8-13.

Lahteenmaki P, Rapeli T, Kaariainen M, et al. Late postcoital treatment against pregnancy with antiprogesterone RU 486. Fertil Steril 1988;50(1):36-8.

Landon MB, Gabbe SG. Antepartum fetal surveillance in gestational diabetes mellitus. Diabetes 1985;34(SUPPL. 2):50-4.

Lang J, Lieberman E. Prolonged pregnancy: The management debate. BMJ 1988;297(6650):715-7.

Langer O, Vega-Rich M, Cohen W. Terminal pattern: Characteristics and management. Am J Perinatol 1985;2(4):300-4.

Lazor LZ, Philipson EH, Ingardia CJ, et al. A randomized comparison of 15- and 40-minute dosing protocols for labor augmentation and induction. Obstet Gynecol 1993;82(6):1009-12.

Leaphart WL, Meyer MC, Capeless EL. Labor induction with a prenatal diagnosis of fetal macrosomia. J Matern Fetal Med 1997;6(2):99-102.

Ledger WJ. Identification of the high risk mother and fetus--does it work? [Review]. Clin Perinatol 1980;7(1):125-34.

Lee HY. A randomised double-blind study of vaginal misoprostol vs dinoprostone for cervical ripening and labour induction in prolonged pregnancy. Singapore Med J 1997;38(7):292-4.

Lee NC. Postmaturity and induction of labour. S Afr Med J 1986;70(11):647.

LeFevre ML, Bain RP, Ewigman BG, et al. A randomized trial of prenatal ultrasonographic screening: impact on maternal management and outcome. RADIUS (Routine Antenatal Diagnostic Imaging with Ultrasound) Study Group. Am J Obstet Gynecol 1993;169(3):483-9.

Leiman G, Harrison NA, Rubin A. Pregnancy following conization of the cervix: complications related to cone size. Am J Obstet Gynecol 1980;136(1):14-8.

Lemancewicz A, Urban R, Skotnicki MZ, et al. Uterine and fetal Doppler flow changes after misoprostol and oxytocin therapy for induction of labor in post-term pregnancies. Int J Gynaecol Obstet 1999;67(3):139-45.

Lemons JA, Vargas P, Delaney JJ. Infant of the diabetic mother: Review of 225 cases. Obstet Gynecol 1981;57(2):187-92.

Lennox CE, Patel NB. Early versus late induction of labour in post-term pregnancy. BMJ 1987;294(6588):1689.

Leppert PC. Anatomy and physiology of cervical ripening. Clin Obstet Gynecol 1995;38(2):267-79.

Leslie GI, Gallery EDM, Arnold JD, et al. Neonatal outcome in a randomized, controlled trial of low-dose aspirin in high-risk pregnancies. J Paediatr Child Health 1995;31(6):549-52.

Leveno KJ, Quirk JG, Cunningham FG, et al. Prolonged pregnancy. I. Observations concerning the

causes of fetal distress. Am J Obstet Gynecol 1984;150(5 Pt 1):465-73.

Levine AB, Lockwood CJ, Brown B, et al. Sonographic diagnosis of the large for gestational age fetus at term: does it make a difference? Obstet Gynecol 1992;79(1):55-8.

Lewis R, Mercer BM, Salama M, et al. Oral terbutaline after parenteral tocolysis: A randomized, double- blind, placebo-controlled trial. Am J Obstet Gynecol 1996;175(4 I):834-7.

Librizzi RJ. Antepartum fetal monitoring: nonstress testing in the evaluation of uteroplacental insufficiency. J Am Osteopath Assoc 1980;79(10):636-40.

Lidegaard O, Bottcher LM, Weber T. Description, evaluation and clinical decision making according to various fetal heart rate patterns. Inter-observer and regional variability. Acta Obstet Gynecol Scand 1992;71(1):48-53.

Lien JM, Morgan MA, Garite TJ, et al. Antepartum cervical ripening: applying prostaglandin E2 gel in conjunction with scheduled nonstress tests in postdate pregnancies. Am J Obstet Gynecol 1998;179(2):453-8.

Lim BH, Mahmood TA, Smith NC, et al. A prospective comparative study of transvaginal ultrasonography and digital examination for cervical assessment in the third trimester of pregnancy. J Clin Ultrasound 1992;20(9):599-603.

Lindhard A, Nielsen PV, Mouritsen LA, et al. The implications of introducing the symphyseal-fundal height-measurement. A prospective randomized controlled trial. Br J Obstet Gynaecol 1990;97(8):675-80.

Lipscomb KR, Gregory K, Shaw K. The outcome of macrosomic infants weighing at least 4500 grams: Los Angeles County + University of Southern California experience. Obstet Gynecol 1995;85(4):558-64.

Losh DP, Duhring JL. Management of the postdates pregnancy. Am Fam Physician 1987;36(2):184-94.

Luckas M, Bricker L. Intravenous prostaglandin for induction of labour (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Luckas M, Buckett W, Alfirevic Z. Comparison of outcomes in uncomplicated term and post-term pregnancy following spontaneous labor. J Perinat Med 1998;26(6):475-9.

Lumbiganon P, Laopaiboon M, Kuchaisit C, et al. Oral prostaglandins (excluding misoprostol) for cervical ripening and labour induction when the baby is alive (Cochrane Review). In: The Cochrane Library, Issue 3, 2000. Oxford: Update Software.

Lumley J, Lester A, Anderson I, et al. A randomized trial of weekly cardiotocography in high-risk obstetric patients. Br J Obstet Gynaecol 1983;90(11):1018-26.

Lurie S, Matzkel A, Weissman A, et al. Outcome of pregnancy in class A1 and A2 gestational diabetic patients delivered beyond 40 weeks' gestation. Am J Perinatol 1992;9(5-6):484-8.

Lurie S, Zalel Y, Hagay ZJ. The evaluation of accelerated fetal growth. [Review]. Curr Opin Obstet Gynecol 1995;7(6):477-81.

Lydon-Rochelle M, Holt VL, Easterling TR, et al. Risk of uterine rupture during labor among women with a prior cesarean delivery. N Engl J Med 2001;345(1):3-8.

Maccato M, McLean W, Riddle G, et al. Isolation of Kingella denitrificans from amniotic fluid in a woman with chorioamnionitis. A case report. J Reprod Med 1991;36(9):685-7.

Macer JA, Macer CL, Chan LS. Elective induction versus spontaneous labor: a retrospective study of complications and outcome. Am J Obstet Gynecol 1992;166(6 Pt 1):1690-6.

MacGillivray I. Twins and other multiple deliveries. [Review]. Clin Obstet Gynaecol 1980;7(3):581-600.

MacKenzie IZ, Burns E. Randomised trial of one versus two doses of prostaglandin E2 for induction of labour: 1. Clinical outcome. Br J Obstet Gynaecol 1997;104(9):1062-7.

Macnaughton MC, Chalmers IG, Dubowitz V, et al. Final report of the Medical Research Council/Royal College of Obstetricians and Gynaecologists Multicentre Randomised Trial of Cervical Cerclage. Br J Obstet Gynaecol 1993;100(6):516-23.

Magann EF, Chauhan SP, Nevils BG, et al.

Management of pregnancies beyond forty-one weeks' gestation with an unfavorable cervix. Am J Obstet Gynecol 1998;178(6):1279-87.

Magann EF, McNamara MF, Whitworth NS, et al. Can we decrease postdatism in women with an unfavorable cervix and a negative fetal fibronectin test result at term by serial membrane sweeping? Am J Obstet Gynecol 1998;179(4):890-4.

Mahmood TA. A prospective comparative study on the use of prostaglandin E2 gel (2 mg) and prostaglandin E2 tablet (3 mg) for the induction of labour in primigravid women with unfavorable cervices. Eur J Obstet Gynecol Reprod Biol 1989;33(2):169-75.

Mancuso S, Ferrazzani S, De Carolis S, et al. Term and postterm low-risk pregnancies: management schemes for the reduction of high rates of cesarean section. Minerva Ginecol 1996;48(3):95-8.

Mandelbrot L, Dommergues M, Dumez Y. Prepartum transabdominal amnio-infusion for severe oligohydramnios. Acta Obstet Gynecol Scand 1992;71(2):124-5.

Mandelbrot L, Verspyck E, Dommergues M, et al. Transabdominal amnioinfusion for the management of nonlaboring postdates with severe oligohydramnios. Fetal Diagn Ther 1993;8(6):412-7.

Mandouvalos H, Alkalai BK, Metallinos C, et al. Considerations on the etiology of postponed labour. Ther Hung 1982;30(2):83-7.

Mandruzzato G, Meir YJ, D'Ottavio G, et al. Computerised evaluation of fetal heart rate in post-term fetuses: long term variation. Br J Obstet Gynaecol 1998;105(3):356-9.

Manning FA. The use of sonography in the evaluation of the high-risk pregnancy. Radiol Clin North Am 1990;28(1):205-16.

Mannino F. Neonatal complications of postterm gestation. J Reprod Med Obstet Gynecol 1988;33(3):271-6.

Manolitsas T, Wein P, Beischer NA, et al. Value of cardiotocography in women with antepartum haemorrhage - Is it too late for Caesarean section when the cardiotocograph shows ominous features? Aust N Z J Obstet Gynaecol 1994;34(4):403-8.

Manor M, Blickstein I, Ben-Arie A, et al. Case series of labor induction in twin gestations with an intrauterine balloon catheter. Gynecol Obstet Invest 1999;47(4):244-6.

Martin JN, Perry KG, Roberts WE, et al. Plasma exchange for preeclampsia: II. Unsuccessful antepartum utilization for severe preeclampsia with or without HELLP syndrome. J Clin Apheresis 1994;9(3):155-61.

Martin JN, Sessums JK, Howard P, et al. Alternative approaches to the management of gravidas with prolonged-postterm-postdate pregnancies. J Miss State Med Assoc 1989;30(4):105-11.

Mathews TJ. Trends in stimulation and induction of labor 1989-1995. Stat Bull Metrop Insur Co 1997;78(4):20-6.

Matijevic R. Outcome of post-term pregnancy: a matched-pair case-control study. Croat Med J 1998;39(4):430-4.

Maymon R, Shulman A, Pomeranz M, et al. Uterine rupture at term pregnancy with the use of intracervical prostaglandin E2 gel for induction of labor. Am J Obstet Gynecol 1991;165(2):368-70.

McColgin SW, Hampton HL, McCaul JF, et al. Stripping membranes at term: can it safely reduce the incidence of post-term pregnancies? Obstet Gynecol 1990;76(4):678-80.

McColgin SW, Patrissi GA, Morrison JC. Stripping the fetal membranes at term. Is the procedure safe and efficacious? Journal of Reproductive Medicine 1990;35(8):811-4.

McLaren M, Greer IA, Smith JR, et al. Maternal plasma bicycling PGE2 levels following vaginal administration of prostaglandin E2 pessaries in full term pregnancies. Prog Clin Biol Res 1987;242:199-203.

McLean FH, Boyd ME, Usher RH, et al. Postterm infants: too big or too small? Am J Obstet Gynecol 1991;164(2):619-24.

McMahon MJ, Kuller JA, Yankowitz J. Assessment of the post-term pregnancy. Am Fam Physician 1996;54(2):631-6, 641-2.

McNally OM, Turner MJ. Induction of labour after 1 previous Caesarean section. Aust N Z J Obstet

Gynaecol 1999;39(4):425-9.

Megafu U, Ozumba BC. Obstetric complications of macrosomic babies in African women. Int J Gynaecol Obstet 1988;26(2):197-202.

Mehl-Madrona L, Madrona MM. Physician- and midwife-attended home births. Effects of breech, twin, and post-dates outcome data on mortality rates. J Nurse-Midwifery 1997;42(2):91-8.

Mercer BM, McNanley T, O'Brien JM, et al. Early versus late amniotomy for labor induction: a randomized trial. Am J Obstet Gynecol 1995;173(4):1321-5.

Merrill PA, Porto M, Lovett SM, et al. Evaluation of the nonreactive positive contraction stress test prior to 32 weeks: The role of the biophysical profile. Am J Perinatol 1995;12(4):229-31.

Milchev N, Pehlivanov B, Paskaleva V, et al. Prostaglandin E2 in preinduction cervical ripening in postdate pregnancy. Folia Med (Plovdiv) 1999;41(3):81-5.

Miller DD. Tocolytic therapy for the prevention of preterm labor. J Intraven Nurs 1995;18(5):233-8.

Miller FC, Read JA. Intrapartum assessment of the postdate fetus. Am J Obstet Gynecol 1981;141(5):516-20.

Minaretzis D, Tsionu C, Papageorgiou I, et al. Intracervical prostaglandin E2 gel for cervical ripening and labor induction: what is the appropriate dose? Gynecol Obstet Invest 1993;35(1):34-7.

Minchom P, Niswander K, Chalmers I, et al. Antecedents and outcome of very early neonatal seizures in infants born at or after term. Br J Obstet Gynaecol 1987;94(5):431-9.

Misra M, Vavre S. Labour induction with intracervical prostaglandin Einf 2 gel and intravenous oxytocin in women with a very unfavourable cervix. Aust N Z J Obstet Gynaecol 1994;34(5):511-5.

Mittendorf R, Williams MA, Berkey CS, et al. The length of uncomplicated human gestation. Obstet Gynecol 1990;75(6):929-32.

Miyakawa I, Ichimaru S, Tayama C, et al. Effect of ACTH infusion on induction of uterine contraction in prolonged human pregnancy in association with endocrine function. Int J Gynecol Obstet 1981;19(2):89-96.

Miyazaki FS, Miyazaki BA. False reactive nonstress tests in postterm pregnancies. Am J Obstet Gynecol 1981;140(3):269-76.

Mizutani S, Itakura A, Kurauchi O, et al. Maternal serum oxytocinase activities in patients with poor obstetric dates. Med Sci Res 1994;22(11):771-5.

Moher D, Jadad AR, Tugwell P. Assessing the quality of randomized controlled trials. Current issues and future directions. Internat J Tech Assess Health Care 1996;12(2):195-208.

Mohide PT. Randomised comparison of early versus late induction of labor in post-term pregnancy. BMJ 1987;295(6594):388-9.

Moldin PG, Sundell G. Induction of labour: a randomised clinical trial of amniotomy versus amniotomy with oxytocin infusion. Br J Obstet Gynaecol 1996;103(4):306-12.

Monaghan J, O'Herlihy C, Boylan P. Ultrasound placental grading and amniotic fluid quantitation in prolonged pregnancy. Obstet Gynecol 1987;70(3 Pt 1):349-52.

Mongan PF, Fadel HE. Postdatism. Am Fam Physician 1981;23(3):208-12.

Monincx WM, Zondervan HA, Birnie E, et al. High risk pregnancy monitored antenatally at home. Eur J Obstet Gynecol Reprod Biol 1997;75(2):147-53.

Montan S, Malcus P. Amniotic fluid index in prolonged pregnancy: A cohort study. J Matern Fetal Invest 1995;5(1):4-7.

Moran C, Carranza-Lira S, Ochoa R, et al. Gastrin levels in mothers and neonates at delivery in various perinatal conditions. Acta Obstet Gynecol Scand 1996;75(7):608-11.

Moran DJ, McGarrigle HH, Lachelin GC. Lack of normal increase in saliva estriol/progesterone ratio in women with labor induced at 42 weeks' gestation. Am J Obstet Gynecol 1992;167(6):1563-4.

Mouw RJ, Egberts J, Kragt H, et al. Cervicovaginal fetal fibronectin concentrations: predictive value of impending birth in postterm pregnancies. Eur J Obstet Gynecol Reprod Biol 1998;80(1):67-70.

Mukherji J, Samaddar JC. How safe is caesarean section. J Obstet Gynaecol 1995;21(1):17-21.

Muller PR, Stubbs TM, Laurent SL. A prospective randomized clinical trial comparing two oxytocin induction protocols. Am J Obstet Gynecol 1992;167(2):373-80.

Mundle WR, Young DC. Vaginal misoprostol for induction of labor: a randomized controlled trial. Obstet Gynecol 1996;88(4 Pt 1):521-5.

Murphy PA, Fullerton J. Outcomes of intended home births in nurse-midwifery practice: a prospective descriptive study. [Review]. Obstet Gynecol 1998;92(3):461-70.

Nageotte MP, Freeman RK, Freeman AG, et al. Short-term variability assessment from abdominal electrocardiogram during the antepartum period. Am J Obstet Gynecol 1983;145(5):566-9.

Nagey DA. Post-term pregnancy. Obstet Gynecol 1981;58(1):135-6.

National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. A clinical trial of induction of labor versus expectant management in postterm pregnancy. The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. Am J Obstet Gynecol 1994;170(3):716-23.

Nationwide Inpatient Sample [NIS]. [electronic database]. Release 6; 1997 data. Agency for Healthcare Research and Quality: Rockville, MD. Available from: http://www.ntis.gov/fcpc/cpn8834.htm. 1997.

Neerhof MG, Cravello C, Haney EI, et al. Timing of labor induction after premature rupture of membranes between 32 and 36 weeks' gestation. Am J Obstet Gynecol 1999;180(2 I):349-52.

Neiger R. Fetal macrosomia in the diabetic patient. Clin Obstet Gynecol 1992;35(1):138-50.

Neilson JP. Symphysis-fundal height measurement in pregnancy (Cochrane Review). In: The Cochrane Library, Issue 3, 2000. Oxford: Update Software.

Neilson JP. Ultrasound for fetal assessment in early pregnancy (Cochrane Review). In: The Cochrane Library, Issue 4, 2000. Oxford: Update Software.

Neilson JP. Mifepristone for induction of labour (Cochrane Review). In: The Cochrane Library, Issue 3, 2001. Oxford: Update Software.

Neilson JP, Alfirevic Z. Doppler ultrasound for fetal assessment in high risk pregnancies (Cochrane Review). In: The Cochrane Library, Issue 3, 2000. Oxford: Update Software.

Neri A, Kaplan B, Rabinerson D, et al. The management of persistent occipito-posterior position. Clin Exp Obstet Gynecol 1995;22(2):126-31.

Newnham JP, Evans SF, Michael CA, et al. Effects of frequent ultrasound during pregnancy: a randomised controlled trial. Lancet 1993;342(8876):887-91.

Nichols CW. Postdate pregnancy: clinical implications... part 2. J Nurse Midwifery 1985;30(5):259-68.

Nichols CW. Postdate pregnancy. Part I. A literature review. [Review]. J Nurse Midwifery 1985;30(4):222-39.

Nocon JJ, McKenzie DK, Thomas LJ, et al. Shoulder dystocia: an analysis of risks and obstetric maneuvers. Am J Obstet Gynecol 1993;168(6 Pt 1):1732-7.

Norman K, Pattinson RC, De SJ, et al. Ampicillin and metronidazole treatment in preterm labour: A multicentre, randomised controlled trial. Br J Obstet Gynaecol 1994;101(5):404-8.

Novakov A, Segedi D, Milasinovic L, et al. Induction of labor by endocervical application of prostaglandins and intravenous infusion of oxytocin in postterm pregnancy. Med Pregl 1998;51(9-10):419-26.

Novy MJ, Walsh SW. Dexamethasone and estradiol treatment in pregnant rhesus macaques: Effects on gestational length, maternal plasma hormones, and fetal growth. Am J Obstet Gynecol 1983;145(8):920-31.

O'Brien JM, Mercer BM, Cleary NT, et al. Efficacy of outpatient induction with low-dose intravaginal prostaglandin E2: a randomized, double-blind, placebo-controlled trial. Am J Obstet Gynecol 1995;173(6):1855-9.

O'Connor R. Induction of labour at term. Evidence on

outcome favours induction. BMJ 1993;306(6889):1413-4.

O'Leary JA, Leonetti HB. Shoulder dystocia: prevention and treatment. Am J Obstet Gynecol 1990;162(1):5-9.

O'Reilly-Green CP, Divon MY. Predictive value of amniotic fluid index for oligohydramnios in patients with prolonged pregnancies. J Matern Fetal Med 1996;5(4):218-26.

O'Reilly-Green CP, Divon MY. Receiver operating characteristic curves of sonographic estimated fetal weight for prediction of macrosomia in prolonged pregnancies. Ultrasound Obstet Gynecol 1997;9(6):403-8.

Ocak V, Sen C, Demirkiran F, et al. FHR monitoring and perinatal mortality in high-risk pregnancies. Eur J Obstet Gynecol Reprod Biol 1992;44(1):59-63.

Ohel G, Rahav D, Rothbart H, et al. Randomised trial of outpatient induction of labor with vaginal PGE2 at 40-41 weeks of gestation versus expectant management. Arch Gynecol Obstet 1996;258(3):109-12.

Ohel G, Yaacobi N, Linder N, et al. Postdate antenatal testing. Int J Gynaecol Obstet 1995;49(2):145-7.

Ojo A, Oronsaye U. Who is the elderly primigravida in Nigeria? Int J Gynecol Obstet 1988;26(1):51-5.

Olofsson P, Saldeen P. The prospects for vaginal delivery in gestations beyond 43 weeks. Acta Obstet Gynecol Scand 1996;75(7):645-50.

Olofsson P, Saldeen P, Marsal K. Association between a low umbilical artery pulsatility index and fetal distress in labor in very prolonged pregnancies. Eur J Obstet Gynecol Reprod Biol 1997;73(1):23-9.

Olsen K. Taken to the limit. Pract Midwife 1999;2(6):46.

Orhue AA. A randomised trial of 45 minutes and 15 minutes incremental oxytocin infusion regimes for the induction of labour in women of high parity. Br J Obstet Gynaecol 1993;100(2):126-9.

Orhue AA. A randomized trial of 30-min and 15-min oxytocin infusion regimen for induction of labor at term in women of low parity. Int J Gynaecol Obstet 1993;40(3):219-25.

Orhue AA. Incremental increases in oxytocin infusion regimens for induction of labor at term in primigravidas: a randomized controlled trial. Obstet Gynecol 1994;83(2):229-33.

Otto C, Platt LD. Fetal growth and development. [Review]. Obstet. Gynecol. Clin. North Am. 1991;18(4):907-31.

Owen J, Henson BV, Hauth JC. A prospective randomized study of saline solution amnioinfusion. Am J Obstet Gynecol 1990;162(5):1146-9.

Owen J, Winkler CL, Harris BA, et al. A randomized, double-blind trial of prostaglandin E2 gel for cervical ripening and meta-analysis. Am J Obstet Gynecol 1991;165(4 Pt 1):991-6.

Oyarzun E, Gomez R, Rioseco A, et al. Antibiotic treatment in preterm labor and intact membranes: A randomized, double-blinded, placebo-controlled trial. J Matern Fetal Med 1998;7(3):105-10.

Papageorgiou I, Tsionou C, Minaretzis D, et al. Labor characteristics of uncomplicated prolonged pregnancies after induction with intracervical prostaglandin E2 gel versus intravenous oxytocin. Gynecol Obstet Invest 1992;34(2):92-6.

Park MI, Hwang JH, Cha KJ, et al. Computerized analysis of fetal heart rate parameters by gestational age. Int J Gynecol Obstet 2001;74:157-64.

Parry E, Parry D, Pattison N. Induction of labour for post term pregnancy: an observational study. Aust N Z J Obstet Gynaecol 1998;38(3):275-80.

Pattison N, McCowan L. Cardiotocography for antepartum fetal assessment (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Pearce JM, Cardozo C. Prolonged pregnancy: the management debate. BMJ 1988;297:715-7.

Peck T. Electronic monitoring evidence of fetal distress in high-risk pregnancies. J Reprod Med 1980;24(3):103-8.

Perkins RP. Sudden fetal death in labor. The significance of antecedent monitoring characteristics and clinical circumstances. J Reprod Med 1980;25(6):309-14.

Persson-Kjerstadius N, Forsgren H, Westgren M. Intrapartum amnioinfusion in women with oligohydramniosis. A prospective randomized trial. Acta Obstet Gynecol Scand 1999;78(2):116-9.

Peterson LP, Kundu N. Endocrine assessment of high-risk pregnancies. Obstet Gynecol Annu 1980; 9:169-94.

Phelan JP. Medical-legal considerations in the postdate pregnancy. Clin Obstet Gynecol 1989;32(2):294-302.

Phelan JP, Platt LD, Yeh SY, et al. The role of ultrasound assessment of amniotic fluid volume in the management of the postdate pregnancy. Am J Obstet Gynecol 1985;151(3):304-8.

Phelan JP, Platt LD, Yeh SY, et al. Continuing role of the nonstress test in the management of postdates pregnancy. Obstet Gynecol 1984;64(5):624-8.

Phelps JY, Higby K, Smyth MH, et al. Accuracy and intraobserver variability of simulated cervical dilatation measurements. Am J Obstet Gynecol 1995;173(3 Pt 1):942-5.

Phillips K, Berry C, Mathers AM. Uterine rupture during second trimester termination of pregnancy using mifepristone and a prostaglandin. Eur J Obstet Gynecol Reprod Biol 1996;65(2):175-6.

Piccolo AR. Clinical management of postdate pregnancy in the small community hospital. J Am Osteopath Assoc 1980;80(4):271-5.

Plaut MM, Schwartz ML, Lubarsky SL. Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. Am J Obstet Gynecol 1999;180(6 Pt 1):1535-42.

Pollack RN, Hauer-Pollack G, Divon MY. Macrosomia in postdates pregnancies: the accuracy of routine ultrasonographic screening. Am J Obstet Gynecol 1992;167(1):7-11.

Porreco RP. Postdates management: induction after 41 weeks, with cervical preparation. Birth 1990;17(4):227.

Porto M. The unfavorable cervix: methods of cervical priming. [Review]. Clin Obstet Gynecol 1989;32(2):262-8.

Ramrekersingh-White P, Farkas AG, Chard T, et al. Self-selected expectant management of post dates pregnancy including the use of Doppler ultrasound. J Obstet Gynaecol 1993;13(1):16-9.

Ramsey P, Harris D, Ogburn P, et al. Comparative efficacy of prostaglandin analogues dinoprostone and misoprostol as labor preinduction agents. Am J Obstet Gynecol 1998;178:S94.

Rand L, Robinson JN, Economy KE, et al. Post-term induction of labor revisited. Obstet Gynecol 2000;96(5):779-83.

Rasmussen S, Dalaker K, Nordbo BL, et al. One-stage ultrasound screening in pregnancy. An evaluation. Acta Obstet Gynecol Scand 1990;69(7-8):581-8.

Rayburn W, Gosen R, Ramadei C, et al. Outpatient cervical ripening with prostaglandin E2 gel in uncomplicated postdate pregnancies. Am J Obstet Gynecol 1988;158(6 Pt 1):1417-23.

Rayburn WF, Chang FE. Management of the uncomplicated postdate pregnancy. J Reprod Med 1981;26(2):93-5.

Rayburn WF, Motley ME, Stempel LE, et al. Antepartum prediction of the postmature infant. Obstet Gynecol 1982;60(2):148-53.

Reilly KEH. Induction of labor. Am Fam Physician 1994;49(6):1427-32.

Rekwot PI, Oyedipe EO, Mukasa-Mugerwa E, et al. Fertility in zebu cattle (Bos indicus) after prostaglandin administration and artificial insemination. Vet J 1999;158(1):53-8.

Ribbert LSM, Visser GHA, Mulder EJH, et al. Changes with time in fetal heart rate variation, movement incidences and haemodynamics in intrauterine growth retarded fetuses: A longitudinal approach to the assessment of fetal well being. Early Hum Dev 1993;31(3):195-208.

Ries M, Beinder E, Gruner C, et al. Rapid development of hydrops fetalis in the donor twin following death of the recipient twin in twin-twin transfusion syndrome. J Perinat Med 1999;27(1):68-73.

Rizzo N, Farina A, Santarsiero G, et al. Amniotic fluid index and labor length of pregnancies induced beyond 41 weeks of gestation with unfavorable

cervix. Gynecol Obstet Invest 2000;49(4):244-8.

Roach VJ, Rogers MS. Pregnancy outcome beyond 41 weeks gestation. Int J Gynaecol Obstet 1997;59(1):19-24.

Roberts CL, Algert CS, March LM. Delayed childbearing—are there any risks? Med J Aust 1994;160(9):539-44.

Roberts CL, Taylor L, Henderson-Smart D. Trends in births at and beyond term: evidence of a change? Br J Obstet Gynaecol 1999;106(9):937-42.

Roberts L, Cook E, Beardsworth SA, et al. Prolonged pregnancy: two years experience of offering women conservative management. J R Army Med Corps 1994;140(1):32-6.

Roberts LJ. Induction of labour. BMJ 1993;307(6895):66-7.

Roberts LJ, Young KR. The management of prolonged pregnancy—an analysis of women's attitudes before and after term. Br J Obstet Gynaecol 1991;98(11):1102-6.

Roberts WE, Morrison JC. Has the use of home monitors, fetal fibronectin, and measurement of cervical length helped predict labor and/or prevent preterm delivery in twins? Clin Obstet Gynecol 1998;41(1):95-102.

Rodriguez MH. Ultrasound evaluation of the postdate pregnancy. [Review]. Clin Obstet Gynecol 1989;32(2):257-61.

Romero R, Sibai BM, Sanchez-Ramos L, et al. An oxytocin receptor antagonist (atosiban) in the treatment of preterm labor: A randomized, double-blind, placebo-controlled trial with tocolytic rescue. Am J Obstet Gynecol 2000;182(5):1173-83.

Ron M, Adoni A, Hochner-Celnikier D, et al. The significance of baseline tachycardia in the postterm fetus. Int J Gynaecol Obstet 1980;18(1):76-7.

Rooks JP, Weatherby NL, Ernst EK. The National Birth Center Study. Part III. Intrapartum and immediate postpartum and neonatal complications and transfers, postpartum and neonatal care, outcomes, and client satisfaction. J Nurse Midwifery 1992;37(6):361-97.

Rose VL. ACOG issues report on the management of post-term pregnancy. American College of Obstetricians and Gynecologists. Am Fam Physician 1998;57(7):1686-7.

Rosen MG, Dickinson JC. Management of post-term pregnancy. N Engl J Med 1992;326(24):1628-9.

Rosser J, Anderson T. Sweeping the membranes. Pract Midwife 1998;1(3):28-9.

Rouse DJ, Owen J, Goldenberg RL, et al. The effectiveness and costs of elective cesarean delivery for fetal macrosomia diagnosed by ultrasound. JAMA 1996;276(18):1480-6.

Roussis P, Cox SM, Campbell BA, et al. Survey on the management of post date pregnancy. J Matern Fetal Invest 1993;2(3):155-7.

Royal College of Obstetricians and Gynaecologists. Induction of labour: evidence-based clinical guideline number 9. London: Royal College of Obstetricians and Gynaecologists, 2001.

Rubinstein TH, Schifrin BS. Prolonged labor with persistent occiput-posterior position in postterm pregnancy. J Perinatol 1992;12(2):181-4.

Ryo E, Kozuma S, Sultana J, et al. Fetal size as a determinant of obstetrical outcome of post-term pregnancy. Gynecol Obstet Invest 1999;47(3):172-6.

Sachs BP, Friedman EA. Results of an epidemiologic study of postdate pregnancy. J Reprod Med 1986;31(3):162-6.

Salamalekis E, Vitoratos N, Kassanos D, et al. Sweeping of the membranes versus uterine stimulation by oxytocin in nulliparous women. A randomized controlled trial. Gynecol Obstet Invest 2000;49(4):240-3.

Salamalekis E, Vitoratos N, Loghis C, et al. The predictive value of a nonstress test taken 24 h before delivery in high-risk pregnancies. Int J Gynaecol Obstet 1994;45(2):105-7.

Sanchez-Ramos L, Kaunitz AM, Del Valle GO, et al. Labor induction with the prostaglandin E1 methyl analogue misoprostol versus oxytocin: a randomized trial. Obstet Gynecol 1993;81(3):332-6.

Sandmire HF, DeMott RK. The Green Bay cesarean section study. IV. The physician factor as a determinant of cesarean birth rates for the large fetus.

Am J Obstet Gynecol 1996;174(5):1557-64.

Sarkar PK, Duthie SJ. The clinical significance of reduced amniotic fluid index in post-term pregnancy: a retrospective study. J Obstet Gynaecol 1997;17(3):274-5.

Sarkar PK, Williams J, Duthie J. Fetal macrosomia (1). J Obstet Gynaecol 1994;14(4):293.

Sarno AP, Hinderstein WN, Staiano RA. Fetal macrosomia in a military hospital: Incidence, risk factors, and outcome. Mil Med 1991;156(2):55-8.

Sarno AP, Polzin WJ, Feinstein SJ, et al. Transabdominal amnioinfusion in preterm pregnancies complicated by fetal growth restriction, oligohydramnios and umbilical cord compression. Fetal Diagn Ther 1995;10(6):408-14.

Satin AJ, Hankins GD. Induction of labor in the postdate fetus. [Review]. Clin Obstet Gynecol 1989;32(2):269-77.

Satin AJ, Hankins GD, Yeomans ER. A prospective study of two dosing regimens of oxytocin for the induction of labor in patients with unfavorable cervices. Am J Obstet Gynecol 1991;165(4 Pt 1):980-4.

Satin AJ, Leveno KJ, Sherman ML, et al. High-dose oxytocin: 20- versus 40-minute dosage interval. Obstet Gynecol 1994;83(2):234-8.

Sawai SK, O'Brien WF, Mastrogiannis DS, et al. Patient-administered outpatient intravaginal prostaglandin E2 suppositories in post-date pregnancies: a double-blind, randomized, placebo-controlled study. Obstet Gynecol 1994;84(5):807-10.

Sawai SK, Williams MC, O'Brien WF, et al. Sequential outpatient application of intravaginal prostaglandin E2 gel in the management of postdates pregnancies. Obstet Gynecol 1991;78(1):19-23.

Schan P. Why?... inter-uterine death at 42 weeks. Midwifery Matters 1997;74:12.

Schreyer P, Bar-Natan N, Sherman DJ, et al. Fetal breathing movements before oxytocin induction in prolonged pregnancies. Am J Obstet Gynecol 1991;165(3):577-81.

Schwartz RW. Pregnancy in physicians: characteristics and complications. Obstet Gynecol 1985;66(5):672-6.

Sciscione AC, Nguyen L, Manley J, et al. A randomized comparison of transcervical Foley catheter to intravaginal misoprostol for preinduction cervical ripening. Obstet Gynecol 2001;97:603-7.

Sciscione AC, Nguyen L, Manley JS, et al. Uterine rupture during preinduction cervical ripening with misoprostol in a patient with a previous Caesarean delivery. Aust N Z J Obstet Gynaecol 1998;38(1):96-7.

Scollo P. Epidemiology of cesarean sections: prolonged pregnancy. [Review]. Clin Exp Obstet Gynecol 1999;26(1):22-6.

Semczuk M, Lopucka M. Evaluation of cervix condition according to Bishop score in post term pregnancy. Ann Univ Mariae Curie Sklodowska [Med] 1986;41:125-31.

Setness P. Prolonged pregnancy. When to wait, when to intervene. Postgrad Med 1988;84(3):61-5, 69.

Seyb ST, Berka RJ, Socol ML, et al. Risk of cesarean delivery with elective induction of labor at term in nulliparous women. Obstet Gynecol 1999;94(4):600-7.

Shapiro H, Lyons E. Late maternal age and postdate pregnancy. Am J Obstet Gynecol 1989;160(4):909-12.

Sharma JB, Smith RJ, Wilkin DJ. Induction of labour at term. Women not for waiting. BMJ 1993;306(6889):1413.

Shaw K, Clark SL. Reliability of intrapartum fetal heart rate monitoring in the postterm fetus with meconium passage. Obstet Gynecol 1988;72(6):886-9.

Shaw KJ, Medearis AL, Horenstein J, et al. Selective labor induction in postterm patients. Observations and outcomes. J Reprod Med 1992;37(2):157-61.

Shea KM, Wilcox AJ, Little RE. Postterm delivery: a challenge for epidemiologic research. Epidemiology 1998;9(2):199-204.

Shearer MH, Estes M. A critical review of the recent literature on postterm pregnancy and a look at women's experiences. Birth 1985;12(2):95-111.

Sherer DM, Onyeije CI, Binder D, et al. Uncomplicated baseline fetal tachycardia or bradycardia in postterm pregnancies and perinatal outcome. Am J Perinatol 1998;15(5):335-8.

Sherman DJ, Frenkel E, Pansky M, et al. Balloon cervical ripening with extra-amniotic infusion of saline or prostaglandin E2: a double-blind, randomized controlled study. Obstet Gynecol 2001;97(3):375-80.

Shime J. Influence of prolonged pregnancy on infant development. J Reprod Med Obstet Gynecol 1988;33(3):277-84.

Shime J, Gare DJ, Andrews J, et al. Prolonged pregnancy: surveillance of the fetus and the neonate and the course of labor and delivery. Am J Obstet Gynecol 1984;148(5):547-52.

Shime J, Librach CL, Gare DJ, et al. The influence of prolonged pregnancy on infant development at one and two years of age: a prospective controlled study. Am J Obstet Gynecol 1986;154(2):341-5.

Sholl JS. Abruptio placentae: Clinical management in nonacute cases. Am J Obstet Gynecol 1987;156(1):40-51.

Shy KK. Roundtable: postterm pregnancy. Antenatal testing and candid reassurance... part 2. Birth 1991;18(2):113.

Sibai BM, Taslimi M, Abdella TN, et al. Maternal and perinatal outcome of conservative management of severe preeclampsia in midtrimester. Am J Obstet Gynecol 1985;152(1):32-7.

Silber M, Larsson B, Uvnas-Moberg K. Oxytocin, somatostatin, insulin and gastrin concentrations vis-a-vis late pregnancy, breastfeeding and oral contraceptives. Acta Obstet Gynecol Scand 1991;70(4-5):283-9.

Sims ME, Walther FJ. Neonatal morbidity and mortality and long-term outcome of postdate infants. [Review]. Clin Obstet Gynecol 1989;32(2):285-93.

Skretek M, Bielecki M, Zdanowicz A, et al. Prognostic value of cardiotocographic acoustic test in post-term pregnancies. Ann Med Univ Bialyst Pol 1993;38(1):79-85.

Small ML, Phelan JP, Smith CV, et al. An active management approach to the postdate fetus with a reactive nonstress test and fetal heart rate decelerations. Obstet Gynecol 1987;70(4):636-40.

Smith CA, Crowther CA. Acupuncture for induction of labour (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Smith GCS. Life-table analysis of the risk of perinatal death at term and post term in singleton pregnancies. Am J Obstet Gynecol 2001;184:489-96.

Snyder SG, King EA. Prostaglandin gel for cervical ripening. J Fam Pract 1996;43(1):19-20.

Soper DE, Mayhall CG, Froggatt JW. Characterization and control of intraamniotic infection in an urban teaching hospital. Am J Obstet Gynecol 1996;175(2):304-9.

Spong CY, Beall M, Rodrigues D, et al. An objective definition of shoulder dystocia: prolonged head-to-body delivery intervals and/or the use of ancillary obstetric maneuvers. Obstet Gynecol 1995;86(3):433-6.

Sporken JMJ, Hein PR, Gerritsen SM. First experiences with Swan-Ganz measurements in patients with severe hypertension during pregnancy from an unselected population. Eur J Obstet Gynecol Reprod Biol 1986;21(3):135-41.

Srisomboon J, Piyamongkol W, Aiewsakul P. Comparison of intracervical and intravaginal misoprostol for cervical ripening and labour induction in patients with an unfavourable cervix. J Med Assoc Thai 1997;80(3):189-94.

Steel SA, Pearce JM. Specific therapy in severe fetal intrauterine growth retardation: failure of prostacyclin. J R Soc Med 1988;81(4):214-6.

Steer PJ. Postmaturity—much ado about nothing. Br J Obstet Gynaecol 1986;93(2):105-8.

Stenlund PM, Ekman G, Aedo A-R, et al. Induction of labor with mifepristone: a randomized, double-blind study versus placebo. Acta Obstet Gynecol Scand 1999;78(9):793-8.

Steyn DW, Odendaal HJ. Computerised cardiotocography in a high-risk unit in a developing country—its influence on inter-observer variation and duration of recording. S Afr Med J 1996;86(2):172-5.

Stroup DF, Berlin JA, Morton SC, et al. Meta-

analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA 2000;283(15):2008-12.

Stubblefield PG, Berek JS. Perinatal mortality in term and post-term births. Obstet Gynecol 1980;56(6):676-82.

Sue-A-Quan AK, Hannah ME, Cohen MM, et al. Effect of labour induction on rates of stillbirth and cesarean section in post-term pregnancies. CMAJ 1999;160(8):1145-9.

Suikkari AM, Jalkanen M, Heiskala H, et al. Prolonged pregnancy: induction or observation. Acta Obstet Gynecol Scand 1983;116(Suppl):58.

Sulik SM, Greenwald JL. Evaluation and management of postdate pregnancy. [Review]. Am Fam Physician 1994;49(5):1177-86, 1191-2.

Summers L. Methods of cervical ripening and labor induction. [Review]. J Nurse Midwifery 1997;42(2):71-85.

Surbek DV, Boesiger H, Hoesli I, et al. A double-blind comparison of the safety and efficacy of intravaginal misoprostol and prostaglandin E2 to induce labor. Am J Obstet Gynecol 1997;177(5):1018-23.

Suzuki S, Otsubo Y, Sawa R, et al. Clinical trial of induction of labor versus expectant management in twin pregnancy. Gynecol Obstet Invest 2000;49(1):24-7.

Tabor B, Anderson J, Stettler B, et al. Misoprostol vs prostaglandin E2 gel for cervical ripening. Am J Obstet Gynecol 1995;172:425.

Tai-Seale M, Rodwin M, Wedig G. Drive-through delivery: where are the "savings"? Med Care Res Rev 1999;56(1):30-46.

Takahashi K, Diamond F, Bieniarz J, et al. Uterine contractility and oxytocin sensitivity in preterm, term, and postterm pregnancy. Am J Obstet Gynecol 1980;136(6):774-9.

Tal Z, Frankel ZN, Ballas S, et al. Breast electrostimulation for the induction of labor. Obstet Gynecol 1988;72(4):671-4.

Tam W-H, Tai SMB, Rogers MS. Prediction of cervical response to prostaglandin Einf 2 using fetal fibronectin. Acta Obstet Gynecol Scand 1999;78(10):861-5.

Tan KH. Fetal manipulation for facilitating tests of fetal wellbeing (Cochrane Protocol). In: The Cochrane Library, Issue 3, 2000. Oxford: Update Software.

Taylor HA. Induction of labor vs. antenatal monitoring in post-term pregnancy. Am Fam Physician 1997;55(7):2428, 2431.

Taylor NF. Review: placental sulphatase deficiency. [Review]. J Inherit Metab Dis 1982;5(3):164-76.

Tchobroutsky C, Vray MM, Altman J-J. Risk/benefit ratio of changing late obstetrical strategies of insulin-dependent diabetic pregnancies. A comparison between 1971-1977 and 1978-1985 periods in 389 pregnancies. Diabetes Metabol 1991;17(2):287-94.

Tessarolo M, Bellino R, Arduino S, et al. Cervical cerclage for the treatment of patients with placenta previa. Clin Exp Obstet Gynecol 1996;23(3):184-7.

Thacker SB, Stroup DF. Continuous electronic heart rate monitoring for fetal assessment during labor (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Thompson G, Newnham JP, Roberman BD, et al. Contraction stress fetal heart rate monitoring at preterm gestational ages. Aust N Z J Obstet Gynaecol 1990;30(2):120-3.

Thornton YS, Yeh SY, Petrie RH. Antepartum fetal heart rate testing and the post-term gestation. J Perinat Med 1982;10(4):196-202.

Todros T, Preve CU, Plazzotta C, et al. Fetal heart rate tracings: observers versus computer assessment. Eur J Obstet Gynecol Reprod Biol 1996;68(1-2):83-6.

Tongsong T, Srisomboon J. Amniotic fluid volume as a predictor of fetal distress in postterm pregnancy. Int J Gynaecol Obstet 1993;40(3):213-7.

Toppozada MK, Anwar MY, Hassan HA, et al. Oral or vaginal misoprostol for induction of labor. Int J Gynaecol Obstet 1997;56(2):135-9.

Toth M, Rehnstrom J, Fuchs AR. Prostaglandins E and F in cervical mucus of pregnant women. Am J

Perinatol 1989;6(2):142-4.

Treacy B, Smith C, Rayburn W. Ultrasound in labor and delivery. [Review]. Obstet Gynecol Surv 1990;45(4):213-9.

Trimmer KJ, Leveno KJ, Peters MT, et al. Observations on the cause of oligohydramnios in prolonged pregnancy. Am J Obstet Gynecol 1990;163(6 I):1900-3.

Trofatter KF. Endocervical prostaglandin E2 gel for preinduction cervical ripening. Clinical trial results. J Reprod Med 1993;38(1 Suppl):78-82.

Tsuei JJ, Lai Y, Sharma SD. The influence of acupuncture stimulation during pregnancy: the induction and inhibition of labor. Obstet Gynecol 1977;50(4):479-8.

Tuffnell DJ, Bryce F, Johnson N, et al. Simulation of cervical changes in labour: reproducibility of expert assessment. Lancet 1989;2(8671):1089-90.

Tuncer R, Erkaya S, Sipahi T, et al. Emergency postpartum hysterectomy. J Gynecol Surg 1995;11(4):209-13.

Tunon K, Eik-Nes SH, Grottum P. Fetal outcome in pregnancies defined as post-term according to the last menstrual period estimate, but not according to the ultrasound estimate. Ultrasound Obstet Gynecol 1999;14(1):12-6.

Turner JE, Burke MS, Porreco RP, et al. Prostaglandin E2 in tylose gel for cervical ripening before induction of labor. J Reprod Med 1987;32(11):815-21.

Turner MJ, Fox R, Gordon H. Induction of labour in primiparae after 41 weeks of pregnancy using vaginal prostaglandins. J Obstet Gynaecol 1988;8(Suppl. 1):S14-S15.

Turnley MJ. Round table: postterm pregnancy. Our moment with Courtney. Birth 1990;17(4):224-6.

Ugwumadu A. Induction of labour confers benefits in prolonged pregnancy. Br J Obstet Gynaecol 1995;102(1):79-81.

Umbach DM, Wilcox AJ. A technique for measuring epidemiologically useful features of birthweight distributions. Stat Med 1996;15(13):1333-48.

Usher RH, Boyd ME, McLean FH, et al. Assessment of fetal risk in postdate pregnancies. Am J Obstet Gynecol 1988;158(2):259-64.

Valensise H, Dell'anna D, Menghini S, et al. Computerized analysis of fetal heart rate in postterm pregnancy. J Matern Fetal Invest 1994;4(4):251-6.

Van de Pas PM, Nijhuis JG, Jongsma HW. Fetal behaviour in uncomplicated pregnancies after 41 weeks of gestation. Early Hum. Dev. 1994;40(1):29-38.

Van GMJC, Major AL, Scherjon SA. Placental anatomy, fetal demise and therapeutic intervention in monochorionic twins and the transfusion syndrome: New hypotheses. Eur J Obstet Gynecol Reprod Biol 1998;78(1):53-62.

Vantanasiri C, Manassakorn J, Thitadilok W, et al. False reactive nonstress tests in postterm pregnancies. J Med Assoc Thai 1985;68(5):266-8.

Varaklis K, Gumina R, Stubblefield PG. Randomized controlled trial of vaginal misoprostol and intracervical prostaglandin E2 gel for induction of labor at term. Obstet Gynecol 1995;86(4 Pt 1):541-4.

Ventura SJ, Martin JA, Curtin SC, et al. Births: Final data for 1998. National Vital Statistics Report, vol 48, no 3. Hyattsville, MD. National Center for Health Statistics, 2000.

Villar J, Gulmezoglu AM, De OM. Nutritional and antimicrobial interventions to prevent preterm birth: An overview of randomized controlled trials. Obstet Gynecol Surv 1998;53(9):575-85.

Voss DH, Cumminsky KC, Cook VD, et al. Effects of three concentrations of intracervical prostaglandin Einf 2 gel for cervical ripening. J Matern Fetal Med 1996;5(4):186-93.

Votta RA, Cibils LA. Active management of prolonged pregnancy. Am J Obstet Gynecol 1993;168(2):557-63.

Wald NJ, Cuckle HS, Boreham J, et al. Maternal serum alpha-fetoprotein and birth weight. Br J Obstet Gynaecol 1980;87(10):860-3.

Waldenstrom U, Nilsson S, Fall O, et al. Effects of routine one-stage ultrasound screening in pregnancy: A randomised controlled trial. Lancet 1988;2(8611):585-8.

Walles B, Tyden T, Herbst A, et al. Maternal health care program and markers for late fetal death. Acta Obstet Gynecol Scand 1994;73(10):773-8.

Webb GW, Raynor BD, Huddleston JF, et al. Induction of labor with an unfavorable cervix: a randomized prospective trial. Am J Obstet Gynecol 1997;176:S22.

Webster LA, Daling JR, McFarlane C, et al. Prevalence and determinants of caesarean section in Jamaica. J Biosoc Sci 1992;24(4):515-25.

Weeks JW, Pitman T, Spinnato JA 2nd. Fetal macrosomia: does antenatal prediction affect delivery route and birth outcome? Am J Obstet Gynecol 1995;173(4):1215-9.

Weerasekera DS, Grant JM. Sweeping of the membranes is an effective method of induction of labour in prolonged pregnancy: a report of a randomised trial (2). Br J Obstet Gynaecol 1993;100(2):193-4.

Weiner Z, Farmakides G, Barnhard Y, et al. Doppler study of the fetal cardiac function in prolonged pregnancies. Obstet Gynecol 1996;88(2):200-2.

Weiner Z, Farmakides G, Schulman H, et al. Central and peripheral haemodynamic changes in post-term fetuses: Correlation with oligohydramnios and abnormal fetal heart rate pattern. Br J Obstet Gynaecol 1996;103(6):541-6.

Weiner Z, Farmakides G, Schulman H, et al. Computerized analysis of fetal heart rate variation in postterm pregnancy: prediction of intrapartum fetal distress and fetal acidosis. Am J Obstet Gynecol 1994;171(4):1132-8.

Weiner Z, Reichler A, Zlozover M, et al. The value of Doppler ultrasonography in prolonged pregnancies. Eur J Obstet Gynecol Reprod Biol 1993;48(2):93-7.

Weinstein D, Ezra Y, Picard R, et al. Expectant management of post-term patients: observations and outcome. J Matern Fetal Med 1996;5(5):293-7.

Weissman A, Yoffe N, Jakobi P, et al. Management of triplet pregnancies in the 1980s - Are we doing better? Am J Perinatol 1991;8(5):333-7.

Wikstrom I, Axelsson O, Bergstrom R, et al. Traumatic injury in large-for-date infants. Acta Obstet Gynecol Scand 1988;67(3):259-64.

Wilcox AJ, Weinberg CR, Baird DD. Timing of sexual intercourse in relation to ovulation. Effects on the probability of conception, survival of the pregnancy, and sex of the baby. N Engl J Med 1995;333(23):1517-21.

Wilcox JA, Nasrallah HA. Perinatal distress and prognosis of psychotic illness. Neuropsychobiology 1987;17(4):173-5.

Wing DA, Fassett MJ, Mishell DR. Mifepristone for preinduction cervical ripening beyond 41 weeks' gestation: a randomized controlled trial. Obstet Gynecol 2000;96(4):543-8.

Wing DA, Jones MM, Rahall A, et al. A comparison of misoprostol and prostaglandin E2 gel for preinduction cervical ripening and labor induction. Am J Obstet Gynecol 1995;172(6):1804-10.

Wing DA, Lovett K, Paul RH. Disruption of prior uterine incision following misoprostol for labor induction in women with previous cesarean delivery. Obstet Gynecol 1998;91(5 Pt 2):828-30.

Wing DA, Ortiz-Omphroy G, Paul RH. A comparison of intermittent vaginal administration of misoprostol with continuous dinoprostone for cervical ripening and labor induction. Am J Obstet Gynecol 1997;177(3):612-8.

Wing DA, Paul RH. A comparison of differing dosing regimens of vaginally administered misoprostol for preinduction cervical ripening and labor induction [published erratum appears in Am J Obstet Gynecol 1997 Jun;176(6):1423]. Am J Obstet Gynecol 1996;175(1):158-64.

Wing DA, Rahall A, Jones MM, et al. Misoprostol: an effective agent for cervical ripening and labor induction. Am J Obstet Gynecol 1995;172(6):1811-6.

Wiriyasirivaj B, Vutyavanich T, Ruangsri RA. A randomized controlled trial of membrane stripping at term to promote labor. Obstet Gynecol 1996;87(5 Pt 1):767-70.

Wischnik A, Zieger W. Tocolytic therapies in development for preterm labour. Expert Opin Invest Drugs 1994;3(6):621-9.

Witter FR, Weitz CM. A randomized trial of induction at 42 weeks gestation versus expectant

management for postdates pregnancies. Am J Perinatol 1987;4(3):206-11.

Witter FR, Weitz CM. Cervical examination prior to induction in postdate pregnancies. Surg Gynecol Obstet 1989;168(3):214-6.

Wood CL. Postdate pregnancy update. [Review]. J Nurse Midwifery 1994;39(2 Suppl):110S-22S.

Wood L. Induction of labour confers benefit in prolonged pregnancy. Br J Obstet Gynaecol 1994;101(12):1102.

Wu MY, Yang YS, Huang SC, et al. Prolongation of pregnancy and survival of both twins after preterm premature rupture of membrane of twin A and 19 weeks' gestation. Acta Obstet Gynecol Scand 1996;75(3):299-302.

Xenakis EM, Piper JM, Conway DL, et al. Induction of labor in the nineties: conquering the unfavorable cervix. Obstet Gynecol 1997;90(2):235-9.

Yates WT, Cibils LA. Why not begin 'active management' of prolonged pregnancy sooner? (2). Am J Obstet Gynecol 1994;170(1 I):253.

Yeast JD, Jones A, Poskin M. Induction of labor and the relationship to cesarean delivery: A review of 7001 consecutive inductions. Am J Obstet Gynecol 1999;180(3 Pt 1):628-33.

Yeh S, Huang X, Phelan JP. Postterm pregnancy after previous cesarean section. J Reprod Med 1984;29(1):41-4.

Yeh SY, Bruce SL, Thornton YS. Intrapartum monitoring and management of the postdate fetus. [Review]. Clin Perinatol 1982;9(2):381-6.

Yeh SY, Read JA. Management of post-term pregnancy in a large obstetric population. Obstet Gynecol 1982;60(3):282-7.

Yeh SY, Read JA. Plasma unconjugated estriol as an indicator of fetal dysmaturity in postterm pregnancy. Obstet Gynecol 1983;62(1):22-5.

Yoonessi M, Wieckowska W, Mariniello D, et al. Cervical intra-epithelial neoplasia in pregnancy. Int J Gynecol Obstet 1982;20(2):111-8.

Yudkin P. Risk of unexplained stillbirth in prolonged pregnancy. Midwife Health Visit Community Nurse 1988;24(10):407-10.

Yudkin PL, Redman CW. Caesarean section dissected, 1978-1983. Br J Obstet Gynaecol 1986;93(2):135-44.

Yudkin PL, Wood L, Redman CWG. Risk of unexplained stillbirth at different gestational ages. Lancet 1987;1:1192-4.

Zelop CM, Shipp TD, Cohen A, et al. Trial of labor after 40 weeks' gestation in women with prior cesarean. Obstet Gynecol 2001;97:391-3.

Zhao YG, Shao YL. The combined tests of overnight 12-hour urine E3/C ratio and fetal heart rate in perinatal monitoring. Chin Med J (Engl) 1989;102(3):188-92.

Zhou W, Sorensen HT, Olsen J. Induced abortion and subsequent pregnancy duration. Obstet Gynecol 1999;94(6):948-53.

Zuazu J, Julia A, Sierra J, et al. Pregnancy outcome in hematologic malignancies. Cancer 1991;67(3):703-9.