

# Index

- abdominal pain, acute
  - analgesics for, 396
  - analgesics studies, 397, 399
- academic detailing for modifying physician behavior, 595, 597
- accreditation methods to promote patient safety practices, 607
- accreditation organizations, 602
- active error, 51
- acute pain services
  - described, 400
  - studies, 401, 403
- acute spinal cord injury, VTE risk, 338
- ADE detection
  - alert monitors, 79
  - computerized alert monitors, 79, 80
  - computerized systems, studies, 83
  - pharmacists, physicians, and, 79
  - physician response to alert monitors, 81
  - research needs, 82
- ADEs
  - anticoagulants and, 87
  - CDSS and, 590
  - clinical pharmacists and, 71, 74, 75
  - computerized detection and alert systems, 79
  - described, 79
  - incident reporting, 42
  - information transfer between inpatient/outpatient pharmacies, 471
  - prevalence, 71
- advance directives for end-of-life care, 556, 560
- Agency for Health Care Policy and Research (AHCPR), 396, 406
- Agency for Healthcare Research and Quality (AHRQ)
  - acknowledgments, 21
  - patient advocacy, 569
  - patient safety, 601
  - patient safety improvement, 13
- American Medical Association (AMA) Code of Medical Ethics, 546
- American Society for Health System Pharmacists (ASHP), antibiotic administration, 221
- American Society of Anesthesiologists (ASA), 259, 265, 407, 604
- amphotericin, topical, 193
- analgesics
  - acute abdominal pain management, 396
  - acute abdominal pain management studies, 397, 399
  - antiemetics for PCA, 404
  - research needs, 398
  - surgeon decision making, impact on, 398
- anesthesia
  - body temperature and, 231
  - intraoperative monitoring and, 265
  - pre-anesthesia checklists, 259
  - simulator-based training, 512
- anesthesia crisis resource management (ACRM), 504
- anesthesia equipment checklists, 466
- antibiotic resistance and pneumococcal vaccines, 390
- antibiotics
  - benefits for controlling nosocomial infections, 143
  - practices to change use, 141
  - prescription errors, 142
  - prophylaxis for SSI, 221, 224, 226
  - urinary catheters and, 149
- anticoagulation
  - ADEs and, 88
  - inpatient services, 94
  - outpatient clinics, 88
  - patient self-management, 89, 96
  - research needs, 91
- antiemetics for pain management
  - described, 404
  - studies, 405
- ASEPSIS, 232, 235
- aspirin to prevent venous thromboembolism, 333
- atelectasis, 185, 234
- audit and feedback for modifying physician behavior, 595, 596
- Australian Incident Monitoring Study (AIMS), 43, 45, 266
- bacteremia and urinary catheters, 149
- bacteriuria and urinary catheters, 149
- bar coding
  - bedside scanning, 491
  - described, 487
  - medical recordkeeping, 489
  - medication dispensing and administration, 488
  - patient identification, 487
  - patient identification studies, 490, 491
  - research needs, 492
  - specimen handling, 488
- barrier precautions, central venous catheter insertion
  - contamination reduction, 165
  - studies, 165
- barrier precautions, VRE and *C. difficile*
  - compliance vs. specific intervention, 127
  - described, 127
  - further research, 130
  - list of, 127
  - studies, 129, 132, 135, 136
- Baxter ATC-212 dispensing system, 111
- bed alarms for fall prevention in older patients
  - described, 291
  - studies, 292, 293
- bed sores, 156
- beds or mattresses for pressure ulcer prevention
  - described, 301
  - studies, 302, 304
- bedsores, 301
- behavior-based safety interventions, 450
- Best Medicine, The*, 570
- beta-blockers, perioperative
  - described, 271
  - research needs, 274
  - studies, 272, 276
- bloodstream infections, catheter-related (CR-BSI)
  - defined, 163
  - risks, 164
- bronchopulmonary secretions, 185
- business community and patient safety, 603

calorimetry, 359  
 Candida species in catheter-related infections, 163  
 capnography, 267  
 cardiac disease, heparin and warfarin monitoring, 88  
 cardiac events  
     hypothermia, 231  
     reducing with beta blockers, 271  
 care delivery and interventions, nurse staffing and, 423, 426, 430  
 catheter colonization, 163  
 catheter related infections  
     central venous, 167  
 catheter-related bloodstream infection (CR-BSI)  
     antibacterial coatings and antiseptic agents, 169  
     CHG solutions and, 175  
     defined, 163, 164  
     intravenous antimicrobial use of, 179  
 catheter-related infections  
     bloodstream, 163  
     central venous, 164  
     urinary tract, 149  
 Centers for Disease Control and Prevention (CDC), 42, 164, 194, 602  
     antibiotic administration, 221  
     pneumococcal vaccines, 385  
 central venous catheters  
     antibacterial coatings or antiseptic agents, 169, 174  
     barrier precautions during insertion, 165  
     CHG and PI studies, 177, 178  
     chlorhexidine gluconate at the insertion site, 175  
     heparin and, 178  
     infection studies, 167  
     placement of, 245  
     prolonged use, infection risks and risk reduction, 165  
     research needs, 183  
     ultrasound guidance for insertion, 246  
     use of, 163  
 children, transporting, 535, 536, 537  
 chlorhexidine gluconate (CHG) solution  
     central venous catheter insertion, 175  
     compared to PI, studies, 177, 178  
     studies, 175  
 chlorhexidine/silver sulfadiazine-impregnated catheters  
     described, 169  
     studies, 169, 172, 174  
 circadian rhythm, 519, 524  
 clinical decision support systems (CDSS)  
     CPOE and, 589  
     described, 589  
     research needs, 593  
     safety practice promotion and implementation, 590  
     studies, 590, 591  
 clinical pharmacists  
     ADEs and, 71  
     ambulatory settings, study results, 73  
     consultation services, 71  
     impact on ADEs, studies, 75  
     outpatient settings, studies, 72  
     roles and activities, 71  
 closed ICU model, 413  
 Clostridium difficile (C. difficile)  
     antibiotic studies, 142, 143  
     antibiotics and, 141  
     barrier precaution studies, 129  
     barrier precautions and, 127  
     relationship to VRE, 128  
     research needs, 131, 144  
     SSI and, 222  
     transferability, 127  
 Cockpit Management Attitudes Questionnaire (CMAQ), 502  
 computerized discharge summaries, 479  
 computerized physician order entry (CPOE) with CDSS  
     CDSS described, 589  
 computerized sign-out program, 476, 479  
 continuous intravenous insulin infusion (CII) during surgery, 238  
 continuous oscillation  
     described, 185, 187  
     studies, 187, 189  
 cranial computed tomography (CT), 376  
 Crew Resource Management (CRM)  
     aviation application vs. health care, 504  
     aviation environment, 501  
     described, 501  
     effectiveness measurement tools, 502  
     health care environment, 504  
     operating room environment, 505  
     peer monitoring, 505  
     research needs, 507  
     studies, 503, 506  
 critical incident technique, 41  
 critical pathways  
     described, 581  
     medical information dissemination, 581  
     research needs, 584  
     studies, 582, 585  
 cross-coverage, 476  
*Crossing the Quality Chasm: A New Health System for the 21<sup>st</sup> Century*, 13  
 decubitus ulcers, 156, 301  
 deep vein thrombosis (DVT), 333, 341  
 delirium in older hospitalized patients  
     described, 307  
     geriatric consultation services and, 313  
     interventions, 307  
     research needs, 309  
     studies, 308, 310  
 diabetes  
     glucose control during surgery, 237  
     RCIN and, 349  
 diarrhea, C. difficile-associated, 128  
 direct mail patient notification, 483  
 discharge summaries  
     computerized, 479  
     research needs, 481  
     structured vs. unstructured format, 479  
     studies, 480, 481  
 disseminating medical information, 581  
 distal DVT, 333  
 domains, defined, 29  
 duplex ultrasonography, 334  
 durable power of attorney for health care (DPOA-HC), 556  
 educational techniques to modify physician behavior

- described, 595
- research needs, 598
- studies, 596, 599
- elastic stockings (ES) to prevent venous thromboembolism, 333
- elderly
  - beta-blockers and, 273
  - delirium, preventing, 307
  - falling, preventing, 281
  - falls, preventing, 283
  - geriatric consultation services, 313
  - pneumococcal infections, 386
  - pneumococcal vaccines and, 387
  - pressure ulcers, preventing, 301
  - preventable deaths and functional decline, 324
- embolism to the pulmonary vasculature, 333
- end-of-life care
  - administrative initiatives, 560
  - advance directives, 560
  - described, 556
  - documenting patient preferences, 557
  - existing programs/contributions, 562
  - forms to promote implementation of patient preferences, 559
  - implementation of patient preferences, 557
  - improving physician/patient communication, 560
  - physician/patient communication, 558
  - preferences in outpatient settings, 560
  - public education, 562
  - research needs, 563
- enteral nutrition (EN)
  - immune-enhanced, 363
  - paths for administering, 359
  - studies, 361
- environmental modifications to prevent delirium, 307
- equipment, anesthesia, checklists for, 259
- error mode prediction, 26
- examination equipment, dedicated or disposable, as a barrier precaution, 127, 136
- extended-spectrum beta-lactamase (ESBL), 128, 143
- falls in hospitalized and institutionalized older patients
  - bed alarms and, 291
  - geriatric consultation services and, 313
  - hip protectors and, 295
  - identification bracelets and, 284
  - interventions, list of, 283
  - physical restraints and, 287
  - problem description, 281
  - research needs, 283, 284, 289, 292, 294, 297
  - risk assessment, 282
  - risk factors, 281
  - special flooring and, 293
- fatigue
  - lighting at work, 524
  - medical errors and, 519
  - medical therapies, 525
  - napping, 524
  - research needs, 526
  - shift limits and error reduction, 521
  - shift rotation and, 520
  - shift rotation, direction and speed, 523
  - sleep hygiene education, 523
  - studies, 522
- fibrinogen leg scanning, 334
- Flight Management Attitudes Questionnaire (FMAQ), 502
- flooring, special, for fall prevention in older patients
  - described, 293
  - studies, 294, 295
- Foley catheters, 149
- Food and Drug Administration (FDA), 42, 51, 259, 262, 459, 489
- gastroesophageal reflux, 185
- gastrointestinal bleeding
  - described, 368
  - nosocomial pneumonia and, studies, 372
  - research needs, 371
  - studies, 369
  - sucralfate and, 199
- gentamicin, topical, 193
- geriatric consultation services
  - described, 313
  - research needs, 316
  - studies, 314, 315, 318
- geriatric evaluation and management (GEM) units
  - described, 313, 323
  - research needs, 326
  - studies, 324, 327
- gloves/gowns as barrier precaution, 127
- glucose control, perioperative
  - described, 237
  - research needs, 240
  - studies, 238, 241
  - surgical infections and, 238
- government agencies and patient safety practices, 602
- gram negative rods in catheter-related infections, 163
- H<sub>2</sub>-receptor antagonists, 198, 368, 369, 372
- hand disinfection, 119
- handwashing
  - compliance difficulties, 119
  - relationship to infection, 119
  - research needs, 122
  - risk factors, 120
  - skin irritation to health care workers, 121
  - strategies to improve compliance, 121
  - studies, 120, 123
- Health Care Financing Administration (HCFA)
  - medical errors, 601
  - physical restraints and, 288
  - pneumococcal vaccines, 287, 385
- health literacy problem, 547
- heart failure
  - ADEs and, 72
  - clinical pharmacists and, 73
- heparin
  - about, 87
  - bleeding during therapy with, 88
  - central venous catheters and, 178
  - dosing and monitor errors, reducing, 87
  - LDUH/LMWH to prevent venous thromboembolism, 333
  - monitoring, 88
- heparin nomograms
  - described, 87
  - PTTs and, 89

- studies, 88, 92
- high reliability theory, 26, 447
- high-volume centers, localizing care to
  - Leapfrog Group's efforts, 208
  - relationship to patient outcomes, 205
  - research needs, 208
  - studies, 206, 210
- hip fractures
  - falls in older patients, 281
  - VTE risk in surgery, 336
- hip protectors for fall prevention in older patients, 296, 298
- hip replacement surgery, VTE risk, 336
- histamine-2 receptor blocking agents (H<sub>2</sub>-receptor antagonists)
  - described, 368
  - studies, 369, 372
  - VAP prevention, 198
- hospitalists and end-of-life care, 561
- human error identification, 26
- human factors engineering (HFE)
  - anaesthesia equipment checklists, 466
  - described, 459
  - medical device alarms, 462
  - medical devices and errors, 459
  - research needs, 467
- hydration protocols to prevent RCIN, 349
- hygienic hand rub, 119
- hypercholesterolemia, clinical pharmacists and, 73
- hypertension, clinical pharmacists and, 73
- hypoxemia and anesthesia, 267
- identification bracelets for fall prevention in older patients
  - study, 285
  - using, 284
- incident reporting
  - AIMS data vs. JCAHO data, 45
  - barriers to linking data with outcome, 46
  - critical incident monitoring, 41
  - event appropriateness, 43
  - intensive care units (ICUs), 42
  - JCAHO surveillance mandates, 43
  - medication errors and, 41
  - physician reporting, 45
  - research needs, 46
  - sentinel event alerts, 47
- industries outside health care
  - aviation, 259, 501
  - behavior-based safety programs, 450
  - corporate cultures, 448
  - critical pathways, 581
  - disasters contributed to by fatigue, 520
  - incident reporting systems, 41, 42
  - safety cultures, 449
  - simulator training, 511
- infection control officers, 602
- information transfer, improving
  - abnormal results notification, 482
  - discharge summaries, 479
  - inpatient/outpatient pharmacies, 471
  - research needs, 474, 486
  - sign-out systems, 476
- informed consent
  - asking patient to recall discussion information, studies, 550
  - form readability studies, 549
  - forms inadequacies, 548
  - legal and ethical obligations, 546
  - physician/patient discussion, studies, 549
  - procedural inefficiencies in obtaining, 547
  - process, 546
  - research needs, 552
  - visual or auditory aids, studies, 550
  - written information to patients, studies, 551
- Institute of Medicine (IOM) reports, 13, 205, 207, 210
- intensive care units (ICUs)
  - incident reporting and, 42
  - intensivist model studies, 418
  - Leapfrog Group recommendations, 413
  - nutritional support for patients, 359
  - research needs, 416
  - structure model studies, 414, 415
  - structure models, 413
- intensivist co-management ICU model, 413
- intensivists in ICUs, 413
- interhospital transportation of critically ill patients, 536, 537
- interhospital transportation of critically ill patients, 534
- intermittent pneumatic compression (IPC) to prevent venous thromboembolism, 333
- International Classification of Diseases (ICD-9) codes, 79
- intra-hospital transportation of critically ill patients, 534, 538
- ischemia, myocardial, 267, 271
- Joint Commission on the Accreditation of Healthcare Organizations (JCAHO)
  - pain management guidelines, 396
  - patient safety advocacy, 602
  - physical restraint mandates, 288
  - root cause analysis mandate, 51
  - safety standards, 606
  - sentinel event alerts, 47
  - Sentinel Event Database, 43
  - surveillance mandate, 43
  - wrong-site surgery, avoiding, recommendations, 495
- knee replacement surgery, VTE risk, 336
- laparoscopic cholecystectomy training
  - complications with procedure, 214
  - learning curve, 213
  - research needs, 216
  - skills needed, 213
  - studies, 215
- latent error, 51
- Leapfrog Group
  - high-volume centers, 205
  - ICUs model recommendations, 413
  - methods to promote patient safety practices, 607
  - patient safety advocacy, 603
  - referral system, 208
- learning curves for new procedures, 213
- legal issues
  - end-of-life care, 556
  - informed consent, 546
- legislation
  - methods to promote patient safety practices, 607

Natural Death Acts, 556  
 nursing issues, 425  
 Omnibus Budget Reconciliation Act of 1987, 288, 302  
 patient safety, 601  
 Patient Self-Determination Act (PSDA), 556  
 publication of performance data, 605  
*Lerner's Consumer Guide*, 569  
 lighting and fatigue, 524  
 LINE/LOS checklist, 502  
 living will for end-of-life care, 556  
 localizing care to high-volume centers, 205  
 low dose unfractionated heparin (LDUH) to prevent VTE, 333, 335  
 low molecular weight heparin (LMWH) to prevent VTE, 333, 335  
 machine-readable automatic identification (ID) systems  
   conventional wrist band use, 489  
   described, 487  
   medical record keeping, 489  
   medication dispensing and administration, 488  
   patient identification, 487  
   patient identification studies, 490, 491  
   research needs, 492  
   specimen handling, 488  
 malnutrition in hospitalized patients, 359, 363  
 McLaughlin dispensing system, 111  
 median sternotomy, 238  
 medical device alarms, 462  
   frequency reduction, 465  
   research needs, 466  
   source identification, 463  
   urgency ranking, 463  
   visual interfaces, 463  
 medical devices  
   design and development, 459  
   device evaluation, 461, 462  
 medical recordkeeping and bar codes, 489  
 medical records, patient access to, 570  
 medication dispensing and bar codes, 488  
 medication dispensing systems, automated  
   Baxter ATC-212 system, 111  
   described, 111  
   harmpotential, 113  
   machine-readable code, 111  
   McLaughlin system, 111  
   research needs, 114  
   studies, 111, 112, 115  
   unit-dose distribution systems and, 112  
 medication errors  
   clinical pharmacists impact on, studies, 75  
   dosage errors, 112  
   identification of ADEs, 41  
   information transfer between inpatient/outpatient pharmacies, 471  
   studies, 106  
   time-constraints and, 101  
   unit-dose dispensing, 102  
 MedTeams behavior-based teamwork system, 505  
 melatonin supplementation for fatigue, 525  
 methicillin-resistant *Staphylococcus aureus* (MRSA), 128  
 minocycline/rifampin-coated catheters, 169  
 minocycline-coated urinary catheters, 149  
 misidentification, avoiding  
   bar coding, 487  
   wrong-site surgery, 494, 496, 497  
   wrong-site surgery, research needs, 498  
 monitoring, intraoperative  
   ASA standards, 268  
   described, 265  
   research needs, 267  
   studies, 266  
 napping, 524  
 National Quality Forum (NQF), 21  
 National Vaccine Advisory Committee, pneumococcal vaccines, 385  
 Natural Death Acts, 556  
 nausea due to PCA, 404  
 neurosurgery, VTE risk, 337  
 non-radiologists  
   radiograph interpretations, 376  
   training, 377, 381  
 normal accident theory, 447  
 normothermia, perioperative  
   described, 231  
   studies, 232, 233  
 nosocomial infections  
   antibiotics and, 141  
   barrier precautions and, 127  
   handwashing and, 119  
   handwashing practices and, 120  
   pathogens other than VRE and *C. difficile*, 128  
   pneumonia and GI bleeding, 368  
 nurse staffing  
   availability studies, 426, 431  
   care delivery models, 430  
   care delivery studies, 426  
   care practices evaluated, 424  
   intervention studies, 427  
   measures, 430  
   nurse intervention and patient outcome studies, 437  
   organization models and patient outcome studies, 435  
   relationship of models to patient outcomes, 423  
   research needs, 428  
   studies, 425  
   unions and, 425  
 nutritional support practices  
   described, 359  
   studies, 360, 365  
 Occupational Safety and Health Administration (OSHA), 601  
 open ICU model, 413  
 opinion leaders for modifying physician behavior, 595, 597  
 opioids, 404  
 orientation techniques to prevent delirium, 307  
 oropharyngeal secretions, 190  
 osmolar iodinated contrast media to prevent RCIN, 349  
 oxygen, perioperative  
   described, 234  
   future research needs, 236  
   studies, 235, 236  
 pain management  
   abdomen, acute, 396  
   about, 396  
   acute pain services, described, 400

- acute pain services, studies, 401, 403
- antiemetics for PCA, 404
- non-pharmacologic intervention studies, 407, 409
- non-pharmacologic interventions, 406
- patient education, 407
- research needs, 408
- palliative care programs, 561
- patient advocacy
  - benefits, 569
  - improving non-compliance with medical advice and treatment, 570
  - medical records, access to, 570
  - research needs, 571
- patient classification systems (PCSs), 423
- patient mobility to prevent delirium, 307
- patient notification of abnormal results
  - direct mail, 483
  - problems and current methods, 482
  - research needs, 484
  - study, 483, 485
- Patient Self-Determination Act (PSDA), 556
- patient to nurse ratios, 424
- peer monitoring, 505
- percutaneous transluminal angioplasty, 271
- pericatheter skin inflammation, 163
- pharmacies
  - automated drug dispensing, 102
  - information transfer, 471, 473, 475
- pharmacologic measures to prevent venous thromboembolism, 333
- physical restraints for fall prevention in older patients
  - described, 287
  - studies, 288, 290
- physician behavior, modifying
  - education techniques, research needs, 598
  - education techniques, studies, 596, 599
  - educational techniques, 595
  - practice guidelines, 576
- pneumococcal infections
  - re-infection risks, 385
  - vaccines and, 385
- pneumococcal vaccines
  - antibiotic resistance, 390
  - delivery method studies, 392
  - delivery methods, studies, 388
  - described, 385
  - effectiveness studies, 386, 387, 391
  - research needs, 390
- pneumothorax, 247
- POLST (Physician Orders for Life-Sustaining Treatment), 559, 564
- polymixin, topical, 193
- povidone-iodine (PI)
  - central venous catheter insertion, 175
  - compared to (CHG), studies, 177, 178
- practice guidelines for modifying physician behavior
  - computer-based, 577
  - described, 575
  - research needs, 578
  - studies, 576, 577
- pre-anesthesia equipment checklists
  - described, 259
  - research needs, 261
  - studies, 260, 262
- pressure ulcers
  - in older patients, 301
  - research needs, 304
  - suprapubic catheters and, 156
- professional societies and patient safety, 604
- professional societies, to promote patient safety practices, 607
- psychosocial support for pain management, 407
- public reporting systems, 605
- pulmonary embolization (PE), 333
- pulse oximetry, 265, 266, 466, 604
- Pyxis Medstations, 112
- radiocontrast-induced nephropathy (RCIN)
  - described, 349
  - prevention practices, 349
  - prevention studies, 353, 354
  - studies, 350, 351
- radiograph interpretation errors by non-radiologists
  - described, 376
  - digitized vs. film-based radiographs, 376
  - reducing, 377, 378, 381
  - research needs, 380
- radiologists and non-radiologists
  - radiograph interpretations, 376
  - training, 377
- reminder systems for modifying physician behavior, 595
- renal failure, RCIN and, 349
- report methodology
  - abstraction elements, 34
  - evidence review, 15
  - exclusion criteria, 31
  - exclusions, 16
  - inclusion criteria, 31
  - inclusions, 16
  - organization, 18
  - outcome measures, 32
  - review methodologies, 29
  - safety practice ratings, 613
  - safety practices evaluation, 33
  - safety practices organization, 29
  - search strategy, 30
  - study design hierarchy, 31
- review methodologies, error-based, 29
- rifampin-coated urinary catheters, 149
- risk assessment to prevent falls in older patients, 282
- root cause analysis (RCA)
  - industrial accidents, 51
  - JCAHO mandate, 51
  - process, 52
  - research needs, 55
  - studies, 53, 54
- safety culture, promoting
  - accident theories, 447
  - behavior-based programs, 450
  - checklist of patient safety concepts, 453
  - methods, medical and non-medical, 448
  - research needs, 451
  - VHA initiative, 449
- safety practice ratings
  - data inputs, 613

- decision rules and judgment considerations, 615
- goals, 613
- impact and evidence overall rating, 616
- rating process, 614
- rating tables, 617
- research priority overall rating, 616
- safety practices
  - all practices, categorical ratings, and comments, 633
  - clinical vs. general issues in content, 17
  - definition and scope, 13, 29
  - further research likely to be beneficial, 630
  - further research likely to be highly beneficial, 627
  - greatest strength of evidence regarding their impact and effectiveness, 620
  - high strength of evidence regarding their impact and effectiveness, 621
  - hospital care and, 14
  - industries outside health care, 25
  - low strength of evidence regarding their impact and effectiveness, 624
  - lowest strength of evidence regarding their impact and effectiveness, 625
  - medium strength of evidence regarding their impact and effectiveness, 622
  - organization of, 29
  - rated by strength of evidence, 619
  - research priority ratings, 627
- selective digestive tract decontamination (SDD)
  - described, 193
  - studies, 194, 196
- semi-recumbent positioning
  - described, 185
  - studies, 186, 189
- Sentinel Event Database, 43
- shift rotation and fatigue, 523
- sign-out systems for cross-coverage
  - described, 476
  - research needs, 478
  - studies, 477, 479
- silicone urethral catheters, 149
- silver alloy catheters
  - bacteriuria reduction, 151
  - described, 149
- simulator-based training
  - advantages, 511
  - anesthesia, 512
  - cardiology, 514
  - CRM and, 512
  - gastroenterology, 514
  - radiology, 513
  - research needs, 514
  - risks, 515
  - surgery, 513
- sleep deprivation
  - lighting at work, 524
  - medical errors and, 519
  - medical therapies speed, 525
  - napping, 524
  - research needs, 526
  - shift limits and error reduction, 521
  - shift rotation and, 520
  - shift rotation direction and speed, 523
  - sleep hygiene education, 523
  - studies, 522
- sponge, sharp, and instrument counts
  - described, 255
  - research needs, 256
  - studies, 256
- staff cohorting as a barrier precaution, 127
- staphylococci and catheter-related infections, 163
- sternal wound infections, 238
- streptococcus pneumoniae, 385
- stress ulcers
  - H<sub>2</sub>-receptor antagonist therapy, 198
  - studies, 201, 202
- subclavian-vein catheters, 246
- subglottic secretions, continuous aspiration of
  - described, 190
  - diagram, 192
  - studies, 193
- sucralfate
  - gastrointestinal bleeding reduction, described, 368
  - gastrointestinal bleeding reduction, studies, 369, 372
  - studies, 201, 202
  - VAP reduction, 198
- suprapubic catheters
  - compared to standard catheters, 156
  - compared to urethral, studies, 159
  - placement of, 155
  - research needs, 157
  - studies, 157
- surgical site infections (SSI)
  - antimicrobial prophylaxis and, 221
  - blood flow and body temperature, 231
  - glucose control for diabetics and, 237
  - normothermia and, 232
  - oxygen and, 234
  - research needs, 223, 233
  - studies, 222, 224, 226
- surgical sponges and instruments, retained, 255
- teicoplanin and central venous catheters, 179
- teleradiology, 376
- thromboembolism
  - heparin and warfarin monitoring, 88
  - preventions, 333
- To Err is Human: Building a Safer Health System*, 13, 25
- tobramycin, topical, 193
- total parental nutrition (TPN), 359, 361
- training
  - end-of-life care, for physicians, 560
  - laparoscopic cholecystectomy, 213
  - nurses, 424
  - radiograph interpretations, 377
  - simulator-based, 511
- transesophageal echocardiography, 265
- transfusion errors, 489
- transport teams, 534
- transportation of critically ill patients
  - about, 534
  - interhospital, 534
  - intra-hospital transport, 534
  - manual vs. mechanical ventilation, studies, 541
  - transport teams vs. standard, studies, 540
  - transporting children, 535, 537

transurethral resection of the prostate, 157  
 trauma patients, VTE risk, 338  
 UCSF-Stanford Evidence-based Practice Center (EPC), 13  
 ultrasound guidance of central vein catheterization  
   described, 245  
   insertion success rates, 246  
   research needs, 249  
   studies, 246, 247, 250  
 unions and nurse staffing, 425  
 unit-dose dispensing  
   automated dispensing systems, 112  
   described, 101  
   dispensing variations, 101  
   JCAHO standards, 102  
   research needs, 105  
   safety targets, 102  
   studies, 103, 104, 106  
 urethral catheters  
   compared to suprapubic, 159  
   latex, 149  
   patient discomfort, 156  
   research needs, 157  
   silicone, 149  
 urethral strictures, 157  
 urinary catheters  
   impregnated with antibiotic, 149  
   silver alloy catheters, 149  
   studies, 150, 153  
 urinary tract infections (UTIs)  
   catheters and, 149  
   prevalence, 149  
   research needs, 151  
 usability testing, 461  
 vancomycin, systemic use with central venous catheters,  
   179  
 vancomycin-resistant enterococci (VRE)  
   antibiotic studies, 142, 143  
   antibiotics and, 141  
   barrier precaution studies, 129  
   barrier precautions and, 127  
   *C. difficile* relationship, 128  
   research needs, 131, 144  
   transferability, 127  
 venography, 334  
 venous thromboembolism (VTE)  
   described, 333  
   preventative measures, mechanical and pharmacologic,  
     343  
   preventions, 333  
   prophylaxis for surgical procedures and medical  
     conditions, 345  
   risk in general surgery, 335  
   risk in hospitalized medical patients, 339  
   risk in neurosurgery, 337  
   risk in orthopedic surgery, 336  
   risk reduction methods, 344  
   studies, 334  
   therapy factors, 337  
   urethral catheters and, 156  
 ventilation when transporting critically ill patients, 538,  
   541  
 ventilator-associated pneumonia (VAP)  
   continuous aspiration of subglottic secretions, 190  
   continuous oscillation, 185, 187  
   described, 185  
   patient positioning studies, 189  
   research needs, 195, 199, 203  
   SDD, 193  
   semi-recumbent positioning, 185, 186  
   sucralfate, 198  
 Veterans Health Administration (VHA), 449  
 vomiting due to PCA, 404  
 ward stock system, 102  
 warfarin  
   about, 87  
   dosing and monitoring errors, reducing, 87  
   monitoring, 88  
   self-management, studies, 89, 96  
   venous thromboembolism prevention, 333  
 warfarin nomograms, studies, 92  
 warming blankets for normothermia during surgery, 231  
 web-based health information, 16, 292, 451, 561, 570  
 web-based programs, 478  
 wrong-site surgery, avoiding  
   described, 494  
   research needs, 498  
   studies, 496, 497