7/1/03 This manual is being completely revised. The drugs listed in SEER Book 8 may not include any antineoplastic agents developed since 1993. Please check our web site under **New Releases** at: http://SEER.CANCER.GOV for the new edition of this database containing a complete and current list of cancer drugs.

SEER Program

Self Instructional Manual for Tumor Registrars

Book Eight

Third Edition



NATIONAL INSTITUTES OF HEALTH National Cancer Institute

SELF-INSTRUCTIONAL MANUAL FOR TUMOR REGISTRARS

Book 8 - ANTINEOPLASTIC DRUGS

Third Edition

Prepared by

Evelyn M. Shambaugh, M.A., CTR
Cancer Statistics Branch
Division of Cancer Prevention and Control
National Cancer Institute

Susan G. Nayfield, M.D., M.Sc.
Community Oncology and Rehabilitation Branch
Division of Cancer Prevention and Control
National Cancer Institute

Terry M. Swenson, M.T. (ASCP) Information Management Services, Inc. (IMS)

> Mary A. Kruse Bethesda, Maryland

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
National Institutes of Health

NIH Publication No. 94-2441

ACKNOWLEDGEMENTS

We are grateful to the following persons who have contributed to and critiqued this publication and to the pharmaceutical companies that responded to our requests for information. Each has given us unending encouragement in the compilation of this list of antineoplastic drugs. Without their assistance, this publication would not have been possible.

Helen F. Lever, Assistant, Drug Information, Drug Management and Authorization Section, Investigational Drug Branch, National Cancer Institute

John F. Waters, B.A., Information Technology Branch, National Cancer Institute

Abbott Pharmaceuticals, Inc. Chicago, Illinois

Adria Laboratories, Columbus, Ohio

Bristol-Myers Squibb Company, Evansville, Indiana

Burroughs Wellcome Company, Research Triangle Park, North Carolina

Calbiochem-Behring Corporation, La Jolla, California

Ciba Pharmaceutical Company, Summit, New Jersey

Herbert Laboratories, Irvine, California

Lederle Laboratories, Pearl River, New York

Lilly Research Laboratories, Indianapolis, Indiana

Mead Johnson Pharmaceutical Division, Evansville, Indiana

Miles Pharmaceutical Division, West Haven, Connecticut

Roche Laboratories, Nutley, New Jersey

Schering Corporation, Kenilworth, New Jersey

Searle Laboratories, Chicago, Illinois

Smith, Kline & Beecham Laboratories, Philadelphia, Pennsylvania

The Upjohn Company, Kalamazoo, Michigan

BOOK 8: ANTINEOPLASTIC DRUGS

CONTENTS

| | Page |
|------|---|
| INTF | RODUCTION |
| A. | Objectives and Content of Instructional Book 8 |
| B. | Section 1. Chemotherapeutic Drugs and Combination Regimens (Introduction) 5 |
| | Chemotherapeutic Drugs |
| | Combination Regimens |
| C. | Section 2. Ancillary Drugs (Introduction) |
| | Ancillary Drugs |
| D. | Section 3. Differentiation-Inducing Agents (Introduction) |
| | Differentiation-Inducing Agents |
| | Agents with both Cytotoxic and Differentiating Properties |
| E. | Section 4. Biologic Therapy of Cancer (Introduction) |
| | Biological Response Modifiers |
| F. | Section 5. Hormones and Agents Acting Via Hormonal Mechanisms (Introduction) 69 |
| | Hormones or Agents Acting Via Hormonal Mechanisms |
| G. | Section 6. Drugs Used in the Treatment of AIDS and Its Complications (Introduction) 85 |
| | Drugs Used To Treat HIV Infection |
| | Drugs Used To Treat Opportunistic Infections and Other Complications Associated With AIDS |
| H. | Protocol Investigators |
| I. | Selected Bibliography |
| ĭ | Index 100 |

Introduction

OBJECTIVES AND CONTENT OF INSTRUCTIONAL BOOK 8

Book 8 was first published in 1981 in response to numerous requests from SEER Program participants for identification of drugs which should be included on the abstract of the medical record as cancer-directed therapy. The demand for this manual has been rewarding, and, therefore, this update is being offered in 1993.

Many new anti-cancer agents have been developed since the original publication of this book, and these have been added to the appropriate sections of the new edition. Experimental drugs proven to be ineffective are listed with a '#' sign following the drug name. Brand names formerly used and drugs not in clinical use at this time are also designated with a '#' sign. The index will assist you in locating a particular drug.

This publication is divided into six sections:

- 1. Chemotherapeutic drugs and combination regimens
- 2. Ancillary drugs
- 3. Differentiation-inducing agents
- 4. Biological response modifiers
- 5. Hormones and agents acting via hormonal mechanisms
- 6. Drugs used in the treatment of AIDS and its complications

Chemical formulae and information about drug metabolism and mechanisms of action are not included in this manual. If you find it necessary to know the drug's formula or pharmacology, please refer to the following resources:

- 1. Fleeger CA, ed. *USAN and the USP Dictionary of Drug Names*. Rockville, MD, United States Pharmacopeial Convention, Inc., 1992.
- 2. Drug Information for the Health Care Professional. USP-DI (12th Edition), Volumes IA and IB. Rockville, MD, United States Pharmacopeial Convention, Inc., 1992.
- 3. Chabner BA, Collins JA. Cancer Chemotherapy: Principles and Practice. JB Lippincott Company, New York, 1990.
- 4. NCI Investigational Drugs Pharmaceutical Data. NIH Publication No. 91-2141. Bethesda, MD, US Department of Health and Human Services, 1990.
- 5. Physicians Desk Reference, Montvale, New Jersey, Medical Economics Data, 1993.

In October 1982, the National Cancer Institute (NCI) announced the establishment of PDQ (Physicians Data Query). PDQ is a clinically oriented computer database developed to make recent cancer information widely available to the medical community. PDQ provides state-of-the-art cancer treatment information and descriptions of NCI-sponsored clinical trials for all major cancer sites. For clinical trials involving antineoplastic agents, protocol summaries can be retrieved using a drug's generic name, acronym, short name, or synonym.

Information on both established drugs and investigational agents can be obtained through PDQ from CANCERLIT searches on the National Library of Medicine (NLM) computer system. CANCERLIT is a bibliographic database that contains approximately 700,000 citations, including journal articles, meeting papers, books, reports, and doctoral theses. It is updated each month with close to 5,500 new citations, most containing abstracts.

Information about access to PDQ can be obtained from the PDQ Information Coordinator, National Cancer Institute, R. A. Bloch International Cancer Information Center, Building 82, Room 105, 9030 Old Georgetown Road, Bethesda, MD 20892. Instructions for accessing CANCERLIT through PDQ are included in the PDQ User's Guide.

Continuous updating is required to make this manual a useful reference. Therefore, suggestions with respect to changes or additions are welcome. With your help, this book can be an up-to-date directory of antineoplastic agents that you may encounter in abstracting a medical record.

Cancer Statistics Branch
Surveillance Program
Division of Cancer Prevention and Control
National Cancer Institute
Executive Plaza North, Room 343J
Bethesda, MD 20892

Telephone: (301) 496-8510

SYMBOLS USED IN THIS MANUAL

- NSC number, a National Service Center number from the National Cancer Institute (NCI). This number is assigned to a drug during its investigational phase, prior to the adoption of a United States Adopted Name (USAN).
- * Published in the Federal Register as a Federal Drug Administration (FDA) registered name.
- # Experimental agent proven to be ineffective, brand name formerly used, or drug thought not to be in clinical use at this time.

<u>Underlined</u> <u>Names</u>

USAN official names (usually of the drug substance, as distinct from the dosage forms), as used in current editions of the *United States Pharmacopeia* and the *National Formulary*.



Section 1

CHEMOTHERAPEUTIC DRUGS AND COMBINATION REGIMENS

Chemotherapy is a relatively new form of cancer treatment. Prior to the 1940's, no drugs were known to be effective against cancer. However, during that decade, observations of the effects of nitrogen mustard gas on the lymphoid system of seamen exposed during World War II resulted in its use in patients with Hodgkin's disease and other lymphomas and led to the development of other alkylating agents. Similarly, studies of the effect of folic acid on the growth of leukemic cells in children with acute lymphoblastic leukemia led to the development of antifolate agents in the early 1950's. This new approach to cancer therapy made it possible to treat disease which had spread beyond its site of origin.

Development and Testing of New Chemotherapeutic Agents

Scientists work continuously to develop new antineoplastic agents and to explore different methods for using agents which are currently available. New drugs undergo rigorous preclinical evaluations to determine antineoplastic effects in cancer cell cultures and animal tumor models. Drugs which demonstrate potential in preclinical studies then must be formulated for human use and go through toxicology testing in laboratory animals before entering systematic clinical evaluation.

Clinical studies of chemotherapeutic agents are conducted according to strict guidelines, or protocols, which provide specific instructions for patient selection, drug administration, and observation of treatment effects. There are three phases to clinical evaluation:

Phase I clinical trials assess the pharmacology and dose-limiting toxicity of the drug in humans. Patients in Phase I clinical trials generally have advanced malignancies which are no longer amenable to conventional cancer therapy.

Phase II studies determine the activity, or therapeutic effect, of a new drug in specific tumor types. Patients in Phase II clinical trials have tumors that are no longer amenable to conventional therapy; their tumor masses must be measurable on clinical examination or by radiographic studies to evaluate tumor response to the new drug. Response rates, duration of response, and frequency and severity of side effects are recorded in Phase II clinical trials.

Phase III clinical trials compare the efficacy of a standard drug (or treatment) to that of a new drug (or treatment) which is expected to be at least as effective as the standard therapy for a specific cancer stage and primary site. In most Phase III studies, patients are randomly assigned to treatment groups, and the response rate and toxicity for the new therapy is compared to that for the standard therapy ("control"). Phase III studies establish the place of a new drug or treatment in clinical oncology practice.

Clinical studies, especially Phase III clinical trials, require large numbers of patients with specific tumor types and stages. To complete these studies in a timely and efficient manner, investigators from different practice locations have joined together as cooperative clinical trials groups. Through this mechanism, patients from diverse geographical areas can enter a clinical study under the care of

their own oncology specialist. Each patient receives therapy according to a defined protocol, and data on response and toxicity for each patient are collected by a central office for monitoring and analysis.

Many cooperative groups evaluate surgical procedures or radiation therapy treatments, often combined with chemotherapy. Some cooperative groups conduct Phase I-II studies as well as Phase III randomized controlled clinical trials. Examples of cooperative clinical trials groups include the Cancer and Leukemia Group B (CALGB), the Southwest Oncology Group (SWOG), the Eastern Cooperative Oncology Group (ECOG), the National Surgical Adjuvant Breast and Bowel Project (NSABP), and the Radiation Therapy Oncology Group (RTOG). See page 95 for a list of the cooperative clinical trials groups.

How Chemotherapy Works

Chemotherapeutic agents inhibit the reproduction of cancer cells by interfering with DNA synthesis and mitosis, causing the cells to stop growing (cytostatic agents) or to die (cytotoxic agents). Cancer cells that are synthesizing DNA and rapidly dividing are more vulnerable to the effects of many drugs than are those in a resting (non-cycling) phase.

In systemic chemotherapy, drugs administered by injection or orally are carried throughout the body in the blood stream. Most drugs will reach cells in all areas of the body. However, certain "sanctuary" sites (such as the testes, eyes, and central nervous system) are protected by physiological barriers. For example, some drugs will not cross the blood-brain barrier and so must be administered directly into the cerebrospinal fluid (CSF) surrounding the brain and spinal cord (intrathecal administration). Carmustine (BCNU) and high-dose cytarabine (Ara-C) are transported across the blood-brain barrier.

In general, chemotherapeutic agents are classified into groups according to their mechanism of action. They may also be classified according to the timing of action in cell division: phase nonspecific agents act throughout the mitotic cycle, while phase-specific drugs act at defined points in the cell division process.

Alkylating agents cause cross-linking and abnormal base pairing that interferes with DNA replication. DNA strands may also break, and synthesis of enzymes and nucleic acids may be inhibited. Most alkylating agents are cell cycle nonspecific and affect both resting and dividing cells. However, rapidly dividing cells are most sensitive to their effects. Because normal cells in the bone marrow and gastrointestinal tract divide rapidly, bone marrow suppression and gastrointestinal disturbances are common toxicities associated with these drugs.

Alkylating agents include:

- 1. Nitrogen mustard and its derivatives: mechlorethamine (Mustargen), phenylalanine mustard (Melphalan), chlorambucil (Leukeran), cyclophosphamide (Cytoxan)
- 2. Ethylenimine derivatives: triethylene-thiophosphoramide (Thio-TEPA)
- 3. Alkyl sulfonates: busulfan (Myleran)

4. Nitrosoureas: carmustine (Lomustine)

5. Triazines: DTIC (Dacarbazine)

Antimetabolites are structurally similar to natural metabolites which are necessary for cell function. These agents replace natural metabolites in important molecules, altering the function of enzymes required for cell metabolism and protein synthesis. This interference is most pronounced during the S (synthesis) phase of cell division when DNA, RNA, and protein synthesis occurs. The enzymatic block produced by methotrexate can be bypassed by administration of folinic acid (leucovorin), which then "rescues" normal cells from methotrexate action. Antimetabolites include:

1. Folic acid analogues: methotrexate (Amethopterin, MTX)

2. Pyrimidine analogues: 5-fluorouracil (5-FU)

3. Purine analogues: 6-mercaptopurine (6-MP)

Natural products include antibiotics, plant alkaloids, and enzymes. Antitumor antibiotics may have antimicrobial properties, but their cytotoxic properties are most prominent. These agents prevent nucleic acid synthesis by a process called intercalation and block DNA and RNA transcription. They act throughout the cell cycle, but some are more effective during S (synthesis) and M (mitosis) phases. Antitumor antibiotics include dactinomycin (Actinomycin D), doxorubicin (Adriamycin), daunorubicin (Daunomycin), bleomycin (Blenoxane), and mitomycin C (Mutamycin).

Plant alkaloids, derived from the periwinkle plant *Vinca rosea*, are also called vinca alkaloids. These drugs interfere with the microtubular spindle proteins necessary for cell division and thus inhibit mitosis (M phase). Vinblastine (Velban, VBL) and vincristine (Oncovin, VCR) are the two vinca alkaloids commonly used in clinical oncology.

L-asparaginase (Elspar) is an **enzyme** which catalyzes the breakdown of asparagine into aspartic acid and ammonia. It inhibits the growth of tumor cells that are unable to synthesize l-asparagine, an amino acid necessary for protein synthesis. Cells that make l-asparagine from aspartic acid are not affected.

Several anticancer drugs are classified as miscellaneous agents, either because they do not fall into one of the distinct categories above or because their action is not fully understood. Cis-diammine dichloroplatinum II (Cisplatin) is a platinum-containing complex which binds to DNA and disrupts its function. Hydroxyurea (Hydrea) inhibits enzymes which are important in repair of DNA damage. Procarbazine (Matulane) is broken down by the body into active metabolites which bind nuclear DNA and inhibit DNA, RNA, and protein synthesis.

Combination Chemotherapy Regimen

Few chemotherapeutic drugs in general use today are administered as single agents. Combinations of chemotherapeutic agents are carefully selected, taking advantage of different mechanisms of drug action to increase tumor response rates, to prevent or delay the development of drug resistance, and to modulate side effects of therapy.

A treatment cycle is a round of a chemotherapy regimen administered according to a specific schedule. Treatment cycles for most regimens are 21 or 28 days. The first day of the cycle is Day 1, the second is Day 2, and so forth. The treatment protocol specifies each drug, dose and method of administration, and schedule (day of cycle). The acronym for the regimen is usually formed from the first letters of the chemotherapeutic agents used in the regimen. Thus MOPP, for the treatment of Hodgkin's disease, is a combination of nitrogen Mustard, vincristine (Oncovin), Procarbazine, and Prednisone. See page 29 for a list of commonly-used combination regimens.

A cycle of the MOPP regimen is shown below. Nitrogen mustard and vincristine are administered intravenously on Day 1 and Day 8, usually in the outpatient setting. In addition, procarbazine and prednisone are taken orally for the first 14 days of the cycle. Note that no chemotherapy is administered on Days 15 through 28; this rest period allows recovery from any side effects that occur. The regimen is repeated every 28 days for a minimum of six cycles, or until complete remission is achieved plus two additional cycles.

While some treatment plans require a specific number of chemotherapy cycles, in other cases, treatment continues until the patient's disease is no longer controlled by the chemotherapeutic regimen.

| MOPP Regimen for Hodgkin's Disease (28-day Cycle) | | | |
|---|-----------------------------------|-------------------|--|
| Drug | Dose* | Schedule | |
| M - Nitrogen mustard | 6.0 mg/m ² IV | Days 1 and 8 | |
| O - Vincristine (Oncovin) | $1.4 \text{ mg/m}^2 \text{ IV}$ | Days 1 and 8 | |
| P - Procarbazine | $100.0 \text{ mg/m}^2 \text{ PO}$ | Days 1 through 14 | |
| P - Prednisone | 40.0 mg/m ² PO | Days 1 through 14 | |
| | | | |

^{*} Dose is usually calculated as milligrams (mg) drug per square meter (m²) body surface area. Body surface area is determined from charts depicting surface area as a function of the patient's height and weight.

When disease is widespread, as in the acute leukemias, intensive chemotherapy may be administered in an attempt to eradicate all evidence of disease. This period of treatment is called **induction**; the absence of clinically demonstrable disease is termed **remission**. Following induction, a second period of intensive treatment (**consolidation**) may be given to increase the remission rate or increase duration of remission. In some clinical situations, a prolonged period of less intense chemotherapy, called **maintenance therapy**, is also administered to delay disease recurrence (relapse).

In certain cancers such as Hodgkin's disease, re-treatment of patients who relapse following a prolonged disease-free interval may be effective. This attempt to achieve another remission is often called salvage therapy.

Administration of Chemotherapy

The most common routes of administration of chemotherapy are oral and intravenous. The drug enters the bloodstream by absorption from the gastrointestinal tract or directly by intravenous injection and is carried to cells throughout the body. However, in specific clinical situations, other methods of administration are required to achieve higher drug concentrations in specific tissues or to access tissues that are isolated by physiologic barriers.

Drugs that do not cross the blood-brain barrier must be administered directly into the cerebrospinal fluid (CSF) surrounding the brain and spinal cord (intrathecal administration). This is accomplished through a lumbar puncture (spinal tap) needle, into an implanted access device (Ommaya reservoir). Intrathecal chemotherapy is used to treat meningeal leukemia or lymphoma or occasionally in other tumors which have meningeal metastases. In patients with acute lymphoblastic leukemia, intrathecal chemotherapy may be given as prophylaxis to prevent the development of meningeal leukemia. Methotrexate and cytosine arabinoside are chemotherapeutic agents commonly used intrathecally.

Chemotherapeutic agents such as bleomycin may be injected directly into the pleural or pericardial space to control malignant effusions.

Intraperitoneal administration of drugs may be used to treat tumors such as ovarian cancer which spread by direct extension through the abdominal cavity. Chemotherapy is injected into the peritoneal cavity using techniques similar to peritoneal dialysis. This approach allows increased concentrations of drug in the abdominal cavity compared to lower levels achieved by intravenous administration.

Specialized infusion methods have been designed to increase drug delivery to affected tissues while sparing normal or uninvolved tissues. These require insertion of a catheter into the artery which supplies blood to the affected area and infusion of the drug through the catheter. Hepatic artery infusion has been used to treat primary liver tumors or liver metastases, and carotid artery infusion has been attempted to treat head and neck cancers or brain tumors. While advances in technology have simplified these infusion procedures, they are still investigational. Their role needs to be determined by careful scientific evaluation through clinical trials.

Monitoring Side Effects

Antineoplastic drugs affect not only cancer cells but also normal cells. For chemotherapeutic drugs, the type and severity of side effects depend on the specific agent, its dose, and its route of administration. Side effects are usually temporary and respond to supportive treatment. However, if toxicity is severe or persistent, drug dosages may be reduced for subsequent cycles to decrease

toxicity, a new drug with similar actions but different side effects may be substituted for the drug causing symptoms, or scheduled therapy may be delayed to allow recovery. Most protocols include specific directions for dose modification due to toxicity.

In patients with chronic kidney or liver disease, metabolism and/or excretion of specific chemotherapeutic agents may be impaired. This results in prolonged exposure of normal tissues to the drugs and their metabolites and often causes an increase in frequency and severity of side effects. Drug doses may be decreased according to protocol guidelines to prevent toxicity in patients who have these medical conditions.

The bone marrow is the site of production of circulating blood cells: red blood cells (RBCs), which contain hemoglobin and transport oxygen to the tissues; white blood cells (WBCs), which fight infection and function in immunity; and platelets, which prevent bleeding when small blood vessels are injured. Bone marrow cells divide rapidly and are very sensitive to most antineoplastic agents, so that hematologic side effects are common in patients receiving these drugs. Toxicity may be expressed as decreases in one or more of the circulating blood cell types.

Most patients receiving chemotherapy develop a decrease in RBCs (anemia), reflected by a fall in laboratory measurements of hemoglobin or hematocrit. Anemia may be associated with generalized symptoms of fatigue and weakness, or it may exacerbate cardiovascular symptoms of shortness of breath (dyspnea) or congestive heart failure. Blood (RBC) transfusions are given if anemia becomes severe or symptomatic.

A decrease in platelet count (thrombocytopenia) may result in the appearance of small redpurple hemorraghic spots (petechiae), particularly on the skin of the lower legs and on the oral mucosa, large bruises (ecchymoses), or blood in the urine (hematuria). More serious internal hemorrhages may develop without apparent cause in patients with very low platelet counts. Platelet transfusions are usually given to patients with severe thrombocytopenia to prevent significant bleeding.

A decrease in the number of white cells (leukopenia), particularly in neutrophils which fight bacteria (neutropenia), makes the patient more susceptible to bacterial and fungal infections. WBC transfusions may be used in patients with life-threatening infections which are unresponsive to antibiotics.

The term pancytopenia is used when all blood cell lines are decreased.

Hematologic toxicity associated with chemotherapy for most cancers usually lasts seven to ten days at the most. However, induction therapy for acute leukemias, intensive chemotherapy for other widespread malignancies, and bone marrow transplantation often result in prolonged periods of pancytopenia. Patients undergoing these treatments require intravenous antibiotics and frequent transfusions of RBCs and platelets for hematologic support until bone marrow recovery occurs.

Gastrointestinal symptoms due to antineoplastic drugs may include nausea, vomiting, diarrhea, mouth ulcers (stomatitis), and irritation of the esophagus (esophagitis) or stomach lining (gastritis). Non-chemotherapeutic medications can be very effective in controlling these side effects. Gastrointestinal toxicity is usually short-lived (one to two days), but, in some situations, may be prolonged (five to six days).

Rapidly-dividing cells at the base of the hair follicles are also sensitive to chemotherapy. Thinning or loss of body hair (alopecia) occurs in many patients receiving chemotherapy. This is usually most pronounced on the scalp, but beard, axillary, pubic, and leg hair may also be affected. Hair growth resumes when chemotherapy is discontinued.

In both men and women, cells responsible for reproductive function may be affected by chemotherapy. In premenopausal women, menstrual cycles may become irregular or stop completely, and hot flashes may occur. Men may experience a decrease in the number of spermatozoa (oligospermia). Patients who complete chemotherapy may produce normal children; however, fertility may be reduced in long-term cancer survivors who have received chemotherapy.

Inflammation of the urinary bladder with bleeding into the urine (hemorrhagic cystitis) may occur with chemotherapeutic agents which are concentrated in the urine. Cyclophosphamide and ifosphamide are examples of drugs which may cause hemorrhagic cystitis. This complication may be prevented by increased fluid intake and/or by ancillary drugs which protect the bladder epithelium. Renal toxicity, manifested by increases in serum creatinine and blood urea nitrogen (BUN) and electrolyte imbalance, may be associated with cisplatin therapy.

Cardiovascular side effects are limited to specific agents. Antitumor antibiotics with anthracycline chemical structures, doxorubicin (Adriamycin) and daunomycin (Daunorubicin), can damage myocardial cells. Because congestive heart failure occurs more frequently in patients who receive high total doses of drug, a total dose limit is required in regimens containing these agents.

Pulmonary toxicity has been described in patients who receive long-term therapy with alkylating agents (busulfan, cyclophosphamide, and BCNU) and antitumor antibiotics (bleomycin and mitomycin). The development of pulmonary fibrosis may present as shortness of breath and has clinical findings of diffuse changes on chest x-ray and decreased oxygen diffusion on pulmonary function studies.

Fatigue and/or muscle weakness may result from anemia or from direct effects of the drugs on muscles or nerves. Patients receiving corticosteriods such as prednisone for prolonged periods may develop muscle weakness (myopathy), particularly involving the muscles of the upper arms and legs. Certain chemotherapeutic agents, such as the vinca alkaloids, cisplatin, and taxol, are associated with damage to the peripheral nervous system. The loss of sensation in the hands and feet (peripheral neuropathy) which develops in many patients receiving these drugs often improves with time when the agent has been discontinued.

Dermatitis associated with chemotherapy is uncommon. Patients receiving 5-fluorouracil by continuous infusion may develop redness and peeling of the skin on palms and soles (Hand/Foot Syndrome). Exfoliative dermatitis may be a dose-limiting toxicity of the drug PALA.

As more cancer patients achieve cure or prolonged survival, attention has focused on late sequelae of chemotherapy and problems of cancer survivors. Second primary malignancies related to treatment, such as acute non-lymphocytic leukemia in patients treated with alkylating agents, may occur years after completion of therapy. Current treatment plans attempt to decrease the potential for late complications whenever possible. For example, alternating cycles of MOPP and ABVD chemotherapy in patients with Hodgkin's disease reduces the total dose of alkylating agents and may decrease the risk of late sequelae.

Monitoring Tumor Response

Assessing response to therapy is of major importance both in day-to-day oncology practice and in clinical research. Many cancer patients will have **measurable disease**, tumor masses which can be measured in two dimensions on physical examination or by radiographic studies such as chest x-ray or CT scan. Other patients will have disease that can be evaluated objectively by examination or studies but cannot be measured directly; this is termed **evaluable disease**. Mediastinal tumors, in which widening of the mediastinum on chest x-ray reflects the extent of tumor involvement but distinct margins of the lesion cannot be determined clearly, are an example of evaluable disease. Some clinical trials require that participating patients have measurable disease, while others accept patients with evaluable disease.

A complete response (CR) occurs when the tumor mass resolves and there is no evidence of residual disease. A partial response (PR) shows definite improvement but disease remains present; specific criteria for partial response may vary from protocol to protocol. Other response categories of stable disease and progressive disease may be used to describe patients who do not achieve complete or partial responses. Response rates and duration of response are used to evaluate the efficacy of a particular regimen in a defined patient population.

The impact of a treatment is also evaluated by **overall survival** of treated patients and, for situations in which all clinically apparent disease is eradicated with treatment, by **disease-free survival** (time until relapse).

Recently, attention has been directed to the impact of cancer treatment, especially chemotherapy, on a patient's health-related quality of life (QOL). This construct represents the patient's physical and mental well-being. For the cancer patient, QOL includes physical performance, emotional and psychologic function, social interactions, and symptoms of disease. QOL is now measured as an outcome in many cancer treatment clinical trials. It is an especially important consideration when treatments have similar response rates and survival but different toxicities, or when one treatment has improved survival but has more severe side effects.

Multimodality Approach to Cancer Therapy

Chemotherapy may be administered with other treatment modalities such as surgery and radiation therapy in an organized treatment plan. This approach is called **multimodality therapy** or **combined modality therapy**. Administration of chemotherapy may precede or follow surgery or radiotherapy; in some cases, chemotherapy or hormonal therapy may be administered during radiation therapy. The use of chemotherapy or hormonal therapy immediately following surgery for early stage breast cancer, all as a part of the initial course of treatment, is an example of combined modality therapy.

Many patients who appear to have localized cancers at diagnosis will develop recurrence at sites distant from that of the original surgery. These patterns of recurrent disease suggest that micrometastases were present at diagnosis but could not be detected clinically. In early stage breast cancer and colon cancer, administration of chemotherapy following initial surgery can prolong disease-free survival and overall survival. Chemotherapy administered following surgical resection in patients without clinically demonstrable disease is called adjuvant chemotherapy.

In certain situations, chemotherapy has been used to reduce the bulk of a locally advanced primary cancer prior to surgical resection or radiotherapy. This approach is called **neoadjuvant** (or primary) chemotherapy. The role of neoadjuvant chemotherapy in management of head and neck carcinomas, lung cancer, esophageal malignancies, and locally advanced breast cancer is currently under investigation.

Progress in Cancer Treatment with Chemotherapy

Although chemotherapy is a relatively new modality in cancer treatment, antineoplastic agents are used at some time during treatment in most patients with cancer. The development of effective combination chemotherapy regimens has made cure possible in many patients with advanced disease and has delayed tumor recurrence and prolonged survival in many patients with early stage cancer.

The following table summarizes advances in cure of advanced cancer with chemotherapy in both adult and pediatric malignancies.

| Curability of Disseminated Cancer with Chemotherapy | | | | |
|--|------------------------------|-----------------|--|--|
| Disease Therapy Cure Rate | | | | |
| Adults | | · | | |
| Diffuse histiocytic lymphoma (Stages III and IV) | CTX | <u>></u> 50% | | |
| Hodgkin's disease (Stages III and IV) | СТХ | <u>≥</u> 50% | | |
| Testicular carcinoma (Stage III) | CTX + SURG | <u>></u> 75% | | |
| Gestational choriocarcinoma | Methotrexate ± actinomycin D | 90% | | |
| Ovarian carcinoma | Alkylating agents or CTX | 10 - 20% | | |
| Acute myelocytic leukemia | CTX | 20% | | |
| Children | | | | |
| Acute lymphocytic leukemia Non-Hodgkin's lymphoma | CTX + cranial XRT | <u>≥</u> 50% | | |
| Burkitt's lymphoma | Cyclophosphamide or CTX | <u>≥</u> 50% | | |
| Wilms's tumor Childhood sarcomas | SURG, CTX, XRT | ≥ 50% | | |
| CTX=Combination chemotherapy; SURG=Surge | ery; XRT=Radiation therapy | | | |
| Adapted from: Chabner BA, Collins JA. <u>Cancer Chemotherapy: Principles and Practice</u> . JB Lippincott Company, New York, 1990. | | | | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|----------------------------|--------------------|---|-------------------|
| | | | |
| Acivicin | | Antibiotic AT-125, U-42,126 | 163501 |
| Aclarubicin | ACM, ACR, ACM-A | Aclacinomycin, Aclacinomycin-A#, Aclacinon, Aclarubein, Antibiotic MA, 144 A1 (anthracycline) | 208734 |
| Acodazole (HCL) | AD-32 | EU-3120 | 305884 |
| Acronine* | | Acronycin#, Acronycine, Compound 42339 | 403169 |
| Adozelesin | | U-73,975 | |
| Alanosine | | L-Alanosine (antitumor antibiotic) | 153353 |
| Alpha-TGdR | | Deoxythioguanosine hydrate, Thioguanosine deoxyriboside | 071851 |
| Altretamine* | нмм | ENT 50852, Hemel, Hexalen (see Hexamethylmelamine) | 013875 |
| <u>Ambomycin</u> | | Antitumor antibiotic | 053397 |
| Ametantrone Acetate | CI-881 | 287513 | |
| Aminopterin | APGA | Aminopteridine, Aminopterin sodium, A-Ninopterin, ENT 26079 (antimetabolite) | 000739 |
| Aminothiadiazole | ATDA, A-TD | TF-128, 2-Amino-1,3,4-thiadiazole | 004728 |
| Amsacrine | AMSA, m-AMSA | Acridinyl anisidide, Amsidyl, CI-880 (miscellaneous agent) | 249992 |
| Anguidine | ANG, DAS | ANG 66, Anguidin, Diacetoxyscirpenol | 141537 |
| Aniline mustard | | Lymphochin, Lymphocin, Lymphoquin, TL 476 | 018429 |
| Anthramycin | | Antramycin, 2-Propenamide | |
| Anthrapyrazole C 19 | 41 | | |
| Aphidicolin Glycina | te | Aphidicolin-17 glycinate, ICI 137233 | 303812 |
| Asaley | | Asalex | 167780 |
| Asperlin# | | U-13,933 | 093158 |
| 5-Aza-2'Deoxy- cytidine | DAC | | |
| 8-Azaguanine | Azan 8-AG, AZG | Azaguanine, B-28, Guanazol, Guanazolo, Pathocidin, Pathocidine, | 000749 |
| Azapicyl | | SF-337 | |
| Azaribine* | AZR | 6-Azauridine triacetate, CB 304, Triazure# | 067239 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC Mo</u> . |
|--------------------|-----------------------|---|-------------------|
| Azaserine*# | AZAS, AZS | Azaserin, CI 337, CN 15,757, Diazoacetylserine, L-Serine diazo- acetate ester, P-165 (antimetabolite) | 000742 |
| Azathioprine* | 6 ATP, AZAT | Azanin, Azathioprin, Azatioprin, Azothioprine, BW 57-322, Imuran, Imurek, Imurel, Muran | 039084 |
| Azetepa# | | Azatepa, CL-25477, Thiatriamide | 064826 |
| Azidothymidine | AZT | 3'-Azido-3' Deoxythymidine, BW A 509U, Retrovir, <u>Zidovudine</u> (used in treatment of Kaposi's sarcoma only) | 602670 |
| Azotomycin | AZOT | Duazomycin-B, Antibiotic 1719 | 056654 |
| 6-Azuridine | Azur | 6-Azauracil riboside, Azauridine, 6-Azur, 6-Azauridine, Ribo-azauracil, Ribo-azuracil, Riboazuracil | 032074 |
| Baker's antifol | BAF, TZT | Baker's antifolante, Ethanesulfonic- acid compound, Triazinate | 139105 |
| <u>Benzodepa</u> | | AB-103, Benzcarbimine, Dualar#, ENT-50451 | 037096 |
| Beta-TGdR | BTGR, TGdR | Thioguanine deoxyriboside, B-2' Deoxythioguanosine | 071261 |
| Bisantrene (HCL) | ADAH, ADC | Bisantrene, CL-216,942, Orange- Crush# (anthracene derivative) | 337766 |
| Bleomycin Sulfate | BLEO, BLE, BLM | Blenoxane, Bleomycin, NDC 0015-3010 (antitumor antibiotic) | 125066 |
| BMY-45622 | | | |
| Brequinar Sodium | | Dup 785 | 368390 |
| Bromacrylide# | | 2-Propenamide | 066248 |
| <u>Bropirimine</u> | | U-54,461 | |
| Bruceantin | BRU | | 165563 |
| Bryostatin | | | 339555 |
| Busulfan* | BSF, BU, BUS, Busu | Busulphan, Joacamine, Mablin, Misulban, Mitostan, Myelosan (Russia), Mylecytan, Myleran (alkylating agent) | 000750 |
| BW A 773U | | Crisnatol | |
| Cactinomycin | | Actinochrysin, Actinomycin C#, H.B.F. 386, Sanamicia, Sanamycin, Sandamycin, (antitumor antibiotic) | 018268 |
| Camptothecin | CAMP | Camptothecine sodium | 100880 |
| <u>Caracemide</u> | | | 253272 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|----------------------------|--|---|------------------|
| Carlossins | | N 477 | |
| Carbetimer | | N-137 | |
| <u>Carboplatin</u> * | CBDCA | JM-8, Paraplatin, | 241240 |
| Carmustine* | BCNU | BiCNU, Carmustinea, (nitrosourea) (alkylating agent) | 409962 |
| Carubicin (HCL) | СММ | Carminomicin I, Carminomycin, Carminomycin HCl#, Carminomycin I, Carubicin, Karminomitsin, Karminomycin HCl | 275649 180024 |
| CB-10-277 | | | 208107 |
| 2-CdA | | 2-Chloro-2'Deoxyadenosine, Cladribine, Leustatin | 105014 |
| <u>Chlorambucil</u> * | CHL, CLB | Ambochlorin, CB-1348, Chloraminophene, Chlorbutin, Chlorbutine, Ecloril, Leukeran, Leukersan, Leukoran, Linfilizin, Linfolysin (alkylating agent) | 003088 |
| Chlorozotocin | CLZ, CZT, DCNU | SRI 5244 (analog of Streptozocin) | 178248 |
| Chlorsulfa- quinoxaline | CQS | Chloroquinoxaline sulfonamide | 339004 |
| Chromomycin A3# | | Aburamycin B, Antibiotic B 599, Toyamicin, Toyomycin | 058514 |
| CI-921 | | | 343499 |
| CI-937 | | | |
| Cilastatin Sodium | | Component of Primaxin (antibiotic) | |
| <u>Cirolemycin</u> * | | U-12,241 | 077950 |
| <u>Cisplatin</u> * | CACP, CPPD, Cis- DDP, CPD, CPDC, CPDD, DDP | Cis-Diamminedichloroplatinum (II), Cis-Platin, Cisplatino (Spanish), Cis-Platinum II#, Cisplatyl, Neoplatin, Peyrone's chloride, Platinex, Platinol, Platinum (miscellaneous agent) | 119875 |
| Clomesone | | reaction (missectaneous agent) | 338947 |
| Colcemid | | Alkaloid H-3, C-12669, Colcemide, Colchamin, Colchicine, Deacetyl-N-methyl- colchicine, Demecolcine, Desacetyl- methylcolchicine, Desmecolcine, Kolchamin, Kolkamin, Omain, Omaine, Santavy's Substance F, Substance F (Reichstein's) | 003096 |
| CPT-11 | | | |
| <u>Crisnatol</u> mesylate | | BW A770U mesylate | |
| Cyclocytidine (HCl) | CYC, | Cyclo-C Ancitabine hydrochloride, Ancytabine, CycloCMP hydrochloride (miscellaneous agent) | 145668 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>MSC No</u> . |
|----------------------------|---|---|-------------------|
| | | | |
| Cyclodisone | | | 348948 |
| Cyclohexanecarboxami | de | Aziridinyl cyclohexane, Ba 24648 | 051915 |
| Cycloleucin | ACPC, CL, CYL | Cycloleucine, CB 1639, WR 14997 | 001026 |
| Cyclopentenyl- cytosine | Ce-Cyd, CPE-C | | 375575 |
| <u>Cyclophosphamide</u> * | CP, CPM, CTX, CTY, CYC, Cyclo, CYT, CYTOX, CYTX | Asta B 518, Clafen, Claphene, Cyclophosphamid, Cyclophosphan, Cyclophosphane, Cytophosphan, Cytoxan, Endoxan, Endoxana, Enduxan, Genoxal, Neosar, Procytox, Sendoxan, Tymtram (alkylating agent) | 026271 |
| Cytarabine* (HCL) | Ara-C, CA, HDA, HDARA-C | AC-1075, Alexan, Arabinocytidine, Ara-cytidine, Beta-Arabinosylcytosine, Cytarabin, Cytarabinoside, Cytosine arabinoside hydrochloride#, Spongocytidine HCL, U-19920A (antimetabolite) | 063878 |
| Cytembena | СТВ | MBBA, Mebryl (Czechoslovakia), sodium bromebrate | 104801 |
| cos | | See Chlorsulfaquinoxaline | 339004 |
| <u>Dacarbazine</u> * | DIC, DTIC, ICDT, | Dimethyl-triazeno-imidazole-carbox- amide, DTIC-Dome, (antimetabolite) | 045388 |
| <u>Dactinomycin</u> * | ACD, ACT, ACT-D, ACTD, Acto-D, AD, Dact | Actinomycin AIV, Actinomycin C1, Actinomycin D, Actinomycindioic D- acid dilactone, Actinomycin I1, Actinomycin IV, Cosmegan, Dactino- mycin D, Meractinomycin, Oncostatin K (antitumor antibiotic) | 003053 |
| Daumorubicin (HCL)* | Daun, Dauno, DNM, DNR, DRB | Acetyladriamycin, Cerubidine, Dauno- blastin, Daunomycin, Daunomycin-HCl, Daunorubicin, Daunorubicine, Leuk- aemomycin C, NDC-0082-4155, RP-13057, Rubidomycin-HCl, Rubomycin C (antitumor antibiotic) | 082151 |
| 3-Deazauridine | DAU | Deazauridine, EO-26 | 126849 |
| Decitabine | DAC | | 127716 |
| Deoxydoxorubicin HCl | DxDx, 4-DxDx | 4'-Deoxyadriamycin hydrochloride, 4'-Deoxydoxorubicin, Esorubicin HCl, 1-Dox, 1M1 58 | 267469 |
| Deoxyspergualin HCl | | 15-Deoxyspergualin trihydrochloride, Heptanamide | 356894 |
| Detorubicin | DTR | | |
| Dexormaplatin | | υ-78,938 | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|---------------------------|-------------------------|---|-------------------|
| | | | |
| Dexrazoxane | | ADR-529, ICRF-187 | 169780 |
| Dezaquanine | | CI-908 | 261726 |
| Dezaguanine Mesylat | <u>e</u> | C1-908 mesylate, PD 90,695-73 | |
| DFMO | DFMO (HCl) | Alpha-DFMO HCl, Alpha-Difluoromethylornithine, Difluoromethylornithine | 337250 |
| DHAC | | 5,6-Dihydro-5-azacytidine hydrochloride, Dihydroazacytidine | 264880 |
| Dianhydrodulcitol | DAG | Dianhydrogalactitol, Dulcitol- diepoxide, Galactitol (miscellaneous agent) | 132313 |
| <u>Diaziquone</u> | AZQ | Aziridinyl Benzoquinone, CI 904 (quinone derivative) | 182986 |
| Dibromodulcitol | DBD | Dibromdulcit, Elobromol, Mitolac, Mitolactol (miscellaneous agent) | 104800 |
| Dibromomannitol | DBM | Dibromannit, Mielobromol, Mitobron- itol, Myebrol, Myelobromol (alkylating agent) | 094100 |
| Dichloroallyl lawsone | DCL | Dichlorolapachol | 126771 |
| Dichlorometho- trexate | DCM, DCMTX | Dichloroamethopterin, DichloroMetho- trexate | 029630 |
| Didemnin B | | | 325319 |
| Diglycoaldehyde | INOX, STGDR | Inosine dialdehyde, Wy-5321 | 118994 |
| Dihydrolenperone | | Dihydro-lenperone | 343513 |
| DON | | 6-Diazo-5-Oxo-L-Norleucine (901) | 007365 |
| Doxorubicin (HCl)* | ADM, ADR, ADRI, Adri | Adriamycin, Adriamycin-TM, Adri- blastina (Italy), FT-106, 14-hydroxy- daunomycin, Rubex (antitumor antibiotic) | 123127 |
| Doxorubicin-DNA com | plex | decision, and cancerdate and brother, | |
| Duazomycin | | (N-Acetyl DON) | 051097 |
| Duborimycin | | | |
| Echinomycin | | Quinomycin A | 526417 |
| Edam-10 | | | 626715 |
| <u>Edatrexate</u> | | CG 30694 | |
| Eflornithine HCl | | DFMO, MDL 71,782A, Ornidyl | |
| Elsamitrucin | | BBM-2478A, BMY-28090 | |
| Emetine | EMET | Emetin, Emetine monohydrochloride | 033669 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|--------------------------|---------------------------|--|-------------------|
| | | | |
| Emofolin (sodium) | Метннғ | Homofolic acid, 5-Methyltetrahydro- homofolate disodium | 139490 |
| Enpromate# | | 59156 | 112682 |
| Epipropidine# | | Eponate, Epoxypropidine, LY 28002 | 056308 |
| Epirubicin (HCL) | EPI | IMI-28, Pharmorubicin | 259642 |
| Ethidium chloride | | Babidium chloride, Homidium chloride, Novidium chloride | 084423 |
| Ethoglucid | | Epodyl, Etoglucid, ICI-32865 | 080439 |
| 9-Ethyl-6-Mercaptop | urine | 9-Ethyl-6-MP, 79T61 | 014575 |
| Etoposide* | EPEG, VP-16, VP-16-213 | Epipodophyllotoxin ethylidene gluco- pyranoside, Ethylidine-Lignan-P, Podophyllotoxin derivative, Vepesid (plant alkaloid) | 141540 |
| Etoprine | | • | |
| Fadrazole HCl | | CGS-16949A | |
| <u>Fazarabine</u> | Ara-AC | 5-azocytosine arabinoside, Kymarabine | 281272 |
| <u>Fenretinide</u> | | McN-R-1967 | |
| flavone acetic acid | FAA, FVA | Flavone-8-Acetic Acid | 347512 |
| F3TDR | | Trifluorothymidine, Trifluridine | 075520 |
| <u>Floxuridine</u> * | FUDR, 5-FUDR, 5-FURD | <pre>Floxuridin, 5-Fluorouracil deoxyriboside, 5-Flurodeoxyuridine, FUDR (antimetabolite)</pre> | 027640 |
| Fludarabine Phosphate | 2-FAMP | Fludara, 2-Fluoroadenine arabinoside- 5-Phosphate, 2-Fluoro-ARA AMP | 312887 |
| 2-Fluoroadenosine# | 2 FAS | SRI-727 | 030605 |
| Fluorodopane | | Fluorodopan, Fluorpan | 073754 |
| Fluorouracil* | 5-FU, FU | Adrucil, Efudex, Fluoroplex, 5-Fluorouracil, Fluracil, 5-Fluracil, Fluril, Oracil, Ro 2-9757 (fluorinated pyrimidine) (antimetabolite) | 019893 |
| Flurocitabine# | | AAFC, Ro 21-0702 | |
| <u>Fosqui done</u> | | GR 63178K | |
| Fostriecin Sodium | | Antibiotic CI-920, Antibiotic CL 1565A, | 339638 |
| Fotemustine | | Pyranone Phosphate | |
| Ftorafur | | See <u>Tegafur</u> | 148958 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|-------------------------------------|----------------------|--|------------|
| Gemcitabine | | LY188011 | |
| Gemcitabine HCl | | LY188011 hydrochloride | |
| GR 63178A | | | |
| Guanazole | | 3,5-Diamino-s-triazole | 001895 |
| Hepsulfam | | 1,7-Heptanediylsulfamate (ester) | 329680 |
| Hexamethylmelamine | нмм, нхм | See <u>Altretamine</u> | 013875 |
| Homoharringtonine | нн | | 141633 |
| Hycanthone (mesylat | e) | Etrenol, Hycanthone methanesulfonate, Hycanthone monomethanesulfonate | 142982 |
| 5-Hydroxypico- linaldehyde thios | 5-HP emicarbazone | | 107392 |
| <u>Hydroxyurea</u> * | HU, HUR, HYD | Carbamohydroxamic acid, Carbamohydroximic acid, Carbamoyl oxime, Hidrix, Hydrea, Hydreia, Hydroxycarbamide, Hydroxycarbamide, Hydroxycarbamine, Hydroxylurea, Hydura, Litaler, Litalir, Onco-Carbide, Oxyurea, SQ 1089 (miscellaneous agent) | 032065 |
| ICRF-187 | | See <u>Dexrazoxane</u> | 169780 |
| <u>Idarubicin HCl</u> * | 4-DMDR, IDA | 4-Demethoxydaunorubicin, Idamycin, IMI-30 | 256439 |
| <u>Ifosfamide</u> * | IFX, IPP | A-4942, Asta Z-4942, Cyfos#, Holoxan 1000, <u>Ifex</u> , Ifosfamid, Iphosphamid, Iphosphamide, Iso- endoxan, Isofosfamide, Isophos- phamide, Mitoxana, MJF 9325, Naxamide#, Z-4942 (alkylating agent) | 109724 |
| Ilmofosine | | BM 41,440 | |
| <u>Imi penem</u> | | Imipemide#, component of Primaxin (antibiotic | :) |
| Indicine-N-oxide | INDI, INO | | 132319 |
| Inproquone | | E-39, Inprochone, RP-6870 | 017261 |
| <u>Iproplatin</u> | | CHIP, JM-9, 2-Propanimine, Platinum-complex (9CI) | 256927 |
| JB-11 | | | 249008 |
| <u>Liposome-Doxorubici</u> | n Kit | (TLC-Dox 99) | 620212 |
| Lometrexol Sodium | | LY 264618 disodium | |
| <u>Lomustine</u> * | CCNU | Belustine (nitrosourea), CeeNU, Chloroethylcyclohexylnitrosourea, ICIG 1109 (alkylating agent) | 079037 |
| Lovastatin | | Mevacor, Mevinolin# | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|---|--------------------------------|--|----------|
| M-Azido-Pyrimetham Ethane Sulphonato | | | |
| Mafosfamide | | Mafosfamide L-lysine | 626122 |
| Magnamycin | | Carbomycin, Carbomycin A, Delta- mycin A4, Magnamycin A (antibiotic) | 051001 |
| Mannosulfan | | R-52, Zitostop | 201289 |
| Marcellomycin | | | 265211 |
| Masoprocol | | Actinex, CHX 100, meso-NDGA | |
| <u>Maytansine</u> | MTS | Ansamacrolide compound (natural product) | 153858 |
| Mechlorethamine (HCl)* | HN2, NH2, NH2-HCl | Chloromethine-HCl, Dichloromethyldiethylamine-HCl, Mechlorethamine, Mustargen, Mustargen-HCl, Mustine, NH2 Nitrogen Mustard (alkylating agent) | 000762 |
| <u>Melphalan</u> * (<u>HCl</u>) | L-PAM, MPL, PAM | Alanine mustard, Alkeran, CB-3025, DL Sarcolysine, L-Phenylalanine- mustard, L-Sarcolysin, Melfalan, Nitrogen mustard, Sarcolysine, Sarkoklorin, Sarkolizin (alkylating agent) | 008806 |
| <u>Menogaril</u> | MEN | 7-OMEN, Tomosar, U-52,047 | 269148 |
| Merbarone | | 5-Carboxyanilino-2-thiobarbituric acid | 336628 |
| Mercaptopurine* | 6-MP, MP | Ismipur, Leukerin, Leupurin, Merca- leukin, 6-Mercaptopurin, 6-Mercapto- purine, Mercapurin, Mern, Purinethiol, Purinethol, 6-Thiohypoxanthine, 6-Thiopurine, 6-Thioxopurine (antimetabolite) | 000755 |
| 6-Mercaptopurine riboside | 6-MPR | 6-Mercaptoinosine, 6-Mercaptopurine- ribonucleoside, 6-Thioinosine | 004911 |
| MeTHHF | | See Enfolin sodium | 139490 |
| <u>Methotrexate</u> * | мтх | Amethopterin#, Antifolan, CL-14377, Folex, Methotrexate sodium, Mexate, Mexate AQ, R 9985, Rheumatrex (antimetabolite) | 000740 |
| Methyl CCNU | | See <u>Semustine</u> | 095441 |
| Methyl-G | MeG, MeGAG, MGA, MGBG, MGGH | Methyl-GAG, Methylglyoxal bisguanyl- hydrazone dihydrochloride monohydrate, Mitoguazone, Mitoguazone dihydrochloride | 032946 |
| Methyl methanesulp | honate | Methyl methanesulfonate, Methyl- methane sulfonate | 050256 |
| Methylene Dimethan | e Sulfonate | | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC</u> No. |
|--------------------------------------|--------------------------------|---|------------------|
| 6-Methylmercapto- purine riboside | 6-MMPR | 6-Methyl MP-riboside, Methylthioinosine, 6-Methylthiopurine riboside, SQ 21,977 (biochemical modulator) | 040774 |
| <u>Metoprine</u> | DDMP | Delcronol (antimetabolite) | 007364 |
| Meturedepa | | AB-132, Dimethylurethimine, Turloc | 051325 |
| Miltefostine | He-PC | Hexadecylphosphocholine | |
| Mithramycin | | Plicamycin | |
| Mitindomide | | CAS 10403-51-7 | 284356 |
| Mitocarcin | | 24281 (antibiotic) | |
| Mitoclomine | | | 114575 |
| Mitocromin | | B 35251 (antibiotic) | 077471 |
| <u>Mitogillin</u> | | | 069529 |
| Mitoguazone | | See Methyl G | |
| Mitomalcin | | Antibiotic | 113233 |
| Mitomycin* | MITC, Mito, MITO-C MMC, MTC | Ametycine, Mitocin-C, Mitomycin-C, Mutamycin (alkylating agent) | 026980 |
| Mitopodozide | | 2-ethylhydrazide, Podophyllic acid, Proresipar, SP-I 77 | 072274 |
| Mitosper | | 31595c | 117032 |
| <u>Mitotane</u> * | o.p'-DDD, o.p'-DDE | CB-313, Chloditan, Chlodithane, Lysodren, Mitotan (miscellaneous agent) | 038721 |
| Mitoxantrone (MCL)* | DHAD, DHAQ | CL-232,315, Dihydroxyanthracenedione dihydrochloride, Novantrone | 301739 |
| Mitozolomide | | Azolastone | 353451 |
| Mopidamol | | RA-233 | |
| Nafazatrom | | BAY G6575 | |
| Nafidimide | BIDA | Amonafide, Amonafide HCL (NSC 621093), Benzisoquinolenedione | 308847 |
| Nitrofurazone | | Furacin | |
| Nocodazole (Belgium |) | R 17,934 | 238159 |
| Nogalamycin# | | υ-1516 7 | 070845 |
| Novatrone | | Synthetic antineoplastic anthracenedione | |
| Oxisuran*# | | W-6495 | 356716 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC Mo</u> . |
|------------------------------|-------------|---|-------------------|
| | | | |
| Paclitaxel | | Taxol | 125973 |
| PALA | | CI-882, N-Phosphonacetyl-L-Aspartate disodium, Sparfosate <u>Sodium</u> | 224131 |
| Pancratistatin | | (miscellaneous agent) | 349156 |
| PCNU | | Nitrosourea | 095466 |
| Peliomycin | | Oligomycin B (antibiotic) | 076455 |
| Penberol | | | |
| Penclomidine | | | 338720 |
| Pentamethyl- melamine HCl | РММ | | 118742 |
| Pentamustine | NCNU | Salisburystin | 324595 |
| Pentosan Polysulfate | Sodium | Pentosan sulfate | 626201 |
| Peplomycin Sulfate | | NK 631, Pepleomycin | |
| Peptochemio | PTC | | |
| Phenesterine | | Fenesterin, Fenestrin, Phenesterin, Phenestrin | 104469 |
| Phetharbital | | Fedibaretta, N-Phenylbarbitol, Phenidiemal, Pyrictal | 085043 |
| Phosphoramide mustard# | PDA | Phosphami de | 069945 |
| Photofrin II | | See <u>Porfimer Sodium</u> | 603062 |
| Phyllanthoside | | | 328426 |
| Pibenzimol HCl | | Bisbenzamide, Bisbenzimidazole, Hoechst No. 33258 | 322921 |
| Piperazinedione compound | PZD | Actinomycete fermentation product, Compound 593A, Crystalline antibiotic (miscellaneous agent) | 135758 |
| Pipobroman* | PIBR | A-8103, Amedel, Vercyte (alkylating agent) | 025154 |
| Piposulfan | PISU | A-20968, Ancyte# | 047774 |
| Piroxantrone HCl | | Anthrapyrazole Dihydrochloride, Oxanthrazole, Oxantrazole HCl | 349174 |
| Plicamycin* | мітн, мтн | A-2371, Antibiotic LA 7017, Aureolic acid, Mithracin, Mithramycin#, PA-144 (antitumor antibiotic) | 024559 |
| Porfimer Sodium | | Dihematoporphyrin Ether, Photofrin II, Photofrin, CL 184,116 | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|----------------------------------|---------------------------------|---|----------|
| | | | |
| Porfiromycin | PORF | ENT-50825, N-Methylmitomycin C, Porfiromycine, Regamycin#, U-14,743 | 056410 |
| Prednimustine | | Sterecyt | |
| Procarbazine* (HCL) | IBZ, MIH, PCB, PCH, PCI, PCZ | Ibenzmethyzin hydrochloride, Matulane, Methylhydrazine, Natulan, Ro4-6467/1 (miscellaneous agent) | 077213 |
| Puromycin (HCL) | | CL-16,536, P-638, <u>Puromycin</u> , Stillo- mycin, Stylomycin, 1-MM, 3123L | 003055 |
| Pyrazine diazohydro | xide | Sodium N-Nitroso-Pyrazinamine | 361456 |
| Pyrazofurin# | PRZF, PZF | Antibiotic A 23813, Pirazofurin, Pyrazomycin, 47599 | 143095 |
| Pyrazoloacridine | | | 366140 |
| Pyrazolo-Imidazole compound | IMPY | Pyrazolo (2,3-a) imadazolidine | 051143 |
| Razoxane | | ICI-59118, ICRF-159, Tepirone | 129943 |
| Rhizoxin | RZN | | 332598 |
| Riboprine* | IPA | SQ 22558 | 105546 |
| Rifampin | | Ba 41166/E, L-5103, Rifadin, Rimactane (antibiotic) | 113926 |
| Rubidazone | | See Zorubicin hydrochloride | 164011 |
| Selenium | | | |
| Semustine | MCCNU, MCNU, Meccnu | Methyl CCNU, Methylcyclohexyl nitro- sourea, trans-Methyl CCNU | 095441 |
| Simtrazene# | | Centrazene, CL 26193 | 083799 |
| Sparfosate Sodium | | See PALA | 224131 |
| <u>Sparsomycin</u> | | บ-19183 | 059729 |
| Spirogermanium (<u>HCl</u>) | SPG | Spiro-32 | 192965 |
| Spiromustine | | <pre>Spirohydantoin mustard# (alkylating agent)</pre> | 172112 |
| Spiroplatin | | TNO-6 | 311056 |
| SR 2508 | | See Ancillary drugs | |
| Streptonigrin | SN, STN, STP | Bruneomycin, Methyl streptonigrin, Nigrin, Rufochromomycin, Rufocromo- mycin, 5278 R.P. | 045383 |
| Streptozocin | STR, Strept, STZ, SZ, SZC | Streptozoticin, Streptozotocin, U-9889, Zanosar, (nitrosourea) (alkylating agent) | 085998 |

| <u>Generic Name</u> | <u>Short</u> <u>Names</u> | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|-----------------------|--|--|-------------------|
| | | | |
| Strontium | | | |
| Sulofenur | | LY 186641 | |
| Suramin Sodium | | Antrypol, Bayer 205, Fourneau 309, Germanin, Moranyl, Naganin, Naganol, Naphuride Sodium, Sodium Suramin | 034936 |
| <u>Talisomycin</u> | | BU-2231A, Tallysomycin A# | |
| Taxol | | (See Paclitaxel) | |
| Taxotere | | | |
| Tazobactam | | YTR 830, (antibiotic, beta lactamase inhibitor) | |
| TCN | | See Triciribine phosphate | 280594 |
| <u>Tegafur</u> | FT | Florafur, Fluaid, Fluorofur, FT-207, Ftorafur, Futraful, MJF-12264 | 148958 |
| <u>Teniposide</u> | EPT, PTG | Epipodophyllotoxin, Thenylidene, Thenylidine-Lignan-P, Vee M-26, Vehem, VM-26, Vumon (plant alkaloid) | 122819 |
| Terephthalamidine | | N,N'- bis (p-N'- methylamidinolphenyl-) terephthalamidine 4 HCl, Symetamine | 057155 |
| <u>Teroxirone</u> | | Alpha TGI, Glycidyl isocyanurate, Henkel's compound, Triazinetrione- triepoxide, XB 2615 | 296934 |
| <u>D</u> -Tetrandrine | | Tetrandirine, Tetrandrin | 077037 |
| Tetraplatin | | Ormaplatin, U-77,233 | 363812 |
| Thalicarpine | тс | Taliblastine, Thaliblastine, Thalicarpin | 068075 |
| Theprubicin | THP | | |
| Thiamiprine | ITG | BW-57-323, Guaneran, Tiamiprine | 038887 |
| <u>Thioguanine</u> * | TG, 6-TG, TT-6, TT-G | 6-Mercaptoguanine, Tabloid, 6-Thioguanine, Tioguanin (antimetabolite) | 000752 |
| <u>Thiotepa</u> * | STEPA, TESP, TESPA, Thio-TEPA, TSP, TSPA | Oncotepa, Oncotiotepa, Tespa, Tespamin, Tespamine, Thiofozil, Thiophosphamide, Thiotef, ThioTepa, Thiotepa, Tifosyl, Tiofosfamid, Tiofosyl(Sweden), Tiofozil, TioTEF, Triethylene thiophosphoramide (alkylating agent) | 006396 |
| Thymidine | DT, DThyd, TdR | Deoxythymidine, 5-Methyldeoxyuridine, Thymidin, Thymine deoxyriboside | 021548 |
| <u>Tiazofurin</u> | TCAR | CI-909, Riboxamide | 286193 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> - |
|--|----------------|--|-------------------|
| TIC-mustard | BIC, BTIC, TIC | Imidazole mustard, Imididazole- carboximide, TIC NH2 (misc. agent) | 082196 |
| Timp-2 | | | |
| TMCA | TNCA | Citostal, Methyl ether, Trimethylcolchicinic acid, methyl ester, L-tartrate (1:1) | 036354 |
| Topotecan HCl | | Hycamptamine, SKF 104864-A | 609699 |
| Toyocamycin | | Antibiotic 1037, Antibiotic E 212, Toyokamycin, Unamycin B, Vengicide | 063701 |
| Treosul fan | | | |
| Triazinate | | See Baker's antifol | 139105 |
| Triazine antifol | | | 127755 |
| <u>Triciribine</u> <u>Phosphate</u> | TCN, TCN-P | Tricyclic nucleoside 5'-phosphate#, Tricycloside phosphate | 280594 |
| Triethylenemelamine | TEM, TET | Ho 1/193, M-9500, Persistol, R-246, SK-1133, Tretamine, Triamelin (alkylating agent) | 009706 |
| Triglycidylurazol | TGU | Anaxirone | 332488 |
| <u>Trimetrexate</u> | TMTX | CI-898, <u>Trimetrexate glucuronate</u> | 352122 |
| Trityl cysteine | | S-trityl-L-cysteine, Tritylthio- alanine | 083265 |
| Trofosfamide | | | |
| Tubercidine | | Antibiotic XK 101-1, Aplisol, 7-Deaza- adenosine, Sparsomycin A, Tubercidin, U-10071, 12,540 | 056408 |
| <u>Uracil</u> <u>Mustard</u> * | UHN2 | U-8344, Uracil, Uramustin, Uramustine (alkylating agent) | 034462 |
| <u>Uredepa</u> | | AB-100, Avinar#, Urethimine | 037095 |
| Urethane | | Ethyl carbamate# | 000746 |
| Uridine | | Uracil riboside, Urd, Uridin | 020256 |
| <u>Vapreotide</u> | | BMY-41606, RC-160 | |
| <u>Vidarabine</u> * | Ara-A | Adenine arabinoside, Ara-adenosine, Arabinosyladenine, CI-673, Spongo- adenosine, Vidarabin, Vidarabin- Thilo, Vira-A, Vitarabin, Vitarin | 404241 |
| <u>Vinblastine</u> * <u>Sulfate</u> | VBL, VELB, VLB | Velban, Vinblastine, Velsar, Vinca- leukoblastine sulfate, 29060-LE (vinca plant alkaloid) (mitotic inhibitor) | 049842 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|--------------------------------------|----------------|--|----------|
| Vincristine* Sulfate | LCR, VCR, VNCR | Leurocristine sulfate, Oncovin, Vincaleukoblastine, Vincasar, Vincristine, Vincrystine, Vinkristin, 37231 (vinca plant alkaloid) (antimetabolite) | 067574 |
| Vindesine | | Compound 112531 | |
| <u>Vindesine</u> <u>Sulfate</u> | DAVA, DVA | Desacetylvinblastine amide sulfate, Eldisine, LY-099094 | 245467 |
| Vinepidine Sulfate | | LY 119863 | |
| Vinglycinate Sulfat | <u>e</u> | 49040 | |
| <u>Vinleurosine</u> Sulfat | e# | Leurosine, Vinleurosine, 32645 | 528004 |
| Vinorelbine | VNB | Navelbine | |
| Vinrosidine Sulfate | # | Leurosidine sulfate, 3678l | |
| <u>Vinzolidine</u> <u>Sulfate</u> | VZL | LY-104208 (vinca plant alkaloid derivative) | |
| WR 2721 | | See Ancillary drugs | |
| Yoshi-864 | | | 102627 |
| <u>Zeniplatin</u> | | CL 286,558 | |
| Zidovudine* | AZT | See Azidothymidine | 602670 |
| Zinostatin | NCS, NZS | Neocarcinostatin, Neocarzinostatin#, Vinostatin | 157365 |
| Zorubicin (HCL) | RBD, RUB | Benzoyl hydrazone daunorubicin, RP-22,050, Rubidazone, Rubidazone- hydrochloride | 164011 |

COMBINATION REGIMENS

There are many combinations of antineoplastic drugs and hormones in general use. Teaching institutions are continuously establishing and testing protocols using new combinations of drugs. Listed in this segment are a few combinations so that you will understand how these combinations are named. As a registrar you should familiarize yourself with the antineoplastic drugs used in your hospital's oncology department to ensure the proper identification of its drug combinations.

ABVD Adriamycin, <u>Bleomycin</u>, <u>Vinblastine</u>, <u>Dacarbazine</u>

AcfuCy Actinomycin, <u>5-Fluorouracil</u>, <u>Cyclophosphamide</u>

ADOC Adriamycin, DDP, Oncovin, Cyclophosphamide

BACOP <u>Bleomycin</u>, Adriamycin, <u>Cyclophosphamide</u>, Oncovin, <u>Prednisone</u>

B-CAVe <u>Bleomycin</u>, CCNU, Adriamycin, <u>Vinblastine</u>

B-DOPA <u>Bleomycin</u>, <u>Dacarbazine</u>, Oncovin, <u>Prednisone</u>, Adriamycin

CAF <u>Cyclophosphamide</u>, Adriamycin, <u>5-Fluorouracil</u>

CAMF <u>Cyclophosphamide</u>, Adriamycin, <u>Methotrexate</u>, <u>5-Fluorouracil</u>

CAMP <u>Cyclophosphamide</u>, Adriamycin, <u>Methotrexate</u>, <u>Procarbazine</u>

CAP <u>Cyclophosphamide</u>, Adriamycin, <u>Cisplatin</u>

CAV <u>Cyclophosphamide</u>, Adriamycin, <u>Vincristine</u>

CAVV <u>Cyclophosphamide</u>, Adriamycin, <u>Vincristine</u>, VP-16

CCVPP CCNU, Cyclophosphamide, Vincristine, Procarbazine, Prednisone

CFP <u>Cyclophosphamide</u>, <u>5-Fluorouracil</u>, <u>Prednisone</u>

CHAD <u>Cyclophosphamide</u>, Hexamethylmelamime, Adriamycin, <u>DDP</u>

CHAP <u>Cyclophosphamide</u>, Hexamethylmelamine, Adriamycin, <u>Cisplatin</u>

CHIVPP Chlorambucil, Vinblastine, Procarbazine, Prednisone

CHOP <u>Cyclophosphamide</u>, 14-Hydroxydaunomycin (Adriamycin), Oncovin, <u>Prednisone</u>

CMC CCNU, Methotrexate, Cyclophosphamide

CMF Cyclophosphamide, Methotrexate, 5-Fluorouracil

CMFVP Cyclophosphamide, Methotrexate, 5-Fluorouracil, Vincristine, Prednisone

COAP <u>Cyclophosphamide</u>, Oncovin, Adriamycin, <u>Prednisone</u>

C-MOPP <u>Cyclophosphamide</u>, <u>Methotrexate</u>, Oncovín, <u>Procarbazine</u>, <u>Prednisone</u>

COM <u>Cyclophosphamide</u>, Oncovin, <u>Methotrexate</u>

COMLA <u>Cyclophosphamide</u>, Oncovin, <u>Methotrexate</u>, <u>Leucovorin</u> rescue, Ara-C

COMP <u>Cyclophosphamide</u>, Oncovin, <u>Methotrexate</u>, <u>Prednisone</u>

COP <u>Cyclophosphamide</u>, Oncovin, <u>Prednisone</u>

COP-BLAM Cyclophosphamide, Oncovin, Prednisone, Bleomycin, Doxorubicin, Procarbazine

CVP <u>Cyclophosphamide</u>, <u>Vincristine</u>, <u>Prednisone</u>

Cyta BOM <u>Vincristine</u>, <u>Methotrexate</u> and citrovorum factor

COMBINATION REGIMENS (Cont'd)

CYVADIC <u>Cyclophosphamide</u>, <u>Vincristine</u>, Adriamycin, <u>Dacarbazine</u>

DDP-CHAD <u>DDP</u>, <u>Cyclophosphamide</u>, Hexamethylmelamine, Adriamycin, <u>DDP</u>

DDP-MECY DDP, Methotrexate with Leucovorin rescue, Cyclophosphamide

DVB <u>Cisplatin</u>, <u>Vindesine</u>, <u>Bleomycin</u>

DVM Cisplatin, Vindesine, Methyl G

D-ZAPO <u>Daunorubicin</u>, <u>5-Azacytidine</u>, Ara-C, <u>Prednisone</u>, Oncovin

EBAP Eldisine, BCNU, Adriamycin, <u>Prednisone</u>

FAC <u>5-Fluorouracil</u>, Adriamycin, <u>Cyclophosphamide</u>

FACP <u>5-Fluorouracil</u>, Amethopterin, <u>Cyclophosphamide</u>, <u>Prednisone</u>

FAM <u>5-Fluorouracil</u>, Adriamycin, <u>Mitomycin</u>

FEC <u>5-Fluorouracil</u>, <u>Epirubicin</u>, <u>Cyclophosphamide</u>

FOMi <u>5-Fluorouracil</u>, Oncovin, <u>Mitomycin</u>

HEXA-CAF Hexamethylmelamine, Cyclophosphamide, Amethopterin, 5-Fluorouracil

MABOP <u>Mechlorethamine</u>, Adriamycin, <u>Bleomycin</u>, Oncovin, <u>Prednisone</u>

MACC <u>Methotrexate</u>, Adriamycin, <u>Cyclophosphamide</u>, CCNU

MACE <u>Methotrexate</u>, Adriamycin, <u>Cyclophosphamide</u>, <u>Etoposide</u>

MACOP-B Methotrexate, Doxorubicin, Cyclophosphamide, Vincristine, Prednisone, Bleomycin

MAID <u>Mesna</u>, <u>Doxirubicin</u>, <u>Ifosfamide</u>, DTIC

M-BACOD Methotrexate, Bleomycin, Adriamycin, Cyclophosphamide, Oncovin, Dexamethasone

MMV <u>Mitomycin</u>, <u>Methotrexate</u>, <u>Vincristine</u>

MOF <u>Mitomycin</u> or MeCCNU, Oncovin, <u>5-Fluorouracil</u>

MOF-S MeCCNU, Oncovin, <u>5-Fluorouracil</u>, <u>Streptozotocin</u>

MOPP <u>Mechlorethamine</u>, Oncovin, <u>Procarbazine</u>, <u>Prednisone</u>

MTX-CF <u>Methotrexate</u> with citrovorum factor rescue

MVAC <u>Methotrexate</u>, <u>Vinblastine</u>, Adriamycin, <u>Cisplatin</u>

PAC <u>Cisplatin</u>, Adriamycin, <u>Cyclophosphamide</u>

PEB <u>Cisplatin</u>, <u>Etoposide</u>, <u>Bleomycin</u>

POC <u>Procarbazine</u>, Oncovin, CCNU

POMP 6-Mercaptopurine, Oncovin, Methotrexate, Prednisone

PRO-MACE <u>Prednisone</u>, <u>Methotrexate</u>, Adriamycin, <u>Cyclophosphamide</u>, <u>Epipodophyllotoxin</u>,

ProMACE-MOPP <u>Prednisone</u>, <u>Methotrexate</u>, <u>Adriamycin</u>, <u>Cyclophosphamide</u>, <u>Epipodophyllotoxin</u>,

followed by Mechlorethamine, Oncovin, Procarbazine, Prednisone

PVB <u>Cisplatin</u>, <u>Vinblastine</u>, <u>Bleomycin</u>

TBP <u>Thiotepa</u>, <u>Bleomycin</u>, <u>Prednisone</u>

COMBINATION REGIMENS (Cont'd)

VAC <u>Vincristine</u>, Actinomycin D, <u>Cyclophosphamide</u>

VAP <u>Vincristine</u>, Adriamycin, <u>Procarbazine</u>

VATH <u>Vinblastine</u>, Adriamycin, <u>Thiotepa</u>, Halotestin

VBAP <u>Vincristine</u>, BCNU, Adriamycin, <u>Prednisone</u>

VBP <u>Vinblastine</u>, <u>Bleomycin</u>, <u>Cisplatin</u>

VMP <u>Vincristine</u>, <u>Methotrexate</u>, <u>Prednisone</u> (systemic consolidation therapy)

NOTES

Section 2

ANCILLARY DRUGS

Ancillary drugs are medications whose actions are not directed at the patient's malignancy per se but which enhance the effects of the cancer-directed therapy. For example, ancillary drugs may modulate the actions of specific chemotherapeutic agents by increasing their effectiveness in destroying tumor cells or by decreasing the potential for certain side effects. Ancillary drugs may be also used to control symptoms related to the patient's cancer or its treatment, or to manage other coexisting medical problems. Ancillary drugs are NOT to be coded as cancer-directed therapy.

Ancillary Drugs Administered with Chemotherapy

Ancillary drugs may be administered with chemotherapeutic agents to enhance their anti-neoplastic effects. The chemotherapeutic agent 5-fluorouracil (5-FU), an antimetabolite, works by inhibiting the enzyme thymidylate synthesis which is necessary for production of one of the components needed for DNA synthesis. Folinic acid (leucovorin) stabilizes the drug-enzyme complex and thus increases the cytotoxic effects of the 5-FU and is frequently administered with 5-FU for this purpose.

Ancillary drugs are also used for protection from side effects of chemotherapeutic agents. Folinic acid (leucovorin) is administered at a prescribed time after administration of methotrexate to decrease the toxic effects of the drug on rapidly-dividing cells such as bone marrow and the cells lining the gastrointestinal tract. Mercaptoethane sulfate (MESNA) is administered to patients receiving the alkylating agents ifosphamide and cyclophosphamide to prevent hemorrhagic cystitis. MESNA metabolizes in urine and binds to the toxic alkylating agent to prevent irritation of the bladder lining. Mannitol, a small sugar alcohol molecule, is administered intravenously in patients receiving cisplatin to increase urine flow (diuresis) and protect against kidney damage associated with chemotherapy.

Cancer cells may become resistant to chemotherapeutic agents through a variety of mechanisms. Multidrug resistance (MDR) occurs through genetic change and allows cancer cells to "pump out" chemotherapeutic agents before they can cause fatal damage to the cell. As part of the MDR phenomenon, cancer cells produce a substance called P-glycoprotein which binds to the chemotherapeutic agents within the cells and participates in the pump process. Calcium channel blocking agents, frequently used to treat cardiovascular disease, bind to P-glycoprotein and inhibit the MDR pump. The calcium channel blocking agent verapamil has been used with vinca alkaloids to reverse MDR in patients who have become resistant to vincristine or vinblastine, but this approach to modulating MDR is still under clinical investigation.

When large numbers of cells are destroyed by chemotherapy, materials from these cells must be cleared from the body. Uric acid, the chemical substance usually associated with gout, is produced as the cellular materials are dissolved by tissue enzymes and made ready for excretion by the kidneys. High plasma concentrations of uric acid may cause kidney damage (uric acid nephropathy) as the uric acid is filtered into the urine. This can be prevented in many patients by allopurinol, an oral medication that prevents the formation of uric acid by inhibiting the enzyme xanthine oxidase. Administration of allopurinol is particularly important in the treatment of hematologic malignancies (leukemia and lymphoma) because treatment of these tumors is most often associated with high levels of uric acid.

Radiosensitizers and Radioprotectants

In some instances, as with chemotherapy, the effects of radiotherapy may be modulated by agents which have no antitumor activity themselves. Radiosensitizing drugs augment the cytotoxicity of radiotherapy, and radioprotective agents diminish damage to normal, non-cancerous tissues. Many antineoplastic drugs, including the anthracyclines, 5-fluorouracil, and methotrexate, can enhance the cell killing effects of radiation; this is the basis for many combined-modality regimens. Other agents have been specifically synthesized as chemical modifiers of radiation effects and are under study in clinical trials.

The energy from therapeutic radiation ejects electrons from molecules in the target tissues. These electrons interact with other molecules in the cell to form free radicals which are unstable and highly reactive. DNA damage resulting from free radical formation contributes to cell injury and death. Thus any process which increases the DNA damage associated with free radical formation or inhibits damage repair results in radiosensitization.

Radiosensitizers are difficult to classify because they often have more than one mechanism of action. BUdR (Bromodeoxyuridine), an agent that is similar to the DNA precursor thymidine, is under study for brain tumors. Agents with electron affinity, such as metronidazole and misonidazole, are also in clinical investigation as radiosensitizers. Other agents in preclinical and clinical studies include SR 2508, Ro 03-8799, RSU 1069, and L-BSO.

Drugs or processes which interfere with free radical formation or facilitate repair of cellular damage result in radioprotection. WR-2721 is an interesting investigational new agent in this category because it appears to have activity not only as a radioprotectant but also in prevention of nephrotoxicity and neurotoxicity associated with cisplatin therapy.

Growth Factors

Suppression of blood cell production by the bone marrow and the resulting cytopenia is the side effect which most often limits the dose of chemotherapy that a patient can receive or necesitates a reduction in standard doses of chemotherapy. This is an important consideration because, for many cancers, tumor response to treatment depends on the amount of drug administered over a course of therapy (dose intensity).

Research studies of blood formation in bone marrow cultures have identified substances (or growth factors) which regulate blood cell growth and development. For example, GM-CSF is a colony stimulating factor (CSF) which promotes growth and development of clusters or "colonies" of blood cells which will become granulocytes and macrophages. G-CSF is a growth factor which acts at a later step in myeloid development and stimulates colonies which will produce predominantly granulocytes. Erythropoietin stimulates the production of red blood cells.

Recombinant DNA technology has allowed scientists to insert the genes responsible for growth factor production into bacteria, so that growth factors for clinical use can be produced by pharmaceutical companies from bacterial cultures. GM-CSF and G-CSF are now administered to patients who develop severe neutropenia as a side effect of chemotherapy, to facilitate bone marrow recovery and shorten the period of neutropenia. They also have been used to lessen bone marrow toxicity and allow administration of higher doses of chemotherapy.

Ancillary Drugs used in Symptom Control

Patients undergoing tumor-related therapy may develop uncomfortable symptoms related to their tumors (such as pain or cachexia), acute and subacute side effects from cancer treatment (such as nausea or mucositis), or more serious complications of therapy (such as systemic infections). Medications which are administered to control symptoms or side effects rather than as tumor-directed therapy are classified as ancillary drugs. Examples include analgesics for pain, antiemetics to prevent or control chemotherapy-induced nausea, clonidine for hot flashes associated with breast cancer therapy, and antibiotics for infection.

Some confusion arises when drugs that have antitumor effects in specific cancers are also used for symptom control. Corticosteroids such as decadron are important therapeutic agents in the treatment of lymphoid leukemias, lymphomas, and multiple myeloma. However, in the general cancer patient population, decadron is more often used for symptom control: to prevent nausea or drug reactions when chemotherapy is administered; to decrease the edema in tissue surrounding brain metastases; and, in some cases, to improve appetite in patients with cancer anorexia-cachexia syndrome.

| Generic Name | Short Names | Symonyms and Brand or Trade Names | + <u>NSC No</u> . |
|---------------------------------|-------------------------------------|---|-------------------|
| A | | (and the same) | |
| Acetaminophen | | (reduces fever) | |
| <u>Acetazolamide</u> | ACZ | Diamox (carbonic anhydrase inhibitor) (diuretic) | |
| <u>Acetylcysteine</u> | | 5052 | 111180 |
| Acyclovir | | Zovirax | |
| Adicillin | | Cephalosprin N (antibiotic) | |
| Albuterol | | Proventil, Ventolin (bronchodilator) | |
| Alendronate Sodium | | MK-217 (suppressant; bone resorption) | |
| Alizapride | | (anti-emetic) | |
| Allopurinol* | ALLO | Allopur, Atisuril, Bloxanth, BW-56-158, Epidropal, Foligan, Gotax, HPP, Lopurin, Milurit, Uricemil, Uriprim, Urosin, Zyloprim, Zyloric | 001390 |
| Alprazolam | | Xanax, U-31,889 (sedative) | |
| <u>Amifostine</u> | | Ethiofos#, Ethyol, Gammaphos, WR-2721 (radioprotector) (prevents nephrotoxicity, neurotoxicity) | 296961 |
| Aminobenzoate Potas | sium | (analgesic) | |
| Aminosyn | | (prevents acivicin-induced toxicity) | |
| <u>Amiodarone</u> | | Cordarone, L-3428, SKF 33134-A (cardiac depressant) | |
| Batanopride HCl | | BMY-25801-01 (Serotonin antagonist; anti-emet | ic) |
| Benznidazole | | (chemosensitizer) | |
| Benzquinamide | | Quantril, P-2647 (anti-emetic) | 064375 |
| Beta Carotene | | Solatene (ultraviolet screen) | |
| Borocaptate Sodium | <u>B 10</u> | Borolife, NASH; BSH (radioactive agent) | |
| Bromodeoxyuridine | BUdR, BDU, 5-BDU, BrdUrd, 5-BUDR | 5-Bromodeoxyuridine, 5-Bromodesoxy- uridine, Bromouracil deoxyriboside, 5-Bromouracil deoxyriboside, Broxuridine (radiosensitizer) | 038297 |
| <u>Calcium</u> <u>Carbonate</u> | | Cal-Sup, Component of Calcitrel, Component of Titralac (antacid) | |
| Calcium Gluconate | | Component of Calcet, (calcium replenisher) | |
| <u>Calcitonin</u> * | | Calcimar, Cibacalcin (regulates calcium) | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|----------------------------|----------------|--|-------------------|
| | | | |
| Chlornaphazine* | | CB-1048, Chlornaftina, Chlornaphazin, Chlornaphthin, Chloronaphthina, Chloronaphthine, Cloronaftina, Erysan, Nafticlorina, Naphthylamine mustard, R-48 | 062209 |
| Chlorpromazine* | | <pre>Chlor-PZ#, Thorazine (anti-emetic; antipsychotic)</pre> | |
| Cimetidine | | Tagamet | |
| Clonidine | | Catapres-TTS (controls hot flashes in breast therapy) | cancer |
| Codeine | | Analgesic (narcotic) | |
| Coenzyme Q | | Reduces myoclonic seizures | |
| Combid spansule | | (See Isopropamide Iodide and Prochlorperazine Maleate) | : |
| CSF-GM | | CSF-GM (H)/Yeast, Recombinant (Immunex), Recombinant Human GM-CSF Granulocyte-Colony Stimulating Factor | 613795 |
| CSF-GM | | CSF-GM (Immunex) Granulocyte-Colony Stimulating Factor (Amgen) | 614629 |
| CSF-GM | | (Schering) Plough CSF-Granulocyte-Macrophage | 617589 |
| CSF-GM | | Granulocyte-Macrophage-Colony Stimulating Factor (Recombinant Human) (E.coli) CSF-GM E.coli (Hoechst) | 622183 |
| CSF-Macrophage | | Macrophage-Colony-Stimulating Factor (M-CSF) Cetus | 625377 |
| CSF-Macrophage | | (H-CSF) Cetus | 635258 |
| Danazol | | Danocrine, Win 17,757 (antipituitary suppressant) | |
| <u>Dexamethasone</u> *∇ | DECA, DM, DSM, | Aeroseb-Dex, Decaderm, Decadron Tablets, Decaspray, Component of Deronil, Component of Dexacidin, Dexasone, Dexone, Hexadrol Elixir, Hexadrol Tablets, Maxidex, Component of Maxitrol, Component of Tobradex (glucocorticoid) | 345211 |
| Deferoxamine | | Prevents cardiac toxicity | 527604 |
| Defosfamide | | Desmophosphamide | 040627 |
| Desmethyl- misonidazole | DES-ME | Demethylmisonidazole, Ro-5-9963, SR-1530, (radiosensitizer) | 261036 |

 $[\]nabla$ If used in the treatment of the malignancies mentioned in the introduction on p. 37 (bold), code as hormonal therapy. For all other cancers, consider ancillary unless specifically stated to be hormonal therapy.

| Generic Name | Short Names | Symonyms and Brand or Trade Names | + <u>NSC No</u> . |
|-----------------------------------|-----------------|--|-------------------|
| | | | |
| Diethyldithio- carbamate | DDC, DDTC, DEDC | Cuprol, Dithiocarb, Thiocarb | 038583 |
| Diltiazem HCL | | Cardizem (vasodilator, calcium influx inhibitor) | |
| <u>Dimepranol</u> <u>Acedoben</u> | | Component of Isoprinosine (immunomodulator) | |
| Diphenidol | | Anti-emetic | |
| Diphenhydramine HCL | | Benadryl, Component of Benylin, Component of Caladryl (antihistaminic) | |
| Dipyridamole | | Persantine, Ra-8 (coronary vasodilator) | 515776 |
| Domperidone | | R-33-812 (anti-emetic) | |
| Droperidol* | | Inapsine, Component of Innovar, McN-JR-4749, R-4749 | |
| Elliot's B solution | | K-4/47 | 614386 |
| EMLA | | | |
| Endorphin | | Analgesic | |
| Erythropoietin | | (See biological response modifiers) | |
| <u>Etanidazole</u> | | Radinyl, SR 2508 (hypoxic cell radiosensitizer) | 301467 |
| Ethacrynic Acid | | Edecrin, MK-595 (diuretic) | 085791 |
| Ethiodized Oil* | | Ethiodol-131# (radio-opaque medium) | |
| Ethiofos | | See <u>Amifostine</u> (radioprotector) | 296961 |
| Etidronate Disodium | | Didronel (calcium regulator) | |
| Fansidar | | Sulfadoxine/Pyrimethamine | |
| Fentanyl Citrate | | Analgesic; narcotic | |
| Fluconazole | | Diflucan (antifungal) | |
| Fluosol-DA | | Radiation-sensitizing agent | |
| Flurbiprofen | | Ansaid, BTS 18,322, U-27,182 (analgesic; anti-inflammatory) | |
| Furosemide* | | Disal, Lasix, LB-502 (diuretic) | |
| G-CSF | | r-met Hu G-CSF (Granulocyte Colony Stimulating Factor) | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|-----------------------|-----------------------------------|--|-----------|
| Ganciclovir | DHPG | Cytovene (therapy for immunosuppressed | |
| | | patients, bone marrow recipients) | |
| Glycine | | Corilin, component of (irrigating solution) | |
| GM-CSF | | Granulocyte macrophage colony stimulating factor | |
| Gold Au 198 | | Aurcoloid-198#, Aureotope# , Auroscan-198# (diagnostic aid, liver imaging; radioactive agent) | |
| Granisetron | | BRL-43694 (serotonin antagonist; anti-emetic) | |
| Guanfacine HCl | | Tenex | |
| Haloperidol | | Haldol, McN-JR-1625, R-1625 (antipsychotic) | |
| Hematoporphyrin De | rivative | Emits fluorescence in neoplastic cells | |
| Heroin HCl | | Diacetylmorphine dihydrochloride, Diamorphine HCl, Heroin hydrochloride | 302357 |
| Hydrazine Sulfate | | (anti-cachexia) | 150014 |
| Hydromorphone HCl | | Dilaudid (analgesic; narcotic) | |
| Hydroxyzine HC1 | | Atarax, Component of Ataxaroid, Component of Cartrax, Component of Enarax, Component of Marax, Quiess, Vistaril, Component of Vistrax (tranquilizer) | |
| Hyperbaric Oxygen | | Radiation-sensitizing agent | |
| Idoxuridine* | IDU, IDUR, IdUrd, IUdR, 5-IUDR | Allergan 211, Dendrid, Emanil, Herpesil, Herpidu, Herplex, Herplex Liquifilm, Idexur, Idoxuridin, Iducher, Idulea, Idu-Oculos, Iduridin, Iododeoxyuridine, Joddeoxyuridin, Kerecid, Ophthalmadine, SK&F 14287, Stoxil, Synmiol (radiosensitizer) | 039661 |
| <u>Indomethacin</u> * | | <pre>Indocin (anti-inflammatory; lowers blood calcium) (analgesic)</pre> | |
| Inosine | | Component of Isoprinosine | |
| Isoprinosine | | Imunovir (Combination product: see Dimepranol Acedoben and Inosine) | |
| Isopropamide Iodio | <u>de</u> | (Used with Prochlorperazine Maleate as Combic (anticholinergic; anti-emetic) | spansule) |
| Itraconazole | ITZ | R-51,211, Sporonax (antifungal) (Rx of Candidiasis) | |
| Ketorolac Trometha | <u>amine</u> | Toradol (anti-inflammatory; analgesic) | |
| L-BSO | | $\underline{L}	ext{-Buthionine}$ sulfoximine (radiosensitizer) | 326231 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|--|-------------|---|-------------------|
| | | | |
| <u>Leucovorin</u> (<u>calcium</u>)* | CF, C.F. | Calcium folinate, Calcium leucovorin, Citrovorum factor, Folinic acid, Lederfoline, Leucosar, Wellcovorin | 003590 |
| Levonantradol (HCl) | | CP-50,556-1 (analgesic) | 331615 |
| Lidocaine | | Anestacon, Lidomantle-HC, component of of, Xylocaine (local anesthetic) | |
| <u>Lithium</u> <u>Carbonate</u> | | C.P. 15,467-61, Eskalith, Lithane, Lithobid, Lithonate, Lithotabs, (antimanic) | 016895 |
| L-Leucovorin | | eventuals, (antimovie) | |
| Lonidamine | | Radiation-sensitizing agent | |
| Lorazepam | | Ativan, Wy-4036 (anti-emetic) | |
| <u>Loxoribine</u> | | Immunostimulant; vaccine adjuvant | |
| Magnesium Sulfate | | Anti-convulsant; laxative; electrolyte replenisher | |
| Mannitol | | Diuretic | |
| Mannomustine | BCM | Degranol, Mannit-Lost, Mannit-Mustard, Mannitol Mustard | 009698 |
| MDL-72222 | | Serotonin antagonist; anti-emetic | |
| MDL-73147EF | | Serotonin antagonist; anti-emetic | |
| Meperidine HCL | | Demerol, Demerol HCl, Mepadin#, Component of Mepergan (analgesic; narcotic) | |
| <u>Mesna</u> * | | D 7093, Mesnex (uromitexan), Mesnum, Mistabron, Mitexan, Mucofluid, Sodium 2-mercaptoethanesulfonate (prevents bladder toxicity) | 113891 |
| Methadone HCl | | Adanon HCl#, Dolophine HCl, Component of Mespergan, Methadose (analgesic; narcotic) | |
| Methoxsalen | | 8-MOP, Oxsoralen (photosensitizer) | |
| Methylphenidate HCl | <u>.</u> | Ritalin | |
| Metoclopramide (<u>HCl</u>)* | MCP | AHR-3070-C, Octamide PFS, Reglan (anti-emetic) | |
| Metronidazole | | Radiosensitizer (See also Drugs Used in Rx of AIDS complications) | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC No</u> . |
|-----------------------|--------------|--|-------------------|
| | | | |
| <u>Metyrosine</u> | | Anti-hypertensive; therapy for malignant pheochromocytoma | |
| Midazolam HCL | | Ro 21-3981/003, Versed (anesthetic; CNS depressant) | |
| Misonidazole | MNI | Ro-7-0582, SR-1354, SRI-1354 (radiosensitizer) | 261037 |
| Molybdenum | | Metabolic therapy | |
| Morphine Sulfate | | Astramorph, Duramorph, MS Contine, MSIR (analgesic; narcotic) | |
| <u>Nabi Lone</u> | | Cesamet, Cpd 109514 (tranquilizer) | |
| Naloxone HCl | | Antagonist to narcotics | |
| Naltrexone | | EN-1639A, Trexan (antagonist to narcotics) | |
| Naproxen | | Naprosyn, RS-3540 (anti-inflammatory; analgesic; antipyretic) | |
| Niacin | | Niac, Nicobid, Nicolar (therapy for patients on parenteral nutrition) | |
| Nifedipine | | Calcium channel blocking agent | |
| Nortriptyline HCl | | Aventyl HCl, Pamelor, 38489 (antidepressant) | |
| Ondansetron HCl | | GRF 38032F, Ondanserin, Zofran (serotonin anta anti-emetic, antischizophrenic, anxiolytic) | agonist; |
| Pamidronate Disodiu | <u>.m</u> | Aredia (suppressant; bone resorption) (Hypocal | cemia) |
| Papaverine HCl | | Cerespann, Component of Copavin, Pavabid (smooth muscle relaxant) | |
| <u>Pentoxifylline</u> | | BL 191, Trental (vasodilator) (radiation-sensitizing agent) | |
| <u>Perphenazine</u> | | Component of Estrafon, Component of Triavil, Trilafon | |
| Pimonidazole | | Ro 03-8799 (radiosensitizer) | |
| Piperazine | | Antiren, Dispermine, Eraverm, Hexahydropyrazine, Lumbrical, Piperazidine, Piperazin, Pipersol, Pyrazine Hexahydride, UN 2579, UN 2685, Uvilon, Vermex, Component of Wazine-34, Wurmirazin (anthelmintic) | 000474 |
| Piroxicam | | CP-16,171, Feldene (anti-inflammatory) | |
| Polyethylene Glyco | <u>l</u> PEG | Carbowax Sentry, Colyte, component of, Pluracol E400, E600, E1450 | |
| <u>Probenecid</u> | | Uricosuric; increases excretion of uric acid in respiratory acidosis | |

| Generic Name | Short Names | Symonyms and Brand or Trade Names | + <u>NSC No</u> . |
|----------------------------------|--------------|---|-------------------|
| Prochlorperazine* | | Compazine, Component of Eskatrol# (anti-emetic) | |
| Prochlorperazine Ma | <u>leate</u> | Used with Isopropamide Iodide as Combid spans (anticholinergic; anti-emetic) | sule |
| Promazine HCL* | | Sparine (antipsychotic) | |
| Promethazine HCL* | | Anergan 25, Anergan 50, Component of Mepergan, Phenergan, Component of Phenergan VC, Remsed (anti-emetic) | |
| <u>Pyrazinamide</u> | PZA | ,, ,, ,, ,, ,, ,, ,, , | |
| Pyridoxine HCL | | Beesix, Hexa-Betalin, Vitamin B6 (reduces neurotoxicity) | |
| Pyrimethamine | | BW 50-63, Chloridin, Chloridine, Darapram, Daraprim, Daraprime, Diaminopyritamin, Erbaprelina, Component of Fansidar, Malocide, Pirimecidan, Pirimetamina, Tindurin | 003061 |
| Quinacrine (HCl) | Atab, QUIN | Acrichin, Acrinamine, Acriquine, Akrichin, Antimalarina, Atabrine HCl, Haffkinine, Italchine, Mepacrine, Quinactine | 014229 |
| Quinidine | | Chemosensitizing agent | |
| R-Verapamil HCl | | Calcium channel blocking agent | 632821 |
| Rhenium-186 | HEDP | Hydroxyethylidene diphosphonate (palliative, skeletal metastases) | |
| Riboflavin | | Cancer preventative | |
| Ro-03-8799 | | Pimonidazole (radiosensitizer) | |
| RSU 1069 | | Radiosensitizer | |
| Serotonin | | GR 38032 F (antagonist) | |
| Sodium Chloride | | Ayr (chemoprotective agent reduces nephrotoxicity) | |
| Sodium Fluoride | | Floridine, Fluorol, Pergantene, Xaridium, Zymafluor | 077385 |
| Sodium Iodide | | I 131 Idotope | |
| Sodium Phosphate P | 32 | Radioactive agent, diagnostic aid | |
| <u>Sodium</u> <u>Thiosulfate</u> | | Sulfactol#, Component of Komed, Component | 045624 |
| Sparfloxacin | | of Tinver (reduces nephrotoxicity) | |
| Spiramycin | | Leucomycin | 055926 |
| SR 2508 | SN | See <u>Etanidazole</u> (radiosensitizer) | 301467 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|---------------------------|-------------|---|----------|
| | | | |
| <u>Sucralfate</u> | | Carafate (anti-ulcerative) | |
| <u>Sulfadiazine</u> | | Sulfonamide Duplex, component of# | |
| Sulfamethoxazole | | Azo Gantanol, component of, Bactrim, component Cotrim, component of, Gantanol, Septra, component, Sulfatrim, component of | |
| <u>Sul indac</u> | | Clinoril (anti-inflammatory) | |
| Tetrahydro- cannabinol | THC | Cannabinol, Delta-9-THC, <u>Dronabinol</u> , Marinol, SP-104 (anti-emetic) | 134454 |
| Theophylline | | Bronchodilator | |
| Triflupromazine HCl | * | Vesprin (antipsychotic) | |
| Trimethobenzamide H | <u>cı</u> | Tigan (anti-emetic) | |
| Trimethoprim | TMP | Bactrim, component of, Cotrim, component of, Proloprim, Septra, component of, Sulfatrim, component of, Trimpex | 106568 |
| Tropisetron | | <pre>ICS-205930 (Serotonin antagonist; anti-emetic)</pre> | |
| Urokinase | | Plasminogen activator; thromboembolytic therapy | |
| Verapamil (HCl)* | | Calan, Isoptin, Isoptin hydrochloride, Izoptin, Verapamil (Isoptrin), Verelan (chemosensitizer) (calcium channel blocker) | 256325 |
| <u>Vitamin E</u> | | Anti-fibrosis therapy | |
| Warfarin Sodium* | | Coumadin, Panwarfin (anticoagulant) | 059813 |
| WR 2026 | | | |
| WR 2721 | | See Amifostine | 296961 |

NOTES

Section 3

DIFFERENTIATION-INDUCING AGENTS

A new approach to cancer therapy seeks to prevent the development of a cancer cell or to reverse its malignant potential by inducing its differentiation.

Most normal cells are differentiated, that is, they have progressed through stages of normal development to have specific characteristics and function. During the process of differentiation, normal cells tend to lose the ability to proliferate. In most tissues, a few progenitor cells (stem cells) remain undifferentiated and retain the ability to produce new cells, to replace the normal loss of mature cells. Most cancers probably develop from these less mature progenitor cells.

A precursor cell that undergoes malignant transformation usually loses its ability to differentiate in an efficient and orderly manner, while it retains the ability to proliferate. However, many tumors have enough differentiation to allow pathologists to determine the tissue of origin or "primary site" of the tumor. In some tissues there is an orderly progression from benign tissue to noninvasive but premalignant lesions to frank malignancy. For example, in the uterine cervix, normal tissue may progress through stages of hyperplasia, metaplasia, anaplasia, carcinoma-in situ, to invasive carcinoma.

The objective of using differentiation-inducing agents is to prevent the onset of frank malignancy and/or to convert a premalignant tumor into a more differentiated growth without invasive and metastatic properties. In contrast to conventional cytotoxic chemotherapeutic agents, which cause the death of tumor cells, differentiation-inducing agents may control the expression of specific genes and thereby prevent the progression of a premalignant cell to a malignant state.

Current experimental studies suggest that non-cytotoxic concentrations of a number of low molecular weight chemicals can induce differentiation of several tumor cell lines in cultures, and several of these agents are under clinical study. These agents include 1,25-dihydroxyvitamin D, dimethyl sulfoxide, hexamethylene bis-acetamide, 4-hydroxyphenyl retinamide and 13-cis retinoic acid.

Some drugs may have different effects at different doses. For example, Ara-C is used as cytotoxic chemotherapy in high doses to treat acute myeloblastic leukemia. It is also used in much lower doses as a differentiating agent in pre-leukemia (myelodysplastic syndromes), to stimulate immature cells to progress to more mature forms.

Outside clinical trials for cancer treatment, differentiating agents are most often used in patients as chemoprevention of a new cancer. For example, a patient who has had an oral cancer removed may take 13 cis-retinoic acid to prevent a second (new) oral tumor. Thus, differentiating agents should not be coded as cancer-directed therapy unless the chart specifically states that it is.

DIFFERENTIATION-INDUCING AGENTS

| Generic Name | Short Names | <u>Synonyms and</u> <u>Brand or Trade Names</u> | + <u>NSC</u> No. |
|----------------------------------|--|--|------------------|
| | Do not code as c | cancer-directed therapy. See Introduction. | |
| <u>Acitretin</u> | | Soriatane | |
| <u>Dimethyl</u> <u>Sulfoxide</u> | | See Biological Response Modifiers | 000763 |
| 1,25-Dihydroxy- vitamin D3 | Vit 03 | | |
| <u>Etretinate</u> | | Tegison | |
| HMBA | | Hexamethylene bisacetamide, Hexamethylene diacetamide | 095580 |
| 4-Hydroxyphenyl retinamide | 4-HPR | | |
| Mycophenolic acid | | Melbex#, 68618 | 129185 |
| Phenyl Acetate | | | |
| Retinoic acid | Vit A | Accutane, <u>Isotretinoin*</u> , Neovitamin A acid, Ro 4-3780, 13-cis-Retinoic acid, 13-cis-Vitamin A acid | 329481 |
| Retinyl Acetate | RA | | |
| Sodium Butyrate | | | |
| <u>Tretinoin</u> * | Vit A | Aberel, Airol, All- <u>trans</u> -retinoic acid, Dermairol, Eudyna, Retin-A, Retinoic acid, Vitamin A acid | 122 7 58 |
| | AGENTS WITH BOTH C | YTOTOXIC AND DIFFERENTIATING PROPERTIES | |
| | Cod | e as chemotherapy. | |
| Cytarabine | Ara-C | Cytosar-U, Tarabine, U-19920, (under investigation for Rx of Progressive Multifocal Leukoencephalopathy (PML)) | |
| Azacitidine | 5-AC, 5-AZA, 5-AZAC, AZC, 5-AZCR | Antibiotic U-18,496, Azacytidine, 5-Azacytidine, Ladakamycin, Mylosar (antimetabolite) | 102816 |

Methylformamide, Monomethylformamide

N-Methylformamide

NMF

003051

NOTES

Section 4

BIOLOGIC THERAPY OF CANCER

Biologic therapy or immunotherapy is defined as treatment that produces its anticancer effect by enhancing the patient's natural defense mechanisms against cancer. This is accomplished by administering agents that modulate the patient's immune response to his/her tumor, by providing additional natural immune molecules made through new technologies, or by transferring immune cells stimulated to act against the patient's cancer. The drugs, chemicals, or biologic agents used in biologic therapy are often referred to as biological response modifiers.

Biologic therapy is a new and rapidly developing area of oncology. Research in this area has been stimulated by increased understanding of the activity of the human immune system and by advances in biotechnology that have made possible the production of antibodies and other immune molecules by recombinant DNA methods.

The Human Immune System

In contrast to most other organ systems, the cells of the immune system are not in constant contact with each other. Lymphocytes, monocytes, macrophages, and other immune cells move freely throughout the lymphatic and circulatory systems. Their specific functions and complex interactions give rise to a highly organized system of host defense against "non-self" substances, organisms, or cells.

Antibodies are proteins that are produced by the body in response to a foreign substance (antigen). These protein complexes consist of units (immunoglobins) which are composed of pairs of heavy (longer) and light (shorter) polypeptide chains. Antibodies have distinct binding sites which react with the specific antigen that stimulated their production. This phenomenon of humoral immunity explains why one rarely has childhood diseases such as measles or mumps twice.

A recent advance in immunology has been the recognition that certain groups of lymphoid cells can secrete a second type of protein molecule that is structurally different from antibodies. These cytokines are produced in small amounts and are usually not detectable in plasma. They function as hormones, acting on a broad spectrum of immune cells and on cells outside the immune system. Identification of specific cytokines and exploration of their activities have led to new and exciting approaches to biotherapy of cancer.

Immune System Cells

Lymphocytes, the central cells in immune function, are classified according to development and function as B cells, T cells, and null cells. In addition, detailed studies of leukocyte cell surface antigens have led to identification of groups of antigens called clusters of differentiation (CDs) which are also used to classify lymphocyte subpopulations.

In humans, B cells develop in the bone marrow. As they mature, they develop immunoglobins on their surface which function as antigen receptors; they also develop cell surface receptors for lymphokines (cytokines produced by lymphocytes). B cells require both the presence of antigen and stimulation by lymphokines to produce an antibody response.

T cells mature and differentiate in the thymus before dispersing to other lymphoid tissue. In contrast to B cells which function in humoral immunity, T cells have prominent roles in cellular immunity. T cells also recognize antigen in a manner quite different from that of B cells. In general, T cells cannot respond to an antigen without the assistance of other cells. Antigen must be "processed" by other cells and presented to the T cell as a combination of antigen components with major histocompatibility complex (MHC) molecules, while B cells can recognize and respond to unprocessed antigen.

Traditionally, T cells have been classified by function as helper, suppressor, or cytotoxic T cells. Currently, two major subsets of T cells are identified by the class of MHC molecules required for antigen recognition. T cells which require Class I MHC molecules (i.e., HLA-A, B, or C) express the cluster of differentiation CD8 on their cell surfaces, while T cells which require Class II MHC molecules (i.e., DP, DQ, and DR) express the cluster of differentiation CD4. While cells with helper and cytotoxic functions can be identified within each subset, CD4+ T cells are more likely to express helper function, and CD8+ cells are more frequently cytotoxic or are more often capable of suppressing the immune response of other lymphoid populations.

Null cells are lymphoid cells which express neither T cell nor B cell surface markers. Subgroups derived from null cells include natural killer (NK) cells, lymphokine-activated killer (LAK) cells, and cells involved in antibody-dependent cellular cytotoxicity.

Reticuloendothelial cells, especially circulating monocytes and tissue macrophages, are also important components of the immune system. Monocytes are produced and mature in the bone marrow, enter the circulation, and then exit into tissue where they may die or mature further to become tissue macrophages. Macrophages are highly phagocytic cells which engulf, process, and present antigen components to T lymphocytes. They also produce monokines (cytokines) which facilitate T-cell activation and interaction with B cells to induce antibody formation.

Immune-mediated Cell Destruction

The immune system can cause the destruction of target cells through a variety of mechanisms. Certain subtypes of antibodies (IgG and IgM immunoglobins), attached to cell surface antigens, can activate **complement**, a system of functionally linked proteins that interact to damage the target cell membrane. Antibodies attached to cell surface antigens can also act as **opsonins**, making the target cells more susceptible to phagocytosis by macrophages.

Target cell lysis can occur by direct actions of immune cells. Cytotoxic T lymphocytes (CTLs) bind to surface antigens on target cells and release cytotoxic substances by the T cell. CTLs are not damaged during this process; they detach from the target cell and recycle to attack other target cells. Antibodies to surface antigens on target cells or lectins can serve as a crosslink between target cell and cytolytic immune cells in antibody (or lectin)-dependent cellular cytotoxicity. NK cells have the ability to lyse cells in cultured cell lines without prior immune sensitization to the cells, but they appear to have little effect on fresh tumor cells. LAK cells develop the ability to kill tumor cells after exposure to the cytokine interleukin-2 (IL-2).

Many cytokines can mediate cell destruction, either by direct toxicity toward target cells or by initiating an inflammatory reaction in target tissue.

Tumor Antigens

When a cell becomes malignant, it undergoes biochemical changes that may result in the production of new cell surface proteins capable of distinguishing it from a normal cell. Such tumor antigens may be seen as foreign by the host immune system and may result in a specific immunologic response to the tumor cells. In animal models, tumor antigens may be highly specific for an individual tumor cell line. However, most tumor antigens studied in naturally-occurring tumors are shared by different tumors and may even be expressed to a lesser extent by normal cells; these are termed tumor-associated antigens.

Tumor infiltrating lymphocytes (TILs) have been isolated from the inflammatory infiltrates of a variety of human cancers. TILs with specific cytolytic function have been isolated from human melanoma and occasionally from renal cell cancer; nonlytic TILs with specific recognition of tumor antigens have been identified in patients with melanoma and breast cancer. TILs can be activated and expanded in culture with IL-2.

Immunotherapy

Agents and substances used in biologic therapy, or immunotherapy, may be classified according to the way in which they modify the host defense mechanism.

Active Immunotherapy

Active immunotherapy refers to immunization of the cancer patient with materials designed to elicit an immune reaction against the patient's tumor. Early attempts at immunotherapy focused on nonspecific stimulation of the immune system by bacterial products, in hopes that general immune stimulation would lead to an increased host immune response to its established tumor. This approach developed from observations of tumor regression following severe bacterial infections or exposure to bacterial toxins. Agents used for nonspecific immune stimulation include BCG, MER, *C. parvum*, and levamisole.

Prospective randomized clinical trials have failed to support a role for nonspecific immunotherapy in the cancer treatment, with a few notable exceptions. The addition of levamisole to adjuvant therapy with 5-fluorouracil has improved survival in patients with Stages B2 and C colon cancer. Intralesional immunotherapy of cutaneous metastases from malignant melanoma with BCG has resulted in good local control, and BCG has been used successfully as instillation therapy for the treatment of superficial bladder cancers.

Other approaches to active nonspecific immunotherapy involve administration of cytokines such as interferon or IL-2 to enhance host immune response. Interferons (INFs) belong to a family of proteins produced by cells in response to viral infection. Three major classes of interferons have been identified: INF-alpha, produced by leukocytes; INF-beta, produced by fibroblasts and epithelial cells; and INF-gamma, produced by activated lymphocytes. In addition to antiviral and immunomodulatory effects, biologic properties of interferons include antiproliferative activity, regulation of differentiation, interaction with other cytokines, and enhancement of tumor-associated antigens.

Recent progress in recombinant DNA technology has made large quantities of recombinant interferons (rINFs) available for clinical studies. Interferons have antitumor activity against a variety

of solid tumors and hematologic malignancies, including hairy cell leukemia, chronic myelogenous leukemia (CML) and other myeloproliferative disorders, cutaneous T-cell lymphoma, and Kaposi's sarcoma.

Interleukin 2 (IL-2), a cytokine produced by activated T lymphocytes, has several actions of interest in immunotherapy. IL-2 stimulates the growth and proliferation of lymphoid cells and causes the release of a variety of other cytokines; it also enhances cytolytic activity of adoptively transferred cells (see below). IL-2 has no direct activity against cancer cells.

IL-2 has been used alone and in conjunction with adoptive transfer of lymphoid cells. IL-2 as a single agent has produced complete and partial responses in patients with advanced renal cell cancer and melanoma. rIL-2 was the first biologic response modifier to receive approval by the U.S. Food and Drug Administration for the treatment of advanced cancer that acted only through immune mechanisms.

Active specific immunotherapy involves immunization to boost the cancer patient's immune response specifically against his/her own tumor. This approach relies on the presence of tumor-associated antigens on the surface of the malignant cells and on the ability of these antigens to produce a host immune response. Development of tumor vaccines is an area of concentrated research effort at the present time.

Passive Immunotherapy

In cancer treatment, passive immunotherapy refers to the administration of specific antibodies or activated immune cells directed toward a patient's tumor. This is an area of intense research effort, and the therapeutic approaches described below represent investigational procedures at cancer research centers rather than standard cancer care.

The recognition of specific tumor antigens and the development of monoclonal antibodies (MOABs) with relatively unique specificity for these antigens have fostered new approaches to cancer diagnosis and therapy. As with antibody-mediated cytotoxicity described above, unmodified MOABs can kill tumor cells through activation of complement or through antibody-dependent cellular cytotoxicity. Tumor responses to unmodified MOABs have been observed in the treatment of B-cell lymphoma and in T-cell leukemia or lymphoma.

Monoclonal antibodies can also be conjugated with nonimmune cytotoxic substances to enhance antitumor effects. Examples of these conjugates include chemotherapeutic drugs (e.g., adriamycin), toxins (e.g., ricin), and radionuclides (e.g., ¹³¹I, ¹¹¹In, ⁹⁰Y).

Adoptive immunotherapy refers to the administration of cells with antitumor activity to the cancer patient. This approach uses either LAK cells or TILs in combination with IL-2 to confer enhanced cell-mediated tumor lysis. In a variety of tumors, the response rate to LAK cells with IL-2 has been relatively low: best objective response rates of 15-30 percent have been observed for melanoma and renal cell cancer. Pilot studies of therapy with TILS and IL-2 in patients with advanced cancer are currently in progress.

Clinical Research in Cancer Immunology

The interaction between the immune response of a tumor-bearing individual and his/her cancer is undergoing intensive investigation. The major goals of immunological research are earlier diagnosis and effective immunization as a mode of cancer therapy. Although much has been learned about the etiology and biology of many tumors, much is yet unknown about tumor immunology. Dramatic expansions in understanding the role of the immune system in the etiology, prevention, and control of cancer offer hope that these goals will eventually be achieved.

References

- 1. DeVita VT Jr, Hellman S, Rosenberg SA (eds). Cancer: Principles & Practice of Oncology. 4th Edition. Philadelphia, J. B. Lippincott Company, 1993.
- 2. Baseler MW, Urba WJ. Immunologic monitoring and clinical trials of biological response modifiers. Cancer Chemotherapy and Biological Response Modifiers 13:231-250, 1992.
- 3. Clark JW. Biological response modifiers. *Cancer Chemotherapy and Biological Response Modifiers* 13:175-190, 1992.
- 4. Hillman GG, Haas GP, Wahl WH, Callewaert DM. Adoptive immunotherapy of cancer: Biological response modifiers and cytotoxic cell therapy. *Biotherapy* 5:119-129, 1992.
- 5. Abbas AK, Lichtman AH, Pober JS. Cellular and Molecular Immunology. Philadelphia, W.B. Saunders Company, 1991.
- 6. DeVita VT Jr, Hellman S, Rosenberg SA (eds). *Biologic Therapy of Cancer*. Philadelphia, J. B. Lippincott Company, 1991.
- 7. Janik JE, Longo DL. Biological response modifiers in breast cancer. *Comprehensive Therapy* 17(12):28-31, 1991.

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|---|----------------------------|--|----------|
| ADA Transduced T-ce | 11 | Autologous Lymphocytes | 635386 |
| ADA TT all succed 1 - Ce | • | Natorogous Lymphocytes | 373364 |
| Allogeneic cells | | Allogeneic therapy, Allogeneic bone marrow transplantation, Bone marrow therapy, Killed tumor cells | |
| Anticea, I-131 | | Antiferritin antibody | |
| Anti-Thymocyte Glob | ulin | | |
| | | | |
| <u>Asparaginase</u> (Escherichia coli) | L-ASP, ASP, ASPA, L-ASE | Asnase, L-Asparaginase, L-Asparagine amidohydrolase, Colaspase, Crasnitin#, EC 3.1.5.1, Elspar, Leunase (enzyme) | 109229 |
| Asparaginase (Erwinia carotovora | L-ASP) | L-Asparaginase, Porton asparaginase (enzyme) | 106977 |
| Asparaginase-PEG (K | -H) | | 644954 |
| Asparaginase-PEG | | (Merck) | 624239 |
| Autologous | | Autol Tumor-Trnsdcd/Gene for IL-2 (GT1), Autol Tumor-Trnsdcd/Gene for TNF (Cetus), Autol Tumor-Trnsdcd/Gene for TNF (GTI), | 373364 |
| | | Autol Tumor-Trnsdcd/Gene for TNF-COMPA (Cetus), Autologous Tumor Cell (BCG) + | 624355 |
| | | <pre>IL-2 (Bionetics), Autologous Tumor Cell (BCG + IL-2 (Cetus), Autologous Tumor Cell Vaccine + IL-2 (Cetus)</pre> | 373364 |
| | | Autologous Tumor Cell Vaccine + BCG (Bionetics) | 624335 |
| Azimexon | AMX | 2-aziridinecarboxamide compound (synthetic chemical, immunomodulator) | |
| BACI | | Bovine anti-Cryptosporidium Immunoglobulin | |
| 8CG | | Bacillus Calmette-Guerin (Bacterial and Fungi, Mycobacteria biological) (Mycobacterium bovis) (immunomodulator) | |
| | | • Connaught | 614389 |
| BCG (Cont'd) | | • Pasteur | B116328 |
| | | • Tice | 614388 |
| | | • TheraCys | |
| BCG | MER | Methanol extracted residual of BCG (Bacteria & Fungi Mycobacterial biological) | 143769 |
| Bestatin | BST | (Bacteria & Fungi Streptomyces biological) (immunomodulator) | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC</u> No. |
|----------------------------------|-------------|---|------------------|
| | | | |
| Beta-Carotene | | Solatene (ultraviolet screen) | |
| Biostim | | RU 41740 | |
| Bromocryptine | | Bromoergocryptine | |
| C-parvum | C.p., CPAR | Coparvax (R), (Merrieux) Corynebacterium parvum, H37Ra (immunomodulator) | 197213 |
| | | Corynebacterium parvum, Wellcome CN6134 (BW) (Bacteria & Fungi biological) | 220537 |
| | | Corynebacterium granulosum (Bacteria & Fungi biological) | |
| CGP 19835A | MTP-PE | | |
| Copovithane | CPV | | |
| Cyclosporine | | Cyclosporin#, 27-400 | |
| <u>Danazol</u> | | Chronogyn#, Danocrine | |
| Dehydroemetine | | | |
| Detox | | | |
| <u>Dimethyl</u> <u>sulfoxide</u> | DMSO | Demasorb#, Demavet, Demeso, Demsodrox, Dermasorb, Dimexide, DMS-70, DMS-90, Dolicur, Dolisur, Domoso, Dromisol, Durasorb, Gamasol 90, Hyadur, Infiltrin Methyl sulfoxide, Somipront, SQ 9453, Sulfinylbismethane, Syntexan | 000763 |
| Dimethylbusulfan | | | |
| Dinitrochlorobenzen | e | | |
| Dinitrofluoro- benzene | DNFB | | |
| Erythropoietin | EPO | Epogen, <u>Epoietin Alfa</u> , Eprox, rHu-EPO, Procrit (stimulates RBC production) | 628281 |
| Ethylchlorformate | | | |
| <u>Filgrastim</u> | | Neupogen, r-met Hu G-CSF | |
| Freund's adjuvant | FCA | (Bacteria & Fungi, Mycobacteria biological) | |
| Gallium Nitrate | GAN | Gallium trinitrate, Ganite | 015200 |
| Ganglioside | GM2 | | |
| HLTV antibody | | | |
| 41-Hydroperoxycyclo | phosphamide | | |

| Generic Name | Short Names | <u>Synonyms and</u> <u>Brand or Trade Names</u> | + <u>MSC</u> <u>Mo</u> . |
|------------------------------------|-----------------|---|--------------------------|
| | 474 | | |
| I 131 Imexon | 131 1 | Antiferritin antibody (see Anticea) 1,3-Diazabicyclo (3.1.0.) Hex-3-en- | |
| Immune RNA | | 2-one, 4-amino (synthetic chemical) (Mammalian biological product) (immunomodulator) | |
| Imuvert | | (| |
| Interferon and Interferon induc | IF, IFN cers | Human fibroblast interferon Human leukocyte interferon (Meloy) IFN: Leukocyte (Meloy) | 335044 |
| | | Human leukocyte interferon (W-L) | 340855 |
| | | Human leukocyte interferon, Beta (Mochida) | 605606 |
| Interferon Alfa-n | 1 | Human lymphoblastoid interferon (BW) Wellferon | 339140 |
| Interferon Alfa-n | <u>3</u> | Alferon N Injection, Leukocyte Interferon#, Alferon Rec Gamma (Biogen) | |
| Interferon | | Immune interferon (Meloy), IFN: Immune | 354655 |
| | | IFN: Rec Gamma (Schering) | 609473 |
| Interferon Alfa-2 | <u>e</u> | Recombinant Leukocyte A interferon, Roferon, Roferon-A, Ro 22-8181 (HLR) | 367982 |
| Interferon Alfa-2 | <u>Þ</u> | Recombinant Alpha 2 interferon, Betaseron, Recombinant interferon Gamma, (B), Gamma Interferon, Intron A (Recombinant analogue, Bacterial) IFN: Rec Gamma (Biogen) | |
| Interferon Gamma- | <u>16</u> | Recombinant interferon Gamma (G), Immune Actimmune (Genentech) | 600662 |
| Interleukin-1 | | <pre>IL-1 Beta (Syntex), Interleukin-1 Beta IL-1 Beta (Syntex) + Il-2 (Cetus)</pre> | 628282 |
| | | Il-1 alpha, Interleukin-1 alpha (Dainìppon) | 621381 |
| | | Il-1 alpha (Immunex) | 640032 |
| Interleukin-2 | IL-2, TCGF | Human Jurkat Tumor Derived, (Dupont) Il-2 Jurkat Derived | 364831 |
| | | IL-2 Recombinant Interleukin-2 | 373364 |
| | | IL-2: Rec (Cetus), Aldesleukin, Proleukin | |
| | | IL-2: Rec (Shionogi), S6820 | |
| | | IL-2: Rec (Cetus) + TIL | |
| | | IL-2 (CT)/MDELAK | |
| Interleukin-2 | | IL-2 /Adherent LAK (CT) | |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>NSC Mo</u> . |
|---|-------------|--|-------------------|
| | | | |
| | | Aut Tum Vac/IL-2/BCG | 373364 |
| | | Aut Tum Vac/BCG/Il-2 | |
| | | Aut Tum Vac/Il-2 /BCG, Tice BCG Vaccine, frozen | 624335 |
| | | IL-2 (CT)/TIL (BiorExp) | |
| | | IL-2: Rec.(HLR) | 600664 |
| | | IL-2/TIL (HLR) | |
| | | IL-2/AutoEduLymph (HLR) | |
| | | IL-2 Recombinant | |
| | | IL-2 (HR) | |
| | | IL-2 (HR)/OKT 3-LAK | 600664 |
| | | IL-2 Polyethylene Glycol (Peg IL-2) | 625376 |
| | | IL:-2 (HR)/OKT 3-LAK | 600664 618843 |
| | | IL-2 Natural (Collaborative) | 600663 |
| | | IL-2 + IL-4 (Cetus & Sterling) | 373364 |
| | | IL-2 + IL-4 (Cetus & Sterling) | 620211 |
| | | IL-2 + IL-4 HLR/STRLNG | 600664 |
| Interleukin-2 lipo | osome | | |
| Interleukin-3 | | IL-3 + GM-CSF | 643496 |
| | | IL-3 + GM-CSF (Sandoz) | 641115 |
| Interleukin-4 | | IL-4, Interleukin-4 (Immunex) Eastman Kodak | 620611 620211 |
| | | IL-4 (K)/IL-2 (CT) | 620211 373364 |
| | | IL-4 (K)/IL-2 (HR) | 620211 |
| Interleukin-6 | | (Sandoz) | 600664 643497 |
| Ipomeanol | | 4-Ipomeanol | 349438 |
| IVIG L A K cells | | Gammaguard (Immune globulin) Lymphokine activated killer cells | |
| Lentinan | | (Bacteria & Fungi Lintinus Edodes biological) (immunomodulator) | |
| <u>Levamisole</u> <u>HCl</u> * Meg-CSF | LEV | Citarin L, Decaris, Ergamisol, Ketrax, Levomysol HCL, R-12,564, Ripercol L, L-Tetramisole HCl, Tetramizole, Tramisol, Tramisole (Synthetic chemical, immunomodulato | 177023 or) |

Synonyms and Brand or Trade Names

Short Names

Generic Name

+NSC No.

| Monoclonal Antibody MoAB | | |
|--|---|--------|
| | developed. Therefore, they will not be | |
| MVE - 2 | Maleic anhydride divinyl ether copolymer (immunomodulator) ¹ | 332586 |
| N2-T1L/ADA Gene | Adenosine Deaminase Transduced Autologous T-lymphocytes | 635386 |
| N2 Transduced TIL | N2 Retroviral Vector with Neo(R) Gene Insert | 622283 |
| Nocardia rubra N-CWS | Cell wall skeleton (Bacteria & Fungi biological) (immunomodulator) | |
| Octreotide Acetate | Sandostatin (Reduces secretory diarrhea) | |
| OK-432 | Beta hemolytic Streptococci (Bacteria & Fungi Streptococcus biological) | |
| Ovine Sialomucin | (immunomodulator) | |
| <u>Pegaspargase</u> | (Enzon) | |
| PEG-IL-2 | (CT) | 625376 |
| Pentostatin DCF, 2'-DCF | CI 825, CL-67310465, Co-Vidarabine, Deoxycoformycin, 2'-Deoxycoformycin, Nipent | 218321 |
| Poly AU | Polyadenylic-polyuridylic acid, Poly A-poly U | |
| 51-Inosinic acid Poly I:C polymers | Poly(rI).poly(rC), Poly I:C, Polyribocytidylic-polyriboinosinic acid | 120949 |
| Poly I:Poly C with Poly IC/LC Poly-L-lysine stabilizer | Polyinosinic-Polycytidylic acid stabilized with Poly-L-lysine and carboxymethylcellulose (Interferons and Interferon inducers) | 301463 |
| Poly I: Poly C12U | Ampligen | |
| Protein A Staphylococcus Aureus | Cowan I | |
| Pseudomonas aeruginosa | Heptavalent lipopolysaccharide (Bacterial & Fungi biological) | |
| PSK | (Bacteria & Fungi biological) | |
| Pyran copolymer | Divema, Pyran, Pyran XA 124-177, XA 146-85-2 | 046015 |

¹Drugs previously known as immunotherapy drugs are now being considered biological response modifiers.

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No. |
|-----------------------------------|-------------|---|----------------------------|
| | | | |
| Recombinant Soluble CD4 | | rCD4 | |
| Recombinant Soluble T4 | | sT4 | |
| Recombinant Tumor Necrosis Factor | | rTNF (produced by recombination) | |
| Sargramostim | | Leukine, Prokine (Recomb. granulocyte-macrophage colony-stimulating factor) (reduces duration and impact of neutropenia) | |
| Thymosin | Alpha 1 | | 337793 |
| Thymosin | Fx5 | Numan thymosin-fraction 5 | 350391 |
| | | Numan thymosin-fraction Alpha 1 (Mammalian biological product) (Thymosins) | 337793 |
| Tilorone | | (Synthetic Chemical) (Immunomodulator and Interferon inducer | |
| TIL/TNF | | Retroviral vector carrying gene coding for tumor necrosis factor (PA 317/LTNFSN) (Cetus) | 636228 |
| | | <pre>TlL transduced with NEOR Gene IL-2 (NHLBI)</pre> | 622283 |
| | | TIL transduced with NEOR Gene IL-2/IL (Cetus/Sterling/GTI) | 622283 |
| | | TIL transduced with NEOR Gene IL-2/IL-4 (Cetus/Sterling) | 373364 |
| | | TIL transduced/Gene coding for TNF (Cetus) | 636228 |
| | | TIL transduced/Gene coding for TNF | 373364 |
| | | Tumor Primed Anti-CD3 activated lymphocytes | 618843 |
| TNF Transd TIL (GTI) | | Tumor infiltrating lymphocytes transduced with the tumor necrosis factor gene (Retrovirus vector TNF-NeoR-GTI) | 637656 373364 |
| Tumor necrosis factor | TNF | (Cetus) (Genentech) (Knoll) | 606515 604175 635257 |
| TP OKT 3 Lymphoct | yes | | 618843 |
| Tumor primed Anti-CD3 | | Activated lymphocytes | 618843 |
| Vibrio cholera neuraminidase | VCN | Malignant cell antigens, Mammalian biological product | |

NOTES

Section 5

HORMONES AND AGENTS ACTING VIA HORMONAL MECHANISMS

Hormones are substances produced by the body which help regulate body mechanisms such as growth, metabolism, and reproduction. Used in the treatment of cancer, administration of hormones, withdrawal of hormones, or interference with hormone function may alter the growth of hormone-responsive malignant neoplasms. Cancer therapy which achieves its anti-tumor effect through changes in hormonal balance is often called endocrine therapy. This includes the administration of hormones, agents acting via hormonal mechanisms, antihormones, steroids, and surgery or radiotherapy directed at hormone-producing tissue.

Cancers in which endocrine therapy causes tumor regression are often referred to as hormone-responsive. In general, hormone-responsive cancers derive from tissue which require hormones for normal development (e.g., breast and prostate cancers). However, some cases of other cancers, particularly melanoma and hypernephroma, may also respond to endocrine therapy.

In recent years, laboratory methods have been developed to identify the presence of specific hormone receptors on the surface of normal and malignant cells. The presence of hormone receptors (particularly of estrogen and progesterone receptors in breast cancer) may predict the response of the patient's tumor to endocrine therapy. A positive response to endocrine therapy observed in some cases of receptor-negative tumors suggests that hormonal agents may exert their effects through other pathways in addition to direct action on hormone receptors of cancer cells.

Hormones

Synthetic hormones may be administered in **pharmacologic** doses which achieve hormone levels far above the natural (**physiologic**) hormone levels, to alter hormone balance in the patient and modify the growth of susceptible cancers. These compounds are usually classified as estrogens, progestins, androgens, and corticosteroids.

Administration of estrogen, usually the oral estrogen diethylstilbestrol, has been a standard therapy for advanced prostate cancer since the 1940's. In men, pharmacologic doses of estrogen suppress luteinizing hormone (LH) release by the pituitary gland and result in decreased testosterone production (see below). This "androgen withdrawal" effect produces a clinical response in most patients with previously untreated disease that is comparable to the therapeutic effect associated with orchiectomy.

Hormone manipulation can provide effective control of breast cancer for many women with slow-growing metastatic disease, particularly in soft tissue or skeleton. Hormone receptor levels in breast cancer tissue are useful predictors of response in breast cancer patients: up to 75% of women with tumors positive for both estrogen receptors and progesterone receptors will respond to endocrine therapy.

Common approaches to hormone therapy for breast cancer comprise the oral administration of pharmacologic doses of an estrogen, progestin, or androgen as a single agent. Patients who have a good initial response to additive hormone therapy but develop disease progression on treatment may experience a second response when the initial agent is discontinued. Breast cancer patients with a

good response to initial hormone therapy often respond to a second or subsequent hormonal approach and may achieve long-term disease control by continuing through a series of hormonal treatments.

Corticosteroids are hormones produced by the adrenal gland which have actions in metabolism and immune modulation. These agents have antitumor activity in lymphomas and in lymphoid leukemias and are included in most treatment regimens for these diseases. Corticosteroids may be used for hormone replacement following adrenalectomy (see below) and in the management of clinical complications associated with malignancy (e.g., hypercalcemia, brain metastases). Steroids (particularly decadron) are often used as antiemetics prior to chemotherapy and may be used in the supportive care of patients with advanced disease to palliate the symptoms of general debility.

Agents acting by Hormonal Mechanisms

Hormone-Releasing Factors

The production and/or release of many hormones is governed by hormone-releasing factors from the pituitary gland. This frequently involves a feed-back loop, so that balance is achieved between the hormone-releasing factor and its target hormone. For example, thyroid-stimulating hormone (TSH, thyrotropin) from the pituitary gland stimulates release of thyroid hormone by the thyroid gland, and increased levels of thyroid hormone result in decreases in TSH.

Drugs have been synthesized which mimic (agonists) or inhibit (antagonists) the actions of specific hormone-releasing factors. Agonists for gonadotropin-releasing factor (LHRH, GnRH), including leuprolide (Leupron), buserelin, and goserelin (Zoladex), have clinical applications in the management of breast and prostate cancers. Normally, LHRH from the hypothalamus controls release of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the pituitary; FSH and LH regulate circulating levels of natural estrogens, progestins, and androgens. Chronic treatment of men with high doses of LHRH agonists results in decreases in testosterone and dihydrotestosterone similar to those observed following castration. Premenopausal women treated with LHRH agonists have estrogen and progesterone levels that are reduced to those of postmenopausal women. This dramatic reduction in circulating hormones is thought to be responsible for the antitumor effects of LHRH agonists in breast and prostate cancer.

Somatostatin is a peptide hormone produced by the hypothalamus which inhibits the release of pituitary hormones (e.g., growth hormone, prolactin, thyrotropin). It also suppresses pancreatic function and inhibits gastrointestinal secretions. Somatostatin (Sandostatin) has been used to treat a broad spectrum of secretory tumors of the gastrointestinal tract, including vipomas, carcinoid syndrome, glucagonomas, insulinomas, gastrinomas, and APUD tumors.

Hormone-Synthesis Inhibitors

One approach to hormone regulation is to block the production of the hormone by inhibiting the actions of enzymes which participate in its synthesis. Most estrogen in postmenopausal women is produced outside the ovary by enzyme actions on steroid substances made by the adrenal gland. The drug aminoglutethimide blocks the action of several enzymes in this pathway, resulting in decreased production of adrenal steroids ("medical adrenalectomy"), especially the androstenedione which is converted to estrogen.

Aminoglutethimide has been used in the management of hormone-responsive breast cancer, usually after relapse during other endocrine therapies. Because the drug interferes with the production of glucocorticoids and mineralocorticoids necessary for normal metabolism and body function, patients who receive aminoglutethimide must also take steroid replacements on a daily basis to avoid developing adrenocorticoid insufficiency (Addison's disease) and orthostasis (see below).

Antihormones

The identification of specific hormone receptors on cancer cell surfaces offered new approaches to regulate the actions of hormones in promoting growth of hormonally-responsive tumors. Since the early 1970's, "antihormone" agents have been developed that interfere with the interaction between the hormone molecule and its receptor site that stimulates cell growth and proliferation.

Tamoxifen (Novaldex), the first antiestrogen to achieve widespread clinical use, has become standard first-line hormonal therapy for metastatic breast cancer and for adjuvant therapy of early stage breast cancer in postmenopausal women. Clinical trials are currently underway to test the efficacy of tamoxifen as therapy of ductal carcinoma-in situ (DCIS) and as a chemopreventive agent in women at high risk of breast cancer.

Tamoxifen acts by attaching to the cell surface estrogen receptor without initiating the intracellular processes which lead to proliferation. Its presence at the receptor site blocks the attachment of natural estrogens and thus interferes with their actions on breast cells and breast cancer cells. Tamoxifen also stimulates the production of growth factors that further modulate the proliferation of breast tissue.

While tamoxifen has antiestrogen actions on breast tissue, it has estrogen-like activities at other organ sites. Women receiving tamoxifen as adjuvant therapy have benefits similar to those of estrogen replacement therapy in lowering blood lipids and slowing bone loss. However, they also have potential side effects associated with estrogen therapy such as endometrial abnormalities and risk of thromboembolism. Other antiestrogens are currently under development that have pure antiestrogen effects (e.g., toremifene) or that may be used in tumors that have become resistant to tamoxifen.

The recent development of antiandrogens has expanded the options for medical management of advanced prostate cancer. The drug flutamide attaches to androgen receptor of prostate cancer cells and blocks the actions of testosterone and dihydrotestosterone in stimulating cell growth. Flutamide may be used as a single agent for initial hormonal treatment of advanced disease, or it may be given with an LHRH agonist (or surgical orchiectomy) as "combined androgen blockade." Casodex and nilutamide, other antiandrogens, are currently in preclinical and early clinical studies.

Surgery and Radiotherapy to Hormone-Producing Tissues

Surgical procedures to remove hormone-producing tissues were among the first attempts at endocrine therapy of cancer (e.g., oophorectomy or adrenalectomy for breast cancer, orchiectomy for prostate cancer). Radiotherapy to hormone-producing organs (e.g., ovaries) has been used to suppress hormone production in patients who could not withstand surgery.

The use of drugs to suppress hormone production has, to a great extent, replaced surgical approaches to endocrine therapy of breast cancer: LHRH agonists for "medical castration", and aminoglutethimide for "medical adrenalectomy." However, orchiectomy remains a standard approach

to endocrine management because of the simplicity of the procedure, its immediate effect, and the lack of side effects associated with drugs used for endocrine therapy of this disease.

Hormone Replacement in Patients with Cancer

Tumor involvement or specific cancer-directed therapy may destroy tissue that produces hormones necessary for normal metabolism and body function. In these clinical situations, hormone replacement therapy is necessary to maintain life. For example, patients receiving aminoglutethimide for breast cancer have suppressed production of glucocorticoids and mineralocorticoids which function in glucose metabolism and fluid and electrolyte balance, in addition to the desired suppression of androstenedione production. Thus patients receiving aminoglutethimide must also take a glucocorticoid (hydrocortisone) and may also require a mineralocorticoid (Florinef) to replace the natural corticoids normally produced by the adrenal gland. In contrast to endocrine treatment of cancer, hormones administered as replacements are given in physiologic doses rather than at the higher, pharmacologic doses used for tumor-directed therapy.

Thyroid hormone replacement is an exception because it inhibits pituitary production of thyroid stimulating hormone (TSH) which could stimulate tumor growth. It should be coded as hormonal therapy.

| Generic Name | Short Names | <u>Synonyms and</u> <u>Brand or Trade Names</u> | + <u>NSC No</u> . |
|----------------------------------|--------------|---|-------------------|
| Adrenocortico- tropic hormone | ACTH | Acethropan, Acthar, Adrenal Cortex hormone, Adrenocorticotrophin, Adrenocorticotropin, Corstiline, Corticotrophin, Corticotrophin*, Cortrophin*, Depo-ACTH* | 025933 |
| Algestone* Acetophe | <u>enide</u> | Deladroxone#, Droxone#, SQ 15,101 (progestin) | 067831 |
| Amadinone Acetate | | RS-2208 (progestin) | |
| <u>Amcinonide</u> | | CL-34699, Cyclocort (glucocorticoid) | |
| Aminoglutethimide* | AG, AGT | Ba-16038, Cytadren, Elipten (adrenocortical suppressant) | 330915 |
| Anagestone Acetate | | Anatropin# (progestin) | 073880 |
| Beclomethasone Dip | ropionate* | Beclometasone, Beclovent, Beconase, Sch-18020W, Vancenase, Vanceril (glucocorticoid) | |
| Benorterone# | | SK&F-7690 (anti-androgen) | |
| Benzestrol | | Estrogen | 408889 |
| Betamethasone* | | Celestone, Sch-4831 (glucocorticoid) | 039470 |
| Betamethasone Acet | ate . | Component of Celestone Soluspan (glucocorticoid) | |
| Betamethasone Benz | oate* | Benisone#, Uticort, W-5975 (glucocorticoid) | |
| Betamethasone Dipr | opionate* | Component of Alphatrex, Diprolene, Diprosone, Component of Lotrisone, Psorion, Sch 11460 (glucocorticoid) | 206525 |
| Betamethasone Sodi Phosphate* | <u>um</u> | B-S-P, Component of Celestone Soluspan (glucocorticoid) | 090616 |
| Betamethasone Vale | rate* | Component of Betatrex, Beta-Val, Valisone (glucocorticoid) | 206540 |
| Bolasterone# | | Myagen#, U-19763 (anabolic) | 066233 |
| Busrelin Acetate | | Hoe 766, Suprafact | |
| Calusterone | CAL, CLS | Dimethyltestosterone, Methosarb#, U-22,550 (androgen) | 088536 |
| Carbenoxolone Sodi | <u>um</u> | (Biorex, England) | 281727 |
| Casodex | | Anti-androgen | |
| Chlormadinone* Acetate | CAP | Bovisynchron, Chlormadinonu (Polish), Clordion, Gestafortin, Lormin, Lutinyl, Matrol, Synchrosyn, Synchrosyn P (progestin) | 092338 |

| Generic Name Short Names | Synonyms and Brand or Trade Names | + <u>NSC</u> No. |
|---|--|------------------|
| <u>Chlorotrianisene*</u> | Anisene, Chlorotrianisine, Chloro- tris, Chlortrianizen, Clorestrolo, Clorotrisin, Hormonisene, Khlortri- anizen, Merbentul, Metace, Rianil, TACE (estrogen) | 010108 |
| Chromic Phosphate P 32 | Chromphosphotope#, Phosphocol P 32 | |
| <u>Cingestol</u> * | Progestin | |
| Clocortolone Acetate | SH 818 (glucocorticoid) | |
| Clocortolone Pivalate* | Clocortolone trimethyl acetate, SH 863 (glucocorticoid) | |
| Clogestone* Acetate | AY-11,440 (progestin) | |
| Clomegestone* Acetate | SH 741 (progestin) | |
| Clometherone# | 38000 (anti-estrogen) | |
| Clomiphene* Citrate) | Cis-Clomiphene citrate, Clomiphene A citrate, Serophene, Zuclomiphene citrate | 151466 035770 |
| Corticotropin* | See Adrenocorticotropic hormone | |
| Cortisone Acetate* | Cortisone, Cortogen acetate#, Cortone acetate, Component of Neosone (glucocorticoid) | 009703 |
| Cortivazol# | Glucocorticoid | 080998 |
| Cosyntropin* | Cortrosyn (adrenocorticotropic hormone) | |
| Cyproterone Acetate | SH 714 (anti-androgen) | 081430 |
| Decapeptyl | D-TrP6 (LH-RH) | |
| Delmadinone Acetate | RS-1301 (progestin, anti-androgen, anti-estrogen) | |
| Descinolone Acetonide# | Cl-27,071 (glucocorticoid) | 044827 |
| Desoxycortico-DCA, DOCA, sterone Acetate DOXO | Cortate acetate, Decortin, Decosterone, Decostrate#, Doca acetate#, Dorcostrin#, Percorten acetate#, Percotol, Primocort, Primocortan, Syncort (adrenocortical steroid) | 009567 |
| Desoxycorticosterone Pivalate* | Percorten pivalate# (adrenocortical steroid) | 095278 |
| <u>Dexamethasone</u> *∇ DECA, DM, DSM, | Aeroseb-Dex, Decaderm, Decadron Tablets, Decaspray, Component of Deronil, Component of Dexacidin, Dexasone, Dexone, Hexadrol Elixir, Hexadrol Tablets, Maxidex, Component of Maxitrol, Component of Tobradex (glucocorticoid) | 345211 |
| <u>Dexamethasone</u> DXM <u>Acetate</u> * | Dalalone D.P., Dalalone L.A., Decadron-LA (adrenocortical steroid) | |

[∇] If used in the treatment of the malignancies mentioned in the introduction on p. 37 (bold), code as hormonal therapy. For all other cancers, consider ancillary unless specifically stated to be hormonal therapy.

Synonyms and +NSC No. Generic Name Short Names Brand or Trade Names ST 12 Dexamethasone <u>Dipropionate</u> Dexamethasone Sodium Phosphate* Dalalone, Hexadrol, Component of NeoDecadron (glucocorticoid) 059809 Dienestrol* Component of Estan#, Synestrol# (estrogen) Diethylstilbestrol* DEB, DES, 003070 Bio-Des, Comestrol, Cyren A. Di-Estryl, Domestrol, Estilbin, Estrobene, Etrosyn, Fonatol, Graf-estrol, Hi-Bestrol, Nicroest, Nilestrol, NCO Oestrogenine, Serral, Sexocretin, Sibol, Stil-Rol, Stilbestrol, Stil-betin#, Stilboestroform, Stilkap, Synth-oestrin, Syntofolin, trans-Diethylstil-bestrol, Vagestrol# (estrogen) Diethylstilbestrol 010481 Fosfestrol, Stilphostrol (estrogen) Diphosphate* Diethylstilbestrol Dipropionate Dibestil# (estrogen) Diflucortolone* Glucocorticoid Diflucortolone* Pivalate SH-968 (glucocorticoid) NDHT Drolban#, Drostanolone propionate, Emdisterone, Masterid, Masteril, 012198 <u>Dromostanolone</u> <u>Propionate</u> Masterone, Nedrotestron propionate, Permastril, 32379 092336 Dydrogesterone* Diphaston, Dufaston, Duphaston, Duvaron, Gestatron, Hydrogesterone, Isopregnenone, Prodel, Retrone, Terolut (progestin) <u>Equilin</u> 7-Dehydroestrone (estrogen) 010971 Estradiol* Component of Androgyn L.A., Aquadiol#, 009895 Diogyn, Diogynets, Estrace, Estraderm TTS, Progynon# (estrogen) Estradiol Benzoate 009566 Progynon B (estrogen) Estradiol Cypionate* DepGynogen, Depo-Estradiol. Component of Depo-Testadiol (estrogen) Estradiol Dipropionate Ovocyclin dipropionate# (estrogen) 003354 Estradiol Mustard 112259 Estradiol Valerate* Component of Deladumone, Delestrogen, 017590 Component of Deluteval 2X#, Component of Ditate#, Gynogen L.A. (estrogen) Leo 275, Ro 21-8837, Ro 22-2296/000 (chemotoxic hormone) Estramustine 089201

Synonyms and Generic Name : Short Names Brand or Trade Names +NSC No. Emcyt, Estracyt, 080100 Estramustine Phosphate Sodium* Estramustine phosphate disodium, Ro 21-8837/001 (chemotoxic hormone) <u>Estriol</u> 012169 Theelol# (estrogen) Estrogens, Conjugated* Component of Cyclogesterin#, Component of Milprem, Component of PMB-200, Component of PMB-400, Component of Premarin with Methyltestosterone, Premarin Estrogens, Esterified* Amnestrogen#, Estratab, Component of Estratest, Menest, Component of Menrium (palliative) 009699 Theelin (estrogen) Estrone* Estropipate* Ogen, Component of Ortho-Cyclen, Piperazine estrone sulfate#, Sulestrex# (estrogen) 010973 Ethinyl Estradiol* Component of Brevicon, Chee-O-Gen, Chee-O-Genf, Component of Demulan, Diogyn E, Dyloform, Esteed, Estigyn, Estinyl, Eston-E, Estoral, Ethidol, Ethinoral#, Eticyclin, Eticylol#, Etinestrol, Feminone, Ginestrene, Halodrin, Inestra, Component of Levlen, Component of Loestrin, Component of Lo/Ovral, Component of Lynoral#, Menolyn, Microfollin, Component of Modicon, Component of Nordette, Component of Norethrin 1/35E, Component of Norlestrin, Novestrol, Oradiol, Orestralyn, Palonyl Perovex, Spanestrin (estrogen) Florinef Acetate (adrenocorticoid steroid), Fludrocortisone Acetate (salt-regulating) 054702 Flumethasone* U-10,974 (glucocorticoid) 033001 Cortilet, Delmeson, Fluoromethalone, Fluorometholone* FML Forte, FML Liquifilm, Component of FML-S Liquifilm, FML S.O.P., Component of Neo-Oxylone#, Oxylone#, U-8614 (glucocorticoid) 012165 FLU, FXM, HAL Fluoxymesterone* Androfluorene, Anrosterolo, Fluotestin, Fluoximesterone, Flusteron, Flutestos, Halodrin, Halotestin, Neo-Ormonal, Oralsterone, Oratestin, Ora-Testryl#, Testoral, Ultandren (androgen; palliative) Methral, P-1742 (glucocorticoid) 067756 Fluperolone Acetate 047439 Alphadrol#, U-7800 (glucocorticoid) Fluprednisolone* Glucocorticoid Fluprednisolone* Valerate# Cordran, Flurandrenolone#, 33379 Flurandrenolide* (glucocorticoid)

SC-9880 (progestin)

Flurogestone Acetate

065411

| Generic Name | Short Names | Synonyms and Brand or Trade Names | +NSC No- |
|--------------------------------------|---------------------|---|----------|
| | | | |
| Flutamide* | | Eulexin, Niftholide, Niftolide, Sch 13521 | 147834 |
| <u>Formocortal</u> | | Deflamene, Fluderma (glucocorticoid) | 150527 |
| Gestaclone | | SH-1040 (progestin) | |
| <u>Gosrelin</u> | | Decapeptide 1, ICI 118,630, Zoladex (LHRH agonist) | 606864 |
| Gestonorone Caproate | | SH-582 (progestin) | 084054 |
| Haloprogesterone | | Progestin | |
| Hexestrol | | Cycloestrol, Estrifar, Estronal, Extra-Plex, Hexanoestrol, Hexestro- fen, Hexoestrol, Hormoestrol, Sinestrol, Synestrol, Synoestrol, Synthovo, Syntrogene | 009894 |
| <u>Hydrocortisone</u> * | нс | Acticort, Aeroseb-HC, Cetacort, Cort-Dome, Cortef, Cortenema, Cortril, (glucocorticoid) | 010483 |
| Hydrocortisone Ace | tate* | Cortef acetate, Cortril acetate-AS (glucocorticoid) | 000741 |
| <u>Hydrocortisone Sod</u> | ium Phosphate* | Hydrocortone Phosphate (glucocorticoid) | |
| Hydrocortisone Sodium Succinate | HDC | A-hydroCort, Solu-Cortef (glucocorticoid) | |
| Hydroxyprogesteron | e <u>Caproate</u> * | Delalutin#, Component of Deluteval 2X#, Gestoral LA 250, Prodrox (progestin) | 017592 |
| <u>Ketoconazole</u> * | | Nizoral, R-41,400 | |
| Leuprolide Acetate | * | Abbott 43818, Lupron TAP, Lupron Depot (TAP), Leuprorelin, TAP-144 | 377526 |
| <u>Levothyroxine</u> Sodi | <u>um</u> | Levothroid, Synthroid (thyroid hormone) | |
| <u>Liothyronine Sodiu</u> | in [±] | Basoprocin, Cynomel, Cytomel, Ibiothyron, Tertroxin, (thyroid hormone) | 080774 |
| <u>Liotrix</u> | | Euthroid, Thyrolar (thyroid hormone) | |
| <u>Lynestrenol</u> # | | Ethynylestrenol, Exluton, Linestrenol, Org 485-50, Orgametril#, Orgametrol, Ovoresta M (progestin) | 037725 |
| Medrogestone | | AY-62022 (progestin) | 123018 |
| Medroxypro- gesterone Acetate* | DMPA, MAP, MPA | Curretab, Cycrin, Depo-Provera, Farlutin, Gestapuran, Lutopolar, Lutoral, Oragest, Perlutex, Prodasone, Progestal, Provera, Proverone, Repromap, Repromix, Sirprogen, U-8839, Veramix (progestin) | 026386 |
| Medrysone* | | Medrocort#, U-8471 (glucocorticoid) | 063278 |

| Generic Name | Short Names | Synonyms and Brand or Trade Names | + <u>MSC</u> <u>No</u> . |
|----------------------------------|------------------|---|--------------------------|
| Megestrol Acetate* | | BDH-1298, Megace, SC-10363, 5071 (progestin) | 071423 |
| Melengestrol Acetate | MGA | BDH-1921, 5373 (antineoplastic progestin) | 070968 |
| Meprednisone* | | Betapar#, Sch-4358 | 527579 |
| Mesterolone | | SH-723 (androgen) | 075054 |
| Mestranol* | | Component of Enovid, Component of Norethin 1/50 M, Component of Norinyl, Component of Norquen, Component of Ortho-Novum, Component of Ovulen (estrogen) | |
| Methandros- tenolone | MA | Abirol, Anabolin, Compound 17309, Crein, Danabol, Dehydromethyltestosterone, Dianabol#, Geabol, Metanabol, Metandienon, Metandrostenolone, Metasol, Methandronolone, Naposim, Nerobol, Protobolin, Stenolon (androgen) | 042722 |
| Methylprednisolone* | , | Dopmedrol, Medrate, Medrol, Medrone, Promacortine, Urbason (glucocorticoid) | 019987 |
| Methylprednisolone | <u>Acetate</u> * | DepMedalone, Depo-Medrol, Component of Neo-Medrol (glucocorticoid) | 048985 |
| Methylprednisolone Phosphate | <u>Sodium</u> | Medrol Stabisol#, U-12,019E (glucocorticoid) | 048989 |
| Methylprednisolone Succinate* | <u>Sodium</u> | A-methaPred, Solu-Medrol (glucocorticoid) | |
| Methyltestosterone* | • | Andrometh, Androsan, Androsten, Anertan, Dumogran, Component of Estratest, Homandren, Malestrone, Metandren, Metrone, Neo-Hombreol-M#, Oreton Methyl, Component of Premarin with Methyl- testosterone, Steronyl, Synandrets, Synandrotabs, Testoviron, Testred (ICN) (androgen) | 009701 |
| Methynodiol Diaceta | ate# | SC-19198 (progestin) | |
| Mibolerone | | Miboleron, U-10,997 (anabolic; androgen) | 072260 |
| Mifepristone | | RU 486 | |
| Nafarelin Acetate | | RS-94991-298, Synarel (LHRH agonist) | |
| Nafoxidine HCL | NAFO, NFX | U-11100A (anti-estrogen) | 070735 |
| Nandrolone Decanoa | <u>te</u> * | Deca-Durabolin, Nandrobolic L.A.# (androgen) | |

| Generic Name | Short Names | Synonyms <u>and</u> Brand or Trade Names | + <u>NSC No</u> . |
|---------------------------------------|-------------------|--|-------------------|
| Nandrolone Phenpropionate* | NPP, NTPP | Durabolin, Fenobolin, Nandrobolic, Nandrolone phenylpropionate, Nerobil, Nortestosterone phenylpropionate, Phenobolin, Superanabolon (androgen) | 023162 |
| Nilutamide | | Anti-androgen | |
| 4-Nitroestrone | | | 321803 |
| Nitromifene Citrate | | Anti-estrogen | 110407 |
| Nivazol | | Win-27,914 (glucocorticoid) | |
| Norethandrolone | | Androgen | 009893 070581 |
| Nylestriol | | 49825 (estrogen) | 101105 |
| Oxandrolone* | | Anavar, Protivar, SC 11585, Vasorome, 8075 C. B. (androgen) | 067068 |
| Oxymetholone* | нмо | Adroyd#, Anadrol, CI-406, Drostanolone, Medrotestron, Metholone, Ora-Testryl (anabolic androgen) | 026198 |
| Oxytocin | | Pitocin, Syntocinon (Tumor marker) | |
| Paramethasone* Aceta | <u>ite</u> | Haldrone, Stemex (glucocorticoid) | |
| Polyestradiol Phosph | ate* | Estradurin, Leo 114 (estrogen) | |
| <u>Prednisolone</u> *∇ | PRDL | component of Ataraxoid, Delta-Cortef, Hydeltra, Component of K Predne-Dome#, Meti-Derm, Paracortol#, Predne-Dome#, Sterane (glucocorticoid) | 009120 |
| Prednisolone Acetate | <u>e</u> *∇ | DepPredalone, Meticortelone acetate, Predate, Sterane IM and IA | 010966 |
| Prednisolone Sodium Hemisuccinate∇ | | Meticortelone Soluble, Predmisolone sodium succinate, Solu-Decortin (glucocorticoid) | 009151 |
| Prednisolone Sodium | Phosphate*∇ | Hydeltrasol, Inflamase, Metreton, Component of Optimyd, Pediapred, Predate-S, PSP-IV, Solu-Predalone, Component of Vasocidin Solution (glucocorticoid) | |
| <u>Prednisolone</u> <u>Tebuta</u> | <u>te</u> *∇ | Hydeltra-T.B.A., Predalone T.B.A. (glucocorticoid) | |
| <u>Prednisone</u> *∇ | PDN, PRD, PRED | Ancortone, Dacortin, Delta-Dome#, Deltasone, Deltra, Meticorten, Orasone, Paracort#, Ultracorten (glucocorticoid) | 010023 |
| Prednival*# | | W-4869 (glucocorticoid) | |

 $[\]nabla$ If used in the treatment of the malignancies mentioned in the introduction on p. 37 (bold), code as hormonal therapy. For all other cancers, consider ancillary unless specifically stated to be hormonal therapy.

Generic Name

Short Names

Synonyms and Brand or Trade Names

+NSC No.

| Formula 405 (Non-hormonal sterol derivative) | |
|---|---|
| Component of Cyclogesterin#, Gesterol 50, Lipo-Lutin#, Walutron#, Syngesterone, Syngestrets (progestin) | 009704 |
| Estrovis, W-3566 (estrogen) | |
| W-4540 (progestin) | |
| W-3399 (progestin) | |
| SC-16148 (androgen) | 095147 |
| Growth hormone release inhibiting factor | |
| Protropin (rHGH, growth hormone) | |
| | |
| Component of Aldactazide, Aldactone, SC-9420, Spiresis, Spiridon, Spirolactone, Spirone, Uractone, Verospirone (diuretic; aldosterone antagonist) | 150399 |
| Neodrot | 010972 |
| WIN 14833, Winstrol (androgen) | 043193 |
| 1CI-46,474, Nolvadex, Tamofen, Tamoxifen (anti-estrogen) | 180973 |
| Fludestrin, SQ-9538, Teolit, Teslac, Teslak, Testolacton | 023759 |
| Andro 100, Component of Androgyn L.A., Mertestate#, Oreton#, Synandrol F, Testosteroid, (androgen) | 009700 |
| Component of Depo-Testadiol, DepAndro 100, DepAndro 200, Depo-Testosterone, Depovirin# (androgen) | 009157 |
| Andro L.A. 200, Androgen L.A., Component of Androgyn L.A., Component of Deladumone, Delatestryl, Component of Ditate# (androgen) | 017591 |
| Androgen | |
| Androgen | |
| Synandrol, Synerone# (androgen), | 009166 |
| Proloid, Thyractin#, Thyroprotein# (thyroid hormone) | |
| | derivative) Component of Cyclogesterin#, Gesterol 50, Lipo-Lutin#, Nalutron#, Syngesterone, Syngestrets (progestin) Estrovis, W-3566 (estrogen) W-4540 (progestin) W-3399 (progestin) SC-16148 (androgen) Growth hormone release inhibiting factor Protropin (rHGH, growth hormone) Component of Aldactazide, Aldactone, SC-9420, Spiresis, Spiridon, Spirolactone, Spirone, Uractone, Verospirone (diuretic; aldosterone antagonist) Neodrol WIN 14833, Winstrol (androgen) 1C1-46,474, Nolvadex, Tamofen, Tamoxifen (anti-estrogen) Fludestrin, SQ-9538, Teolit, Teslac, Teslak, Testolacton Andro 100, Component of Androgyn L.A., Mertestate#, Oreton#, Synandrol F, Testosteroid, (androgen) Component of Depo-Testadiol, DepAndro 100, DepAndro 200, Depo-Testosterone, Depovirin# (androgen) Andro L.A. 200, Androgen L.A., Component of Androgyn L.A., Component of Deladumone, Delatestryl, Component of Ditate# (androgen) Androgen Synandrol, Synerone# (androgen), Proloid, Thyractin#, Thyroprotein# |

| Generic Name Short Names | Symonyms and Brand or Trade Names | + <u>NSC No</u> . |
|--|--|-------------------|
| | | |
| <u>Thyroid</u> | Armour Thyroid, Thyrar (thyroid hormone) | 026492 |
| Thyrotropin | | |
| <u>Tigestol</u> * | Progestin | |
| Toremifene Citrate | FC 1157a (anti-estrogen) | |
| <u>Tralonide</u> # | Glucocorticoid | |
| Trestolone Acetate# | U-15,614 (antineoplastic; androgen) | 069948 |
| <u>Triamcinolone</u> * | Aristocort, Kenacort (glucocorticoid) | 013397 |
| <u>Triamcinolone</u> <u>Acetonide</u> * | Azmacort, Flutone#, Kenalog, Component of Mycolog II, Component of Myco-Triacet II, Component of Mytrex, TAC-3, TAC-40#, Tramacin#, Triacet, Triamonide 40, Trymex# (glucocorticoid) | 021916 |
| <u> Triamcinolone Acetonide</u> <u>Sodium Phosphate</u> | CL-61965, CL-106359 (glucocorticoid) | |
| <u>Triamcinolone</u> <u>Diacetate</u> * | Aristocort, Aristocort forte parenteral, Kenacort Diacetate syrup, TAC-D#, Triamolone 40 (glucocorticoid) | |
| <u>Triamcinolone</u> TATBA <u>Hexacetonide</u> * | Aristospan, CL-34433 (glucocorticoid) | |
| <u>Irilostane</u> * | Modrenal, Win 24,540 (adrenocortical suppressant) | |
| Trioxifene Mesylate | Compound 133314 (anti-estrogen) | |
| <u>Triptorelin</u> | CL 118,532 | |
| Vitamin D | <u>Cholecalciferol</u> (functions as a hormone) | |
| Zoladex (see Gosrelin) | | 606864 |

NOTES

Section 6

DRUGS USED IN THE TREATMENT OF AIDS AND ITS COMPLICATIONS

It has long been recognized that immunocompromised patients are at increased risk for certain cancers. Patients treated with immunosuppressant drugs for organ transplantation have an increased incidence of squamous cell carcinomas of the skin and lip, malignant melanoma, non-Hodgkin's lymphoma (including primary lymphoma of the brain), hepatobiliary cancers, cervical carcinoma and carcinomas of the vulva and perineum, and Kaposi's sarcoma.

AIDS was first described as a new and distinct clinical entity in 1981 because of the unusual occurrence of Kaposi's sarcoma and pneumocystis pneumonia in young homosexual men. Thus, the first recognition of cancers associated with human immunodeficiency virus (HIV) infection was closely tied to the initial recognition of AIDS as a disease. Today the relationship between immunocompromise from HIV infection and the development of malignancy is widely accepted. In fact, the occurrence of Kaposi's sarcoma or non-Hodgkin's lymphoma in persons seropositive for antibodies to HIV is considered to be diagnostic of AIDS.

Studies linking population-based tumor registries with population-based AIDS registry files have provided greater understanding of the problem of cancer in patients with antibodies to HIV. In addition to Kaposi's sarcoma and non-Hodgkin's lymphoma, increased incidence of Hodgkin's disease, cancer of the anus and rectum, skin cancers other than melanoma, and cancers of the nose, nasal cavity and middle ear have been demonstrated among AIDS registry cohorts compared to the general population. Longer follow-up of these cohorts may lead to identification of other tumors as late complications of HIV infection.

Treatment of the underlying virus infection is an important aspect of cancer care for patients with AIDS-related malignancies. Most patients with an AIDS-related cancer receive therapy directed at modulating the HIV infection and/or improving immune function in addition to tumor-directed therapy. They may also receive treatment for opportunistic infections or other complications of AIDS.

Drugs for the Treatment of AIDS

Identification of the HIV retrovirus as the causative agent of AIDS and scientific studies of the virus life cycle have opened the door to pharmacologic research in antiviral drugs with potential efficacy against this agent. One step in the reproductive cycle of HIV involves the transfer of genetic information from RNA to DNA, a reverse or "retro" direction from the replication process in human cells. This transfer of genetic code is catalyzed by the enzyme reverse transcriptase. The first drugs tested for activity against HIV were developed to interfere with this part of the viral replication process, either by providing imperfect substrates for the reverse transcriptase enzyme or by inhibiting the formation or activity of reverse transcriptase.

Nucleoside analogs are drugs whose chemical structure is similar to that of the nucleosides (building blocks for DNA and RNA) which are incorporated into viral DNA by reverse transcriptase. In general, the slight differences in structure from normal nucleosides interfere with formation of the

DNA chain, terminating DNA synthesis and preventing viral replication. Well-known nucleoside analogues include azidothymidine (AZT), dideoxycytidine (ddC), and dideoxyinosine (ddI). Other nucleoside analogues that were in clinical trials by early 1992 include D4T, 3'-fluoro-ddT (a dideoxythymidine [ddT] derivative), and 3TC. Still others, such as AzdU, FddC, FlddA, and FLT are in earlier stages of the drug development process.

Some non-nucleoside agents are thought to interfere with the formation of reverse transcriptase from large proenzyme proteins or with the activity of reverse transcriptase itself. For example, both TIBO and BI-RG-587 bind to HIV reverse transcriptase, while hypericin and its related compounds may form free radicals that interact with viral DNA, in addition to their effects on reverse transcriptase.

A number of compounds under investigation as potential antiretroviral drugs appear to work at other stages in viral replication. AL 721, a dietary agent composed of various phospholipids, may inhibit the binding of HIV to target cells by distorting the virus capsule or the cell surface attachment site. Soluble CD4 and specific monoclonal antibodies act at the site of virus attachment to susceptible cells, blocking the receptor for the virus and preventing infection; dextran sulfate may also act in this manner. Compounds such as GLQ 223, castanospermine, MDL 28574, and Deoxynojirimycin (N-butyl DNJ) interfere with the synthesis and glycosylation of viral proteins. Myristic acid analogs prevent the assembly of new virus from polyproteins and viral RNA, while alpha interferon may interfere with the release of virus from infected cells. Agents such as cytotoxic lymphocytes (CTLs), immunomodulators, CD4-toxin, and other antibody-toxin complexes impact the HIV disease process by other mechanisms.

As with chemotherapy, many of the antiviral drugs are under study as combination therapy. Hopefully, synergistic or additive therapeutic effects of antiviral agents will give improved results in HIV treatment, and the use of agents with different toxicities will reduce the morbidity associated with treatment. Agents that have been tested in combination with AZT include acyclovir, alpha interferon, ampligen, ddC, dextran sulfate, and growth factors. Drug resistance is also a problem in managing patients with HIV infection, and major research efforts are directed at developing new agents for use as alternative therapies in patients with HIV infections that have become resistant to standard antiretroviral drugs.

Opportunistic Infections

Opportunistic infections are a major cause of morbidity among HIV-infected persons and represent the cause of death in approximately 90 percent of AIDS patients. Patients with AIDS are at risk for infections that are common in the general population, as well as for opportunistic infections that take advantage of the immunosuppressive effects of HIV.

Primary Pulmonary Infections

Pneumocystis carinii pneumonia (PCP) occurs in approximately 80 percent of patients with AIDS who are not receiving prophylaxis at some time during the course of their illness. Clinical presentation includes fever, respiratory symptoms, and exercise intolerance. Diagnosis depends on demonstrating the organism in sputum or bronchial brushings. Two antibiotic regimens with comparable efficacy in treatment of PCP are trimethoprim/sulfamethoxazole (oral or intravenous

administration) and pentamidine (intravenous administration). For patients who cannot tolerate either of these regimens, investigational approaches of dapsone and trimethoprim or of primaquine plus clindamycin are often used. The addition of corticosteroids (e.g., prednisone) to specific antimicrobial therapy has resulted in marked decrease in respiratory failure and death in patients with moderate or severe PCP. Effective strategies to prevent PCP include prophylaxis with oral trimethoprim/sulfamethoxazole or inhaled (aerosolized) pentamidine.

Patients with AIDS may also develop respiratory symptoms from infections caused by the fungus Coccidioides immitis. Pulmonary findings range from focal lung disease to diffuse pulmonary infiltrates and may be associated with skin lesions and lymph node or liver involvement. Amphotericin B is standard treatment for coccidioidosis in AIDS patients.

Histoplasmosis is endemic in the central United States. In immunocompetent persons, primary infection by *Histoplasma capsulatum* seldom advances beyond the lungs or hilar lymph nodes. However, patients with HIV infection frequently present with disseminated disease manifest by fever and weight loss. Amphotericin B is standard, effective treatment for disseminated disease; maintenance therapy may be required to prevent relapse. Disseminated histoplasmosis in a patient seropositive for HIV is considered diagnostic of AIDS.

Patients with AIDS are at high risk for reactivation of latent tuberculosis (*Mycobacterium tuberculosis*). Early in the course of HIV disease, pulmonary tuberculosis predominates; as HIV disease progresses, atypical presentations and disseminated tuberculosis become more common. Drug regimens used for tuberculosis in AIDS patients include isoniazid, rifampin, pyrazinamide, and ethambutol.

Disseminated infections with nontuberculous mycobacteria, particularly Mycobacterium avium complex, are among the most commonly reported bacterial infections in AIDS patients with severe immunocompromise. In contrast to tuberculosis, pulmonary symptoms are infrequent; fever, weight loss, anemia, and diarrhea are usual manifestations of infection. Antituberculous drug regimens have been disappointing in the treatment of M. avium complex. Prolonged therapy with combination drug regimens is used to reduce symptoms. Agents include ethambutol, clofazamine, ciprofloxacin, rifampin or rifabutin, and amikacin.

Bacterial pneumonia is also common in AIDS patients. Infections acquired in the community are often due to *Streptococcus pneumonia* or *Hemophilus influenza*. Infections acquired during hospitalization (nosocomial infections) are frequently due to Gram negative bacteria or to *Staphylococcus aureus*. Management includes antibiotic regimens appropriate for the sensitivities of the infecting organism.

Central Nervous System Infections

Clinical infections with the protozoa *Toxoplasma gondii* in patients with AIDS usually involve the central nervous system, although pneumonia and disseminated disease have been reported. The lesions demonstrated by computed tomography (CT) or magnetic resonance imaging (MRI) may resemble lymphoma or infection by other opportunistic organisms; in some cases, brain biopsy may be required to confirm the clinical diagnosis. Toxoplasmosis responds well to pyrimethamine in

combination with sulfadiazine; corticosteroids may be used in patients with cerebral edema. Alternative therapeutic approaches are under investigation for patients who are intolerant of pyrimethamine or sulfadiazine. These include combinations of pyrimethamine with clindamycin, high-dose pyrimethamine, azithromycin, clarithromycin, dapsone, and 566C80.

Cryptococcus neoformans, a yeast-like fungus, is the major cause of meningitis in AIDS patients. The organism usually infects the brain as well as the meninges, producing a diffuse encephalitis in many patients. The organism or its antigen is usually detected in spinal fluid obtained by lumbar puncture. Initial therapy for cryptococcal meningitis in AIDS patients comprises amphotericin B alone or in combination with flucytosine; following initial treatment, suppressive therapy with fluconazole is given for the life of the patient. Fungal meningitis in AIDS patients may also be caused by Coccidioides immitis, also treated with amphotericin B.

Central nervous system infections in patients with AIDS may also be due to *Mycobacterium tuberculosis*, presenting with clinical features of meningitis or localized lesions (tuberculomas).

Infections of the Gastrointestinal Tract

Gastrointestinal infections in patients with AIDS are a major cause of morbidity; malabsorption, diarrhea, and wasting can be linked to infection in most patients with advanced disease.

Oropharyngeal candidiasis (thrush) and vaginal infections with Candida species are common in immunocompromised patients. Candidal infections of the esophagus or the respiratory tract are recognized as indicator infections for AIDS diagnosis. These infections are seldom invasive or disseminated unless patients are neutropenic or have other complicating factors. Oropharyngeal candidiasis is treated by topical administration of nystatin, clotrimazole, ketoconazole, or fluconazole. Esophageal candidiasis usually requires ketoconazole; amphotericin B may be administered intravenously to patients with candidal esophagitis that is refractory to therapy with the azole drugs.

Enteric infections with the protozoa Cryptosporidia, Microsporidia, and Isospora may cause diarrhea, weight loss, nausea or vomiting, and abdominal pain. Therapy is predominantly supportive, emphasizing fluid balance, electrolytes, and nutrition. No specific therapies have been shown to be effective for cryptosporidiosis or microsporidiosis, although metronidazole may have some efficacy in microsporidiosis. Isosporiasis is common in Haitian or African patients with AIDS: it can be treated effectively with trimethoprim/sulfamethoxazole, pyrimethamine, metronidazole, or quinacrine.

Bacterial infections of the gastrointestinal tract are also common in AIDS patients. Salmonella, Campylobacter, and Shigella infections, usually self-limited in immunocompetent individuals, often require antibiotic therapy in patients with AIDS.

Viral Infections

Cytomegalovirus (CMV) infection is common among adult patients with AIDS. Retinitis due to CMV is the most frequent presentation, manifest as painless progressive visual impairment in patients with advanced disease. CMV infection at other sites may cause stomatitis, esophagitis, gastritis, and colitis in the gastrointestinal tract; hepatitis and cholecystitis; and adrenalitis with adrenal insufficiency. Antiviral therapy with gancyclovir may be limited by bone marrow toxicity in AIDS patients; foscarnet is an alternative therapy for gancyclovir-resistant infections or in patients who

cannot tolerate gancyclovir.

Progressive infections at oral and genital sites in AIDS patients are usually due to herpes simplex virus (HSV). HSV is also a common cause of esophagitis in patients with advanced disease. Acyclovir is the drug of choice in initial treatment of HSV infections as topical, oral, or intravenous therapy. Foscarnet or vidarabine may be used in acyclovir-resistant HSV.

Common viral infections due to varicella-zoster virus (VZV), such as chicken pox and shingles, may be severe and progressive in immunocompromised patients and may require treatment with acyclovir.

Other Complications of AIDS

Neurologic complications of AIDS are diverse, affecting both the central and peripheral nervous system, and may occur at all stages of disease. The clinical syndrome of AIDS dementia complex is among the most common central nervous system complications of untreated AIDS and is one of the diagnostic criteria for AIDS. Cognitive and behavioral findings (e.g., slowing of thought processes, impaired concentration, apathy, personality changes) are accompanied by impaired motor function. The clinical course of the complex is variable in progression. Accumulating evidence suggests that the clinical syndrome is due to effects of HIV on nervous system tissue, and therapy with antiretroviral agents is directed at the underlying HIV infection. Symptomatic management may include the use of neuroleptic drugs.

Progressive multifocal leukoencephalitis (PML) represents an opportunistic infection by the Jakob-Creutzfeldt (JC) virus. Focal neurologic deficits are caused by areas of demyelination, i.e., loss of the myelin sheath covering nerve fibers. While deficits are usually progressive, the clinical course is more protracted than that of other central nervous system opportunistic infections. There is no proven therapy for PML, although approaches including interferons and antiviral agents are under investigation.

Many agents used to treat AIDS and its complications have broad applications in cancer therapy. For example, antibiotics used to treat infections in AIDS patients are frequently used to treat similar infections in patients who are immunosuppressed by cancer or its treatment. Also, many chemotherapeutic agents and biological response modifiers used in cancer therapy have applications in the management of HIV infection. The inclusion of a drug in this section of Book 8 does <u>not</u> imply that the medication is used <u>only</u> for treatment of AIDS or its complications. Check sections 1-5 first for cancer-directed treatment. If not found in sections 1-5, do not code for cancer patients.

References

- 1. American Foundation for AIDS Research (AmFAR). AIDS/HIV Treatment Directory. Compiled and published on a regular basis by AmFAR, 1993 (Updated quarterly). Available through CDC National AIDS Clearing House (800) 458-5231.
- 2. Johnston MI, Hoth DF. Present status and future prospects for HIV therapies. *Science* **260**:1286-1293, 1993.
- 3. Sanford JP, Sande MA, Gilbert DN, Gerberding JL. *Guide to HIV/AIDS Therapy*. Dallas, Antimicrobial Therapy, Inc., 1993.
- 4. DeVita VT Jr, Hellman S, Rosenberg SA (eds). AIDS: Etiology, Diagnosis, Treatment, and Prevention, Third Edition. Philadelphia, J.B. Lippincott Company, 1992.
- 5. Matthews SJ, Cersosimo RJ, Spivak ML. Zidovudine and other reverse transcriptase inhibitors in the management of human immunodeficiency virus-related disease. *Pharmacotherapy* 11(6):419-449, 1991.

Generic Name

Short Names

Synonyms and Brand or Trade Names

+MSC No.

602670

Drugs Used To Treat HIV Infection

A-77003

A 80967

AL 721 Inhibits binding of HIV to target cells

All-<u>trans</u>-retinoic acid See Tretinoin (under investigation for Rx of

lesions in Kaposi's sarcoma

Alpha-APA derivatives R 18893, R 89439 (alpha-anilino-phenylacetamide)

(Non-nucleoside reverse transcriptase inhibitor)

Alpha Interferon INF May interfere with release of virus from infected

cells

Ampligen Poly 1: Poly C12U (see Biological Response

Modifiers)

Antabuse Disulfiram

AS-101 Arsanilic acid

Autologous CD8 Infusion CD8+ T-lymphocytes from an HIV-infected

person are stimulated outside the body to replicate and then are reinfused

into the patient of origin.

Azidothymidine AZT 3'-Azido-3' Deoxythymidine, BW A 509U,

Retrovir, <u>Zidovudine</u> (inhibits reverse transcriptase) (used in the treatment

of Kaposi's sarcoma)

Azidouridine

BI-RG-587 Nevirapine (non-nucleoside reverse

transcriptase inhibitor)

BRL 61063 Exerts TNF-alpha (cachectin) inhibition.

Castanospermine

CD4-IgG

CD4-PE40 Recomb. protein; consists of HIV-binding

region of CD4 linked to Pseudomonas

Aeruginosa exotoxin A

Generic Name Short Names

Synonyms and Brand or Trade Names

+NSC No.

Drugs Used To Treat HIV Infection (Cont'd)

<u>Cimetidine</u> Tagamet (immunomodulatory effect,

interacts with H₂ receptors on cells

Compound Q (See GLQ223)

Curdian Sulfate

Didehydrodideoxythymidine, Dideoxynucleoside

(thymidine) analogue, Stavudine

(Non-nucleoside reverse transcriptase

inhibitor)

Deoxynojirimycin Butyl-DNJ, N-butyl-DNJ (Protein N-

glycosylation inhibitor)

Dideoxyadenosine ddA 098700

Dideoxycytidine ddC 2',3'-dideoxycytidine, HIVID, 606170

Zalcitabine, Ro 24-2027

Dideoxyinosine ddI BMY-40900, <u>Didanosine</u>, 612049

2',3'-dideoxyinosine, Videx

Fluorothymidine FLT Synthesized fluor-substituted derivative

of dideoxythymidine

GEM 91 May inhibit HIV replication

GLQ223 Compound Q, Trichosanthin

GP120 vaccine Recombinant, an envelope protein from HIV

ENV 2,3 (Biocine)

GP160 vaccine Recombinant, an envelope protein from HIV

HIV immunogen Inactivated virus incorporated in modified

Freund's adjuvant

HPA-23

Human Growth Hormone Protropin (recombinant human growth hormone)

Therapy for HIV-related wasting disease

<u>Hydroxychloroguine</u> <u>Sulfate</u> Plaquenil Sulfate

Hypericin St Johns Wort

Generic Name

Short Names

Synonyms and Brand or Trade Names

+NSC No.

Drugs Used To Treat HIV Infection (Cont'd)

Insulin-like Growth Factor

IGF-1

Imreg-1

Imuthiol

DTC

ACTG 166

Interferon Beta Ser 17

<u>Interferon</u> <u>Gamma</u>

Interleukin-2,

IL-2,

Lymphokine; results in proliferation and

expansion of activated T-lymphocytes

Interleukin-2 Fusion Toxin

Interleukin-3

IL-3

Regulates proliferation and differentiation

of hematopoietic and lymphoid cells

L-697639

L-697661

Pyridinone (Inhibits HIV-1 reverse transcriptase)

Lent inan

MDL 28574

MN rgp120/HIV-1

Myristic acid analogs

Prevents assembly of new virus from

polyproteins and viral RNA

NAC

Mucomyst, N-acetyl cysteine (Cysteine precursor;

may indirectly inhibit HIV replication)

OTC

Procysteine, cysteine precursor

PEG-IL-2

Polyethylene-glycol derivatized IL-2

Pentamidine Isethionate

Pentam 300, RP 2512

620107

<u>Pentoxifylline</u>

Trental (see also Ancillary Drugs)

Peptide T

PMEA

Inhibits HIV replication

Pneumovax

Synonyms and

Generic Name Short Names Brand or Trade Names +NSC No.

Drugs Used To Treat HIV Infection (Cont'd)

Protease inhibitor Ro 31-8959 (Inhibits the protease

enzyme necessary for formation of functional, infectious virions

rCD4 Receptin (Recombinant Soluble CD4)

rgp120 CHO Biocine

rgp120/HIV-1 (IIIB) Genetically engineered form of envelope

glycoprotein gp120 derived from HIV-1

strain IIIB

<u>Ribavirin</u> Virazole

Ro 24-7429 Code name for a benzodiazepine derivative with

anti-HIV activity (TAT gene inhibitor)

sT4 Recombinant Soluble T4

3TC Lamivudine, Analogue of cytidine; inhibits

HIV replication through viral DNA chain

 ${\tt termination}$

Thymic Humoral THF Increases number of T-lymphocytes and augments

Factor cell-mediated immunity

<u>Thymopentin</u> TP-5 Regulates and enhances function of peripheral

T cells

TIBO derivatives R 62913 (Non-nucleoside reverse transcriptase

inhibitor)

U-85961 (bis(heteroaryl)piperazine compound (BHAP)

U-87201E Non-nucleoside reverse transcriptase inhibitor

(bis(heteroaryl)piperazine compound (BHAP)

U-90152 BHAP compound

VaxSyn Recombinant gp160 (See GP160 vaccine)

Wobenzym Suppresses inflammatory response in AIDS

patients

Zidovudine* AZT (see Azidothymidine, Section 1.) 602670

935U83 Inhibits the HIV reverse transcriptase

Syrnonyms and

Generic Name

Short Names

Brand or Trade Names

+NSC No.

Drugs Used To Treat Opportunistic Infections And Other Complications Associated With AIDS

Acyclovir Sodium Zovirax (Rx of HSV and VZV infections: antiviral)

Albendazole Antifungal agent; Rx of microsporidiosis

All-trans retinoic acid Under investigation for Rx of Kaposi's sarcoma

Alpha Interferon Rx of Kaposi's sarcoma

Amikacin Sulfate Amikin (antimycobacterial drug)

Amphotericin B **AMB** Fungizone, Component of Mysteclin-F#

(antifungal)

Amphotericin B Lipid Complex (ABLC)

Anti-B4-blocked Ricin

Ara-C Cytarabine (under investigation for Rx of

Progressive Multifocal Leukoencephalopathy (PML))

Atovaguone Mepron, 566 (therapy for Pneumocystis carinii

pneumonia; under investigation for Rx of

cryptosporidiosis)

Azithromycin Zithromax (Rx of Chlamydia; under investigation

for Rx of Mycobacterium Avium Complex (MAC),

toxoplasmosis, and cryptosporidiosis

BACI Bovine anti-Cryptosporidium Immunoglobulin

(Under investigation for Rx of cryptosporidiosis)

Bactrim/Septra TMP/SMX Cotrim, Trimethoprim (TMP)-sulfamethoxazole (SMX)

Butoconazole Femstat (topical antifungal agent)

BV ara-U

BW 256U87

BW 348U87

BW 882C87

Capreomycin Sulfate Capastal sulfate (antimycobacterial drug)

<u>Ceftriaxone</u> Sodium Rocephin

Ciprofloxacin HCL Cipro (antimycobacterial drug)

Generic Name

Short Names

Symonyms and Brand or Trade Names

+NSC No.

Drugs Used To Treat Opportunistic Infections And Other Complications Associated With AIDS

Clarithromycin Biaxin (antimycobacterial drug)

<u>Clindamycin</u> Cleocin (antiparasitic; Rx of PCP)

<u>Clofazimine</u> Lamprene (antimycobacterial drug) 141046

<u>Clotrimazole</u> Canestin, Component of Lotrimax, Lotrimin,

Component of Lotrisone, Mycelex, Mycelex G (Rx of candidiasis) (topical antifungal

antibiotic)

CMVIg Cytomegalovirus Immune Globulin

Corticosteroids, adjunctive Rx of Pneumocystis pneumonia carinii

Cycloserine Seromycin (antimycobacterial drug)

<u>Dapsone</u> Antiparasitic 006091

Daunorubicin, liposome-encapsulated Daunoxone

Dextran Sulfate Uendex

Dianoxide Furoate Antibiotic; Rx of amebiasis

Doxorubicin, liposome-encapsulated Under investigation for Rx of Kaposi's

Sarcoma

<u>Dronabinol</u> See Tetrahydrocannabinol (appetite stimulant) 134454

Econazole Spectazole (antifungal)

Eflornithine HCl DFMO Ornidyl (Rx of Pneumocystis pneumonia carinii)

Erythropoietin EPO Epogen, Epoietin Alfa, Eprox, rHu-EPO, 628281

Procrit (stimulates RBC production)

Ethambutol HCl Myambutol (Under investigation for Rx of

Mycobacterium Avium Complex (MAC; Rx of

tuberculosis)

Ethionamide Trecator-SC (antimycobacterial drug)

Famciclovir BRL 39123A

Fansidar Prophylaxis for PCP and toxoplasmosis

FIAU (Rx for cytomegalovirus)

Synonyms and

Generic Name Short Names Brand or Trade Names +NSC No.

Drugs Used To Treat Opportunistic Infections And Other Complications Associated With AIDS

<u>Fluconazole</u> Diflucan (antifungal; Rx of Candidiasis)

Flucytosine 5-FC Ancobon (antifungal; adjunct to Amphotericin B)

Foscarnet Sodium PFA EHB 776, Foscavir, Trisodium phosphonoformate

(Rx of cytomegalovirus retinitis and HSV

infections)

<u>Furazolidone</u> Furoxone (Antimicrobial used in the oral RX

of giardiasis

G-CSF Filgrastim, Neupogen (Granulocyte Colony-

stimulating factor; Adjunct of Ganciclovir

therapy; treatment of neutropenia

Gamma Interferon Actimmune

Ganciclovir DHPG, 2 NDG BIOLF-62, BW 759U, Cytovene, RS-21592

(Rx of AIDS-related retinopathies)

GM-CSF Leukine, Sargramostim (Granulocyte macrophage

colony-stimulating factor

Guanfacine Rx of AIDS Dementia Complex

HPMPC Under investigation of asymptomatic CMV

infections

<u>Imipenem-Cilastatin</u> Primaxin, component of (antibiotic;

antimycobacterial)

Iodoquinol Yodoxin (Antibiotic; Rx of amebiasis)

<u>Isoniazid*</u> INH Cotinazin, Dinacrin#, Ditubin#, Isolyn#,

Isonicotinic acid hydrazide, Laniazid, Niconyl#, Nidaton#, Nydrazid, Rimifon#,

Teebaconin, Tisin#, Tyvid#

Isoprinosine

Itraconazole ITZ R-52,211, Sporonax (antifungal) (Rx of Candidiasis)

IVIG Intravenous immunoglobulin; provides

passive immunity (viral infections)

Ketoconazole (antifungal)

L-693,989 Activity against PCP and Candida

Generic Name

Short Names

Synonyms and Brand or Trade Names

+NSC No.

Drugs Used To Treat Opportunistic Infections And Other Complications Associated With AIDS

Letrazuril

Rx of Cryptosporidiosis

Lomotil

Anti-diarrhea

Malathion

Ovid (Rx for ectoparasites)

Megestrol Acetate

Megace (appetite stimulant) (see Hormones)

<u>Metronidazole</u>

Flagyl, Component of Flagyl I.V. RTU,

050364

Metro I.V., Protostat, Satric

(Rx of microsporidiosis)

Mexiletine HCl

Mexitil (Therapy for HIV-related neuropathy)

Miconazole

Micatin, Monistat (topical antifungal agent)

MSL-109

Antiviral monoclonal antibody with activity

against CMV strains

Mucopirocin

Bactroban

Nimodipine

Nimotop (Oral calcium-channel blocking agent under investigation for the Rx of

AIDS dementia complex

Nystatin

Mycostatin, Nilstat, Nystex, component of Mycolog II, component of Myco-Triacet II, component of Mytrex, component of Terrastatin, component of Tetrastatin (topical antifungal antibiotic)

(Rx of Candidiasis)

Ofloxacin

Floxin (antimycobacterial drug)

Para-aminosalicylic acid

PAS, Teebacin (antimycobacterial drug)

Paromomycin Sulfate

Humatin (antibiotic) (Under investigation

for Rx of cryptosporidiosis

<u>Pefloxacin</u>

Under investigation for Rx of PCP

Generic Name

Short Names

Synonyms and Brand or Trade Names

+NSC No.

Drugs Used To Treat Opportunistic Infections And Other Complications Associated With AIDS

Pentamidine Lomidine, NebuPent, Pentam 300 (Rx or

prophylaxis of PCP) (antiparasitic)

Peptide T Under investigation for Rx of HIV-associated

neuropsychiatric impairment and neuopathy

Peridex Chlorhexidine gluconate (Under investigation

for the prevention of oral candidiasis

Permethrin (topical agent used for ectoparasites)

<u>Piritrexim Isethionate</u> Rx of PCP

Platelet Factor 4 Under investigation for Rx of Kaposis's sarcoma

Podofilox Condylox (Rx of human papilloma virus (HPV)

<u>Primaguine Phosphate</u> Used in combination with Clindamycin for Rx of

PCP (antiparasitic)

Pyrantel Pamoate (antihelminthic)

Pyrazinamide PZA Rx of tuberculosis

<u>Pyrimethamine</u> Daraprim, Component of Fansidar, Malocide

(Rx of toxoplasmosis) (antiparasitic)

Quinacrine Atabrine (antiparasitic)

Ranitidine HCl Zantac

Retinoic Acid (Rx of molluscum contagiosum virus)

Rifabutin Ansamycin, Mycobutin 113926

(Rx of Mycobacterium Avium complex (MAC))

<u>Rifampin</u> Rifadin, Rimactane

(Under investigation for Rx of MAC; Rx of

tuberculosis)

SP-PG Inhibits angiogenesis; under investigation

for Rx of Kaposis's sarcoma

Sparfloxacin (antimycobacterial)

Spiramycin Rx of cryptosporidiosis 055926

Generic Name

Short Names

Synonyms and Brand or Trade Names

+NSC No.

Drugs Used To Treat Opportunistic Infections And Other Complications Associated With AIDS

Spiramycin HCL 064393

Streptomycin

Sulfadiazine (antiparasitic)

Sulfamethoxazole Rx of PCP

Terconazole Terazol (topical antifungal agent)

Thiabendazole Mintezol (Rx of strongyloidiasis)

Tioconazole Vagistat (antifungal)

TLC G-65 Rx of Mycobacterium Avium Complex

TNP-470 AGM-1470# (inhibits angiogenesis)

<u>Irifluridine</u> IFI Viroptic, See F3TDR 075520

<u>Trimethoprim</u> TMP Trimpex, Proloprim (Rx or prophylaxis of PCP) 106568

(antiparasitic)

<u>Trimetrexate</u> Salvage therapy for PCP

<u>Trospectomycin Sulfate</u> Antibiotic for Rx of oral chlamydia and

gonorrhea; under investigation for Rx of

Mycobacterium Avium Complex

Vidarabine

WR 6026 Therapy for Pneumocystis carinii pneumonia

566C80

NOTES

PROTOCOL INVESTIGATORS

AFIP Armed Forces Institute of Pathology

AJC American Joint Commission, College of Surgeons (ACOS)

AMLI Acute Myelocytic Leukemia Intergroup

BTSG Brain Tumor Study Group
BTS British Testicular Study

CALGA Cancer and Leukemia Group A
CALGB Cancer and Acute Leukemia Group B
CCSG Children's Cancer Study Group

CCDEP Central Clinical Drug Evaluation Program

COG Central Oncology Group

CRC Cancer Research Campaign United Kingdom

DFCC Dana-Farber Cancer Center

ECOG Eastern Cooperative Oncology Group

EORTC European Organization for Research and Treatment of

Cancer

ESSG Ewing's Sarcoma Study Group FDA Federal Drug Administration

FIGO International Federation of Gynecology and Obstetrics

GITSG Gastrointestinal Tumor Study Group
GOG Gynecologic Oncology Group
HNCP Head and Neck Contracts Program
HTSG Hepatic Tumor Study Group

IARC International Agency for Research on Cancer IFAC International Federation Against Cancer

INTERG Intergroup (Other)

IRS Intergroup Rhabdomyosarcoma Study

LCSG Lung Cancer Study Group
MAOP Mid-Atlantic Oncology Program

MARCOG Mid-Atlantic Regional Co-Op Oncology Group
NABMTG North American Bone Marrow Treatment Group

NCCTG North Central Cancer Treatment Group

NCI National Cancer Institute

NCOG Northern California Oncology Group

NETDC New England Trophoblastic Disease Center, Harvard

Medical School

NORCA Nutrition Oncology Research Cooperative Association

NPCTG National Prostatic Cancer Treatment Group

NSABP National Surgical Adjuvant Project for Breast and Bowel

Cancers

NWTS National Wilms' Tumor Study
UORG Uro-Oncology Research Group
POA Piedmont Oncology Association
POG Pediatric Oncology Group

PVACCG Pacific VA Cancer Chemotherapy Group

PVSG Polycythemia Vera Study Group

PROTOCOL INVESTIGATORS (cont'd)

RMS Intergroup Rhabdomyosarcoma Study, Children's Cancer

Study Group

RTOG Radiation Therapy Oncology Group SECSG Southeastern Cancer Study Group

SIOP International Society of Pediatrics Oncology

SWOG Southwest Oncology Group
TPNG Total Parenteral Nutrition Group
UICC International Union Against Cancer

VACG Veterans Administration Chemotherapy Group

VALG Veterans Administration Lung Group

VASAG Veterans Administration Surgical Adjuvant Group VASOG Veterans Administration Surgical Oncology Group

WCCG Western Cancer Chemotherapy Group

WCG Weski Cancer Group
WHO World Health Organization
WTSG Wilms' Tumor Study Group

SELECTED BIBLIOGRAPHY

SELECTED BIBLIOGRAPHY

- American Index of Drugs. Philadelphia: J.B. Lippincott Co., 1981.
- Cancer Treatment Reports, National Cancer Institute. Washington, D.C.: Superintendent of Documents, U. S. Government Printing Office.
- Chemicals for Which Investigational New Drugs Applications Have Been Filed by NCI. Information Technology Branch, DCT, NCI, 1991
- Chabner, D.E. *The Language of Medicine*, Work Text in Terminology. 2nd ed. Philadelphia: Saunders, 1981.
- Compendia-Based Drug Bulletin. Association of Community Cancer Centers. Rockville, MD, 1992.
- Department of Health and Human Services, National Institutes of Health, Division of Cancer Treatment. *Bulletins*, 1979-1981.
- DeVita, Vincent T. Jr., M.D.; Hellman, Samuel, M.D.; and Rosenberg, Steven A., M.D., Ph.D. Cancer: Principles and Practice of Oncology. Philadelphia: J. P. Lipincott, 1982.
- Diseases of the Lymphatic System, Diagnosis and Therapy. Edited by D.W. Molander, M.D. New York: Springer-Verlag, 1984.
- Dorr, R. T. and Fritz, W.L. Cancer Chemotherapy Handbook. New York: Elsevier, 1980.
- Frei, Emil, III, M.D., and Holland, James F., M.D. Cancer Medicine. Philadelphia: Lea & Febiger, 1973.
- Griffiths, M.C.; Dickerman, M.J.; and Miller, L.C., (editors). USAN and the USP Dictionary of Drug Names. Rockville, MD: U.S. Pharmacopeial Convention, Inc., 1976.
- International Classification of Diseases for Oncology. Geneva: World Health Organization, 1976.
- Knobf, Mary K. et al. Cancer Chemotherapy Treatment and Care. Edited by M.E. Morra. New Haven: Communications Office, Yale Comprehensive Cancer Center, 1979.
- Mondell, Walter, ed. Drugs of Choice. St. Louis: C.V. Mosby Company, 1978-1979.
- Physician's Desk Reference. Montvale, NJ: Medical Economics Data, 1993.
- Rubin, Philip, M.D., ed. Clinical Oncology for Medical Students and Physicians. 6th ed. Rochester: The University of Rochester School of Medicine and Dentistry and the American Cancer Society, Inc., 1983.

SELECTED BIBLIOGRAPHY (cont'd)

- USAN and the USP Dictionary of Drug Names. Edited by Carolyn A. Fleeger. United States Pharmacopeial Convention, Inc. Rockville, Maryland, 1992
- WHO Technical Report Series 605, "Chemotherapy of Solid Tumors; A Report of a WHO Expert Committee." Albany: WHO Publications Centre, USA, 1977.

DRUG INDEX

BOOK 8 INDEX

```
8075 C.B. (see Oxandrolone*), 79
016895 (see Lithium Carbonate), 43
085791 (see Ethacrynic Acid), 41
111180 (see Acetylcysteine), 39
12,540 (see Tubercidine), 27
131 I (see I 131), 63
144 A1 (see Aclarubicin), 15
24281 (antibiotic) (see Mitocarcin), 23
27-400 (see Cyclosporine), 62
29060-LE (see Vinblastine* Sulfate), 27
296961 (see WR 2721), 41, 46
3123L (see Puromycin (HCl)), 25
31595C (see Mitosper), 23
32379 (see Dromostanolone Propionate), 75
32645 (see Vinleurosine Sulfate#), 28
33379 (see Flurandrenolide), 76
3678l (see Vinrosidine Sulfate#), 28
37231 (see Vincristine* Sulfate), 28
38000 (see Clometherone#), 74
38489 (see Nortriptyline HCl), 44
47599 (see Pyrazofurin#), 25
49040 (see Vinglycinate Sulfate), 28
49825 (see Nylestriol), 79
5052 (see Acetyclysteine), 39
5071 (see Megestrol Acetate), 78
515776 (see Dipyridamole), 41
5278 R.P. (see Streptonigrin), 25
5373 (see Melengestrol Acetate), 78
566 (see Atovaquone), 95
566C80, 100
59156 (see Enpromate#), 20
630 (see Gosrelin), 77
68618 (see Mycophenolic acid), 51
79T61 (see 9-Ethyl-6-Mercaptopurine), 20
Α
A-20968 (see Piposulfan), 24
A-2371 (see Plicamycin*), 24
A-4942 (see Ifosfamide*), 21
A-8103 (see Pipobroman*), 24
A-hydroCort (see Hydrocortisone Sodium Succinate). 77
A-methaPred (see Methylprednisolone Sodium Succinate), 78
```

A-Ninopterin (see Aminopterin), 15 A-TD (see Aminothiadiazole), 15 AAFC (see Flurocitabine#), 20 AB-100 (see Uredepa), 27 AB-103 (see Benzodepa), 16 AB-132 (see Meturedepa), 23 Abbott 43818 (see Leuprolide Acetate), 77 Aberel (see Tretinoin), 51 Abirol (see Methandrostenolone), 78 ABLC (see Amphotericin B Lipid Complex), 95 Aburamycin B (see Chromomycin A3#), 17 AC-1075 (see Cytarabine HCl), 18 5-AC (see Azacitidine), 51 Accutane (see Retinoic acid), 51 ACD (see Dactinomycin*), 18 Acethropan (see Adrenocorticotropic hormone), 73 Acetyladriamycin (see Daunorubicin (HCl)*), 18 Acetylcysteine, 39 Acitretin, 51 Acivicin, 15 Aclacinomycin (see Aclarubicin), 15 Aclacinomycin-A# (see Aclarubicin), 15 Aclacinon (see Aclarubicin), 15 Aclarubein (see Aclarubicin), 15 Aclarubicin, 15 ACM (see Aclarubicin), 15 ACM-A (see Aclarubicin), 15 Acodazole (HCl), 15 ACPC (see Cycloleucin), 18 ACR (see Aclarubicin), 15 Acrichin (see Quinacrine (HCl)), 45 Acridinyl anisidide (see Amsacrine), 15 Acrinamine (see Quinacrine (HCl)), 45 Acriquine (see Quinacrine (HCl)), 45 Acronine*, 15 Acronycin# (see Acronine*), 15 Acronycine (see Acronine*), 15 ACT (see Dactinomycin*), 18 ACT-D (see Dactinomycin*), 18 ACTD (see Dactinomycin*), 18 ACTG (see Imuthil), 93 ACTH (see Adrenocorticotropic hormone), 73 ACTH (see Corticotropin), 74 Acthar (see Adrenocorticotropic hormone), 73

Acticort (see Hydrocortisone), 77 Actimmune (Genentech) (see Interferon Gamma-1b), 63 Actimmune (see Gamma Interferon), 97 Actinex (see Masoprocol), 22 Actinochrysin (see Cactinomycin), 16 Actinomycete fermentation product (see Piperazinedione compound), 24 Actinomycin AIV (see Dactinomycin*), 18 Actinomycin C# (see Cactinomycin), 16 Actinomycin C1 (see Dactinomycin*), 18 Actinomycin D (see Dactinomycin*), 18 Actinomycin I1 (see Dactinomycin*), 18 Actinomycin IV (see Dactinomycin*), 18 Actinomycindioic D acid dilactone (see Dactinomycin*), 18 Acto-D (see Dactinomycin*), 18 Acyclovir Sodium, 95 Acyclovir, 39, 94 ACZ (see Acetazolamide), 39 AD (see Dactinomycin*), 18 ADA Transduced T-cell, 61 ADAH (see Bisantrene (HCl)), 16 Adanon HCl# (see Methadone HCl), 43 ADC (see Bisantrene (HCl)), 16 Adenine arabinoside (see Vidarabine*), 27 Adicillin, 39 ADM (see Doxorubicin (HCl)*), 19 Adozelesin, 15 ADR (see Doxorubicin (HCl)*), 19 ADR-529 (see Dexrazoxane), 19 Adrenal Cortex hormone (see Adrenocorticotropic hormone), 73 Adrenocorticotrophin (see Adrenocorticotropic hormone), 73 Adrenocorticotropic hormone, 73 Adrenocorticotropin (see Adrenocorticotropic hormone), 73 ADRI (see Doxorubicin (HCl)*), 19 Adriamycin (see Doxorubicin (HCl)*), 19 Adriamycin-TM (see Doxorubicin (HCl)*), 19 Adriblastina (Italy) (see Doxorubicin (HCl)*), 19 Adroyd# (see Oxymetholone*), 79 Adrucil (see Fluorouracil*), 20 Aeroseb-Dex (see Dexamethasone), 40, 74 Aeroseb-HC (see Hydrocortisone), 77 AG (see Aminoglutethimide), 73 8-AG (see 8-Azaguanine), 15 AGM-1470, 94 AGT (see Aminoglutethimide), 73

AHR-3070-C (see Metoclopramide (HCl)*), 43 Airol (see Tretinoin), 51 Akrichin (see Quinacrine (HCl)), 45 AL 721, 94 Alanine mustard (see Melphalan* (HCl)), 22 Alanosine, 15 Albendazole, 95 Albuterol, 39 Aldactazide (see Spironolactone), 80 Aldactone (see Spironolactone), 80 Aldesleukin (see Interleukin-2), 63 Alendronate Sodium, 39 Alexan (see Cytarabine HCl), 18 Alferon N Injection (see Interferon Alfa-n3), 63 Algestone* Acetophenide, 73 Alizapride, 39 Alkaloid H-3 (see Colcemid), 17 Alkeran (see Melphalan* (HCl)), 22 All-trans retinoic acid (see Tretinoin), 51, 91 Allergan 211 (see Idoxuridine*), 42 ALLO (see Allopurinol*), 39 Allogeneic bone marrow transplantation (see Allogeneic cells), 61 Allogeneic cells, 61 Allopur (see Allopurinol*), 39 Allopurinol*, 39 Alpha 1 (see Thymosin), 66 Alpha Interferon, 91 Alpha TGI (see Teroxirone), 26 Alpha-DFMO HCl (see DFMO), 19 Alpha-Difluoromethylornithine (see DFMO), 19 Alpha-TGdR, 15 Alphadrol# (see Fluprednisolone*), 76 Alphatrex (see Betamethasone Dipropionate), 73 Alprazolam, 39 Altretamine*, 15 Amadinone Acetate, 73 AMB (see Amphotericin B), 95 Ambochlorin (see Chlorambucil*), 17 Ambomycin, 15 Amcinonide, 73 Amedel (see Pipobroman*), 24 Ametantrone Acetate, 15

Amethopterin# (see Methotrexate*), 22

Amidohydrolase (see (Escherichia coli)), 61

Ametycine (see Mitomycin*), 23

Amifostine (see WR 2721), 46

Amifostine, 39

Amikacin Sulfate, 95

Amikin (see Amikacin Sulfate), 95

Aminobenzoate Potassium, 39

Aminoglutethimide, 73

Aminopteridine (see Aminopterin), 15

Aminopterin, 15

Aminopterin sodium (see Aminopterin), 15

Aminosyn, 39

2-Amino-1,3,4-thiadiazole (see Aminothiadiazole), 15

Aminothiadiazole, 15

Amiodarone, 39

Amnestrogen#, (see Estrogens, Esterified), 76

Amonafide (see Nafidimide), 23

Amonafide HCl (see Nafidimide), 23

Amphotericin B, 91

Amphotericin B Lipid Complex, 95

Ampligen (see Poly I: Poly C12U), 65

Ampligen, 91

AMSA (see Amsacrine), 15

Amsacrine, 15

Amsidyl (see Amsacrine), 15

AMX (see Azimexon), 61

Anabolin (see Methandrostenolone), 78

Anadrol (see Oxymetholone*), 79

Anagestone Acetate, 73

Anatropin# (see Anagestone Acetate), 73

Anavar (see Oxandrolone*), 79

Anaxirone (see Triglycidylurazol), 27

Ancitabine hydrochloride (see Cyclocytidine (HCl)), 17

Ancobon (see Flucytosine), 97

Ancortone (see Prednisone), 79

Ancyte# (see Piposulfan), 24

Andro 100 (see Testosterone), 80

Andro L.A. 200 (see Testosterone Enanthate), 80

Androfluorene (see Fluoxymesterone), 76

Androgen L.A. (see Testosterone Enanthate), 80

Androgyn L.A. (see Estradiol), 75

Androgyn L.A. (see Testosterone Enanthate), 80

Androgyn L.A. (see Testosterone), 80

Andrometh (see Methyltestosterone), 78

Androsan (see Methyltestosterone), 78

Androsten (see Methyltestosterone), 78

Anergan 25 (see Promethazine HCl*), 45 Anergan 50 (see Promethazine HCl*), 45 Anertan (see Methyltestosterone), 78 Anestacon (see Lidocaine), 43 ANG (see Anguidine), 15 ANG 66 (see Anguidine), 15 Anguidin (see Anguidine), 15 Anguidine, 15 Aniline mustard, 15 Anisene (see Chlorotrianisene), 74 Anrosterolo (see Fluoxymesterone), 76 Ansaid (see Flurbiprofen), 41 Ansamacrolide compound (see Maytansine), 22 Ansamycin (see Rifabutin), 99 Antabuse, 91 Anthramycin, 15 Anthrapyrazole C 1941, 15 Anthrapyrazole Dihydrochloride (see Piroxantrone HCl), 24 Anti-B4-blocked Ricin, 95 Anti-Thymocyte Globulin, 61 Antibiotic 1037 (see Toyocamycin), 27 Antibiotic 1719 (see Azotomycin), 16 Antibiotic A 23813 (see Pyrazofurin#), 25 Antibiotic AT-125 (see Acivicin), 15 Antibiotic B 599 (see Chromomycin A3#), 17 Antibiotic CI-920 (see Fostriecin Sodium), 20 Antibiotic CL 1565A (see Fostriecin Sodium), 20 Antibiotic E 212 (see Toyocamycin), 27 Antibiotic LA 7017 (see Plicamycin*), 24 Antibiotic MA (see Aclarubicin), 15 Antibiotic U-18,496 (see Azacitidine), 51 Antibiotic XK 101-1 (see Tubercidine), 27 Anticea, I-131, 61 Antiferritin antibodies, I-131, 61 Antiferritin antibody (see Anticea), 63 Antifolan (see Methotrexate*), 22 Antimalarina (see Quinacrine (HCl)), 45 Antiren (see Piperazine), 44 Antramycin (see Anthramycin), 15 Antrypol (see Suramin Sodium), 26 APGA (see Aminopterin), 15 Aphidicolin Glycinate, 15 Aphidicolin-17 glycinate (see Aphidicolin Glycinate), 15 Aplisol (see Tubercidine), 27

Aquadiol# (see Estradiol), 75 Ara-A (see Vidarabine*), 27 Ara-AC (see Fazarabine), 20 Ara-adenosine (see Vidarabine*), 27 Ara-C (see Cytarabine HCl), 18 Ara-C (see Cytarabine), 51 Ara-C, 51, 95 Ara-cytidine (see Cytarabine HCl), 18 Arabinocytidine (see Cytarabine HCl), 18 Arabinosyladenine (see Vidarabine*), 27 Aristocort (see Triamcinolone Diacetate), 81 Aristocort (see Triamcinolone), 81 Aristocort forte parenteral (see Triamcinolone Diacetate), 81 Aristospan (see Triamcinolone Hexacetonide), 81 Armour Thyroid (see Thyroid), 81 Arsanilic acid (see AS-101), 91 AS-101, 91 Asalex (see Asaley), 15 Asaley, 15 Asnase (see Asparaginase), 61 ASP (see Asparaginase), 61 ASPA (see Asparaginase), 61 Asparaginase (Erwinia carotovora), 61 Asparaginase, 61 Asparaginase-PEG (K-H), 61 Asperlin#, 15 Asta B 518 (see Cyclophosphamide*), 18 Asta Z-4942 (see Ifosfamide*), 21 Autologous, 61 Autologous Lymphocytes (see ADA Transduced T-cell), 61 Autologous Tumor Cell Vaccine (see Autologous), 61 Aventyl (see Nortriptyline HCl), 44 Avinar# (see Uredepa), 27 AY-11,440 (progestin) (see Clogestone* Acetate), 74 AY-62022 (progestin) (see Medrogestone), 77 Ayr (see Sodium Chloride), 45 5-AZA (see Azacitidine), 51 5-AZAC (see Azacitidine), 51 Azacitidine, 51 Azacytidine (see Azacitidine), 51 5-Azacytidine (see Azacitidine), 51 5-Aza-2'Deoxycytidine, 15 Azaguanine (see 8-Azaguanine), 15

Azan (see 8-Azaguanine), 15

Azanin (see Azathioprine*), 16 Azapicyl. 15 Azaribine*, 15 AZAS (see Azaserine*#), 16 Azaserin (see Azaserine*#), 16 Azaserine*#, 16 AzAT (see Azathioprine*), 16 Azatepa (see Azetepa#), 16 Azathioprin (see Azathioprine*), 16 Azathioprine*, 16 Azatioprin (see Azathioprine*), 16 6-Azauracil riboside (see 6-Azuridine), 16 Azauridine (see 6-Azuridine), 16 6-Azauridine (see 6-Azuridine), 16 6-Azauridine triacetate (see Azaribine*), 15 AZC (see Azacitidine), 51 5-AZCR (see Azacitidine), 51 Azetepa#, 16 AZG (see 8-Azaguanine), 15 3'-Azido-3' deoxythymidine (see Zidovudine*), 28 Azidothymidine, 16, 91 Azidouridine, 91 Azimexon, 61 2-aziridinecarboxamide compound (see Azimexon), 61 Aziridinyl Benzoquinone (see Diaziquone), 19 Aziridinyl cyclohexane (see Cyclohexanecarboxamide), 18 Azithromycin, 95 Azmacort (see Triamcinolone Acetonide), 81 5-azocytosine arabinoside (see Fazarabine), 20 Azo Gantanol, component of (see Sulfamethoxazole), 46 Azolastone (see Mitozolomide), 23 AZOT (see Azotomycin), 16 Azothioprine (see Azathioprine*), 16 Azotomycin, 16 AZQ (see Diaziquone), 19 AZR (see Azaribine*), 15 AZS (see Azaserine*#), 16 AZT (see Azidothymidine), 16, 91 AZT (see Zidovudine*), 28, 94 AzUR (see 6-Azuridine), 16 6-Azur (see 6-Azuridine), 16 6-Azuridine, 16

B 35251 (antibiotic) (see Mitocromin), 23

B-2' Deoxythioguanosine (see Beta-TGdR), 16

B-28 (see 8-Azaguanine), 15

B-S-P (see Betamethasone Sodium Phosphate), 73

Ba 24648 (see Cyclohexanecarboxamide), 18

Ba 41166/E (see Rifampin), 25

Ba-16038 (see Aminoglutethimide), 73

Babidium chloride (see Ethidium chloride), 20

BACI, 61

Bacillus Calmette-Guerin (see BCG), 61

Bactrim, component of (see Sulfamethoxazole), 46

Bactrim, component of (see Trimethoprim), 46

Bactrim/Septra, 39, 95

Bactroban (see Mucopirocin), 98

BAF (see Baker's antifol), 16

Baker's antifol, 16

Baker's antifolante (see Baker's antifol), 16

Bamide (see Hydroxyurea*), 21

Basoprocin (see Liothyronine Sodium), 77

Batanopride HCl, 39

BAY G6575 (see Nafazatrom), 23

Bayer 205 (see Suramin Sodium), 26

BBM-2478A (see Elsamitrucin), 19

BCG, 61

BCM (see Mannomustine), 43

BCNU (see Carmustine*), 17

BDH-1298 (see Megestrol Acetate), 78

BDH-1921 (see Melengestrol Acetate), 78

BDU (see Bromodeoxyuridine), 39

5-BDU (see Bromodeoxyuridine), 39

Beclometasone (see Beclomethasone Dipropionate), 73

Beclomethasone Dipropionate, 73

Beclovent (see Beclomethasone Dipropionate), 73

Beconase (see Beclomethasone Dipropionate), 73

Beesix (see Pyridoxine HCl), 45

Belustine (nitrosourea) (see Lomustine*), 21

Benadryl (see Diphenhydramine HCI), 41

Benisone# (see Betamethasone Benzoate), 73

Benorterone#, 73

Benylin, component of (see Diphenhydramine HCl), 41

Benzcarbimine (see Benzodepa), 16

Benzestrol, 73

Benzisoquinolenedione (see Nafidimide), 23

Benznidazole, 39

Benzodepa, 16

Benzoyl hydrazone daunorubicin (see Zorubicin (HCl)), 28

Benzquinamide, 39

Bestatin, 61

Beta Carotene, 39

Beta-Arabinosylcytosine (see Cytarabine HCl), 18

Beta-TGdR, 16

Beta-Val (see Betamethasone Valerate), 73

Betamethasone, 73

Betamethasone Acetate, 73

Betamethasone Benzoate, 73

Betamethasone Dipropionate, 73

Betamethasone Sodium Phosphate, 73

Betamethasone Valerate, 73

Betapar# (see Meprednisone*), 78

Betaseron (see Interferon Alfa-2b), 63

Betatrex (see Betamethasone Valerate), 73

BI-RG-587, 91

Biaxin (see Clarithromycin), 96

BIC (see TIC-mustard), 27

BiCNU (see Carmustine*), 17

BIDA (see Nafidimide), 23

Bio-Des (see Diethylstilbestrol), 75

Biocine (see rgp120 CHO), 94

BIOLF-62 (see Ganciclovir), 97

Biorex, England (see Carbenoxolone Sodium), 73

Biostim, 62

Bisantrene (HCl), 16

Bisantrene (see Bisantrene (HCl)), 16

Bisbenzamide (see Pibenzimol HCl), 24

Bisbenzimidazole (see Pibenzimol HCl), 24

BL 191 (see Pentoxifylline), 44

BLE (see Bleomycin Sulfate), 16

Blenoxane (see Bleomycin Sulfate), 16

BLEO (see Bleomycin Sulfate), 16

Bleomycin (see Bleomycin Sulfate), 16

Bleomycin Sulfate, 16

BLM (see Bleomycin Sulfate), 16

Bloxanth (see Allopurinol*), 39

BM 41.440 (see Ilmofosine), 21 BMY-25801-01 (anti-emetic) (see Batanopride HCI), 39 BMY-28090 (see Elsamitrucin), 19 BMY-40900 (see Dideoxyinosine), BMY-41606 (see Vapreotide), 27 Bolasterone#, 73 Borocaptate Sodium B 10, 39 Borolife (see Borocaptate Sodium B 10), 39 Bovisynchron (see Chlormadinone* Acetate), 73 BrdUrd (see Bromodeoxyuridine), 39 Brequinar Sodium, 16 Brevicon, component of (see Ethinyl Estradiol), 76 BRL 39123A (see Famciclovir), 96 BRL 61063 91 BRL-43694 (see Granisetron), 42 Bromacrylide#, 16 Bromocryptine, 62 Bromodeoxyuridine, 39 5-Bromodeoxyuridine (see Bromodeoxyuridine), 39 5-Bromodesoxy-uridine (see Bromodeoxyuridine), 39 Bromoergocryptine (see Bromocryptine), 62 Bromouracil deoxyriboside (see Bromodeoxyuridine), 39 Bropirimine, 16 Broxuridine (radiosensitizer) (see Bromodeoxyuridine), 39 BRU (see Bruceantin), 16 Bruceantin, 16 Bruneomycin (see Streptonigrin), 25 Bryostatin, 16 BSF (see Busulfan*), 16 BSO, 42 BST (see Bestatin), 61 BTGR (see Beta-TGdR), 16 BTIC (see TIC-mustard), 27 BTS 18,322 (see Flurbiprofen), 41 BU (see Busulfan*), 16 BU-2231A (see Talisomycin), 26 BUdR (see Bromodeoxyuridine), 39 5-BUDR (see Bromodeoxyuridine), 39 BUS (see Busulfan*), 16 Busrelin Acetate, 73 BUSU (see Busulfan*), 16 Busulfan*, 16 Busulphan (see Busulfan*), 16 Butoconazole, 95

Butyl-DNJ (see Deoxynojirimycin), 92 BV ara-U, 95 BW 256U87, 95 BW 348U87, 95 BW 50-63 (see Pyrimethamine), 45 BW 57-322 (see Azathioprine*), 16 BW 759U (see Ganciclovir), 97 BW 882C87, 95 BW A 509U (see Azidothymidine), 16, 91 BW A 773U, 16 BW A509U (see Zidovudine*), 28 BW A770U mesylate (see Crisnatol mesylate), 17 BW-56-158 (see Allopurinol*), 39 BW-57-323 (see Thiamiprine), 26 C C.F. (see Leucovorin (calcium)*), 43 C.p. (see C-parvum), 62 C.P. 15,467-61 (see Lithium Carbonate), 43 C-12669 (see Colcemid), 17 C-parvum, 62 CA (see Cytarabine HCl), 18 CACP (see Cisplatin*), 17 Cactinomycin, 16 CAL (see Calusterone), 73 Cal-Sup (see Calcium Carbonate), 39 Caladryl, component of (see Diphenhydramine HCl), 41 Calan (see Verapamil (HCl)*), 46 Calcimar (see Calcitonin*), 39 Calcitonin*, 39 Calcium Carbonate, 39 Calcium folinate (see Leucovorin (calcium)*), 43 Calcium Gluconate, 39 Calcium leucovorin (see Leucovorin (calcium)*), 43 Calusterone, 73 CAMP (see Camptothecin), 16 Camptothecin, 16 Camptothecine sodium (see Camptothecin), 16 Canestin (see Clotrimazole), 96 Cannabinol (see Tetrahydrocannabinol), 46 CAP (see Chlormadinone* Acetate), 73 Capastal sulfate (see Capreomycin Sulfate), 95 Capreomycin Sulfate, 95

Caracemide, 16 Carafate (anti-ulcerative) (see Sucralfate), 46 Carbamohydroxamic acid (see Hydroxyurea*), 21 Carbamohydroximic acid (see Hydroxyurea*), 21 Carbamoyl oxime (see Hydroxyurea*), 21 Carbenoxolone Sodium, 73 Carbetimer, 17 Carbomycin (see Magnamycin), 22 Carbomycin A (see Magnamycin), 22 Carboplatin*, 17 Carbowax Sentry (see Polyethylene Glycol), 44 5-Carboxyanilino-2-thiobarbituric acid (see Merbarone), 22 Cardizem (vasodilator, calcium influx inhibitor) (see Diltiazem HCL), 41 Carminomicin I (see Carubicin (HCl)), 17 Carminomycin (see Carubicin (HCl)), 17 Carminomycin HCl# (see Carubicin (HCl)), 17 Carminomycin I (see Carubicin (HCl)), 17 Carmustine*, 17 Carmustinea (see Carmustine*), 17 Cartrax (see Hydroxyzine HCI), 42 Carubicin (HCl), 17 Carubicin (see Carubicin (HCl)), 17 CAS 10403-51-7 (see Mitindomide), 23 Casodex, 73 Castanospermine, 91 Catapres-TTS (see Clonidine), 40 CB 1639 (see Cycloleucin), 18 CB 304 (see Azaribine*), 15 CB-10-277, 17 CB-1048 (see Chlornaphazine*), 40 CB-1348 (see Chlorambucil*), 17 CB-3025 (see Melphalan* (HCl)), 22 CB-313 (see Mitotane*), 23 CBDCA (see Carboplatin*), 17 CCNU (see Lomustine*), 21 2-CdA, 17, 62 CD4-IgG 91 CD4-PE40, 91 Ce-Cyd (see Cyclopentenylcytosine), 18 CeeNU (see Lomustine*), 21 Ceftriaxone, 95 Celestone (see Betamethasone), 73 Celestone Soluspan (see Betamethasone Acetate), 73 Celestone Soluspan (see Betamethasone Sodium Phosphate), 73

Centrazene (see Simtrazene#), 25 Cephalosprin N (see Adicillin), 39 Cerespann (see Papaverine Disodium), 44 Cerubidine (see Daunorubicin (HCl)*), 18 Cesamet (see Nabilone), 44 Cetacort (see Hydrocortisone), 77 CF (see Leucovorin (calcium)*), 43 CG 30694 (see Edatrexate), 19 CGP 19835A, 62 CGS-16949A (see Fadrazole HCl), 20 Chee-O-Gen (see Ethinyl Estradiol), 76 Chee-O-Genf (see Ethinyl Estradiol), 76 CHIP (see Iproplatin), 21 CHL (see Chlorambucil*), 17 Chloditan (see Mitotane*), 23 Chlodithane (see Mitotane*), 23 Chlor-PZ# (see Chlorpromazine*), 40 Chlorambucil*, 17 Chloraminophene (see Chlorambucil*), 17 Chlorbutin (see Chlorambucil*), 17 Chlorbutine (see Chlorambucil*), 17 Chlorhexidine gluconate (see Peridex), 99 Chloridin (see Pyrimethamine), 45 Chloridine (see Pyrimethamine), 45 Chlormadinone* Acetate, 73 Chlormadinonu (see Chlormadinone* Acetate), 73 Chlornaftina (see Chlornaphazine*), 40 Chlornaphazin (see Chlornaphazine*), 40 Chlornaphazine*, 96 Chlornaphthin (see Chlornaphazine*), 40 2-Chloro-2'Deoxyadenosine (see 2 CdA), 17 Chloroethylcyclohexylnitrosourea (see Lomustine*), 21 Chloromethine-HCl (see Mechlorethamine (HCl)*), 22 Chloronaphthina (see Chlornaphazine*), 40 Chloronaphthine (see Chlornaphazine*), 40 Chloroquinoxaline sulfonamide (see Chlorsulfaquinoxaline), 17 Chlorotrianisene, 74 Chlorotrianisine (see Chlorotrianisene), 74 Chlorotris (see Chlorotrianisene), 74 Chlorozotocin, 17 Chlorpromazine*, 40 Chlorsulfaquinoxaline, 17 Chlortrianizen (see Chlorotrianisene), 74 Cholecalciferol (see Vitamin D), 81 Chromic Phosphate P 32, 74

```
Chromomycin A3#, 17
Chromphosphotope# (see Chromic Phosphate P 32), 74
Chronogyn# (see Danazol), 62
CHX 100 (see Masoprocol), 22
CI 337 (see Azaserine*#), 16
CI 825 (see Pentostatin), 65
CI 904 (see Diaziquone), 19
CI-27,071 (see Descinolone Acetonide#), 74
CI-406 (see Oxymetholone*), 79
CI-673 (see Vidarabine*), 27
CI-880 (miscellaneous agent) (see Amsacrine), 15
CI-881 (see Ametantrone Acetate), 15
CI-882 (see PALA), 24
CI-898 (see Trimetrexate), 27
CI-908 (see Dezaquanine), 19
CI-908 mesylate (see Dezaguanine Mesylate), 19
CI-909 (see Tiazofurin), 26
CI-921, 17
CI-937, 17
Cibacalcin (see Calcitonin*), 39
Cilastatin Sodium, 17
Cimetidine, 40, 92
Cingestol*, 74
Cipro (see Ciprofloxacin), 95
Ciprofloxacin, 40
Cirolemycin*, 17
Cis-Clomiphene citrate (see Clomiphene* Citrate), 74
Cis-DDP (see Cisplatin*), 17
Cis-Diamminedichloroplatinum (II) (see Cisplatin*), 17
Cis-Platin (see Cisplatin*), 17
Cis-Platinum II# (see Cisplatin*), 17
Cisplatin*, 17
Cisplatino (Spanish) (see Cisplatin*), 17
Cisplatyl (see Cisplatin*), 17
Citarin L (see Levamisole HCl*), 64
Citostal (see TMCA), 27
Citrovorum factor (see Leucovorin (calcium)*), 43
 CL (see Cycloleucin), 18
 CL (see Leucovorin (calcium)*), 43
 CL 118,532 (see Triptorelin), 81
 CL 184,116 (see Porfimer Sodium), 24
 CL 26193 (see Simtrazene#), 25
 CL 286,558 (see Zeniplatin), 28
 CL-106359 (see Triamcinolone Acetonide Sodium Phosphate), 81
```

CL-14377 (see Methotrexate*), 22 CL-16,536 (see Puromycin (HCl)), 25 CL-216,942 (see Bisantrene (HCl)), 16 CL-232,315 (see Mitoxantrone (HCl)*), 23 CL-25477 (see Azetepa#), 16 CL-34433 (see Triamcinolone Hexacetonide), 81 CL-34699 (see Amcinonide), 73 CL-61965 (see Triamcinolone Acetonide Sodium Phosphate), 81 CL-67310465 (see Pentostatin), 65 Cladribine (see 2-CdA), 17 Clafen (see Cyclophosphamide*), 18 Claphene (see Cyclophosphamide*), 18 Clarithromycin, 96 CLB (see Chlorambucil*), 17 Clindamycin, 39 Clinoril (anti-inflammatory) (see Sulindac), 46 Clocortolone Acetate, 74 Clocortolone Pivalate, 74 Clocortolone trimethyl acetate (see Clocortolone Pivalate), 74 Clofazimine, 96 Clogestone* Acetate, 74 Clomegestone* Acetate, 74 Clomesone, 17 Clometherone#, 74 Clomiphene A citrate (see Clomiphene* Citrate), 74 Clomiphene* Citrate, 74 Clordion (see Chlormadinone* Acetate), 73 Clorestrolo (see Chlorotrianisene), 74 Cloronaftina (see Chlornaphazine*), 40 Clorotrisin (see Chlorotrianisene), 74 Clotrimazole, 40 CLS (see Calusterone), 73 CLZ (see Chlorozotocin), 17 CMM (see Carubicin (HCl)), 17 CMVIg, 96 CN 15,757 (see Azaserine*#), 16 Co-Vidarabine (see Pentostatin), 65 Codeine, 40 Coenzyme Q, 40 Colaspase (see Asparaginase), 61 Colcemid, 17 Colcemide (see Colcemid), 17 Colchamin (see Colcemid), 17

Colchicine (see Colcemid), 17

Colyte, component of (see Polyethylene Glycol), 44 Combid spansule, 40 Comestrol (see Diethylstilbestrol), 75 Compazine (see Prochlorperazine*), 45 Component of Calcitrel (see Calcium Carbonate), 39 Component of Titralac (see Calcium Carbonate), 39 Compound 112531 (see Vindesine), 28 Compound 133314 (see Trioxifene Mesvlate), 81 Compound 17309 (see Methandrostenolone), 78 Compound 42339 (see Acronine*), 15 Compound 593A (see Piperazinedione compound), 24 Compound Q (see GLQ223), 92 Compound Q, 92 Condylox (see Podofilox), 99 Coparvax (R) (see C-parvum), 62 Copavin (see Papaverine Disodium), 44 Copovithane, 62 Cordarone (see Amiodarone), 39 Cordran (see Flurandrenolide), 76 Corilin, component of (see Glycine), 42 Corstiline (see Adrenocorticotropic hormone), 73 Cort-Dome (see Hydrocortisone), 77 Cortate acetate (see Desoxycorticosterone Acetate), 74 Cortef (see Hydrocortisone), 77 Cortef acetate (see Hydrocortisone Acetate), 77 Cortenema (see Hydrocortisone), 77 Corticotrophin (see Adrenocorticotropic hormone), 73 Corticotropin (see Adrenocorticotropic hormone), 73 Corticotropin, 74 Cortilet (see Fluorometholone), 76 Cortisone (see Cortisone Acetate), 74 Cortisone Acetate, 74 Cortivazol#, 74 Cortogen acetate# (see Cortisone Acetate), 74 Cortone acetate (see Cortisone Acetate), 74 Cortril (see Hydrocortisone), 77 Cortril acetate-AS (see Hydrocortisone Acetate), 77 Cortrophin# (see Adrenocorticotropic hormone), 73 Cortrosyn (see Cosyntropin*), 74 Corynebacterium granulosum (see C-parvum), 62 Corynebacterium parvum (see C-parvum), 62 Cosmegan (see Dactinomycin*), 18 Cosyntropin*, 74

Cotrim (see Bactrim/Septra), 95

Cotrim, component of (see Sulfamethoxazole, Trimethoprim), 46 Coumadin (see Warfarin Sodium*), 46 Cowan I (see Protein A Staphylococcus Aureus), 65 C.P. 15,467-61 (see Lithium Carbonate), 43 CP (see Cyclophosphamide*), 18 CP-16,171 (see Prioxicam), 44 CP-50,556-1 (analgesic) (see Levonantradol HCI), 43 CPAR (see C-parvum), 62 CPD (see Cisplatin*), 17 Cpd 109514 (tranquilizer) (see Nabilone), 44 CPDC (see Cisplatin*), 17 CPDD (see Cisplatin*), 17 CPE-C (see Cyclopentenylcytosine), 18 CPM (see Cyclophosphamide*), 18 CPPD (see Cisplatin*), 17 CPT-11, 17 CPV (see Copovithane), 62 CQS (see Chlorsulfaquinoxaline), 17 CQS, 18 Crasnitin (see Asparaginase), 61 Crein (see Methandrostenolone), 78 Crisnatol (see BW A 773U), 16 Crisnatol mesylate, 17 CSF-GM (H)/Yeast (see CSF-GM), 40 CSF-GM E.coli (Hoechst) (see CSF-GM), 40 CSF-GM, 40, 62 CSF-Granulocyte-Macrophage (see CSF-GM), 40 CSF-Macrophage, 40 CTB (see Cytembena), 18 CTX (see Cyclophosphamide*), 18 CTY (see Cyclophosphamide*), 18 Cuprol (see Diethyldithiocarbamate), 41 Curdlan Sulfate 92 Curretab (see Medroxyprogesterone Acetate), 77 CYC (see Cyclocytidine (HCl)), 17 CYC (see Cyclophosphamide*), 18 Cyclo (see Cyclophosphamide*), 18 Cyclo-C (see Cyclocytidine (HCl)), 17 CycloCMP hydrochloride (see Cyclocytidine (HCl)), 17 Cyclocort (glucocorticoid) (see Amcinonide), 73 Cyclocytidine (HCl), 17 Cyclodisone, 18 Cycloestrol (see Hexestrol), 77 Cyclogesterin# (see Estrogens, Conjugated), 76

Cyclogesterin# (see Progesterone), 80 Cyclohexanecarboxamide, 18 Cycloleucin, 18 Cycloleucine (see Cycloleucin), 18 Cyclopentenylcytosine, 18 Cyclophosphamid (see Cyclophosphamide*), 18 Cyclophosphamide*, 18 Cyclophosphan (see Cyclophosphamide*), 18 Cyclophosphane (see Cyclophosphamide*), 18 Cycloserine, 96 Cyclosporin# (see Cyclosporine), 62 Cyclosporine, 62 Cycrin (see Medroxyprogesterone Acetate), 77 Cyfos# (see Ifosfamide*), 21 CYL (see Cycloleucin), 18 Cynomel (see Liothyronine Sodium), 77 Cyproterone Acetate, 74 Cyren A (see Diethylstilbestrol), 75 CYT (see Cyclophosphamide*), 18 Cytadren (see Aminoglutethimide), 73 Cytarabin (see Cytarabine HCl), 18 Cytarabine HCl, 18, 62 Cytarabine, 51 Cytarabinoside (see Cytarabine HCl), 18 Cytembena, 18 Cytomel (see Liothyronine Sodium), 77 Cytophosphan (see Cyclophosphamide*), 18 Cytosar-U (see Cytarabine HCl), 18 Cytosar-U (see Cytarabine), 51 Cytosine arabinoside hydrochloride# (see Cytarabine HCl), 18 Cytovene (see Ganciclovir), 42, 97 CYTOX (see Cyclophosphamide*), 18 Cytoxan (see Cyclophosphamide*), 18 CYTX (see Cyclophosphamide*), 18 CZT (see Chlorozotocin), 17 D D 7093 (see Mesna*), 43 D-Tetrandrine, 26 D-TrP6 (LH-RH) (see Decapeptyl), 74 D4T, 92 DAC (see 5-Aza-2'Deoxycytidine), 15 DAC (see Decitabine), 18

Dacarbazine*, 18 Dacortin (see Prednisone), 79 Dact (see Dactinomycin*), 18 Dactinomycin D (see Dactinomycin*), 18 Dactinomycin*, 18 DAG (see Dianhydrodulcitol), 19 Dalalone (see Dexamethasone Sodium Phosphate), 75 Dalalone D.P. (see Dexamethasone Acetate), 74 Dalalone L.A. (see Dexamethasone Acetate), 74 Danabol (see Methandrostenolone), 78 Danazol, 62 Danocrine (see Danazol), 62 Dapsone, 95 Darapram (see Pyrimethamine), 45 Daraprim (see Pyrimethamine), 45 Daraprime (see Pyrimethamine), 45 DAS (see Anguidine), 15 DAU (see 3-Deazauridine), 18 Daun (see Daunorubicin (HCl)*), 18 Dauno (see Daunorubicin (HCl)*), 18 Daunoblastin (see Daunorubicin (HCl)*), 18 Daunomycin (see Daunorubicin (HCl)*), 18 Daunomycin-HCl (see Daunorubicin (HCl)*), 18 Daunorubicin (HCl)*, 18 Daunorubicin (liposome encapsulated), 96 Daunorubicin (see Daunorubicin (HCl)*), 18 Daunorubicine (see Daunorubicin (HCl)*), 18 Daunoxone (see Daunorubicin, liposome encapsulated), 96 DAVA (see Vindesine Sulfate), 28 DBD (see Dibromodulcitol), 19 DBM (see Dibromomannitol), 19 DCA (see Desoxycorticosterone Acetate), 74 DCF (see Pentostatin), 65 2'-DCF (see Pentostatin), 65 DCL (see Dichloroallyl lawsone), 19 DCM (see Dichloromethotrexate), 19 DCMTX (see Dichloromethotrexate), 19 DCNU (see Chlorozotocin), 17 DDA (see Dideoxyadenosine). 92 DDC (see Dideoxycytidine), 92 DDC (see Diethyldithiocarbamate), 41 DDI (see Dideoxyinosine), 92 DDMP (see Metoprine), 23

DDP (see Cisplatin*), 17

DDTC (see Diethyldithiocarbamate), 41

Deacytl-N-methylcolchicine (see Colcemid), 17

7-Deaza-adenosine (see Tubercidine), 27

Deazauridine (see 3-Deazauridine), 18

3-Deazauridine, 18

DEB (see Diethylstilbestrol), 75

DECA (see Dexamethasone), 40, 74

Deca-Durabolin (see Nandrolone Decanoate), 78

Decaderm (see Dexamethasone), 40, 74

Decadron Tablets (see Dexamethasone), 40, 74

Decadron-LA Acet (see Dexamethasone Acetate), 74

Decapeptide 1 (Gosrelin) (see Zoladex (see Gosrelin)), 81

Decapeptide 1 (see Gosrelin), 77

Decapeptyl, 74

Decaris (see Levamisole HCl*), 64

Decaspray (see Dexamethasone), 40, 74

Decitabine, 18

Decortin (see Desoxycorticosterone Acetate), 74

Decosterone (see Desoxycorticosterone Acetate), 74

Decostrate# (see Desoxycorticosterone Acetate), 74

DEDC (see Diethyldithiocarbamate), 41

Deferoxamine, 40

Deflamene (see Formocortal), 77

Defosfamide, 40

Degranol (see Mannomustine), 43

Dehydroemetine, 62

7-Dehydroestrone (estrogen) (see Equilin), 75

Dehydromethyltestosterone (see Methandrostenolone), 78

Deladroxone# (see Algestone* Acetophenide), 73

Deladumone (see Estradiol Valerate), 75

Deladumone (see Testosterone Enanthate), 80

Delalutin# (see Hydroxyprogesterone Caproate), 77

Delatestryl (see Testosterone Enanthate), 80

Delcronol (antimetabolite) (see Metoprine), 23

Delestrogen (see Estradiol Valerate), 75

Delmadinone Acetate, 74

Delmeson (see Fluorometholone), 76

Delta-9-THC (see Tetrahydrocannabinol), 46

Delta-Cortef (see Prednisolone), 79

Delta-Dome# (see Prednisone), 79

Deltamycin A4 (see Magnamycin), 22

Deltasone (see Prednisone), 79

Deltra (see Prednisone), 79

Deluteval 2X# (see Estradiol Valerate), 75

Deluteval 2X# (see Hydroxyprogesterone Caproate), 77

Demecolcine (see Colcemid), 17

Demecolcine, 17

Demerol (see Meperidine HC1), 43

Demerol HCl (see Meperidine HCl), 43

4-Demethoxydaunorubicin (see Idarubicin HCl*), 21

Demethylmisonidazole (see Desmethylmisonidazole), 40

Demulan, component of (see Ethinyl Estradiol), 76

Dendrid (see Idoxuridine*), 42

4'-Deoxyadriamycin hydrochloride (see Deoxydoxorubicin HCl), 18

Deoxycoformycin (see Pentostatin), 65

2'-Deoxycoformycin (see Pentostatin), 65

Deoxydoxorubicin HCl, 18

4'-Deoxydoxorubicin (see Deoxydoxorubicin HCl), 18

Deoxynojirimycin, 92

Deoxyspergualin (HCl), 18

15-Deoxyspergualin trihydrochloride (see Deoxyspergualin HCl), 18

Deoxythioguanosine hydrate (see Alpha-TGdR), 15

Deoxythymidine (see Thymidine), 26

DepAndro 100 (see Testosterone Cypionate), 80

DepAndro 200 (see Testosterone Cypionate), 80

DepGynogen (see Estradiol Cypionate), 75

DepMedalone (see Methylprednisolone Acetate), 78

Depo-ACTH# (see Adrenocorticotropic hormone), 73

Depo-ACTH# (see Corticotropin), 74

Depo-Estradiol (see Estradiol Cypionate), 75

Depo-Medrol (see Methylprednisolone Acetate), 78

Depo-Provera (see Medroxyprogesterone Acetate), 77

Depo-Testadiol (see Estradiol Cypionate), 75

Depo-Testadiol (see Testosterone Cypionate), 80

Depo-Testadiol, component of (see Testosterone Cypionate), 80

Depoepovirin# (see Testosterone Cypionate), 80

DepPredalone (see Prednisolone Acetate), 79

Dermairol (see Tretinoin), 51

Deronil (see Dexamethasone), 40, 74

DES (see Diethylstilbestrol), 75

DES-ME (see Desmethylmisonidazole), 40

Desacetylmethylcolchicine (see Colcemid), 17

Desacetylvinblastine amide sulfate (see Vindesine Sulfate), 28

Descinolone Acetonide#, 74

Desmecolcine (see Colcemid), 17

Desmethylmisonidazole, 40

Desmophosphamide (see Defosfamide), 40

Desoxycorticosterone Acetate, 74

Desoxycorticosterone Pivalate, 74

Detorubicin, 18

Detox, 62

Dexacidin (see Dexamethasone), 40, 74

Dexamethasone Acetate, 74

Dexamethasone Dipropionate, 75

Dexamethasone Sodium Phosphate, 75

Dexamethasone, 40, 74

Dexasone (see Dexamethasone), 40, 74

Dexone (see Dexamethasone), 40, 74

Dexormaplatin, 18

Dexrazoxane, 19

Dezaguanine Mesylate, 19

Dezaquanine, 19

DFMO (HCl) (see DFMO), 19

DFMO (see Eflornithine HCl), 96

DFMO, 19

DHAC, 19

DHAD (see Mitoxantrone (HCl)*), 23

DHAQ (see Mitoxantrone (HCl)*), 23

DHPG (see Ganciclovir), 42, 97

Di-Estryl (see Diethylstilbestrol), 75

Diacetoxyscirpenol (see Anguidine), 15

Diacetylmorphine dihydrochloride (see Heroin), 42

Diaminopyritamin (see Pyrimethamine), 45

3,5-Diamino-s-triazole (see Guanazole), 21

Diamorphine HCl (see Heroin), 42

Diamox (carbonic anydrase inhibitor) (see Acetazolamide), 39

Dianabol# (see Methandrostenolone), 78

Dianhydrodulcitol, 19

Dianhydrogalactitol (see Dianhydrodulcitol), 19

Dianoxide Furoate, 96

Diaziquone, 19

Diazoacetylserine (see Azaserine*#), 16

6-Diazo-5-Oxo-L-Norleucine (9CI) (see DON), 19

Dibestil# (see Diethylstilbestrol Dipropionate), 75

Dibromannit (see Dibromomannitol), 19

Dibromdulcit (see Dibromodulcitol), 19

Dibromodulcitol, 19

Dibromomannitol, 19

DIC (see Dacarbazine*), 18

Dichloroallyl lawsone, 19

Dichloroamethopterin (see Dichloromethotrexate), 19

Dichlorolapachol (see Dichloroallyl lawsone), 19

DichloroMethotrexate (see Dichloromethotrexate), 19

Dichloromethotrexate, 19

Dichloromethyl-diethylamine-HCl (see Mechlorethamine (HCl)*), 22

Didanosine (see Dideoxyinosine), 92

Didemnin B, 19

Dideoxyadenosine, 92

Dideoxycytidine, 92

2',3'-dideoxycytidine (see Dideoxycytidine), 92

2',3'-dideoxyinosine (see Dideoxyinosine), 92

Dideoxyinosine, 92

Dideoxynucleoside (thymidine) analogue (see D4T), 92

Didronel (see Etidronate Disodium), 41

Dienestrol, 75

Diethyldithiocarbamate, 41

Diethylstilbestrol Diphosphate, 75

Diethylstilbestrol Dipropionate, 75

Diethylstilbestrol, 75

Diflucan (see Fluconazole), 41, 97

Diflucortolone* Pivalate, 75

Diflucortolone*, 75

Difluoromethylornithine (see DFMO), 19

Diglycoaldehyde, 19

Dihematoporphyrin Ether (see Porfimer Sodium), 24

Dihydro-5-Azacytidine HCl, 19

5,6-Dihydro-5-azacytidine hydrochloride (see DHAC), 19

Dihydroazacytidine (see DHAC), 19

Dihydrolenperone, 19

Dihydroxyanthracenedione dihydrochloride (see Mitoxantrone (HCl), 23

1,25-Dihydroxy-vitamin D3, 51

Dihydro-lenperone (see Dihydrolenperone), 19

Dilaudid (analgesic; narcotic) (see Hydroxyzine HCI), 42

Diltiazem HCl. 41

Dimepranol Acedoben, 41

Dimethyl Sulfoxide, 51

Dimethyl-triazeno-imidazole-carboxamide (see Dacarbazine*), 18

Dimethylbusulfan, 62

Dimethyltestosterone (see Calusterone), 73

Dimethylurethimine (see Meturedepa), 23

Dimonidazole, 41

Dinacrin# (see Isoniazid*), 97

Dinitrochlorobenzene, 62

Dinitrofluorobenzene, 62

Diogyn (see Estradiol), 75

Diogyn E (see Ethinyl estradiol), 76

Diogynets (see Estradiol), 75

Diphaston (see Dydrogesterone*), 75

Diphenhydramine HCl, 41

Diphenidol, 41

Diprolene (see Betamethasone Dipropionate), 73

Diprosone (see Betamethasone Dipropionate), 73

Dipyridamole, 41

Disal (see Furosemide*), 41

Dispermine (see Piperazine), 44

Disulfiram (see Antabuse), 91

Ditate# (see Estradiol Valerate), 75

Ditate# (see Testosterone Enanthate), 80

Dithiocarb (see Diethyldithiocarbamate), 41

Ditubin# (see Isoniazid*), 97

Divema (see Pyran copolymer), 65

DL Sarcolysine (see Melphalan* (HCl)), 22

DM (see Dexamethasone), 40, 74

4-DMDR (see Idarubicin HCl*), 21

DMPA (see Medroxyprogesterone Acetate), 77

DNFB (see Dinitrofluorobenzene), 62

DNM (see Daunorubicin (HCl)*), 18

DNR (see Daunorubicin (HCl)*), 18

DOCA (see Desoxycorticosterone Acetate), 74

Doca acetate# (see Desoxycorticosterone Acetate), 74

Dolophine HCl (see Methadone HCl), 43

Domestrol (see Diethylstilbestrol), 75

Domperidone, 41

DON, 19

Dopmedrol (see Methylprednisolone), 78

Dorcostrin# (see Desoxycorticosterone Acetate), 74

DOXO (see Desoxycorticosterone Acetate), 74

Doxorubicin (HCl)*, 19

Doxorubicin (liposome-encapsulated), 96

Doxorubicin-DNA complex, 19

DRB (see Daunorubicin (HCl)*), 18

Drolban# (see Dromostanolone Propionate), 75

Dromostanolone Propionate, 75

Dronabinol, 96

Droperidol*, 41

Drostanolone (see Oxymetholone*), 79

Drostanolone propionate (see Dromostanolone Propionate), 75

Droxone# (see Algestone* Acetophenide), 73

DSM (see Dexamethasone), 40, 74

DT (see Thymidine), 26

DThyd (see Thymidine), 26 DTIC (see Dacarbazine*), 18 DTIC-Dome (see Dacarbazine*), 18 DTR (see Detorubicin), 18 Dualar# (see Benzodepa), 16 Duazomycin, 19 Duazomycin-B (see Azotomycin), 16 Duborimycin, 19 Dufaston, (see Dydrogesterone*), 75 Dulcitol diepoxide (see Dianhydrodulcitol), 19 Dumogran (see Methyltestosterone), 78 Dup 785 (see Brequinar Sodium), 16 Duphaston (see Dydrogesterone*), 75 Durabolin (see Nandrolone Phenpropionate), 79 Duramorph (see Morphine Sulfate), 44 Duvaron (see Dydrogesterone*), 75 DV (see Dienestrol), 75 DVA (see Vindesine Sulfate), 28 DxDx (see Deoxydoxorubicin HCl), 18 4-DxDx (see Deoxydoxorubicin HCl), 18 DXM (see Dexamethasone Acetate), 74 Dydrogesterone*, 75 Dyloform (see Ethinyl estradiol), 76

E

E-39 (see Inproquone), 21 EC 3.1.5.1 (see (Escherichia coli)), 61 EC 3.1.5.1 (see Asparaginase), 61 Echinomycin, 19 Ecloril (see Chlorambucil*), 17 Econazole, 96 Edam-10, 19 Edatrexate, 19 Edecrin (see Ethacrynic Acid), 41 Eflornithine HCl, 19, 96 Efudex (see Fluorouracil*), 20 EHB 776 (see Foscarnet Sodium), 97 Eldisine (see Vindesine Sulfate), 28 Elipten (see Aminoglutethimide), 73 Elliot's B solution, 41 Elobromol (see Dibromodulcitol), 19 Elsamitrucin, 19 Elspar (see Asparaginase), 61

Emanil (see Idoxuridine*), 42 Emcyt (see Estramustine Phosphate Sodium), 76 Emdisterone (see Dromostanolone Propionate), 75 EMET (see Emetine), 19 Emetin (see Emetine), 19 Emetine monohydrochloride (see Emetine), 19 Emetine, 19 Emofolin (sodium), 20 EN-1639A (see Naltrexone), 44 Enarax (see Hydroxyzine HC1), 42 Endorphin, 41 Endoxan (see Cyclophosphamide*), 18 Endoxana (see Cyclophosphamide*), 18 Enduxan (see Cyclophosphamide*), 18 Enovid (see Mestranol), 78 Enpromate#, 20 ENT 26079 (see Aminopterin), 15 ENT 50852 (see Altretamine), 15 ENT-50451 (see Benzodepa), 16 ENT-50825 (see Porfiromycin), 25 EO-26 (see 3-Deazauridine), 18 EPEG (see Etoposide*), 20 EPI (see Epirubicin (HCl)), 20 Epidropal (see Allopurinol*), 39 Epipodophyllotoxin (see Teniposide), 26 Epipodophyllotoxin ethylidene glucopyranoside (see Etoposide*) 20 Epipropidine#, 20 Epirubicin (HCl), 20 EPO (see Erythropoietin), 62, 96 Epodyl (see Ethoglucid), 20 Epoetin Alfa, 62, 96 Epogen (see Erythropoietin), 62, 96 Eponate (see Epipropidine#), 20 Epoxypropidine (see Epipropidine#), 20 Eprox (see Erythropoietin), 62, 96 EPT (see Teniposide), 26 Equilin, 75 Eraverm (see Piperazine), 44 Erbaprelina (see Pyrimethamine), 45 Ergamisol (see Levamisole HCl*), 64 Erysan (see Chlornaphazine*), 40 Erythropoietin, 62, 96 Eskalith (see Lithium Carbonate), 43

Eskatrol (see Prochlorperazine*), 45

Ethiodized Oil*, 41

Esorubicin HCl (see Deoxydoxorubicin HCl), 18 Estan# (see Dienestrol), 75 Esteed (see Ethinyl estradiol), 76 Estigyn (see Ethinyl estradiol), 76 Estilbin (see Diethylstilbestrol), 75 Estinyl (see Ethinyl estradiol), 76 Eston-E (see Ethinyl estradiol), 76 Estoral (see Ethinyl estradiol), 76 Estrace (see Estradiol), 75 Estracyt (see Estramustine Phosphate Sodium), 76 Estraderm TTS (see Estradiol), 75 Estradiol Benzoate, 75 Estradiol Cypionate, 75 Estradiol Dipropionate, 75 Estradiol Mustard, 75 Estradiol Valerate, 75 Estradiol, 75 Estradurin (see Polyestradiol phosphate), 79 Estrafon (see Perphenazine), 44 Estramustine phosphate disodium (see Estramustine Phosphate Sodium), 76 Estramustine Phosphate Sodium, 76 Estramustine, 75 Estratab (see Estrogens, Esterified), 76 Estratest (see Methyltestosterone), 78 Estratest, (see Estrogens, Esterified), 76 Estrifar (see Hexestrol), 77 Estriol, 76 Estrobene (see Diethylstilbestrol), 75 Estrogens, Conjugated, 76 Estrogens, Esterified, 76 Estronal (see Hexestrol), 77 Estrone, 76 Estropipate, 76 Estrovis (see Quinestrol), 80 Etanidazole, 41 Ethacrynic Acid, 41 Ethambutol HCl, 96 Ethanesulfonic acid compound (see Baker's antifol), 16 Ethidium chloride, 20 Ethidol (see Ethinyl estradiol), 76 Ethinoral# (see Ethinyl estradiol), 76 Ethinyl Estradiol, 76

Ethiodol-131# (radio-opaque medium) (see Ethiodized Oil*), 41

Ethiofos# (see Amifostine), 39 Ethionamide, 96 Ethoglucid, 20 Ethyl carbamate# (see Urethane), 27 9-Ethyl-6-Mercaptopurine, 20 9-Ethyl-6-MP (see 9-Ethyl-6-Mercaptopurine), 20 Ethylchlorformate, 62 2-ethylhydrazide (see Mitopodozide), 23 Ethylidine-Lignan-P (see Etoposide*), 20 Ethynylestrenol (see Lynestrenol#), 77 Ethyol (see Amifostine), 39 Eticyclin (see Ethinyl estradiol), 76 Eticylol# (see Ethinyl estradiol), 76 Etidronate Disodium, 41 Etinestrol (see Ethinyl estradiol), 76 Etoglucid (see Ethoglucid), 20 Etoposide*, 20 Etoprine, 20 Etrenol (see Hycanthone (mesylate)), 21 Etretinate, 51 Etrosyn (see Diethylstilbestrol), 75 EU-3120 (see Acodazole (HCl)), 15 Eudyna (see Tretinoin), 51 Eulexin (see Flutamide), 77 Euthroid (see Liotrix), 77 Exluton (see Lynestrenol#), 77 Extra-Plex (see Hexestrol), 77

F

F3TDR, 20
FAA (see Flavone acetic acid), 20
Fadrazole HCl, 20
Famciclovir, 96
2-FAMP (see Fludarabine Phosphate), 20
Fansidar, component of (see Pyrimethamine), 45
Fansidar, 41
Farlutin (see Medroxyprogesterone Acetate), 77
FAS (see 2-Fluoroadenosine#), 20
2 FAS (see 2-Fluoroadenosine#), 20
Fazarabine, 20
5-FC (see Flucytosine), 97
FC 1157a (antiglucocorticoid) (see Toremifene Citrate), 81
FCA (see Freund's adjuvant), 62

Fedibaretta (see Phetharbital), 24

Feldene (anti-inflammatory) (see Piroxicam), 44

Feminone (see Ethinyl estradiol), 76

Femstat (see Butoconazole), 95

Fenesterin (see Phenesterine), 24

Fenestrin (see Phenesterine), 24

Fenobolin (see Nandrolone Phenpropionate), 79

Fenretinide, 20

Fentanyl Citrate, 41

FIAU 96

Filgrastim (see G-CSF), 97

Filgrastim, 62

Flagyl (see Metronidazole), 98

Flagyl, component of, I.V. RTU (see Metronidazole), 98

Flavone acetic acid, 20

Flavone-8-Acetic Acid (see Flavone acetic acid), 20

Florafur (see Tegafur), 26

Floridine (see Sodium Fluoride), 45

Florinef Acetate (see Fludrocortisone Acetate), 76

Floxin (see Ofloxacin), 98

Floxuridin (see Floxuridine*), 20

Floxuridine*, 20

FLT (see Fluorothymidine), 92

FLU (see Fluoxymesterone), 76

Fluaid (see Tegafur), 26

Fluconazole, 41, 97

Flucytosine, 97

Fludara (see Fludarabine Phosphate), 20

Fludarabine Phosphate, 20

Fluderma (see Formocortal), 77

Fludestrin (see Testolactone*), 80

Fludrocortisone Acetate, 76

Flumethasone*, 76

2-Fluoroadenine arabinoside 5-Phosphate (see Fludarabine Phosphate), 20

2-Fluoroadenosine#, 20

2-Fluoro-ARA AMP (see Fludarabine Phosphate), 20

Fluorodopan (see Fluorodopane), 20

Fluorodopane, 20

Fluorofur (see Tegafur), 26

Fluorol (see Sodium Fluoride), 45

Fluoromethalone (see Fluorometholone), 76

Fluorometholone, 76

Fluoroplex (see Fluorouracil*), 20

Fluorothymidine, 92

Fluorouracil*, 20

5-Fluorouracil (see Fluorouracil*), 20

5-Fluorouracil deoxyriboside (see Floxuridine*), 20

Fluorpan (see Fluorodopane), 20

Fluosol-DA, 41

Fluotestin (see Fluoxymesterone), 76

Fluoximesterone (see Fluoxymesterone), 76

Fluoxymesterone, 76

Fluperolone Acetate, 76

Fluprednisolone* Valerate#, 76

Fluprednisolone*, 76

Fluracil (see Fluorouracil*), 20

5-Fluracil (see Fluorouracil*), 20

Flurandrenolide, 76

Flurandrenolone# (see Flurandrenolide), 76

Flurbiprofen, 41

Fluril (see Fluorouracil*), 20

Flurocitabine#, 20

5-Flurodeoxyuridine (see Floxuridine*), 20

Flurogestone Acetate, 76

Flusteron (see Fluoxymesterone), 76

Flutamide, 77

Flutestos (see Fluoxymesterone), 76

Flutone# (see Triamcinolone Acetonide), 81

FML Forte (see Fluorometholone), 76

FML Liquifilm (see Fluorometholone), 76

FML S.O.P. (see Fluorometholone), 76

FML-S Liquifilm (see Fluorometholone), 76

Folex (see Methotrexate*), 22

Foligan (see Allopurinol*), 39

Folinic acid (see Leucovorin (calcium*)), 43

Fonatol (see Diethylstilbestrol), 75

Formocortal, 77

Formula 405 (see Pregnenolone* Succinate), 80

Foscarnet Sodium, 97

Foscavir (see Foscarnet Sodium), 97

Fosfestrol (see Diethylstilbestrol Diphosphate), 75

Fosquidone, 20

Fostriecin Sodium, 20

Fotemustine, 20

Fourneau 309 (see Suramin Sodium), 26

Freund's adjuvant, 62

FT (see Tegafur), 26

FT-106 (see Doxorubicin (HCl)*), 19

FT-207 (see Tegafur), 26

Ftorafur (see Tegafur), 20, 26
FU (see Fluorouracil*), 20
5-FU (see Fluorouracil*), 20
FUDR (see Floxuridine*), 20
5-FUDR (see Floxuridine*), 20
Fungizone (see Amphotericin B), 95
Furacin (see Nitrofurazone), 23
Furazolidone, 97
5-FURD (see Floxuridine*), 20
Furosemide*, 41
Furoxone (see Furazolidone), 97
Futraful (see Tegafur), 26
FVA (see Flavone acetic acid), 20
Fx5 (see Thymosin), 66
FXM (see Fluoxymesterone), 76

G

G-CSF (Granulocyte Colony-Stimulating Factor), 41, 97 G-CSF, 41, 97 Galactitol (see Dianhydrodulcitol), 19 Gallium Nitrate, 62 Gallium trinitrate (see Gallium Nitrate), 62 Gamma Interferon, 97 Gammaguard (see IVIG), 64 Gammaphos (see Amifostine), 39 GAN (see Gallium Nitrate), 62 Ganciclovir, 42, 97 Ganglioside, 62 Ganite (see Gallium Nitrate), 62 Gantanol, component of (see Sulfamethoxazole), 46 Geabol (see Methandrostenolone), 78 Gemcitabine HCl, 21 Gemcitabine, 21 Genoxal (see Cyclophosphamide*), 18 Germanin (see Suramin Sodium), 26 Gestaclone, 77 Gestafortin (see Chlormadinone* Acetate), 73 Gestapuran (see Medroxyprogesterone Acetate), 77 Gestatron (see Dydrogesterone*), 75 Gesterol 50 (see Progesterone), 80 Gestonorone Caproate, 77 Gestoral LA 250 (see Hydroxyprogesterone Caproate), 77 Ginestrene (see Ethinyl estradiol), 76

GLQ223, 92 Glycidyl isocyanurate (see Teroxirone), 26 Glycine, 42 GM-CSF, 42, 92 GM2 (see Ganglioside), 62 Gold Au 198, 42 Gosrelin, 77 Gotax (see Allopurinol*), 39 GP160 vaccine, 92 GR 38032 F (antagonist) (see Serotonin), 45 GR 38032F (see Ondansetron HCl), 44 GR 63178A, 21 GR 63178K (see Fosquidone), 20 Grafestrol (see Diethylstilbestrol), 75 Granisetron, 42 Granulocyte colony stimulating factor (see G-CSF), 41 Granulocyte macrophage colony stimulating factor (see GM-CSF), 42 Granulocyte-colony stimulating factor (see CSF-GM), 40 GRF 38032F (see Ondansetron HCl), 44 Guanazol (see 8-Azaguanine), 15 Guanazole, 21 Guanazolo (see 8-Azaguanine), 15 Guaneran (see Thiamiprine), 26 Guanfacine HCl, 42 Guanfacine, 97 Gynogen L.A. (estrogen) (see Estradiol Valerate), 75

Н

H.B.F. 386 (see Cactinomycin), 16
H37Ra (see C-parvum), 62
Haffkinine (see Quinacrine (HCl)), 45
HAL (see Fluoxymesterone), 76
Haldol (see Haloperidol), 42
Haldrone (see Paramethasone* Acetate), 79
Halodrin (see Ethinyl estradiol), 76
Halodrin (see Fluoxymesterone), 76
Haloperidol, 42
Haloprogesterone, 77
Halotestin (see Fluoxymesterone), 76
HC (see Hydrocortisone), 77
HDA (see Cytarabine HCl), 18
HDARA-C (see Cytarabine HCl), 18
HDC (see Hydrocortisone Sodium Succinate), 77

He-PC (see Miltefostine), 23 HEDP (see Rhenium-186), 45 Hematoporphyrin Derivative, 42 Hemel (see Altretamine), 15 Henkel's compound (see Teroxirone), 26 Hepsulfam, 21 Heptanamide (see Deoxyspergualin (HCI)), 18 1.7-Heptanedivlsulfamate ester (see Hepsulfam), 21 Heroin HCl, 42 Heroin hydrochloride (see Heroin), 42 Heroin, 42 Herpesil (see Idoxuridine*), 42 Herpidu (see Idoxuridine*), 42 Herplex (see Idoxuridine*), 42 Hexa-Betalin (see Pyridixone HCl), 45 Hexadecylphosphocholine (see Miltefostine), 23 Hexadrol (see Dexamethasone Sodium Phosphate), 75 Hexadrol Elixir (see Dexamethasone), 40, 74 Hexadrol Tablets (see Dexamethasone), 40, 74 Hexahydropyrazine (see Piperazine), 44 Hexalen (see Altretamine*), 15 Hexamethylene bisacetamide (see HMBA), 51 Hexamethylene diacetamide (see HMBA), 51 Hexamethylmelamine, 21 Hexanoestrol (see Hexestrol), 77 Hexestrofen (see Hexestrol), 77 Hexestrol, 77 Hexoestrol (see Hexestrol), 77 HH (see Homoharringtonine), 21 Hi-Bestrol (see Diethylstilbestrol), 75 Hidrix (see Hydroxyurea*), 21 HIV immunogen, 92 Hivid (see Dideoxycytidine), 92 HLTV antibody, 62 HMBA, 51 HMD (see Oxymetholone*), 79 HMM (see Altretamine), 15 HMM (see Hexamethylmelamine), 21 HN2 (see Mechlorethamine (HCl)*), 22 Ho 1/193 (see Triethylenemelamine), 27 Hoe 33258 (see Pibenzimol HCl), 24 Hoe 766 (see Busrelin Acetate), 73 Hoechst No. 33258 (see Pibenzimol HCl), 24 Holoxan 1000 (see Ifosfamide*), 21

Homandren (see Methyltestosterone), 78 Homidium chloride (see Ethidium chloride), 20 Homofolic acid (see Emofolin (sodium)), 20 Homoharringtonine, 21 Hormoestrol (see Hexestrol), 77 Hormonisene (see Chlorotrianisene), 74 5-HP (see 5-Hydroxypicolinaldehyde thiosemicarbazone), 21 HPA-23, 93, 94 HPMPC, 97 HPP (see Allopurinol*), 39 4-HPR (see 4-Hydroxyphenyl retinamide), 51 HU (see Hydroxyurea*). 21 Human fibroblast interferon (see Immune RNA), 63 Human fibroblast interferon (see Interferon and interferon inducers), 63 Human Jurkat Tumor Derived (see Interleukin-2), 63 Human leukocyte interferon (see Interferon and interferon inducers), 63 Human lymphoblastoid interferon (BW) (see Interferon Alfa-n1), 63 Humatin (see Paromomycin Sulfate), 98 HUR (see Hydroxyurea*), 21 HXM (see Hexamethylmelamine), 21 Hycamptamine (see Topotecan HCl), 27 Hycanthone (mesylate), 21 Hycanthone methanesulfonate (see Hycanthone (mesylate)), 21 Hycanthone monomethanesulfonate (see Hycanthone (mesylate)), 21 HYD (see Hydroxyurea*), 21 Hydeltra (see Prednisolone), 79 Hydeltra-T.B.A. (see Prednisolone Tebutate), 79 Hydeltrasol (see Prednisolone Sodium Phosphate), 79 Hydrazine Sulfate, 42 Hydrea (see Hydroxyurea*), 21 Hydreia (see Hydroxyurea*), 21 Hydrocortisone Acetate, 77 Hydrocortisone Sodium Phosphate, 77 Hydrocortisone Sodium Succinate, 77 Hydrocortisone, 77 Hydrocortone Phosphate (see Hydrocortisone Sodium Phosphate), 77 Hydrogesterone (see Dydrogesterone*), 75 Hydromorphone HCl, 42 4'-Hydroperoxycyclophosphamide, 62 Hydroximic acid (see Hydroxyurea*), 21 Hydroxycarbamide (see Hydroxyurea*), 21 Hydroxycarbamine (see Hydroxyurea*), 21 Hydroxychloroquine, 92 14-hydroxydaunomycin (see Doxorubicin (HCI)*), 19

Hydroxyethylidene diphosphonate (see Rhenium-186), 45 Hydroxylurea (see Hydroxyurea*), 21 4-Hydroxyphenyl retinamide, 51 5-Hydroxypicolinaldehyde thiosemicarbazone, 21 Hydroxyprogesterone Caproate, 77 Hydroxyurea*, 21 Hydroxyzine HC1, 42 Hydura (see Hydroxyurea*), 21 Hyperbaric Oxygen, 42 Hypericin, 92 I I 131, 63 I 131 Idotope (see Sodium Iodide), 45 I-Dox (see Deoxydoxorubicin HCl), 18 Ibenzmethyzin hydrochloride (see Procarbazine* (HCl)), 25 Ibiothyron (see Liothyronine Sodium), 77 IBZ (see Procarbazine* (HCl)), 25 ICDT (see Dacarbazine*), 18 ICI 118 (see Gosrelin), 77 ICI 137233 (see Aphidicolin Glycinate), 15 ICI-32865 (see Ethoglucid), 20 ICI-46,474 (see Tamoxifen Citrate), 80 ICI-59118 (see Razoxane), 25 ICIG 1109 (alkylating agent) (see Lomustine*), 21 ICRF Isomer (see ICRF-187), 21 ICRF-159 (see Razoxane), 25 ICRF-187 (see Dexrazoxane), 19 ICRF-187, 21 ICS-205930 (see Tropisetron), 46 ICT (see Dacarbazine*), 18 IDA (see Idarubicin HCl*), 21 Idamycin (see Idarubicin HCl*), 21 Idarubicin HCl*, 21 Idexur (see Idoxuridine*), 42 Idoxuridin (see Idoxuridine*), 42 Idoxuridine*, 42 IDU (see Idoxuridine*), 42 Idu-Oculos (see Idoxuridine*), 42 Idulea (see Idoxuridine*), 42 IDUR (see Idoxuridine*), 42 5-IUDR (see Idoxuridine*), 42 IdUrd (see Idoxuridine*), 42

Iduridin (see Idoxuridine*), 42 IF (see Interferon and interferon inducers), 63 Ifex (see Ifosfamide*), 21 IFN (see Immune RNA), 63 IFN (see Interferon and interferon inducers), 63 Ifosfamid (see Ifosfamide*), 21 Ifosfamide*, 21 IFX (see Ifosfamide*), 21 IL-2 (see Interleukin-2), 63, 93 IL-2 + IL-4 (Cetus & Sterling) (see Interleukin-2), 64 Il-2 Jurkat Derived (see Interleukin-2), 63 IL-2 Recombinant Interleukin-2 (see Interleukin-2), 63 Ilmofosine, 21 Imexon, 63 IMI 58 (see Deoxydoxorubicin HCl), 18 IMI-28 (see Epirubicin (HCl)), 20 IMI-30 (see Idarubicin HCl*), 21 Imidazole mustard (see TIC-mustard), 27 Imidazole-1-Acetamide (see SR 2508), 45 Imididazole carboximide (see TIC-mustard), 27 Imipemide (see Imipenem), 21 Imipenem, 21 Imipenem-Cilastatin, 42, 97 Immune globulin IV, 93 Immune interferon (see Interferon Alfa-n3), 63 Immune RNA, 63 IMPY (see Pyrazolo-Imidazole compound), 25 Imreg-1, 92 Imunovir (see Isoprinosine), 42 Imuran (see Azathioprine*), 16 Imurek (see Azathioprine*), 16 Imurel (see Azathioprine*), 16 Imuthiol, 92 Imuvert, 63 Inapsine (see Droperidol*), 41 INDI (see Indicine-N-oxide), 21 Indicine-N-oxide, 21 Indocin (anti-inflammatory (see Indomethacin*), 42 Indomethacin*, 42 Inestra (see Ethinyl estradiol), 76 Inflamase (see Prednisolone Sodium Phosphate), 79 INH (see Isoniazid*), 97 INO (see Indicine-N-oxide), 21 Inosine, 42

Inosine dialdehyde (see Diglycoaldehyde), 19 5'-Inosinic acid polymers, 65 INOX (see Diglycoaldehyde), 19 Inprochone (see Inproquone), 21 Inproquone, 21 Interferon Alfa-2a, 63 Interferon Alfa-2b, 63 Interferon Alfa-n1, 63 Interferon Alfa-n3, 63 Interferon and interferon inducers, 63 Interferon Gamma-1b, 63 Interleukin, 64 Interleukin-1 alpha (see Interleukin-1), 63 Interleukin-1 beta (see Interleukin-1), 63 Interleukin-1, 63 Interleukin-2 Fusion Toxin, 93 Interleukin-2 liposome, 64 Interleukin-2, 63, 92 Interleukin-4, 64 Interleukin-6, 64 Intron A (see Interferon Alfa-2b), 63 Iododeoxyuridine (see Idoxuridine*), 42 Iodoquinol, 97 IPA (see Riboprine*), 25 Iphosphamid (see Ifosfamide*), 21 Iphosphamide (see Ifosfamide*), 21 Ipomeanol, 64 4-Ipomeanol (see Ipomeanol), 64 IPP (see Ifosfamide*), 21 Iproplatin, 21 Ismipur (see Mercaptopurine*), 22 Isoendoxan (see Ifosfamide*), 21 Isofosfamide (see Ifosfamide*), 21 Isolyn# (see Isoniazid*), 97 Isoniazid*, 97 Isophosphamide (see Ifosfamide*), 21 Isopregnenone (see Dydrogesterone), 75 Isoprinosine, component of (see Dimepranol Acedoben), 41 Isoprinosine, component of (see Inosine), 42 Isoprinosine, 42, 97 Isopropamide Iodide, 42 Isoptin (see Verapamil (HCl)*), 46 Isoptin hydrochloride (see Verapamil (HCl)*), 46 Isotretinoin* (see Retinoic acid), 51

Italchine (see Quinacrine (HCl)), 45 ITG (see Thiamiprine), 26 Itraconazole, 42, 97 ITZ (see Itraconazole), 42, 97 IUdR (see Idoxuridine*), 42 IVIG, 64, 97 Izoptin (see Verapamil (HCl)*), 46

J

JB-11, 21 JM-8 (see Carboplatin*), 17 JM-9 (see Iproplatin), 21 Joacamine (see Busulfan*), 16 Joddeoxyuridin (see Idoxuridine*), 42

K

K Predne-Dome# (see Prednisolone), 79
Karminomitsin (see Carubicin (HCl)), 17
Karminomycin HCl (see Chlorambucil*), 17
Kenacort (see Triamcinolone), 81
Kenacort Diacetate syrup (see Triamcinolone Diacetate), 81
Kenalog (see Triamcinolone Acetonide), 81
Kerecid (see Idoxuridine*), 42
Ketoconazole, 77, 97
Ketorolac Tromethamine, 42
Ketrax (see Levamisole HCl*), 64
Khlortrianizen (see Chlorotrianisene), 74
Kolchamin (see Colcemid), 17
Kolkamin (see Colcemid), 17
Komed, component of (see Sodium Thiosulfate), 45
Kymarabine (see Fazarabine), 20

L

L-3428 (see Amiodarone), 39 L-5103 (see Rifampin), 25 L-693,989, 97 L-69739, 93 L-697661, 93 L-Alanosine (see Alanosine), 15 L-ASE (see (Escherichia coli)), 61

L-ASE (see Asparaginase), 61 L-ASP (see Asparaginase), 61 L-Asparaginase (see Asparaginase), 61 L-Asparagine amidohydrolase (see Asparaginase), 61 L-Buthionine sulfoximine (see BSO), 42 L-Leucovorin, 43 L-PAM (see Melphalan* (HCl)), 22 L-Phenylalanine mustard (see Melphalan* (HCl)), 22 L-Sarcolysin (see Melphalan* (HCl)), 22 L-Serine diazo-acetate ester (see Azaserine*#), 16 L-Tetramisole HCL (see Levamisole HCI*), 64 Ladakamycin (see Azacitidine), 51 LAK cells, 64 Lamivudine (see 3TC), 94 Laniazid (see Isoniazid), 97 Lasix (see Furosemide*), 41 LB-502 (diuretic) (see Furosemide*), 41 LCR (see Vincristine* Sulfate), 28 Lederfoline (see Leucovorin (calcium)*), 43 Lentinan, 64, 93 Leo 114 (see Polyestradiol phosphate), 79 Leo 275 (see Estramustine), 75 Letrazuril, 98 Leucomycin (see Spiramycin), 45 Leucovorin (calcium)*, 43 Leukaemomycin C (see Daunorubicin (HCl)*), 18 Leukeran (see Chlorambucil*), 17 Leukerin (see Mercaptopurine*), 22 Leukersan (see Chlorambucil*), 17 Leukine (see GM-CSF), 97 Leukine (see Sargramostim), 66 Leukoran (see Chlorambucil*), 17 Leunase (see Asparaginase), 61 Leuprolide Acetate, 77 Leuprorelin (see Leuprolide Acetate), 77 Leupurin (see Mercaptopurine*), 22 Leurocristine sulfate (see Vincristine* Sulfate), 28 Leurosidine sulfate (see Vinrosidine Sulfate#), 28 Leurosine (see Vinleurosine Sulfate#), 28 Leustatin (see 2-CdA), 17 LEV (see Levamisole HCl*), 64 Levamisole HCl*, 64 Levlen (see Ethinyl estradiol), 76

Levomysol HCL (see Levamisole HCl*), 64

Levothroid (see Levothyroxine Sodium), 77 Levothyroxine Sodium, 77 Lidocaine, 43 Lidomantle-HC, component of (see Lidocaine), 43 Linestrenol (see Lynestrenol#), 77 Linfilizin (see Chlorambucil*), 17 Linfolysin (see Chlorambucil*), 17 Liothyronine Sodium, 77 Liotrix, 77 Lipo-Lutin# (see Progesterone), 80 Liposome-Doxorubicin Kit, 21 Liquifilm (see Idoxuridine*), 42 Litaler (see Hydroxyurea*), 21 Litalir (see Hydroxyurea*), 21 Lithane (see Lithium Carbonate), 43 Lithium Carbonate, 43 Lithobid (see Lithium Carbonate), 43 Lithonate (see Lithium Carbonate), 43 Lithotabs (see Lithium Carbonate), 43 Lo/Ovral (see Ethinyl estradiol), 76 Loestrin (see Ethinyl estradiol), 76 Lometrexol Sodium, 21 Lomidine (see Pentamidine), 98 Lomustine*, 21 Lonidamine, 43 Lopurin (see Allopurinol*), 39 Lorazepam, 43 Lormin (see Chlormadinone* Acetate), 73 Lotrimax, component of (see Clotrimazole), 96 Lotrimin (see Clotrimazole), 96 Lotrisone (see Betamethasone Dipropionate), 73 Lotrisone, component of (see Clotrimazole), 96 Lovastatin, 21 Lowers blood (see Indomethacin*), 42 Loxoribine, 43 Lumbrical (see Piperazine), 44 Lupron Depot (TAP) (see Leuprolide Acetate), 77 Lupron TAP (see Leuprolide Acetate), 77 Lutinyl (see Chlormadinone* Acetate), 73 Lutopolar (see Medroxyprogesterone Acetate), 77 Lutoral, (see Medroxyprogesterone Acetate), 77 LY 119863 (see Vinepidine Sulfate). 28 LY 186641 (see Sulofenur), 26 LY 264618 disodium (see Lometrexol Sodium), 21

LY 28002 (see Epipropidine#), 20
LY-099094 (see Vindesine Sulfate), 28
LY-104208 (see Vinzolidine Sulfate), 28
LY188011 (see Gemcitabine), 21
LY188011 hydrochloride (see Gemcitabine HCl), 21
Lymphochin (see Aniline mustard), 15
Lymphocin (see Aniline mustard), 15
Lymphokine activated killer cells (see LAK cells), 64
Lymphoquin (see Aniline mustard), 15
Lynestrenol#, 77
Lynoral# (see Ethinyl estradiol), 76
Lysodren (see Mitotane*), 23

M

M-9500 (see Triethylenemelamine), 27 M-AMSA (see Amsacrine), 15 M-Azido-Pyrimethamine Ethane Sulphonate, 21 MA (see Methandrostenolone), 78 Mablin (see Busulfan*), 16 Mafosfamide L-lysine (see Mafosfamide), 22 Mafosfamide, 21 Magnamycin A (see Magnamycin), 22 Magnamycin, 22 Magnesium Sulfate, 43 Malathion, 98 Malestrone (see Methyltestosterone), 78 Malocide (see Pyrimethamine), 45, 99 Mannit-Lost (see Mannomustine), 43 Mannit-Mustard (see Mannomustine), 43 Mannitol Mustard (see Mannomustine), 43 Mannitol, 43 Mannomustine, 43 Mannosulfan, 22 MAP (see Medroxyprogesterone Acetate), 77 Marax (see Hydroxyzine HC1), 42 Marcellomycin, 22 Marinol (see Dronabinol, Tetrahydrocannabinol), 46, 96 Masoprocol, 22 Masterid (see Dromostanolone Propionate), 75 Masteril (see Dromostanolone Propionate), 75 Masterone (see Dromostanolone Propionate), 75 Matrol (see Chlormadinone* Acetate), 73 Matulane (see Procarbazine* (HCl)), 25

Maxidex (see Dexamethasone), 40, 74

Maxitrol (see Dexamethasone), 40, 74

Maytansine, 22

MBBA (see Cytembena), 18

MCCNU (see Semustine), 25

McN-JR-1625 (see Haloperidol), 42

McN-JR-4749 (see Droperidol*), 41

McN-R-1967 (see Fenretinide), 20

MCNU (see Semustine), 25

MCP (see Metoclopramide (HCl)*), 43

MDL 28574, 93

MDL 71,782A (see Effornithine HCl), 19

MDL-72222, 43

MDL-73147EF, 43

Mebryl (Czechoslovakia) (see Cytembena), 18

MeCCNU (see Semustine), 25

Mechlorethamine (HCl)*, 22

Mechlorethamine (see Mechlorethamine (HCl)*), 22

Medrate (see Methylprednisolone), 78

Medrocort# (see Medrysone*), 77

Medrogestone, 77

Medrol (see Methylprednisolone), 78

Medrol Stabisol# (see Methylprednisolone Sodium Phosphate), 78

Medrone (see Methylprednisolone), 78

Medrotestron (see Oxymetholone*), 79

Medroxyprogesterone Acetate, 77

Medrysone*, 77

MeG (see Methyl-G), 22

Meg-CSF, 64

Megace (see Megestrol Acetate), 78, 98

MeGAG (see Methyl-G), 22

Megestrol Acetate, 78, 98

Melbex# (see Mycophenolic acid), 51

Melengestrol Acetate, 78

Melfalan (see Melphalan* (HCl)), 22

Melphalan* (HCl), 22

MEN (see Menogaril), 22

Menest (see Estrogens, Esterified), 76

Menogaril, 22

Menolyn (see Ethinyl estradiol), 76

Menrium, component of (see Estrogens, Esterified), 76

Mepacrine (see Quinacrine (HCl)), 45

Mepadin (see Meperidine HC1), 43

Meperga (see Meperidine HC1), 43

Mepergan, component of (see Promethazine HCl*), 45

Meperidine HCI, 43

Meprednisone*, 78

MER (see BCG), 61

Meractinomycin (see Dactinomycin*), 18

Merbarone, 22

Merbentul (see Chlorotrianisene), 74

Mercaleukin (see Mercaptopurine*), 22

Mercaptopurine*, 22

6-Mercaptoguanine (see Thioguanine*), 26

6-Mercaptoinosine (see 6-Mercaptopurine riboside), 22

6-Mercaptopurin (see Mercaptopurine*), 22

6-Mercaptopurine (see Mercaptopurine*), 22

6-Mercaptopurine ribonucleoside (see 6-Mercaptopurine riboside), 22

6-Mercaptopurine riboside, 22

Mercapurin (see Mercaptopurine*), 22

Mern (see Mercaptopurine*), 22

Mertestate# (see Testosterone), 80

Mesna*, 43

Mesnex (uromitexan) (see Mesna*), 43

Mesnum (see Mesna*), 43

Meso-NDGA (see Masoprocol), 22

Mespergan (see Methadone HCl), 43

Mesterolone, 78

Mestranol, 78

Metace (see Chlorotrianisene), 74

Metanabol (see Methandrostenolone), 78

Metandienon (see Methandrostenolone), 78

Metandren (see Methyltestosterone), 78

Metandrostenolone (see Methandrostenolone), 78

Metasol (see Methandrostenolone), 78

Methadone HCl, 43

Methadose (analgesic; narcotic) (see Methadone HCl), 43

Methandronolone (see Methandrostenolone), 78

Methandrostenolone, 78

Methanol extracted residual of BCG (see BCG), 61

MeTHHF (see Emofolin (sodium)), 20

Methionyl Interferon Consensus, 65

Metholone (see Oxymetholone*), 79

Methosarb# (see Calusterone), 73

Methotrexate sodium (see Methotrexate*), 22

Methotrexate*, 22

Methoxsalen, 43

Methral (see Fluperolone Acetate), 76

Methyl CCNU (see Semustine), 25

Methyl ether (see TMCA), 27

Methyl methanesulfonate (see Methyl methanesulphonate), 22

Methyl methanesulphonate, 22

Methyl streptonigrin (see Streptonigrin), 25

Methyl-G, 22

Methyl-GAG (see Methyl-G), 22

6-Methyl MP-riboside (see 6-Methylmercaptopurine riboside), 23

Methylcyclohexyl nitrosourea (see Semustine), 25

5-Methyldeoxyuridine (see Thymidine), 26

Methylene Dimethane Sulfonate, 22

Methylformamide (see N-Methylformamide), 51

Methylglyoxal bisguanyl-hydrazone dihydrochloride monohydrate, 22

Methylhydrazine (see Procarbazine* (HCl)), 25

6-Methylmercaptopurine riboside, 23

Methylmethane sulfonate (see Methyl methanesulphonate), 22

Methylphenidate, 43

Methylprednisolone, 78

Methylprednisolone Acetate, 78

Methylprednisolone Sodium Phosphate, 78

Methylprednisolone Sodium Succinate, 78

Methyltestosterone, 78

5-Methyltetrahydrohomofolate disodium (see Emofolin (sodium)), 20

Methylthioinosine (see 6-Methylmercaptopurine riboside), 23

6-Methylthiopurine riboside (see 6-Methylmercaptopurine riboside), 23

Methynodiol Diacetate#, 78

Meti-Derm (see Prednisolone), 79

Meticortelone acetate (see Prednisolone Acetate), 79

Meticortelone Soluble (see Prednisolone Hemisuccinate), 79

Meticorten (see Prednisone), 79

Metoclopramide (HCl)*, 43

Metoprine, 23

Metreton (see Prednisolone Sodium Phosphate), 79

Metro I.V. (see Metronidazole), 98

Metrone (see Methyltestosterone), 78

Metronidazole, 98

Meturedepa, 23

Metyrosine, 44

Mevacor (see Lovastatin), 21

Mevinolin# (see Lovastatin), 21

Mexate (see Methotrexate*), 22

Mexate AQ (see Methotrexate*), 22

Mexiletine HCl. 98

Mexitil (see Mexiletine HCl), 98

MGA (see Melengestrol Acetate), 78 MGA (see Methyl-G), 22 MGBG (see Methyl-G), 22 MGGH (see Methyl-G), 22 Miboleron (see Mibolerone), 78 Mibolerone, 78 Micatin (see Miconazole), 98 Miconazole, 98 Microfollin (see Ethinyl estradiol), 76 Midazolam HCl, 44 Mielobromol (see Dibromomannitol), 19 Mifepristone, 78 MIH (see Procarbazine* (HCl)), 25 Milprem (see Estrogens, Conjugated), 76 Miltefostine, 23 Milurit (see Allopurinol*), 39 Mintezol (see Thiabendazole), 100 Misonidazole, 44 Mistabron (see Mesna*), 43 Misulban (see Busulfan*), 16 MITC (see Mitomycin*), 23 Mitexan (see Mesna*), 43 MITH (see Plicamycin*), 24 Mithracin (see Plicamycin*), 24 Mithramycin# (see Plicamycin*), 24 Mitindomide, 23 Mito (see Mitomycin*), 23 MITO-C (see Mitomycin*), 23 Mitobronitol (see Dibromomannitol), 19 Mitocarcin, 23 Mitocin-C (see Mitomycin*), 23 Mitoclomine, 23 Mitocromin, 23 Mitogillin, 23 Mitoguazone (see Methyl-G), 22 Mitoguazone dihydrochloride (see Methyl-G), 22 Mitolac (see Dibromodulcitol), 19 Mitolactol (see Dibromodulcitol), 19 Mitomalcin, 23 Mitomycin-C (see Mitomycin*), 23 Mitomycin*, 23 Mitopodozide, 23 Mitosper, 23 Mitostan (see Busulfan*), 16

Mitotan (see Mitotane*), 23

Mitotane*, 23

Mitoxana (see Ifosfamide*), 21

Mitoxantrone (HCl)*, 23

Mitozolomide, 23

MJF 9325 (see Ifosfamide*), 21

MJF-12264, (see Tegafur), 26

MK-217 (see Alendronate Sodium), 39

MK-595 (diuretic) (see Ethacrynic Acid), 41

1-MM (see Puromycin (HCl)), 25

MMC (see Mitomycin*), 23

6-MMPR (see 6-Methylmercaptopurine riboside), 23

MN rgp120/HIV-1, 93

MNI (see Misonidazole), 44

Modicon (see Ethinyl estradiol), 76

Modrenal (see Trilostane), 81

Molybdenum, 44

Monistat (see Miconazole), 98

Monoclonal Antibody, 65

Monomethylformamide (see N-Methylformamide), 51

8-MOP (see Methoxsalen), 43

Mopidamol, 23

Moranyl (see Suramin Sodium), 26

Morphine Sulfate, 44

MP (see Mercaptopurine*), 22

6-MP (see Mercaptopurine*), 22

MPA (see Medroxyprogesterone Acetate), 77

MPL (see Melphalan* (HCl)), 22

6-MPR (see 6-Mercaptopurine riboside), 22

MS Contine (see Morphine Sulfate), 44

MSI (see Morphine Sulfate), 44

MSL-109, 98

MTC (see Mitomycin*), 23

MTH (see Plicamycin*), 24

MTP-PE (see CGP 19835A), 62

MTS (see Maytansine), 22

MTX (see Methotrexate*), 22

Mucofluid (see Mesna*), 43

Mucopirocin, 98

Muran (see Azathioprine*), 16

Mustard (see Chlornaphazine*), 40

Mustargen (see Mechlorethamine (HCl)*), 22

Mustargen-HCl (see Mechlorethamine (HCl)*), 22

Mustine (see Mechlorethamine (HCl)*), 22

Mutamycin (see Mitomycin*), 23 MVE-2, 65 Myagen# (see Bolasterone#), 73 Mycelex (see Clotrimazole), 96 Mycelex G (see Clotrimazole), 96 Myco-Triacet II (see Triamcinolone Acetonide), 81 Myco-Triacet II, component of (see Nystatin), 98 Mycobutin (see Rifabutin), 99 Mycolog II (see Triamcinolone Acetonide), 81 Mycolog II, component of (see Nystatin), 98 Mycophenolic acid, 51 Mycostatin (see Nystatin), 98 Myebrol (see Dibromomannitol), 19 Myelobromol (see Dibromomannitol), 19 Myelosan (see Busulfan*), 16 Mylecytan (see Busulfan*), 16 Myleran (see Busulfan*), 16 Mylosar (see Azacitidine), 51 Myristic acid analogs, 93 Mysteclin-F#, component of (see Amphotericin B), 95 Mytrex (see Triamcinolone Acetonide), 81 Mytrex, component of (see Nystatin), 98

Ν

N-137 (see Carbetimer), 17 N-acetyl cysteine (see NAC), 93 N-Acetyl DON (see Duazomycin), 19 N-butyl-DNJ (see Deoxynojirimycin), 92 N-CWS (see Nocardia rubra), 65 N-CWS, 65 N-Methylformamide, 51 N-Methylmitomycin C (see Porfiromycin), 25 N-Phenylbarbitol (see Phetharbital), 24 N-Phosphonacetyl-L-Aspartate disodium (see PALA), 24 N2 Transduced TIL, 65 N2-TIL/ADA Gene, 65 Nabilone, 44 NAC, 93 Nafarelin Acetate, 78 Nafazatrom, 23 Nafidimide, 23 NAFO (see Nafoxidine HCL), 78 Nafoxidine HCL, 78

Nafticlorina (see Chlornaphazine*), 40

Naganin (see Suramin Sodium), 26

Naganol HCl (see Suramin Sodium), 26

Naloxone, 44

Naltrexone, 44

Nalutron# (see Progesterone), 80

Nandrobolic (see Nandrolone Phenpropionate), 79

Nandrobolic L.A.# (see Nandrolone Decanoate), 78

Nandrolone Decanoate, 78

Nandrolone Phenpropionate, 79

Nandrolone phenylpropionate (see Nandrolone Phenpropionate), 79

Naphthylamine (see Chlornaphazine*), 40

Naphuride Sodium (see Suramin Sodium), 26

Naposim (see Methandrostenolone), 78

Naprosyn (see Naproxen), 44

Naproxen, 44

NASH; BSH (radioactive agent) (see Borocaptate Sodium B 10), 39

Natulan (see Procarbazine* (HCl)), 25

Navelbine (see Vinorelbine), 28

Naxamide# (see Ifosfamide*), 21

NCNU (see Pentamustine), 24

NCO (see Diethylstilbestrol), 75

NCS (see Zinostatin), 28

NDC 0015-3010 (see Bleomycin Sulfate), 16

NDC-0082-4155 (see Daunorubicin (HCl)*), 18

2'NDG (see Ganciclovir), 97

NDHT (see Dromostanolone Propionate), 75

NebuPent (see Pentamidine), 98

Nedrotestron propionate (see Dromostanolone Propionate), 75

Neo-Hombreol-M# (see Methyltestosterone), 78

Neo-Medrol (see Methylprednisolone Acetate), 78

Neo-Ormonal (see Fluoxymesterone), 76

Neo-Oxylone# (see Fluorometholone), 76

Neocarcinostatin (see Zinostatin), 28

Neocarzinostatin# (see Zinostatin), 28

NeoDecadron (see Dexamethasone Sodium Phosphate), 75

Neodrol (see Stanolone), 80

Neoplatin (see Cisplatin*), 17

Neosar (see Cyclophosphamide*), 18

Neosone, component of (see Cortisone Acetate), 74

Neovitamin A acid (see Retinoic acid), 51

Nerobil (see Nandrolone Phenpropionate), 79

Nerobol (see Methandrostenolone), 78

Neupogen (see Filgrastim), 62, 97

Neupogen (see G-CSF), 62, 97

NFX (see Nafoxidine HCL), 78

NH2 (see Mechlorethamine (HCl)*), 22

NH2 Nitrogen Mustard (see Mechlorethamine (HCl)*), 22

NH2-HCl (see Mechlorethamine (HCl)*), 22

Niac (see Niacin), 44

Niacin, 44

Nicobid (see Niacin), 44

Nicolar (see Niacin), 44

Niconyl# (see Isoniazid*), 97

Nicroest (see Diethylstilbestrol), 75

Nidaton# (see Isoniazid*), 97

Nifedipine, 44

Niftholid (see Flutamide), 77

Niftolid (see Flutamide), 77

Nigrin (see Streptonigrin), 25

Nilestrol (see Diethylstilbestrol), 75

Nilstat (see Nystatin), 98

Nilutamide, 79

Nimodipine, 98

Nimotop (see Nimodipine), 98

Nipent (see Pentostatin), 65

4-Nitroestrone, 79

Nitrofurazone, 23

Nitrogen mustard (see Melphalan* (HCl)), 22

Nitromifene Citrate, 79

Nivazol, 79

Nizoral (see Ketoconazole), 77

NK 631 (see Peplomycin Sulfate), 24

NMF (see N-Methylformamide), 51

Nocardia rubra, 65

Nocodazole (Belgium), 23

Nogalamycin#, 23

Nolvadex (see Tamoxifen Citrate), 80

Nordette (see Ethinyl estradiol), 76

Norethandrolone, 79

Norethin 1/50 M (see Mestranol), 78

Norethrin 1/35E (see Ethinyl estradiol), 76

Norinyl (see Mestranol), 78

Norlestrin (see Ethinyl estradiol), 76

Norquen (see Mestranol), 78

Nortestosterone phenylpropionate, (see Nandrolone Phenpropionate), 79

Nortriptyline HCl, 44

Novantrone (see Mitoxantrone (HCl)*), 23

Novatrone, 23 Novestrol (see Ethinyl estradiol), 76 Novidium chloride (see Ethidium chloride), 20 NPP (see Nandrolone Phenpropionate), 79 NTPP (see Nandrolone Phenpropionate), 79 Nydrazid (see Isoniazid*), 97 Nylestriol, 79 Nystatin, 98 Nystex (see Nystatin), 98 NZS (see Zinostatin), 28 0 O.p'-DDD (see Mitotane*), 23 O.p'-DDE (see Mitotane*), 23 Octamide PFS (see Metoclopramide (HCl)*), 43 Octreotide Acetate, 65 Oestrogenine (see Diethylstilbestrol), 75 Ofloxacin, 98 Ogen (see Estropipate), 76 OK-432, 65 Oligomycin B (antibiotic) (see Peliomycin), 24 Omain (see Colcemid), 17 Omaine (see Colcemid), 17 7-OMEN (see Menogaril), 22 Onco-Carbide (see Hydroxyurea*), 21 Oncostatin K (see Dactinomycin*), 18 Oncotepa (see Thiotepa*), 26 Oncotiotepa (see Thiotepa*), 26 Oncovin (see Vincristine* Sulfate), 28 Ondanserin (see Ondansetron HCl), 44 Ondansetron HCl, 44 Ophthalmadine (see Idoxuridine*), 42 Optimyd (see Prednisolone Sodium Phosphate), 79 Ora-Testryl (see Oxymetholone*), 79 Ora-Testryl# (see Fluoxymesterone), 76 Oracil (see Fluorouracil*), 20 Oradiol (see Ethinyl estradiol), 76 Oragest (see Medroxyprogesterone Acetate), 77 Oralsterone (see Fluoxymesterone), 76 Orange Crush# (anthracene derivative) (see Bisantrene (HCl)), 16 Orasone (see Prednisone), 79 Oratestin (see Fluoxymesterone), 76

Orestralyn (see Ethinyl estradiol), 76

Oreton (see Methyltestosterone), 78 Oreton# (see Testosterone), 80 Org 485-50 (see Lynestrenol#), 77 Orgametril# (see Lynestrenol#), 77 Orgametrol (see Lynestrenol#), 77 Ormaplatin (see Tetraplatin), 26 Ormaplatin, 23 Ornidyl (see Eflornithine HCl), 19, 96 Ortho-Cyclen (see Estropipate), 76 Ortho-Novum (see Mestranol), 78 OTC, 93 Ovid (see Malathion), 98 Ovocyclin dipropionate# (see Estradiol Dipropionate), 75 Ovoresta M (progestin) (see Lynestrenol#), 77 Ovulen (see Mestranol), 78 Oxandrolone*, 79 Oxanthrazole (see Piroxantrone HCl), 24 Oxantrazole HCl (see Piroxantrone HCl), 24 Oxisuran*#, 23 Oxsoralen (see Methoxsalen), 43 Oxylone# (see Fluorometholone), 76 Oxymetholone*, 79 Oxytocin, 79 Oxyurea (see Hydroxyurea*), 21 P P-165 (antimetabolite) (see Azaserine*#), 16 P-1742 (glucocorticoid) (see Fluperolone Acetate), 76 P-2647 (anti-emetic) (see Benzquinamide), 39 P-638 (see Puromycin (HCl)), 25 PA-144 (see Plicamycin*), 24 Paclitaxel, 24 PALA, 24 Palonyl Perovex (see Ethinyl estradiol), 76 PAM (see Melphalan* (HCl)), 22 Pamelor (see Nortriptyline HCl), 44 Pamidronate Disodium, 44 Pancratistatin, 24 Panwarfin (anticoagulant) (see Warfarin Sodium*), 46 Papaverine HCl, 44 Para-aminosalicylic acid, 98 Paracort# (see Prednisone), 79 Paracortol# (see Prednisolone), 79

Paramethasone* Acetate, 79 Paraplatin (see Carboplatin*), 17 Paromomycin Sulfate, 98 PAS (see Para-aminosalicylic acid), 98 Pathocidin (see 8-Azaguanine), 15 Pathocidine (see 8-Azaguanine), 15 Pavabid (see Papaverine Disodium), 44 PCB (see Procarbazine* (HCl)), 25 PCH (see Procarbazine* (HCl)), 25 PCI (see Procarbazine* (HCl)), 25 PCNU, 24 PCZ (see Procarbazine* (HCl)), 25 PD 90,695-73 (see Dezaguanine Mesylate), 19 PDA (see Phosphoramide mustard#), 24 PDN (see Prednisone), 79 Pediapred (see Prednisolone Sodium Phosphate), 79 PEG (see Polyethylene Glycol), 44 PEG-IL-2, 65, 93 Pegaspargase, 65 Peliomycin, 24 Penberol, 24 Penclomidine, 24 Pentam 300 (see Pentamidine Isethionate), 93 Pentam 300 (see Pentamidine), 98 Pentamethylmelamine HCl. 24 Pentamidine Isethionate, 93 Pentamidine, 98 Pentamustine, 24 Pentosan Polysulfate Sodium, 24 Pentosan sulfate (see Pentosan Polysulfate Sodium), 24 Pentostatin, 65 Pentoxifylline, 44 Pepleomycin (see Peplomycin Sulfate), 24 Peplomycin Sulfate, 24 Peptide T, 93, 99 Peptochemio, 24 Percorten acetate# (see Desoxycorticosterone Acetate), 74 Percorten pivalate# (see Desoxycorticosterone Pivalate), 74 Percotol (see Desoxycorticosterone Acetate), 74 Pergantene (see Sodium Fluoride), 45 Peridex, 99

Perlutex (see Medroxyprogesterone Acetate), 77 Permastril (see Dromostanolone Propionate), 75

Permethrin, 99

Persantine (see Dipyridamole), 41 Persistol (see Triethylenemelamine), 27 Peyrone's chloride (see Cisplatin*), 17 PFA (see Foscarnet Sodium), 97 Pharmorubicin (see Epirubicin (HCl)), 20 Phenergan (see Promethazine HCl*), 45 Phenergan-D, component of (see Promethazine HCl), 45 Phenergan VC, component of (see Promethazine HCl), 45 Phenesterin (see Phenesterine), 24 Phenesterine, 24 Phenestrin (see Phenesterine), 24 Phenidiemal (see Phetharbital), 24 Phenobolin (see Nandrolone Phenpropionate), 79 Phenyl acetate, 51 Phetharbital, 24 Phosphamide (see Phosphoramide mustard#), 24 Phosphocol P 32 (see Chromic Phosphate P 32), 74 Phosphoramide mustard#, 24 Photofrin (see Porfimer Sodium), 24 Photofrin II (see Porfimer Sodium), 24 Phyllanthoside, 24 Pibenzimol HCl, 24 PIBR (see Pipobroman*), 24 Pimonidazole (see Ro-03-8799), 45 Piperazidine (see Piperazine), 44 Piperazin (see Piperazine), 44 Piperazine estrone sulfate# (see Estropipate), 76 Piperazine, 44 Piperazinedione compound, 24 Pipersol (see Piperazine), 44 Pipobroman*, 24 Piposulfan, 24 Pirazofurin (see Pyrazofurin#), 25 Pirimecidan (see Pyrimethamine), 45 Pirimetamina (see Pyrimethamine), 45 Piritrexim Isethionate, 99 Piroxantrone HCl, 24 Piroxicam, 44 PISU (see Piposulfan), 24 Pitocin (see Oxytocin), 79 Plaquenil Sulfate (see Hydroxychloroquine Sulfate), 92 Platelet Factor 4, 99

Platinex (see Cisplatin*), 17 Platinol (see Cisplatin*), 17

Platinum (miscellaneous agent) (see Cisplatin*), 17

Platinum complex (9CI) (see Iproplatin), 21

Plicamycin*, 24

Pluracol E400, E600, E1450 (see Polyethylene Glycol), 44

PMB-200 (see Estrogens, Conjugated), 76

PMB-400, component of (see Estrogens, Conjugated), 76

PMEA, 93

PMM (see Pentamethylmelamine), 24

Pneumovax, 93

Podofilox, 99

Podophyllic acid (see Mitopodozide), 23

Podophyllotoxin derivative (see Etoposide*), 20

Poly AU, 65

Poly I: Poly C12U, 65, 91

Poly I:C (see 5'-Inosinic acid polymers), 65

Poly I:Poly C with Poly-I-lysine stabilizer, 65

Poly IC/LC (see Poly I:Poly C with Poly-I-lysine stabilizer), 65

Polyestradiol Phosphate, 79

Polyethylene Glycol, 44

PolyI: Poly C12U (see Ampligen), 65, 91

PORF (see Porfiromycin), 25

Porfimer Sodium, 24

Porfiromycin, 25

Porfiromycine (see Porfiromycin), 25

Porton asparaginase (see Asparaginase), 61

PRD (see Prednisone), 79

PRDL (see Prednisolone), 79

PRED (see Prednisone), 79

Predalone T.B.A. (see Prednisolone Tebutate), 79

Predate (see Prednisolone Acetate), 79

Predate-S (see Prednisolone Sodium Phosphate), 79

Predne-Dome# (see Prednisolone), 79

Prednimustine, 25

Prednisolone, 79

Prednisolone Acetate, 79

Prednisolone Sodium Hemisuccinate, 79

Prednisolone Sodium Phosphate, 79

Prednisolone sodium succinate (see Prednisolone Hemisuccinate) 79

Prednisolone Tebutate, 79

Prednisone, 79

Prednival*#, 79

Pregnenolone* Succinate, 80

Premarin (see Estrogens, Conjugated), 76

Premarin with Methyltestosterone, component of (see Methyltestosterone), 78

Premarin with Methyltestosterone, component of (see Estrogens, Conjugated), 76

Primaquine Phosphate, 99

Primaxin, component of (see Cilastatin Sodium), 17

Primaxin, component of (see Imipenem), 21

Primaxin, component of (see Imipenem-Cilastatin), 97

Primocort (see Desoxycorticosterone Acetate), 74

Primocortan (see Desoxycorticosterone Acetate), 74

Probenecid, 44

Procarbazine* (HCl), 25

Prochlorperazine Maleate, 45

Prochlorperazine*, 45

Procrit (see Erythropoietin), 62, 96

Procytox (see Cyclophosphamide*), 18

Prodasone (see Medroxyprogesterone Acetate), 77

Prodel (see Dydrogesterone*), 75

Prodrox (progestin) (see Hydroxyprogesterone Caproate), 77

Progestal (see Medroxyprogesterone Acetate), 77

Progesterone*, 80

Progynon B (estrogen) (see Estradiol Benzoate), 75

Progynon# (estrogen) (see Estradiol), 75

Prokine (see Sargramostim), 66

Proleukin (see Interleukin-2), 63

Proloid (see Thyroglobulin), 80

Proloprim (see Trimethoprim), 46, 100

Promacortine (see Methylprednisolone), 78

Promazine HCl*, 45

Promethazine HCl*, 45

2-Propanimine (see Iproplatin), 21

2-Propenamide (see Anthramycin), 15

2-Propenamide (see Bromacrylide#), 16

Proresipar (see Mitopodozide), 23

Protease inhibitor, 93

Protein A Staphylococcus Aureus, 65

Protivar (see Oxandrolone*), 79

Protobolin (see Methandrostenolone), 78

Protostat (see Metronidazole), 98

Protropin (see Human Growth Hormone), 92

Protropin (see Somatrem), 80

Proventil (see Albuterol), 39

Prover (see Medroxyprogesterone Acetate), 77

Provera (see Medroxyprogesterone Acetate), 77

Proverone (see Medroxyprogesterone Acetate), 77

PRZF (see Pyrazofurin#), 25

Pseudomonas aeruginosa, 65

PSK, 65 Psorion (see Betamethasone Dipropionate), 73 PSP-IV (see Prednisolone Sodium Phosphate), 79 PTC (see Peptochemio), 24 PTG (see Teniposide), 26 Purinethiol (see Mercaptopurine*), 22 Purinethol (see Mercaptopurine*), 22 Puromycin (HCl), 25 Puromycin (see Puromycin (HCl)), 25 Pyran (see Pyran copolymer), 65 Pyran copolymer, 65 Pyran XA 124-177 (see Pyran copolymer), 65 Pyranone Phosphate (see Fostriecin Sodium), 20 Pyrantel Pamoate, 99 Pyrazinamide, 45, 99 Pyrazine diazohydroxide, 25 Pyrazine Hexahydride (see Piperazine), 44 Pyrazofurin#, 25 Pyrazolo (2,3-a)imadazolidine (see Pyrazolo-Imidazole compound), 25 Pyrazolo-Imidazole compound, 25 Pyrazoloacridine, 25 Pyrazomycin (see Pyrazofurin#), 25 Pyrictal (see Phetharbital), 24 Pyridoxine, 45 Pyrimethamine, 45 PZA (see Pyrazinamide), 45 PZD (see Piperazinedione compound), 24 PZF (see Pyrazofurin#), 25

Q

Quantril (see Benzquinamide), 39
Quiess (see Hydroxyzine HC1), 42
QUIN (see Quinacrine (HCl)), 45
Quinacrine (HCl), 45
Quinacrine, 99
Quinactine (see Quinacrine (HCl)), 45
Quinestrol, 80
Quingestanol* Acetate, 80
Quingestrone#, 80
Quinidine, 45
Quinomycin A (see Echinomycin), 19

R 17,934 (see Nocodazole (Belgium)), 23 R 9985 (see Methotrexate*), 22 R-12,564 (see Levamisole HCl*), 64 R-1625 (antipsychotic) (see Haloperidol), 42

R-246 (see Triethylenemelamine), 27

R-33-812 (anti-emetic) (see Domperidone), 41

R-41,400 (see Ketoconazole), 77

R-4749 (see Droperidol*), 41

R-48 (see Chlornaphazine*), 40

R-52,211 (see Itraconazole), 42, 97

R-52 (see Mannosulfan), 22

R-met Hu G-CSF (see G-CSF), 41

R-met Hu G-CSF (see Filgrastim), 62

R-Verapamil HCl, 45

RA (see Retinyl Acetate), 51

RA-233 (see Mopidamol), 23

Ra-8 (coronary vasodilator) (see Dipyridamole), 41

Radinyl (see Etanidazole), 41

Ranitidine HCl, 99

Razoxane, 25

RBD (see Zorubicin (HCl)), 28

RC-160 (see Vapreotide), 27

Recombinant Alpha 2 interferon (see Interferon Alfa-2b), 63

Recombinant Human GM-CSF (see CSF-GM), 40

Recombinant human growth hormone, 94

Recombinant interferon Gamma (see Interferon Alfa-2b), 63

Recombinant Leukocyte A interferon (see Interferon Alfa-2b), 63

Recombinant Soluble CD4, 66

Recombinant Soluble T4, 66

Recominbant Tumor Necrosis Factor, 66

Regamycin# (see Porfiromycin), 25

Reglan (anti-emetic) (see Metoclopramide (HCl)*), 43

Remsed (anti-emetic) (see Promethazine HCl*), 45

Repromap (see Medroxyprogesterone Acetate), 77

Repromix (see Medroxyprogesterone Acetate), 77

Retin-A (see Tretinoin), 51

Retinoic acid (see Tretinoin), 51

13-cis-Retinoic acid (see Retinoic acid), 51

Retinyl Acetate, 51

Retrone (see Dydrogesterone*), 75

Retrovir (see Azidothymidine, Zidovudine*), 16, 28, 91

rgp120 CHO, 94

rgp120/HIV-1 (IIIB), 94

Rhenium-186, 45

Rheumatrex (antimetabolite) (see Methotrexate*), 22

rHGH, 93

Rhizoxin, 25

rHu-EPO (see Erythropoietin), 62, 96

Rianil (see Chlorotrianisene), 74

Ribavirin, 94

Ribo-azauracil (see 6-Azuridine), 16

Ribo-azuracil (see 6-Azuridine), 16

Riboazuracil (see 6-Azuridine), 16

Riboflavin, 45

Riboprine*, 25

Riboxamide (see Tiazofurin), 26

Rifabutin, 99

Rifadin (see Rifampin), 25, 99

Rifampin, 25, 99

Rimactane (see Rifampin), 25, 99

Rimifon# (see Isoniazid*), 97

Ritalin (see Methylphenidate HCl), 43

Ro 03-8799 (see Pimonidazole), 44

Ro 2-9757 (see Fluorouracil*), 20

Ro 21-0702 (see Flurocitabine#), 20

Ro 21-3981/003 (see Midazolam HCl), 44

Ro 21-8837 (see Estramustine), 75

Ro 21-8837/001 (chemotoxic hormone) (see Estramustine Phosphate), 76

Ro 22-2296/000 (see Estramustine), 75

Ro 22-8181 (see Interferon Alfa-2a), 63

Ro 24-2027 (see Dideoxycytidine), 92

Ro 4-3780 (see Retinoic acid), 51

Ro 5-9963 (see Desmethylmisonidazole), 40

Ro-03-8799, 45

Ro-7-0582 (see Misonidazole), 44

Ro4-6467/1 (see Procarbazine* (HCl)), 25

Rocephin (see Ceftriaxone), 95

Roferon (see Interferon Alfa-2a), 63

Roferon-A (see Interferon Alfa-2a), 63

RP 2512 (see Pentamidine Isethionate), 93

RP-13057 (see Daunorubicin (HCl)*), 18

RP-22,050 (see Zorubicin (HCl)), 28

RP-6870 (see Inproquone), 21

RS-1301 (see Delmadinone Acetate), 74

RS-21592 (see Ganciclovir), 97

RS-2208 (progestin) (see Amadinone Acetate), 73

RS-3540 (see Naproxen), 44

RS-94991-298 (see Nafarelin Acetate), 78
RU 41740 (see Biostim), 62
RU 486 (see Mifepristone), 78
RUB (see Zorubicin (HCl)), 28
Rubex (see Doxorubicin (HCl)*), 19
Rubidazone (see Zorubicin (HCl)), 28
Rubidazone hydrochloride (see Zorubicin (HCl)), 28
Rubidomycin-HCl (see Daunorubicin (HCl)*), 18
Rubomycin C (see Daunorubicin (HCl)*), 18
Rufochromomycin (see Streptonigrin), 25
Rufocromomycin (see Streptonigrin), 25

S

S-trityl-L-cysteine (see Trityl cysteine), 27 S6820 (Shionogi) (see IL-2), 63 Salisburystin (see Pentamustine), 24 Sanamicia (see Cactinomycin), 16 Sanamycin (see Cactinomycin), 16 Sandamycin (see Cactinomycin), 16 Sandostatin (see Octreotide Acetate), 65 Sandostatin, 45 Santavy's Substance F (see Colcemid), 17 Sarcolysine (see Melphalan* (HCl)), 22 Sargramostim (see GM-CSF), 66, 97 Sarkoklorin (see Melphalan* (HCl)), 22 Sarkolizin (see Melphalan* (HCl)), 22 Satric (see Metronidazole), 98 SC 11585 (see Oxandrolone*), 79 SC-10363 (see Megestrol Acetate), 78 SC-16148 (see Silandrone*#), 80 SC-19198 (see Methynodiol Diacetate#), 78 SC-9420 (see Spironolactone), 80 SC-9880 (see Flurogestone Acetate), 76 Sch 11460 (see Betamethasone Dipropionate), 73 Sch 13521 (see Flutamide), 77 Sch-18020W (see Beclomethasone Dipropionate), 73 Sch-4358 (see Meprednisone*), 78 Sch-4831 (see Betamethasone), 73 Semustine, 25 Sendoxan (see Cyclophosphamide*), 18 Septra, component of (see Sulfamethoxazole), 46 Septra, component of (see Trimethoprim), 46 Seromycin (see Cycloserine), 96

Somatuline, 80

Serophene (see Clomiphene* Citrate), 74 Serotonin, 45 Serral (see Diethylstilbestrol), 75 Sexocretin (see Diethylstilbestrol), 75 SF-337 (see 8-Azaguanine), 15 SH 714 (see Cyproterone Acetate), 74 SH 741 (progestin) (see Clomegestone* Acetate), 74 SH 818 (glucocorticoid) (see Clocortolone Acetate), 74 SH 863 (see Clocortolone Pivalate), 74 SH-1040 (see Gestaclone), 77 SH-582 (see Gestonorone Caproate), 77 SH-723 (see Mesterolone), 78 SH-968 (glucocorticoid) (see Diflucortolone* Pivalate), 75 Sibol (see Diethylstilbestrol), 75 Silandrone*#, 80 Simtrazene#, 25 Sinestrol (see Hexestrol), 77 Sirprogen (see Medroxyprogesterone Acetate), 77 SK-1133 (see Triethylenemelamine), 27 SK&F 14287 (see Idoxuridine*), 42 SK&F-7690 (see Benorterone#), 73 SKF 104864-A (see Topotecan HCl), 27 SKF 33134-A (cardiac depressant) (see Amiodarone), 39 SN (see SR 2508), 45 SN (see Streptonigrin), 25 Sodium 2-mercaptoethane (see Mesna*), 43 Sodium 2-mercaptoethanesulfonate (see Mesna*), 43 Sodium bromebrate (see Cytembena), 18 Sodium Butyrate, 51 Sodium Chloride, 45 Sodium Fluoride, 45 Sodium Iodide, 45 Sodium N-Nitroso-Pyrazinamine (see Pyrazine diazohydroxide), 25 Sodium Phosphate P 32, 45 Sodium Suramin (see Suramin Sodium), 26 Sodium Thiosulfate, 45 Solatene (ultraviolet screen) (see Beta Carotene), 39, 62 Solu-Cortef (see Hydrocortisone Sodium Succinate), 77 Solu-Decortin (see Prednisolone Hemisuccinate), 79 Solu-Medrol (see Methylprednisolone Sodium Succinate), 78 Solu-Predalone (see Prednisolone Sodium Phosphate), 79 Somatostatin, 80 Somatrem, 80

Soriatane (see Acitretin), 51 SP-104 (anti-emetic) (see Tetrahydrocannabinol), 46 SP-I 77 (see Mitopodozide), 23 SP-PG, 94, 99 Spanestrin (see Ethinyl estradiol), 76 Sparfloxacin, 45, 99 Sparfosate Sodium (see PALA), 24 Sparine (antipsychotic) (see Promazine HCl*), 45 Sparsomycin A (see Tubercidine), 27 Sparsomycin, 25 Spectazole (see Econazole), 96 SPG (see Spirogermanium (HCl)), 25 Spiramycin, 45, 99 Spiresis (see Spironolactone), 80 Spiridon (see Spironolactone), 80 Spiro-32 (see Spirogermanium (HCl)), 25 Spirogermanium (HCl), 25 Spirohydantoin mustard# (see Spiromustine), 25 Spirolactone (see Spironolactone), 80 Spiromustine, 25 Spirone (see Spironolactone), 80 Spironolactone, 80 Spiroplatin, 25 SPL (see Spironolactone), 80 Spongo-adenosine (see Vidarabine*), 27 Spongocytidine HCl (see Cytarabine HCl), 18 Sporonax (antifungal) (see Itraconazole), 42, 97 SQ 1089 (see Hydroxyurea*), 21 SQ 15,101 (see Algestone* Acetophenide), 73 SQ 21,977 (see 6-Methylmercaptopurine riboside), 23 SQ 22558 (see Riboprine*), 25 SQ-9538 (see Testolactone), 80 SR 2508 (see Etanidazole), 41 SR-1354 (see Misonidazole), 44 SR-1530 (see Desmethylmisonidazole), 40 SRI 5244 (see Chlorozotocin), 17 SRI-1354 (see Misonidazole), 44 SRI-727 (see 2-Fluoroadenosine#), 20 ST 12 (see Dexamethasone Dipropionate), 75 St Johns Wort (see Hypericin), 92 Stanolone, 80 Stanozolol*, 80 Stavudine (see D4T), 92 Stemex (see Paramethasone* Acetate), 79

Stenolon (see Methandrostenolone), 78

STEPA (see Thiotepa*), 26

Sterane (see Prednisolone), 79 Sterane IM and IA (see Prednisolone Acetate), 79 Sterecyt (see Prednimustine), 25 Steronyl (see Methyltestosterone), 78 STGDR (see Diglycoaldehyde), 19 Stil-Rol (see Diethylstilbestrol), 75 Stilbestrol (see Diethylstilbestrol), 75 Stilbetin (see Diethylstilbestrol), 75 Stilboestroform (see Diethylstilbestrol), 75 Stiletin# (see Diethylstilbestrol), 75 Stilkap (see Diethylstilbestrol), 75 Stillomycin (see Puromycin (HCl)), 25 Stilphostrol (estrogen) (see Diethylstilbestrol Diphosphate), 75 STN (see Streptonigrin), 25 Stoxil (see Idoxuridine*), 42 STP (see Streptonigrin), 25 STR (see Streptozocin), 25 Strept (see Streptozocin), 25 Streptonigrin, 25 Streptozocin, 25 Streptozoticin (see Streptozocin), 25 Streptozotocin (see Streptozocin), 25 Strontium, 26 Stylomycin (see Puromycin (HCl)), 25 STZ (see Streptozocin), 25 Substance F (Reichstein's) (see Colcemid), 17 Sucralfate, 46 Sulestrex# (see Estropipate), 76 Sulfactol# (see Sodium Thiosulfate), 45 Sulfadiazine, 46, 100 Sulfadoxine/Pyrimethamine (see Fansidar), 41 Sulfamethoxazole (see Bactrim/Septra), 95 Sulfamethoxazole, 100 Sulfatrim, component of (see Sulfamethoxazole), 46 Sulfatrim, component of (see Trimethoprim), 46 Sulfonamide Duplex, component of# (see Sulfadiazine), 46 Sulfonate (see Mesna*), 43 Sulindac, 46 Sulofenur, 26 Superanabolon (see Nandrolone Phenpropionate), 79 Suprafact (see Busrelin Acetate), 73 Suramin Sodium, 26

Symetamine (see Terephthalamidine), 26 Synandrets (see Methyltestosterone), 78 Synandrol (see Testosterone Propionate), 80 Synandrol F (see Testostosterone), 80 Synandrotabs (see Methyltestosterone), 78 Synarel (see Nafarelin Acetate), 78 Synchrosyn (see Chlormadinone* Acetate), 73 Synchrosyn P (see Chlormadinone* Acetate), 73 Syncort (adrenocortical steroid) (see Desoxycorticosterone Acetate), 74 Synerone# (see Testosterone Propionate), 80 Synestrol (see Hexestrol), 77 Synestrol# (see Dienestrol), 75 Syngesterone (see Progesterone), 80 Syngestrets (see Progesterone), 80 Synmiol (radiosensitizer) (see Idoxuridine*), 42 Synoestrol (see Hexestrol), 77 Synthoestrin (see Diethylstilbestrol), 75 Synthovo (see Hexestrol), 77 Synthroid (see Levothyroxine Sodium), 77 Syntocinon (see Oxytocin), 79 Syntofolin (see Diethylstilbestrol), 75 Syntrogene (see Hexestrol), 77 SZ (see Streptozocin), 25 SZC (see Streptozocin), 25

T

Tabloid (see Thioguanine*), 26 TAC-3 (see Triamcinolone Acetonide), 81 TAC-40# (see Triamcinolone Acetonide). 81 TAC-D# (see Triamcinolone Diacetate), 81 TACE (see Chlorotrianisene), 74 Tagamet (see Cimetidine), 40 Taliblastine (see Thalicarpine), 26 Talisomycin, 26 Tallysomycin A# (see Talisomycin), 26 TAM (see Tamoxifen Citrate), 80 Tamofen (see Tamoxifen Citrate*) 80 Tamoxifen (see Tamoxifen Citrate), 80 Tamoxifen Citrate, 80 TAP-144 (see Leuprolide Acetate), 77 Tarabine (see Cytarabine), 51 TATBA (see Triamcinolone Hexacetonide), 81 Taxol (see Paclitaxel), 24

Taxotere, 26 Tazobactam, 26 TC (see Thalicarpine), 26 3TC, 94 TCAR (see Tiazofurin), 26 TCGF (see Interleukin-2), 63 TCN (see Triciribine phosphate), 27 TCN-P (see Triciribine Phosphate), 27 TdR (see Thymidine), 26 Teebacin (see Para-aminosalicylic acid), 98 Teebaconin (see Isoniazid), 97 Tegafur, 26 Tegison (see Etretinate), 51 TEM (see Triethylenemelamine), 27 Tenex (see Guanfacine HCl), 42 Teniposide, 26 Teolit (see Testolactone*), 80 Tepirone (see Razoxane), 25 Terazol (see Terconazole), 100 Terconazole, 100 Terephthalamidine, 26 Terolut (see Dydrogesterone*), 75 Teroxirone, 26 Terrastatin, component of (see Nystatin), 98 Tertroxin (see Liothyronine Sodium), 77 Teslac (see Testolactone*), 80 Teslak (see Testolactone*), 80 TESP (see Thiotepa*), 26 Tespa (see Thiotepa*), 26 Tespamin (see Thiotepa*), 26 Tespamine (see Thiotepa*), 26 Testolacton (see Testolactone*), 80 Testolactone, 80 Testoral (see Fluoxymesterone), 76 Testosteroid (see Testosterone), 80 Testosterone, 80 Testosterone Cypionate, 80 Testosterone Enanthate, 80 Testosterone Ketolaurate, 80 Testosterone Phenylacetate#, 80 Testosterone Propionate, 80 Testoviron (see Methyltestosterone), 78 Testred (ICN) (see Methyltestosterone), 78

Testryl# (see Fluoxymesterone), 76

TET (see Triethylenemelamine), 27 Tetrahydrocannabinol, 46 Tetramizole (see Levamisole HCl), 64 Tetrandirine (see D-Tetrandrine), 26 Tetrandrin (see D-Tetrandrine), 26 Tetraplatin, 26 Tetrastatin, component of (see Nystatin), 98 TF-128 (see Aminothiadiazole), 15 TFT (see Trifluridine), 100 TG (see Thioguanine*), 26 6-TG (see Thioguanine*), 26 TGdR (see Beta-TGdR), 16 TGU (see Triglycidylurazol), 27 Thaliblastine (see Thalicarpine), 26 Thalicarpin (see Thalicarpine), 26 Thalicarpine, 26 THC (see Tetrahydrocannabinol), 46 Theelin (see Estrone), 76 Theelol# (see Estriol), 76 Thenylidene (see Teniposide), 26 Thenylidine-Lignan-P (see Teniposide), 26 Theophylline, 46 Theprubicin, 26 THF (see Thymic Humoral Factor). 94 Thiabendazole, 100 Thiamiprine, 26 Thiatriamide (see Azetepa#), 16 Thio-TEPA (see Thiotepa*), 26 Thiocarb (see Diethyldithiocarbamate), 41 Thiofozil (see Thiotepa*), 26 Thioguanine*, 26 6-Thioguanine (see Thioguanine*), 26 Thioguanine deoxyriboside (see Beta-TGdR), 16 Thioguanosine deoxyriboside (see Alpha-TGdR), 15 6-Thiohypoxanthine (see Mercaptopurine*), 22 6-Thioinosine (see 6-Mercaptopurine riboside), 22 Thiophosphamide (see Thiotepa*), 26 6-Thiopurine (see Mercaptopurine*), 22 Thiotef (see Thiotepa*), 26 Thiotepa*, 26 6-Thioxopurine (see Mercaptopurine*), 22 Thorazine (anti-emetic (see Chlorpromazine*), 40 THP (see Theprubicin), 26 Thymic Humoral Factor, 94

Thymidin (see Thymidine), 26 Thymidine, 26 Thymine deoxyriboside (see Thymidine), 26 Thymopentin, 94 Thymosin, 66 Thyractin# (see Thyroglobulin), 80 Thyrar (see Thyroid), 81 Thyroglobulin, 80 Thyroid, 81 Thyrolar (see Liotrix), 77 Thyroprotein# (see Thyroglobulin), 80 Thyrotropin, 81 Tiamiprine (see Thiamiprine), 26 Tiazofurin, 26 TIBO derivatives, 94 TIC (see TIC-mustard), 27 TIC NH2 (misc. agent) (see TIC-mustard), 27 TIC-mustard, 27 Tice BCG Vaccine (see Interleukin-2), 64 Tifosyl (see Thiotepa*), 26 Tigan (anti-emetic) (see Trimethobenzamide HCl), 46 Tigestol*, 81 TIL/TNF, 66 Tilorone, 66 Timp-2, 27 Tindurin (see Pyrimethamine), 45 Tinver, component of (see Sodium Thiosulfate), 45 Tio-TEF (see Thiotepa*), 26 Tioconazole, 100 Tiofosfamid (see Thiotepa*), 26 Tiofosyl (Sweden) (see Thiotepa*), 26 Tiofozil (see Thiotepa*), 26 Tioguanin (see Thioguanine*), 26 Tisin# (see Isoniazid*), 97 TL (see Testolactone*), 80 TL 476 (see Aniline mustard), 15 TLC G-65, 100 TLC-Dox 99 (see Liposome-Doxorubicin Kit), 21 TMCA, 27 TMP (see Trimethoprim), 46 TMP/SMX (see Bactrim/Septra), 95 TMTX (see Trimetrexate), 27

TMX (see Tamoxifen Citrate), 80

TNCA (see TMCA), 27

TNF (see Tumor necrosis factor), 66

TNF Transd TIL (GTI), 66

TNO-6 (see Spiroplatin), 25

TNP-470, 94, 100

Tobradex, component of (see Dexamethasone), 40, 74

Tomosar (see Menogaril), 22

Topotecan HCl, 27

Toradol (anti-inflammatory; analgesic) (see Ketorolac Tromethamine), 42

Toremifene Citrate, 81

Toyamicin (see Chromomycin A3#), 17

Toyocamycin, 27

Toyokamycin (see Toyocamycin), 27

Toyomycin (see Chromomycin A3#), 17

TP (see Testosterone Propionate), 80

TP OKT 3 Lymphoctyes, 66

TP-5 (see Thymopentin), 94

Tralonide#, 81

Tramacin# (see Triamcinolone Acetonide), 81

Tramisole (see Levamisole HCl), 64

trans-Diethylstilbestrol (see Diethylstibestrol), 75

Trans-Methyl CCNU (see Semustine), 25

Trecator-SC (see Ethionamide), 96

Trental (vasodilator) (see Pentoxifylline), 44

Treosulfan, 27

Trestolone Acetate#, 81

Tretamine (see Triethylenemelamine), 27

Tretinoin, 51

Trexan (see Naltrexone), 44

Triacet (see Triamcinolone Acetonide), 81

Triamcinolone Acetonide Sodium Phosphate, 81

Triamcinolone Acetonide, 81

Triamcinolone Diacetate, 81

Triamcinolone Hexacetonide, 81

Triamcinolone, 81

Triamelin (see Triethylenemelamine), 27

Triamolone 40 (see Triamcinolone Diacetate), 81

Triamonide 40 (see Triamcinolone Acetonide), 81

Triavil (see Perphenazine), 44

Triazinate (see Baker's antifol), 16

Triazine antifol, 27

Triazinetrione triepoxide (see Teroxirone), 26

Triazure# (see Azaribine*), 15

Trichosanthin (see GLQ223), 92

Triciribine Phosphate, 27

Tricyclic nucleoside 5'-phosphate# (see Triciribine Phosphate), 27 Tricycloside phosphate (see Triciribine Phosphate), 27 Triethylene thiophosphoramide (see Thiotepa*), 26 Triethylenemelamine, 27 Trifluorothymidine (see F3TDR), 20 Triflupromazine HCl*, 46 Trifluridine (see F3TDR), 20 Trifluridine, 94, 100 Triglycidylurazol, 27 Trilafon (see Perphenazine), 44 Trilostane, 81 Trimethobenzamide HCl, 46 Trimethoprim (see Bactrim/Septra), 95 Trimethoprim, 46, 100 Trimethylcolchicinic acid, methyl ester, L-tartrate (1:1) (see TMCA), 27 Trimetrexate glucuronate (see Trimetrexate), 27 Trimetrexate, 27, 100 Trimpex (see Trimethoprim), 46, 100 Trioxifene Mesylate, 81 Triptorelin, 81 Trisodium phosphonoformate (see Foscarnet Sodium), 97 Trityl cysteine, 27 Tritylthioalanine (see Trityl cysteine), 27 Trofosfamide, 27 Tropisetron, 46 Trymex# (see Triamcinolone Acetonide), 81 TSP (see Thiotepa*), 26 TSPA (see Thiotepa*), 26 TT-6 (see Thioguanine*), 26 TT-G (see Thioguanine*), 26 Tubercidin (see Tubercidine), 27 Tubercidine, 27 Tumor necrosis factor, 66 Tumor primed Anti-CD3, 66 Turloc (see Meturedepa), 23 Tymtram (alkylating agent) (see Cyclophosphamide*), 18 TZT (see Baker's antifol), 16 U U-10,974 (see Flumethasone*), 76 U-10,997 (see Mibolerone), 78 U-10071 (see Tubercidine), 27

U-11100A (see Nafoxidine HCL), 78

```
U-12,019E (see Methylprednisolone Sodium Phosphate), 78
U-12,241 (see Circlemycin*), 17
U-13,933 (see Asperlin#), 15
U-14,743 (see Porfiromycin), 25
U-15,614 (see Trestolone Acetate#), 81
U-15167 (see Nogalamycin#), 23
U-19183 (see Sparsomycin), 25
U-19763 (see Bolasterone#), 73
U-19920 (see Cytarabine), 51
U-19920A (see Cytarabine HCl),
U-22,550 (see Cabasterone), 73
U-27,182 (see Flurbiprofen), 41
U-31,889 (see Alprazolam), 39
U-42,126 (see Acivicin), 15
U-52,047 (see Menogaril), 22
U-54,461 (see Bropirimine), 16
U-73,975 (see Adozelesin), 15
U-77,233 (see Tetraplatin), 26
U-78,938 (see Dexormaplatin), 18
U-7800 (glucocorticoid) (see Fluprednisolone*), 76
U-8344 (see Uracil Mustard*), 27
U-8471 (see Medrysone*), 77
U-8614 (see Fluorometholone), 76
U-87201, 100
U-8839 (see Medroxyprogesterone Acetate), 77
U-90152, 94
U-9889 (see Streptozocin), 25
Uendex (see Dextran Sulfate), 96
UHN2 (see Uracil Mustard*), 27
Ultandren (see Fluoxymesterone), 76
Ultracorten (see Prednisone), 79
UN 2579 (see Piperazine), 44
UN 2685 (see Piperazine), 44
Unamycin B (see Toyocamycin), 27
Uracil (see Uracil Mustard*), 27
Uracil Mustard*, 27
Uracil riboside (see Uridine), 27
Uractone (see Spironolactone), 80
Uramustin (see Uracil Mustard*), 27
Uramustine (see Uracil Mustard*), 27
Urbason (see Methylprednisolone), 78
Urd (see Uridine), 27
Urea (see Hydroxyurea*), 21
Uredepa, 27
```

Urethane, 27
Urethimine (see Uredepa), 27
Uricemil (see Allopurinol*), 39
Uridin (see Uridine), 27
Uridine, 27
Uriprim (see Allopurinol*), 39
Urokinase, 46
Urosin (see Allopurinol*), 39
Uticort (see Betamethasone Benzoate), 73
Uvilon (see Piperazine), 44

V

Vagestrol# (see Diethylstilbestrol), 75 Vagistat (see Tioconazole), 100 Valisone (see Betamethasone Valerate), 73 Vancenase (see Beclomethasone Dipropionate), 73 Vanceril (see Beclomethasone Dipropionate), 73 Vapreotide, 27 Vasocidin Solution (see Prednisolone Sodium Phosphate), 79 Vasorome (see Oxandrolone*), 79 VaxSyn, 94 VBL (see Vinblastine* Sulfate), 27 VCN (see Vibrio cholera neuraminidase), 66 VCR (see Vincristine* Sulfate), 28 Vee M-26 (see Teniposide), 26 Vehem (see Teniposide), 26 VELB (see Vinblastine* Sulfate), 27 Velban (see Vinblastine* Sulfate), 27 Velsar (see Vinblastine Sulfate), 27 Vengicide (see Toyocamycin), 27 Ventolin (see Albuterol), 39 Vepesid (see Etoposide*), 20 Veramix (see Medroxyprogesterone Acetate), 77 Verapamil (HCl)*, 46 Verapamil (Isoptrin) (see Verapamil (HCl)*), 46 Vercyte (see Pipobroman*), 24 Verelan (see Verapamil (HCl)*), 46 Vermex (see Piperazine), 44 Verospirone (see Spironolactone), 80 Versed (anesthetic; CNS depressant) (see Midazolam HCI), 44 Vesprin (see Triflupromazine HCl), 46 Vibrio cholera neuraminidase, 66 Vidarabin (see Vidarabine*), 27

Vidarabin-Thilo (see Vidarabine), 27

Vidarabine, 100

Vidarabine*, 27

Videx (see Dideoxyinosine), 92

Vinblastine (see Vinblastine* Sulfate), 27

Vinblastine* Sulfate, 27

Vincaleukoblastine (see Vincristine* Sulfate), 28

Vincaleukoblastine sulfate (see Vinblastine* Sulfate), 27

Vincasar (see Vincristine* Sulfate), 28

Vincristine (see Vincristine* Sulfate), 28

Vincristine* Sulfate, 28

Vincrystine (see Vincristine* Sulfate), 28

Vindesine Sulfate, 28

Vindesine, 28

Vinepidine Sulfate, 28

Vinglycinate Sulfate, 28

Vinkristin (see Vincristine* Sulfate), 28

Vinleurosine (see Vinleurosine Sulfate#), 28

Vinleurosine Sulfate#, 28

Vinorelbine, 28

Vinostatin (see Zinostatin), 28

Vinrosidine Sulfate#, 28

Vinzolidine Sulfate, 28

Vira-A (see Vidarabine*), 27

Virazole (see Ribavirin), 94

Vistaril (see Hydroxyzine HC1), 42

Vistrax (tranquilizer) (see Hydroxyzine HC1), 42

Vit A (see Retinoic acid), 51

Vit A (see Tretinoin), 51

Vit D3 (see 1,25-Dihydroxy-vitamin D3), 51

Vitamin A acid (see Tretinoin), 51

13-cis-Vitamin A acid (see Retinoic acid), 51

Vitamin B6 (see Pyridoxine HCl), 45

Vitamin D, 81

Vitamin E, 46

Vitarabin (see Vidarabine*), 27

Vitarin (see Vidarabine*), 27

VLB (see Vinblastine* Sulfate), 27

VM-26 (see Teniposide), 26

VNB (see Vinorelbine), 28

VNCR (see Vincristine* Sulfate), 28

VP-16 (see Etoposide*), 20

VP-16-213 (see Etoposide*), 20

Vumon (see Teniposide), 26

VZL (see Vinzolidine Sulfate), 28

W

W-3399 (see Quingestrone#), 80 W-3566 (see Quinestrol), 80 W-4540 (see Quingestanol* Acetate), 80 W-4869 (see Prednival*#), 79 W-5975 (see Betamethasone Benzoate), 73 W-6495 (see Oxisuran*#), 23 Warfarin Sodium*, 46 Wazine-34 (see Piperazine), 44 Wellcome CN6134 (BW) (see C-parvum), 62 Wellcovorin (see Leucovorin (calcium)*), 43 WIN 14833 (see Stanozolol*), 80 Win 24,540 (see Trilostane), 81 Win-27,914 (see Nivazol), 79 Winstrol (see Stanozolol*), 80 Wobenzym, 94 WR 14997 (see Cycloleucin), 18 WR 2026, 46 WR 2721 (see Ancillary drugs), 28 WR 2721, 28, 46 WR 6026, 100 WR-2721 (radioprotector) (see Amifostine), 39 Wurmirazin (anthelmintic) (see Piperazine), 44 Wy-4036 (anti-emetic) (see Lorazepam), 43 Wy-5321 (see Diglycoaldehyde), 19 X XA 146-85-2 (see Pyran copolymer), 65 Xanax (sedative) (see Alprazolam), 39 Xaridium (see Sodium Fluoride), 45 XB 2615 (see Teroxirone), 26 Xylocaine (see Lidocaine), 43 Y Yodoxin (see Iodoquinol), 97 Yoshi-864, 28 YTR 830 (see Tazobactam), 26 \mathbf{Z} Z-4942 (alkylating agent) (see Ifosfamide*), 21 Zanosar (see Streptozocin), 25 Zantac (see Ranitidine HCl), 99

Zeniplatin, 28
Zidovudine (see Azidothymidine), 16, 91
Zidovudine*, 28
Zinostatin, 28
Zitostop (see Mannosulfan), 22
Zofran (see Ondansetron HCl), 44
Zoladex (see Gosrelin), 77, 81
Zorubicin (HCl), 28
Zovirax (see Acyclovir Sodium), 39, 95
Zuclomiphene citrate (see Clometherone#), 74
Zuclomiphene citrate (see Clomiphene* Citrate), 74
Zyloprim (see Allopurinol*), 39
Zyloric (see Allopurinol*), 39
Zymafluor (see Sodium Fluoride), 45