

**Table 1.51A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	46,558	14,680	6,210
<b>AGE</b>			
12-17	3,389	2,283	979
18-25	8,587	4,411	1,683
26 or Older	34,582	7,986	3,549
<b>GENDER</b>			
Male	24,371	7,472	3,086
Female	22,187	7,208	3,125
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	41,814	12,635	5,366
White	36,210	10,881	4,616
Black or African American	3,626	1,122	526
American Indian or Alaska Native	395	99	47
Native Hawaiian or Other Pacific Islander	171	82	31
Asian	962	283	59
Two or More Races	450	167	88
Hispanic or Latino	4,744	2,046	844

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.51B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	19.8	6.2	2.6
<b>AGE</b>			
12-17	13.7	9.2	4.0
18-25	27.7	14.2	5.4
26 or Older	19.3	4.5	2.0
<b>GENDER</b>			
Male	21.5	6.6	2.7
Female	18.3	5.9	2.6
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	20.3	6.1	2.6
White	21.9	6.6	2.8
Black or African American	13.5	4.2	2.0
American Indian or Alaska Native	26.7	6.7	3.2
Native Hawaiian or Other Pacific Islander	21.0	10.0	3.8
Asian	10.6	3.1	0.7
Two or More Races	17.9	6.7	3.5
Hispanic or Latino	16.3	7.0	2.9

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.52A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	3,389	2,283	979
<b>GENDER</b>			
Male	1,641	1,091	456
Female	1,748	1,192	523
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	2,829	1,924	826
White	2,288	1,592	678
Black or African American	370	224	108
American Indian or Alaska Native	40	27	8
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	63	38	22
Two or More Races	47	33	8
Hispanic or Latino	560	359	153
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	1,113	754	314
Female - White	1,175	838	363
Male - Black or African American	163	97	39
Female - Black or African American	207	128	68
Male - Hispanic or Latino	285	192	87
Female - Hispanic or Latino	276	167	66

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.52B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	13.7	9.2	4.0
<b>GENDER</b>			
Male	13.0	8.6	3.6
Female	14.4	9.8	4.3
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	13.6	9.2	4.0
White	14.7	10.3	4.4
Black or African American	10.2	6.2	3.0
American Indian or Alaska Native	21.2	14.3	4.4
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	6.3	3.7	2.2
Two or More Races	12.0	8.3	2.0
Hispanic or Latino	14.3	9.1	3.9
<b>GENDER/RACE/HISPANIC ORIGIN</b>			
Male - White	14.0	9.5	3.9
Female - White	15.5	11.1	4.8
Male - Black or African American	9.0	5.3	2.2
Female - Black or African American	11.5	7.1	3.8
Male - Hispanic or Latino	14.2	9.6	4.4
Female - Hispanic or Latino	14.4	8.7	3.4

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.53A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	8,587	4,411	1,683
<b>GENDER</b>			
Male	4,600	2,371	912
Female	3,987	2,040	770
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	7,459	3,865	1,467
White	6,461	3,397	1,271
Black or African American	574	275	125
American Indian or Alaska Native	*	41	*
Native Hawaiian or Other Pacific Islander	*	15	*
Asian	172	64	11
Two or More Races	125	74	32
Hispanic or Latino	1,127	546	215
<b>EDUCATION</b>			
< High School	1,930	1,048	403
High School Graduate	2,945	1,433	575
Some College	2,816	1,515	551
College Graduate	896	415	153
<b>CURRENT EMPLOYMENT</b>			
Full-Time	4,178	2,030	759
Part-Time	2,049	1,084	395
Unemployed	764	447	179
Other <sup>1</sup>	1,596	850	350

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.53B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	27.7	14.2	5.4
<b>GENDER</b>			
Male	29.6	15.3	5.9
Female	25.7	13.2	5.0
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	29.2	15.1	5.7
White	33.5	17.6	6.6
Black or African American	14.1	6.7	3.1
American Indian or Alaska Native	*	18.8	*
Native Hawaiian or Other Pacific Islander	*	9.1	*
Asian	12.5	4.7	0.8
Two or More Races	31.1	18.3	7.9
Hispanic or Latino	20.5	9.9	3.9
<b>EDUCATION</b>			
< High School	28.1	15.2	5.9
High School Graduate	27.8	13.5	5.4
Some College	28.8	15.5	5.6
College Graduate	23.6	10.9	4.0
<b>CURRENT EMPLOYMENT</b>			
Full-Time	28.8	14.0	5.2
Part-Time	25.9	13.7	5.0
Unemployed	31.1	18.2	7.3
Other <sup>1</sup>	25.9	13.8	5.7

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.54A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	34,582	7,986	3,549
<b>GENDER</b>			
Male	18,130	4,010	1,717
Female	16,452	3,976	1,831
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	31,526	6,845	3,073
White	27,461	5,892	2,667
Black or African American	2,682	624	293
American Indian or Alaska Native	270	32	12
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	727	181	26
Two or More Races	278	61	48
Hispanic or Latino	3,056	1,141	476
<b>EDUCATION</b>			
< High School	4,353	1,406	692
High School Graduate	10,487	2,702	1,259
Some College	10,085	2,153	945
College Graduate	9,658	1,724	652
<b>CURRENT EMPLOYMENT</b>			
Full-Time	23,501	5,159	2,126
Part-Time	4,154	995	491
Unemployed	1,435	532	233
Other <sup>1</sup>	5,493	1,300	699

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.54B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	19.3	4.5	2.0
<b>GENDER</b>			
Male	21.2	4.7	2.0
Female	17.5	4.2	1.9
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	19.7	4.3	1.9
White	21.0	4.5	2.0
Black or African American	14.0	3.3	1.5
American Indian or Alaska Native	25.1	2.9	1.1
Native Hawaiian or Other Pacific Islander	*	*	*
Asian	10.9	2.7	0.4
Two or More Races	16.2	3.5	2.8
Hispanic or Latino	15.5	5.8	2.4
<b>EDUCATION</b>			
< High School	14.3	4.6	2.3
High School Graduate	18.3	4.7	2.2
Some College	23.6	5.0	2.2
College Graduate	19.8	3.5	1.3
<b>CURRENT EMPLOYMENT</b>			
Full-Time	23.0	5.1	2.1
Part-Time	21.3	5.1	2.5
Unemployed	28.0	10.4	4.5
Other <sup>1</sup>	10.4	2.5	1.3

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.



**Table 1.55A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	43,169	12,397	5,231
<b>GENDER</b>			
Male	22,730	6,381	2,630
Female	20,439	6,016	2,601
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	38,985	10,711	4,540
White	33,922	9,289	3,938
Black or African American	3,256	898	418
American Indian or Alaska Native	355	72	39
Native Hawaiian or Other Pacific Islander	150	72	28
Asian	899	245	37
Two or More Races	403	135	80
Hispanic or Latino	4,184	1,686	691
<b>EDUCATION</b>			
< High School	6,282	2,454	1,095
High School Graduate	13,432	4,135	1,834
Some College	12,901	3,669	1,496
College Graduate	10,553	2,140	805
<b>CURRENT EMPLOYMENT</b>			
Full-Time	27,680	7,189	2,885
Part-Time	6,202	2,079	886
Unemployed	2,198	979	411
Other <sup>1</sup>	7,089	2,150	1,049

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.

**Table 1.55B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2002**

Demographic Characteristic	TIME PERIOD		
	Lifetime	Past Year	Past Month
<b>TOTAL</b>	20.5	5.9	2.5
<b>GENDER</b>			
Male	22.5	6.3	2.6
Female	18.7	5.5	2.4
<b>HISPANIC ORIGIN AND RACE</b>			
Not Hispanic or Latino	21.0	5.8	2.5
White	22.6	6.2	2.6
Black or African American	14.0	3.9	1.8
American Indian or Alaska Native	27.5	5.6	3.0
Native Hawaiian or Other Pacific Islander	20.6	9.8	3.9
Asian	11.2	3.0	0.5
Two or More Races	19.0	6.3	3.8
Hispanic or Latino	16.6	6.7	2.7
<b>EDUCATION</b>			
< High School	16.9	6.6	2.9
High School Graduate	19.8	6.1	2.7
Some College	24.5	7.0	2.8
College Graduate	20.1	4.1	1.5
<b>CURRENT EMPLOYMENT</b>			
Full-Time	23.8	6.2	2.5
Part-Time	22.6	7.6	3.2
Unemployed	29.0	12.9	5.4
Other <sup>1</sup>	12.0	3.7	1.8

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002.