

APPENDIX A

ATSDR MINIMAL RISK LEVEL AND WORKSHEETS

The Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA) [42 U.S.C. 9601 et seq.], as amended by the Superfund Amendments and Reauthorization Act (SARA) [Pub. L. 99-499], requires that the Agency for Toxic Substances and Disease Registry (ATSDR) develop jointly with the U.S. Environmental Protection Agency (EPA), in order of priority, a list of hazardous substances most commonly found at facilities on the CERCLA National Priorities List (NPL); prepare toxicological profiles for each substance included on the priority list of hazardous substances; and assure the initiation of a research program to fill identified data needs associated with the substances.

The toxicological profiles include an examination, summary, and interpretation of available toxicological information and epidemiologic evaluations of a hazardous substance. During the development of toxicological profiles, Minimal Risk Levels (MRLs) are derived when reliable and sufficient data exist to identify the target organ(s) of effect or the most sensitive health effect(s) for a specific duration for a given route of exposure. An MRL is an estimate of the daily human exposure to a hazardous substance that is likely to be without appreciable risk of adverse noncancer health effects over a specified duration of exposure. MRLs are based on noncancer health effects only and are not based on a consideration of cancer effects. These substance-specific estimates, which are intended to serve as screening levels, are used by ATSDR health assessors to identify contaminants and potential health effects that may be of concern at hazardous waste sites. It is important to note that MRLs are not intended to define clean-up or action levels.

MRLs are derived for hazardous substances using the no-observed-adverse-effect level/uncertainty factor approach. They are below levels that might cause adverse health effects in the people most sensitive to such chemical-induced effects. MRLs are derived for acute (1–14 days), intermediate (15–364 days), and chronic (365 days and longer) durations and for the oral and inhalation routes of exposure. Currently, MRLs for the dermal route of exposure are not derived because ATSDR has not yet identified a method suitable for this route of exposure. MRLs are generally based on the most sensitive chemical-induced end point considered to be of relevance to humans. Serious health effects (such as irreparable damage to the liver or kidneys, or birth defects) are not used as a basis for establishing MRLs. Exposure to a level above the MRL does not mean that adverse health effects will occur.

APPENDIX A

MRLs are intended only to serve as a screening tool to help public health professionals decide where to look more closely. They may also be viewed as a mechanism to identify those hazardous waste sites that are not expected to cause adverse health effects. Most MRLs contain a degree of uncertainty because of the lack of precise toxicological information on the people who might be most sensitive (e.g., infants, elderly, nutritionally or immunologically compromised) to the effects of hazardous substances. ATSDR uses a conservative (i.e., protective) approach to address this uncertainty consistent with the public health principle of prevention. Although human data are preferred, MRLs often must be based on animal studies because relevant human studies are lacking. In the absence of evidence to the contrary, ATSDR assumes that humans are more sensitive to the effects of hazardous substance than animals and that certain persons may be particularly sensitive. Thus, the resulting MRL may be as much as a hundredfold below levels that have been shown to be nontoxic in laboratory animals.

Proposed MRLs undergo a rigorous review process: Health Effects/MRL Workgroup reviews within the Division of Toxicology, expert panel peer reviews, and agencywide MRL Workgroup reviews, with participation from other federal agencies and comments from the public. They are subject to change as new information becomes available concomitant with updating the toxicological profiles. Thus, MRLs in the most recent toxicological profiles supersede previously published levels. For additional information regarding MRLs, please contact the Division of Toxicology, Agency for Toxic Substances and Disease Registry, 1600 Clifton Road, Mailstop E-29, Atlanta, Georgia 30333.

APPENDIX A

MINIMAL RISK LEVEL WORKSHEET

Chemical Name: Ammonia
CAS Number: 7664-41-7
Date: September 2002
Profile Status: Draft 3 Pre Public
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 14
Species: Human

Minimal Risk Level: 1.7 mg/kg/day ppm

Reference: Verberk MM. 1977. Effects of ammonia in volunteers. Int Arch Occup Environ Health 39:73-81.

Experimental design: Sixteen volunteers (8 science faculty with knowledge of the effects of ammonia and 8 non-science university students not familiar with ammonia health effects) were exposed 4 at a time to 50, 80, 110, or 140 ppm ammonia for 2 hours. Each group was exposed to each exposure level with 1 week in between exposures. Immediately before and after exposure, respiratory function tests (vital capacity [VC], forced expiratory volume in the first second [FEV₁], and forced inspiratory volume in the first second [FIV₁]) were done. During exposure, each participant recorded subjective effect levels for smell, taste, irritation of eyes, irritation of nose, irritation of throat, irritation of breast, urge to cough, headache, and general discomfort. The scale used was: 0=no sensation, 1=just perceptible, 2=distinctly perceptible, 3=nuisance, 4=offensive, and 5=unbearable. A (+) or (-) could be used to interpolate between the levels. A few weeks after the experiments, the histamine threshold was determined for 13 of the 16 volunteers as a measure of pre-existing non-specific reactivity of the airways to exogenous stimuli.

Effects noted in study and corresponding doses: None of the participants was hypersusceptible to non-specific irritants. No participant had a decrease of more than 10% of pre-exposure values for VC, FEV₁, or FIV₁. There was a difference between the science faculty group (experts) and the students for the subjective scoring. Students consistently scored higher for smell and there was little increase in score with concentration. Score for irritation of the eyes increased with concentration and there was no difference between groups. Irritation of the throat had a sharp increase in score with concentration and scores were higher for students. All students left the exposure chamber between 0.5 and 1.25 hours in the 140 ppm exposure because of severe irritation. Scores for urge to cough and general discomfort were low in the expert group, but increased with concentration in the student group. All students left the chamber before 2 hours of exposure to 140 ppm.

Dose and end point used for MRL derivation: 50 ppm for mild irritation to the eyes, nose, and throat in humans exposed to ammonia gas for 2 hours.

NOAEL LOAEL

Uncertainty Factors used in MRL derivation:

- 3 for use of a minimal LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

APPENDIX A

Was a conversion used from ppm in food or water to a mg/body weight dose?

If so, explain: None needed.

If an inhalation study in animals, list the conversion factors used in determining human equivalent dose:

N/A

Other additional studies or pertinent information which lend support to this MRL: The MRL is supported by other observations of respiratory effects associated with acute- and intermediate-duration exposure including transient irritation of the nose and throat of humans exposed to 100 ppm (Ferguson et al. 1977); nasal discharge in rats at 376 ppm (Coon et al. 1970); nasal lesions in rats at 150 ppm (Broderick et al. 1976); and nasal inflammation and lesions in rats at 500 ppm (Richard et al. 1978a). A study of piggerie workers exposed to a mean level of 7.9 ppm ammonia measured lung function change over a workshift; a small but borderline significant decrease in lung function was noted (Heederik et al. 1990). This was not used as a basis for MRL derivation because the workers were also exposed to other potential respiratory toxicants (dust and endotoxins).

Agency Contact (Chemical Manager): Nickolette Roney, MPH

APPENDIX A

MINIMAL RISK LEVEL WORKSHEET

Chemical Name: Ammonia
 CAS Number: 7664-41-7
 Date: September 2002
 Profile Status: Draft 3 Pre Public
 Route: Inhalation Oral
 Duration: Acute Intermediate Chronic
 Graph Key: 49
 Species: Human

Minimal Risk Level: 0.3 mg/kg/day ppm

Reference: Holness DL, Purdham JT, Nethercott JR. 1989. Acute and chronic respiratory effects of occupational exposure to ammonia. Am Ind Hyg Assoc J 50:646–650.

Experimental design: Fifty-two workers and 6 maintenance workers at a soda ash facility were evaluated on 2 days within a week for sense of smell (using detection of pyridine) and lung function parameters (FVC, FEV₁, FEV₁/FVC, FEF₅₀, and FEF₇₅). Each participant filled out a questionnaire regarding prevalence of respiratory symptoms (cough, bronchitis, wheeze, dyspnea, and others), eye and throat irritation, past occupational exposures, working conditions, and smoking history. Thirty-five controls from the plant were tested in the same way. Each worker wore a personal ammonia level monitor during their workshift; average workshift was 8.4 hours. Mean age of the workers was 40.5 years and average length of employment was 15 years. The TWA exposure level for the exposed group was 9.2±1.4 ppm, while the TWA for the controls was 0.3±0.1 ppm. Analysis was performed using each worker's personal exposure and his change in lung function over the workweek. The cohort was also divided into groups that were exposed to low (<6.25 ppm), medium (6.25–12.5 ppm), and high (>12.5 ppm) ammonia levels and analyzed for change in lung function. Differences due to number of years of ammonia exposure was also assessed.

Effects noted in study and corresponding doses: No difference in the prevalence of reporting of symptoms was observed between the control and exposed groups, and the detection threshold for pyridine was similar between the groups. Baseline lung functions were similar between controls and exposed individuals, and no differences in change in lung function over the workweek were seen between the groups. No statistically significant differences were seen between the level of personal exposure and change in lung function or in lung function between low, medium, and high exposed groups. No association was evident between years of exposure and lung function changes.

Dose and end point used for MRL derivation: 12.5 ppm for sense of smell, prevalence of respiratory symptoms (cough, bronchitis, wheeze, dyspnea, and others), eye and throat irritation, and lung function parameters (FVC, FEV₁, FEV₁/FVC, FEF₅₀, and FEF₇₅) in humans exposed for an average of 15 years in a soda ash plant.

NOAEL LOAEL

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

APPENDIX A

Was a conversion used from ppm in food or water to a mg/body weight dose?

If so, explain: None needed. The NOAEL was adjusted for continuous exposure as follows:
 $12.5 \text{ ppm} \times 8.4/24 \text{ hours} \times 5/7 \text{ days} = 3.1 \text{ ppm}$

If an inhalation study in animals, list the conversion factors used in determining human equivalent dose:
N/A

Other additional studies or pertinent information which lend support to this MRL: The MRL is supported by other observations of respiratory effects associated with chronic-duration exposure including an association between exposure to pollutants, including ammonia, in livestock confinement buildings and an increase in respiratory symptoms (such as bronchial reactivity/hyperresponsiveness, inflammation, cough, wheezing, or shortness of breath) and/or a decrease in lung function (such as forced expiratory volume in the first second [FEV_{1.0}], maximum expiratory flow rates [MEF₅₀ and MEF₇₅], and maximal mid-expiratory flow rate [MMEF]) in farmers exposed to ammonia levels of 2.3–20.7 ppm (Choudat et al. 1994; Cormier et al. 2000; Donham et al. 1995, 2000; Heederik et al. 1990; Reynolds et al. 1996; Vogelzang et al. 1997, 2000). The farmers were also exposed to other possible respiratory toxins, such as dust and endotoxins. A cross-sectional study of male workers at two fertilizer factories in Saudi Arabia showed a significant association between exposure to ammonia gas and respiratory symptoms and bronchial asthma (Ballal et al. 1998). No continuous exposure levels could be calculated for these workers because the number of days worked per week was not provided.

Agency Contact (Chemical Manager): Nickolette Roney, MPH

APPENDIX A

MINIMAL RISK LEVEL WORKSHEET

Chemical Name: Ammonia and Ammonium Compounds
CAS Number: 7664-41-7
Date: September 2002
Profile Status: Draft 3 Pre Public
Route: Inhalation Oral
Duration: Acute Intermediate Chronic
Graph Key: 11
Species: Rat

Minimal Risk Level: 0.3 mg NH₄/kg/day ppm

Reference: Gupta BN, Khanna RN, Datta KK. 1979. Toxicological studies of ammonium sulfamate in rat after repeated oral administration. Toxicology 13:45–49.

Experimental design: Groups of 20 adult female albino rats and weanling albino rats of each sex were given a standard diet ad libitum and administered 0, 100, 250, or 500 mg ammonium sulfamate/kg/day in the drinking water 6 days/week for 90 days. Food and water intake were recorded over the 24 hours prior to weighing; animals were weighed twice a week during the first 2 months, then once weekly. Dose of ammonium sulfamate was adjusted to body weight. At the end of 30, 60, and 90 days exposure, six rats from each group were killed, and blood was analyzed for hemoglobin content, packed cell volume, total red cell count, and total and differential white cell counts. A necropsy was performed and histological examination was performed on the heart, liver, stomach, spleen, kidneys, thyroid, adrenal glands, gonads, intestine, lung, and lymph nodes.

Effects noted in study and corresponding doses: The general condition and health of all rats remained good throughout the study, except for one adult in the 250 mg/kg/day group and one male weanling in the 500 mg/kg/day group died on days 64 and 76, respectively, of bronchopneumonia. No differences in body weights were noted except for adult females exposed to 500 mg/kg/day, which were statistically significantly reduced (by 16%; p<0.05). Relative food intake decreased in all groups, but was only statistically significant in the weanlings in the 500 mg/kg/day groups. Similarly, water intake increased generally and was statistically significant in the weanlings in the 500 mg/kg/day groups. No differences were noted in hematology, relative organ weights, or histology.

Dose and end point used for MRL derivation: 39.5 mg/kg/day for weight loss in rats exposed to ammonium sulfamate in drinking water 6 days/week for 90 days.

NOAEL LOAEL

Uncertainty Factors used in MRL derivation:

- 10 for use of a LOAEL
- 10 for extrapolation from animals to humans
- 10 for human variability

APPENDIX A

Was a conversion used from ppm in food or water to a mg/body weight dose?

If so, explain: None needed. The NOAEL was based on mg NH₄/mg/kg, so 100 mg ammonium sulfamate/kg/day=18.04/114.119x100=15.8 mg NH₄/kg/day, adjusted for continuous exposure=15.8x6/7=13.5 mg NH₄/kg/day; 250 mg ammonium sulfamate/kg/day=39.5, adjusted for continuous exposure=33.9 mg NH₄/kg/day; 500 mg ammonium sulfamate/kg/day=79, adjusted for continuous exposure=67.8 mg NH₄/kg/day

If an inhalation study in animals, list the conversion factors used in determining human equivalent dose:

N/A

Other additional studies or pertinent information which lend support to this MRL: Decreased body weight or body weight gain has also been seen in rats exposed orally to 991 mg/kg/day for 330 days (Barzel and Jowsey 1969), to 960 mg/kg/day for 5 days (Noda and Chikamori 1976), or to 3,102 mg/kg/day for 7 or 15 days (Boyano-Adánez et al. 1996). No true controls were included in the Boyano-Adánez et al. (1996) study; a group of rats that received a standard diet that contained a small amount of ammonium (equivalent to 22 mg NH₄⁺/kg/day) was used as the control. It is impossible to tell where the actual NOAEL is from this study. Animals exposed to ammonia vapor via inhalation have also had decreased body weight or weight gain (Diekman et al. 1993; Drummond et al. 1980; Gustin et al. 1994; Richard et al. 1978a; Stombaugh et al. 1969).

Agency Contact (Chemical Manager): Nickolette Roney, MPH

APPENDIX B

USER'S GUIDE

Chapter 1

Public Health Statement

This chapter of the profile is a health effects summary written in non-technical language. Its intended audience is the general public especially people living in the vicinity of a hazardous waste site or chemical release. If the Public Health Statement were removed from the rest of the document, it would still communicate to the lay public essential information about the chemical.

The major headings in the Public Health Statement are useful to find specific topics of concern. The topics are written in a question and answer format. The answer to each question includes a sentence that will direct the reader to chapters in the profile that will provide more information on the given topic.

Chapter 2

Relevance to Public Health

This chapter provides a health effects summary based on evaluations of existing toxicologic, epidemiologic, and toxicokinetic information. This summary is designed to present interpretive, weight-of-evidence discussions for human health end points by addressing the following questions.

1. What effects are known to occur in humans?
2. What effects observed in animals are likely to be of concern to humans?
3. What exposure conditions are likely to be of concern to humans, especially around hazardous waste sites?

The chapter covers end points in the same order they appear within the Discussion of Health Effects by Route of Exposure section, by route (inhalation, oral, dermal) and within route by effect. Human data are presented first, then animal data. Both are organized by duration (acute, intermediate, chronic). *In vitro* data and data from parenteral routes (intramuscular, intravenous, subcutaneous, etc.) are also considered in this chapter. If data are located in the scientific literature, a table of genotoxicity information is included.

The carcinogenic potential of the profiled substance is qualitatively evaluated, when appropriate, using existing toxicokinetic, genotoxic, and carcinogenic data. ATSDR does not currently assess cancer potency or perform cancer risk assessments. Minimal risk levels (MRLs) for noncancer end points (if derived) and the end points from which they were derived are indicated and discussed.

Limitations to existing scientific literature that prevent a satisfactory evaluation of the relevance to public health are identified in the Chapter 3 Data Needs section.

APPENDIX B

Interpretation of Minimal Risk Levels

Where sufficient toxicologic information is available, we have derived minimal risk levels (MRLs) for inhalation and oral routes of entry at each duration of exposure (acute, intermediate, and chronic). These MRLs are not meant to support regulatory action; but to acquaint health professionals with exposure levels at which adverse health effects are not expected to occur in humans. They should help physicians and public health officials determine the safety of a community living near a chemical emission, given the concentration of a contaminant in air or the estimated daily dose in water. MRLs are based largely on toxicological studies in animals and on reports of human occupational exposure.

MRL users should be familiar with the toxicologic information on which the number is based. Chapter 2, "Relevance to Public Health," contains basic information known about the substance. Other sections such as Chapter 3 Section 3.9, "Interactions with Other Substances," and Section 3.10, "Populations that are Unusually Susceptible" provide important supplemental information.

MRL users should also understand the MRL derivation methodology. MRLs are derived using a modified version of the risk assessment methodology the Environmental Protection Agency (EPA) provides (Barnes and Dourson 1988) to determine reference doses for lifetime exposure (RfDs).

To derive an MRL, ATSDR generally selects the most sensitive end point which, in its best judgement, represents the most sensitive human health effect for a given exposure route and duration. ATSDR cannot make this judgement or derive an MRL unless information (quantitative or qualitative) is available for all potential systemic, neurological, and developmental effects. If this information and reliable quantitative data on the chosen end point are available, ATSDR derives an MRL using the most sensitive species (when information from multiple species is available) with the highest NOAEL that does not exceed any adverse effect levels. When a NOAEL is not available, a lowest-observed-adverse-effect level (LOAEL) can be used to derive an MRL, and an uncertainty factor (UF) of 10 must be employed. Additional uncertainty factors of 10 must be used both for human variability to protect sensitive subpopulations (people who are most susceptible to the health effects caused by the substance) and for interspecies variability (extrapolation from animals to humans). In deriving an MRL, these individual uncertainty factors are multiplied together. The product is then divided into the inhalation concentration or oral dosage selected from the study. Uncertainty factors used in developing a substance-specific MRL are provided in the footnotes of the LSE Tables.

Chapter 3**Health Effects****Tables and Figures for Levels of Significant Exposure (LSE)**

Tables (3-1, 3-2, and 3-3) and figures (3-1 and 3-2) are used to summarize health effects and illustrate graphically levels of exposure associated with those effects. These levels cover health effects observed at increasing dose concentrations and durations, differences in response by species, minimal risk levels (MRLs) to humans for noncancer end points, and EPA's estimated range associated with an upper-bound individual lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. Use the LSE tables and figures for a quick review of the health effects and to locate data for a specific exposure scenario. The LSE tables and figures should always be used in conjunction with the text. All entries in these tables and figures represent studies that provide reliable, quantitative estimates of No-Observed-Adverse-Effect Levels (NOAELs), Lowest-Observed-Adverse-Effect Levels (LOAELs), or Cancer Effect Levels (CELs).

APPENDIX B

The legends presented below demonstrate the application of these tables and figures. Representative examples of LSE Table 3-1 and Figure 3-1 are shown. The numbers in the left column of the legends correspond to the numbers in the example table and figure.

LEGEND**See LSE Table 3-1**

- (1) Route of Exposure One of the first considerations when reviewing the toxicity of a substance using these tables and figures should be the relevant and appropriate route of exposure. When sufficient data exists, three LSE tables and two LSE figures are presented in the document. The three LSE tables present data on the three principal routes of exposure, i.e., inhalation, oral, and dermal (LSE Table 3-1, 3-2, and 3-3, respectively). LSE figures are limited to the inhalation (LSE Figure 3-1) and oral (LSE Figure 3-2) routes. Not all substances will have data on each route of exposure and will not therefore have all five of the tables and figures.
- (2) Exposure Period Three exposure periods - acute (less than 15 days), intermediate (15–364 days), and chronic (365 days or more) are presented within each relevant route of exposure. In this example, an inhalation study of intermediate exposure duration is reported. For quick reference to health effects occurring from a known length of exposure, locate the applicable exposure period within the LSE table and figure.
- (3) Health Effect The major categories of health effects included in LSE tables and figures are death, systemic, immunological, neurological, developmental, reproductive, and cancer. NOAELs and LOAELs can be reported in the tables and figures for all effects but cancer. Systemic effects are further defined in the "System" column of the LSE table (see key number 18).
- (4) Key to Figure Each key number in the LSE table links study information to one or more data points using the same key number in the corresponding LSE figure. In this example, the study represented by key number 18 has been used to derive a NOAEL and a Less Serious LOAEL (also see the 2 "18r" data points in Figure 3-1).
- (5) Species The test species, whether animal or human, are identified in this column. Chapter 2, "Relevance to Public Health," covers the relevance of animal data to human toxicity and Section 3.4, "Toxicokinetics," contains any available information on comparative toxicokinetics. Although NOAELs and LOAELs are species specific, the levels are extrapolated to equivalent human doses to derive an MRL.
- (6) Exposure Frequency/Duration The duration of the study and the weekly and daily exposure regimen are provided in this column. This permits comparison of NOAELs and LOAELs from different studies. In this case (key number 18), rats were exposed to 1,1,2,2-tetrachloroethane via inhalation for 6 hours per day, 5 days per week, for 3 weeks. For a more complete review of the dosing regimen refer to the appropriate sections of the text or the original reference paper, i.e., Nitschke et al. 1981.
- (7) System This column further defines the systemic effects. These systems include: respiratory, cardiovascular, gastrointestinal, hematological, musculoskeletal, hepatic, renal, and dermal/ocular. "Other" refers to any systemic effect (e.g., a decrease in body weight) not covered in these systems. In the example of key number 18, 1 systemic effect (respiratory) was investigated.

APPENDIX B

- (8) NOAEL A No-Observed-Adverse-Effect Level (NOAEL) is the highest exposure level at which no harmful effects were seen in the organ system studied. Key number 18 reports a NOAEL of 3 ppm for the respiratory system which was used to derive an intermediate exposure, inhalation MRL of 0.005 ppm (see footnote "b").
- (9) LOAEL A Lowest-Observed-Adverse-Effect Level (LOAEL) is the lowest dose used in the study that caused a harmful health effect. LOAELs have been classified into "Less Serious" and "Serious" effects. These distinctions help readers identify the levels of exposure at which adverse health effects first appear and the gradation of effects with increasing dose. A brief description of the specific end point used to quantify the adverse effect accompanies the LOAEL. The respiratory effect reported in key number 18 (hyperplasia) is a Less serious LOAEL of 10 ppm. MRLs are not derived from Serious LOAELs.
- (10) Reference The complete reference citation is given in Chapter 9 of the profile.
- (11) CEL A Cancer Effect Level (CEL) is the lowest exposure level associated with the onset of carcinogenesis in experimental or epidemiologic studies. CELs are always considered serious effects. The LSE tables and figures do not contain NOAELs for cancer, but the text may report doses not causing measurable cancer increases.
- (12) Footnotes Explanations of abbreviations or reference notes for data in the LSE tables are found in the footnotes. Footnote "b" indicates the NOAEL of 3 ppm in key number 18 was used to derive an MRL of 0.005 ppm.

LEGEND**See Figure 3-1**

LSE figures graphically illustrate the data presented in the corresponding LSE tables. Figures help the reader quickly compare health effects according to exposure concentrations for particular exposure periods.

- (13) Exposure Period The same exposure periods appear as in the LSE table. In this example, health effects observed within the intermediate and chronic exposure periods are illustrated.
- (14) Health Effect These are the categories of health effects for which reliable quantitative data exists. The same health effects appear in the LSE table.
- (15) Levels of Exposure concentrations or doses for each health effect in the LSE tables are graphically displayed in the LSE figures. Exposure concentration or dose is measured on the log scale "y" axis. Inhalation exposure is reported in mg/m³ or ppm and oral exposure is reported in mg/kg/day.
- (16) NOAEL In this example, the open circle designated 18r identifies a NOAEL critical end point in the rat upon which an intermediate inhalation exposure MRL is based. The key number 18 corresponds to the entry in the LSE table. The dashed descending arrow indicates the extrapolation from the exposure level of 3 ppm (see entry 18 in the Table) to the MRL of 0.005 ppm (see footnote "b" in the LSE table).
- (17) CEL Key number 38r is 1 of 3 studies for which Cancer Effect Levels were derived. The diamond symbol refers to a Cancer Effect Level for the test species-mouse. The number 38 corresponds to the entry in the LSE table.

APPENDIX B

- (18) Estimated Upper-Bound Human Cancer Risk Levels This is the range associated with the upper-bound for lifetime cancer risk of 1 in 10,000 to 1 in 10,000,000. These risk levels are derived from the EPA's Human Health Assessment Group's upper-bound estimates of the slope of the cancer dose response curve at low dose levels (q_1^*).
- (19) Key to LSE Figure The Key explains the abbreviations and symbols used in the figure.

SAMPLE

1 6

Table 3-1. Levels of Significant Exposure to [Chemical x] – Inhalation

Key to figure ^a	Species	Exposure frequency/ duration	System	NOAEL (ppm)	LOAEL (effect)		Reference
					Less serious (ppm)	Serious (ppm)	
INTERMEDIATE EXPOSURE							
	5	6	7	8	9		10
3 6	Systemic	9	9	9	9		9
4 6	18	Rat	13 wk 5 d/wk 6 hr/d	Resp	3 ^b	10 (hyperplasia)	Nitschke et al. 1981
CHRONIC EXPOSURE							
						11	
	Cancer					9	
38	Rat	18 mo 5 d/wk 7 hr/d				20	(CEL, multiple organs) Wong et al. 1982
39	Rat	89–104 wk 5 d/wk 6 hr/d				10	(CEL, lung tumors, nasal tumors) NTP 1982
40	Mouse	79–103 wk 5 d/wk 6 hr/d				10	(CEL, lung tumors, hemangiosarcomas) NTP 1982

12 6

^a The number corresponds to entries in Figure 3-1.


^b Used to derive an intermediate inhalation Minimal Risk Level (MRL) of 5×10^{-3} ppm; dose adjusted for intermittent exposure and divided by an uncertainty factor of 100 (10 for extrapolation from animal to humans, 10 for human variability).

DRAFT FOR PUBLIC COMMENT

APPENDIX B

SAMPLE

Figure 3-1. Levels of Significant Exposure to [Chemical X] - Inhalation



DRAFT FOR PUBLIC COMMENT

APPENDIX B

APPENDIX C

ACRONYMS, ABBREVIATIONS, AND SYMBOLS

ACOEM	American College of Occupational and Environmental Medicine
ACGIH	American Conference of Governmental Industrial Hygienists
ADI	acceptable daily intake
ADME	absorption, distribution, metabolism, and excretion
AED	atomic emission detection
AOEC	Association of Occupational and Environmental Clinics
AFID	alkali flame ionization detector
AFOSH	Air Force Office of Safety and Health
ALT	alanine aminotransferase
AML	acute myeloid leukemia
AOAC	Association of Official Analytical Chemists
AP	alkaline phosphatase
APHA	American Public Health Association
AST	aspartate aminotransferase
atm	atmosphere
ATSDR	Agency for Toxic Substances and Disease Registry
AWQC	Ambient Water Quality Criteria
BAT	best available technology
BCF	bioconcentration factor
BEI	Biological Exposure Index
BSC	Board of Scientific Counselors
C	centigrade
CAA	Clean Air Act
CAG	Cancer Assessment Group of the U.S. Environmental Protection Agency
CAS	Chemical Abstract Services
CDC	Centers for Disease Control and Prevention
CEL	cancer effect level
CELDS	Computer-Environmental Legislative Data System
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
CFR	Code of Federal Regulations
Ci	curie
CI	confidence interval
CL	ceiling limit value
CLP	Contract Laboratory Program
cm	centimeter
CML	chronic myeloid leukemia
CPSC	Consumer Products Safety Commission
CWA	Clean Water Act
DHEW	Department of Health, Education, and Welfare
DHHS	Department of Health and Human Services
DNA	deoxyribonucleic acid
DOD	Department of Defense
DOE	Department of Energy
DOL	Department of Labor
DOT	Department of Transportation

APPENDIX C

DOT/UN/ NA/IMCO	Department of Transportation/United Nations/ North America/International Maritime Dangerous Goods Code
DWEL	drinking water exposure level
ECD	electron capture detection
ECG/EKG	electrocardiogram
EEG	electroencephalogram
EEGL	Emergency Exposure Guidance Level
EPA	Environmental Protection Agency
F	Fahrenheit
F ₁	first-filial generation
FAO	Food and Agricultural Organization of the United Nations
FDA	Food and Drug Administration
FEMA	Federal Emergency Management Agency
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FPD	flame photometric detection
fpm	feet per minute
FR	<i>Federal Register</i>
FSH	follicle stimulating hormone
g	gram
GC	gas chromatography
gd	gestational day
GLC	gas liquid chromatography
GPC	gel permeation chromatography
HPLC	high-performance liquid chromatography
HRGC	high resolution gas chromatography
HSDB	Hazardous Substance Data Bank
IARC	International Agency for Research on Cancer
IDLH	immediately dangerous to life and health
ILO	International Labor Organization
IRIS	Integrated Risk Information System
K _d	adsorption ratio
kg	kilogram
K _{oc}	organic carbon partition coefficient
K _{ow}	octanol-water partition coefficient
L	liter
LC	liquid chromatography
LC _{Lo}	lethal concentration, low
LC ₅₀	lethal concentration, 50% kill
LD _{Lo}	lethal dose, low
LD ₅₀	lethal dose, 50% kill
LDH	lactic dehydrogenase
LH	lutinizing hormone
LT ₅₀	lethal time, 50% kill
LOAEL	lowest-observed-adverse-effect level
LSE	Levels of Significant Exposure
m	meter
MA	<i>trans,trans</i> -muconic acid
MAL	maximum allowable level
mCi	millicurie
MCL	maximum contaminant level
MCLG	maximum contaminant level goal

APPENDIX C

MFO	mixed function oxidase
mg	milligram
mL	milliliter
mm	millimeter
mmHg	millimeters of mercury
mmol	millimole
mppcf	millions of particles per cubic foot
MRL	Minimal Risk Level
MS	mass spectrometry
NAAQS	National Ambient Air Quality Standard
NAS	National Academy of Science
NATICH	National Air Toxics Information Clearinghouse
NATO	North Atlantic Treaty Organization
NCE	normochromatic erythrocytes
NCEH	National Center for Environmental Health
NCI	National Cancer Institute
ND	not detected
NFPA	National Fire Protection Association
ng	nanogram
NIEHS	National Institute of Environmental Health Sciences
NIOSH	National Institute for Occupational Safety and Health
NIOSHTIC	NIOSH's Computerized Information Retrieval System
NLM	National Library of Medicine
nm	nanometer
NHANES	National Health and Nutrition Examination Survey
nmol	nanomole
NOAEL	no-observed-adverse-effect level
NOES	National Occupational Exposure Survey
NOHS	National Occupational Hazard Survey
NPD	nitrogen phosphorus detection
NPDES	National Pollutant Discharge Elimination System
NPL	National Priorities List
NR	not reported
NRC	National Research Council
NS	not specified
NSPS	New Source Performance Standards
NTIS	National Technical Information Service
NTP	National Toxicology Program
ODW	Office of Drinking Water, EPA
OERR	Office of Emergency and Remedial Response, EPA
OHM/TADS	Oil and Hazardous Materials/Technical Assistance Data System
OPP	Office of Pesticide Programs, EPA
OPPTS	Office of Prevention, Pesticides and Toxic Substances, EPA
OPPT	Office of Pollution Prevention and Toxics, EPA
OR	odds ratio
OSHA	Occupational Safety and Health Administration
OSW	Office of Solid Waste, EPA
OW	Office of Water
OWRS	Office of Water Regulations and Standards, EPA
PAH	polycyclic aromatic hydrocarbon
PBPD	physiologically based pharmacodynamic

APPENDIX C

PBPK	physiologically based pharmacokinetic
PCE	polychromatic erythrocytes
PEL	permissible exposure limit
PID	photo ionization detector
pg	picogram
pmol	picomole
PHS	Public Health Service
PMR	proportionate mortality ratio
ppb	parts per billion
ppm	parts per million
ppt	parts per trillion
PSNS	pretreatment standards for new sources
RBC	red blood cell
REL	recommended exposure level/limit
RfC	reference concentration
RfD	reference dose
RNA	ribonucleic acid
RTECS	Registry of Toxic Effects of Chemical Substances
RQ	reportable quantity
SARA	Superfund Amendments and Reauthorization Act
SCE	sister chromatid exchange
SGOT	serum glutamic oxaloacetic transaminase
SGPT	serum glutamic pyruvic transaminase
SIC	standard industrial classification
SIM	selected ion monitoring
SMCL	secondary maximum contaminant level
SMR	standardized mortality ratio
SNARL	suggested no adverse response level
SPEGL	Short-Term Public Emergency Guidance Level
STEL	short term exposure limit
STORET	Storage and Retrieval
TD ₅₀	toxic dose, 50% specific toxic effect
TLV	threshold limit value
TOC	total organic carbon
TPQ	threshold planning quantity
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
TWA	time-weighted average
UF	uncertainty factor
U.S.	United States
USDA	United States Department of Agriculture
USGS	United States Geological Survey
VOC	volatile organic compound
WBC	white blood cell
WHO	World Health Organization
>	greater than
≥	greater than or equal to
=	equal to
<	less than

APPENDIX C

#	less than or equal to
%	percent
α	alpha
β	beta
γ	gamma
δ	delta
μm	micrometer
μg	microgram
q_1^*	cancer slope factor
-	negative
+	positive
(+)	weakly positive result
(-)	weakly negative result

APPENDIX D

INDEX

acute inhalation exposure	40
adenocarcinoma	41, 49
adrenal gland	46, 98
adrenal glands	59
adsorption	121, 130, 131
aerobic	132
air	2-4, 8, 9, 11-14, 39, 42, 43, 48, 66, 67, 71, 74, 76, 79, 80, 87, 88, 90, 91, 93, 103, 109, 110, 124-126, 129, 131-133, 138-140, 142, 143, 150, 153, 154, 161, 165
ambient air	139
anaerobic	120, 132
aspartate aminotransferase	61
bioaccumulation	143
bioavailability	142
biomarker	89, 90, 102, 155
blood	7, 16, 42, 44-46, 48, 50, 57-59, 69, 71, 72, 74, 75, 79, 87, 89-92, 98, 102, 147, 148
body weight effects	47, 60, 63
breast cancer	62
breast milk	6, 139
cancer	5, 9, 20, 49, 62, 63, 68, 87, 98, 99, 108, 166
carcinogen	62, 91
carcinogenic	5, 9, 19, 20, 49, 62, 63, 68, 91, 99, 101, 160
carcinogenicity	99, 161, 164
carcinoma	41, 49, 68
cardiovascular effects	44, 45, 57, 65, 97
catfish	130
chronic inhalation exposure	79
Clean Water Act	162-164
deoxyribonucleic acid (see DNA)	70
Department of Health and Human Services (see DHHS)	5
dermal effects	13, 47, 59, 65
DHHS	5, 9
DNA	69, 70, 89
dog	72
endocrine effects	46, 59
FDA	8, 159, 163, 164, 166
FEDRIP	105, 144, 155, 156
fetus	6, 48, 86, 99
FEV1/FVC	16, 159
fish	130
Food and Drug Administration (see FDA)	8, 166
friable	50, 57, 66, 85
gastrointestinal effects	45, 57, 65
general population	11, 89, 121, 138, 141
groundwater	88, 125, 131, 132, 136, 137
half-life	11, 12, 89, 120, 121, 131, 132
hematological effects	45, 58
hepatic effects	46, 59
hydrolysis	128, 149
hydroxyl radical	131
immune system	47, 67, 100
immunological effects	60, 67
Integrated Risk Information System	166
kidney	12, 22, 46, 59, 74, 75, 92, 95, 104, 147
LD50	47

APPENDIX D

liver	12, 14, 22, 46, 71-76, 78, 79, 82, 88, 89, 92-95, 102, 147
lung	5, 13, 16, 23, 40, 41, 43, 50, 62, 65, 88, 89, 91
lymph	44
lymphoreticular effects	47, 60, 67
milk	6, 139, 144, 145
Minimal Risk Levels (see MRL)	14, 20
MRL	14-17, 20, 22, 23, 60, 97, 98, 159
MRLs	14, 17, 20, 21, 23
Muscarinic receptors	83
musculoskeletal effects	45, 58
National Priorities List (see NPL)	1, 117
neoplastic	41
neurobehavioral	86
neurochemical	100
neurotransmitter	88, 100
NIOSH	8, 9, 108, 138, 147, 150, 155, 161, 166
NOAEL	16, 17, 19, 60, 159, 160
NOAELs	19, 20, 44, 45, 59, 60
NOES	138
NPL	1, 12, 117, 118, 121, 125, 137
ocean	130, 133, 137
ocular effects	14, 47, 59, 63, 66, 67, 97
particulate	39, 88, 131, 132, 153, 154
particulates	133, 150
partition coefficients	109
PBPD	80
PBPK	80-83
pharmacodynamic	80
pharmacokinetic	80-83, 90
physiologically based pharmacodynamic (see PBPD)	80
physiologically based pharmacokinetic (see PBPK)	80, 82
precipitation	137
public health	1, 5, 8, 11, 19, 20, 85, 94, 95, 141, 147, 149, 154
reference dose (see RfD)	166
reflexes	14, 48
regulations	8, 159, 161, 162, 165, 166
renal effects	12, 46, 59, 65, 97, 98
reportable quantity	163-165
RfD	164, 166
sea	133
sediment	11, 125, 130, 132, 137
selenium	89
serum	43, 58, 69, 147, 148
soil	2-4, 8, 11, 12, 111, 117, 121, 125, 126, 128-133, 137, 142, 143, 152, 153
solubility	13, 65, 83, 94, 107, 109
Superfund	165
surface water	125, 128, 130, 132, 137
time-weighted average (see TWA)	16, 166
toxicokinetic	19, 103
Toxics Release Inventory (see TRI)	121
transport	4, 14, 61, 83, 86, 104, 117, 127, 129, 130, 136, 144
TRI	121, 124, 125, 142
tumors	62, 63, 91, 99
TWA	16, 161, 162, 166
U.S. Department of Agriculture (see USDA)	144, 145
USDA	144, 145
vapor pressure	109
volatility	107
volatilization	120, 127, 130-133, 135, 139, 145, 153, 154
water	2-5, 8, 11-13, 17, 49, 50, 57-62, 65, 79, 83, 84, 87-89, 91, 93, 94, 98, 107, 109, 110, 115-117, 120-125, 127-130, 132, 133, 135-137, 142, 143, 149, 151, 153, 154, 159, 161-166