

Table 1.108A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	22,870	22,995	2,084	2,075	635	570
GEOGRAPHIC DIVISION						
Northeast	4,476	4,675	379	393	98	104
New England	1,330	1,294	151	112	41	26
Middle Atlantic	3,147	3,381	229	280	56	78
Midwest	5,061	5,377	507	484	150	134
East North Central	3,735	3,808	349	339	114	81
West North Central	1,326	1,570	158	145	36	53
South	7,318	7,067	685	646	203	179
South Atlantic	4,035	3,908	393	386	137	100
East South Central	1,103	1,023	108	78	14	17
West South Central	2,179	2,136	185	181	52	62
West	6,015	5,876	512	553	184	153
Mountain	1,828	1,831	156	172	44	46
Pacific	4,187	4,045	356	380	140	107
COUNTY TYPE¹						
Large Metro	12,694	12,931	1,109	1,100	341	300
Small Metro	6,941	6,750	645	642	192	192
250K – 1 Mil. Pop.	4,425	4,555	422	424	146	130
< 250K Pop.	2,516	2,196	223	218	46	61
Nonmetro	3,236	3,314	329	334	102	78
Urbanized	1,653	1,704	166	149	53	42
Less Urbanized	1,277	1,375	120	160	39	25
Completely Rural	305	235	43	25	9	12

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.108B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	9.7	9.7	0.9	0.9	0.3	0.2
GEOGRAPHIC DIVISION						
Northeast	9.9	10.3	0.8	0.9	0.2	0.2
New England	11.3	10.9	1.3	0.9	0.3	0.2
Middle Atlantic	9.5	10.1	0.7	0.8	0.2	0.2
Midwest	9.5	10.0	0.9	0.9	0.3	0.2
East North Central	10.0	10.1	0.9	0.9	0.3	0.2
West North Central	8.3	9.7	1.0	0.9	0.2	0.3
South	8.7	8.3	0.8	0.8	0.2	0.2
South Atlantic	9.2	8.8	0.9	0.9	0.3	0.2
East South Central	7.8	7.2	0.8	0.5	0.1	0.1
West South Central	8.4	8.2	0.7	0.7	0.2	0.2
West	11.4	10.9	1.0	1.0	0.3	0.3
Mountain	11.9	11.7	1.0	1.1	0.3	0.3
Pacific	11.1	10.6	0.9	1.0	0.4	0.3
COUNTY TYPE¹						
Large Metro	10.1	10.3	0.9	0.9	0.3	0.2
Small Metro	10.0	9.5	0.9	0.9	0.3	0.3
250K – 1 Mil. Pop.	9.7	9.7	0.9	0.9	0.3	0.3
< 250K Pop.	10.6	9.0	0.9	0.9	0.2	0.3
Nonmetro	8.1	8.2	0.8	0.8	0.3	0.2
Urbanized	10.3	9.5	1.0	0.8	0.3	0.2
Less Urbanized	6.5	7.5	0.6 ^a	0.9	0.2	0.1
Completely Rural	6.9	5.7	1.0	0.6	0.2	0.3

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.109A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	2,605	2,670	1,078	1,119	301	320
GEOGRAPHIC DIVISION						
Northeast	441	429	178	191	47	59
New England	124	118	61	54	18	11
Middle Atlantic	317	311	118	137	30	48
Midwest	597	621	231	258	52	63
East North Central	412	453	163	185	42	43
West North Central	185	167	68	73	10	20
South	926	889	400	370	125	108
South Atlantic	514	448	232	201	82 ^a	44
East South Central	143	140	53	40	14	13
West South Central	268	301	116	129	29	52
West	641	732	269	301	77	89
Mountain	213	219	83	88	23	26
Pacific	428	513	185	213	54	63
COUNTY TYPE¹						
Large Metro	1,344	1,348	550	567	142	153
Small Metro	773	833	325	343	104	115
250K – 1 Mil. Pop.	508	547	206	221	71	76
< 250K Pop.	265	285	119	122	34	39
Nonmetro	488	489	203	209	55	52
Urbanized	190	223	77	92	18	22
Less Urbanized	237	204	95	97	30	22
Completely Rural	62	62	31	20	6	7

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.109B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	10.5	10.7	4.4	4.5	1.2	1.3
GEOGRAPHIC DIVISION						
Northeast	9.9	9.6	4.0	4.3	1.1	1.3
New England	10.7	10.0	5.2	4.6	1.5	0.9
Middle Atlantic	9.6	9.4	3.6	4.1	0.9	1.4
Midwest	10.5	10.9	4.1	4.5	0.9	1.1
East North Central	10.3	11.3	4.1	4.6	1.0	1.1
West North Central	10.8	9.8	4.0	4.3	0.6	1.2
South	10.6	10.0	4.6	4.1	1.4	1.2
South Atlantic	11.7	9.9	5.3	4.5	1.9 ^b	1.0
East South Central	10.1	9.6	3.7	2.8	1.0	0.9
West South Central	9.1	10.1	3.9	4.3	1.0	1.7
West	11.0	12.4	4.6	5.1	1.3	1.5
Mountain	12.5	12.7	4.9	5.1	1.3	1.5
Pacific	10.4	12.4	4.5	5.1	1.3	1.5
COUNTY TYPE¹						
Large Metro	10.2	10.1	4.2	4.3	1.1	1.1
Small Metro	10.7	11.3	4.5	4.6	1.4	1.6
250K – 1 Mil. Pop.	10.4	11.0	4.2	4.4	1.4	1.5
< 250K Pop.	11.3	11.8	5.1	5.0	1.4	1.6
Nonmetro	11.3	11.4	4.7	4.9	1.3	1.2
Urbanized	11.0	12.0	4.5	5.0	1.1	1.2
Less Urbanized	11.3	10.3	4.5	4.9	1.4	1.1
Completely Rural	13.0	14.0	6.6	4.6	1.3	1.6

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.110A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	4,867	4,729	685	662	150	134
GEOGRAPHIC DIVISION						
Northeast	945	919	152	156	20	33
New England	296	249	60	39	5	8
Middle Atlantic	649	670	92	117	15	25
Midwest	1,229	1,142	174	162	33	25
East North Central	900	816	121	119	19	20
West North Central	329	325	53	43	14	5
South	1,485	1,450	213	164	53	39
South Atlantic	804	768	132	98	38	30
East South Central	231	229	26	14	*	*
West South Central	450	452	55	53	15	10
West	1,208	1,219	147	180	44	36
Mountain	404	417	48	57	10	7
Pacific	803	801	99	123	34	29
COUNTY TYPE¹						
Large Metro	2,439	2,375	383	341	96	66
Small Metro	1,717	1,603	208	215	32	46
250K – 1 Mil. Pop.	1,062	1,082	124	142	20	33
< 250K Pop.	655 ^a	522	84	73	12	13
Nonmetro	712	750	94	105	22	22
Urbanized	362	400	57	53	11	20
Less Urbanized	295	315	24	51	8	2
Completely Rural	54	34	12 ^a	1	3	0

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.110B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	15.7	14.9	2.2	2.1	0.5	0.4
GEOGRAPHIC DIVISION						
Northeast	17.2	16.3	2.8	2.8	0.4	0.6
New England	20.6	16.9	4.2	2.7	0.3	0.5
Middle Atlantic	15.9	16.1	2.3	2.8	0.4	0.6
Midwest	17.1	15.5	2.4	2.2	0.5	0.3
East North Central	18.1 ^a	16.1	2.4	2.3	0.4	0.4
West North Central	14.8	14.3	2.4	1.9	0.6	0.2
South	13.5	12.8	1.9	1.4	0.5	0.3
South Atlantic	14.9	13.8	2.4	1.8	0.7	0.5
East South Central	12.1	11.9	1.3	0.7	*	*
West South Central	12.1	11.9	1.5	1.4	0.4	0.3
West	16.6	16.4	2.0	2.4	0.6	0.5
Mountain	18.5	18.7	2.2	2.5	0.5	0.3
Pacific	15.7	15.5	1.9	2.4	0.7	0.6
COUNTY TYPE¹						
Large Metro	15.4	14.3	2.4	2.1	0.6	0.4
Small Metro	16.9	15.7	2.0	2.1	0.3	0.4
250K – 1 Mil. Pop.	16.5	16.3	1.9	2.1	0.3	0.5
< 250K Pop.	17.6 ^a	14.7	2.3	2.1	0.3	0.4
Nonmetro	14.3	15.3	1.9	2.1	0.4	0.5
Urbanized	16.1	16.9	2.6	2.2	0.5	0.8
Less Urbanized	13.0	14.4	1.1 ^a	2.3	0.4	0.1
Completely Rural	11.7	9.5	2.6 ^a	0.2	0.6	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.111A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	15,398	15,596	320	295	183	117
GEOGRAPHIC DIVISION						
Northeast	3,090	3,328	49	45	30	12
New England	910	928	30	19	18	7
Middle Atlantic	2,180	2,399	19	27	12	5
Midwest	3,235	3,615	103	65	66	46
East North Central	2,422	2,538	66	35	53	18
West North Central	812 ^a	1,077	37	30	12	27
South	4,906	4,728	72	112	25	31
South Atlantic	2,717	2,692	29	87	17	27
East South Central	729	654	29	24	*	4
West South Central	1,461	1,382	14	*	8	*
West	4,167	3,925	96	73	63	28
Mountain	1,211	1,195	24	28	11	13
Pacific	2,956	2,730	72	45	52	15
COUNTY TYPE¹						
Large Metro	8,911	9,208	176	191	103	82
Small Metro	4,451	4,314	112	84	55	31
250K – 1 Mil. Pop.	2,855	2,926	92	61	55	21
< 250K Pop.	1,596	1,388	20	22	*	10
Nonmetro	2,036	2,075	32	20	25	4
Urbanized	1,102	1,081	31	4	24	*
Less Urbanized	745	855	1	12	1	*
Completely Rural	189	139	*	4	*	4

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.111B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	8.6	8.6	0.2	0.2	0.1	0.1
GEOGRAPHIC DIVISION						
Northeast	8.8	9.5	0.1	0.1	0.1	0.0
New England	9.9	10.0	0.3	0.2	0.2	0.1
Middle Atlantic	8.4	9.2	0.1	0.1	0.0	0.0
Midwest	8.0	8.9	0.3	0.2	0.2	0.1
East North Central	8.5	8.9	0.2	0.1	0.2	0.1
West North Central	6.7 ^a	8.9	0.3	0.2	0.1	0.2
South	7.7	7.3	0.1	0.2	0.0	0.0
South Atlantic	8.0	7.8	0.1	0.3	0.1	0.1
East South Central	6.7	6.0	0.3	0.2	*	0.0
West South Central	7.6	7.1	0.1	*	0.0	*
West	10.5	9.7	0.2	0.2	0.2	0.1
Mountain	10.6	10.3	0.2	0.2	0.1	0.1
Pacific	10.4	9.5	0.3	0.2	0.2	0.1
COUNTY TYPE¹						
Large Metro	9.2	9.6	0.2	0.2	0.1	0.1
Small Metro	8.6	8.0	0.2	0.2	0.1	0.1
250K – 1 Mil. Pop.	8.4	8.3	0.3	0.2	0.2	0.1
< 250K Pop.	9.0	7.5	0.1	0.1	*	0.1
Nonmetro	6.6	6.7	0.1	0.1	0.1	0.0
Urbanized	9.1	7.9	0.3	0.0	0.2	*
Less Urbanized	4.9	6.1	0.0	0.1	0.0	*
Completely Rural	5.4	4.2	*	0.1	*	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.112A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	20,265	20,325	1,005	956	334	250
GEOGRAPHIC DIVISION						
Northeast	4,035	4,246	201	202	50	45
New England	1,206	1,177	90	58	23	15
Middle Atlantic	2,830	3,070	111	144	27	30
Midwest	4,464	4,757	277	227	98	71
East North Central	3,322	3,355	187	154	72	38
West North Central	1,141 ^a	1,402	90	73	26	33
South	6,392	6,178	285	276	78	70
South Atlantic	3,521	3,460	161	185	55	56
East South Central	960	883	55	38	*	4
West South Central	1,911	1,835	69	53	23	10
West	5,374	5,144	243	252	107	64
Mountain	1,615	1,612	72	85	21	20
Pacific	3,759	3,531	170	168	86	43
COUNTY TYPE¹						
Large Metro	11,350	11,583	559	533	199	147
Small Metro	6,168	5,917	320	299	88	76
250K – 1 Mil. Pop.	3,917	4,007	216	203	76	54
< 250K Pop.	2,251	1,910	104	96	12	22
Nonmetro	2,747	2,825	126	125	47	27
Urbanized	1,464	1,481	89	57	35	20
Less Urbanized	1,040	1,170	25 ^a	63	9	2
Completely Rural	243	173	12	5	3	4

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.112B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	9.6	9.6	0.5	0.4	0.2	0.1
GEOGRAPHIC DIVISION						
Northeast	9.9	10.4	0.5	0.5	0.1	0.1
New England	11.3	11.0	0.8	0.5	0.2	0.1
Middle Atlantic	9.4	10.2	0.4	0.5	0.1	0.1
Midwest	9.4	9.9	0.6	0.5	0.2	0.1
East North Central	9.9	10.0	0.6	0.5	0.2	0.1
West North Central	8.0	9.7	0.6	0.5	0.2	0.2
South	8.5	8.1	0.4	0.4	0.1	0.1
South Atlantic	8.9	8.7	0.4	0.5	0.1	0.1
East South Central	7.5	6.9	0.4	0.3	*	0.0
West South Central	8.3	7.9	0.3	0.2	0.1	0.0
West	11.4	10.7	0.5	0.5	0.2	0.1
Mountain	11.9	11.6	0.5	0.6	0.2	0.1
Pacific	11.2	10.4	0.5	0.5	0.3	0.1
COUNTY TYPE¹						
Large Metro	10.1	10.3	0.5	0.5	0.2	0.1
Small Metro	9.9	9.2	0.5	0.5	0.1	0.1
250K – 1 Mil. Pop.	9.6	9.6	0.5	0.5	0.2	0.1
< 250K Pop.	10.5	8.7	0.5	0.4	0.1	0.1
Nonmetro	7.7	7.9	0.4	0.3	0.1	0.1
Urbanized	10.2	9.3	0.6	0.4	0.2	0.1
Less Urbanized	5.9	7.2	0.1 ^a	0.4	0.1	0.0
Completely Rural	6.1	4.7	0.3	0.1	0.1	0.1

*Low precision; no estimate reported.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.