

**Table 1.19A Any Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	108,255	110,205	35,132	34,993	19,522	19,470
12	509	503	290	255	136	112
13	752	811	469	516	212	211
14	1,165	1,086	824	772	399	349
15	1,450	1,414	1,074	1,043	548	551
16	1,769	1,835	1,318	1,391	710	780
17	1,997	1,981	1,519	1,478	874	807
18	2,379	2,464	1,758	1,761	974	1,042
19	2,350	2,265	1,620	1,551	913	914
20	2,388	2,497	1,533	1,623	910	967
21	2,705 <sup>a</sup>	2,415	1,593 <sup>a</sup>	1,372	877	812
22	2,335 <sup>a</sup>	2,570	1,304	1,347	751	786
23	2,358	2,538	1,276	1,287	741	719
24	2,103	2,265	1,028	1,095	600	626
25	1,944 <sup>a</sup>	2,181	888	939	499	569
26-29	8,769	8,555	3,421	3,483	1,958	1,982
30-34	11,738	11,463	3,429	3,345	1,742	1,771
35-39	13,426	12,903	3,414	3,141	1,890	1,757
40-44	14,573	14,821	3,166	3,175	1,719	1,831
45-49	12,908	13,349	2,600	2,691	1,592	1,463
50-54	9,604	10,389	1,218	1,474	635	784
55-59	5,139	5,716	483	663	272	298
60-64	2,754	2,798	474	336	293	127
65 or Older	3,139	3,387	432	254	280	209

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.19B Any Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	46.0	46.4	14.9	14.7	8.3	8.2
12	12.4	12.2	7.1	6.2	3.3	2.7
13	18.0	18.7	11.2	11.9	5.1	4.9
14	26.7	26.3	18.9	18.7	9.1	8.5
15	35.3	34.2	26.1	25.2	13.3	13.3
16	43.7	43.8	32.6	33.2	17.5	18.6
17	50.6	48.4	38.5	36.1	22.1	19.7
18	54.5	53.5	40.3	38.2	22.3	22.6
19	57.5	58.3	39.6	39.9	22.3	23.5
20	59.7	62.0	38.4	40.3	22.8	24.0
21	66.8 <sup>b</sup>	61.6	39.4 <sup>b</sup>	35.0	21.7	20.7
22	60.4 <sup>a</sup>	64.0	33.7	33.5	19.4	19.6
23	61.9	63.4	33.5	32.2	19.4	18.0
24	59.7	62.3	29.2	30.1	17.0	17.2
25	58.4	60.1	26.7	25.9	15.0	15.7
26-29	57.4	57.9	22.4	23.6	12.8	13.4
30-34	59.0	56.8	17.2	16.6	8.8	8.8
35-39	61.0	61.7	15.5	15.0	8.6	8.4
40-44	66.4	65.3	14.4	14.0	7.8	8.1
45-49	60.9	62.3	12.3	12.6	7.5	6.8
50-54	51.1	52.0	6.5	7.4	3.4	3.9
55-59	35.0	38.3	3.3	4.4	1.9	2.0
60-64	23.7	23.8	4.1	2.9	2.5	1.1
65 or Older	9.2	9.9	1.3	0.7	0.8	0.6

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.20A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	94,946	96,611	25,755	25,231	14,584	14,638
12	86	59	58	38	24	21
13	271	225	195	159	95 <sup>a</sup>	62
14	641	578	498	459	239	218
15	1,023	947	835	759	409	378
16	1,381	1,434	1,057	1,114	534 <sup>a</sup>	638
17	1,703	1,645	1,262	1,216	722	654
18	2,046	2,109	1,462	1,459	836	897
19	2,086	1,953	1,379	1,290	807	795
20	2,156	2,255	1,311	1,385	791	799
21	2,511 <sup>b</sup>	2,200	1,360 <sup>a</sup>	1,168	743	699
22	2,074 <sup>a</sup>	2,342	1,103	1,135	644	668
23	2,146	2,276	1,055	1,001	614	574
24	1,918	2,015	839	857	515	506
25	1,744 <sup>a</sup>	1,954	728	749	426	462
26-29	7,803	7,547	2,531	2,556	1,477	1,543
30-34	10,537	10,287	2,466	2,488	1,245	1,396
35-39	12,164	11,840	2,260	2,113	1,315	1,256
40-44	13,496	13,751	2,109	2,227	1,161	1,357
45-49	11,940	12,349	1,759	1,609	1,135	884
50-54	8,659	9,488	812	910	464	489
55-59	4,380	4,896	237	418	108 <sup>a</sup>	258
60-64	2,230	2,225	274 <sup>a</sup>	74	173	51
65 or Older	1,952	2,236	165	48	108	33

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.20B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	40.4	40.6	11.0	10.6	6.2	6.2
12	2.1	1.4	1.4	0.9	0.6	0.5
13	6.5	5.2	4.6	3.7	2.3 <sup>a</sup>	1.4
14	14.7	14.0	11.4	11.1	5.5	5.3
15	24.9	22.9	20.3	18.4	9.9	9.2
16	34.1	34.2	26.1	26.6	13.2 <sup>a</sup>	15.2
17	43.1 <sup>a</sup>	40.2	31.9	29.7	18.3 <sup>a</sup>	16.0
18	46.9	45.8	33.5	31.7	19.2	19.5
19	51.0	50.2	33.7	33.2	19.7	20.5
20	53.9	56.0	32.8	34.4	19.8	19.8
21	62.0 <sup>b</sup>	56.1	33.6 <sup>a</sup>	29.8	18.4	17.8
22	53.6 <sup>b</sup>	58.3	28.5	28.3	16.7	16.6
23	56.3	56.8	27.7	25.0	16.1	14.3
24	54.4	55.4	23.8	23.6	14.6	13.9
25	52.4	53.8	21.9	20.6	12.8	12.7
26-29	51.1	51.1	16.6	17.3	9.7	10.4
30-34	53.0	50.9	12.4	12.3	6.3	6.9
35-39	55.3	56.6	10.3	10.1	6.0	6.0
40-44	61.5	60.6	9.6	9.8	5.3	6.0
45-49	56.4	57.7	8.3	7.5	5.4	4.1
50-54	46.1	47.5	4.3	4.5	2.5	2.4
55-59	29.8	32.8	1.6	2.8	0.7 <sup>a</sup>	1.7
60-64	19.2	18.9	2.4 <sup>a</sup>	0.6	1.5	0.4
65 or Older	5.7	6.5	0.5	0.1	0.3	0.1

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.21A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	33,910	34,891	5,902	5,908	2,020	2,281
12	10	8	8	7	3	2
13	15	22	8	9	1 <sup>a</sup>	7
14	48	62	43	41	15	14
15	101	99	80	74	27	28
16	187	187	136	129	45	42
17	309	270	233 <sup>a</sup>	178	56	60
18	355	445	233	285	68 <sup>a</sup>	115
19	497	468	280	283	89	92
20	616	567	308	278	86	67
21	752 <sup>a</sup>	610	323	271	81	81
22	671	678	280	273	92	104
23	706	728	278	272	109	82
24	646	660	209	233	58	78
25	544	598	175	203	48	77
26-29	2,330	2,590	640	769	184	263
30-34	3,859	3,722	639	650	245	256
35-39	5,975	5,454	666	627	253	323
40-44	6,528	6,781	525	591	220	264
45-49	5,329	5,440	486	419	248	144
50-54	2,670 <sup>b</sup>	3,614	271	255	68	167
55-59	959	1,208	65	47	24	*
60-64	490	410	14	*	*	*
65 or Older	315	269	*	14	*	14

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.21B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	14.4	14.7	2.5	2.5	0.9	1.0
12	0.2	0.2	0.2	0.2	0.1	0.1
13	0.4	0.5	0.2	0.2	0.0 <sup>a</sup>	0.2
14	1.1	1.5	1.0	1.0	0.3	0.3
15	2.4	2.4	2.0	1.8	0.7	0.7
16	4.6	4.5	3.4	3.1	1.1	1.0
17	7.8	6.6	5.9 <sup>a</sup>	4.3	1.4	1.5
18	8.1	9.7	5.4	6.2	1.6 <sup>a</sup>	2.5
19	12.2	12.0	6.8	7.3	2.2	2.4
20	15.4	14.1	7.7	6.9	2.2	1.7
21	18.6 <sup>a</sup>	15.5	8.0	6.9	2.0	2.1
22	17.4	16.9	7.2	6.8	2.4	2.6
23	18.5	18.2	7.3	6.8	2.9	2.1
24	18.3	18.2	5.9	6.4	1.6	2.1
25	16.3	16.5	5.3	5.6	1.4	2.1
26-29	15.3	17.5	4.2	5.2	1.2	1.8
30-34	19.4	18.4	3.2	3.2	1.2	1.3
35-39	27.2	26.1	3.0	3.0	1.2	1.5
40-44	29.8	29.9	2.4	2.6	1.0	1.2
45-49	25.2	25.4	2.3	2.0	1.2	0.7
50-54	14.2 <sup>a</sup>	18.1	1.4	1.3	0.4	0.8
55-59	6.5	8.1	0.4	0.3	0.2	*
60-64	4.2	3.5	0.1	*	*	*
65 or Older	0.9	0.8	*	0.0	*	0.0

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.22A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	8,402	7,949	1,554	1,406	567	604
12	3	*	2	*	2	*
13	7	2	4	2	*	2
14	13	18	10	11	9	2
15	25	28	17	18	3	7
16	45	53	25	33	5	14
17	72	57	42	31	7	7
18	70	91	35	40	5	18
19	127	114	42	44	10	7
20	130	148	27	40	2	8
21	158	128	46	28	10	6
22	151	170	31	29	5	1
23	198	187	33	29	7	6
24	193	184	27	37	10	5
25	155	171	26	29	4	4
26-29	723	776	137	151	19	60
30-34	1,168	1,152	230	165	55	82
35-39	1,583	1,547	205	200	127	113
40-44	1,601	1,515	213	224	119	122
45-49	1,203	990	220	172	134	50
50-54	520	457	147	113	12	90
55-59	159	124	21	11	21	*
60-64	95 <sup>a</sup>	5	14	*	*	*
65 or Older	4	32	*	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.22B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	3.6	3.3	0.7	0.6	0.2	0.3
12	0.1	*	0.1	*	0.0	*
13	0.2	0.0	0.1	0.0	*	0.0
14	0.3	0.4	0.2	0.3	0.2	0.1
15	0.6	0.7	0.4	0.4	0.1	0.2
16	1.1	1.3	0.6	0.8	0.1	0.3
17	1.8	1.4	1.1	0.8	0.2	0.2
18	1.6	2.0	0.8	0.9	0.1	0.4
19	3.1	2.9	1.0	1.1	0.2	0.2
20	3.3	3.7	0.7	1.0	0.0	0.2
21	3.9	3.3	1.1	0.7	0.3	0.1
22	3.9	4.2	0.8	0.7	0.1	0.0
23	5.2	4.7	0.9	0.7	0.2	0.2
24	5.5	5.1	0.8	1.0	0.3	0.1
25	4.6	4.7	0.8	0.8	0.1	0.1
26-29	4.7	5.3	0.9	1.0	0.1	0.4
30-34	5.9	5.7	1.2	0.8	0.3	0.4
35-39	7.2	7.4	0.9	1.0	0.6	0.5
40-44	7.3	6.7	1.0	1.0	0.5	0.5
45-49	5.7	4.6	1.0	0.8	0.6 <sup>a</sup>	0.2
50-54	2.8	2.3	0.8	0.6	0.1	0.4
55-59	1.1	0.8	0.1	0.1	0.1	*
60-64	0.8 <sup>a</sup>	0.0	0.1	*	*	*
65 or Older	0.0	0.1	*	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.23A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	34,314	34,363	4,749 <sup>b</sup>	3,936	1,196	1,042
12	35	44	21	21	14	7
13	73	63	54	41	21	12
14	148	140	109	101	32	39
15	240	217	164	127	36	47
16	387	321	257	207	69	70
17	537	465	340	282	82	84
18	783	667	439	366	116	108
19	886	780	423 <sup>a</sup>	319	101	87
20	976	963	421	414	91	95
21	1,148 <sup>b</sup>	949	392 <sup>b</sup>	279	82	55
22	991	1,062	282	220	67	68
23	1,035	1,050	270	207	54	55
24	911	998	221	183	48	46
25	787	910	162	123	29	30
26-29	3,255	3,406	409	475	112	125
30-34	4,004	3,679	319	273	78	40
35-39	4,302	4,305	173	102	51	41
40-44	5,098	5,300	136	69	32	33
45-49	4,748	4,486	80	56	27	*
50-54	2,612	3,158	59	70	53	*
55-59	717	1,006	17	*	*	*
60-64	268	287	*	*	*	*
65 or Older	373 <sup>a</sup>	109	*	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.23B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	14.6	14.5	2.0 <sup>b</sup>	1.7	0.5	0.4
12	0.8	1.1	0.5	0.5	0.3	0.2
13	1.8	1.5	1.3	0.9	0.5	0.3
14	3.4	3.4	2.5	2.4	0.7	0.9
15	5.8	5.2	4.0	3.1	0.9	1.1
16	9.6 <sup>a</sup>	7.7	6.3 <sup>a</sup>	4.9	1.7	1.7
17	13.6 <sup>a</sup>	11.4	8.6 <sup>a</sup>	6.9	2.1	2.1
18	18.0 <sup>b</sup>	14.5	10.1 <sup>a</sup>	7.9	2.7	2.3
19	21.7	20.1	10.3 <sup>a</sup>	8.2	2.5	2.2
20	24.4	23.9	10.5	10.3	2.3	2.4
21	28.4 <sup>b</sup>	24.2	9.7 <sup>b</sup>	7.1	2.0	1.4
22	25.6	26.4	7.3 <sup>a</sup>	5.5	1.7	1.7
23	27.2	26.2	7.1 <sup>a</sup>	5.2	1.4	1.4
24	25.8	27.5	6.3	5.0	1.4	1.3
25	23.6	25.1	4.9 <sup>a</sup>	3.4	0.9	0.8
26-29	21.3	23.1	2.7	3.2	0.7	0.8
30-34	20.1	18.2	1.6	1.4	0.4	0.2
35-39	19.6	20.6	0.8	0.5	0.2	0.2
40-44	23.2	23.4	0.6	0.3	0.1	0.1
45-49	22.4	20.9	0.4	0.3	0.1	*
50-54	13.9	15.8	0.3	0.4	0.3	*
55-59	4.9	6.7	0.1	*	*	*
60-64	2.3	2.4	*	*	*	*
65 or Older	1.1 <sup>a</sup>	0.3	*	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.24A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	22,870	22,995	2,084	2,075	635	570
12	266	289	137	107	58	44
13	419	461	185	231	56	76
14	498	473	224	235	72	57
15	498	489	192	193	65	60
16	458	485	163	186	27	48
17	466	473	176	166	23	35
18	504	577	149	171	22	35
19	596	524	145	109	24	16
20	620	632	102	100	28	28
21	627	584	75	60	26 <sup>b</sup>	4
22	656	571	63	63	17	12
23	709	640	76	79	15	17
24	602	609	48	50	12	14
25	553	592	26	31	6	7
26-29	2,145	2,052	95	65	38	4
30-34	2,800	2,707	34	25	5	8
35-39	3,018	2,677	25	34	18	18
40-44	2,985	2,988	60	73	51	54
45-49	2,379	2,619	63	69	36	23
50-54	1,052 <sup>a</sup>	1,606	*	12	*	9
55-59	430	522	*	*	*	*
60-64	275	189	*	*	*	*
65 or Older	312	238	43	17	35	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.24B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	9.7	9.7	0.9	0.9	0.3	0.2
12	6.5	7.0	3.3	2.6	1.4	1.1
13	10.0	10.6	4.4	5.3	1.3	1.8
14	11.4	11.5	5.1	5.7	1.6	1.4
15	12.1	11.8	4.7	4.7	1.6	1.4
16	11.3	11.6	4.0	4.4	0.7	1.1
17	11.8	11.6	4.5	4.1	0.6	0.9
18	11.6	12.5	3.4	3.7	0.5	0.8
19	14.6	13.5	3.5	2.8	0.6	0.4
20	15.5	15.7	2.6	2.5	0.7	0.7
21	15.5	14.9	1.9	1.5	0.7 <sup>b</sup>	0.1
22	17.0 <sup>a</sup>	14.2	1.6	1.6	0.4	0.3
23	18.6 <sup>a</sup>	16.0	2.0	2.0	0.4	0.4
24	17.1	16.8	1.4	1.4	0.3	0.4
25	16.6	16.3	0.8	0.8	0.2	0.2
26-29	14.0	13.9	0.6	0.4	0.3	0.0
30-34	14.1	13.4	0.2	0.1	0.0	0.0
35-39	13.7	12.8	0.1	0.2	0.1	0.1
40-44	13.6	13.2	0.3	0.3	0.2	0.2
45-49	11.2	12.2	0.3	0.3	0.2	0.1
50-54	5.6 <sup>a</sup>	8.0	*	0.1	*	0.0
55-59	2.9	3.5	*	*	*	*
60-64	2.4	1.6	*	*	*	*
65 or Older	0.9	0.7	0.1	0.1	0.1	*

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.25A Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	46,558	47,882	14,680	14,986	6,210	6,336
12	231	237	118	132	58	60
13	285	330	184	211	83	90
14	483	452	338	310	165	134
15	643	612	433	442	175	205
16	820	808	557	555	249	240
17	926	913	653	640	250	268
18	1,062	1,180	655	706	253	285
19	1,137	1,161	692	675	259	266
20	1,087	1,181	599	701	238 <sup>a</sup>	317
21	1,214	1,153	605	554	227	233
22	1,129	1,176	576	489	234	215
23	1,084 <sup>a</sup>	1,245	492	566	195	225
24	975	1,097	439	516	143	198
25	898	1,017	352	392	133	174
26-29	3,749	3,824	1,557	1,491	667	597
30-34	4,827	4,795	1,350	1,454	607	595
35-39	5,531	5,518	1,559	1,377	656	544
40-44	6,370	6,591	1,349	1,187	564	508
45-49	5,736	5,873	951	1,288	438	609
50-54	3,620	4,320	474	616	186	280
55-59	2,084	2,120	264	216	175 <sup>a</sup>	40
60-64	1,130	1,056	249	263	120	76
65 or Older	1,536	1,221	232	206	136	176

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.25B Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	19.8	20.1	6.2	6.3	2.6	2.7
12	5.6	5.7	2.9	3.2	1.4	1.5
13	6.8	7.6	4.4	4.9	2.0	2.1
14	11.1	11.0	7.8	7.5	3.8	3.3
15	15.6	14.8	10.5	10.7	4.2	5.0
16	20.2	19.3	13.7	13.2	6.1	5.7
17	23.5	22.3	16.5	15.6	6.3	6.5
18	24.3	25.6	15.0	15.3	5.8	6.2
19	27.8	29.9	16.9	17.4	6.3	6.9
20	27.2	29.3	15.0	17.4	6.0 <sup>a</sup>	7.9
21	30.0	29.4	14.9	14.1	5.6	5.9
22	29.2	29.3	14.9 <sup>a</sup>	12.2	6.0	5.3
23	28.5	31.1	12.9	14.1	5.1	5.6
24	27.7	30.2	12.5	14.2	4.1	5.4
25	27.0	28.0	10.6	10.8	4.0	4.8
26-29	24.6	25.9	10.2	10.1	4.4	4.0
30-34	24.3	23.7	6.8	7.2	3.0	2.9
35-39	25.1	26.4	7.1	6.6	3.0	2.6
40-44	29.0	29.0	6.1	5.2	2.6	2.2
45-49	27.1	27.4	4.5	6.0	2.1	2.8
50-54	19.3	21.6	2.5	3.1	1.0	1.4
55-59	14.2	14.2	1.8	1.4	1.2 <sup>a</sup>	0.3
60-64	9.7	9.0	2.1	2.2	1.0	0.6
65 or Older	4.5	3.6	0.7	0.6	0.4	0.5

\*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.26A Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	29,611 <sup>a</sup>	31,207	10,992	11,671	4,377	4,693
12	175	199	94	110	48	48
13	222 <sup>a</sup>	286	151	186	76	76
14	396	369	279	260	138	111
15	531	521	354	371	147	170
16	698	654	460	458	196	191
17	753	768	531	533	188	204
18	882	987	555	618	204	226
19	939	960	571	543	199	208
20	873 <sup>a</sup>	1,014	476 <sup>a</sup>	583	178 <sup>b</sup>	270
21	980	929	478	450	173	170
22	893	944	449	393	169	167
23	845 <sup>a</sup>	1,009	392 <sup>a</sup>	484	143	174
24	776 <sup>a</sup>	895	351	417	104	146
25	664 <sup>a</sup>	775	267	315	93	125
26-29	2,716	2,947	1,200	1,155	469	389
30-34	3,112	3,342	947	1,095	405	491
35-39	3,355	3,360	1,064	1,033	416	388
40-44	3,475	3,615	925	866	312	316
45-49	3,030	3,230	618 <sup>a</sup>	949	311	449
50-54	1,910	2,079	406	394	153	175
55-59	1,045	970	151	120	112 <sup>a</sup>	17
60-64	589	648	139	189	73	51
65 or Older	751	704	132	149	70	131

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.26B Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	12.6	13.1	4.7	4.9	1.9	2.0
12	4.3	4.8	2.3	2.7	1.2	1.2
13	5.3	6.6	3.6	4.3	1.8	1.8
14	9.1	8.9	6.4	6.3	3.2	2.7
15	12.9	12.6	8.6	9.0	3.6	4.1
16	17.2	15.6	11.4	10.9	4.8	4.6
17	19.1	18.8	13.5	13.0	4.8	5.0
18	20.2	21.4	12.7	13.4	4.7	4.9
19	23.0	24.7	14.0	14.0	4.9	5.3
20	21.8 <sup>a</sup>	25.2	11.9 <sup>a</sup>	14.5	4.4 <sup>b</sup>	6.7
21	24.2	23.7	11.8	11.5	4.3	4.3
22	23.1	23.5	11.6	9.8	4.4	4.2
23	22.2 <sup>a</sup>	25.2	10.3	12.1	3.7	4.3
24	22.0	24.6	10.0	11.5	2.9	4.0
25	19.9	21.4	8.0	8.7	2.8	3.5
26-29	17.8	20.0	7.9	7.8	3.1	2.6
30-34	15.6	16.5	4.8	5.4	2.0	2.4
35-39	15.3	16.1	4.8	4.9	1.9	1.9
40-44	15.8	15.9	4.2	3.8	1.4	1.4
45-49	14.3	15.1	2.9 <sup>a</sup>	4.4	1.5	2.1
50-54	10.2	10.4	2.2	2.0	0.8	0.9
55-59	7.1	6.5	1.0	0.8	0.8 <sup>a</sup>	0.1
60-64	5.1	5.5	1.2	1.6	0.6	0.4
65 or Older	2.2	2.1	0.4	0.4	0.2	0.4

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.27A Any Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Numbers in Thousands, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	70,300	71,128	20,423	20,305	8,777	8,849
12	461	471	250	229	116	95
13	623	695	357	416	140	168
14	826	805	519	519	239	202
15	966	948	599	600	248	285
16	1,155	1,143	750	744	331	309
17	1,255	1,258	864	844	347	358
18	1,533	1,552	943	968	356	427
19	1,514	1,490	919	878	359	368
20	1,578	1,630	894	966	350	427
21	1,783	1,601	884	762	350	296
22	1,606	1,700	792	718	341	325
23	1,644	1,731	737	780	311	318
24	1,449	1,574	600	689	213	278
25	1,329	1,492	509	505	179	227
26-29	5,798	5,592	2,016	2,062	871	859
30-34	7,867	7,589	1,868	1,933	826	773
35-39	9,161	8,810	2,112	1,779	900	807
40-44	9,852	9,817	1,828	1,715	817	821
45-49	8,426	8,738	1,399	1,615	688	768
50-54	5,240 <sup>a</sup>	6,527	725	839	305	434
55-59	2,800	2,990	330	263	199 <sup>a</sup>	40
60-64	1,452	1,404	264	263	120	76
65 or Older	1,983	1,573	267	220	171	190

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 1.27B Any Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Percentages, 2002 and 2003**

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	29.9	29.9	8.7	8.5	3.7	3.7
12	11.2	11.4	6.1	5.6	2.8	2.3
13	14.9	16.0	8.5	9.6	3.3	3.9
14	18.9	19.5	11.9	12.6	5.5	4.9
15	23.5	22.9	14.6	14.5	6.0	6.9
16	28.5	27.3	18.5	17.8	8.2	7.4
17	31.8	30.7	21.9	20.6	8.8	8.7
18	35.1	33.7	21.6	21.0	8.2	9.3
19	37.0	38.3	22.5	22.6	8.8	9.5
20	39.5	40.5	22.4	24.0	8.8	10.6
21	44.1	40.8	21.8	19.4	8.6	7.5
22	41.5	42.3	20.5	17.9	8.8	8.1
23	43.2	43.2	19.3	19.5	8.2	7.9
24	41.1	43.3	17.0	19.0	6.0	7.7
25	39.9	41.1	15.3	13.9	5.4	6.3
26-29	38.0	37.9	13.2	14.0	5.7	5.8
30-34	39.5	37.6	9.4	9.6	4.2	3.8
35-39	41.6	42.1	9.6	8.5	4.1	3.9
40-44	44.9	43.3	8.3	7.6	3.7	3.6
45-49	39.8	40.8	6.6	7.5	3.2	3.6
50-54	27.9 <sup>a</sup>	32.6	3.9	4.2	1.6	2.2
55-59	19.1	20.0	2.2	1.8	1.4 <sup>a</sup>	0.3
60-64	12.5	11.9	2.3	2.2	1.0	0.6
65 or Older	5.8	4.6	0.8	0.6	0.5	0.6

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.