

Table 1.103C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 602 | 589 | 151 | 147 | 80 | 72 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 251 | 257 | 59 | 52 | 30 | 26 |
| New England | 156 | 125 | 24 | 22 | 11 | 11 |
| Middle Atlantic | 194 | 227 | 54 | 46 | 28 | 23 |
| Midwest | 232 | 220 | 63 | 44 | 28 | 24 |
| East North Central | 171 | 174 | 50 | 37 | 23 | 21 |
| West North Central | 156 | 134 | 38 | 22 | 16 | 11 |
| South | 327 | 325 | 90 | 86 | 47 | 46 |
| South Atlantic | 267 | 265 | 77 | 73 | 38 | 36 |
| East South Central | 111 | 72 | 27 | 20 | 13 | 14 |
| West South Central | 161 | 175 | 38 | 41 | 23 | 24 |
| West | 362 | 345 | 87 | 97 | 50 | 43 |
| Mountain | 143 | 151 | 37 | 45 | 19 | 14 |
| Pacific | 333 | 310 | 79 | 87 | 46 | 41 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 488 | 453 | 120 | 120 | 68 | 63 |
| Small Metro | 303 | 304 | 82 | 66 | 39 | 29 |
| 250K – 1 Mil. Pop. | 229 | 234 | 64 | 52 | 35 | 20 |
| < 250K Pop. | 192 | 183 | 53 | 37 | 19 | 20 |
| Nonmetro | 197 | 191 | 48 | 45 | 19 | 17 |
| Urbanized | 136 | 125 | 32 | 34 | 9 | 11 |
| Less Urbanized | 122 | 126 | 31 | 28 | 15 | 13 |
| Completely Rural | 39 | 65 | 16 | 13 | 5 | 1 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.103D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.26 | 0.25 | 0.06 | 0.06 | 0.03 | 0.03 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.56 | 0.57 | 0.13 | 0.11 | 0.07 | 0.06 |
| New England | 1.32 | 1.06 | 0.20 | 0.19 | 0.10 | 0.09 |
| Middle Atlantic | 0.58 | 0.68 | 0.16 | 0.14 | 0.08 | 0.07 |
| Midwest | 0.43 | 0.41 | 0.12 | 0.08 | 0.05 | 0.04 |
| East North Central | 0.46 | 0.46 | 0.13 | 0.10 | 0.06 | 0.06 |
| West North Central | 0.98 | 0.83 | 0.24 | 0.14 | 0.10 | 0.07 |
| South | 0.39 | 0.38 | 0.11 | 0.10 | 0.06 | 0.05 |
| South Atlantic | 0.61 | 0.60 | 0.17 | 0.16 | 0.09 | 0.08 |
| East South Central | 0.79 | 0.50 | 0.19 | 0.14 | 0.09 | 0.10 |
| West South Central | 0.62 | 0.67 | 0.15 | 0.16 | 0.09 | 0.09 |
| West | 0.69 | 0.64 | 0.16 | 0.18 | 0.09 | 0.08 |
| Mountain | 0.93 | 0.96 | 0.24 | 0.29 | 0.13 | 0.09 |
| Pacific | 0.89 | 0.81 | 0.21 | 0.23 | 0.12 | 0.11 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.39 | 0.36 | 0.10 | 0.10 | 0.05 | 0.05 |
| Small Metro | 0.44 | 0.43 | 0.12 | 0.09 | 0.06 | 0.04 |
| 250K – 1 Mil. Pop. | 0.50 | 0.50 | 0.14 | 0.11 | 0.08 | 0.04 |
| < 250K Pop. | 0.81 | 0.75 | 0.22 | 0.15 | 0.08 | 0.08 |
| Nonmetro | 0.49 | 0.47 | 0.12 | 0.11 | 0.05 | 0.04 |
| Urbanized | 0.85 | 0.70 | 0.20 | 0.19 | 0.06 | 0.06 |
| Less Urbanized | 0.62 | 0.69 | 0.16 | 0.15 | 0.08 | 0.07 |
| Completely Rural | 0.89 | 1.59 | 0.36 | 0.32 | 0.11 | 0.03 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.103P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.6951 | | 0.0000 | | 0.1149 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.2642 | | 0.3854 | | 0.2947 | |
| New England | 0.9789 | | 0.8191 | | 0.0137 | |
| Middle Atlantic | 0.1577 | | 0.3972 | | 0.8176 | |
| Midwest | 0.5796 | | 0.0014 | | 0.4270 | |
| East North Central | 0.4823 | | 0.0040 | | 0.3748 | |
| West North Central | 0.9464 | | 0.1380 | | 0.9673 | |
| South | 0.2037 | | 0.0001 | | 0.2289 | |
| South Atlantic | 0.1822 | | 0.0010 | | 0.3021 | |
| East South Central | 0.3899 | | 0.0088 | | 0.4539 | |
| West South Central | 0.8489 | | 0.3409 | | 0.7665 | |
| West | 0.4458 | | 0.4010 | | 0.7631 | |
| Mountain | 0.7733 | | 0.2282 | | 0.3253 | |
| Pacific | 0.4803 | | 0.7641 | | 0.9271 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.8505 | | 0.0132 | | 0.5060 | |
| Small Metro | 0.4769 | | 0.0003 | | 0.0774 | |
| 250K – 1 Mil. Pop. | 0.8823 | | 0.0186 | | 0.0517 | |
| < 250K Pop. | 0.3060 | | 0.0044 | | 0.8139 | |
| Nonmetro | 0.6845 | | 0.0935 | | 0.4734 | |
| Urbanized | 0.8533 | | 0.4309 | | 0.8634 | |
| Less Urbanized | 0.5566 | | 0.2539 | | 0.6061 | |
| Completely Rural | 0.2889 | | 0.1538 | | 0.0300 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.104C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 49 | 48 | 40 | 37 | 20 | 20 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 17 | 19 | 14 | 16 | 7 | 9 |
| New England | 7 | 11 | 6 | 9 | 3 | 5 |
| Middle Atlantic | 15 | 16 | 12 | 13 | 6 | 7 |
| Midwest | 20 | 18 | 14 | 15 | 7 | 9 |
| East North Central | 16 | 15 | 12 | 13 | 6 | 7 |
| West North Central | 11 | 10 | 8 | 8 | 3 | 5 |
| South | 33 | 26 | 29 | 22 | 14 | 13 |
| South Atlantic | 27 | 18 | 25 | 16 | 9 | 9 |
| East South Central | 10 | 8 | 8 | 6 | 5 | 3 |
| West South Central | 17 | 16 | 12 | 14 | 10 | 8 |
| West | 24 | 30 | 19 | 21 | 11 | 10 |
| Mountain | 13 | 15 | 9 | 8 | 5 | 5 |
| Pacific | 20 | 26 | 17 | 19 | 10 | 8 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 36 | 38 | 30 | 29 | 15 | 15 |
| Small Metro | 28 | 24 | 24 | 19 | 11 | 11 |
| 250K – 1 Mil. Pop. | 23 | 21 | 20 | 15 | 7 | 9 |
| < 250K Pop. | 16 | 13 | 13 | 11 | 9 | 7 |
| Nonmetro | 17 | 17 | 13 | 13 | 7 | 7 |
| Urbanized | 13 | 10 | 11 | 9 | 5 | 4 |
| Less Urbanized | 11 | 12 | 9 | 8 | 6 | 6 |
| Completely Rural | 5 | 5 | 5 | 5 | 1 | 1 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.104D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.20 | 0.19 | 0.16 | 0.15 | 0.08 | 0.08 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.38 | 0.43 | 0.30 | 0.35 | 0.16 | 0.19 |
| New England | 0.64 | 0.91 | 0.54 | 0.77 | 0.28 | 0.44 |
| Middle Atlantic | 0.46 | 0.48 | 0.36 | 0.39 | 0.18 | 0.21 |
| Midwest | 0.35 | 0.31 | 0.25 | 0.27 | 0.12 | 0.15 |
| East North Central | 0.41 | 0.37 | 0.30 | 0.32 | 0.16 | 0.17 |
| West North Central | 0.63 | 0.58 | 0.48 | 0.47 | 0.16 | 0.32 |
| South | 0.38 | 0.29 | 0.33 | 0.25 | 0.16 | 0.14 |
| South Atlantic | 0.60 | 0.40 | 0.58 | 0.36 | 0.20 | 0.20 |
| East South Central | 0.68 | 0.55 | 0.53 | 0.44 | 0.34 | 0.24 |
| West South Central | 0.57 | 0.54 | 0.41 | 0.47 | 0.32 | 0.28 |
| West | 0.41 | 0.51 | 0.33 | 0.35 | 0.18 | 0.17 |
| Mountain | 0.75 | 0.86 | 0.55 | 0.44 | 0.27 | 0.30 |
| Pacific | 0.50 | 0.63 | 0.41 | 0.46 | 0.24 | 0.20 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.27 | 0.28 | 0.23 | 0.22 | 0.11 | 0.11 |
| Small Metro | 0.39 | 0.33 | 0.33 | 0.25 | 0.16 | 0.15 |
| 250K – 1 Mil. Pop. | 0.47 | 0.41 | 0.41 | 0.31 | 0.15 | 0.17 |
| < 250K Pop. | 0.70 | 0.53 | 0.56 | 0.44 | 0.38 | 0.28 |
| Nonmetro | 0.39 | 0.39 | 0.31 | 0.31 | 0.17 | 0.17 |
| Urbanized | 0.77 | 0.55 | 0.67 | 0.48 | 0.27 | 0.23 |
| Less Urbanized | 0.53 | 0.59 | 0.43 | 0.39 | 0.26 | 0.29 |
| Completely Rural | 1.06 | 1.23 | 1.03 | 1.13 | 0.24 | 0.18 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.104P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0066 | | 0.0011 | | 0.9461 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.1422 | | 0.3622 | | 0.7350 | |
| New England | 0.7999 | | 0.6727 | | 0.2194 | |
| Middle Atlantic | 0.1126 | | 0.4124 | | 0.2740 | |
| Midwest | 0.0750 | | 0.2452 | | 0.9793 | |
| East North Central | 0.0842 | | 0.2475 | | 0.3387 | |
| West North Central | 0.5509 | | 0.7527 | | 0.0750 | |
| South | 0.0044 | | 0.0016 | | 0.5030 | |
| South Atlantic | 0.0000 | | 0.0006 | | 0.2256 | |
| East South Central | 0.4187 | | 0.5371 | | 0.5964 | |
| West South Central | 0.3864 | | 0.6393 | | 0.6119 | |
| West | 0.9155 | | 0.4038 | | 0.5280 | |
| Mountain | 0.7432 | | 0.0071 | | 0.1982 | |
| Pacific | 0.9545 | | 0.8609 | | 0.9829 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.0490 | | 0.0452 | | 0.9642 | |
| Small Metro | 0.2211 | | 0.0973 | | 0.6353 | |
| 250K – 1 Mil. Pop. | 0.4535 | | 0.1679 | | 0.5319 | |
| < 250K Pop. | 0.2755 | | 0.3579 | | 0.9509 | |
| Nonmetro | 0.1333 | | 0.0214 | | 0.5074 | |
| Urbanized | 0.2258 | | 0.1932 | | 0.1557 | |
| Less Urbanized | 0.5708 | | 0.1324 | | 0.6684 | |
| Completely Rural | 0.4292 | | 0.3372 | | 0.3851 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.105C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 140 | 134 | 81 | 77 | 38 | 37 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 56 | 54 | 31 | 33 | 18 | 15 |
| New England | 31 | 25 | 16 | 19 | 10 | 7 |
| Middle Atlantic | 44 | 47 | 26 | 27 | 16 | 13 |
| Midwest | 54 | 46 | 34 | 26 | 12 | 16 |
| East North Central | 46 | 39 | 30 | 22 | 11 | 13 |
| West North Central | 26 | 24 | 17 | 14 | 6 | 8 |
| South | 88 | 80 | 50 | 39 | 25 | 21 |
| South Atlantic | 66 | 57 | 37 | 27 | 17 | 16 |
| East South Central | 33 | 29 | 20 | 14 | 7 | 7 |
| West South Central | 45 | 48 | 24 | 24 | 17 | 12 |
| West | 78 | 80 | 46 | 52 | 18 | 22 |
| Mountain | 32 | 37 | 21 | 22 | 9 | 8 |
| Pacific | 70 | 71 | 40 | 47 | 15 | 20 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 109 | 107 | 59 | 61 | 30 | 31 |
| Small Metro | 78 | 70 | 46 | 43 | 21 | 19 |
| 250K – 1 Mil. Pop. | 59 | 61 | 35 | 38 | 17 | 15 |
| < 250K Pop. | 51 | 37 | 30 | 22 | 12 | 13 |
| Nonmetro | 42 | 38 | 31 | 19 | 12 | 10 |
| Urbanized | 30 | 27 | 26 | 15 | 7 | 7 |
| Less Urbanized | 26 | 24 | 13 | 13 | 7 | 6 |
| Completely Rural | 9 | 9 | 7 | 2 | 5 | 1 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.105D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.45 | 0.42 | 0.26 | 0.24 | 0.12 | 0.12 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 1.01 | 0.95 | 0.56 | 0.58 | 0.33 | 0.26 |
| New England | 2.17 | 1.69 | 1.14 | 1.31 | 0.67 | 0.50 |
| Middle Atlantic | 1.09 | 1.14 | 0.63 | 0.65 | 0.38 | 0.31 |
| Midwest | 0.74 | 0.63 | 0.48 | 0.36 | 0.17 | 0.21 |
| East North Central | 0.93 | 0.77 | 0.60 | 0.43 | 0.21 | 0.26 |
| West North Central | 1.17 | 1.08 | 0.74 | 0.62 | 0.27 | 0.37 |
| South | 0.80 | 0.71 | 0.45 | 0.34 | 0.23 | 0.19 |
| South Atlantic | 1.22 | 1.02 | 0.69 | 0.48 | 0.32 | 0.28 |
| East South Central | 1.74 | 1.52 | 1.07 | 0.70 | 0.34 | 0.38 |
| West South Central | 1.22 | 1.26 | 0.65 | 0.64 | 0.45 | 0.33 |
| West | 1.07 | 1.08 | 0.62 | 0.70 | 0.24 | 0.29 |
| Mountain | 1.46 | 1.66 | 0.97 | 0.98 | 0.40 | 0.35 |
| Pacific | 1.38 | 1.37 | 0.79 | 0.90 | 0.30 | 0.39 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.68 | 0.64 | 0.37 | 0.36 | 0.19 | 0.18 |
| Small Metro | 0.77 | 0.69 | 0.46 | 0.43 | 0.20 | 0.19 |
| 250K – 1 Mil. Pop. | 0.91 | 0.92 | 0.54 | 0.57 | 0.26 | 0.22 |
| < 250K Pop. | 1.37 | 1.03 | 0.81 | 0.61 | 0.32 | 0.35 |
| Nonmetro | 0.84 | 0.76 | 0.61 | 0.39 | 0.24 | 0.20 |
| Urbanized | 1.36 | 1.16 | 1.14 | 0.62 | 0.31 | 0.29 |
| Less Urbanized | 1.13 | 1.08 | 0.59 | 0.58 | 0.30 | 0.29 |
| Completely Rural | 1.88 | 2.47 | 1.46 | 0.55 | 1.06 | 0.31 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.105P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0978 | | 0.0000 | | 0.2807 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.9777 | | 0.3880 | | 0.1548 | |
| New England | 0.9834 | | 0.7556 | | 0.0272 | |
| Middle Atlantic | 0.9806 | | 0.4077 | | 0.7679 | |
| Midwest | 0.2786 | | 0.0044 | | 0.1805 | |
| East North Central | 0.1226 | | 0.0076 | | 0.2860 | |
| West North Central | 0.6004 | | 0.2616 | | 0.4135 | |
| South | 0.2960 | | 0.0001 | | 0.2268 | |
| South Atlantic | 0.2930 | | 0.0003 | | 0.2727 | |
| East South Central | 0.5537 | | 0.0009 | | 0.3464 | |
| West South Central | 0.9076 | | 0.8135 | | 0.7898 | |
| West | 0.3077 | | 0.0378 | | 0.7954 | |
| Mountain | 0.4180 | | 0.0422 | | 0.1306 | |
| Pacific | 0.4593 | | 0.1801 | | 0.6672 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.3080 | | 0.0007 | | 0.7300 | |
| Small Metro | 0.0926 | | 0.0014 | | 0.3582 | |
| 250K – 1 Mil. Pop. | 0.9965 | | 0.2143 | | 0.2813 | |
| < 250K Pop. | 0.0050 | | 0.0001 | | 0.9796 | |
| Nonmetro | 0.9918 | | 0.0329 | | 0.2485 | |
| Urbanized | 0.7260 | | 0.2704 | | 0.8289 | |
| Less Urbanized | 0.8991 | | 0.1524 | | 0.2104 | |
| Completely Rural | 0.9847 | | 0.0025 | | 0.0754 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.106C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 572 | 549 | 109 | 109 | 67 | 54 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 239 | 247 | 40 | 40 | 22 | 20 |
| New England | 152 | 128 | 10 | 10 | 1 | 3 |
| Middle Atlantic | 182 | 214 | 38 | 39 | 22 | 20 |
| Midwest | 223 | 209 | 43 | 28 | 26 | 14 |
| East North Central | 159 | 166 | 33 | 25 | 21 | 14 |
| West North Central | 156 | 128 | 27 | 12 | 15 | * |
| South | 314 | 294 | 64 | 63 | 37 | 34 |
| South Atlantic | 263 | 243 | 53 | 56 | 32 | 28 |
| East South Central | 101 | 66 | 19 | 16 | 10 | 11 |
| West South Central | 147 | 153 | 30 | 24 | 17 | 16 |
| West | 336 | 324 | 67 | 75 | 44 | 34 |
| Mountain | 135 | 134 | 25 | 34 | 16 | 9 |
| Pacific | 308 | 295 | 61 | 66 | 41 | 33 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 459 | 426 | 96 | 94 | 57 | 49 |
| Small Metro | 288 | 281 | 49 | 39 | 32 | 18 |
| 250K – 1 Mil. Pop. | 217 | 218 | 42 | 28 | 30 | 10 |
| < 250K Pop. | 184 | 164 | 26 | 28 | 11 | 15 |
| Nonmetro | 178 | 178 | 25 | 37 | 13 | 12 |
| Urbanized | 126 | 115 | 9 | 26 | 1 | 7 |
| Less Urbanized | 112 | 117 | 22 | 22 | 13 | 10 |
| Completely Rural | 36 | 65 | 9 | 11 | 1 | * |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.106D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.32 | 0.30 | 0.06 | 0.06 | 0.04 | 0.03 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.68 | 0.70 | 0.11 | 0.11 | 0.06 | 0.06 |
| New England | 1.65 | 1.38 | 0.11 | 0.11 | 0.01 | 0.03 |
| Middle Atlantic | 0.70 | 0.82 | 0.15 | 0.15 | 0.08 | 0.08 |
| Midwest | 0.55 | 0.51 | 0.11 | 0.07 | 0.06 | 0.03 |
| East North Central | 0.56 | 0.58 | 0.12 | 0.09 | 0.07 | 0.05 |
| West North Central | 1.29 | 1.06 | 0.22 | 0.10 | 0.12 | * |
| South | 0.49 | 0.46 | 0.10 | 0.10 | 0.06 | 0.05 |
| South Atlantic | 0.77 | 0.71 | 0.16 | 0.16 | 0.09 | 0.08 |
| East South Central | 0.94 | 0.61 | 0.18 | 0.15 | 0.10 | 0.10 |
| West South Central | 0.77 | 0.79 | 0.16 | 0.12 | 0.09 | 0.08 |
| West | 0.85 | 0.80 | 0.17 | 0.18 | 0.11 | 0.08 |
| Mountain | 1.18 | 1.15 | 0.22 | 0.30 | 0.14 | 0.07 |
| Pacific | 1.09 | 1.02 | 0.22 | 0.23 | 0.14 | 0.11 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.48 | 0.44 | 0.10 | 0.10 | 0.06 | 0.05 |
| Small Metro | 0.56 | 0.52 | 0.10 | 0.07 | 0.06 | 0.03 |
| 250K – 1 Mil. Pop. | 0.64 | 0.62 | 0.12 | 0.08 | 0.09 | 0.03 |
| < 250K Pop. | 1.04 | 0.89 | 0.15 | 0.15 | 0.06 | 0.08 |
| Nonmetro | 0.58 | 0.58 | 0.08 | 0.12 | 0.04 | 0.04 |
| Urbanized | 1.04 | 0.85 | 0.07 | 0.19 | 0.01 | 0.05 |
| Less Urbanized | 0.74 | 0.83 | 0.15 | 0.16 | 0.08 | 0.07 |
| Completely Rural | 1.04 | 1.97 | 0.27 | 0.35 | 0.02 | * |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.106P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.8669 | | 0.2990 | | 0.1714 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.2170 | | 0.7358 | | 0.6913 | |
| New England | 0.9826 | | 0.9267 | | 0.4413 | |
| Middle Atlantic | 0.1194 | | 0.7103 | | 0.6321 | |
| Midwest | 0.3661 | | 0.0574 | | 0.0546 | |
| East North Central | 0.2099 | | 0.1621 | | 0.1425 | |
| West North Central | 0.9926 | | 0.1864 | | * | |
| South | 0.3973 | | 0.2112 | | 0.5760 | |
| South Atlantic | 0.3934 | | 0.4872 | | 0.7260 | |
| East South Central | 0.5318 | | 0.4426 | | 0.8456 | |
| West South Central | 0.9040 | | 0.3127 | | 0.6471 | |
| West | 0.5783 | | 0.5104 | | 0.6956 | |
| Mountain | 0.9346 | | 0.7075 | | 0.3454 | |
| Pacific | 0.5743 | | 0.5830 | | 0.9307 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.9035 | | 0.5028 | | 0.4351 | |
| Small Metro | 0.8533 | | 0.1679 | | 0.0766 | |
| 250K – 1 Mil. Pop. | 0.9148 | | 0.1168 | | 0.0425 | |
| < 250K Pop. | 0.8605 | | 0.8347 | | 0.9348 | |
| Nonmetro | 0.5419 | | 0.6009 | | 0.6513 | |
| Urbanized | 0.6508 | | 0.2636 | | 0.0955 | |
| Less Urbanized | 0.5340 | | 0.7655 | | 0.8717 | |
| Completely Rural | 0.2156 | | 0.7856 | | * | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.107C Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 599 | 584 | 144 | 140 | 78 | 68 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 248 | 260 | 54 | 51 | 29 | 25 |
| New England | 153 | 128 | 23 | 21 | 11 | 9 |
| Middle Atlantic | 192 | 229 | 49 | 47 | 27 | 23 |
| Midwest | 227 | 219 | 60 | 40 | 28 | 22 |
| East North Central | 167 | 175 | 49 | 36 | 23 | 20 |
| West North Central | 152 | 133 | 34 | 19 | 16 | 8 |
| South | 329 | 319 | 84 | 80 | 45 | 44 |
| South Atlantic | 271 | 261 | 71 | 69 | 37 | 35 |
| East South Central | 111 | 70 | 25 | 20 | 12 | 13 |
| West South Central | 160 | 172 | 38 | 36 | 23 | 22 |
| West | 361 | 343 | 86 | 93 | 49 | 41 |
| Mountain | 144 | 152 | 36 | 44 | 19 | 12 |
| Pacific | 331 | 307 | 78 | 82 | 46 | 39 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 486 | 451 | 114 | 115 | 66 | 61 |
| Small Metro | 301 | 301 | 77 | 63 | 38 | 26 |
| 250K – 1 Mil. Pop. | 229 | 232 | 59 | 50 | 34 | 18 |
| < 250K Pop. | 192 | 181 | 51 | 36 | 17 | 19 |
| Nonmetro | 194 | 187 | 45 | 42 | 17 | 16 |
| Urbanized | 135 | 122 | 30 | 33 | 8 | 10 |
| Less Urbanized | 119 | 125 | 28 | 26 | 14 | 12 |
| Completely Rural | 38 | 65 | 13 | 12 | 5 | 1 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.107D Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.28 | 0.27 | 0.07 | 0.07 | 0.04 | 0.03 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.61 | 0.64 | 0.13 | 0.13 | 0.07 | 0.06 |
| New England | 1.44 | 1.20 | 0.22 | 0.20 | 0.11 | 0.09 |
| Middle Atlantic | 0.64 | 0.76 | 0.16 | 0.16 | 0.09 | 0.08 |
| Midwest | 0.47 | 0.46 | 0.13 | 0.08 | 0.06 | 0.05 |
| East North Central | 0.50 | 0.52 | 0.15 | 0.11 | 0.07 | 0.06 |
| West North Central | 1.06 | 0.92 | 0.24 | 0.13 | 0.11 | 0.06 |
| South | 0.44 | 0.42 | 0.11 | 0.11 | 0.06 | 0.06 |
| South Atlantic | 0.69 | 0.65 | 0.18 | 0.17 | 0.09 | 0.09 |
| East South Central | 0.87 | 0.55 | 0.20 | 0.16 | 0.09 | 0.10 |
| West South Central | 0.70 | 0.74 | 0.16 | 0.16 | 0.10 | 0.10 |
| West | 0.77 | 0.71 | 0.18 | 0.19 | 0.10 | 0.09 |
| Mountain | 1.06 | 1.10 | 0.26 | 0.32 | 0.14 | 0.08 |
| Pacific | 0.99 | 0.90 | 0.23 | 0.24 | 0.14 | 0.11 |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.43 | 0.40 | 0.10 | 0.10 | 0.06 | 0.05 |
| Small Metro | 0.48 | 0.47 | 0.12 | 0.10 | 0.06 | 0.04 |
| 250K – 1 Mil. Pop. | 0.56 | 0.55 | 0.14 | 0.12 | 0.08 | 0.04 |
| < 250K Pop. | 0.89 | 0.82 | 0.24 | 0.16 | 0.08 | 0.09 |
| Nonmetro | 0.54 | 0.52 | 0.12 | 0.12 | 0.05 | 0.04 |
| Urbanized | 0.94 | 0.76 | 0.21 | 0.20 | 0.05 | 0.06 |
| Less Urbanized | 0.68 | 0.77 | 0.16 | 0.16 | 0.08 | 0.07 |
| Completely Rural | 0.96 | 1.77 | 0.33 | 0.32 | 0.12 | 0.03 |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.107P Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Geographic Characteristic | TIME PERIOD | | | | | |
|--------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.8638 | | 0.0004 | | 0.1025 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.2218 | | 0.5129 | | 0.2404 | |
| New England | 0.9956 | | 0.9326 | | 0.0622 | |
| Middle Atlantic | 0.1265 | | 0.4986 | | 0.5975 | |
| Midwest | 0.4858 | | 0.0025 | | 0.4281 | |
| East North Central | 0.3893 | | 0.0096 | | 0.5629 | |
| West North Central | 0.9203 | | 0.1140 | | 0.5543 | |
| South | 0.3052 | | 0.0024 | | 0.2928 | |
| South Atlantic | 0.3109 | | 0.0216 | | 0.4340 | |
| East South Central | 0.4325 | | 0.0100 | | 0.5298 | |
| West South Central | 0.9199 | | 0.3930 | | 0.6306 | |
| West | 0.4261 | | 0.5015 | | 0.6418 | |
| Mountain | 0.7391 | | 0.4503 | | 0.1328 | |
| Pacific | 0.4689 | | 0.7225 | | 0.9278 | |
| COUNTY TYPE¹ | | | | | | |
| Large Metro | 0.9880 | | 0.0397 | | 0.4875 | |
| Small Metro | 0.5314 | | 0.0010 | | 0.0456 | |
| 250K – 1 Mil. Pop. | 0.9176 | | 0.0431 | | 0.0304 | |
| < 250K Pop. | 0.3506 | | 0.0071 | | 0.7695 | |
| Nonmetro | 0.5797 | | 0.3051 | | 0.6411 | |
| Urbanized | 0.7819 | | 0.7189 | | 0.3380 | |
| Less Urbanized | 0.6027 | | 0.4208 | | 0.4471 | |
| Completely Rural | 0.2483 | | 0.1866 | | 0.0498 | |

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.