

Table 1.19C Any Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	963	909	588	558	388	404
12	27	30	21	20	15	13
13	34	35	26	29	18	19
14	39	37	35	34	26	22
15	43	43	38	38	29	30
16	45	46	43	42	31	35
17	41	46	39	43	35	36
18	49	54	49	55	40	46
19	52	51	50	52	39	47
20	54	50	56	51	47	41
21	49	48	49	44	42	38
22	48	53	41	48	37	36
23	48	49	44	45	34	34
24	48	44	39	45	32	38
25	42	45	38	41	31	34
26-29	191	180	170	157	127	138
30-34	204	210	160	159	120	120
35-39	254	228	195	166	144	131
40-44	230	247	182	163	132	130
45-49	256	225	160	183	126	138
50-54	324	328	154	160	114	126
55-59	260	259	104	105	69	64
60-64	224	217	107	85	83	47
65 or Older	300	296	118	78	97	74

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.19D Any Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.41	0.38	0.25	0.23	0.17	0.17
12	0.67	0.72	0.52	0.48	0.36	0.33
13	0.81	0.81	0.62	0.67	0.44	0.44
14	0.90	0.90	0.79	0.82	0.60	0.53
15	1.05	1.03	0.94	0.91	0.71	0.72
16	1.12	1.10	1.06	1.00	0.77	0.84
17	1.04	1.12	1.00	1.04	0.90	0.89
18	1.12	1.17	1.13	1.19	0.92	1.00
19	1.27	1.32	1.23	1.34	0.96	1.21
20	1.36	1.24	1.41	1.26	1.18	1.01
21	1.20	1.22	1.22	1.12	1.04	0.96
22	1.24	1.33	1.07	1.18	0.97	0.90
23	1.25	1.22	1.16	1.13	0.88	0.86
24	1.36	1.22	1.12	1.23	0.92	1.05
25	1.27	1.24	1.15	1.14	0.92	0.93
26-29	1.25	1.22	1.11	1.07	0.83	0.93
30-34	1.03	1.04	0.80	0.79	0.60	0.60
35-39	1.15	1.09	0.89	0.79	0.66	0.62
40-44	1.05	1.09	0.83	0.72	0.60	0.57
45-49	1.21	1.05	0.76	0.86	0.59	0.65
50-54	1.72	1.64	0.82	0.80	0.60	0.63
55-59	1.77	1.74	0.71	0.71	0.47	0.43
60-64	1.93	1.84	0.92	0.73	0.71	0.40
65 or Older	0.88	0.86	0.35	0.23	0.28	0.22

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.19P Any Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.5445		0.5217		0.6300	
12	0.8360		0.2072		0.2265	
13	0.5312		0.4396		0.7597	
14	0.7435		0.8705		0.3650	
15	0.4847		0.5089		0.9795	
16	0.9599		0.6789		0.3440	
17	0.1654		0.1115		0.0613	
18	0.5437		0.2200		0.8360	
19	0.6439		0.8801		0.4470	
20	0.2282		0.3034		0.4183	
21	0.0024		0.0087		0.4977	
22	0.0414		0.9014		0.9199	
23	0.3926		0.4083		0.2321	
24	0.1498		0.5656		0.8948	
25	0.3331		0.6220		0.5863	
26-29	0.7675		0.4435		0.6365	
30-34	0.1093		0.5523		0.9854	
35-39	0.6904		0.6755		0.8302	
40-44	0.4588		0.6913		0.7785	
45-49	0.3763		0.8046		0.4592	
50-54	0.7105		0.4462		0.5263	
55-59	0.1695		0.2303		0.8163	
60-64	0.9758		0.3097		0.0795	
65 or Older	0.6150		0.2042		0.5560	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.20C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	912	868	471	455	320	327
12	12	10	10	8	6	7
13	22	20	18	17	13	9
14	32	30	29	27	22	19
15	37	37	34	33	24	25
16	40	45	38	40	28	34
17	41	42	38	39	32	32
18	47	52	45	54	38	44
19	52	55	49	50	38	45
20	57	52	54	49	44	38
21	50	51	48	43	40	35
22	48	53	38	44	34	34
23	49	49	41	40	32	31
24	47	46	38	42	31	35
25	42	46	36	39	29	32
26-29	199	188	146	141	113	120
30-34	206	208	148	140	104	100
35-39	260	224	150	135	113	109
40-44	232	255	151	134	105	108
45-49	257	230	133	136	107	95
50-54	325	318	126	127	98	93
55-59	253	252	70	83	39	62
60-64	205	200	88	33	64	30
65 or Older	244	238	86	21	65	18

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.20D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.39	0.37	0.20	0.19	0.14	0.14
12	0.29	0.25	0.24	0.20	0.14	0.16
13	0.54	0.47	0.44	0.39	0.30	0.21
14	0.75	0.72	0.65	0.64	0.50	0.45
15	0.89	0.90	0.83	0.81	0.59	0.61
16	1.00	1.06	0.93	0.94	0.69	0.80
17	1.03	1.03	0.97	0.96	0.82	0.79
18	1.09	1.14	1.03	1.17	0.87	0.94
19	1.27	1.41	1.19	1.30	0.94	1.15
20	1.42	1.28	1.36	1.21	1.09	0.95
21	1.23	1.29	1.19	1.10	0.99	0.90
22	1.25	1.32	0.98	1.09	0.88	0.84
23	1.30	1.22	1.07	1.00	0.84	0.77
24	1.35	1.27	1.07	1.16	0.89	0.95
25	1.27	1.26	1.09	1.06	0.88	0.88
26-29	1.30	1.28	0.96	0.95	0.74	0.81
30-34	1.03	1.03	0.74	0.70	0.52	0.49
35-39	1.18	1.07	0.68	0.65	0.51	0.52
40-44	1.06	1.12	0.69	0.59	0.48	0.48
45-49	1.21	1.07	0.63	0.63	0.51	0.44
50-54	1.73	1.59	0.67	0.64	0.52	0.46
55-59	1.72	1.69	0.47	0.56	0.27	0.42
60-64	1.76	1.70	0.75	0.28	0.55	0.26
65 or Older	0.72	0.69	0.25	0.06	0.19	0.05

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.20P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.6011		0.2191		0.8202	
12	0.0799		0.1177		0.7077	
13	0.0737		0.0947		0.0236	
14	0.5199		0.7440		0.7759	
15	0.1237		0.0888		0.3562	
16	0.9448		0.7296		0.0482	
17	0.0459		0.1022		0.0433	
18	0.4857		0.2448		0.8178	
19	0.6599		0.7488		0.6315	
20	0.2834		0.3760		0.9765	
21	0.0010		0.0176		0.6865	
22	0.0088		0.8602		0.9794	
23	0.7738		0.0652		0.1333	
24	0.5913		0.8839		0.5849	
25	0.4141		0.4315		0.9556	
26-29	0.9939		0.6010		0.4865	
30-34	0.1351		0.9413		0.3434	
35-39	0.4083		0.8513		0.9755	
40-44	0.5360		0.8247		0.3074	
45-49	0.4285		0.3905		0.0784	
50-54	0.5495		0.7983		0.9743	
55-59	0.2018		0.1026		0.0459	
60-64	0.9073		0.0316		0.0837	
65 or Older	0.4469		0.1912		0.2765	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.21C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	583	565	222	229	128	147
12	4	5	4	5	2	2
13	5	7	3	3	1	3
14	10	11	10	9	7	6
15	13	14	11	12	8	7
16	18	20	16	16	11	9
17	24	23	21	17	10	10
18	32	31	29	24	12	16
19	31	32	23	24	12	15
20	36	33	25	23	13	12
21	38	34	27	25	12	13
22	41	37	26	26	14	15
23	39	38	24	26	15	14
24	35	35	19	25	10	14
25	32	34	19	20	10	12
26-29	131	149	78	97	37	50
30-34	175	155	80	74	49	52
35-39	207	196	83	78	51	55
40-44	226	228	73	76	52	49
45-49	208	202	80	73	55	37
50-54	223	237	87	73	36	66
55-59	160	144	36	28	22	*
60-64	105	99	14	*	*	*
65 or Older	102	81	*	14	*	14

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.21D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.25	0.24	0.09	0.10	0.05	0.06
12	0.11	0.12	0.09	0.11	0.05	0.04
13	0.11	0.15	0.07	0.08	0.02	0.07
14	0.24	0.26	0.23	0.21	0.16	0.15
15	0.31	0.33	0.27	0.30	0.19	0.18
16	0.46	0.47	0.40	0.38	0.26	0.21
17	0.61	0.55	0.53	0.42	0.26	0.24
18	0.74	0.67	0.67	0.52	0.27	0.35
19	0.75	0.83	0.56	0.63	0.30	0.38
20	0.90	0.83	0.63	0.58	0.32	0.29
21	0.94	0.87	0.66	0.63	0.31	0.34
22	1.06	0.93	0.66	0.66	0.37	0.38
23	1.02	0.94	0.62	0.65	0.39	0.35
24	0.98	0.97	0.54	0.70	0.30	0.40
25	0.96	0.95	0.57	0.56	0.29	0.33
26-29	0.86	1.01	0.51	0.66	0.24	0.34
30-34	0.88	0.77	0.40	0.37	0.25	0.26
35-39	0.94	0.94	0.38	0.37	0.23	0.26
40-44	1.03	1.01	0.33	0.34	0.23	0.22
45-49	0.98	0.94	0.38	0.34	0.26	0.17
50-54	1.19	1.18	0.46	0.37	0.19	0.33
55-59	1.09	0.97	0.25	0.19	0.15	*
60-64	0.91	0.84	0.12	*	*	*
65 or Older	0.30	0.24	*	0.04	*	0.04

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.21P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4415		0.8514		0.2069	
12	0.8039		0.9883		0.8722	
13	0.4760		0.8298		0.0468	
14	0.2508		0.9822		0.9940	
15	0.8983		0.6819		0.9316	
16	0.7896		0.6066		0.7493	
17	0.1277		0.0162		0.8790	
18	0.1557		0.3569		0.0256	
19	0.9115		0.6136		0.6998	
20	0.2797		0.3499		0.2392	
21	0.0193		0.2568		0.8954	
22	0.7345		0.6591		0.7270	
23	0.8067		0.5674		0.1307	
24	0.9065		0.5938		0.3175	
25	0.9053		0.6994		0.1073	
26-29	0.0887		0.2425		0.1808	
30-34	0.3882		0.9921		0.9149	
35-39	0.4102		0.9523		0.2742	
40-44	0.9273		0.6582		0.6188	
45-49	0.8564		0.4939		0.1091	
50-54	0.0200		0.7807		0.2118	
55-59	0.2782		0.6783		*	
60-64	0.5695		*		*	
65 or Older	0.7120		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.22C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	306	264	124	124	68	91
12	2	*	2	*	2	*
13	3	2	2	2	*	2
14	7	5	6	4	6	2
15	6	7	5	6	2	4
16	8	9	6	7	3	5
17	12	11	9	7	4	3
18	11	16	8	9	2	7
19	17	18	9	10	3	3
20	18	17	7	9	1	5
21	17	17	9	8	5	3
22	20	18	8	7	3	1
23	19	21	8	7	3	3
24	19	21	7	10	4	2
25	18	20	8	8	2	3
26-29	81	83	33	41	9	27
30-34	104	101	53	36	18	26
35-39	124	114	48	44	38	35
40-44	133	114	49	48	38	34
45-49	120	101	45	47	36	22
50-54	114	93	69	51	7	49
55-59	79	44	21	11	21	*
60-64	43	5	14	*	*	*
65 or Older	4	23	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.22D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.13	0.11	0.05	0.05	0.03	0.04
12	0.05	*	0.05	*	0.04	*
13	0.08	0.05	0.06	0.05	*	0.05
14	0.15	0.13	0.15	0.10	0.15	0.04
15	0.16	0.18	0.12	0.14	0.06	0.09
16	0.20	0.21	0.16	0.18	0.07	0.13
17	0.31	0.26	0.22	0.17	0.09	0.06
18	0.26	0.34	0.18	0.20	0.05	0.15
19	0.42	0.45	0.22	0.25	0.08	0.08
20	0.44	0.43	0.19	0.22	0.03	0.14
21	0.42	0.44	0.22	0.21	0.12	0.07
22	0.51	0.44	0.21	0.17	0.08	0.02
23	0.51	0.51	0.21	0.17	0.09	0.07
24	0.54	0.59	0.19	0.27	0.12	0.06
25	0.54	0.55	0.25	0.23	0.07	0.07
26-29	0.53	0.56	0.22	0.28	0.06	0.18
30-34	0.52	0.50	0.26	0.18	0.09	0.13
35-39	0.56	0.55	0.22	0.21	0.17	0.17
40-44	0.61	0.50	0.22	0.21	0.17	0.15
45-49	0.57	0.47	0.21	0.22	0.17	0.10
50-54	0.61	0.47	0.37	0.25	0.04	0.24
55-59	0.54	0.29	0.15	0.07	0.15	*
60-64	0.37	0.04	0.12	*	*	*
65 or Older	0.01	0.07	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.22P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.1637		0.3425		0.7889	
12	*		*		*	
13	0.2242		0.5259		*	
14	0.5001		0.7687		0.3031	
15	0.8161		0.8988		0.4076	
16	0.5879		0.4805		0.1396	
17	0.2882		0.2819		0.9251	
18	0.3974		0.8284		0.0931	
19	0.7724		0.7657		0.5479	
20	0.4911		0.2658		0.2587	
21	0.3112		0.1649		0.4150	
22	0.6328		0.7694		0.2251	
23	0.4779		0.5969		0.8005	
24	0.5993		0.4883		0.3280	
25	0.9172		0.9941		0.9120	
26-29	0.5128		0.7160		0.1351	
30-34	0.8176		0.2790		0.4098	
35-39	0.7999		0.9368		0.8661	
40-44	0.4238		0.9611		0.9765	
45-49	0.1668		0.4380		0.0464	
50-54	0.5041		0.6241		0.1177	
55-59	0.6869		0.6661		*	
60-64	0.0379		*		*	
65 or Older	0.2312		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.23C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	602	589	151	147	80	72
12	7	9	5	6	4	3
13	10	10	9	7	6	4
14	20	13	18	12	8	7
15	20	21	16	16	7	9
16	25	23	21	18	13	11
17	28	28	24	22	11	12
18	40	36	29	28	15	16
19	40	43	26	28	14	14
20	44	43	31	30	14	14
21	44	43	30	24	11	12
22	43	45	25	22	13	13
23	45	41	24	20	12	10
24	42	41	23	23	12	11
25	38	42	17	16	7	8
26-29	160	169	60	71	30	41
30-34	184	163	53	54	28	19
35-39	179	185	45	27	27	18
40-44	206	209	34	27	17	22
45-49	202	185	31	22	19	*
50-54	228	225	40	37	39	*
55-59	124	142	14	*	*	*
60-64	82	79	*	*	*	*
65 or Older	119	49	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.23D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.26	0.25	0.06	0.06	0.03	0.03
12	0.18	0.21	0.13	0.14	0.11	0.08
13	0.25	0.24	0.21	0.17	0.14	0.08
14	0.45	0.32	0.40	0.29	0.17	0.18
15	0.48	0.51	0.39	0.39	0.17	0.22
16	0.62	0.55	0.51	0.44	0.32	0.26
17	0.72	0.67	0.60	0.55	0.27	0.28
18	0.91	0.77	0.67	0.61	0.35	0.34
19	0.97	1.10	0.64	0.71	0.33	0.35
20	1.11	1.08	0.79	0.73	0.34	0.35
21	1.08	1.10	0.75	0.62	0.28	0.30
22	1.12	1.13	0.64	0.55	0.33	0.32
23	1.18	1.02	0.62	0.51	0.31	0.26
24	1.18	1.13	0.67	0.64	0.33	0.31
25	1.15	1.16	0.52	0.45	0.21	0.22
26-29	1.05	1.14	0.39	0.48	0.20	0.28
30-34	0.92	0.81	0.27	0.27	0.14	0.09
35-39	0.81	0.89	0.20	0.13	0.12	0.08
40-44	0.94	0.92	0.16	0.12	0.08	0.10
45-49	0.95	0.86	0.15	0.10	0.09	*
50-54	1.21	1.12	0.21	0.19	0.21	*
55-59	0.85	0.95	0.10	*	*	*
60-64	0.70	0.67	*	*	*	*
65 or Older	0.35	0.14	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.23P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.6951		0.0000		0.1149	
12	0.4444		0.9225		0.1977	
13	0.3856		0.1968		0.1479	
14	0.9840		0.9102		0.4220	
15	0.3974		0.1075		0.3611	
16	0.0187		0.0334		0.9318	
17	0.0237		0.0309		0.9710	
18	0.0042		0.0192		0.5187	
19	0.2767		0.0331		0.5961	
20	0.7330		0.8169		0.8466	
21	0.0074		0.0078		0.1274	
22	0.6098		0.0356		0.9217	
23	0.5438		0.0152		0.9348	
24	0.3215		0.1708		0.8256	
25	0.3792		0.0295		0.9344	
26-29	0.2534		0.3952		0.7468	
30-34	0.0911		0.5143		0.2441	
35-39	0.3938		0.2115		0.8215	
40-44	0.9262		0.1079		0.9764	
45-49	0.2514		0.5210		*	
50-54	0.2517		0.8965		*	
55-59	0.1392		*		*	
60-64	0.8882		*		*	
65 or Older	0.0391		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.24C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	474	441	95	83	64	45
12	19	22	14	12	9	8
13	27	28	16	21	9	11
14	30	27	20	22	12	12
15	30	27	19	18	12	10
16	28	28	17	19	7	9
17	26	26	17	17	6	7
18	31	34	17	20	6	8
19	34	33	18	14	6	5
20	38	35	16	14	9	9
21	35	35	11	11	7	2
22	37	34	12	12	7	5
23	38	35	13	15	6	8
24	31	34	10	12	6	7
25	30	32	6	9	4	5
26-29	128	135	29	25	21	4
30-34	149	138	15	13	3	8
35-39	167	150	12	16	11	12
40-44	169	154	29	25	29	24
45-49	148	169	32	25	23	12
50-54	148	171	*	9	*	9
55-59	90	101	*	*	*	*
60-64	83	53	*	*	*	*
65 or Older	95	76	36	17	35	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.24D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.20	0.19	0.04	0.03	0.03	0.02
12	0.46	0.53	0.35	0.30	0.23	0.20
13	0.64	0.64	0.39	0.47	0.21	0.26
14	0.69	0.65	0.46	0.54	0.27	0.28
15	0.72	0.65	0.45	0.44	0.28	0.24
16	0.68	0.68	0.43	0.47	0.17	0.22
17	0.67	0.64	0.44	0.42	0.16	0.18
18	0.72	0.73	0.39	0.43	0.13	0.18
19	0.82	0.85	0.45	0.36	0.15	0.14
20	0.94	0.88	0.40	0.35	0.22	0.21
21	0.85	0.90	0.27	0.28	0.17	0.05
22	0.94	0.85	0.31	0.29	0.17	0.11
23	1.00	0.87	0.34	0.37	0.16	0.20
24	0.88	0.94	0.29	0.32	0.16	0.20
25	0.91	0.89	0.19	0.24	0.12	0.14
26-29	0.84	0.92	0.19	0.17	0.14	0.03
30-34	0.75	0.69	0.08	0.06	0.02	0.04
35-39	0.76	0.72	0.05	0.07	0.05	0.06
40-44	0.77	0.68	0.13	0.11	0.13	0.11
45-49	0.70	0.79	0.15	0.12	0.11	0.06
50-54	0.79	0.85	*	0.05	*	0.04
55-59	0.61	0.68	*	*	*	*
60-64	0.72	0.45	*	*	*	*
65 or Older	0.28	0.22	0.11	0.05	0.10	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.24P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.8484		0.8088		0.3622	
12	0.4176		0.1035		0.2257	
13	0.4938		0.1248		0.2178	
14	0.9666		0.4104		0.4876	
15	0.7744		1.0000		0.7101	
16	0.7935		0.5018		0.0753	
17	0.7770		0.4961		0.1875	
18	0.3425		0.6121		0.2674	
19	0.3423		0.1907		0.4091	
20	0.8940		0.8658		0.9723	
21	0.6315		0.4104		0.0016	
22	0.0282		0.8538		0.4922	
23	0.0429		0.9714		0.8901	
24	0.8105		0.9847		0.8573	
25	0.8126		0.8492		0.9258	
26-29	0.9050		0.4642		0.1216	
30-34	0.5113		0.6432		0.7286	
35-39	0.3724		0.6257		0.9471	
40-44	0.6657		0.7847		0.9645	
45-49	0.3386		0.8875		0.6102	
50-54	0.0260		*		*	
55-59	0.5223		*		*	
60-64	0.3677		*		*	
65 or Older	0.5274		0.5230		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.25C Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	737	641	377	354	229	250
12	21	21	14	15	11	9
13	21	23	16	18	11	12
14	31	25	25	23	19	15
15	32	30	27	28	16	20
16	37	35	30	29	20	20
17	35	37	31	33	21	23
18	42	44	35	36	24	25
19	44	43	32	34	21	25
20	44	44	33	38	21	26
21	45	45	34	33	23	22
22	42	44	34	31	23	20
23	45	45	32	31	21	20
24	41	42	29	34	16	24
25	39	42	27	28	16	19
26-29	160	164	119	108	80	78
30-34	209	181	108	119	67	84
35-39	223	201	151	110	103	74
40-44	228	219	122	107	79	67
45-49	213	210	107	134	70	109
50-54	255	266	103	110	60	79
55-59	206	189	78	62	57	18
60-64	150	140	82	79	53	37
65 or Older	211	161	74	75	61	72

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.25D Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.31	0.27	0.16	0.15	0.10	0.11
12	0.50	0.50	0.34	0.36	0.26	0.22
13	0.51	0.54	0.39	0.43	0.27	0.28
14	0.70	0.61	0.58	0.56	0.44	0.37
15	0.78	0.72	0.65	0.67	0.40	0.47
16	0.90	0.84	0.74	0.70	0.49	0.48
17	0.88	0.91	0.78	0.81	0.52	0.55
18	0.97	0.95	0.80	0.78	0.55	0.54
19	1.07	1.11	0.79	0.87	0.52	0.65
20	1.09	1.08	0.81	0.93	0.53	0.65
21	1.11	1.14	0.84	0.84	0.58	0.57
22	1.08	1.10	0.88	0.77	0.59	0.51
23	1.19	1.13	0.84	0.78	0.54	0.50
24	1.16	1.15	0.82	0.93	0.45	0.65
25	1.17	1.15	0.80	0.76	0.47	0.53
26-29	1.05	1.11	0.78	0.73	0.52	0.53
30-34	1.05	0.90	0.54	0.59	0.33	0.41
35-39	1.01	0.96	0.68	0.53	0.47	0.35
40-44	1.04	0.97	0.56	0.47	0.36	0.30
45-49	1.00	0.98	0.50	0.63	0.33	0.51
50-54	1.35	1.33	0.55	0.55	0.32	0.40
55-59	1.40	1.27	0.53	0.41	0.39	0.12
60-64	1.29	1.19	0.70	0.68	0.46	0.31
65 or Older	0.62	0.47	0.22	0.22	0.18	0.21

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.25P Nonmedical Use of Any Prescription-Type Psychotherapeutic in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.3887		0.7744		0.8646	
12	0.8884		0.5014		0.9268	
13	0.3015		0.4433		0.8082	
14	0.8935		0.7543		0.3640	
15	0.4250		0.8657		0.2514	
16	0.4176		0.6131		0.5469	
17	0.3738		0.4326		0.7904	
18	0.3446		0.7827		0.5953	
19	0.1895		0.7139		0.4990	
20	0.1861		0.0589		0.0277	
21	0.7115		0.4984		0.7056	
22	0.9617		0.0170		0.3620	
23	0.1097		0.2853		0.5004	
24	0.1175		0.1609		0.0831	
25	0.5027		0.8446		0.2406	
26-29	0.3904		0.9270		0.6582	
30-34	0.7049		0.6138		0.8522	
35-39	0.3820		0.5688		0.5198	
40-44	0.9936		0.2162		0.4774	
45-49	0.8013		0.0628		0.2254	
50-54	0.2070		0.4697		0.4195	
55-59	0.9843		0.6009		0.0230	
60-64	0.6654		0.9319		0.4895	
65 or Older	0.2347		0.7897		0.6828	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.26C Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	569	517	300	299	186	207
12	18	19	13	14	10	8
13	19	22	15	18	11	11
14	28	25	23	22	18	14
15	29	28	24	26	15	18
16	33	31	27	27	18	18
17	33	34	29	30	19	19
18	39	42	32	35	22	22
19	41	40	31	32	20	23
20	40	42	31	35	18	24
21	42	42	30	31	20	19
22	40	43	30	28	20	19
23	42	42	27	30	17	18
24	36	39	25	32	13	19
25	36	36	23	24	13	16
26-29	137	154	106	94	70	65
30-34	167	161	88	108	52	80
35-39	182	162	120	99	74	65
40-44	183	173	100	94	56	56
45-49	175	160	81	114	62	92
50-54	202	185	100	89	57	64
55-59	146	137	44	48	39	12
60-64	121	119	49	72	39	31
65 or Older	156	129	54	66	39	63

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.26D Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.24	0.22	0.13	0.13	0.08	0.09
12	0.44	0.46	0.31	0.34	0.25	0.19
13	0.46	0.51	0.36	0.41	0.26	0.26
14	0.65	0.60	0.54	0.53	0.41	0.34
15	0.72	0.68	0.60	0.62	0.37	0.44
16	0.81	0.75	0.67	0.63	0.45	0.42
17	0.84	0.83	0.73	0.73	0.47	0.47
18	0.90	0.92	0.74	0.75	0.50	0.47
19	1.00	1.02	0.76	0.82	0.48	0.58
20	1.01	1.05	0.77	0.87	0.44	0.60
21	1.05	1.06	0.74	0.79	0.49	0.48
22	1.03	1.06	0.77	0.70	0.52	0.47
23	1.10	1.06	0.71	0.74	0.44	0.46
24	1.03	1.07	0.72	0.87	0.38	0.52
25	1.08	1.00	0.68	0.67	0.40	0.44
26-29	0.90	1.04	0.70	0.64	0.46	0.44
30-34	0.84	0.80	0.44	0.53	0.26	0.39
35-39	0.83	0.78	0.54	0.47	0.34	0.31
40-44	0.83	0.76	0.46	0.42	0.25	0.25
45-49	0.83	0.75	0.38	0.53	0.29	0.43
50-54	1.07	0.93	0.53	0.44	0.30	0.32
55-59	0.99	0.92	0.30	0.32	0.27	0.08
60-64	1.04	1.01	0.42	0.61	0.33	0.27
65 or Older	0.46	0.37	0.16	0.19	0.11	0.18

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.26P Nonmedical Use of Any Pain Reliever in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.0894		0.1726		0.3328	
12	0.3508		0.3920		0.9911	
13	0.0685		0.2107		0.8788	
14	0.8657		0.8705		0.3769	
15	0.7605		0.6794		0.3552	
16	0.1338		0.6291		0.6542	
17	0.8181		0.6823		0.7446	
18	0.3409		0.5118		0.7453	
19	0.2309		0.9969		0.5036	
20	0.0258		0.0298		0.0035	
21	0.7184		0.7515		0.9385	
22	0.7896		0.0757		0.7702	
23	0.0487		0.0817		0.3598	
24	0.0760		0.1817		0.0946	
25	0.3055		0.4824		0.2420	
26-29	0.1250		0.9742		0.4959	
30-34	0.4463		0.3522		0.4027	
35-39	0.4928		0.8923		0.9344	
40-44	0.9363		0.5192		0.9293	
45-49	0.4983		0.0241		0.2463	
50-54	0.8703		0.7787		0.8913	
55-59	0.6330		0.6111		0.0201	
60-64	0.7633		0.5843		0.6483	
65 or Older	0.7943		0.8527		0.4191	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.27C Any Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	815	756	432	420	276	289
12	27	29	20	19	14	12
13	31	33	22	27	14	17
14	36	33	29	29	22	18
15	38	36	31	32	20	23
16	41	41	35	34	24	22
17	38	43	35	36	24	25
18	49	49	43	43	28	30
19	48	49	37	41	26	29
20	49	50	42	42	26	29
21	48	49	41	38	28	25
22	51	52	39	38	27	26
23	48	48	37	37	25	25
24	46	46	33	39	19	27
25	44	46	31	32	18	21
26-29	187	175	136	123	93	83
30-34	215	206	122	129	83	91
35-39	245	229	162	127	112	89
40-44	244	237	141	127	98	90
45-49	247	235	132	144	90	115
50-54	287	321	128	127	80	101
55-59	228	216	85	68	60	18
60-64	168	164	83	79	53	37
65 or Older	246	195	81	76	71	73

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.27D Any Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.35	0.32	0.18	0.18	0.12	0.12
12	0.66	0.70	0.48	0.47	0.34	0.29
13	0.74	0.77	0.53	0.61	0.34	0.40
14	0.83	0.79	0.67	0.70	0.50	0.45
15	0.91	0.88	0.75	0.77	0.49	0.55
16	1.02	0.98	0.85	0.82	0.59	0.52
17	0.97	1.04	0.88	0.88	0.62	0.62
18	1.13	1.06	0.99	0.94	0.64	0.65
19	1.16	1.26	0.91	1.06	0.63	0.75
20	1.24	1.23	1.05	1.04	0.65	0.72
21	1.19	1.24	1.02	0.96	0.68	0.63
22	1.33	1.29	1.00	0.96	0.70	0.65
23	1.27	1.19	0.98	0.92	0.64	0.62
24	1.31	1.25	0.95	1.07	0.53	0.76
25	1.32	1.27	0.93	0.88	0.53	0.57
26-29	1.22	1.19	0.89	0.83	0.61	0.56
30-34	1.08	1.02	0.61	0.64	0.42	0.45
35-39	1.11	1.09	0.74	0.61	0.51	0.42
40-44	1.11	1.04	0.64	0.56	0.45	0.40
45-49	1.16	1.10	0.62	0.67	0.42	0.54
50-54	1.53	1.60	0.68	0.63	0.42	0.50
55-59	1.55	1.45	0.58	0.45	0.41	0.12
60-64	1.45	1.40	0.71	0.68	0.46	0.31
65 or Older	0.72	0.57	0.24	0.22	0.21	0.21

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.27P Any Illicit Drug Use Other Than Marijuana in Lifetime, Past Year, and Past Month, by Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.9491		0.5682		0.9537	
12	0.8370		0.4283		0.2434	
13	0.3068		0.1932		0.3077	
14	0.6185		0.4719		0.3654	
15	0.6522		0.9526		0.2412	
16	0.3526		0.5131		0.3007	
17	0.4770		0.3297		0.9692	
18	0.3475		0.6648		0.2259	
19	0.4476		0.9442		0.4663	
20	0.5774		0.2870		0.0642	
21	0.0605		0.0982		0.2601	
22	0.6566		0.0602		0.4354	
23	0.9595		0.9112		0.7986	
24	0.2383		0.1889		0.0778	
25	0.5091		0.2640		0.2299	
26-29	0.9560		0.5356		0.8933	
30-34	0.1676		0.8374		0.6014	
35-39	0.7699		0.2625		0.7217	
40-44	0.2910		0.3668		0.8543	
45-49	0.5346		0.3125		0.6325	
50-54	0.0338		0.7123		0.4062	
55-59	0.6351		0.5155		0.0118	
60-64	0.7830		0.9715		0.4895	
65 or Older	0.1903		0.6556		0.8674	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.