

Table 1.1C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 963 | 909 | 588 | 558 | 388 | 404 |
| Marijuana and Hashish | 912 | 868 | 471 | 455 | 320 | 327 |
| Cocaine | 583 | 565 | 222 | 229 | 128 | 147 |
| Crack | 306 | 264 | 124 | 124 | 68 | 91 |
| Heroin | 223 | 205 | 55 | 48 | 42 | 28 |
| Hallucinogens | 602 | 589 | 151 | 147 | 80 | 72 |
| LSD | 514 | 494 | 63 | 44 | 18 | 18 |
| PCP | 284 | 271 | 25 | 24 | 15 | 13 |
| Ecstasy | 297 | 285 | 126 | 105 | 64 | 55 |
| Inhalants | 474 | 441 | 95 | 83 | 64 | 45 |
| Nonmedical Use of Any Psychotherapeutic ² | 737 | 641 | 377 | 354 | 229 | 250 |
| Pain Relievers | 569 | 517 | 300 | 299 | 186 | 207 |
| Tranquilizers | 497 | 435 | 225 | 195 | 136 | 118 |
| Stimulants | 495 | 459 | 149 | 128 | 90 | 93 |
| Methamphetamine | 369 | 369 | 109 | 94 | 73 | 75 |
| Sedatives | 370 | 325 | 116 | 85 | 64 | 55 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 815 | 756 | 432 | 420 | 276 | 289 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.1D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.41 | 0.38 | 0.25 | 0.23 | 0.17 | 0.17 |
| Marijuana and Hashish | 0.39 | 0.37 | 0.20 | 0.19 | 0.14 | 0.14 |
| Cocaine | 0.25 | 0.24 | 0.09 | 0.10 | 0.05 | 0.06 |
| Crack | 0.13 | 0.11 | 0.05 | 0.05 | 0.03 | 0.04 |
| Heroin | 0.10 | 0.09 | 0.02 | 0.02 | 0.02 | 0.01 |
| Hallucinogens | 0.26 | 0.25 | 0.06 | 0.06 | 0.03 | 0.03 |
| LSD | 0.22 | 0.21 | 0.03 | 0.02 | 0.01 | 0.01 |
| PCP | 0.12 | 0.11 | 0.01 | 0.01 | 0.01 | 0.01 |
| Ecstasy | 0.13 | 0.12 | 0.05 | 0.04 | 0.03 | 0.02 |
| Inhalants | 0.20 | 0.19 | 0.04 | 0.03 | 0.03 | 0.02 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.31 | 0.27 | 0.16 | 0.15 | 0.10 | 0.11 |
| Pain Relievers | 0.24 | 0.22 | 0.13 | 0.13 | 0.08 | 0.09 |
| Tranquilizers | 0.21 | 0.18 | 0.10 | 0.08 | 0.06 | 0.05 |
| Stimulants | 0.21 | 0.19 | 0.06 | 0.05 | 0.04 | 0.04 |
| Methamphetamine | 0.16 | 0.16 | 0.05 | 0.04 | 0.03 | 0.03 |
| Sedatives | 0.16 | 0.14 | 0.05 | 0.04 | 0.03 | 0.02 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.35 | 0.32 | 0.18 | 0.18 | 0.12 | 0.12 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.1P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.5445 | | 0.5217 | | 0.6300 | |
| Marijuana and Hashish | 0.6011 | | 0.2191 | | 0.8202 | |
| Cocaine | 0.4415 | | 0.8514 | | 0.2069 | |
| Crack | 0.1637 | | 0.3425 | | 0.7889 | |
| Heroin | 0.9050 | | 0.2101 | | 0.3327 | |
| Hallucinogens | 0.6951 | | 0.0000 | | 0.1149 | |
| LSD | 0.6127 | | 0.0000 | | 0.4406 | |
| PCP | 0.3193 | | 0.5680 | | 0.8982 | |
| Ecstasy | 0.1102 | | 0.0000 | | 0.0112 | |
| Inhalants | 0.8484 | | 0.8088 | | 0.3622 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.3887 | | 0.7744 | | 0.8646 | |
| Pain Relievers | 0.0894 | | 0.1726 | | 0.3328 | |
| Tranquilizers | 0.2567 | | 0.6069 | | 0.9722 | |
| Stimulants | 0.4432 | | 0.0215 | | 0.7636 | |
| Methamphetamine | 0.6807 | | 0.1106 | | 0.9757 | |
| Sedatives | 0.2717 | | 0.2684 | | 0.0860 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.9491 | | 0.5682 | | 0.9537 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.2C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 104 | 107 | 95 | 91 | 71 | 67 |
| Marijuana and Hashish | 88 | 91 | 79 | 79 | 58 | 60 |
| Cocaine | 36 | 36 | 32 | 28 | 18 | 16 |
| Crack | 17 | 17 | 14 | 13 | 9 | 8 |
| Heroin | 14 | 11 | 11 | 8 | 5 | 6 |
| Hallucinogens | 49 | 48 | 40 | 37 | 20 | 20 |
| LSD | 34 | 26 | 25 | 15 | 12 | 10 |
| PCP | 20 | 20 | 13 | 12 | 6 | 6 |
| Ecstasy | 40 | 32 | 32 | 23 | 16 | 12 |
| Inhalants | 67 | 68 | 44 | 48 | 22 | 25 |
| Nonmedical Use of Any Psychotherapeutic ² | 75 | 76 | 64 | 64 | 42 | 42 |
| Pain Relievers | 69 | 70 | 59 | 59 | 39 | 39 |
| Tranquilizers | 38 | 40 | 34 | 33 | 18 | 22 |
| Stimulants | 46 | 40 | 35 | 30 | 20 | 19 |
| Methamphetamine | 27 | 22 | 21 | 17 | 12 | 11 |
| Sedatives | 22 | 22 | 16 | 16 | 9 | 10 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 89 | 97 | 76 | 79 | 52 | 48 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.2D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.42 | 0.43 | 0.38 | 0.36 | 0.29 | 0.27 |
| Marijuana and Hashish | 0.36 | 0.37 | 0.32 | 0.31 | 0.24 | 0.24 |
| Cocaine | 0.15 | 0.15 | 0.13 | 0.11 | 0.07 | 0.06 |
| Crack | 0.07 | 0.07 | 0.06 | 0.05 | 0.03 | 0.03 |
| Heroin | 0.05 | 0.04 | 0.04 | 0.03 | 0.02 | 0.02 |
| Hallucinogens | 0.20 | 0.19 | 0.16 | 0.15 | 0.08 | 0.08 |
| LSD | 0.14 | 0.10 | 0.10 | 0.06 | 0.05 | 0.04 |
| PCP | 0.08 | 0.08 | 0.05 | 0.05 | 0.02 | 0.02 |
| Ecstasy | 0.16 | 0.13 | 0.13 | 0.09 | 0.06 | 0.05 |
| Inhalants | 0.27 | 0.27 | 0.18 | 0.19 | 0.09 | 0.10 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.30 | 0.30 | 0.26 | 0.26 | 0.17 | 0.17 |
| Pain Relievers | 0.28 | 0.28 | 0.24 | 0.24 | 0.16 | 0.15 |
| Tranquilizers | 0.16 | 0.16 | 0.14 | 0.13 | 0.07 | 0.09 |
| Stimulants | 0.18 | 0.16 | 0.14 | 0.12 | 0.08 | 0.08 |
| Methamphetamine | 0.11 | 0.09 | 0.09 | 0.07 | 0.05 | 0.04 |
| Sedatives | 0.09 | 0.09 | 0.07 | 0.06 | 0.03 | 0.04 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.36 | 0.39 | 0.31 | 0.32 | 0.21 | 0.19 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.2P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.5614 | | 0.4872 | | 0.3311 | |
| Marijuana and Hashish | 0.0356 | | 0.0780 | | 0.3878 | |
| Cocaine | 0.5530 | | 0.0730 | | 0.8041 | |
| Crack | 0.7236 | | 0.7667 | | 0.6129 | |
| Heroin | 0.4102 | | 0.2709 | | 0.4199 | |
| Hallucinogens | 0.0066 | | 0.0011 | | 0.9461 | |
| LSD | 0.0000 | | 0.0000 | | 0.5120 | |
| PCP | 0.2649 | | 0.2074 | | 0.7363 | |
| Ecstasy | 0.0000 | | 0.0000 | | 0.0940 | |
| Inhalants | 0.6767 | | 0.6422 | | 0.6467 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.5012 | | 0.8584 | | 0.8951 | |
| Pain Relievers | 0.9726 | | 0.7254 | | 0.9876 | |
| Tranquilizers | 0.5772 | | 0.7938 | | 0.2041 | |
| Stimulants | 0.2004 | | 0.0562 | | 0.4929 | |
| Methamphetamine | 0.2342 | | 0.0490 | | 0.7262 | |
| Sedatives | 0.7615 | | 0.7149 | | 0.9458 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.8845 | | 0.8608 | | 0.8018 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.3C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 156 | 155 | 143 | 153 | 116 | 128 |
| Marijuana and Hashish | 157 | 157 | 135 | 146 | 110 | 116 |
| Cocaine | 114 | 107 | 74 | 72 | 37 | 42 |
| Crack | 51 | 54 | 23 | 23 | 9 | 13 |
| Heroin | 35 | 37 | 17 | 14 | 10 | 7 |
| Hallucinogens | 140 | 134 | 81 | 77 | 38 | 37 |
| LSD | 115 | 108 | 44 | 30 | 8 | 13 |
| PCP | 45 | 49 | 14 | 19 | 6 | 10 |
| Ecstasy | 112 | 109 | 67 | 50 | 31 | 23 |
| Inhalants | 111 | 104 | 41 | 39 | 18 | 19 |
| Nonmedical Use of Any Psychotherapeutic ² | 135 | 135 | 96 | 103 | 60 | 70 |
| Pain Relievers | 119 | 130 | 86 | 97 | 51 | 61 |
| Tranquilizers | 94 | 98 | 57 | 66 | 35 | 36 |
| Stimulants | 89 | 94 | 56 | 54 | 30 | 30 |
| Methamphetamine | 61 | 65 | 38 | 36 | 20 | 21 |
| Sedatives | 39 | 37 | 18 | 18 | 9 | 11 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 157 | 151 | 116 | 127 | 75 | 82 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.3D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.50 | 0.49 | 0.46 | 0.48 | 0.37 | 0.40 |
| Marijuana and Hashish | 0.51 | 0.50 | 0.43 | 0.46 | 0.36 | 0.37 |
| Cocaine | 0.37 | 0.34 | 0.24 | 0.23 | 0.12 | 0.13 |
| Crack | 0.16 | 0.17 | 0.08 | 0.07 | 0.03 | 0.04 |
| Heroin | 0.11 | 0.12 | 0.05 | 0.04 | 0.03 | 0.02 |
| Hallucinogens | 0.45 | 0.42 | 0.26 | 0.24 | 0.12 | 0.12 |
| LSD | 0.37 | 0.34 | 0.14 | 0.09 | 0.03 | 0.04 |
| PCP | 0.15 | 0.16 | 0.04 | 0.06 | 0.02 | 0.03 |
| Ecstasy | 0.36 | 0.34 | 0.22 | 0.16 | 0.10 | 0.07 |
| Inhalants | 0.36 | 0.33 | 0.13 | 0.12 | 0.06 | 0.06 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.43 | 0.43 | 0.31 | 0.33 | 0.19 | 0.22 |
| Pain Relievers | 0.39 | 0.41 | 0.28 | 0.30 | 0.16 | 0.19 |
| Tranquilizers | 0.30 | 0.31 | 0.18 | 0.21 | 0.11 | 0.11 |
| Stimulants | 0.29 | 0.29 | 0.18 | 0.17 | 0.10 | 0.09 |
| Methamphetamine | 0.20 | 0.20 | 0.12 | 0.11 | 0.06 | 0.07 |
| Sedatives | 0.13 | 0.12 | 0.06 | 0.06 | 0.03 | 0.04 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.51 | 0.47 | 0.37 | 0.40 | 0.24 | 0.26 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.3P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.3244 | | 0.1830 | | 0.8639 | |
| Marijuana and Hashish | 0.8370 | | 0.0365 | | 0.5307 | |
| Cocaine | 0.3762 | | 0.7314 | | 0.3692 | |
| Crack | 0.8333 | | 0.9486 | | 0.9523 | |
| Heroin | 0.7162 | | 0.1958 | | 0.2161 | |
| Hallucinogens | 0.0978 | | 0.0000 | | 0.2807 | |
| LSD | 0.0001 | | 0.0000 | | 0.0632 | |
| PCP | 0.1595 | | 0.3990 | | 0.2155 | |
| Ecstasy | 0.5186 | | 0.0000 | | 0.0004 | |
| Inhalants | 0.1087 | | 0.5097 | | 0.4771 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.0228 | | 0.5291 | | 0.0300 | |
| Pain Relievers | 0.0030 | | 0.1570 | | 0.0103 | |
| Tranquilizers | 0.0081 | | 0.1061 | | 0.5747 | |
| Stimulants | 0.9700 | | 0.4592 | | 0.9053 | |
| Methamphetamine | 0.0843 | | 0.5480 | | 0.4563 | |
| Sedatives | 0.0732 | | 0.7193 | | 0.5864 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.8105 | | 0.3854 | | 0.1587 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.4C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 935 | 874 | 514 | 489 | 353 | 363 |
| Marijuana and Hashish | 888 | 831 | 407 | 387 | 279 | 284 |
| Cocaine | 574 | 552 | 200 | 207 | 119 | 137 |
| Crack | 303 | 257 | 121 | 120 | 66 | 90 |
| Heroin | 220 | 201 | 52 | 45 | 40 | 26 |
| Hallucinogens | 572 | 549 | 109 | 109 | 67 | 54 |
| LSD | 496 | 469 | 35 | 28 | 11 | 8 |
| PCP | 282 | 262 | 16 | 6 | 13 | * |
| Ecstasy | 247 | 255 | 93 | 86 | 53 | 46 |
| Inhalants | 457 | 407 | 66 | 54 | 56 | 32 |
| Nonmedical Use of Any Psychotherapeutic ² | 707 | 617 | 345 | 331 | 215 | 237 |
| Pain Relievers | 539 | 477 | 275 | 275 | 172 | 193 |
| Tranquilizers | 481 | 414 | 210 | 175 | 127 | 112 |
| Stimulants | 475 | 447 | 126 | 107 | 82 | 83 |
| Methamphetamine | 356 | 365 | 95 | 85 | 67 | 70 |
| Sedatives | 366 | 316 | 113 | 81 | 63 | 53 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 789 | 728 | 384 | 375 | 259 | 274 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.4D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.52 | 0.48 | 0.29 | 0.27 | 0.20 | 0.20 |
| Marijuana and Hashish | 0.50 | 0.46 | 0.23 | 0.21 | 0.16 | 0.16 |
| Cocaine | 0.32 | 0.30 | 0.11 | 0.11 | 0.07 | 0.08 |
| Crack | 0.17 | 0.14 | 0.07 | 0.07 | 0.04 | 0.05 |
| Heroin | 0.12 | 0.11 | 0.03 | 0.02 | 0.02 | 0.01 |
| Hallucinogens | 0.32 | 0.30 | 0.06 | 0.06 | 0.04 | 0.03 |
| LSD | 0.28 | 0.26 | 0.02 | 0.02 | 0.01 | 0.00 |
| PCP | 0.16 | 0.14 | 0.01 | 0.00 | 0.01 | * |
| Ecstasy | 0.14 | 0.14 | 0.05 | 0.05 | 0.03 | 0.03 |
| Inhalants | 0.25 | 0.22 | 0.04 | 0.03 | 0.03 | 0.02 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.39 | 0.34 | 0.19 | 0.18 | 0.12 | 0.13 |
| Pain Relievers | 0.30 | 0.26 | 0.15 | 0.15 | 0.10 | 0.11 |
| Tranquilizers | 0.27 | 0.23 | 0.12 | 0.10 | 0.07 | 0.06 |
| Stimulants | 0.26 | 0.25 | 0.07 | 0.06 | 0.05 | 0.05 |
| Methamphetamine | 0.20 | 0.20 | 0.05 | 0.05 | 0.04 | 0.04 |
| Sedatives | 0.20 | 0.17 | 0.06 | 0.04 | 0.04 | 0.03 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.44 | 0.40 | 0.21 | 0.21 | 0.14 | 0.15 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.4P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.6311 | | 0.7321 | | 0.6286 | |
| Marijuana and Hashish | 0.4913 | | 0.6227 | | 0.9637 | |
| Cocaine | 0.3191 | | 0.8974 | | 0.3141 | |
| Crack | 0.1751 | | 0.3408 | | 0.8281 | |
| Heroin | 0.9105 | | 0.4610 | | 0.4002 | |
| Hallucinogens | 0.8669 | | 0.2990 | | 0.1714 | |
| LSD | 0.4706 | | 0.1884 | | 1.0000 | |
| PCP | 0.2355 | | 0.2782 | | * | |
| Ecstasy | 0.0093 | | 0.0714 | | 0.4844 | |
| Inhalants | 0.9175 | | 0.7414 | | 0.2953 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.6417 | | 0.9305 | | 0.6434 | |
| Pain Relievers | 0.3039 | | 0.4035 | | 0.8065 | |
| Tranquilizers | 0.5802 | | 0.9608 | | 0.7066 | |
| Stimulants | 0.4931 | | 0.0520 | | 0.5706 | |
| Methamphetamine | 0.9580 | | 0.2110 | | 0.7995 | |
| Sedatives | 0.3590 | | 0.3061 | | 0.0630 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.9959 | | 0.6805 | | 0.6475 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.5C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 44 | 46 | 34 | 35 | 24 | 24 |
| Marijuana and Hashish | 25 | 24 | 21 | 19 | 14 | 11 |
| Cocaine | 6 | 8 | 5 | 6 | 2 | 3 |
| Crack | 4 | 2 | 3 | 2 | 2 | 2 |
| Heroin | 5 | 3 | 1 | 2 | * | * |
| Hallucinogens | 12 | 13 | 10 | 9 | 7 | 5 |
| LSD | 5 | 5 | 4 | 4 | 3 | 3 |
| PCP | 6 | 3 | 4 | 2 | 3 | 1 |
| Ecstasy | 8 | 8 | 7 | 5 | 5 | 2 |
| Inhalants | 33 | 36 | 21 | 24 | 12 | 14 |
| Nonmedical Use of Any Psychotherapeutic ² | 30 | 31 | 22 | 23 | 16 | 15 |
| Pain Relievers | 26 | 29 | 20 | 22 | 15 | 14 |
| Tranquilizers | 11 | 9 | 8 | 8 | 5 | 7 |
| Stimulants | 14 | 13 | 9 | 9 | 4 | 7 |
| Methamphetamine | 5 | 7 | 3 | 4 | 2 | 3 |
| Sedatives | 9 | 8 | 5 | 7 | 3 | 2 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 41 | 43 | 30 | 32 | 21 | 21 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.5D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.53 | 0.54 | 0.41 | 0.42 | 0.29 | 0.28 |
| Marijuana and Hashish | 0.31 | 0.28 | 0.25 | 0.23 | 0.17 | 0.13 |
| Cocaine | 0.08 | 0.10 | 0.06 | 0.07 | 0.02 | 0.04 |
| Crack | 0.05 | 0.02 | 0.04 | 0.02 | 0.02 | 0.02 |
| Heroin | 0.06 | 0.04 | 0.01 | 0.03 | * | * |
| Hallucinogens | 0.15 | 0.16 | 0.12 | 0.11 | 0.08 | 0.06 |
| LSD | 0.06 | 0.05 | 0.05 | 0.04 | 0.04 | 0.03 |
| PCP | 0.07 | 0.03 | 0.05 | 0.02 | 0.04 | 0.02 |
| Ecstasy | 0.09 | 0.09 | 0.08 | 0.06 | 0.06 | 0.02 |
| Inhalants | 0.39 | 0.42 | 0.26 | 0.29 | 0.15 | 0.17 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.36 | 0.37 | 0.26 | 0.27 | 0.19 | 0.18 |
| Pain Relievers | 0.31 | 0.34 | 0.24 | 0.26 | 0.18 | 0.17 |
| Tranquilizers | 0.13 | 0.11 | 0.09 | 0.09 | 0.06 | 0.08 |
| Stimulants | 0.17 | 0.16 | 0.10 | 0.11 | 0.05 | 0.08 |
| Methamphetamine | 0.06 | 0.08 | 0.04 | 0.05 | 0.02 | 0.03 |
| Sedatives | 0.11 | 0.09 | 0.06 | 0.08 | 0.04 | 0.02 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.49 | 0.51 | 0.36 | 0.38 | 0.25 | 0.25 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.5P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.6677 | | 0.9489 | | 0.3678 | |
| Marijuana and Hashish | 0.0211 | | 0.0306 | | 0.0308 | |
| Cocaine | 0.6600 | | 0.9058 | | 0.1486 | |
| Crack | 0.0947 | | 0.2358 | | 0.9447 | |
| Heroin | 0.2643 | | 0.6687 | | * | |
| Hallucinogens | 0.8501 | | 0.2642 | | 0.0440 | |
| LSD | 0.5536 | | 0.6440 | | 0.6714 | |
| PCP | 0.0380 | | 0.0919 | | 0.2573 | |
| Ecstasy | 0.0887 | | 0.0075 | | 0.0269 | |
| Inhalants | 0.2829 | | 0.7724 | | 0.8574 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.3842 | | 0.2911 | | 0.7925 | |
| Pain Relievers | 0.0474 | | 0.1256 | | 0.9158 | |
| Tranquilizers | 0.4182 | | 0.9173 | | 0.7722 | |
| Stimulants | 0.8218 | | 0.8544 | | 0.1103 | |
| Methamphetamine | 0.1715 | | 0.4306 | | 0.1896 | |
| Sedatives | 0.7615 | | 0.3984 | | 0.2466 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.3228 | | 0.5727 | | 0.9381 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.6C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 56 | 58 | 52 | 51 | 38 | 37 |
| Marijuana and Hashish | 50 | 49 | 45 | 43 | 32 | 32 |
| Cocaine | 16 | 18 | 15 | 15 | 10 | 10 |
| Crack | 9 | 9 | 8 | 7 | 7 | 4 |
| Heroin | 7 | 6 | 5 | 4 | * | 2 |
| Hallucinogens | 26 | 25 | 23 | 20 | 10 | 12 |
| LSD | 16 | 15 | 12 | 10 | 7 | 5 |
| PCP | 10 | 13 | 7 | 8 | 5 | 0 |
| Ecstasy | 18 | 16 | 16 | 12 | 6 | 7 |
| Inhalants | 42 | 40 | 26 | 30 | 16 | 16 |
| Nonmedical Use of Any Psychotherapeutic ² | 41 | 41 | 36 | 36 | 25 | 25 |
| Pain Relievers | 38 | 39 | 34 | 34 | 23 | 23 |
| Tranquilizers | 20 | 21 | 20 | 18 | 10 | 13 |
| Stimulants | 24 | 21 | 21 | 17 | 10 | 10 |
| Methamphetamine | 14 | 12 | 12 | 10 | 7 | 5 |
| Sedatives | 12 | 13 | 9 | 9 | 5 | 7 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 49 | 50 | 41 | 43 | 29 | 29 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.6D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.67 | 0.70 | 0.61 | 0.62 | 0.45 | 0.44 |
| Marijuana and Hashish | 0.60 | 0.59 | 0.53 | 0.52 | 0.38 | 0.39 |
| Cocaine | 0.19 | 0.22 | 0.18 | 0.18 | 0.12 | 0.12 |
| Crack | 0.11 | 0.11 | 0.09 | 0.08 | 0.08 | 0.05 |
| Heroin | 0.08 | 0.07 | 0.05 | 0.05 | * | 0.03 |
| Hallucinogens | 0.31 | 0.30 | 0.27 | 0.25 | 0.12 | 0.14 |
| LSD | 0.19 | 0.18 | 0.14 | 0.12 | 0.08 | 0.06 |
| PCP | 0.12 | 0.15 | 0.08 | 0.09 | 0.05 | 0.00 |
| Ecstasy | 0.22 | 0.19 | 0.19 | 0.14 | 0.08 | 0.08 |
| Inhalants | 0.49 | 0.49 | 0.31 | 0.37 | 0.18 | 0.20 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.49 | 0.49 | 0.43 | 0.43 | 0.30 | 0.30 |
| Pain Relievers | 0.45 | 0.47 | 0.40 | 0.41 | 0.28 | 0.27 |
| Tranquilizers | 0.24 | 0.25 | 0.23 | 0.22 | 0.12 | 0.15 |
| Stimulants | 0.28 | 0.25 | 0.24 | 0.20 | 0.12 | 0.12 |
| Methamphetamine | 0.16 | 0.14 | 0.15 | 0.12 | 0.08 | 0.06 |
| Sedatives | 0.14 | 0.15 | 0.11 | 0.11 | 0.06 | 0.08 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.58 | 0.60 | 0.49 | 0.52 | 0.34 | 0.35 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.6P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.5251 | | 0.6257 | | 0.6591 | |
| Marijuana and Hashish | 0.1653 | | 0.1781 | | 0.4380 | |
| Cocaine | 0.5126 | | 0.8033 | | 0.9238 | |
| Crack | 0.5113 | | 0.7482 | | 0.7135 | |
| Heroin | 0.9214 | | 0.9034 | | * | |
| Hallucinogens | 0.5346 | | 0.2099 | | 0.2087 | |
| LSD | 0.0117 | | 0.0128 | | 0.7767 | |
| PCP | 0.5127 | | 0.9628 | | 0.0067 | |
| Ecstasy | 0.5707 | | 0.0597 | | 0.7375 | |
| Inhalants | 0.8691 | | 0.5459 | | 0.4338 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.5556 | | 0.9922 | | 0.8154 | |
| Pain Relievers | 0.7993 | | 0.7953 | | 0.9307 | |
| Tranquilizers | 0.8718 | | 0.9909 | | 0.4091 | |
| Stimulants | 0.2246 | | 0.1865 | | 0.8964 | |
| Methamphetamine | 0.0731 | | 0.1873 | | 0.8722 | |
| Sedatives | 0.7009 | | 0.9424 | | 0.8622 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.9401 | | 0.6177 | | 0.7587 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.7C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 61 | 66 | 58 | 62 | 48 | 53 |
| Marijuana and Hashish | 57 | 63 | 55 | 58 | 43 | 48 |
| Cocaine | 31 | 31 | 27 | 23 | 15 | 13 |
| Crack | 14 | 14 | 11 | 10 | 4 | 6 |
| Heroin | 11 | 9 | 10 | 6 | 5 | 6 |
| Hallucinogens | 38 | 36 | 31 | 28 | 16 | 16 |
| LSD | 27 | 21 | 20 | 12 | 10 | 8 |
| PCP | 16 | 15 | 10 | 9 | 2 | 6 |
| Ecstasy | 32 | 26 | 26 | 20 | 13 | 10 |
| Inhalants | 39 | 38 | 26 | 27 | 9 | 12 |
| Nonmedical Use of Any Psychotherapeutic ² | 50 | 52 | 43 | 45 | 30 | 30 |
| Pain Relievers | 47 | 47 | 41 | 40 | 28 | 27 |
| Tranquilizers | 30 | 32 | 25 | 26 | 14 | 16 |
| Stimulants | 32 | 30 | 25 | 22 | 16 | 15 |
| Methamphetamine | 22 | 18 | 17 | 13 | 10 | 9 |
| Sedatives | 14 | 15 | 11 | 10 | 6 | 7 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 58 | 60 | 50 | 49 | 36 | 33 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.7D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.76 | 0.80 | 0.73 | 0.75 | 0.60 | 0.63 |
| Marijuana and Hashish | 0.71 | 0.76 | 0.68 | 0.70 | 0.53 | 0.57 |
| Cocaine | 0.38 | 0.37 | 0.34 | 0.28 | 0.18 | 0.15 |
| Crack | 0.18 | 0.17 | 0.13 | 0.12 | 0.06 | 0.07 |
| Heroin | 0.14 | 0.11 | 0.13 | 0.08 | 0.06 | 0.07 |
| Hallucinogens | 0.48 | 0.44 | 0.38 | 0.34 | 0.20 | 0.20 |
| LSD | 0.34 | 0.26 | 0.25 | 0.14 | 0.12 | 0.09 |
| PCP | 0.20 | 0.18 | 0.12 | 0.10 | 0.03 | 0.07 |
| Ecstasy | 0.40 | 0.31 | 0.32 | 0.24 | 0.17 | 0.13 |
| Inhalants | 0.49 | 0.46 | 0.33 | 0.32 | 0.11 | 0.14 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.63 | 0.62 | 0.54 | 0.54 | 0.38 | 0.37 |
| Pain Relievers | 0.59 | 0.57 | 0.51 | 0.49 | 0.35 | 0.33 |
| Tranquilizers | 0.37 | 0.39 | 0.31 | 0.31 | 0.18 | 0.20 |
| Stimulants | 0.41 | 0.37 | 0.32 | 0.27 | 0.20 | 0.18 |
| Methamphetamine | 0.27 | 0.21 | 0.21 | 0.15 | 0.13 | 0.11 |
| Sedatives | 0.18 | 0.18 | 0.14 | 0.12 | 0.07 | 0.08 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.73 | 0.72 | 0.62 | 0.59 | 0.44 | 0.40 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.7P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.3612 | | 0.4236 | | 0.4652 | |
| Marijuana and Hashish | 0.1807 | | 0.3677 | | 0.8792 | |
| Cocaine | 0.1865 | | 0.0381 | | 0.9046 | |
| Crack | 0.5923 | | 0.7241 | | 0.2567 | |
| Heroin | 0.4989 | | 0.1976 | | 0.7975 | |
| Hallucinogens | 0.0016 | | 0.0019 | | 0.9297 | |
| LSD | 0.0000 | | 0.0000 | | 0.6063 | |
| PCP | 0.1839 | | 0.2567 | | 0.0315 | |
| Ecstasy | 0.0000 | | 0.0000 | | 0.1256 | |
| Inhalants | 1.0000 | | 0.9805 | | 0.0297 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.2443 | | 0.3675 | | 0.8364 | |
| Pain Relievers | 0.2503 | | 0.5431 | | 0.9428 | |
| Tranquilizers | 0.4801 | | 0.7832 | | 0.3743 | |
| Stimulants | 0.3953 | | 0.1307 | | 0.9056 | |
| Methamphetamine | 0.4141 | | 0.0787 | | 0.8771 | |
| Sedatives | 0.9312 | | 0.3106 | | 0.5389 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.2657 | | 0.2606 | | 0.4794 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.8C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 89 | 96 | 92 | 97 | 79 | 80 |
| Marijuana and Hashish | 93 | 99 | 91 | 96 | 76 | 76 |
| Cocaine | 57 | 60 | 42 | 41 | 22 | 25 |
| Crack | 28 | 32 | 14 | 18 | 4 | 12 |
| Heroin | 17 | 18 | 11 | 9 | 8 | 6 |
| Hallucinogens | 75 | 75 | 53 | 51 | 26 | 28 |
| LSD | 59 | 53 | 32 | 21 | 6 | 11 |
| PCP | 28 | 29 | 12 | 12 | 3 | 4 |
| Ecstasy | 63 | 62 | 45 | 34 | 20 | 15 |
| Inhalants | 63 | 62 | 32 | 28 | 12 | 14 |
| Nonmedical Use of Any Psychotherapeutic ² | 74 | 81 | 58 | 65 | 38 | 45 |
| Pain Relievers | 69 | 77 | 57 | 62 | 33 | 41 |
| Tranquilizers | 49 | 55 | 34 | 42 | 22 | 21 |
| Stimulants | 53 | 53 | 38 | 34 | 19 | 20 |
| Methamphetamine | 32 | 36 | 22 | 21 | 11 | 13 |
| Sedatives | 22 | 19 | 10 | 11 | 6 | 6 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 85 | 89 | 71 | 76 | 46 | 53 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.8D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.72 | 0.77 | 0.74 | 0.77 | 0.63 | 0.64 |
| Marijuana and Hashish | 0.75 | 0.79 | 0.73 | 0.77 | 0.61 | 0.61 |
| Cocaine | 0.46 | 0.48 | 0.34 | 0.33 | 0.18 | 0.20 |
| Crack | 0.22 | 0.26 | 0.11 | 0.14 | 0.03 | 0.10 |
| Heroin | 0.14 | 0.15 | 0.09 | 0.07 | 0.07 | 0.05 |
| Hallucinogens | 0.61 | 0.60 | 0.43 | 0.41 | 0.21 | 0.22 |
| LSD | 0.48 | 0.43 | 0.26 | 0.17 | 0.05 | 0.09 |
| PCP | 0.23 | 0.23 | 0.09 | 0.10 | 0.02 | 0.03 |
| Ecstasy | 0.51 | 0.50 | 0.36 | 0.27 | 0.16 | 0.12 |
| Inhalants | 0.50 | 0.49 | 0.25 | 0.23 | 0.10 | 0.11 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.60 | 0.64 | 0.47 | 0.52 | 0.30 | 0.36 |
| Pain Relievers | 0.55 | 0.62 | 0.46 | 0.49 | 0.27 | 0.33 |
| Tranquilizers | 0.40 | 0.44 | 0.27 | 0.33 | 0.18 | 0.17 |
| Stimulants | 0.43 | 0.43 | 0.30 | 0.27 | 0.15 | 0.16 |
| Methamphetamine | 0.26 | 0.29 | 0.17 | 0.17 | 0.09 | 0.10 |
| Sedatives | 0.18 | 0.15 | 0.08 | 0.09 | 0.05 | 0.05 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.68 | 0.71 | 0.57 | 0.61 | 0.37 | 0.42 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.8P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.6041 | | 0.9722 | | 0.3370 | |
| Marijuana and Hashish | 0.9499 | | 0.7328 | | 0.6907 | |
| Cocaine | 0.9659 | | 0.7380 | | 0.3684 | |
| Crack | 0.5688 | | 0.4103 | | 0.2173 | |
| Heroin | 0.7051 | | 0.1783 | | 0.5025 | |
| Hallucinogens | 0.0193 | | 0.0102 | | 0.5826 | |
| LSD | 0.0001 | | 0.0003 | | 0.0466 | |
| PCP | 0.9888 | | 0.8269 | | 0.3823 | |
| Ecstasy | 0.1161 | | 0.0000 | | 0.0195 | |
| Inhalants | 0.9792 | | 0.6426 | | 0.7704 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.0521 | | 0.1638 | | 0.0507 | |
| Pain Relievers | 0.0159 | | 0.1216 | | 0.0260 | |
| Tranquilizers | 0.0861 | | 0.1394 | | 0.8426 | |
| Stimulants | 0.3052 | | 0.3731 | | 0.6705 | |
| Methamphetamine | 0.2652 | | 0.3706 | | 0.9289 | |
| Sedatives | 0.1956 | | 0.7229 | | 0.8779 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.8757 | | 0.7038 | | 0.0324 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.9C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 123 | 113 | 108 | 105 | 84 | 88 |
| Marijuana and Hashish | 122 | 114 | 100 | 99 | 76 | 79 |
| Cocaine | 88 | 87 | 55 | 58 | 28 | 34 |
| Crack | 42 | 45 | 19 | 17 | 8 | 5 |
| Heroin | 29 | 30 | 12 | 10 | 5 | 4 |
| Hallucinogens | 105 | 104 | 60 | 52 | 24 | 26 |
| LSD | 90 | 88 | 24 | 19 | 6 | 8 |
| PCP | 34 | 38 | 7 | 13 | 4 | 10 |
| Ecstasy | 85 | 86 | 47 | 38 | 21 | 17 |
| Inhalants | 81 | 77 | 25 | 25 | 14 | 13 |
| Nonmedical Use of Any Psychotherapeutic ² | 107 | 99 | 76 | 73 | 46 | 47 |
| Pain Relievers | 96 | 93 | 64 | 68 | 38 | 41 |
| Tranquilizers | 76 | 78 | 45 | 49 | 27 | 28 |
| Stimulants | 67 | 70 | 40 | 38 | 23 | 23 |
| Methamphetamine | 53 | 52 | 31 | 28 | 16 | 17 |
| Sedatives | 31 | 30 | 14 | 14 | 7 | 9 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 121 | 111 | 89 | 88 | 55 | 57 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.9D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.66 | 0.59 | 0.58 | 0.55 | 0.45 | 0.46 |
| Marijuana and Hashish | 0.66 | 0.59 | 0.54 | 0.52 | 0.41 | 0.41 |
| Cocaine | 0.47 | 0.45 | 0.29 | 0.30 | 0.15 | 0.18 |
| Crack | 0.22 | 0.23 | 0.10 | 0.09 | 0.04 | 0.03 |
| Heroin | 0.16 | 0.16 | 0.06 | 0.05 | 0.03 | 0.02 |
| Hallucinogens | 0.56 | 0.54 | 0.32 | 0.27 | 0.13 | 0.13 |
| LSD | 0.48 | 0.46 | 0.13 | 0.10 | 0.03 | 0.04 |
| PCP | 0.18 | 0.20 | 0.04 | 0.07 | 0.02 | 0.05 |
| Ecstasy | 0.46 | 0.45 | 0.25 | 0.20 | 0.11 | 0.09 |
| Inhalants | 0.44 | 0.40 | 0.13 | 0.13 | 0.08 | 0.07 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.58 | 0.52 | 0.41 | 0.38 | 0.25 | 0.25 |
| Pain Relievers | 0.52 | 0.48 | 0.35 | 0.35 | 0.21 | 0.21 |
| Tranquilizers | 0.41 | 0.41 | 0.24 | 0.26 | 0.14 | 0.15 |
| Stimulants | 0.36 | 0.36 | 0.22 | 0.20 | 0.13 | 0.12 |
| Methamphetamine | 0.28 | 0.27 | 0.16 | 0.15 | 0.09 | 0.09 |
| Sedatives | 0.17 | 0.16 | 0.07 | 0.07 | 0.04 | 0.05 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.65 | 0.58 | 0.48 | 0.46 | 0.30 | 0.30 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.9P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.4206 | | 0.1011 | | 0.5544 | |
| Marijuana and Hashish | 0.8086 | | 0.0189 | | 0.2407 | |
| Cocaine | 0.2080 | | 0.4959 | | 0.6375 | |
| Crack | 0.4915 | | 0.5085 | | 0.1343 | |
| Heroin | 0.5518 | | 0.5757 | | 0.2210 | |
| Hallucinogens | 0.6392 | | 0.0000 | | 0.3212 | |
| LSD | 0.0193 | | 0.0060 | | 0.5776 | |
| PCP | 0.0710 | | 0.0975 | | 0.3103 | |
| Ecstasy | 0.7478 | | 0.0000 | | 0.0044 | |
| Inhalants | 0.0272 | | 0.6659 | | 0.2320 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.1663 | | 0.7544 | | 0.2051 | |
| Pain Relievers | 0.0613 | | 0.5525 | | 0.1486 | |
| Tranquilizers | 0.0451 | | 0.3752 | | 0.5812 | |
| Stimulants | 0.4035 | | 0.8569 | | 0.6138 | |
| Methamphetamine | 0.1909 | | 0.9116 | | 0.3420 | |
| Sedatives | 0.1662 | | 0.4590 | | 0.5765 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.8935 | | 0.1479 | | 0.9739 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.10C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 278 | 285 | 239 | 233 | 184 | 183 |
| Marijuana and Hashish | 279 | 290 | 216 | 210 | 158 | 153 |
| Cocaine | 223 | 225 | 110 | 127 | 65 | 70 |
| Crack | 124 | 134 | 61 | 55 | 20 | 37 |
| Heroin | 70 | 68 | 21 | 22 | 17 | 14 |
| Hallucinogens | 247 | 239 | 77 | 87 | 39 | 46 |
| LSD | 217 | 192 | 28 | 20 | 11 | 5 |
| PCP | 112 | 74 | 16 | 6 | 13 | 4 |
| Ecstasy | 159 | 184 | 69 | 72 | 30 | 41 |
| Inhalants | 205 | 188 | 33 | 28 | 22 | 9 |
| Nonmedical Use of Any Psychotherapeutic ² | 284 | 241 | 172 | 160 | 105 | 116 |
| Pain Relievers | 220 | 214 | 142 | 141 | 89 | 101 |
| Tranquilizers | 174 | 180 | 89 | 82 | 56 | 42 |
| Stimulants | 193 | 162 | 82 | 62 | 52 | 53 |
| Methamphetamine | 163 | 139 | 69 | 52 | 42 | 44 |
| Sedatives | 89 | 71 | 35 | 33 | 27 | 26 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 291 | 279 | 183 | 183 | 128 | 127 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.10D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.79 | 0.82 | 0.68 | 0.67 | 0.52 | 0.52 |
| Marijuana and Hashish | 0.79 | 0.83 | 0.61 | 0.60 | 0.45 | 0.44 |
| Cocaine | 0.63 | 0.64 | 0.31 | 0.36 | 0.19 | 0.20 |
| Crack | 0.35 | 0.38 | 0.17 | 0.16 | 0.06 | 0.11 |
| Heroin | 0.20 | 0.20 | 0.06 | 0.06 | 0.05 | 0.04 |
| Hallucinogens | 0.70 | 0.68 | 0.22 | 0.25 | 0.11 | 0.13 |
| LSD | 0.62 | 0.55 | 0.08 | 0.06 | 0.03 | 0.02 |
| PCP | 0.32 | 0.21 | 0.05 | 0.02 | 0.04 | 0.01 |
| Ecstasy | 0.45 | 0.53 | 0.20 | 0.21 | 0.09 | 0.12 |
| Inhalants | 0.58 | 0.54 | 0.09 | 0.08 | 0.06 | 0.03 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.81 | 0.69 | 0.49 | 0.46 | 0.30 | 0.33 |
| Pain Relievers | 0.63 | 0.61 | 0.40 | 0.40 | 0.25 | 0.29 |
| Tranquilizers | 0.49 | 0.52 | 0.25 | 0.24 | 0.16 | 0.12 |
| Stimulants | 0.55 | 0.46 | 0.23 | 0.18 | 0.15 | 0.15 |
| Methamphetamine | 0.46 | 0.40 | 0.20 | 0.15 | 0.12 | 0.12 |
| Sedatives | 0.25 | 0.20 | 0.10 | 0.09 | 0.08 | 0.07 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.83 | 0.80 | 0.52 | 0.52 | 0.36 | 0.36 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.10P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.3408 | | 0.9596 | | 0.7678 | |
| Marijuana and Hashish | 0.3102 | | 0.8028 | | 0.2874 | |
| Cocaine | 0.6059 | | 0.3710 | | 0.3013 | |
| Crack | 0.7965 | | 0.5451 | | 0.1042 | |
| Heroin | 0.5957 | | 0.9682 | | 0.9099 | |
| Hallucinogens | 0.6844 | | 0.8346 | | 0.6866 | |
| LSD | 0.7357 | | 0.2912 | | 0.6550 | |
| PCP | 0.1585 | | 0.2117 | | 0.2705 | |
| Ecstasy | 0.0213 | | 0.2690 | | 0.9551 | |
| Inhalants | 0.5741 | | 0.3679 | | 0.1900 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.7980 | | 0.8124 | | 0.6414 | |
| Pain Relievers | 0.1068 | | 0.5625 | | 0.9334 | |
| Tranquilizers | 0.0998 | | 0.3286 | | 0.2113 | |
| Stimulants | 0.1516 | | 0.0858 | | 0.7638 | |
| Methamphetamine | 0.3659 | | 0.4911 | | 0.9134 | |
| Sedatives | 0.2404 | | 0.3560 | | 0.1069 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.3011 | | 0.5964 | | 0.7566 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 869 | 793 | 426 | 410 | 298 | 300 |
| Marijuana and Hashish | 822 | 761 | 338 | 305 | 236 | 228 |
| Cocaine | 531 | 520 | 164 | 153 | 96 | 113 |
| Crack | 265 | 220 | 106 | 104 | 63 | 80 |
| Heroin | 207 | 183 | 47 | 40 | 37 | 22 |
| Hallucinogens | 491 | 458 | 77 | 57 | 54 | 28 |
| LSD | 426 | 410 | 21 | 20 | * | * |
| PCP | 261 | 253 | * | * | * | * |
| Ecstasy | 167 | 168 | 61 | 45 | 43 | 23 |
| Inhalants | 382 | 344 | 58 | 47 | 52 | 31 |
| Nonmedical Use of Any Psychotherapeutic ² | 609 | 567 | 273 | 278 | 180 | 195 |
| Pain Relievers | 460 | 426 | 212 | 234 | 139 | 157 |
| Tranquilizers | 433 | 366 | 184 | 154 | 110 | 104 |
| Stimulants | 432 | 412 | 95 | 86 | 62 | 63 |
| Methamphetamine | 315 | 334 | 62 | 66 | 51 | 55 |
| Sedatives | 347 | 305 | 107 | 75 | 57 | 46 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 703 | 664 | 313 | 304 | 210 | 227 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.60 | 0.54 | 0.30 | 0.28 | 0.21 | 0.21 |
| Marijuana and Hashish | 0.57 | 0.52 | 0.23 | 0.21 | 0.16 | 0.16 |
| Cocaine | 0.37 | 0.36 | 0.11 | 0.11 | 0.07 | 0.08 |
| Crack | 0.18 | 0.15 | 0.07 | 0.07 | 0.04 | 0.06 |
| Heroin | 0.14 | 0.13 | 0.03 | 0.03 | 0.03 | 0.02 |
| Hallucinogens | 0.34 | 0.31 | 0.05 | 0.04 | 0.04 | 0.02 |
| LSD | 0.30 | 0.28 | 0.01 | 0.01 | * | * |
| PCP | 0.18 | 0.17 | * | * | * | * |
| Ecstasy | 0.12 | 0.11 | 0.04 | 0.03 | 0.03 | 0.02 |
| Inhalants | 0.26 | 0.24 | 0.04 | 0.03 | 0.04 | 0.02 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.42 | 0.39 | 0.19 | 0.19 | 0.13 | 0.13 |
| Pain Relievers | 0.32 | 0.29 | 0.15 | 0.16 | 0.10 | 0.11 |
| Tranquilizers | 0.30 | 0.25 | 0.13 | 0.11 | 0.08 | 0.07 |
| Stimulants | 0.30 | 0.28 | 0.07 | 0.06 | 0.04 | 0.04 |
| Methamphetamine | 0.22 | 0.23 | 0.04 | 0.05 | 0.04 | 0.04 |
| Sedatives | 0.24 | 0.21 | 0.07 | 0.05 | 0.04 | 0.03 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.49 | 0.45 | 0.22 | 0.21 | 0.15 | 0.16 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.3668 | | 0.7361 | | 0.4919 | |
| Marijuana and Hashish | 0.2404 | | 0.4933 | | 0.5616 | |
| Cocaine | 0.3908 | | 0.6506 | | 0.5586 | |
| Crack | 0.0922 | | 0.4602 | | 0.6681 | |
| Heroin | 0.9414 | | 0.4247 | | 0.3763 | |
| Hallucinogens | 0.6506 | | 0.0757 | | 0.1408 | |
| LSD | 0.2998 | | 0.4400 | | * | |
| PCP | 0.4575 | | * | | * | |
| Ecstasy | 0.0943 | | 0.1791 | | 0.2933 | |
| Inhalants | 0.6072 | | 0.8774 | | 0.5385 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.6612 | | 0.9786 | | 0.8033 | |
| Pain Relievers | 0.6667 | | 0.4854 | | 0.7811 | |
| Tranquilizers | 0.9259 | | 0.6215 | | 0.8501 | |
| Stimulants | 0.8839 | | 0.2774 | | 0.6470 | |
| Methamphetamine | 0.7180 | | 0.2872 | | 0.8292 | |
| Sedatives | 0.4736 | | 0.4573 | | 0.2122 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.6361 | | 0.4708 | | 0.7422 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.12C Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 963 | 902 | 565 | 540 | 379 | 396 |
| Marijuana and Hashish | 915 | 865 | 459 | 437 | 313 | 317 |
| Cocaine | 584 | 564 | 217 | 225 | 126 | 146 |
| Crack | 307 | 263 | 124 | 122 | 67 | 91 |
| Heroin | 222 | 204 | 54 | 47 | 41 | 27 |
| Hallucinogens | 599 | 584 | 144 | 140 | 78 | 68 |
| LSD | 513 | 494 | 58 | 41 | 13 | 15 |
| PCP | 281 | 270 | 21 | 20 | 14 | 11 |
| Ecstasy | 296 | 281 | 122 | 101 | 62 | 53 |
| Inhalants | 472 | 435 | 80 | 66 | 59 | 37 |
| Nonmedical Use of Any Psychotherapeutic ² | 730 | 635 | 367 | 351 | 226 | 249 |
| Pain Relievers | 565 | 510 | 293 | 298 | 182 | 205 |
| Tranquilizers | 493 | 433 | 219 | 192 | 133 | 116 |
| Stimulants | 488 | 457 | 142 | 124 | 88 | 90 |
| Methamphetamine | 368 | 369 | 103 | 93 | 70 | 73 |
| Sedatives | 370 | 322 | 115 | 83 | 64 | 54 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 813 | 748 | 415 | 408 | 270 | 286 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.12D Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.46 | 0.42 | 0.27 | 0.25 | 0.18 | 0.19 |
| Marijuana and Hashish | 0.44 | 0.41 | 0.22 | 0.21 | 0.15 | 0.15 |
| Cocaine | 0.28 | 0.27 | 0.10 | 0.11 | 0.06 | 0.07 |
| Crack | 0.15 | 0.12 | 0.06 | 0.06 | 0.03 | 0.04 |
| Heroin | 0.11 | 0.10 | 0.03 | 0.02 | 0.02 | 0.01 |
| Hallucinogens | 0.28 | 0.27 | 0.07 | 0.07 | 0.04 | 0.03 |
| LSD | 0.24 | 0.23 | 0.03 | 0.02 | 0.01 | 0.01 |
| PCP | 0.13 | 0.13 | 0.01 | 0.01 | 0.01 | 0.01 |
| Ecstasy | 0.14 | 0.13 | 0.06 | 0.05 | 0.03 | 0.03 |
| Inhalants | 0.22 | 0.20 | 0.04 | 0.03 | 0.03 | 0.02 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.35 | 0.30 | 0.17 | 0.17 | 0.11 | 0.12 |
| Pain Relievers | 0.27 | 0.24 | 0.14 | 0.14 | 0.09 | 0.10 |
| Tranquilizers | 0.23 | 0.20 | 0.10 | 0.09 | 0.06 | 0.05 |
| Stimulants | 0.23 | 0.21 | 0.07 | 0.06 | 0.04 | 0.04 |
| Methamphetamine | 0.17 | 0.17 | 0.05 | 0.04 | 0.03 | 0.03 |
| Sedatives | 0.18 | 0.15 | 0.05 | 0.04 | 0.03 | 0.03 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.39 | 0.35 | 0.20 | 0.19 | 0.13 | 0.13 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.12P Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.5013 | | 0.5876 | | 0.7561 | |
| Marijuana and Hashish | 0.4596 | | 0.3442 | | 0.9449 | |
| Cocaine | 0.4223 | | 0.9550 | | 0.2107 | |
| Crack | 0.1706 | | 0.3562 | | 0.8271 | |
| Heroin | 0.8670 | | 0.2823 | | 0.2646 | |
| Hallucinogens | 0.8638 | | 0.0004 | | 0.1025 | |
| LSD | 0.9113 | | 0.0001 | | 0.1444 | |
| PCP | 0.3558 | | 0.9285 | | 0.9865 | |
| Ecstasy | 0.0312 | | 0.0000 | | 0.0279 | |
| Inhalants | 0.7991 | | 0.5630 | | 0.2223 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.3485 | | 0.7429 | | 0.8811 | |
| Pain Relievers | 0.0866 | | 0.1889 | | 0.3265 | |
| Tranquilizers | 0.2756 | | 0.6296 | | 0.8539 | |
| Stimulants | 0.5113 | | 0.0513 | | 0.6452 | |
| Methamphetamine | 0.7383 | | 0.1943 | | 0.9817 | |
| Sedatives | 0.2616 | | 0.2878 | | 0.0857 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.9374 | | 0.5770 | | 0.9904 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.13C Illicit Drug Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 608 | 614 | 393 | 384 | 291 | 284 |
| Marijuana and Hashish | 581 | 604 | 326 | 331 | 247 | 249 |
| Cocaine | 417 | 449 | 188 | 191 | 106 | 126 |
| Crack | 238 | 223 | 112 | 105 | 56 | 79 |
| Heroin | 203 | 159 | 52 | 44 | 40 | 23 |
| Hallucinogens | 430 | 441 | 117 | 121 | 69 | 56 |
| LSD | 381 | 384 | 49 | 34 | 14 | 16 |
| PCP | 223 | 219 | 20 | 17 | 14 | 11 |
| Ecstasy | 211 | 221 | 99 | 88 | 54 | 36 |
| Inhalants | 386 | 377 | 77 | 65 | 58 | 36 |
| Nonmedical Use of Any Psychotherapeutic ² | 494 | 468 | 252 | 236 | 155 | 155 |
| Pain Relievers | 396 | 367 | 208 | 207 | 130 | 135 |
| Tranquilizers | 333 | 307 | 160 | 137 | 79 | 71 |
| Stimulants | 364 | 349 | 111 | 85 | 69 | 59 |
| Methamphetamine | 277 | 297 | 83 | 69 | 59 | 55 |
| Sedatives | 260 | 259 | 91 | 43 | 39 | 20 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 549 | 559 | 309 | 283 | 206 | 196 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.13D Illicit Drug Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.54 | 0.53 | 0.35 | 0.33 | 0.26 | 0.25 |
| Marijuana and Hashish | 0.51 | 0.53 | 0.29 | 0.29 | 0.22 | 0.22 |
| Cocaine | 0.37 | 0.39 | 0.17 | 0.17 | 0.09 | 0.11 |
| Crack | 0.21 | 0.19 | 0.10 | 0.09 | 0.05 | 0.07 |
| Heroin | 0.18 | 0.14 | 0.05 | 0.04 | 0.04 | 0.02 |
| Hallucinogens | 0.38 | 0.38 | 0.10 | 0.10 | 0.06 | 0.05 |
| LSD | 0.34 | 0.33 | 0.04 | 0.03 | 0.01 | 0.01 |
| PCP | 0.20 | 0.19 | 0.02 | 0.01 | 0.01 | 0.01 |
| Ecstasy | 0.19 | 0.19 | 0.09 | 0.08 | 0.05 | 0.03 |
| Inhalants | 0.34 | 0.33 | 0.07 | 0.06 | 0.05 | 0.03 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.44 | 0.41 | 0.22 | 0.20 | 0.14 | 0.13 |
| Pain Relievers | 0.35 | 0.32 | 0.18 | 0.18 | 0.11 | 0.12 |
| Tranquilizers | 0.29 | 0.27 | 0.14 | 0.12 | 0.07 | 0.06 |
| Stimulants | 0.32 | 0.30 | 0.10 | 0.07 | 0.06 | 0.05 |
| Methamphetamine | 0.24 | 0.26 | 0.07 | 0.06 | 0.05 | 0.05 |
| Sedatives | 0.23 | 0.23 | 0.08 | 0.04 | 0.03 | 0.02 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.48 | 0.49 | 0.27 | 0.25 | 0.18 | 0.17 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.13P Illicit Drug Use in Lifetime, Past Year, and Past Month among Males: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.5033 | | 0.4871 | | 0.3720 | |
| Marijuana and Hashish | 0.4525 | | 0.2798 | | 0.9479 | |
| Cocaine | 0.4846 | | 0.6979 | | 0.3145 | |
| Crack | 0.6410 | | 0.3237 | | 0.5132 | |
| Heroin | 0.4888 | | 0.2873 | | 0.2114 | |
| Hallucinogens | 0.8492 | | 0.0115 | | 0.1769 | |
| LSD | 0.8619 | | 0.0000 | | 0.1571 | |
| PCP | 0.5257 | | 0.8298 | | 0.9731 | |
| Ecstasy | 0.3535 | | 0.0000 | | 0.0016 | |
| Inhalants | 0.5586 | | 0.1630 | | 0.2400 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.7947 | | 0.9904 | | 1.0000 | |
| Pain Relievers | 0.4749 | | 0.6428 | | 0.6462 | |
| Tranquilizers | 0.9742 | | 0.7174 | | 0.8379 | |
| Stimulants | 0.2588 | | 0.0560 | | 0.7714 | |
| Methamphetamine | 0.7531 | | 0.4410 | | 0.5376 | |
| Sedatives | 0.7176 | | 0.0887 | | 0.0270 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.7255 | | 0.5080 | | 0.8116 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.14C Illicit Drug Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 638 | 625 | 366 | 329 | 232 | 244 |
| Marijuana and Hashish | 607 | 587 | 279 | 241 | 175 | 178 |
| Cocaine | 383 | 353 | 107 | 109 | 63 | 64 |
| Crack | 179 | 144 | 61 | 59 | 37 | 31 |
| Heroin | 101 | 112 | 18 | 18 | 12 | 14 |
| Hallucinogens | 389 | 348 | 85 | 74 | 43 | 42 |
| LSD | 324 | 283 | 35 | 26 | 11 | 8 |
| PCP | 164 | 149 | 14 | 14 | 7 | 7 |
| Ecstasy | 171 | 165 | 71 | 55 | 32 | 38 |
| Inhalants | 247 | 231 | 47 | 47 | 28 | 27 |
| Nonmedical Use of Any Psychotherapeutic ² | 485 | 448 | 257 | 247 | 168 | 175 |
| Pain Relievers | 368 | 365 | 206 | 209 | 130 | 144 |
| Tranquilizers | 312 | 307 | 150 | 139 | 105 | 95 |
| Stimulants | 319 | 313 | 86 | 85 | 56 | 62 |
| Methamphetamine | 225 | 226 | 61 | 58 | 41 | 41 |
| Sedatives | 239 | 204 | 71 | 72 | 51 | 50 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 539 | 527 | 278 | 269 | 181 | 186 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.14D Illicit Drug Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.52 | 0.51 | 0.30 | 0.27 | 0.19 | 0.20 |
| Marijuana and Hashish | 0.50 | 0.48 | 0.23 | 0.20 | 0.14 | 0.14 |
| Cocaine | 0.32 | 0.29 | 0.09 | 0.09 | 0.05 | 0.05 |
| Crack | 0.15 | 0.12 | 0.05 | 0.05 | 0.03 | 0.03 |
| Heroin | 0.08 | 0.09 | 0.02 | 0.01 | 0.01 | 0.01 |
| Hallucinogens | 0.32 | 0.28 | 0.07 | 0.06 | 0.04 | 0.03 |
| LSD | 0.27 | 0.23 | 0.03 | 0.02 | 0.01 | 0.01 |
| PCP | 0.13 | 0.12 | 0.01 | 0.01 | 0.01 | 0.01 |
| Ecstasy | 0.14 | 0.13 | 0.06 | 0.04 | 0.03 | 0.03 |
| Inhalants | 0.20 | 0.19 | 0.04 | 0.04 | 0.02 | 0.02 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.40 | 0.37 | 0.21 | 0.20 | 0.14 | 0.14 |
| Pain Relievers | 0.30 | 0.30 | 0.17 | 0.17 | 0.11 | 0.12 |
| Tranquilizers | 0.26 | 0.25 | 0.12 | 0.11 | 0.09 | 0.08 |
| Stimulants | 0.26 | 0.26 | 0.07 | 0.07 | 0.05 | 0.05 |
| Methamphetamine | 0.19 | 0.18 | 0.05 | 0.05 | 0.03 | 0.03 |
| Sedatives | 0.20 | 0.17 | 0.06 | 0.06 | 0.04 | 0.04 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.44 | 0.43 | 0.23 | 0.22 | 0.15 | 0.15 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.14P Illicit Drug Use in Lifetime, Past Year, and Past Month among Females: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.8097 | | 0.7854 | | 0.7633 | |
| Marijuana and Hashish | 0.9791 | | 0.4179 | | 0.7237 | |
| Cocaine | 0.7219 | | 0.7575 | | 0.3853 | |
| Crack | 0.1014 | | 0.8692 | | 0.5010 | |
| Heroin | 0.1362 | | 0.4127 | | 0.6241 | |
| Hallucinogens | 0.3856 | | 0.0001 | | 0.4484 | |
| LSD | 0.2906 | | 0.0001 | | 0.4701 | |
| PCP | 0.3728 | | 0.4907 | | 0.8332 | |
| Ecstasy | 0.1292 | | 0.0000 | | 0.8885 | |
| Inhalants | 0.5869 | | 0.1101 | | 0.8203 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.3316 | | 0.6673 | | 0.8094 | |
| Pain Relievers | 0.0837 | | 0.1424 | | 0.3572 | |
| Tranquilizers | 0.0801 | | 0.7157 | | 0.8367 | |
| Stimulants | 0.8818 | | 0.1235 | | 0.8774 | |
| Methamphetamine | 0.7907 | | 0.0627 | | 0.4229 | |
| Sedatives | 0.1791 | | 0.8931 | | 0.5131 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.7763 | | 0.8488 | | 0.8736 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.15C Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 75 | 71 | 65 | 64 | 52 | 47 |
| Marijuana and Hashish | 64 | 63 | 55 | 53 | 44 | 43 |
| Cocaine | 26 | 24 | 23 | 19 | 12 | 10 |
| Crack | 11 | 13 | 8 | 9 | 4 | 7 |
| Heroin | 7 | 8 | 5 | 6 | 3 | 5 |
| Hallucinogens | 35 | 33 | 29 | 25 | 14 | 15 |
| LSD | 25 | 19 | 18 | 11 | 8 | 7 |
| PCP | 13 | 13 | 8 | 9 | 4 | 5 |
| Ecstasy | 27 | 21 | 22 | 15 | 10 | 8 |
| Inhalants | 46 | 49 | 31 | 33 | 16 | 16 |
| Nonmedical Use of Any Psychotherapeutic ² | 51 | 49 | 42 | 43 | 29 | 28 |
| Pain Relievers | 47 | 47 | 38 | 40 | 26 | 26 |
| Tranquilizers | 27 | 26 | 22 | 22 | 11 | 14 |
| Stimulants | 30 | 27 | 24 | 20 | 15 | 13 |
| Methamphetamine | 16 | 16 | 14 | 11 | 11 | 8 |
| Sedatives | 13 | 16 | 10 | 13 | 3 | 8 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 63 | 63 | 52 | 53 | 36 | 32 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.15D Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.59 | 0.56 | 0.52 | 0.50 | 0.41 | 0.37 |
| Marijuana and Hashish | 0.50 | 0.49 | 0.43 | 0.42 | 0.35 | 0.34 |
| Cocaine | 0.21 | 0.18 | 0.18 | 0.15 | 0.10 | 0.08 |
| Crack | 0.09 | 0.10 | 0.06 | 0.07 | 0.03 | 0.05 |
| Heroin | 0.05 | 0.06 | 0.04 | 0.05 | 0.02 | 0.04 |
| Hallucinogens | 0.27 | 0.26 | 0.23 | 0.20 | 0.11 | 0.11 |
| LSD | 0.20 | 0.15 | 0.15 | 0.09 | 0.07 | 0.06 |
| PCP | 0.10 | 0.10 | 0.06 | 0.07 | 0.03 | 0.04 |
| Ecstasy | 0.21 | 0.16 | 0.18 | 0.12 | 0.08 | 0.06 |
| Inhalants | 0.37 | 0.39 | 0.25 | 0.26 | 0.13 | 0.13 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.41 | 0.38 | 0.33 | 0.33 | 0.23 | 0.22 |
| Pain Relievers | 0.37 | 0.37 | 0.30 | 0.31 | 0.21 | 0.20 |
| Tranquilizers | 0.21 | 0.20 | 0.17 | 0.17 | 0.08 | 0.11 |
| Stimulants | 0.24 | 0.21 | 0.19 | 0.15 | 0.12 | 0.11 |
| Methamphetamine | 0.13 | 0.12 | 0.11 | 0.09 | 0.08 | 0.06 |
| Sedatives | 0.10 | 0.13 | 0.08 | 0.10 | 0.02 | 0.06 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.50 | 0.50 | 0.41 | 0.41 | 0.28 | 0.25 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.15P Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.3000 | | 0.3141 | | 0.0963 | |
| Marijuana and Hashish | 0.0570 | | 0.0932 | | 0.2487 | |
| Cocaine | 0.5256 | | 0.1385 | | 0.6086 | |
| Crack | 0.1219 | | 0.3526 | | 0.1580 | |
| Heroin | 0.9921 | | 0.7331 | | 0.8097 | |
| Hallucinogens | 0.0068 | | 0.0016 | | 0.7345 | |
| LSD | 0.0000 | | 0.0000 | | 0.3535 | |
| PCP | 0.6976 | | 0.8539 | | 0.7032 | |
| Ecstasy | 0.0003 | | 0.0000 | | 0.0889 | |
| Inhalants | 0.4671 | | 0.4217 | | 0.3599 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.7036 | | 0.8245 | | 0.6636 | |
| Pain Relievers | 0.8111 | | 0.8779 | | 0.8853 | |
| Tranquilizers | 0.7999 | | 0.8315 | | 0.0258 | |
| Stimulants | 0.9174 | | 0.1633 | | 0.8670 | |
| Methamphetamine | 0.6645 | | 0.4608 | | 0.7280 | |
| Sedatives | 0.0461 | | 0.1909 | | 0.0111 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.6417 | | 0.4162 | | 0.5389 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.16C Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 66 | 73 | 63 | 64 | 44 | 45 |
| Marijuana and Hashish | 59 | 62 | 53 | 55 | 35 | 39 |
| Cocaine | 24 | 26 | 23 | 19 | 13 | 12 |
| Crack | 13 | 10 | 11 | 9 | 7 | 3 |
| Heroin | 12 | 8 | 10 | 5 | 5 | 4 |
| Hallucinogens | 34 | 35 | 27 | 26 | 15 | 13 |
| LSD | 22 | 18 | 16 | 11 | 9 | 7 |
| PCP | 14 | 14 | 10 | 8 | 4 | 3 |
| Ecstasy | 28 | 24 | 22 | 17 | 12 | 10 |
| Inhalants | 45 | 45 | 30 | 31 | 16 | 19 |
| Nonmedical Use of Any Psychotherapeutic ² | 52 | 53 | 43 | 46 | 30 | 31 |
| Pain Relievers | 48 | 50 | 40 | 42 | 28 | 28 |
| Tranquilizers | 27 | 28 | 24 | 24 | 14 | 16 |
| Stimulants | 34 | 30 | 24 | 23 | 12 | 14 |
| Methamphetamine | 21 | 17 | 16 | 12 | 6 | 8 |
| Sedatives | 17 | 15 | 13 | 11 | 8 | 6 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 59 | 66 | 51 | 56 | 35 | 37 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.16D Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.54 | 0.60 | 0.52 | 0.52 | 0.37 | 0.37 |
| Marijuana and Hashish | 0.49 | 0.51 | 0.44 | 0.45 | 0.29 | 0.32 |
| Cocaine | 0.20 | 0.21 | 0.19 | 0.16 | 0.11 | 0.10 |
| Crack | 0.11 | 0.08 | 0.09 | 0.07 | 0.06 | 0.03 |
| Heroin | 0.10 | 0.07 | 0.08 | 0.04 | 0.04 | 0.03 |
| Hallucinogens | 0.28 | 0.28 | 0.23 | 0.22 | 0.12 | 0.11 |
| LSD | 0.18 | 0.15 | 0.13 | 0.09 | 0.07 | 0.06 |
| PCP | 0.12 | 0.12 | 0.08 | 0.06 | 0.03 | 0.02 |
| Ecstasy | 0.23 | 0.20 | 0.18 | 0.14 | 0.10 | 0.08 |
| Inhalants | 0.37 | 0.36 | 0.25 | 0.25 | 0.13 | 0.15 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.43 | 0.44 | 0.36 | 0.38 | 0.25 | 0.25 |
| Pain Relievers | 0.40 | 0.41 | 0.33 | 0.34 | 0.23 | 0.23 |
| Tranquilizers | 0.23 | 0.23 | 0.20 | 0.19 | 0.11 | 0.13 |
| Stimulants | 0.28 | 0.24 | 0.20 | 0.19 | 0.10 | 0.11 |
| Methamphetamine | 0.17 | 0.14 | 0.13 | 0.10 | 0.05 | 0.06 |
| Sedatives | 0.14 | 0.12 | 0.10 | 0.09 | 0.07 | 0.05 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.49 | 0.54 | 0.42 | 0.46 | 0.29 | 0.30 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.16P Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.8371 | | 0.9924 | | 0.7092 | |
| Marijuana and Hashish | 0.2646 | | 0.3523 | | 0.9754 | |
| Cocaine | 0.8386 | | 0.2835 | | 0.4438 | |
| Crack | 0.0302 | | 0.2349 | | 0.4954 | |
| Heroin | 0.3141 | | 0.2702 | | 0.3930 | |
| Hallucinogens | 0.2678 | | 0.1274 | | 0.7918 | |
| LSD | 0.0000 | | 0.0043 | | 0.9965 | |
| PCP | 0.2273 | | 0.1109 | | 0.2597 | |
| Ecstasy | 0.0076 | | 0.0002 | | 0.4267 | |
| Inhalants | 0.1538 | | 0.1115 | | 0.1244 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.5501 | | 0.9657 | | 0.8124 | |
| Pain Relievers | 0.7850 | | 0.7189 | | 0.8725 | |
| Tranquilizers | 0.5753 | | 0.8627 | | 0.8567 | |
| Stimulants | 0.1011 | | 0.1962 | | 0.4064 | |
| Methamphetamine | 0.0487 | | 0.0485 | | 0.2899 | |
| Sedatives | 0.1516 | | 0.0742 | | 0.0348 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.7988 | | 0.5681 | | 0.8045 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.17C Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 106 | 103 | 100 | 107 | 87 | 94 |
| Marijuana and Hashish | 112 | 105 | 95 | 104 | 83 | 87 |
| Cocaine | 77 | 80 | 54 | 55 | 30 | 36 |
| Crack | 41 | 42 | 19 | 19 | 8 | 12 |
| Heroin | 28 | 28 | 15 | 11 | 9 | 5 |
| Hallucinogens | 98 | 95 | 63 | 56 | 31 | 29 |
| LSD | 81 | 75 | 34 | 25 | 4 | 12 |
| PCP | 35 | 40 | 8 | 14 | 3 | 9 |
| Ecstasy | 81 | 74 | 51 | 34 | 25 | 14 |
| Inhalants | 82 | 81 | 33 | 31 | 15 | 16 |
| Nonmedical Use of Any Psychotherapeutic ² | 95 | 91 | 70 | 73 | 45 | 54 |
| Pain Relievers | 86 | 88 | 65 | 68 | 41 | 46 |
| Tranquilizers | 67 | 68 | 42 | 48 | 27 | 29 |
| Stimulants | 64 | 65 | 39 | 39 | 21 | 24 |
| Methamphetamine | 45 | 48 | 27 | 25 | 14 | 18 |
| Sedatives | 30 | 26 | 14 | 12 | 8 | 8 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 106 | 106 | 81 | 91 | 57 | 63 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.17D Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.68 | 0.65 | 0.65 | 0.67 | 0.56 | 0.59 |
| Marijuana and Hashish | 0.72 | 0.66 | 0.61 | 0.65 | 0.53 | 0.55 |
| Cocaine | 0.50 | 0.50 | 0.35 | 0.35 | 0.20 | 0.22 |
| Crack | 0.26 | 0.27 | 0.12 | 0.12 | 0.05 | 0.08 |
| Heroin | 0.18 | 0.18 | 0.09 | 0.07 | 0.06 | 0.03 |
| Hallucinogens | 0.63 | 0.60 | 0.41 | 0.35 | 0.20 | 0.18 |
| LSD | 0.52 | 0.47 | 0.22 | 0.15 | 0.03 | 0.08 |
| PCP | 0.23 | 0.25 | 0.05 | 0.09 | 0.02 | 0.06 |
| Ecstasy | 0.52 | 0.47 | 0.33 | 0.22 | 0.16 | 0.09 |
| Inhalants | 0.53 | 0.51 | 0.21 | 0.20 | 0.10 | 0.10 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.61 | 0.57 | 0.45 | 0.46 | 0.29 | 0.34 |
| Pain Relievers | 0.55 | 0.55 | 0.42 | 0.43 | 0.27 | 0.29 |
| Tranquilizers | 0.43 | 0.43 | 0.27 | 0.30 | 0.17 | 0.18 |
| Stimulants | 0.41 | 0.41 | 0.25 | 0.24 | 0.14 | 0.15 |
| Methamphetamine | 0.29 | 0.30 | 0.18 | 0.16 | 0.09 | 0.11 |
| Sedatives | 0.19 | 0.16 | 0.09 | 0.08 | 0.05 | 0.05 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.68 | 0.66 | 0.52 | 0.57 | 0.37 | 0.39 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.17P Illicit Drug Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.2623 | | 0.4792 | | 0.9409 | |
| Marijuana and Hashish | 0.2810 | | 0.3376 | | 0.9708 | |
| Cocaine | 0.3424 | | 0.7701 | | 0.3932 | |
| Crack | 0.7166 | | 0.9218 | | 0.9305 | |
| Heroin | 0.3631 | | 0.2607 | | 0.1053 | |
| Hallucinogens | 0.4830 | | 0.0030 | | 0.4000 | |
| LSD | 0.0614 | | 0.0105 | | 0.0009 | |
| PCP | 0.0769 | | 0.1426 | | 0.1016 | |
| Ecstasy | 0.3247 | | 0.0000 | | 0.0000 | |
| Inhalants | 0.2216 | | 0.1731 | | 0.5930 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.2040 | | 0.7592 | | 0.0305 | |
| Pain Relievers | 0.0147 | | 0.3556 | | 0.0902 | |
| Tranquilizers | 0.2039 | | 0.3430 | | 0.5002 | |
| Stimulants | 0.8544 | | 0.9348 | | 0.1972 | |
| Methamphetamine | 0.8815 | | 0.6751 | | 0.0240 | |
| Sedatives | 0.0728 | | 0.3750 | | 0.9676 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.9895 | | 0.6436 | | 0.2466 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.18C Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Standard Errors of Numbers in Thousands, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 97 | 103 | 88 | 93 | 68 | 72 |
| Marijuana and Hashish | 95 | 105 | 81 | 88 | 63 | 64 |
| Cocaine | 68 | 65 | 44 | 41 | 20 | 22 |
| Crack | 29 | 31 | 13 | 13 | 4 | 4 |
| Heroin | 20 | 20 | 8 | 8 | 4 | 4 |
| Hallucinogens | 84 | 82 | 46 | 42 | 19 | 21 |
| LSD | 67 | 64 | 22 | 14 | 7 | 4 |
| PCP | 27 | 28 | 10 | 10 | 5 | 5 |
| Ecstasy | 67 | 70 | 40 | 34 | 15 | 17 |
| Inhalants | 63 | 62 | 19 | 23 | 10 | 10 |
| Nonmedical Use of Any Psychotherapeutic ² | 84 | 88 | 60 | 67 | 39 | 40 |
| Pain Relievers | 76 | 81 | 54 | 61 | 34 | 37 |
| Tranquilizers | 53 | 65 | 36 | 41 | 21 | 20 |
| Stimulants | 57 | 60 | 33 | 35 | 21 | 16 |
| Methamphetamine | 42 | 41 | 23 | 24 | 14 | 9 |
| Sedatives | 24 | 26 | 10 | 12 | 5 | 7 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 98 | 98 | 71 | 75 | 45 | 47 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.18D Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Standard Errors of Percentages, 2002 and 2003

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.63 | 0.65 | 0.57 | 0.59 | 0.44 | 0.46 |
| Marijuana and Hashish | 0.61 | 0.66 | 0.53 | 0.55 | 0.41 | 0.41 |
| Cocaine | 0.44 | 0.41 | 0.29 | 0.26 | 0.13 | 0.14 |
| Crack | 0.18 | 0.20 | 0.08 | 0.08 | 0.03 | 0.03 |
| Heroin | 0.13 | 0.13 | 0.05 | 0.05 | 0.03 | 0.03 |
| Hallucinogens | 0.54 | 0.52 | 0.30 | 0.27 | 0.12 | 0.13 |
| LSD | 0.43 | 0.40 | 0.14 | 0.09 | 0.04 | 0.02 |
| PCP | 0.17 | 0.18 | 0.07 | 0.06 | 0.03 | 0.03 |
| Ecstasy | 0.43 | 0.45 | 0.26 | 0.22 | 0.10 | 0.11 |
| Inhalants | 0.41 | 0.39 | 0.12 | 0.14 | 0.06 | 0.06 |
| Nonmedical Use of Any Psychotherapeutic ² | 0.54 | 0.56 | 0.39 | 0.43 | 0.25 | 0.25 |
| Pain Relievers | 0.49 | 0.51 | 0.35 | 0.39 | 0.22 | 0.23 |
| Tranquilizers | 0.34 | 0.41 | 0.23 | 0.26 | 0.13 | 0.12 |
| Stimulants | 0.37 | 0.38 | 0.21 | 0.22 | 0.13 | 0.10 |
| Methamphetamine | 0.27 | 0.26 | 0.15 | 0.15 | 0.09 | 0.06 |
| Sedatives | 0.15 | 0.16 | 0.07 | 0.08 | 0.03 | 0.05 |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.63 | 0.62 | 0.46 | 0.47 | 0.29 | 0.30 |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.18P Illicit Drug Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

| Drug | TIME PERIOD | | | | | |
|----------------------------------------------------------|-------------|------|-----------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| ANY ILLICIT DRUG¹ | 0.7357 | | 0.1654 | | 0.8709 | |
| Marijuana and Hashish | 0.4108 | | 0.0214 | | 0.2716 | |
| Cocaine | 0.7142 | | 0.8111 | | 0.7421 | |
| Crack | 0.8992 | | 0.8020 | | 0.9515 | |
| Heroin | 0.5243 | | 0.5043 | | 0.6963 | |
| Hallucinogens | 0.0600 | | 0.0000 | | 0.4276 | |
| LSD | 0.0000 | | 0.0000 | | 0.0817 | |
| PCP | 0.9708 | | 0.7800 | | 0.8998 | |
| Ecstasy | 0.9334 | | 0.0000 | | 0.6342 | |
| Inhalants | 0.2329 | | 0.4659 | | 0.5704 | |
| Nonmedical Use of Any Psychotherapeutic ² | 0.0348 | | 0.5389 | | 0.4429 | |
| Pain Relievers | 0.0671 | | 0.2565 | | 0.0709 | |
| Tranquilizers | 0.0087 | | 0.1788 | | 0.9930 | |
| Stimulants | 0.8797 | | 0.2715 | | 0.1762 | |
| Methamphetamine | 0.0064 | | 0.1747 | | 0.0902 | |
| Sedatives | 0.4590 | | 0.6553 | | 0.3802 | |
| ANY ILLICIT DRUG OTHER THAN MARIJUANA¹ | 0.7392 | | 0.3458 | | 0.4314 | |

*Low precision; no estimate reported.

¹ Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.