

Table 1.93C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	583	565	222	229	128	147
GEOGRAPHIC DIVISION						
Northeast	264	258	102	92	59	56
New England	168	126	52	56	12	28
Middle Atlantic	205	227	88	73	58	49
Midwest	231	207	89	88	59	51
East North Central	174	174	65	74	47	44
West North Central	150	111	60	48	36	27
South	316	324	129	132	66	101
South Atlantic	241	262	105	107	53	85
East South Central	154	90	54	41	29	27
West South Central	152	164	52	66	28	46
West	339	317	119	137	71	75
Mountain	132	140	41	79	31	55
Pacific	312	285	112	111	64	51
COUNTY TYPE¹						
Large Metro	476	471	173	183	111	117
Small Metro	291	282	116	104	50	57
250K – 1 Mil. Pop.	232	238	89	78	38	45
< 250K Pop.	189	147	73	66	32	36
Nonmetro	187	181	62	88	37	66
Urbanized	137	117	38	54	27	21
Less Urbanized	122	123	46	72	23	63
Completely Rural	52	64	20	10	10	9

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.93D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.25	0.24	0.09	0.10	0.05	0.06
GEOGRAPHIC DIVISION						
Northeast	0.59	0.57	0.23	0.20	0.13	0.12
New England	1.42	1.06	0.44	0.47	0.10	0.23
Middle Atlantic	0.62	0.68	0.26	0.22	0.18	0.15
Midwest	0.43	0.38	0.17	0.16	0.11	0.10
East North Central	0.47	0.46	0.17	0.20	0.12	0.12
West North Central	0.94	0.69	0.37	0.30	0.23	0.17
South	0.38	0.38	0.15	0.16	0.08	0.12
South Atlantic	0.55	0.59	0.24	0.24	0.12	0.19
East South Central	1.09	0.63	0.38	0.29	0.20	0.19
West South Central	0.59	0.63	0.20	0.25	0.11	0.18
West	0.64	0.59	0.23	0.25	0.13	0.14
Mountain	0.86	0.90	0.27	0.51	0.20	0.35
Pacific	0.83	0.74	0.30	0.29	0.17	0.13
COUNTY TYPE¹						
Large Metro	0.38	0.37	0.14	0.15	0.09	0.09
Small Metro	0.42	0.39	0.17	0.15	0.07	0.08
250K – 1 Mil. Pop.	0.51	0.51	0.20	0.17	0.08	0.10
< 250K Pop.	0.80	0.60	0.31	0.27	0.13	0.15
Nonmetro	0.47	0.45	0.15	0.22	0.09	0.16
Urbanized	0.85	0.65	0.24	0.30	0.17	0.12
Less Urbanized	0.62	0.68	0.23	0.39	0.12	0.34
Completely Rural	1.18	1.57	0.45	0.25	0.23	0.22

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.93P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4415		0.8514		0.2069	
GEOGRAPHIC DIVISION						
Northeast	0.1222		0.5705		0.2986	
New England	0.8555		0.7939		0.2508	
Middle Atlantic	0.0530		0.6144		0.5355	
Midwest	0.2602		0.1168		0.0273	
East North Central	0.2184		0.4384		0.2286	
West North Central	0.8068		0.1246		0.0318	
South	0.3809		0.9731		0.0287	
South Atlantic	0.2586		0.7203		0.1538	
East South Central	0.4509		0.7570		0.8722	
West South Central	0.5243		0.3808		0.0225	
West	0.8000		0.7785		0.3879	
Mountain	0.6124		0.4862		0.6395	
Pacific	0.9436		0.9655		0.4717	
COUNTY TYPE¹						
Large Metro	0.4195		0.7705		0.5225	
Small Metro	0.4494		0.1099		0.3537	
250K – 1 Mil. Pop.	0.8939		0.2488		0.2057	
< 250K Pop.	0.1272		0.2693		0.9211	
Nonmetro	0.1140		0.1842		0.3313	
Urbanized	0.5729		0.1385		0.9959	
Less Urbanized	0.2933		0.4473		0.2905	
Completely Rural	0.3658		0.0642		0.9106	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.94C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	36	36	32	28	18	16
GEOGRAPHIC DIVISION						
Northeast	12	12	11	9	8	4
New England	6	6	6	4	5	3
Middle Atlantic	10	10	9	8	7	3
Midwest	13	15	12	12	7	5
East North Central	12	12	10	10	6	4
West North Central	6	8	6	6	3	4
South	26	22	23	17	14	11
South Atlantic	21	14	19	12	10	8
East South Central	7	6	6	5	4	2
West South Central	13	14	12	12	8	7
West	18	22	15	16	6	9
Mountain	8	11	7	9	4	5
Pacific	15	19	13	13	5	7
COUNTY TYPE¹						
Large Metro	25	26	22	18	12	11
Small Metro	21	20	18	16	11	10
250K – 1 Mil. Pop.	17	16	16	13	8	9
< 250K Pop.	13	10	10	9	8	3
Nonmetro	16	16	12	15	9	6
Urbanized	10	12	8	11	6	4
Less Urbanized	11	10	8	9	5	5
Completely Rural	4	5	4	5	2	0

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.94D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.15	0.15	0.13	0.11	0.07	0.06
GEOGRAPHIC DIVISION						
Northeast	0.26	0.26	0.25	0.19	0.19	0.10
New England	0.53	0.52	0.49	0.37	0.41	0.23
Middle Atlantic	0.30	0.29	0.28	0.23	0.20	0.10
Midwest	0.23	0.26	0.20	0.21	0.12	0.10
East North Central	0.29	0.31	0.25	0.25	0.14	0.10
West North Central	0.36	0.50	0.35	0.36	0.21	0.22
South	0.29	0.24	0.27	0.19	0.16	0.12
South Atlantic	0.48	0.32	0.43	0.26	0.23	0.17
East South Central	0.50	0.44	0.43	0.36	0.29	0.17
West South Central	0.44	0.49	0.42	0.39	0.28	0.24
West	0.30	0.38	0.25	0.28	0.11	0.15
Mountain	0.50	0.62	0.41	0.53	0.23	0.29
Pacific	0.37	0.47	0.32	0.32	0.12	0.17
COUNTY TYPE¹						
Large Metro	0.19	0.20	0.17	0.14	0.09	0.08
Small Metro	0.29	0.27	0.25	0.22	0.15	0.13
250K – 1 Mil. Pop.	0.34	0.32	0.32	0.25	0.17	0.19
< 250K Pop.	0.53	0.42	0.44	0.36	0.36	0.12
Nonmetro	0.37	0.38	0.28	0.35	0.20	0.15
Urbanized	0.61	0.65	0.45	0.60	0.37	0.23
Less Urbanized	0.54	0.49	0.40	0.44	0.25	0.24
Completely Rural	0.89	1.09	0.88	1.09	0.52	0.04

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.94P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.5530		0.0730		0.8041	
GEOGRAPHIC DIVISION						
Northeast	0.4375		0.2712		0.3103	
New England	0.1958		0.0571		0.4684	
Middle Atlantic	0.9508		0.8585		0.4631	
Midwest	0.5102		0.9882		0.4113	
East North Central	0.7984		0.8839		0.1737	
West North Central	0.4202		0.7950		0.6723	
South	0.0959		0.0476		0.7676	
South Atlantic	0.0467		0.0244		0.5663	
East South Central	0.2717		0.3941		0.5486	
West South Central	0.7658		0.9478		0.1228	
West	0.3311		0.9461		0.1289	
Mountain	0.0284		0.3652		0.2683	
Pacific	0.8108		0.4216		0.2861	
COUNTY TYPE¹						
Large Metro	0.8435		0.2472		0.1988	
Small Metro	0.1603		0.0510		0.6377	
250K – 1 Mil. Pop.	0.1241		0.0374		0.7244	
< 250K Pop.	0.8192		0.7757		0.2004	
Nonmetro	0.8636		0.7919		0.4807	
Urbanized	0.3476		0.4907		0.8331	
Less Urbanized	0.5528		0.8804		0.8039	
Completely Rural	0.8159		0.7441		0.0999	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.95C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	114	107	74	72	37	42
GEOGRAPHIC DIVISION						
Northeast	41	37	29	30	16	16
New England	18	19	16	18	7	8
Middle Atlantic	37	32	24	24	15	14
Midwest	38	39	31	27	16	14
East North Central	33	33	26	23	14	12
West North Central	20	22	17	15	9	6
South	69	68	43	42	24	24
South Atlantic	53	46	32	30	18	18
East South Central	26	24	18	15	9	5
West South Central	34	43	23	25	11	14
West	69	62	43	42	17	27
Mountain	26	33	20	24	12	15
Pacific	65	52	38	35	13	22
COUNTY TYPE¹						
Large Metro	79	85	52	57	27	34
Small Metro	73	56	45	37	23	21
250K – 1 Mil. Pop.	59	48	38	33	17	20
< 250K Pop.	43	31	25	17	15	9
Nonmetro	33	33	24	23	11	13
Urbanized	22	25	16	17	8	11
Less Urbanized	24	21	14	15	7	5
Completely Rural	8	8	7	5	3	3

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.95D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.37	0.34	0.24	0.23	0.12	0.13
GEOGRAPHIC DIVISION						
Northeast	0.75	0.65	0.52	0.53	0.29	0.29
New England	1.25	1.27	1.11	1.21	0.49	0.52
Middle Atlantic	0.92	0.77	0.59	0.59	0.36	0.34
Midwest	0.53	0.53	0.43	0.37	0.23	0.19
East North Central	0.66	0.64	0.53	0.44	0.27	0.24
West North Central	0.88	0.95	0.75	0.66	0.41	0.28
South	0.62	0.60	0.39	0.37	0.21	0.21
South Atlantic	0.98	0.83	0.59	0.54	0.34	0.33
East South Central	1.37	1.25	0.93	0.77	0.49	0.28
West South Central	0.92	1.13	0.62	0.65	0.31	0.37
West	0.95	0.83	0.59	0.57	0.24	0.37
Mountain	1.18	1.48	0.90	1.06	0.55	0.69
Pacific	1.26	1.00	0.75	0.67	0.24	0.43
COUNTY TYPE¹						
Large Metro	0.50	0.51	0.32	0.34	0.17	0.20
Small Metro	0.72	0.55	0.45	0.36	0.23	0.21
250K – 1 Mil. Pop.	0.91	0.71	0.58	0.49	0.27	0.30
< 250K Pop.	1.15	0.87	0.67	0.48	0.39	0.24
Nonmetro	0.67	0.67	0.48	0.47	0.23	0.26
Urbanized	0.97	1.04	0.72	0.73	0.35	0.46
Less Urbanized	1.05	0.95	0.64	0.69	0.33	0.22
Completely Rural	1.67	2.09	1.52	1.31	0.69	0.95

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.95P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.3762		0.7314		0.3692	
GEOGRAPHIC DIVISION						
Northeast	0.9218		0.2171		0.5344	
New England	0.1140		0.5964		0.9076	
Middle Atlantic	0.4503		0.2638		0.4684	
Midwest	0.9758		0.0699		0.7198	
East North Central	0.2226		0.1096		0.8604	
West North Central	0.1043		0.3649		0.3789	
South	0.3425		0.3824		0.6100	
South Atlantic	0.2535		0.4566		0.7435	
East South Central	1.0000		0.5102		0.2985	
West South Central	0.8468		0.8241		0.8960	
West	0.6742		0.5238		0.0425	
Mountain	0.6805		0.9730		0.8802	
Pacific	0.5216		0.4851		0.0098	
COUNTY TYPE¹						
Large Metro	0.6540		0.7263		0.3675	
Small Metro	0.1578		0.0948		0.6552	
250K – 1 Mil. Pop.	0.6690		0.6844		0.2955	
< 250K Pop.	0.0632		0.0066		0.0088	
Nonmetro	0.2922		0.2038		0.1722	
Urbanized	0.4640		0.2193		0.0193	
Less Urbanized	0.5769		0.5015		0.1307	
Completely Rural	0.7799		0.6793		0.5271	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.96C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	574	552	200	207	119	137
GEOGRAPHIC DIVISION						
Northeast	259	253	92	81	53	51
New England	160	120	45	46	9	22
Middle Atlantic	204	224	80	67	52	46
Midwest	227	204	81	83	56	49
East North Central	167	171	58	68	44	42
West North Central	152	110	56	48	35	25
South	310	310	115	122	59	97
South Atlantic	243	254	91	99	48	83
East South Central	149	86	47	39	26	26
West South Central	142	154	52	59	23	44
West	337	318	108	122	69	65
Mountain	132	135	39	67	28	45
Pacific	310	288	101	102	64	47
COUNTY TYPE¹						
Large Metro	467	461	159	165	106	107
Small Metro	277	272	102	95	42	53
250K – 1 Mil. Pop.	227	231	77	72	33	41
< 250K Pop.	179	139	65	62	25	34
Nonmetro	183	178	59	83	33	64
Urbanized	131	114	33	46	24	16
Less Urbanized	118	120	46	70	20	62
Completely Rural	50	64	17	8	8	8

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.96D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.32	0.30	0.11	0.11	0.07	0.08
GEOGRAPHIC DIVISION						
Northeast	0.74	0.72	0.26	0.23	0.15	0.15
New England	1.73	1.30	0.49	0.50	0.10	0.24
Middle Atlantic	0.79	0.86	0.31	0.26	0.20	0.18
Midwest	0.56	0.50	0.20	0.20	0.14	0.12
East North Central	0.59	0.60	0.20	0.24	0.15	0.15
West North Central	1.26	0.91	0.46	0.39	0.29	0.21
South	0.49	0.48	0.18	0.19	0.09	0.15
South Atlantic	0.71	0.74	0.27	0.29	0.14	0.24
East South Central	1.38	0.80	0.44	0.36	0.24	0.24
West South Central	0.74	0.80	0.27	0.30	0.12	0.23
West	0.85	0.78	0.27	0.30	0.17	0.16
Mountain	1.16	1.16	0.34	0.58	0.25	0.39
Pacific	1.09	1.00	0.36	0.35	0.22	0.16
COUNTY TYPE¹						
Large Metro	0.48	0.48	0.16	0.17	0.11	0.11
Small Metro	0.53	0.50	0.20	0.18	0.08	0.10
250K – 1 Mil. Pop.	0.66	0.66	0.23	0.20	0.10	0.11
< 250K Pop.	1.01	0.75	0.37	0.33	0.14	0.19
Nonmetro	0.59	0.57	0.19	0.27	0.11	0.21
Urbanized	1.09	0.83	0.27	0.33	0.20	0.11
Less Urbanized	0.78	0.86	0.30	0.50	0.13	0.44
Completely Rural	1.43	1.95	0.48	0.25	0.23	0.25

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.96P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.3191		0.8974		0.3141	
GEOGRAPHIC DIVISION						
Northeast	0.1054		0.7579		0.3003	
New England	0.7807		0.7624		0.1083	
Middle Atlantic	0.0363		0.8520		0.6245	
Midwest	0.2787		0.2931		0.0312	
East North Central	0.1469		0.7441		0.2342	
West North Central	0.9309		0.2220		0.0374	
South	0.5693		0.4676		0.0151	
South Atlantic	0.4524		0.7978		0.0975	
East South Central	0.4640		0.9680		0.8392	
West South Central	0.4842		0.2683		0.0304	
West	0.7862		0.9486		0.9895	
Mountain	0.8189		0.5530		0.6731	
Pacific	0.8429		0.8122		0.7874	
COUNTY TYPE¹						
Large Metro	0.3445		0.8331		0.8323	
Small Metro	0.6956		0.5327		0.1608	
250K – 1 Mil. Pop.	0.7635		0.4890		0.3939	
< 250K Pop.	0.2458		0.9091		0.2420	
Nonmetro	0.1648		0.3468		0.4380	
Urbanized	0.7321		0.2179		0.3285	
Less Urbanized	0.3086		0.5709		0.2020	
Completely Rural	0.3674		0.0931		0.9849	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.97C Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	584	564	217	225	126	146
GEOGRAPHIC DIVISION						
Northeast	263	258	100	91	59	56
New England	165	126	51	54	11	27
Middle Atlantic	205	227	86	74	58	49
Midwest	229	205	87	86	58	51
East North Central	173	173	64	72	45	44
West North Central	149	110	59	47	36	26
South	318	321	126	130	64	100
South Atlantic	242	260	101	106	52	84
East South Central	154	88	51	40	27	27
West South Central	155	165	55	65	27	46
West	340	321	117	134	70	75
Mountain	132	140	42	79	29	55
Pacific	314	288	110	108	64	50
COUNTY TYPE¹						
Large Metro	477	473	169	182	110	117
Small Metro	290	280	115	100	48	56
250K – 1 Mil. Pop.	232	238	87	76	37	43
< 250K Pop.	188	146	73	65	31	35
Nonmetro	183	181	60	86	34	65
Urbanized	136	117	37	52	26	20
Less Urbanized	119	123	44	72	21	63
Completely Rural	52	64	17	9	9	9

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.97D Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.28	0.27	0.10	0.11	0.06	0.07
GEOGRAPHIC DIVISION						
Northeast	0.65	0.63	0.25	0.22	0.14	0.14
New England	1.55	1.18	0.48	0.51	0.11	0.25
Middle Atlantic	0.69	0.75	0.29	0.24	0.19	0.16
Midwest	0.48	0.43	0.18	0.18	0.12	0.11
East North Central	0.52	0.51	0.19	0.21	0.13	0.13
West North Central	1.04	0.76	0.41	0.33	0.25	0.18
South	0.42	0.42	0.17	0.17	0.09	0.13
South Atlantic	0.61	0.65	0.26	0.26	0.13	0.21
East South Central	1.21	0.69	0.40	0.31	0.21	0.21
West South Central	0.68	0.71	0.24	0.28	0.12	0.20
West	0.72	0.67	0.25	0.28	0.15	0.16
Mountain	0.97	1.01	0.31	0.57	0.21	0.40
Pacific	0.94	0.85	0.33	0.32	0.19	0.15
COUNTY TYPE¹						
Large Metro	0.42	0.42	0.15	0.16	0.10	0.10
Small Metro	0.47	0.44	0.18	0.16	0.08	0.09
250K – 1 Mil. Pop.	0.57	0.57	0.21	0.18	0.09	0.10
< 250K Pop.	0.88	0.66	0.34	0.29	0.14	0.16
Nonmetro	0.51	0.50	0.17	0.24	0.10	0.18
Urbanized	0.95	0.73	0.26	0.32	0.18	0.12
Less Urbanized	0.68	0.76	0.25	0.44	0.12	0.39
Completely Rural	1.31	1.74	0.43	0.26	0.22	0.24

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 1.97P Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	0.4223		0.9550		0.2107	
GEOGRAPHIC DIVISION						
Northeast	0.1097		0.4872		0.2420	
New England	0.9094		0.6399		0.1868	
Middle Atlantic	0.0513		0.5934		0.4844	
Midwest	0.2825		0.1115		0.0305	
East North Central	0.2260		0.4220		0.2735	
West North Central	0.8493		0.1212		0.0264	
South	0.4574		0.7050		0.0279	
South Atlantic	0.3325		0.9888		0.1246	
East South Central	0.4793		0.8339		0.9241	
West South Central	0.5507		0.3679		0.0399	
West	0.8554		0.7717		0.4960	
Mountain	0.7492		0.5803		0.7506	
Pacific	0.9475		0.9718		0.5479	
COUNTY TYPE¹						
Large Metro	0.4157		0.6667		0.6076	
Small Metro	0.4943		0.1847		0.3004	
250K – 1 Mil. Pop.	0.8384		0.4093		0.2165	
< 250K Pop.	0.1306		0.2791		0.8912	
Nonmetro	0.1180		0.1867		0.2753	
Urbanized	0.6515		0.1653		0.9674	
Less Urbanized	0.2584		0.4195		0.2741	
Completely Rural	0.3595		0.0422		0.8512	

*Low precision; no estimate reported.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.