

Table 2.1A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	171,838	172,843	84,731	83,415	71,499	70,757
Cigarettes	162,553	163,240	71,310	69,853	61,136	60,434
Smokeless Tobacco	46,870	46,065	10,577	10,347	7,787	7,725
Cigars	88,053	88,096	25,928	25,386	12,751	12,837
Pipe Tobacco ²	40,003	40,064	--	--	1,816	1,619
ALCOHOL	195,452	197,533	155,476	154,540	119,820	118,965
Binge Alcohol Use ³	--	--	--	--	53,787	53,770
Heavy Alcohol Use ³	--	--	--	--	15,860	16,144

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.1B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	73.1	72.7	36.0	35.1	30.4	29.8
Cigarettes	69.1	68.7	30.3	29.4	26.0	25.4
Smokeless Tobacco	19.9	19.4	4.5	4.4	3.3	3.3
Cigars	37.4	37.1	11.0	10.7	5.4	5.4
Pipe Tobacco ²	17.0	16.9	--	--	0.8	0.7
ALCOHOL	83.1	83.1	66.1 ^a	65.0	51.0	50.1
Binge Alcohol Use ³	--	--	--	--	22.9	22.6
Heavy Alcohol Use ³	--	--	--	--	6.7	6.8

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.2A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	9,113 ^a	8,631	5,839	5,634	3,767	3,610
Cigarettes	8,242 ^a	7,752	5,037 ^a	4,737	3,209	3,044
Smokeless Tobacco	1,975	1,897	1,063	1,034	487	511
Cigars	4,033 ^a	3,776	2,511	2,487	1,117	1,115
Pipe Tobacco ²	687	640	--	--	137	160
ALCOHOL	10,747	10,713	8,564	8,563	4,365	4,425
Binge Alcohol Use ³	--	--	--	--	2,641	2,658
Heavy Alcohol Use ³	--	--	--	--	630	652

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.2B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	36.8 ^b	34.5	23.6 ^a	22.5	15.2	14.4
Cigarettes	33.3 ^b	31.0	20.3 ^b	19.0	13.0	12.2
Smokeless Tobacco	8.0	7.6	4.3	4.1	2.0	2.0
Cigars	16.3 ^a	15.1	10.1	10.0	4.5	4.5
Pipe Tobacco ²	2.8	2.6	--	--	0.6	0.6
ALCOHOL	43.4	42.9	34.6	34.3	17.6	17.7
Binge Alcohol Use ³	--	--	--	--	10.7	10.6
Heavy Alcohol Use ³	--	--	--	--	2.5	2.6

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.3A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	23,426	23,723	17,028	17,073	14,049	14,201
Cigarettes	22,099	22,267	15,214	15,113	12,650	12,764
Smokeless Tobacco	7,367	6,977	2,489	2,485	1,502	1,486
Cigars	14,148	14,327	7,034	7,196	3,417	3,623
Pipe Tobacco ²	2,495	2,441	--	--	332	292
ALCOHOL	26,889	27,622	24,153	24,778	18,759	19,466
Binge Alcohol Use ³	--	--	--	--	12,699	13,188
Heavy Alcohol Use ³	--	--	--	--	4,625	4,788

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.3B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	75.5	74.8	54.9	53.8	45.3	44.8
Cigarettes	71.2	70.2	49.0 ^a	47.6	40.8	40.2
Smokeless Tobacco	23.7 ^b	22.0	8.0	7.8	4.8	4.7
Cigars	45.6	45.2	22.7	22.7	11.0	11.4
Pipe Tobacco ²	8.0	7.7	--	--	1.1	0.9
ALCOHOL	86.7	87.1	77.9	78.1	60.5	61.4
Binge Alcohol Use ³	--	--	--	--	40.9	41.6
Heavy Alcohol Use ³	--	--	--	--	14.9	15.1

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.4A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	139,299	140,489	61,864	60,708	53,683	52,946
Cigarettes	132,212	133,221	51,060	50,004	45,278	44,626
Smokeless Tobacco	37,528	37,190	7,025	6,827	5,797	5,728
Cigars	69,872	69,992	16,382	15,702	8,217	8,099
Pipe Tobacco ²	36,821	36,983	--	--	1,347	1,166
ALCOHOL	157,816	159,198	122,759	121,200	96,696	95,073
Binge Alcohol Use ³	--	--	--	--	38,447	37,924
Heavy Alcohol Use ³	--	--	--	--	10,605	10,704

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.4B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	77.7	77.6	34.5	33.5	29.9	29.3
Cigarettes	73.7	73.6	28.5	27.6	25.2	24.7
Smokeless Tobacco	20.9	20.6	3.9	3.8	3.2	3.2
Cigars	39.0	38.7	9.1	8.7	4.6	4.5
Pipe Tobacco ²	20.5	20.4	--	--	0.8	0.6
ALCOHOL	88.0	88.0	68.4 ^a	67.0	53.9	52.5
Binge Alcohol Use ³	--	--	--	--	21.4	21.0
Heavy Alcohol Use ³	--	--	--	--	5.9	5.9

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.5A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	1,355 ^b	1,181	699 ^a	598	314	273
Cigarettes	1,160 ^b	984	599 ^b	471	265	214
Smokeless Tobacco	236	251	106	104	43	44
Cigars	420	372	193	192	56	66
Pipe Tobacco ²	88	96	--	--	27	31
ALCOHOL	1,501	1,538	1,004	1,028	356	382
Binge Alcohol Use ³	--	--	--	--	150	132
Heavy Alcohol Use ³	--	--	--	--	22	13

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.5B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	16.4 ^b	14.0	8.4 ^a	7.1	3.8	3.2
Cigarettes	14.0 ^b	11.6	7.2 ^b	5.6	3.2 ^a	2.5
Smokeless Tobacco	2.8	3.0	1.3	1.2	0.5	0.5
Cigars	5.1	4.4	2.3	2.3	0.7	0.8
Pipe Tobacco ²	1.1	1.1	--	--	0.3	0.4
ALCOHOL	18.1	18.2	12.1	12.2	4.3	4.5
Binge Alcohol Use ³	--	--	--	--	1.8	1.6
Heavy Alcohol Use ³	--	--	--	--	0.3	0.1

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.6A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	3,177 ^a	2,934	1,940	1,868	1,133	1,099
Cigarettes	2,874 ^b	2,595	1,669	1,544	949	909
Smokeless Tobacco	608	568	326	335	148	155
Cigars	1,283	1,153	778	762	319	322
Pipe Tobacco ²	256	208	--	--	55	44
ALCOHOL	3,823	3,644	2,966	2,882	1,405	1,406
Binge Alcohol Use ³	--	--	--	--	780	773
Heavy Alcohol Use ³	--	--	--	--	159	180

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.6B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	37.5	35.5	22.9	22.6	13.4	13.3
Cigarettes	33.9 ^a	31.4	19.7	18.7	11.2	11.0
Smokeless Tobacco	7.2	6.9	3.9	4.1	1.7	1.9
Cigars	15.1	14.0	9.2	9.2	3.8	3.9
Pipe Tobacco ²	3.0	2.5	--	--	0.7	0.5
ALCOHOL	45.1	44.1	35.0	34.9	16.6	17.0
Binge Alcohol Use ³	--	--	--	--	9.2	9.4
Heavy Alcohol Use ³	--	--	--	--	1.9	2.2

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.7A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	4,581	4,516	3,200	3,168	2,320	2,238
Cigarettes	4,208	4,174	2,769	2,721	1,995	1,921
Smokeless Tobacco	1,131	1,078	630	595	296	312
Cigars	2,331	2,251	1,541	1,533	742	728
Pipe Tobacco ²	343	336	--	--	55	85
ALCOHOL	5,422	5,532	4,594	4,653	2,604	2,638
Binge Alcohol Use ³	--	--	--	--	1,711	1,753
Heavy Alcohol Use ³	--	--	--	--	449	459

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.7B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	57.3 ^a	54.5	40.0	38.2	29.0 ^a	27.0
Cigarettes	52.6	50.4	34.6	32.9	24.9	23.2
Smokeless Tobacco	14.1	13.0	7.9	7.2	3.7	3.8
Cigars	29.1 ^a	27.2	19.3	18.5	9.3	8.8
Pipe Tobacco ²	4.3	4.1	--	--	0.7	1.0
ALCOHOL	67.8	66.8	57.4	56.2	32.6	31.8
Binge Alcohol Use ³	--	--	--	--	21.4	21.2
Heavy Alcohol Use ³	--	--	--	--	5.6	5.5

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.8A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	9,010	8,884	6,760	6,624	5,414	5,340
Cigarettes	8,422	8,264	5,967	5,802	4,811	4,771
Smokeless Tobacco	2,602 ^a	2,347	1,140	1,097	636	581
Cigars	5,409	5,183	3,271	3,190	1,624	1,654
Pipe Tobacco ²	909	809	--	--	180	153
ALCOHOL	10,164	10,223	8,928	8,991	6,348	6,451
Binge Alcohol Use ³	--	--	--	--	4,534	4,531
Heavy Alcohol Use ³	--	--	--	--	1,671	1,645

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.8B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	72.4	70.9	54.3	52.9	43.5	42.6
Cigarettes	67.7	66.0	47.9	46.3	38.7	38.1
Smokeless Tobacco	20.9 ^b	18.7	9.2	8.8	5.1	4.6
Cigars	43.5 ^a	41.4	26.3	25.5	13.0	13.2
Pipe Tobacco ²	7.3	6.5	--	--	1.4	1.2
ALCOHOL	81.7	81.6	71.7	71.8	51.0	51.5
Binge Alcohol Use ³	--	--	--	--	36.4	36.2
Heavy Alcohol Use ³	--	--	--	--	13.4	13.1

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.9A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	14,416	14,839	10,268	10,449	8,635	8,861
Cigarettes	13,677	14,003	9,246	9,311	7,839	7,994
Smokeless Tobacco	4,765	4,630	1,349	1,388	866	904
Cigars	8,738	9,144	3,763	4,006	1,794	1,969
Pipe Tobacco ²	1,586	1,632	--	--	152	139
ALCOHOL	16,725	17,399	15,225	15,787	12,411	13,016
Binge Alcohol Use ³	--	--	--	--	8,165	8,657
Heavy Alcohol Use ³	--	--	--	--	2,954	3,144

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.9B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	77.6	77.3	55.3	54.4	46.5	46.1
Cigarettes	73.6	72.9	49.8	48.5	42.2	41.6
Smokeless Tobacco	25.7 ^a	24.1	7.3	7.2	4.7	4.7
Cigars	47.0	47.6	20.3	20.9	9.7	10.3
Pipe Tobacco ²	8.5	8.5	--	--	0.8	0.7
ALCOHOL	90.0	90.6	82.0	82.2	66.8	67.8
Binge Alcohol Use ³	--	--	--	--	44.0	45.1
Heavy Alcohol Use ³	--	--	--	--	15.9	16.4

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.10A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	27,306	27,147	15,948	15,867	13,423	13,561
Cigarettes	25,661	25,266	13,429	13,460	11,494	11,667
Smokeless Tobacco	10,326	9,841	2,340	2,362	1,863	1,843
Cigars	14,679	14,903	5,052	4,775	2,320	2,420
Pipe Tobacco ²	3,349	3,261	--	--	175	172
ALCOHOL	31,725	31,575	27,393	27,306	21,594	21,060
Binge Alcohol Use ³	--	--	--	--	11,642	11,498
Heavy Alcohol Use ³	--	--	--	--	3,179	3,289

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.10B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	77.7	77.7	45.4	45.4	38.2	38.8
Cigarettes	73.0	72.3	38.2	38.5	32.7	33.4
Smokeless Tobacco	29.4	28.2	6.7	6.8	5.3	5.3
Cigars	41.7	42.6	14.4	13.7	6.6	6.9
Pipe Tobacco ²	9.5	9.3	--	--	0.5	0.5
ALCOHOL	90.2	90.3	77.9	78.1	61.4	60.2
Binge Alcohol Use ³	--	--	--	--	33.1	32.9
Heavy Alcohol Use ³	--	--	--	--	9.0	9.4

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.11A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	111,993	113,342	45,916	44,841	40,260	39,385
Cigarettes	106,551	107,955	37,631	36,544	33,783	32,960
Smokeless Tobacco	27,202	27,349	4,685	4,465	3,934	3,885
Cigars	55,192	55,089	11,330	10,927	5,897	5,679
Pipe Tobacco ²	33,472	33,723	--	--	1,172	994
ALCOHOL	126,091	127,623	95,365	93,894	75,102	74,013
Binge Alcohol Use ³	--	--	--	--	26,806	26,426
Heavy Alcohol Use ³	--	--	--	--	7,427	7,415

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.11B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	77.7	77.6	31.8	30.7	27.9	27.0
Cigarettes	73.9	73.9	26.1	25.0	23.4	22.6
Smokeless Tobacco	18.9	18.7	3.2	3.1	2.7	2.7
Cigars	38.3	37.7	7.9	7.5	4.1	3.9
Pipe Tobacco ²	23.2	23.1	--	--	0.8	0.7
ALCOHOL	87.4	87.4	66.1 ^a	64.3	52.1	50.7
Binge Alcohol Use ³	--	--	--	--	18.6	18.1
Heavy Alcohol Use ³	--	--	--	--	5.2	5.1

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.12A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	162,725	164,212	78,892	77,781	67,732	67,147
Cigarettes	154,311	155,488	66,273	65,116	57,928	57,391
Smokeless Tobacco	44,895	44,167	9,514	9,312	7,299	7,214
Cigars	84,020	84,320	23,417	22,898	11,634	11,722
Pipe Tobacco ²	39,316	39,424	--	--	1,679	1,458
ALCOHOL	184,705	186,820	146,912	145,978	115,455	114,539
Binge Alcohol Use ³	--	--	--	--	51,146	51,112
Heavy Alcohol Use ³	--	--	--	--	15,231	15,492

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.12B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	77.3	77.2	37.5	36.6	32.2	31.6
Cigarettes	73.3	73.1	31.5	30.6	27.5	27.0
Smokeless Tobacco	21.3	20.8	4.5	4.4	3.5	3.4
Cigars	39.9	39.6	11.1	10.8	5.5	5.5
Pipe Tobacco ²	18.7	18.5	--	--	0.8	0.7
ALCOHOL	87.8	87.8	69.8 ^a	68.6	54.9	53.9
Binge Alcohol Use ³	--	--	--	--	24.3	24.0
Heavy Alcohol Use ³	--	--	--	--	7.2	7.3

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.13A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	90,952	91,731	49,558	48,800	41,991	41,288
Cigarettes	83,790	84,200	37,813	37,212	32,636	32,263
Smokeless Tobacco	39,102	38,331	9,703	9,391	7,242	7,096
Cigars	65,644	65,506	20,837	19,943	10,669	10,372
Pipe Tobacco ²	34,971	34,600	--	--	1,487	1,400
ALCOHOL	98,138	99,314	80,308	79,999	65,210	65,927
Binge Alcohol Use ³	--	--	--	--	35,456	35,565
Heavy Alcohol Use ³	--	--	--	--	12,216	11,958

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.13B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	80.1	79.8	43.6	42.4	37.0	35.9
Cigarettes	73.8	73.2	33.3	32.4	28.7	28.1
Smokeless Tobacco	34.4	33.3	8.5	8.2	6.4	6.2
Cigars	57.8	57.0	18.3	17.3	9.4	9.0
Pipe Tobacco ²	30.8	30.1	--	--	1.3	1.2
ALCOHOL	86.4	86.4	70.7	69.6	57.4	57.3
Binge Alcohol Use ³	--	--	--	--	31.2	30.9
Heavy Alcohol Use ³	--	--	--	--	10.8	10.4

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.14A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	80,886	81,112	35,173	34,615	29,509	29,469
Cigarettes	78,763	79,040	33,497	32,641	28,500	28,171
Smokeless Tobacco	7,767	7,734	874	955	545	628
Cigars	22,409	22,590	5,091	5,442	2,082 ^a	2,465
Pipe Tobacco ²	5,032	5,464	--	--	330	219
ALCOHOL	97,314	98,219	75,168	74,542	54,610	53,038
Binge Alcohol Use ³	--	--	--	--	18,331	18,205
Heavy Alcohol Use ³	--	--	--	--	3,645 ^a	4,186

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.14B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	66.6	66.1	28.9	28.2	24.3	24.0
Cigarettes	64.8	64.4	27.6	26.6	23.4	23.0
Smokeless Tobacco	6.4	6.3	0.7	0.8	0.4	0.5
Cigars	18.4	18.4	4.2	4.4	1.7 ^a	2.0
Pipe Tobacco ²	4.1	4.5	--	--	0.3	0.2
ALCOHOL	80.1	80.1	61.8	60.8	44.9 ^a	43.2
Binge Alcohol Use ³	--	--	--	--	15.1	14.8
Heavy Alcohol Use ³	--	--	--	--	3.0 ^a	3.4

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.15A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	4,855	4,609	3,142	3,047	2,019	1,988
Cigarettes	4,193	3,944	2,499	2,354	1,556	1,514
Smokeless Tobacco	1,576	1,533	888	892	436	479
Cigars	2,620 ^a	2,414	1,663	1,608	790	789
Pipe Tobacco ²	481	462	--	--	89	121
ALCOHOL	5,490	5,346	4,211	4,102	2,200	2,185
Binge Alcohol Use ³	--	--	--	--	1,441	1,421
Heavy Alcohol Use ³	--	--	--	--	395	374

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.15B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 to 17: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	38.4 ^b	36.1	24.8	23.9	16.0	15.6
Cigarettes	33.2 ^b	30.9	19.8	18.4	12.3	11.9
Smokeless Tobacco	12.5	12.0	7.0	7.0	3.4	3.7
Cigars	20.7 ^b	18.9	13.1	12.6	6.2	6.2
Pipe Tobacco ²	3.8	3.6	--	--	0.7	0.9
ALCOHOL	43.4	41.9	33.3	32.1	17.4	17.1
Binge Alcohol Use ³	--	--	--	--	11.4	11.1
Heavy Alcohol Use ³	--	--	--	--	3.1	2.9

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.16A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	4,258	4,022	2,697	2,587	1,748	1,622
Cigarettes	4,049	3,808	2,538	2,383	1,653	1,530
Smokeless Tobacco	399	364	175	142	51	32
Cigars	1,413	1,362	848	879	327	326
Pipe Tobacco ²	206	178	--	--	49	40
ALCOHOL	5,257	5,368	4,353	4,460	2,165	2,241
Binge Alcohol Use ³	--	--	--	--	1,200	1,237
Heavy Alcohol Use ³	--	--	--	--	235	278

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.16B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 to 17: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	35.2 ^b	32.9	22.3	21.2	14.4 ^a	13.3
Cigarettes	33.4 ^b	31.2	21.0 ^a	19.5	13.6 ^a	12.5
Smokeless Tobacco	3.3	3.0	1.4	1.2	0.4	0.3
Cigars	11.7	11.1	7.0	7.2	2.7	2.7
Pipe Tobacco ²	1.7	1.5	--	--	0.4	0.3
ALCOHOL	43.4	43.9	36.0	36.5	17.9	18.3
Binge Alcohol Use ³	--	--	--	--	9.9	10.1
Heavy Alcohol Use ³	--	--	--	--	1.9	2.3

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.17A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	12,462	12,662	9,618	9,760	8,097	8,226
Cigarettes	11,446	11,577	8,185	8,208	6,897	7,040
Smokeless Tobacco	5,882	5,562	2,297	2,271	1,455	1,418
Cigars	9,081	9,260	5,185	5,281	2,605	2,758
Pipe Tobacco ²	2,014	1,992	--	--	262	228
ALCOHOL	13,660	13,944	12,438	12,790	10,129 ^a	10,652
Binge Alcohol Use ³	--	--	--	--	7,787	8,163
Heavy Alcohol Use ³	--	--	--	--	3,282	3,372

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.17B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 18 to 25: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	80.3	79.5	61.9	61.3	52.1	51.7
Cigarettes	73.7	72.7	52.7	51.6	44.4	44.2
Smokeless Tobacco	37.9 ^b	34.9	14.8	14.3	9.4	8.9
Cigars	58.5	58.2	33.4	33.2	16.8	17.3
Pipe Tobacco ²	13.0	12.5	--	--	1.7	1.4
ALCOHOL	88.0	87.6	80.1	80.3	65.2	66.9
Binge Alcohol Use ³	--	--	--	--	50.2	51.3
Heavy Alcohol Use ³	--	--	--	--	21.1	21.2

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.18A Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Numbers in Thousands, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	10,964	11,061	7,410	7,313	5,952	5,975
Cigarettes	10,653	10,690	7,029	6,905	5,753	5,724
Smokeless Tobacco	1,485	1,415	192	214	47	67
Cigars	5,067	5,068	1,849	1,915	813	865
Pipe Tobacco ²	481	449	--	--	69	65
ALCOHOL	13,229	13,678	11,716	11,987	8,630	8,814
Binge Alcohol Use ³	--	--	--	--	4,911	5,025
Heavy Alcohol Use ³	--	--	--	--	1,343	1,417

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.18B Tobacco and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 18 to 25: Percentages, 2002 and 2003

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
ANY TOBACCO¹	70.8	70.0	47.8	46.3	38.4	37.8
Cigarettes	68.7	67.6	45.4	43.7	37.1	36.2
Smokeless Tobacco	9.6	8.9	1.2	1.4	0.3	0.4
Cigars	32.7	32.1	11.9	12.1	5.2	5.5
Pipe Tobacco ²	3.1	2.8	--	--	0.4	0.4
ALCOHOL	85.4	86.5	75.6	75.8	55.7	55.8
Binge Alcohol Use ³	--	--	--	--	31.7	31.8
Heavy Alcohol Use ³	--	--	--	--	8.7	9.0

*Low precision; no estimate reported.

-- Not available.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.