

Table 2.70A Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	171,838	172,843	84,731	83,415	71,499	70,757
GEOGRAPHIC DIVISION						
Northeast	32,717	32,290	16,096	15,406	13,265	13,019
New England	8,692	8,801	4,014	3,974	3,240	3,164
Middle Atlantic	24,025	23,488	12,082	11,432	10,025	9,856
Midwest	40,565	40,826	20,714	20,799	17,509	17,514
East North Central	28,254	28,536	14,447	14,406	12,368	12,121
West North Central	12,312	12,291	6,267	6,393	5,141	5,393
South	61,660	61,829	30,858	30,989	26,675	26,946
South Atlantic	32,370	32,948	16,082	15,442	13,840	13,301
East South Central	10,905	10,578	5,488	5,783	4,899	5,046
West South Central	18,385	18,303	9,288	9,764	7,935	8,599
West	36,896	37,897	17,064	16,221	14,050	13,277
Mountain	11,101	11,287	5,496	5,609	4,696	4,759
Pacific	25,794	26,610	11,567	10,611	9,354	8,518
COUNTY TYPE¹						
Large Metro	89,925	88,928	43,737 ^a	41,457	36,422	34,681
Small Metro	51,825	53,120	25,309	26,369	21,377	22,441
250K – 1 Mil. Pop.	33,437	35,023	15,979	17,146	13,396	14,486
< 250K Pop.	18,388	18,097	9,331	9,223	7,980	7,955
Nonmetro	30,089	30,795	15,684	15,589	13,701	13,635
Urbanized	11,886 ^a	13,720	6,100 ^a	6,990	5,286 ^a	6,161
Less Urbanized	14,739	13,916	7,777	6,997	6,749	6,039
Completely Rural	3,464	3,159	1,808	1,602	1,667	1,435

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.70B Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	73.1	72.7	36.0	35.1	30.4	29.8
GEOGRAPHIC DIVISION						
Northeast	72.6	71.3	35.7	34.0	29.4	28.7
New England	73.6	74.0	34.0	33.4	27.4	26.6
Middle Atlantic	72.2	70.3	36.3	34.2	30.1	29.5
Midwest	75.9	75.9	38.8	38.7	32.8	32.6
East North Central	75.5	75.8	38.6	38.2	33.1	32.2
West North Central	76.9	76.3	39.1	39.7	32.1	33.5
South	73.6	72.9	36.8	36.6	31.8	31.8
South Atlantic	74.0	74.2	36.8	34.8	31.6	29.9
East South Central	77.0	74.4	38.8	40.7	34.6	35.5
West South Central	71.1	70.1	35.9	37.4	30.7	32.9
West	69.8	70.4	32.3	30.1	26.6	24.7
Mountain	72.5	72.3	35.9	35.9	30.7	30.5
Pacific	68.7	69.6	30.8 ^a	27.8	24.9	22.3
COUNTY TYPE¹						
Large Metro	71.5	70.5	34.8 ^b	32.9	29.0 ^a	27.5
Small Metro	74.8	74.4	36.5	36.9	30.9	31.4
250K – 1 Mil. Pop.	73.5	74.6	35.1	36.5	29.4	30.9
< 250K Pop.	77.3 ^a	74.0	39.2	37.7	33.5	32.5
Nonmetro	75.0	76.6	39.1	38.8	34.2	33.9
Urbanized	74.2	76.8	38.1	39.1	33.0	34.5
Less Urbanized	75.1	76.4	39.6	38.4	34.4	33.1
Completely Rural	78.1	77.1	40.8	39.1	37.6	35.0

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.71A Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	9,113 ^a	8,631	5,839	5,634	3,767	3,610
GEOGRAPHIC DIVISION						
Northeast	1,581	1,502	1,048	997	665	635
New England	402	384	268	265	181	172
Middle Atlantic	1,179	1,118	780	732	484	463
Midwest	2,208 ^a	2,016	1,460	1,373	1,003	923
East North Central	1,480	1,377	984	939	669	613
West North Central	729	639	476	434	334	310
South	3,468	3,271	2,247	2,112	1,484	1,366
South Atlantic	1,805 ^a	1,594	1,195 ^a	1,007	787 ^a	645
East South Central	623	590	387	386	271	267
West South Central	1,040	1,087	666	719	427	455
West	1,856	1,841	1,083	1,152	615	686
Mountain	618	621	382	400	241	252
Pacific	1,239	1,220	701	752	374	433
COUNTY TYPE¹						
Large Metro	4,498 ^b	4,051	2,830 ^a	2,590	1,741	1,588
Small Metro	2,723	2,752	1,785	1,815	1,173	1,187
250K – 1 Mil. Pop.	1,808	1,804	1,186	1,168	765	746
< 250K Pop.	915	948	599	647	408	441
Nonmetro	1,892	1,827	1,225	1,229	853	835
Urbanized	705	749	445	503	310	345
Less Urbanized	951	908	621	608	426	401
Completely Rural	236	170	159	118	117	89

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.71B Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	36.8 ^b	34.5	23.6 ^a	22.5	15.2	14.4
GEOGRAPHIC DIVISION						
Northeast	35.4	33.5	23.5	22.2	14.9	14.1
New England	34.8	32.8	23.1	22.6	15.6	14.7
Middle Atlantic	35.7	33.7	23.6	22.1	14.6	13.9
Midwest	38.8 ^b	35.4	25.7	24.1	17.6	16.2
East North Central	37.1 ^a	34.5	24.7	23.5	16.8	15.4
West North Central	42.8 ^a	37.6	27.9	25.5	19.6	18.2
South	39.5 ^b	36.6	25.6 ^a	23.7	16.9 ^a	15.3
South Atlantic	41.1 ^b	35.4	27.2 ^b	22.4	17.9 ^b	14.3
East South Central	43.7	40.6	27.1	26.6	19.0	18.4
West South Central	35.2	36.6	22.6	24.2	14.5	15.3
West	31.8	31.3	18.6	19.6	10.6	11.7
Mountain	36.2	35.9	22.4	23.2	14.1	14.6
Pacific	30.0	29.4	17.0	18.1	9.1	10.4
COUNTY TYPE¹						
Large Metro	34.1 ^b	30.4	21.4 ^b	19.5	13.2 ^a	11.9
Small Metro	37.6	37.2	24.6	24.5	16.2	16.0
250K – 1 Mil. Pop.	36.9	36.2	24.2	23.5	15.6	15.0
< 250K Pop.	39.1	39.2	25.6	26.7	17.4	18.2
Nonmetro	44.0	42.6	28.5	28.7	19.8	19.5
Urbanized	40.9	40.3	25.8	27.1	18.0	18.6
Less Urbanized	45.2	45.7	29.5	30.6	20.3	20.2
Completely Rural	49.6 ^b	38.4	33.4	26.7	24.7	20.1

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.72A Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	23,426	23,723	17,028	17,073	14,049	14,201
GEOGRAPHIC DIVISION						
Northeast	4,200	4,215	3,147	3,117	2,586	2,523
New England	1,122	1,138	807	816	665	649
Middle Atlantic	3,078	3,077	2,339	2,301	1,921	1,874
Midwest	5,710	5,819	4,343	4,318	3,621	3,573
East North Central	3,926	3,976	2,984	2,938	2,500	2,431
West North Central	1,784	1,843	1,360	1,380	1,121	1,141
South	8,376	8,367	6,037	6,083	5,033	5,212
South Atlantic	4,039	4,054	2,902	2,942	2,424	2,479
East South Central	1,525	1,448	1,121	1,071	949	949
West South Central	2,811	2,865	2,013	2,069	1,659	1,784
West	5,140	5,322	3,501	3,556	2,809	2,894
Mountain	1,609	1,640	1,127	1,100	933	915
Pacific	3,532	3,683	2,374	2,456	1,876	1,978
COUNTY TYPE¹						
Large Metro	11,661	11,979	8,353	8,433	6,824	6,921
Small Metro	7,796	7,792	5,615	5,684	4,604	4,755
250K – 1 Mil. Pop.	4,897	5,045	3,469	3,670	2,866	3,057
< 250K Pop.	2,899	2,747	2,146	2,014	1,738	1,697
Nonmetro	3,969	3,951	3,060	2,956	2,621	2,526
Urbanized	1,779	1,890	1,344	1,410	1,151	1,194
Less Urbanized	1,819	1,755	1,437	1,318	1,223	1,124
Completely Rural	371	305	279	227	247	208

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.72B Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	75.5	74.8	54.9	53.8	45.3	44.8
GEOGRAPHIC DIVISION						
Northeast	76.3	74.8	57.2	55.3	47.0	44.8
New England	78.3	77.4	56.3	55.5	46.4	44.1
Middle Atlantic	75.6	73.9	57.4	55.3	47.2	45.0
Midwest	79.3	79.1	60.3	58.7	50.3	48.6
East North Central	78.9	78.2	60.0	57.8	50.2	47.8
West North Central	80.2	81.0	61.1	60.7	50.4	50.2
South	76.0 ^a	73.9	54.8	53.7	45.7	46.0
South Atlantic	74.9	72.6	53.8	52.7	45.0	44.4
East South Central	79.8 ^a	74.8	58.6	55.4	49.7	49.0
West South Central	75.6	75.3	54.1	54.4	44.6	46.9
West	70.5	71.8	48.0	48.0	38.5	39.0
Mountain	73.7	73.3	51.6	49.2	42.8	40.9
Pacific	69.1	71.2	46.4	47.5	36.7	38.2
COUNTY TYPE¹						
Large Metro	73.4	72.1	52.6	50.8	43.0	41.7
Small Metro	76.7	76.3	55.2	55.7	45.3	46.6
250K – 1 Mil. Pop.	76.0	75.9	53.9	55.2	44.5	46.0
< 250K Pop.	77.8	77.3	57.6	56.7	46.7	47.7
Nonmetro	79.7	80.3	61.5	60.1	52.7	51.4
Urbanized	79.2	80.0	59.8	59.7	51.2	50.5
Less Urbanized	80.3	80.1	63.4	60.1	54.0	51.3
Completely Rural	79.7	84.0	59.9	62.6	53.0	57.2

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.73A Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	139,299	140,489	61,864	60,708	53,683	52,946
GEOGRAPHIC DIVISION						
Northeast	26,936	26,573	11,901	11,292	10,015	9,862
New England	7,168	7,280	2,939	2,893	2,394	2,343
Middle Atlantic	19,768	19,293	8,963	8,399	7,621	7,519
Midwest	32,647	32,992	14,910	15,108	12,885	13,018
East North Central	22,848	23,182	10,479	10,529	9,200	9,076
West North Central	9,799	9,809	4,431	4,579	3,686	3,942
South	49,817	50,191	22,574	22,795	20,158	20,368
South Atlantic	26,526	27,300	11,985	11,493	10,629	10,177
East South Central	8,757	8,540	3,979	4,326	3,679	3,831
West South Central	14,534	14,351	6,609	6,976	5,849	6,360
West	29,899	30,734	12,479	11,513	10,626	9,698
Mountain	8,875	9,026	3,987	4,109	3,522	3,591
Pacific	21,024	21,707	8,492	7,403	7,104	6,107
COUNTY TYPE¹						
Large Metro	73,766	72,897	32,555 ^a	30,434	27,857	26,172
Small Metro	41,305	42,575	17,909	18,870	15,599	16,500
250K – 1 Mil. Pop.	26,731	28,173	11,324	12,308	9,765	10,683
< 250K Pop.	14,574	14,402	6,586	6,562	5,834	5,817
Nonmetro	24,228	25,017	11,400	11,404	10,227	10,274
Urbanized	9,402 ^a	11,080	4,311	5,077	3,825 ^a	4,622
Less Urbanized	11,969	11,252	5,719	5,070	5,100	4,514
Completely Rural	2,857	2,684	1,370	1,256	1,302	1,139

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.73B Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	77.7	77.6	34.5	33.5	29.9	29.3
GEOGRAPHIC DIVISION						
Northeast	76.7	75.5	33.9	32.1	28.5	28.0
New England	77.8	78.8	31.9	31.3	26.0	25.3
Middle Atlantic	76.4	74.4	34.6	32.4	29.4	29.0
Midwest	80.5	81.0	36.8	37.1	31.8	32.0
East North Central	80.3	81.1	36.8	36.8	32.3	31.7
West North Central	81.1	80.9	36.7	37.8	30.5	32.5
South	77.9	77.8	35.3	35.3	31.5	31.6
South Atlantic	78.1	79.5	35.3	33.5	31.3	29.6
East South Central	81.0	78.9	36.8	39.9	34.0	35.4
West South Central	75.8	74.2	34.5	36.1	30.5	32.9
West	75.2	75.8	31.4	28.4	26.7	23.9
Mountain	77.7	77.5	34.9	35.3	30.8	30.8
Pacific	74.2	75.1	30.0 ^a	25.6	25.1 ^a	21.1
COUNTY TYPE¹						
Large Metro	76.3	75.8	33.7 ^a	31.6	28.8	27.2
Small Metro	79.6	79.2	34.5	35.1	30.1	30.7
250K – 1 Mil. Pop.	78.3	79.8	33.2	34.9	28.6	30.3
< 250K Pop.	82.2 ^b	77.9	37.1	35.5	32.9	31.5
Nonmetro	78.6	80.7	37.0	36.8	33.2	33.2
Urbanized	78.0	81.2	35.8	37.2	31.7	33.9
Less Urbanized	78.4	80.1	37.5	36.1	33.4	32.1
Completely Rural	81.8	81.5	39.2	38.1	37.3	34.6

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.74A Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	162,725	164,212	78,892	77,781	67,732	67,147
GEOGRAPHIC DIVISION						
Northeast	31,136	30,787	15,048	14,409	12,601	12,385
New England	8,290	8,417	3,746	3,709	3,059	2,992
Middle Atlantic	22,846	22,370	11,302	10,700	9,542	9,393
Midwest	38,357	38,810	19,253	19,426	16,506	16,591
East North Central	26,774	27,159	13,463	13,467	11,699	11,508
West North Central	11,583	11,652	5,790	5,959	4,807	5,083
South	58,193	58,558	28,611	28,878	25,191	25,580
South Atlantic	30,565	31,354	14,888	14,435	13,054	12,656
East South Central	10,282	9,988	5,101	5,397	4,629	4,780
West South Central	17,345	17,216	8,622	9,045	7,508	8,144
West	35,040	36,056	15,980	15,068	13,435	12,592
Mountain	10,484	10,666	5,114	5,209	4,455	4,507
Pacific	24,556	25,390	10,866	9,859	8,980	8,085
COUNTY TYPE¹						
Large Metro	85,427	84,876	40,908	38,867	34,681	33,093
Small Metro	49,101	50,368	23,525	24,554	20,204	21,254
250K – 1 Mil. Pop.	31,628	33,219	14,793	15,978	12,632	13,740
< 250K Pop.	17,473	17,149	8,732	8,576	7,572	7,514
Nonmetro	28,197	28,968	14,460	14,359	12,848	12,800
Urbanized	11,181 ^a	12,971	5,655	6,487	4,976 ^a	5,816
Less Urbanized	13,788	13,008	7,156	6,388	6,323	5,638
Completely Rural	3,228	2,989	1,649	1,484	1,549	1,347

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

Table 2.74B Any Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2002 and 2003

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2002	2003	2002	2003	2002	2003
TOTAL	77.3	77.2	37.5	36.6	32.2	31.6
GEOGRAPHIC DIVISION						
Northeast	76.7	75.4	37.1	35.3	31.0	30.3
New England	77.8	78.6	35.2	34.6	28.7	27.9
Middle Atlantic	76.3	74.3	37.7	35.5	31.9	31.2
Midwest	80.4	80.7	40.3	40.4	34.6	34.5
East North Central	80.1	80.7	40.3	40.0	35.0	34.2
West North Central	80.9	80.9	40.5	41.4	33.6	35.3
South	77.6	77.2	38.1	38.1	33.6	33.7
South Atlantic	77.6	78.5	37.8	36.2	33.2	31.7
East South Central	80.8	78.3	40.1	42.3	36.4	37.4
West South Central	75.7	74.4	37.6	39.1	32.8	35.2
West	74.5	75.2	34.0	31.4	28.6	26.3
Mountain	77.0	76.8	37.6	37.5	32.7	32.5
Pacific	73.4	74.5	32.5 ^a	28.9	26.8	23.7
COUNTY TYPE¹						
Large Metro	75.9	75.3	36.3 ^a	34.5	30.8 ^a	29.3
Small Metro	79.1	78.7	37.9	38.4	32.6	33.2
250K – 1 Mil. Pop.	77.9	79.2	36.4	38.1	31.1	32.8
< 250K Pop.	81.4 ^b	77.8	40.7	38.9	35.3	34.1
Nonmetro	78.8	80.7	40.4	40.0	35.9	35.7
Urbanized	78.2	81.0	39.5	40.5	34.8	36.3
Less Urbanized	78.7	80.1	40.8	39.3	36.1	34.7
Completely Rural	81.5	81.8	41.6	40.6	39.1	36.8

*Low precision; no estimate reported.

NOTE: Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^aDifference between estimate and 2003 estimate is statistically significant at the 0.05 level.

^bDifference between estimate and 2003 estimate is statistically significant at the 0.01 level.

¹ Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.