

**Table 5.66A Locations Received Any Illicit Drug Treatment in the Past Year among Persons Who Received Any Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2002 and 2003**

Location of Treatment <sup>1</sup>	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Hospital as an Inpatient	286	187	*	17	46	54	*	*
Rehabilitation Facility - Inpatient	441	318	*	29	80	88	*	*
Rehabilitation Facility - Outpatient	568	427	70 <sup>a</sup>	39	99	126	399	*
Mental Health Center - Outpatient	328	207	*	23	73	44	*	139
Emergency Room	88	40	2	6	14	9	72	25
Private Doctor's Office	*	61	6	7	21 <sup>b</sup>	4	*	50
Self-Help Group	615	410	*	33	97	119	462	*
Prison/Jail	89	71	8	7	21	13	59	*

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Received Any Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug use, or for medical problems associated with drug use.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 5.66B Locations Received Any Illicit Drug Treatment in the Past Year among Persons Who Received Any Illicit Drug Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2002 and 2003**

Location of Treatment <sup>1</sup>	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Hospital as an Inpatient	20.2	17.0	*	15.1	16.1	15.9	*	*
Rehabilitation Facility - Inpatient	31.2	28.9	*	25.4	27.9	25.9	*	*
Rehabilitation Facility - Outpatient	40.2	38.7	49.4	34.4	34.4	37.0	40.5	*
Mental Health Center - Outpatient	23.3	18.7	*	20.2	25.4 <sup>b</sup>	13.0	*	21.5
Emergency Room	6.2	3.6	1.2	4.9	4.9	2.7	7.3	3.9
Private Doctor's Office	*	5.5	4.1	6.3	7.2 <sup>b</sup>	1.3	*	7.6
Self-Help Group	43.5	37.2	*	29.0	33.9	35.0	47.0	*
Prison/Jail	6.3	6.4	5.9	6.3	7.4	3.9	6.0	*

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Received Any Illicit Drug Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug use, or for medical problems associated with drug use.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 5.67A Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2002 and 2003**

Location of Treatment <sup>1</sup>	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Hospital as an Inpatient	275	216	*	*	45	44	*	151
Rehabilitation Facility - Inpatient	396	303	*	*	75	79	280	*
Rehabilitation Facility - Outpatient	544	384	55	*	88	112	401	235
Mental Health Center - Outpatient	316	193	*	*	67	43	*	129
Emergency Room	92 <sup>a</sup>	27	*	*	12	6	70	16
Private Doctor's Office	151	68	*	*	17 <sup>a</sup>	4	*	56
Self-Help Group	508	422	*	*	88	103	378	291
Prison/Jail	95	80	*	*	24	14	60	62

\*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 5.67B Locations Received Alcohol Treatment in the Past Year among Persons Who Received Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2002 and 2003**

Location of Treatment <sup>1</sup>	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Hospital as an Inpatient	17.8	16.6	*	*	16.6	14.4	*	16.8
Rehabilitation Facility - Inpatient	25.5	23.3	*	*	27.7	25.9	24.2	*
Rehabilitation Facility - Outpatient	35.1	29.5	45.1	*	32.7	36.8	34.7	26.2
Mental Health Center - Outpatient	20.4	14.9	*	*	24.8 <sup>a</sup>	14.0	*	14.4
Emergency Room	6.0 <sup>a</sup>	2.1	*	*	4.5	2.1	6.1	1.8
Private Doctor's Office	9.8	5.3	*	*	6.4 <sup>b</sup>	1.2	*	6.3
Self-Help Group	32.8	32.5	*	*	32.6	33.8	32.6	32.5
Prison/Jail	6.2	6.2	*	*	9.0	4.5	5.2	6.9

\*Low precision; no estimate reported.

NOTE: Received Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop alcohol use, or for medical problems associated with alcohol use.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 5.68A Locations Received Any Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Any Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Numbers in Thousands, 2002 and 2003**

Location of Treatment <sup>1</sup>	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Hospital as an Inpatient	859 <sup>a</sup>	587	72	57	117	123	671 <sup>a</sup>	407
Rehabilitation Facility - Inpatient	1,094 <sup>a</sup>	752	83	65	160	185	852 <sup>b</sup>	502
Rehabilitation Facility - Outpatient	1,536	1,243	127	111	269	316	1,140 <sup>a</sup>	815
Mental Health Center - Outpatient	1,016 <sup>a</sup>	729	79	73	169	159	768 <sup>a</sup>	496
Emergency Room	458 <sup>b</sup>	237	28	47	57	58	372 <sup>b</sup>	132
Private Doctor's Office	401 <sup>a</sup>	235	26	35	58	40	316 <sup>a</sup>	160
Self-Help Group	1,527 <sup>a</sup>	1,212	113	82	246	292	1,167 <sup>a</sup>	838
Prison/Jail	198	168	29	21	47	46	123	101

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Received Any Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 5.68B Locations Received Any Illicit Drug or Alcohol Treatment in the Past Year among Persons Who Received Any Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year, by Age Group: Percentages, 2002 and 2003**

Location of Treatment <sup>1</sup>	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
Hospital as an Inpatient	36.6	31.3	38.6	34.3	26.8	25.3	38.9	33.3
Rehabilitation Facility - Inpatient	46.7	40.1	44.5	38.9	36.8	38.1	49.4	41.1
Rehabilitation Facility - Outpatient	65.5	66.3	68.4	66.3	61.7	65.0	66.1	66.8
Mental Health Center - Outpatient	43.3	38.9	42.7	43.8	38.9	32.7	44.5	40.7
Emergency Room	19.5 <sup>a</sup>	12.6	15.3 <sup>a</sup>	27.8	13.1	11.9	21.6 <sup>b</sup>	10.8
Private Doctor's Office	17.1	12.5	14.2	21.0	13.4	8.3	18.3	13.1
Self-Help Group	65.1	64.6	61.1	49.0	56.6	60.1	67.7	68.6
Prison/Jail	8.5	9.0	15.5	12.8	10.8	9.4	7.1	8.3

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Received Any Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level.

<sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents could indicate multiple locations of treatment; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.