| Table 5.84A | Needed and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic |
|-------------|--|
|             | Characteristics: Numbers in Thousands, 2002 and 2003   |

|  | NEEDED           | TREATMENT F | Percentage Who Received                       |       |                  |                                       |  |      |
|--|------------------|-------------|---|-------|------------------|---------------------------------------|--|------|
|  | Total            |             | Received Treatment at a<br>Specialty Facility |       | Treatment        | t Receive<br>at a Specialty<br>cility | Treatment at a Specialty<br>Facility among Persons<br>Who Needed Treatment |      |
| Demographic Characteristic                   | 2002             | 2003        | 2002  | 2003  | 2002             | 2003                                  | 2002   | 2003 |
| TOTAL  | 18,638           | 18,215      | 1,549   | 1,298 | 17,089           | 16,917                                | 8.3  | 7.1  |
| AGE  |                  |             |   |       |                  |                                       |  |      |
| 12-17  | 1,495            | 1,501       | 121   | 95    | 1,374            | 1,406                                 | 8.1  | 6.3  |
| 18-25  | 5,584            | 5,589       | 270   | 305   | 5,313            | 5,284                                 | 4.8  | 5.5  |
| 26 or Older                                  | 11,559           | 11,125      | 1,157   | 898   | 10,402           | 10,228                                | 10.0   | 8.1  |
| GENDER                                       |                  |             |   |       |                  |                                       |  |      |
| Male   | 12,686           | 12,118      | 1,100   | 897   | 11,586           | 11,221                                | 8.7  | 7.4  |
| Female                                       | 5,952            | 6,097       | 449   | 401   | 5,503            | 5,696                                 | 7.5  | 6.6  |
| HISPANIC ORIGIN AND RACE                     |                  |             |   |       |                  |                                       |  |      |
| Not Hispanic or Latino                       | 16,131           | 15,761      | 1,410   | 1,159 | 14,721           | 14,602                                | 8.7  | 7.4  |
| White  | 13,306           | 12,955      | 1,120 <sup>a</sup>                            | 832   | 12,187           | 12,123                                | 8.4 <sup>a</sup>   | 6.4  |
| Black or African American                    | 1,995            | 1,793       | 236   | 251   | 1,759            | 1,542                                 | 11.8   | 14.0 |
| American Indian or Alaska<br>Native          | 186              | 201         | 33  | 38    | 153              | 163                                   | *  | *    |
| Native Hawaiian or Other<br>Pacific Islander | 58               | 48          | 0   | 5     | 57               | 43                                    | *  | *    |
| Asian  | 327 <sup>a</sup> | 541         | 4   | 7     | 324 <sup>a</sup> | 534                                   | *  | *    |
| Two or More Races                            | 258              | 223         | 17  | 26    | 241              | 197                                   | 6.7  | *    |
| Hispanic or Latino                           | 2,507            | 2,455       | 139   | 140   | 2,368            | 2,315                                 | 5.5  | 5.7  |

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level. <sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

# Table 5.84BNeeded and Received Treatment for an Alcohol Problem in the Past Year among Persons Aged 12 or Older, by Demographic<br/>Characteristics: Percentages, 2002 and 2003

|  | NEEDED TI | Percentage Who Received |   |      |                               |               |  |      |
|--|-----------|-------------------------|---|------|-------------------------------|---------------|--|------|
|  | Total     |                         | Received Treatment at a<br>Specialty Facility |      | Did Not<br>Treatment a<br>Fac | t a Specialty | Treatment at a Specialty<br>Facility among Persons<br>Who Needed Treatment |      |
| Demographic Characteristic                   | 2002      | 2003                    | 2002  | 2003 | 2002                          | 2003          | 2002   | 2003 |
| TOTAL  | 7.9       | 7.7                     | 0.7   | 0.5  | 7.3                           | 7.1           | 8.3  | 7.1  |
| AGE  |           |                         |   |      |                               |               |  |      |
| 12-17  | 6.0       | 6.0                     | 0.5   | 0.4  | 5.6                           | 5.6           | 8.1  | 6.3  |
| 18-25  | 18.0      | 17.6                    | 0.9   | 1.0  | 17.1                          | 16.7          | 4.8  | 5.5  |
| 26 or Older                                  | 6.4       | 6.1                     | 0.6   | 0.5  | 5.8                           | 5.7           | 10.0   | 8.1  |
| GENDER                                       |           |                         |   |      |                               |               |  |      |
| Male   | 11.2      | 10.5                    | 1.0   | 0.8  | 10.2                          | 9.8           | 8.7  | 7.4  |
| Female                                       | 4.9       | 5.0                     | 0.4   | 0.3  | 4.5                           | 4.6           | 7.5  | 6.6  |
| HISPANIC ORIGIN AND RACE                     |           |                         |   |      |                               |               |  |      |
| Not Hispanic or Latino                       | 7.8       | 7.6                     | 0.7   | 0.6  | 7.1                           | 7.0           | 8.7  | 7.4  |
| White  | 8.0       | 7.8                     | $0.7^{\mathrm{a}}$                            | 0.5  | 7.4                           | 7.3           | 8.4 <sup>a</sup>   | 6.4  |
| Black or African American                    | 7.4       | 6.5                     | 0.9   | 0.9  | 6.6                           | 5.6           | 11.8   | 14.0 |
| American Indian or Alaska<br>Native          | 12.6      | 16.0                    | 2.2   | 3.0  | 10.4                          | 13.0          | *  | *    |
| Native Hawaiian or Other<br>Pacific Islander | 7.1       | 9.8                     | 0.1   | 1.1  | 7.1                           | 8.7           | *  | *    |
| Asian  | 3.6       | 5.5                     | 0.0   | 0.1  | 3.6                           | 5.5           | *  | *    |
| Two or More Races                            | 10.3      | 8.7                     | 0.7   | 1.0  | 9.6                           | 7.7           | 6.7  | *    |
| Hispanic or Latino                           | 8.6       | 8.2                     | 0.5   | 0.5  | 8.1                           | 7.7           | 5.5  | 5.7  |

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an alcohol problem if they met at least one of three criteria during the past year: (1) dependent on alcohol; (2) abuse of alcohol; or (3) received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level. <sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

| Table 5.85A | Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older |
|-------------|--|
|             | Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Numbers in Thousands,  |
|             | 2002 and 2003  |

|                            | Total Needing But<br>Not Receiving<br>Treatment <sup>1</sup> |           |       |      |             |      |                |      |   |        |
|----------------------------|--|-----------|-------|------|-------------|------|----------------|------|---|--------|
|                            |  |           | Total |      | Made Effort |      | Made No Effort |      | Did Not Feel Need<br>for Treatment <sup>2</sup> |        |
| Demographic Characteristic | 2002   | 2002 2003 |       | 2003 | 2002        | 2003 | 2002           | 2003 | 2002  | 2003   |
| TOTAL                      | 17,089   | 16,917    | 761   | 642  | 266         | 173  | 495            | 469  | 16,328  | 16,275 |
| AGE                        |  |           |       |      |             |      |                |      |   |        |
| 12-17                      | 1,374  | 1,406     | 27    | 32   | *           | *    | *              | *    | 1,347   | 1,373  |
| 18-25                      | 5,313  | 5,284     | 148   | 145  | *           | 31   | *              | 114  | 5,166   | 5,138  |
| 26 or Older                | 10,402   | 10,228    | 587   | 464  | *           | *    | *              | *    | 9,815   | 9,764  |
| GENDER                     |  |           |       |      |             |      |                |      |   |        |
| Male                       | 11,586   | 11,221    | 468   | 409  | *           | *    | *              | *    | 11,118  | 10,813 |
| Female                     | 5,503  | 5,696     | 293   | 233  | *           | *    | *              | *    | 5,210   | 5,463  |

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level. <sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup>Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

<sup>2</sup> Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

## Table 5.85B Perceived Need for Alcohol Treatment and Whether Made an Effort to Get Treatment in the Past Year among Persons Aged 12 or Older Classified as Needing But Not Receiving Treatment for an Alcohol Problem, by Demographic Characteristics: Percentages, 2002 and 2003

|                            | Total Needing But<br>Not Receiving<br>Treatment <sup>1</sup> |       | FELT NEED FOR TREATMENT <sup>2</sup> |      |             |      |                |      |   |      |
|----------------------------|--|-------|--------------------------------------|------|-------------|------|----------------|------|---|------|
|                            |  |       | Total                                |      | Made Effort |      | Made No Effort |      | Did Not Feel Need<br>for Treatment <sup>2</sup> |      |
| Demographic Characteristic | haracteristic 2002 2003                                      |       | 2002                                 | 2003 | 2002        | 2003 | 2002           | 2003 | 2002  | 2003 |
| TOTAL                      | 100.0  | 100.0 | 4.5                                  | 3.8  | 35.0        | 26.9 | 65.0           | 73.1 | 95.5  | 96.2 |
| AGE                        |  |       |                                      |      |             |      |                |      |   |      |
| 12-17                      | 100.0  | 100.0 | 1.9                                  | 2.3  | *           | *    | *              | *    | 98.1  | 97.7 |
| 18-25                      | 100.0  | 100.0 | 2.8                                  | 2.8  | *           | 21.5 | *              | 78.5 | 97.2  | 97.2 |
| 26 or Older                | 100.0  | 100.0 | 5.6                                  | 4.5  | *           | *    | *              | *    | 94.4  | 95.5 |
| GENDER                     |  |       |                                      |      |             |      |                |      |   |      |
| Male                       | 100.0  | 100.0 | 4.0                                  | 3.6  | *           | *    | *              | *    | 96.0  | 96.4 |
| Female                     | 100.0  | 100.0 | 5.3                                  | 4.1  | *           | *    | *              | *    | 94.7  | 95.9 |

\*Low precision; no estimate reported.

<sup>a</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.05 level. <sup>b</sup>Difference between estimate and 2003 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Needing But Not Receiving Treatment refers to respondents classified as needing treatment for alcohol, but have not received treatment for an alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers).

<sup>2</sup> Felt Need for Treatment includes persons who did not receive but felt they needed treatment for an alcohol problem, as well as persons who received treatment at a location other than a specialty facility but felt they needed additional treatment.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.