| | SUBST | ANCE FOR WI | HICH TREATM | ENT AT A SPE | CIALTY FACIL | ITY WAS REC | EIVED IN PAST | YEAR | |
|--------------|----------|-------------|-------------|--------------|---------------------|-----------------------|--|-------------|--|
| | Any Illi | cit Drug | Alc | ohol | Both Any Ill Alc | icit Drug and ohol | Any Illicit Drug or Alcohol ¹ | | |
| Age Category | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | |
| TOTAL | 121 | 91 | 124 | 102 | 88 | 71 | 151 | 118 | |
| 12 | 1 | 2 | 2 | 2 | 1 | 2 | 2 | 3 | |
| 13 | 2 | 1 | 0 | 2 | 0 | * | 2 | 2 | |
| 14 | 2 | 5 | 3 | 1 | 1 | 1 | 3 | 5 | |
| 15 | 6 | 5 | 6 | 6 | 5 | 4 | 7 | 7 | |
| 16 | 10 | 8 | 11 | 6 | 10 | 6 | 12 | 10 | |
| 17 | 11 | 7 | 10 | 7 | 9 | 5 | 12 | 9 | |
| 18 | 10 | 11 | 10 | 10 | 8 | 8 | 11 | 13 | |
| 19 | 7 | 11 | 9 | 7 | 6 | 6 | 11 | 11 | |
| 20 | 9 | 9 | 7 | 7 | 6 | 6 | 10 | 10 | |
| 21 | 10 | 11 | 9 | 8 | 7 | 7 | 11 | 12 | |
| 22 | 10 | 9 | 9 | 10 | 7 | 7 | 12 | 12 | |
| 23 | 8 | 10 | 8 | 10 | 5 | 9 | 10 | 12 | |
| 24 | 6 | 11 | 7 | 11 | 3 | 9 | 9 | 12 | |
| 25 | 6 | 8 | 8 | 8 | 5 | 5 | 9 | 10 | |
| 26-29 | 36 | 31 | 37 | 33 | 24 | 25 | 46 | 37 | |
| 30-34 | 32 | 21 | 32 | 27 | 20 | 14 | 42 | 32 | |
| 35-39 | 48 | 44 | 47 | 45 | 29 | 31 | 61 | 55 | |
| 40-44 | 43 | 34 | 50 | 42 | 32 | 28 | 58 | 48 | |
| 45-49 | 46 | 21 | 54 | 35 | 36 | 15 | 61 | 38 | |
| 50-54 | 67 | 33 | 60 | 30 | 53 | 28 | 77 | 36 | |
| 55-59 | * | 11 | 24 | 15 | * | 11 | 24 | 15 | |
| 60-64 | * | * | 16 | * | * | * | 16 | * | |
| 65 or Older | 22 | * | 14 | 19 | * | * | 26 | 19 | |

 Table 5.59C
 Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s).

| | SUBST | ANCE FOR WI | HICH TREATM | ENT AT A SPE | CIALTY FACIL | LITY WAS REC | EIVED IN PAS | Г YEAR | |
|--------------|----------|-------------|-------------|--------------|--------------|-----------------------|--|--------|--|
| | Any Illi | cit Drug | Alc | ohol | | icit Drug and ohol | Any Illicit Drug or Alcohol ¹ | | |
| Age Category | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | |
| TOTAL | 0.05 | 0.04 | 0.05 | 0.04 | 0.04 | 0.03 | 0.06 | 0.05 | |
| 12 | 0.01 | 0.05 | 0.05 | 0.05 | 0.01 | 0.05 | 0.05 | 0.07 | |
| 13 | 0.04 | 0.01 | 0.01 | 0.04 | 0.01 | * | 0.04 | 0.04 | |
| 14 | 0.04 | 0.12 | 0.06 | 0.03 | 0.03 | 0.03 | 0.07 | 0.12 | |
| 15 | 0.15 | 0.12 | 0.15 | 0.14 | 0.13 | 0.09 | 0.16 | 0.17 | |
| 16 | 0.26 | 0.20 | 0.27 | 0.15 | 0.24 | 0.13 | 0.29 | 0.23 | |
| 17 | 0.28 | 0.18 | 0.25 | 0.18 | 0.22 | 0.13 | 0.30 | 0.22 | |
| 18 | 0.22 | 0.24 | 0.22 | 0.21 | 0.19 | 0.17 | 0.26 | 0.27 | |
| 19 | 0.18 | 0.27 | 0.23 | 0.17 | 0.14 | 0.16 | 0.26 | 0.27 | |
| 20 | 0.22 | 0.22 | 0.18 | 0.18 | 0.15 | 0.14 | 0.25 | 0.25 | |
| 21 | 0.25 | 0.29 | 0.21 | 0.20 | 0.17 | 0.17 | 0.28 | 0.32 | |
| 22 | 0.25 | 0.22 | 0.23 | 0.26 | 0.19 | 0.18 | 0.32 | 0.29 | |
| 23 | 0.20 | 0.26 | 0.21 | 0.25 | 0.14 | 0.21 | 0.25 | 0.30 | |
| 24 | 0.17 | 0.29 | 0.19 | 0.30 | 0.09 | 0.25 | 0.26 | 0.33 | |
| 25 | 0.19 | 0.23 | 0.23 | 0.21 | 0.16 | 0.14 | 0.26 | 0.27 | |
| 26-29 | 0.24 | 0.21 | 0.24 | 0.22 | 0.16 | 0.17 | 0.30 | 0.25 | |
| 30-34 | 0.16 | 0.11 | 0.16 | 0.13 | 0.10 | 0.07 | 0.21 | 0.16 | |
| 35-39 | 0.22 | 0.21 | 0.21 | 0.22 | 0.13 | 0.15 | 0.28 | 0.26 | |
| 40-44 | 0.20 | 0.15 | 0.23 | 0.19 | 0.15 | 0.12 | 0.26 | 0.21 | |
| 45-49 | 0.21 | 0.10 | 0.26 | 0.16 | 0.17 | 0.07 | 0.29 | 0.18 | |
| 50-54 | 0.36 | 0.16 | 0.32 | 0.15 | 0.28 | 0.14 | 0.41 | 0.18 | |
| 55-59 | * | 0.07 | 0.16 | 0.10 | * | 0.07 | 0.16 | 0.10 | |
| 60-64 | * | * | 0.13 | * | * | * | 0.13 | * | |
| 65 or Older | 0.07 | * | 0.04 | 0.06 | * | * | 0.08 | 0.06 | |

 Table 5.59D
 Received Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s).

| | SUBST | ANCE FOR W | HICH TREATM | ENT AT A SPI | ECIALTY FACIL | ITY WAS REC | CEIVED IN PAST | YEAR |
|--------------|-----------|------------|-------------|--------------|---------------|-----------------------|--|------|
| | Any Illio | cit Drug | Alco | ohol | | icit Drug and phol | Any Illicit Drug or Alcohol ¹ | |
| Age Category | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0394 | | 0.1101 | | 0.2940 | | 0.0110 | |
| 12 | 0.3143 | | 0.6865 | | 0.4911 | | 0.4889 | |
| 13 | 0.0693 | | 0.3088 | | * | | 0.7184 | |
| 14 | 0.1629 | | 0.0728 | | 0.5602 | | 0.5855 | |
| 15 | 0.5893 | | 0.9050 | | 0.3599 | | 0.7014 | |
| 16 | 0.6812 | | 0.1904 | | 0.3195 | | 0.6913 | |
| 17 | 0.0307 | | 0.6079 | | 0.2888 | | 0.1082 | |
| 18 | 0.5845 | | 0.7463 | | 0.8231 | | 0.6176 | |
| 19 | 0.4154 | | 0.2615 | | 0.9954 | | 0.8509 | |
| 20 | 0.8479 | | 0.6887 | | 0.9054 | | 0.9730 | |
| 21 | 0.6721 | | 0.2269 | | 0.2650 | | 0.6264 | |
| 22 | 0.9392 | | 0.1410 | | 0.6753 | | 0.6608 | |
| 23 | 0.4007 | | 0.5400 | | 0.2868 | | 0.6065 | |
| 24 | 0.0672 | | 0.1856 | | 0.0275 | | 0.3285 | |
| 25 | 0.5643 | | 0.8591 | | 0.8648 | | 0.5875 | |
| 26-29 | 0.4176 | | 0.2884 | | 0.5319 | | 0.2209 | |
| 30-34 | 0.1884 | | 0.5543 | | 0.3867 | | 0.2204 | |
| 35-39 | 0.6188 | | 0.7591 | | 0.5130 | | 0.8672 | |
| 40-44 | 0.3663 | | 0.9149 | | 0.9430 | | 0.4878 | |
| 45-49 | 0.0014 | | 0.1634 | | 0.0196 | | 0.0267 | |
| 50-54 | 0.4925 | | 0.5323 | | 0.7552 | | 0.2842 | |
| 55-59 | * | | 0.6405 | | * | | 0.6405 | |
| 60-64 | * | | * | | * | | * | |
| 65 or Older | * | | 0.8300 | | * | | 0.4898 | |

Table 5.59PReceived Substance Use Treatment at a Specialty Facility in the Past Year, by Detailed Age Categories: P-Values from Tests of
Differences of Percentages, 2003 Versus 2002

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s).

| Table 5.60C | Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic |
|-------------|--|
| | Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003 |

| | SUBSTANCE | E FOR WHIC | H TREATME | NT AT A SPE | CIALTY FAC | ILITY WAS R | ECEIVED IN | PAST YEAR |
|--|-----------|------------|-----------|-------------|------------|------------------------|---|-----------|
| | Any Illi | cit Drug | Alcohol | | | Illicit Drug Icohol | Any Illicit Drug or Alcohol ¹ | |
| Demographic Characteristic | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 121 | 91 | 124 | 102 | 88 | 71 | 151 | 118 |
| AGE | | | | | | | | |
| 12-17 | 18 | 14 | 17 | 12 | 16 | 9 | 19 | 17 |
| 18-25 | 25 | 29 | 24 | 26 | 18 | 22 | 30 | 33 |
| 26 or Older | 116 | 84 | 121 | 96 | 84 | 67 | 145 | 110 |
| GENDER | | | | | | | | |
| Male | 95 | 75 | 110 | 85 | 82 | 60 | 123 | 97 |
| Female | 77 | 50 | 53 | 55 | 30 | 38 | 89 | 65 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 116 | 89 | 120 | 97 | 85 | 69 | 146 | 113 |
| White | 102 | 69 | 109 | 73 | 74 | 45 | 129 | 90 |
| Black or African American | 53 | 50 | 48 | 57 | 40 | 49 | 66 | 59 |
| American Indian or Alaska Native | 11 | 9 | 11 | 13 | 11 | 9 | 11 | 13 |
| Native Hawaiian or Other Pacific Islander | 1 | * | 0 | 4 | 0 | * | 1 | 4 |
| Asian | 4 | 6 | 4 | 6 | 4 | 6 | 4 | 7 |
| Two or More Races | 11 | 9 | 7 | 9 | 6 | 8 | 12 | 10 |
| Hispanic or Latino | 36 | 21 | 30 | 33 | 23 | 14 | 41 | 36 |

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

40715

| Table 5.60D | Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic |
|-------------|--|
| | Characteristics: Standard Errors of Percentages, 2002 and 2003 |

| | SUBSTANC | E FOR WHIC | H TREATME | NT AT A SPE | CIALTY FAC | ILITY WAS R | ECEIVED IN | PAST YEAR |
|--|----------|------------------|-----------|-------------|------------|------------------------|---|-----------|
| | Any Illi | Any Illicit Drug | | Alcohol | | Illicit Drug Icohol | Any Illicit Drug or Alcohol ¹ | |
| Demographic Characteristic | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.05 | 0.04 | 0.05 | 0.04 | 0.04 | 0.03 | 0.06 | 0.05 |
| AGE | | | | | | | | |
| 12-17 | 0.07 | 0.05 | 0.07 | 0.05 | 0.06 | 0.04 | 0.08 | 0.07 |
| 18-25 | 0.08 | 0.09 | 0.08 | 0.08 | 0.06 | 0.07 | 0.10 | 0.10 |
| 26 or Older | 0.06 | 0.05 | 0.07 | 0.05 | 0.05 | 0.04 | 0.08 | 0.06 |
| GENDER | | | | | | | | |
| Male | 0.08 | 0.06 | 0.10 | 0.07 | 0.07 | 0.05 | 0.11 | 0.08 |
| Female | 0.06 | 0.04 | 0.04 | 0.04 | 0.02 | 0.03 | 0.07 | 0.05 |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.06 | 0.04 | 0.06 | 0.05 | 0.04 | 0.03 | 0.07 | 0.05 |
| White | 0.06 | 0.04 | 0.07 | 0.04 | 0.04 | 0.03 | 0.08 | 0.05 |
| Black or African American | 0.20 | 0.18 | 0.18 | 0.21 | 0.15 | 0.18 | 0.24 | 0.21 |
| American Indian or Alaska Native | 0.72 | 0.74 | 0.74 | 1.00 | 0.71 | 0.71 | 0.75 | 1.04 |
| Native Hawaiian or Other Pacific Islander | 0.08 | * | 0.04 | 0.78 | 0.04 | * | 0.15 | 0.78 |
| Asian | 0.04 | 0.07 | 0.04 | 0.06 | 0.04 | 0.06 | 0.04 | 0.07 |
| Two or More Races | 0.43 | 0.34 | 0.29 | 0.36 | 0.26 | 0.33 | 0.47 | 0.37 |
| Hispanic or Latino | 0.12 | 0.07 | 0.10 | 0.11 | 0.08 | 0.05 | 0.14 | 0.12 |

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

40715

| Table 5.60P | Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Demographic |
|-------------|--|
| | Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002 |

| | SUBSTANCE | E FOR WHIC | H TREATME | NT AT A SPE | CIALTY FAC | ILITY WAS R | RECEIVED IN | PAST YEAR |
|--|------------------|------------|-----------|-------------|------------|------------------------|---|-----------|
| | Any Illicit Drug | | Alcohol | | • | Illicit Drug Icohol | Any Illicit Drug or Alcohol ¹ | |
| Demographic Characteristic | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0394 | | 0.1101 | | 0.2940 | | 0.0110 | |
| AGE | | | | | | | | |
| 12-17 | 0.1751 | | 0.2060 | | 0.1329 | | 0.4461 | |
| 18-25 | 0.2247 | | 0.4130 | | 0.3471 | | 0.3730 | |
| 26 or Older | 0.0218 | | 0.0933 | | 0.2738 | | 0.0054 | |
| GENDER | | | | | | | | |
| Male | 0.3973 | | 0.1246 | | 0.3539 | | 0.1190 | |
| Female | 0.0192 | | 0.5341 | | 0.5867 | | 0.0299 | |
| HISPANIC ORIGIN AND RACE | | | | | | | | |
| Not Hispanic or Latino | 0.1216 | | 0.1031 | | 0.4684 | | 0.0176 | |
| White | 0.2619 | | 0.0327 | | 0.2371 | | 0.0357 | |
| Black or African American | 0.2331 | | 0.8929 | | 0.8191 | | 0.1858 | |
| American Indian or Alaska Native | 0.9725 | | 0.5149 | | 0.9761 | | 0.4639 | |
| Native Hawaiian or Other Pacific Islander | * | | 0.1840 | | * | | 0.2846 | |
| Asian | 0.9366 | | 0.7138 | | 0.7653 | | 0.8015 | |
| Two or More Races | 0.8464 | | 0.5010 | | 0.5021 | | 0.7865 | |
| Hispanic or Latino | 0.0363 | | 0.9426 | | 0.1442 | | 0.3839 | |

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

40715

| | SUBSTANC | SUBSTANCE FOR WHICH TREATMENT AT A SPECIALTY FACILITY WAS RECEIVED IN PAS | | | | | | |
|---------------------------|----------|---|------|---------|------|--------------------------------------|------|--------------------------------|
| | | Any Illicit Drug | | Alcohol | | Both Any Illicit Drug and Alcohol | | t Drug or bhol ¹ |
| Geographic Characteristic | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 121 | 91 | 124 | 102 | 88 | 71 | 151 | 118 |
| GEOGRAPHIC DIVISION | | | | | | | | |
| Northeast | 79 | 40 | 76 | 45 | 59 | 31 | 91 | 52 |
| New England | 57 | 17 | 64 | 20 | 52 | 10 | 68 | 24 |
| Middle Atlantic | 54 | 36 | 40 | 41 | 28 | 30 | 61 | 46 |
| Midwest | 44 | 35 | 55 | 49 | 40 | 28 | 65 | 55 |
| East North Central | 40 | 26 | 45 | 38 | 36 | 17 | 49 | 44 |
| West North Central | 19 | 24 | 31 | 32 | 17 | 22 | 43 | 33 |
| South | 58 | 64 | 69 | 62 | 43 | 52 | 79 | 72 |
| South Atlantic | 43 | 56 | 58 | 52 | 34 | 47 | 63 | 60 |
| East South Central | 27 | 26 | 14 | 27 | 8 | 20 | 30 | 31 |
| West South Central | 27 | 18 | 36 | 21 | 25 | 11 | 37 | 27 |
| West | 56 | 36 | 40 | 46 | 29 | 24 | 62 | 53 |
| Mountain | 17 | 18 | 18 | 19 | 7 | 8 | 24 | 25 |
| Pacific | 53 | 31 | 36 | 42 | 28 | 23 | 57 | 47 |
| COUNTY TYPE ² | | | | | | | | |
| Large Metro | 88 | 77 | 87 | 79 | 54 | 61 | 112 | 92 |
| Small Metro | 77 | 35 | 81 | 46 | 66 | 20 | 91 | 56 |
| 250K - 1 Mil. Pop. | 48 | 33 | 56 | 41 | 38 | 18 | 65 | 51 |
| < 250K Pop. | 60 | 15 | 58 | 22 | 55 | 9 | 64 | 25 |
| Nonmetro | 29 | 33 | 36 | 44 | 26 | 31 | 38 | 46 |
| Urbanized | 19 | 29 | 24 | 37 | 18 | 28 | 25 | 38 |
| Less Urbanized | 21 | 14 | 26 | 21 | 18 | 11 | 28 | 23 |
| Completely Rural | 4 | 9 | 7 | 10 | 4 | 9 | 9 | 10 |

 Table 5.61C
 Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s). ² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

| | SUBSTANC | E FUK WHIC | n ikeaime. T | NI AI A SPI | ECIALTY FAC | | | |
|---------------------------|----------|------------------|-----------------|-------------|-------------|--------------------------------------|------|--------------------------------|
| Geographic Characteristic | Any Illi | Any Illicit Drug | | Alcohol | | Both Any Illicit Drug and Alcohol | | t Drug or bhol ¹ |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.05 | 0.04 | 0.05 | 0.04 | 0.04 | 0.03 | 0.06 | 0.05 |
| GEOGRAPHIC DIVISION | | | | | | | | |
| Northeast | 0.17 | 0.09 | 0.17 | 0.10 | 0.13 | 0.07 | 0.20 | 0.11 |
| New England | 0.48 | 0.14 | 0.55 | 0.16 | 0.44 | 0.08 | 0.58 | 0.20 |
| Middle Atlantic | 0.16 | 0.11 | 0.12 | 0.12 | 0.08 | 0.09 | 0.18 | 0.14 |
| Midwest | 0.08 | 0.07 | 0.10 | 0.09 | 0.07 | 0.05 | 0.12 | 0.10 |
| East North Central | 0.11 | 0.07 | 0.12 | 0.10 | 0.10 | 0.05 | 0.13 | 0.12 |
| West North Central | 0.12 | 0.15 | 0.20 | 0.20 | 0.10 | 0.14 | 0.27 | 0.21 |
| South | 0.07 | 0.08 | 0.08 | 0.07 | 0.05 | 0.06 | 0.09 | 0.09 |
| South Atlantic | 0.10 | 0.13 | 0.13 | 0.12 | 0.08 | 0.11 | 0.14 | 0.13 |
| East South Central | 0.19 | 0.19 | 0.10 | 0.19 | 0.06 | 0.14 | 0.21 | 0.22 |
| West South Central | 0.10 | 0.07 | 0.14 | 0.08 | 0.10 | 0.04 | 0.14 | 0.10 |
| West | 0.11 | 0.07 | 0.08 | 0.09 | 0.05 | 0.05 | 0.12 | 0.10 |
| Mountain | 0.11 | 0.12 | 0.12 | 0.12 | 0.05 | 0.05 | 0.16 | 0.16 |
| Pacific | 0.14 | 0.08 | 0.10 | 0.11 | 0.07 | 0.06 | 0.15 | 0.12 |
| COUNTY TYPE ² | | | | | | | | |
| Large Metro | 0.07 | 0.06 | 0.07 | 0.06 | 0.04 | 0.05 | 0.09 | 0.07 |
| Small Metro | 0.11 | 0.05 | 0.12 | 0.07 | 0.10 | 0.03 | 0.13 | 0.08 |
| 250K - 1 Mil. Pop. | 0.11 | 0.07 | 0.12 | 0.09 | 0.08 | 0.04 | 0.14 | 0.11 |
| < 250K Pop. | 0.25 | 0.06 | 0.24 | 0.09 | 0.23 | 0.04 | 0.27 | 0.10 |
| Nonmetro | 0.07 | 0.08 | 0.09 | 0.11 | 0.06 | 0.08 | 0.10 | 0.11 |
| Urbanized | 0.12 | 0.16 | 0.15 | 0.21 | 0.11 | 0.16 | 0.16 | 0.21 |
| Less Urbanized | 0.11 | 0.07 | 0.13 | 0.12 | 0.09 | 0.06 | 0.14 | 0.13 |
| Completely Rural | 0.09 | 0.21 | 0.15 | 0.24 | 0.08 | 0.21 | 0.20 | 0.24 |

 Table 5.61D
 Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s). ² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

| Geographic Characteristic | | | IICH TREATMENT AT A SP | | | | | |
|---------------------------|------------------|------|------------------------|------|--------------------------------------|------|---|------|
| | Any Illicit Drug | | Alcohol | | Both Any Illicit Drug and Alcohol | | Any Illicit Drug or Alcohol ¹ | |
| | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 | 2002 | 2003 |
| TOTAL | 0.0394 | | 0.1101 | | 0.2940 | | 0.0110 | |
| GEOGRAPHIC DIVISION | | | | | | | | |
| Northeast | 0.2333 | | 0.3630 | | 0.8890 | | 0.1034 | |
| New England | 0.3705 | | 0.1612 | | 0.4033 | | 0.1229 | |
| Middle Atlantic | 0.4240 | | 0.8077 | | 0.3851 | | 0.3906 | |
| Midwest | 0.2295 | | 0.2083 | | 0.1114 | | 0.1596 | |
| East North Central | 0.0893 | | 0.1798 | | 0.0183 | | 0.3088 | |
| West North Central | 0.8135 | | 0.7185 | | 0.8178 | | 0.3300 | |
| South | 0.5543 | | 0.1663 | | 0.6660 | | 0.1704 | |
| South Atlantic | 0.7569 | | 0.2511 | | 0.7839 | | 0.2874 | |
| East South Central | 0.9406 | | 0.4070 | | 0.1943 | | 0.9895 | |
| West South Central | 0.2835 | | 0.1295 | | 0.1327 | | 0.2121 | |
| West | 0.1818 | | 0.3729 | | 0.8799 | | 0.5827 | |
| Mountain | 0.1655 | | 0.3479 | | 0.2155 | | 0.1327 | |
| Pacific | 0.0414 | | 0.6010 | | 0.7949 | | 0.1703 | |
| COUNTY TYPE ² | | | | | | | | |
| Large Metro | 0.3030 | | 0.3902 | | 0.9140 | | 0.0651 | |
| Small Metro | 0.0099 | | 0.0557 | | 0.0351 | | 0.0263 | |
| 250K - 1 Mil. Pop. | 0.0193 | | 0.0861 | | 0.0908 | | 0.0408 | |
| < 250K Pop. | 0.1764 | | 0.3423 | | 0.1702 | | 0.3217 | |
| Nonmetro | 0.6048 | | 0.7740 | | 0.3878 | | 0.9114 | |
| Urbanized | 0.3905 | | 0.4912 | | 0.4137 | | 0.3425 | |
| Less Urbanized | 0.5091 | | 0.4822 | | 0.8795 | | 0.2845 | |
| Completely Rural | 0.3390 | | 0.5948 | | 0.3263 | | 0.9928 | |

 Table 5.61P
 Received Substance Use Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older, by Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

*Low precision; no estimate reported.

NOTE: Received Substance Use Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient), a rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Estimates include persons who received treatment specifically for any illicit drug or alcohol, as well as persons who received treatment but did not specify for what substance(s). ² Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.