

**Table 6.1C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003**

Age Category	Total		GENDER			
			Male		Female	
	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	436	432	252	264	328	345
18	32	37	20	27	25	25
19	31	37	19	22	25	28
20	33	34	22	23	23	25
21	32	33	21	22	24	24
22	30	31	21	20	23	23
23	31	31	20	20	24	25
24	27	30	18	23	21	24
25	29	30	18	16	23	24
26-29	105	120	67	74	76	94
30-34	128	116	80	69	98	93
35-39	152	126	116	73	99	109
40-44	140	139	78	80	114	107
45-49	123	146	78	96	95	111
50-54	165	180	95	121	138	138
55-59	138	126	83	90	112	93
60-64	83	101	54	71	63	73
65 or Older	179	195	87	112	153	156

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.1D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003**

Age Category	Total		GENDER			
			Male		Female	
	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.21	0.20	0.25	0.26	0.30	0.31
18	0.72	0.80	0.92	1.08	1.19	1.17
19	0.77	0.95	0.93	1.18	1.23	1.37
20	0.83	0.86	1.05	1.14	1.19	1.26
21	0.80	0.85	1.05	1.16	1.20	1.22
22	0.77	0.78	1.05	0.99	1.21	1.15
23	0.82	0.77	1.04	1.03	1.30	1.19
24	0.78	0.84	1.08	1.21	1.17	1.33
25	0.88	0.84	1.16	0.90	1.26	1.33
26-29	0.68	0.81	0.90	0.97	0.97	1.30
30-34	0.65	0.58	0.80	0.71	0.98	0.89
35-39	0.69	0.60	1.07	0.70	0.89	1.03
40-44	0.64	0.61	0.74	0.72	1.00	0.93
45-49	0.58	0.68	0.75	0.92	0.89	1.01
50-54	0.88	0.90	1.06	1.24	1.41	1.35
55-59	0.94	0.85	1.23	1.22	1.39	1.23
60-64	0.72	0.86	0.92	1.30	1.10	1.15
65 or Older	0.53	0.57	0.59	0.77	0.79	0.79

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.1P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Gender and Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Age Category	Total		GENDER			
			Male		Female	
	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.0017		0.0361		0.0119	
18	0.1658		0.3218		0.2451	
19	0.0056		0.0669		0.0386	
20	0.1985		0.3673		0.3690	
21	0.8359		0.7952		0.9037	
22	0.2931		0.5135		0.3969	
23	0.5680		0.9708		0.3560	
24	0.0083		0.1157		0.0375	
25	0.3353		0.1085		0.8356	
26-29	0.0476		0.4036		0.0392	
30-34	0.2562		0.2102		0.5076	
35-39	0.0622		0.8216		0.0095	
40-44	0.7362		0.8734		0.8658	
45-49	0.0097		0.0758		0.0750	
50-54	0.3556		0.1009		0.9439	
55-59	0.6958		0.6866		0.7856	
60-64	0.4444		0.5509		0.6824	
65 or Older	0.1817		0.1703		0.4591	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.2C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Demographic Characteristic	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	436	432	93	100	290	280	289	307
<b>GENDER</b>								
Male	252	264	57	64	186	175	160	194
Female	328	345	70	71	216	223	235	238
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	414	415	82	88	279	269	271	287
White	372	388	73	76	244	243	240	275
Black or African American	154	143	28	36	102	98	108	88
American Indian or Alaska Native	34	21	9	5	22	18	*	*
Native Hawaiian or Other Pacific Islander	14	15	*	*	*	*	*	*
Asian	112	83	21	19	93	73	*	21
Two or More Races	54	48	11	10	30	22	*	*
Hispanic or Latino	146	151	43	41	93	112	98	103
<b>EDUCATION</b>								
< High School	209	223	44	46	109	138	167	166
High School Graduate	250	244	52	54	166	159	179	181
Some College	208	212	53	52	153	153	127	133
College Graduate	179	186	29	31	146	139	78	130
<b>CURRENT EMPLOYMENT</b>								
Full-Time	269	291	60	62	234	224	122	182
Part-Time	146	151	46	47	105	108	88	101
Unemployed	99	85	28	31	69	64	*	*
Other <sup>1</sup>	282	271	40	45	125	135	245	222

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.2D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Demographic Characteristic	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	0.30	0.32	0.29	0.28	0.37	0.38
<b>GENDER</b>								
Male	0.25	0.26	0.37	0.40	0.38	0.36	0.44	0.52
Female	0.30	0.31	0.45	0.45	0.42	0.44	0.55	0.54
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.22	0.22	0.32	0.34	0.32	0.31	0.37	0.38
White	0.25	0.26	0.38	0.39	0.36	0.36	0.38	0.43
Black or African American	0.67	0.60	0.68	0.84	0.86	0.83	1.48	1.16
American Indian or Alaska Native	2.60	1.92	4.09	2.57	3.56	3.20	*	*
Native Hawaiian or Other Pacific Islander	1.97	3.53	*	*	*	*	*	*
Asian	1.39	0.94	1.55	1.34	2.03	1.49	*	0.87
Two or More Races	2.53	2.22	2.78	2.80	2.97	2.18	*	*
Hispanic or Latino	0.58	0.58	0.78	0.73	0.65	0.78	1.81	1.81
<b>EDUCATION</b>								
< High School	0.56	0.59	0.64	0.68	0.79	0.94	1.01	1.02
High School Graduate	0.37	0.36	0.49	0.49	0.54	0.53	0.68	0.69
Some College	0.40	0.40	0.54	0.52	0.59	0.60	0.76	0.78
College Graduate	0.34	0.34	0.77	0.78	0.49	0.46	0.40	0.61
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.23	0.25	0.42	0.42	0.32	0.31	0.42	0.58
Part-Time	0.53	0.53	0.58	0.58	0.99	0.96	0.98	1.10
Unemployed	1.31	1.05	1.15	1.16	1.75	1.56	*	*
Other <sup>1</sup>	0.48	0.46	0.65	0.71	0.95	1.01	0.62	0.56

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.2P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Demographic Characteristic	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.0723		0.0157		0.0631	
<b>GENDER</b>								
Male	0.0361		0.2809		0.6245		0.0240	
Female	0.0119		0.1278		0.0075		0.5071	
<b>HISPANIC ORIGIN AND RACE</b>								
Not Hispanic or Latino	0.0186		0.1369		0.1181		0.1030	
White	0.0025		0.4015		0.0477		0.0122	
Black or African American	0.6960		0.3545		0.8879		0.3100	
American Indian or Alaska Native	0.2710		0.4062		0.8833		*	
Native Hawaiian or Other Pacific Islander	0.0821		*		*		*	
Asian	0.4183		0.2625		0.6489		*	
Two or More Races	0.7539		0.1087		0.3369		*	
Hispanic or Latino	0.0056		0.2419		0.0180		0.3342	
<b>EDUCATION</b>								
< High School	0.0451		0.1707		0.0533		0.4837	
High School Graduate	0.0708		0.1553		0.1802		0.4574	
Some College	0.1760		0.9002		0.1222		0.6463	
College Graduate	0.1676		0.3304		0.8110		0.0079	
<b>CURRENT EMPLOYMENT</b>								
Full-Time	0.0050		0.0988		0.1891		0.0052	
Part-Time	0.7102		0.7826		0.6218		0.9251	
Unemployed	0.5553		0.7984		0.6960		*	
Other <sup>1</sup>	0.1186		0.0864		0.0320		0.8102	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.3C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	436	432	93	100	290	280	289	307
<b>GEOGRAPHIC DIVISION</b>								
Northeast	206	194	37	48	138	117	130	133
New England	104	87	22	23	73	63	71	43
Middle Atlantic	176	173	29	41	118	99	109	125
Midwest	192	168	38	42	130	120	127	116
East North Central	166	144	29	36	115	106	106	100
West North Central	97	86	23	22	61	56	70	59
South	271	269	57	52	171	162	183	191
South Atlantic	224	198	38	38	128	116	153	143
East South Central	106	105	28	17	78	74	61	74
West South Central	111	150	32	32	82	86	80	103
West	194	220	53	57	136	155	131	164
Mountain	79	102	25	26	58	63	47	89
Pacific	178	196	46	50	124	142	122	137
<b>COUNTY TYPE<sup>1</sup></b>								
Large Metro	320	300	69	80	229	193	196	228
Small Metro	233	233	49	49	152	162	156	150
250K - 1 Mil. Pop.	182	189	39	39	116	127	124	127
< 250K Pop.	134	122	30	30	90	84	93	85
Nonmetro	183	183	35	33	96	109	147	133
Urbanized	101	135	23	22	62	68	76	100
Less Urbanized	135	113	25	24	67	78	114	84
Completely Rural	61	47	8	7	38	29	49	33

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.3D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	0.30	0.32	0.29	0.28	0.37	0.38
<b>GEOGRAPHIC DIVISION</b>								
Northeast	0.51	0.47	0.67	0.84	0.72	0.62	0.82	0.82
New England	0.98	0.81	1.57	1.59	1.43	1.26	1.71	1.03
Middle Atlantic	0.59	0.58	0.72	0.99	0.83	0.71	0.92	1.05
Midwest	0.40	0.35	0.52	0.57	0.58	0.54	0.70	0.63
East North Central	0.50	0.43	0.59	0.70	0.73	0.68	0.84	0.78
West North Central	0.68	0.60	1.04	0.99	0.92	0.85	1.29	1.06
South	0.36	0.36	0.51	0.46	0.48	0.46	0.64	0.66
South Atlantic	0.57	0.50	0.70	0.68	0.69	0.63	0.99	0.90
East South Central	0.84	0.82	1.47	0.86	1.32	1.26	1.24	1.48
West South Central	0.49	0.65	0.85	0.83	0.74	0.78	0.98	1.24
West	0.41	0.46	0.72	0.77	0.59	0.67	0.79	0.95
Mountain	0.58	0.74	1.15	1.17	0.89	0.96	0.95	1.75
Pacific	0.53	0.58	0.90	0.97	0.74	0.85	1.04	1.13
<b>COUNTY TYPE<sup>1</sup></b>								
Large Metro	0.28	0.27	0.43	0.48	0.41	0.34	0.49	0.57
Small Metro	0.38	0.36	0.49	0.48	0.53	0.56	0.67	0.60
250K - 1 Mil. Pop.	0.45	0.45	0.61	0.58	0.61	0.65	0.81	0.81
< 250K Pop.	0.62	0.55	0.81	0.85	0.92	0.89	1.18	0.93
Nonmetro	0.51	0.51	0.71	0.68	0.63	0.73	0.94	0.83
Urbanized	0.71	0.84	1.04	0.92	0.99	1.00	1.31	1.45
Less Urbanized	0.77	0.70	1.12	1.08	0.91	1.17	1.44	1.15
Completely Rural	1.54	1.28	1.66	1.96	2.37	1.90	2.61	1.87

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.3P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Age Group and Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	Total		AGE GROUP (Years)					
			18-25		26-49		50 or Older	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.0723		0.0157		0.0631	
<b>GEOGRAPHIC DIVISION</b>								
Northeast	0.8462		0.6730		0.2205		0.4196	
New England	0.6059		0.9172		0.8964		0.3414	
Middle Atlantic	0.9735		0.6404		0.1381		0.1576	
Midwest	0.3933		0.7843		0.1781		0.8929	
East North Central	0.6564		0.9630		0.6001		0.8581	
West North Central	0.3681		0.6070		0.0695		0.6262	
South	0.0166		0.0830		0.0626		0.1841	
South Atlantic	0.3875		0.0981		0.5500		0.7223	
East South Central	0.5375		0.5236		0.8300		0.5645	
West South Central	0.0036		0.5218		0.0136		0.0772	
West	0.0038		0.3196		0.0083		0.1428	
Mountain	0.0082		0.5235		0.0083		0.1613	
Pacific	0.0501		0.4264		0.0774		0.3326	
<b>COUNTY TYPE<sup>1</sup></b>								
Large Metro	0.0123		0.0497		0.3279		0.0515	
Small Metro	0.0692		0.2185		0.0463		0.1598	
250K - 1 Mil. Pop.	0.0631		0.3847		0.0835		0.1513	
< 250K Pop.	0.5859		0.4118		0.2014		0.6464	
Nonmetro	0.3858		0.0115		0.1137		0.5558	
Urbanized	0.2456		0.1865		0.5155		0.4257	
Less Urbanized	0.5679		0.0697		0.0537		0.2327	
Completely Rural	0.4827		0.1315		0.5805		0.4904	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.4C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>1</sup></b>	186	169	380	402
Marijuana and Hashish	158	133	395	413
Cocaine	95	78	420	423
Crack	64	45	427	428
Heroin	*	*	434	432
Hallucinogens	51	49	434	425
LSD	20	14	435	431
PCP	*	*	436	432
Ecstasy	44	33	432	430
Inhalants	31	26	436	430
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	132	129	404	420
Pain Relievers	110	104	408	423
Tranquilizers	88	83	412	423
Stimulants	72	51	430	430
Methamphetamine	54	39	434	429
Sedatives	48	*	425	429
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>1</sup></b>	154	138	395	409

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.4D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: Standard Errors of Percentages, 2002 and 2003**

Drug	ILLICIT DRUG USE			
	Yes		No	
	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>1</sup></b>	0.63	0.57	0.21	0.22
Marijuana and Hashish	0.72	0.62	0.21	0.22
Cocaine	1.77	1.42	0.20	0.20
Crack	4.42	3.42	0.20	0.20
Heroin	*	*	0.21	0.20
Hallucinogens	1.35	1.55	0.21	0.20
LSD	2.99	3.36	0.21	0.20
PCP	*	*	0.21	0.20
Ecstasy	1.68	1.83	0.21	0.20
Inhalants	3.08	2.67	0.21	0.20
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	1.07	1.01	0.20	0.21
Pain Relievers	1.21	1.07	0.20	0.21
Tranquilizers	2.05	1.87	0.20	0.20
Stimulants	2.85	2.32	0.21	0.20
Methamphetamine	4.11	3.38	0.21	0.20
Sedatives	5.69	*	0.20	0.20
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>1</sup></b>	0.90	0.82	0.20	0.21

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.4P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Use of Illicit Drugs: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	ILLCIT DRUG USE			
	Yes		No	
	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>1</sup></b>	0.2332		0.0025	
Marijuana and Hashish	0.6050		0.0011	
Cocaine	0.3146		0.0007	
Crack	0.3705		0.0009	
Heroin	*		0.0013	
Hallucinogens	0.6116		0.0008	
LSD	0.6116		0.0015	
PCP	*		0.0016	
Ecstasy	0.1308		0.0006	
Inhalants	0.5755		0.0017	
Nonmedical Use of Any Psychotherapeutic <sup>2</sup>	0.9546		0.0009	
Pain Relievers	0.6437		0.0011	
Tranquilizers	0.9165		0.0012	
Stimulants	0.6583		0.0008	
Methamphetamine	0.6112		0.0010	
Sedatives	*		0.0016	
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>1</sup></b>	0.9005		0.0007	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.5C Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	587	574	201	195	540	506
Marijuana and Hashish	464	457	167	150	435	410
Cocaine	221	232	110	84	191	213
Crack	124	123	77	51	104	107
Heroin	54	47	27	16	47	45
Hallucinogens	145	141	57	54	131	128
LSD	58	41	23	15	53	38
PCP	21	20	10	7	18	19
Ecstasy	123	102	49	34	110	96
Inhalants	80	67	34	26	73	63
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	383	361	154	143	333	323
Pain Relievers	297	304	130	119	253	267
Tranquilizers	222	195	104	93	187	173
Stimulants	144	124	87	60	119	105
Methamphetamine	105	93	61	48	93	74
Sedatives	115	83	68	53	81	67
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	438	425	175	153	385	382

\*Low precision; no estimate reported.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.5D Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Percentages, 2002 and 2003**

Drug	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	0.27	0.25	1.00	0.95	0.27	0.25
Marijuana and Hashish	0.22	0.21	0.86	0.75	0.23	0.20
Cocaine	0.10	0.11	0.60	0.42	0.10	0.11
Crack	0.06	0.06	0.43	0.26	0.05	0.06
Heroin	0.03	0.02	0.15	0.08	0.02	0.02
Hallucinogens	0.07	0.07	0.33	0.27	0.07	0.07
LSD	0.03	0.02	0.13	0.07	0.03	0.02
PCP	0.01	0.01	0.06	0.04	0.01	0.01
Ecstasy	0.06	0.05	0.28	0.17	0.06	0.05
Inhalants	0.04	0.03	0.19	0.13	0.04	0.03
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	0.17	0.17	0.81	0.72	0.17	0.16
Pain Relievers	0.14	0.14	0.69	0.60	0.13	0.14
Tranquilizers	0.10	0.09	0.56	0.47	0.10	0.09
Stimulants	0.07	0.06	0.49	0.31	0.06	0.05
Methamphetamine	0.05	0.04	0.35	0.25	0.05	0.04
Sedatives	0.05	0.04	0.38	0.27	0.04	0.03
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.20	0.19	0.90	0.74	0.19	0.19

\*Low precision; no estimate reported.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.5P Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	0.5876		0.2165		0.5681	
Marijuana and Hashish	0.3442		0.0800		0.4655	
Cocaine	0.9550		0.0739		0.5198	
Crack	0.3562		0.1077		0.8108	
Heroin	0.2823		0.1330		0.5788	
Hallucinogens	0.0004		0.0023		0.0074	
LSD	0.0001		0.0535		0.0003	
PCP	0.9285		0.3825		0.6554	
Ecstasy	0.0000		0.0000		0.0000	
Inhalants	0.5630		0.7870		0.5140	
Nonmedical Use of Any Psychotherapeutic <sup>3</sup>	0.7429		0.1806		0.6025	
Pain Relievers	0.1889		0.3119		0.1220	
Tranquilizers	0.6296		0.4632		0.5670	
Stimulants	0.0513		0.0505		0.1867	
Methamphetamine	0.1943		0.1107		0.4631	
Sedatives	0.2878		0.7620		0.1541	
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.5770		0.0676		0.7811	

\*Low precision; no estimate reported.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>3</sup> Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.6C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco and Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	TOBACCO OR ALCOHOL USE			
	Yes		No	
	2002	2003	2002	2003
<b>PAST YEAR</b>				
Any Tobacco <sup>1</sup>	310	287	302	318
Cigarettes	290	274	312	326
Smokeless Tobacco	77	77	424	426
Cigars	142	146	399	420
Pipe Tobacco <sup>2</sup>	--	--	--	--
Alcohol	345	339	269	259
Binge Alcohol Use <sup>3</sup>	--	--	--	--
Heavy Alcohol Use <sup>3</sup>	--	--	--	--
<b>PAST MONTH</b>				
Any Tobacco <sup>1</sup>	298	273	315	330
Cigarettes	280	258	326	341
Smokeless Tobacco	68	62	424	425
Cigars	101	107	423	422
Pipe Tobacco	47	59	432	431
Alcohol	288	291	323	320
Binge Alcohol Use <sup>3</sup>	195	195	384	383
Heavy Alcohol Use <sup>3</sup>	110	116	418	414

\*Low precision; no estimate reported.

-- Not available.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.6D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco and Alcohol Use: Standard Errors of Percentages, 2002 and 2003**

Drug	TOBACCO OR ALCOHOL USE			
	Yes		No	
	2002	2003	2002	2003
<b>PAST YEAR</b>				
Any Tobacco <sup>1</sup>	0.39	0.37	0.23	0.24
Cigarettes	0.44	0.42	0.22	0.22
Smokeless Tobacco	0.81	0.82	0.21	0.21
Cigars	0.61	0.64	0.21	0.22
Pipe Tobacco <sup>2</sup>	--	--	--	--
Alcohol	0.23	0.23	0.42	0.39
Binge Alcohol Use <sup>3</sup>	--	--	--	--
Heavy Alcohol Use <sup>3</sup>	--	--	--	--
<b>PAST MONTH</b>				
Any Tobacco <sup>1</sup>	0.44	0.41	0.22	0.23
Cigarettes	0.48	0.45	0.21	0.22
Smokeless Tobacco	0.93	0.87	0.21	0.21
Cigars	0.87	0.91	0.21	0.21
Pipe Tobacco	2.82	4.06	0.21	0.20
Alcohol	0.25	0.25	0.34	0.33
Binge Alcohol Use <sup>3</sup>	0.38	0.38	0.24	0.24
Heavy Alcohol Use <sup>3</sup>	0.72	0.75	0.21	0.21

\*Low precision; no estimate reported.

-- Not available.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.6P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year and Past Month Tobacco and Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	TOBACCO OR ALCOHOL USE			
	Yes		No	
	2002	2003	2002	2003
<b>PAST YEAR</b>				
Any Tobacco <sup>1</sup>	0.0013		0.1033	
Cigarettes	0.0011		0.0779	
Smokeless Tobacco	0.4868		0.0009	
Cigars	0.3231		0.0022	
Pipe Tobacco <sup>2</sup>	--		--	
Alcohol	0.0004		0.5434	
Binge Alcohol Use <sup>3</sup>	--		--	
Heavy Alcohol Use <sup>3</sup>	--		--	
<b>PAST MONTH</b>				
Any Tobacco <sup>1</sup>	0.0204		0.0200	
Cigarettes	0.0215		0.0177	
Smokeless Tobacco	0.3373		0.0008	
Cigars	0.6821		0.0015	
Pipe Tobacco	0.3498		0.0022	
Alcohol	0.0018		0.1660	
Binge Alcohol Use <sup>3</sup>	0.0168		0.0171	
Heavy Alcohol Use <sup>3</sup>	0.3337		0.0028	

\*Low precision; no estimate reported.

-- Not available.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.7C Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>PAST YEAR</b>						
Any Tobacco <sup>2</sup>	1,096	1,042	328	325	1,034	935
Cigarettes	1,003	956	311	313	936	849
Smokeless Tobacco	321	314	78	80	314	302
Cigars	495	498	149	158	470	460
Pipe Tobacco <sup>3</sup>	--	--	--	--	--	--
Alcohol	1,796	1,725	350	371	1,742	1,610
Binge Alcohol Use <sup>4</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>4</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Any Tobacco <sup>2</sup>	1,002	972	321	307	933	876
Cigarettes	936	901	306	294	859	801
Smokeless Tobacco	282	288	69	64	276	280
Cigars	348	353	106	115	331	327
Pipe Tobacco	164	130	48	66	159	115
Alcohol	1,531	1,507	293	316	1,486	1,413
Binge Alcohol Use <sup>4</sup>	838	787	199	214	813	727
Heavy Alcohol Use <sup>4</sup>	400	401	113	124	386	374

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.7D Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Percentages, 2002 and 2003**

Drug	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>PAST YEAR</b>						
Any Tobacco <sup>2</sup>	0.42	0.38	1.30	1.18	0.43	0.39
Cigarettes	0.39	0.38	1.26	1.18	0.40	0.38
Smokeless Tobacco	0.15	0.15	0.43	0.40	0.16	0.15
Cigars	0.23	0.21	0.79	0.76	0.23	0.23
Pipe Tobacco <sup>3</sup>	--	--	--	--	--	--
Alcohol	0.43	0.42	1.29	1.07	0.45	0.45
Binge Alcohol Use <sup>4</sup>	--	--	--	--	--	--
Heavy Alcohol Use <sup>4</sup>	--	--	--	--	--	--
<b>PAST MONTH</b>						
Any Tobacco <sup>2</sup>	0.39	0.37	1.32	1.17	0.40	0.37
Cigarettes	0.38	0.36	1.30	1.16	0.38	0.36
Smokeless Tobacco	0.13	0.13	0.39	0.32	0.14	0.14
Cigars	0.16	0.16	0.59	0.56	0.17	0.16
Pipe Tobacco	0.08	0.06	0.27	0.34	0.08	0.06
Alcohol	0.46	0.43	1.33	1.16	0.47	0.45
Binge Alcohol Use <sup>4</sup>	0.35	0.32	1.03	0.96	0.37	0.33
Heavy Alcohol Use <sup>4</sup>	0.19	0.18	0.63	0.60	0.19	0.18

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.7P Tobacco and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Mental Illness: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>PAST YEAR</b>						
Any Tobacco <sup>2</sup>	0.0973		0.8835		0.0314	
Cigarettes	0.1001		0.8864		0.0258	
Smokeless Tobacco	0.4865		0.1035		0.7785	
Cigars	0.2509		0.4345		0.2733	
Pipe Tobacco <sup>3</sup>	--		--		--	
Alcohol	0.0414		0.7310		0.0253	
Binge Alcohol Use <sup>4</sup>	--		--		--	
Heavy Alcohol Use <sup>4</sup>	--		--		--	
<b>PAST MONTH</b>						
Any Tobacco <sup>2</sup>	0.2451		0.6190		0.1556	
Cigarettes	0.2930		0.6271		0.1733	
Smokeless Tobacco	0.6821		0.1023		0.9926	
Cigars	0.9369		0.5812		0.9827	
Pipe Tobacco	0.2325		0.8710		0.1703	
Alcohol	0.0971		0.8512		0.0725	
Binge Alcohol Use <sup>4</sup>	0.5452		0.9417		0.4443	
Heavy Alcohol Use <sup>4</sup>	0.8533		0.8849		0.8899	

\*Low precision; no estimate reported.

-- Not available.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Any Tobacco use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>3</sup> Information about past year use of pipe tobacco was not collected.

<sup>4</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.8C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Any Illicit Drug Use and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Demographic Characteristic	Total		ANY ILLICIT DRUG USE <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	436	432	186	169	380	402
<b>AGE</b>						
18-25	93	100	61	67	67	71
26-49	290	280	150	140	243	254
50 or Older	289	307	86	71	277	300
<b>GENDER</b>						
Male	252	264	141	109	211	245
Female	328	345	117	133	305	319
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	414	415	177	159	362	379
White	372	388	162	147	322	354
Black or African American	154	143	64	60	142	131
American Indian or Alaska Native	34	21	*	*	31	19
Native Hawaiian or Other Pacific Islander	14	15	*	*	9	10
Asian	112	83	16	*	110	76
Two or More Races	54	48	*	13	50	46
Hispanic or Latino	146	151	64	60	124	145
<b>EDUCATION</b>						
< High School	209	223	84	83	182	201
High School Graduate	250	244	106	98	219	221
Some College	208	212	103	92	177	198
College Graduate	179	186	77	73	165	182
<b>CURRENT EMPLOYMENT</b>						
Full-Time	269	291	130	131	233	262
Part-Time	146	151	70	60	124	143
Unemployed	99	85	65	46	70	73
Other <sup>2</sup>	282	271	92	86	260	251

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.8D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Any Illicit Drug Use and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Demographic Characteristic	Total		ANY ILLICIT DRUG USE <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	0.63	0.57	0.21	0.22
<b>AGE</b>						
18-25	0.30	0.32	0.55	0.61	0.33	0.34
26-49	0.29	0.28	0.94	0.89	0.29	0.30
50 or Older	0.37	0.38	3.31	2.59	0.36	0.38
<b>GENDER</b>						
Male	0.25	0.26	0.82	0.64	0.25	0.29
Female	0.30	0.31	0.94	1.07	0.31	0.33
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.22	0.22	0.68	0.61	0.23	0.24
White	0.25	0.26	0.77	0.69	0.25	0.27
Black or African American	0.67	0.60	1.68	1.71	0.74	0.65
American Indian or Alaska Native	2.60	1.92	*	*	2.87	2.11
Native Hawaiian or Other Pacific Islander	1.97	3.53	*	*	1.46	2.89
Asian	1.39	0.94	2.80	*	1.47	0.92
Two or More Races	2.53	2.22	*	2.98	2.95	2.64
Hispanic or Latino	0.58	0.58	1.80	1.69	0.58	0.65
<b>EDUCATION</b>						
< High School	0.56	0.59	1.49	1.50	0.58	0.62
High School Graduate	0.37	0.36	1.12	1.02	0.37	0.38
Some College	0.40	0.40	1.19	1.07	0.40	0.45
College Graduate	0.34	0.34	1.30	1.27	0.35	0.37
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.23	0.25	0.74	0.76	0.24	0.26
Part-Time	0.53	0.53	1.37	1.13	0.56	0.61
Unemployed	1.31	1.05	2.92	1.97	1.31	1.26
Other <sup>2</sup>	0.48	0.46	1.92	1.86	0.48	0.46

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.8P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Any Illicit Drug Use and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Demographic Characteristic	Total		ANY ILLICIT DRUG USE <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.2332		0.0025	
<b>AGE</b>						
18-25	0.0723		0.0125		0.6744	
26-49	0.0157		0.4723		0.0192	
50 or Older	0.0631		0.5882		0.0394	
<b>GENDER</b>						
Male	0.0361		0.4391		0.0047	
Female	0.0119		0.0175		0.0824	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.0186		0.1705		0.0384	
White	0.0025		0.5033		0.0024	
Black or African American	0.6960		0.2213		0.4227	
American Indian or Alaska Native	0.2710		*		0.1694	
Native Hawaiian or Other Pacific Islander	0.0821		*		0.1396	
Asian	0.4183		*		0.2116	
Two or More Races	0.7539		*		0.9944	
Hispanic or Latino	0.0056		0.7774		0.0013	
<b>EDUCATION</b>						
< High School	0.0451		0.5313		0.0410	
High School Graduate	0.0708		0.2292		0.1763	
Some College	0.1760		0.7148		0.0956	
College Graduate	0.1676		0.4162		0.2227	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.0050		0.1496		0.0124	
Part-Time	0.7102		0.8767		0.7270	
Unemployed	0.5553		0.3389		0.1248	
Other <sup>2</sup>	0.1186		0.4014		0.1327	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.9C Any Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Demographic Characteristic	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	587	574	201	195	540	506
<b>AGE</b>						
18-25	222	212	70	81	196	180
26-49	463	448	168	164	416	389
50 or Older	250	229	88	76	243	212
<b>GENDER</b>						
Male	399	404	149	122	379	367
Female	351	337	130	153	314	293
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	548	527	189	184	509	463
White	458	468	174	168	424	413
Black or African American	230	218	73	75	205	183
American Indian or Alaska Native	40	28	*	*	38	25
Native Hawaiian or Other Pacific Islander	29	17	*	*	27	12
Asian	84	77	15	*	83	72
Two or More Races	71	53	*	*	67	51
Hispanic or Latino	222	195	71	65	202	180
<b>EDUCATION</b>						
< High School	217	229	94	99	193	195
High School Graduate	296	287	114	110	275	256
Some College	287	277	117	105	254	244
College Graduate	286	259	78	77	277	244
<b>CURRENT EMPLOYMENT</b>						
Full-Time	427	424	138	151	400	372
Part-Time	237	189	72	66	224	173
Unemployed	137	114	79	47	110	110
Other <sup>2</sup>	209	214	94	96	202	189

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.9D Any Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Demographic Characteristic	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.27	0.25	1.00	0.95	0.27	0.25
<b>AGE</b>						
18-25	0.46	0.48	1.14	1.24	0.49	0.49
26-49	0.42	0.42	1.45	1.40	0.42	0.41
50 or Older	0.32	0.28	2.18	1.57	0.33	0.27
<b>GENDER</b>						
Male	0.38	0.37	2.05	1.72	0.39	0.36
Female	0.32	0.29	1.11	1.14	0.32	0.29
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.28	0.27	1.04	1.01	0.29	0.26
White	0.30	0.30	1.18	1.12	0.31	0.29
Black or African American	0.82	0.78	3.23	3.40	0.84	0.72
American Indian or Alaska Native	2.58	2.06	*	*	2.84	2.06
Native Hawaiian or Other Pacific Islander	4.26	4.05	*	*	4.21	3.53
Asian	1.03	0.88	2.79	*	1.10	0.88
Two or More Races	2.99	2.38	*	*	3.27	2.60
Hispanic or Latino	0.81	0.67	3.60	2.64	0.78	0.70
<b>EDUCATION</b>						
< High School	0.59	0.59	2.35	2.12	0.58	0.57
High School Graduate	0.41	0.40	1.68	1.59	0.43	0.40
Some College	0.53	0.49	2.01	1.86	0.51	0.49
College Graduate	0.52	0.46	2.31	2.16	0.55	0.46
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.35	0.34	1.41	1.43	0.35	0.33
Part-Time	0.75	0.61	2.33	2.33	0.79	0.63
Unemployed	1.54	1.33	5.02	3.48	1.50	1.47
Other <sup>2</sup>	0.36	0.36	1.73	1.55	0.38	0.35

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.9P Any Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Demographic Characteristic	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.5876		0.2165		0.5681	
<b>AGE</b>						
18-25	0.1830		0.4582		0.0489	
26-49	0.8021		0.4991		0.7984	
50 or Older	0.8715		0.3139		0.6849	
<b>GENDER</b>						
Male	0.5903		0.0126		0.9913	
Female	0.7734		0.5466		0.2958	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.6766		0.6300		0.5441	
White	0.6810		0.2108		0.6471	
Black or African American	0.1215		0.4793		0.0482	
American Indian or Alaska Native	0.9034		*		0.7044	
Native Hawaiian or Other Pacific Islander	0.8629		*		0.8252	
Asian	0.6276		*		0.2878	
Two or More Races	0.7925		*		0.9872	
Hispanic or Latino	0.6485		0.0251		0.9612	
<b>EDUCATION</b>						
< High School	0.6537		0.3091		0.6903	
High School Graduate	0.5760		0.7116		0.8309	
Some College	0.9261		0.2182		0.6702	
College Graduate	0.2802		0.6692		0.2620	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.6395		0.5550		0.5187	
Part-Time	0.9492		0.8901		0.9514	
Unemployed	0.7879		0.0848		0.6551	
Other <sup>2</sup>	0.4424		0.5382		0.3931	

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.10C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Standard Errors of Numbers in Thousands, 2002 and 2003**

Dependence/Abuse	Total		DEPENDENCE <sup>1</sup>				DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No		Yes		No	
	2002	2003	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	458	481	99	99	443	459	119	112	436	458
Marijuana	458	481	49	61	457	471	86	76	448	466
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	458	481	88	78	444	467	102	89	441	467
<b>ALCOHOL</b>	458	481	136	130	438	452	166	173	429	440
<b>ANY ILLICIT DRUG OR ALCOHOL<sup>2</sup></b>	458	481	151	156	429	435	185	189	418	427
<b>BOTH ANY ILLICIT DRUG AND ALCOHOL<sup>2</sup></b>	458	481	72	45	452	475	86	82	447	470

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.10D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: Standard Errors of Percentages, 2002 and 2003**

Dependence/Abuse	Total		DEPENDENCE <sup>1</sup>				DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No		Yes		No	
	2002	2003	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	0.21	0.20	2.05	2.16	0.21	0.20	1.71	1.66	0.20	0.20
Marijuana	0.21	0.20	2.13	2.41	0.21	0.20	2.16	1.93	0.21	0.20
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.21	0.20	3.32	3.15	0.20	0.20	2.63	2.46	0.20	0.20
<b>ALCOHOL</b>	0.21	0.20	1.51	1.57	0.20	0.20	0.90	0.91	0.21	0.20
<b>ANY ILLICIT DRUG OR ALCOHOL<sup>2</sup></b>	0.21	0.20	1.24	1.33	0.20	0.20	0.84	0.83	0.21	0.20
<b>BOTH ANY ILLICIT DRUG AND ALCOHOL<sup>2</sup></b>	0.21	0.20	4.68	3.90	0.21	0.20	2.63	2.50	0.20	0.20

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.10P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Substance Dependence or Abuse: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Dependence/Abuse	Total		DEPENDENCE <sup>1</sup>				DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No		Yes		No	
	2002	2003	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY ILLICIT DRUG<sup>2</sup></b>	0.0017		0.3918		0.0017		0.5608		0.0012	
Marijuana	0.0017		0.2791		0.0023		0.6604		0.0017	
<b>ANY ILLICIT DRUG OTHER THAN MARIJUANA<sup>2</sup></b>	0.0017		0.8062		0.0013		0.6752		0.0012	
<b>ALCOHOL</b>	0.0017		0.5446		0.0009		0.2556		0.0026	
<b>ANY ILLICIT DRUG OR ALCOHOL<sup>2</sup></b>	0.0017		0.2613		0.0012		0.3369		0.0014	
<b>BOTH ANY ILLICIT DRUG AND ALCOHOL<sup>2</sup></b>	0.0017		0.8044		0.0012		0.3207		0.0021	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

<sup>2</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.11C Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Numbers in Thousands, 2002 and 2003**

Dependence/Abuse	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>DEPENDENCE OR ABUSE<sup>2</sup></b>						
Any Illicit Drug <sup>3</sup>	229	205	119	112	190	169
Marijuana	154	138	86	76	125	114
Any Illicit Drug Other Than Marijuana <sup>3</sup>	182	161	102	89	147	131
Alcohol	436	397	166	173	395	344
Any Illicit Drug or Alcohol <sup>3</sup>	471	428	185	189	422	368
Both Any Illicit Drug and Alcohol <sup>3</sup>	143	134	86	82	112	103
<b>DEPENDENCE<sup>2</sup></b>						
Any Illicit Drug <sup>3</sup>	179	159	99	99	143	128
Marijuana	104	101	49	61	90	80
Any Illicit Drug Other Than Marijuana <sup>3</sup>	140	131	88	78	108	103
Alcohol	288	269	136	130	247	230
Any Illicit Drug or Alcohol <sup>3</sup>	330	303	151	156	282	254
Both Any Illicit Drug and Alcohol <sup>3</sup>	104	74	72	45	75	58

\*Low precision; no estimate reported.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>3</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.11D Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: Standard Errors of Percentages, 2002 and 2003**

Dependence/Abuse	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>DEPENDENCE OR ABUSE<sup>2</sup></b>						
Any Illicit Drug <sup>3</sup>	0.11	0.09	0.65	0.55	0.10	0.09
Marijuana	0.07	0.06	0.48	0.37	0.06	0.06
Any Illicit Drug Other Than Marijuana <sup>3</sup>	0.09	0.08	0.56	0.44	0.08	0.07
Alcohol	0.19	0.18	0.90	0.81	0.19	0.17
Any Illicit Drug or Alcohol <sup>3</sup>	0.21	0.19	0.98	0.86	0.20	0.18
Both Any Illicit Drug and Alcohol <sup>3</sup>	0.07	0.06	0.48	0.41	0.06	0.05
<b>DEPENDENCE<sup>2</sup></b>						
Any Illicit Drug <sup>3</sup>	0.08	0.07	0.55	0.48	0.07	0.07
Marijuana	0.05	0.05	0.29	0.31	0.05	0.04
Any Illicit Drug Other Than Marijuana <sup>3</sup>	0.07	0.06	0.49	0.38	0.06	0.05
Alcohol	0.13	0.12	0.75	0.62	0.12	0.12
Any Illicit Drug or Alcohol <sup>3</sup>	0.15	0.14	0.82	0.71	0.14	0.13
Both Any Illicit Drug and Alcohol <sup>3</sup>	0.05	0.03	0.41	0.23	0.04	0.03

\*Low precision; no estimate reported.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>3</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.11P Substance Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Dependence/Abuse	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>DEPENDENCE OR ABUSE<sup>2</sup></b>						
Any Illicit Drug <sup>3</sup>	0.3827		0.2740		0.3804	
Marijuana	0.8767		0.5992		0.8053	
Any Illicit Drug Other Than Marijuana <sup>3</sup>	0.3545		0.2932		0.3948	
Alcohol	0.3540		0.4024		0.2828	
Any Illicit Drug or Alcohol <sup>3</sup>	0.2436		0.1612		0.2379	
Both Any Illicit Drug and Alcohol <sup>3</sup>	0.5839		0.8506		0.3295	
<b>DEPENDENCE<sup>2</sup></b>						
Any Illicit Drug <sup>3</sup>	0.3491		0.4169		0.2669	
Marijuana	0.9446		0.9287		0.5889	
Any Illicit Drug Other Than Marijuana <sup>3</sup>	0.5166		0.3452		0.5808	
Alcohol	0.0578		0.1121		0.0796	
Any Illicit Drug or Alcohol <sup>3</sup>	0.0992		0.1592		0.0804	
Both Any Illicit Drug and Alcohol <sup>3</sup>	0.0558		0.1737		0.1097	

\*Low precision; no estimate reported.

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>3</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana includes cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.12C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Any Illicit Drug Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003**

Gender/Age	Total		ANY ILLICIT DRUG DEPENDENCE OR ABUSE <sup>1,2</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	458	481	119	112	436	458
18-25	106	117	38	45	97	104
26-49	313	308	94	95	295	294
50 or Older	298	320	*	*	291	317
<b>MALE</b>	258	282	102	72	240	273
18-25	62	68	26	29	57	59
26-49	197	184	76	65	182	179
50 or Older	165	202	*	*	153	202
<b>FEMALE</b>	352	374	60	79	347	357
18-25	79	85	27	30	72	80
26-49	230	246	54	67	226	236
50 or Older	245	247	*	*	245	245

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.12D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Any Illicit Drug Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2002 and 2003**

Gender/Age	Total		ANY ILLICIT DRUG DEPENDENCE OR ABUSE <sup>1,2</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	1.71	1.66	0.20	0.20
18-25	0.30	0.32	1.27	1.46	0.30	0.31
26-49	0.29	0.28	2.77	2.76	0.28	0.28
50 or Older	0.37	0.38	*	*	0.36	0.38
<b>MALE</b>	0.25	0.26	2.22	1.82	0.24	0.26
18-25	0.37	0.40	1.46	1.60	0.39	0.39
26-49	0.38	0.36	3.39	3.15	0.36	0.36
50 or Older	0.44	0.52	*	*	0.42	0.53
<b>FEMALE</b>	0.30	0.31	2.80	2.78	0.30	0.30
18-25	0.45	0.45	2.27	2.40	0.44	0.45
26-49	0.42	0.44	4.89	4.68	0.43	0.43
50 or Older	0.55	0.54	*	*	0.55	0.54

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.12P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Any Illicit Drug Dependence or Abuse, Gender, and Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Gender/Age	Total		ANY ILLICIT DRUG DEPENDENCE OR ABUSE <sup>1,2</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.5608		0.0012	
18-25	0.0723		0.1883		0.1140	
26-49	0.0157		0.9244		0.0149	
50 or Older	0.0631		*		0.0464	
<b>MALE</b>	0.0361		0.4594		0.0109	
18-25	0.2809		0.3905		0.3251	
26-49	0.6245		0.5470		0.4312	
50 or Older	0.0240		*		0.0083	
<b>FEMALE</b>	0.0119		0.0930		0.0241	
18-25	0.1278		0.3759		0.1741	
26-49	0.0075		0.5755		0.0156	
50 or Older	0.5071		*		0.5672	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

<sup>2</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.13C Any Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003**

Gender/Age	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	229	205	119	112	190	169
18-25	82	79	38	45	70	64
26-49	171	173	94	95	139	142
50 or Older	106	61	61	28	86	55
<b>MALE</b>	183	166	102	72	147	147
18-25	65	64	26	29	59	55
26-49	142	141	76	65	116	122
50 or Older	89	55	*	*	65	54
<b>FEMALE</b>	126	116	60	79	110	83
18-25	45	48	27	30	34	36
26-49	91	101	54	67	73	74
50 or Older	60	26	11	24	59	11

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.13D Any Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2002 and 2003**

Gender/Age	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.11	0.09	0.65	0.55	0.10	0.09
18-25	0.24	0.23	0.84	0.89	0.24	0.22
26-49	0.17	0.17	0.93	0.87	0.15	0.16
50 or Older	0.13	0.08	1.55	0.57	0.12	0.07
<b>MALE</b>	0.18	0.16	1.57	1.04	0.15	0.15
18-25	0.40	0.37	1.56	1.48	0.41	0.37
26-49	0.28	0.28	2.12	1.82	0.25	0.27
50 or Older	0.25	0.15	*	*	0.19	0.15
<b>FEMALE</b>	0.11	0.10	0.53	0.59	0.11	0.08
18-25	0.27	0.29	0.98	1.04	0.25	0.26
26-49	0.18	0.20	0.86	0.92	0.16	0.17
50 or Older	0.14	0.06	0.40	0.80	0.15	0.03

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.13P Any Illicit Drug Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Gender/Age	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.3827		0.2740		0.3804	
18-25	0.2286		0.8195		0.1243	
26-49	0.7282		0.7297		0.7445	
50 or Older	0.0821		0.2975		0.1386	
<b>MALE</b>	0.2043		0.0435		0.5341	
18-25	0.1680		0.6470		0.1462	
26-49	0.7207		0.3700		0.9860	
50 or Older	0.2897		*		0.7710	
<b>FEMALE</b>	0.8002		0.4246		0.4789	
18-25	0.9425		0.9148		0.6226	
26-49	0.2631		0.5239		0.6007	
50 or Older	0.1192		0.3683		0.0355	

\*Low precision; no estimate reported.

NOTE: Any Illicit Drug includes marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or any prescription-type psychotherapeutic used nonmedically.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.14C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003**

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	458	481	166	173	429	440
18-25	106	117	53	58	88	96
26-49	313	308	133	132	302	286
50 or Older	298	320	91	95	283	305
<b>MALE</b>	258	282	131	126	224	253
18-25	62	68	34	39	49	56
26-49	197	184	97	90	177	161
50 or Older	165	202	85	80	139	186
<b>FEMALE</b>	352	374	95	121	346	344
18-25	79	85	35	40	68	72
26-49	230	246	83	98	226	225
50 or Older	245	247	*	*	243	241

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.14D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2002 and 2003**

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	0.90	0.91	0.21	0.20
18-25	0.30	0.32	0.80	0.94	0.32	0.32
26-49	0.29	0.28	1.35	1.31	0.30	0.29
50 or Older	0.37	0.38	3.56	3.75	0.36	0.38
<b>MALE</b>	0.25	0.26	1.07	1.02	0.24	0.26
18-25	0.37	0.40	0.85	1.03	0.40	0.42
26-49	0.38	0.36	1.50	1.33	0.38	0.36
50 or Older	0.44	0.52	4.11	4.31	0.40	0.51
<b>FEMALE</b>	0.30	0.31	1.60	1.84	0.31	0.30
18-25	0.45	0.45	1.53	1.60	0.46	0.46
26-49	0.42	0.44	2.53	2.92	0.44	0.43
50 or Older	0.55	0.54	*	*	0.55	0.54

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.14P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Past Year Alcohol Dependence or Abuse, Gender, and Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Gender/Age	Total		ALCOHOL DEPENDENCE OR ABUSE <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.2556		0.0026	
18-25	0.0723		0.3139		0.1099	
26-49	0.0157		0.4299		0.0261	
50 or Older	0.0631		0.7126		0.0630	
<b>MALE</b>	0.0361		0.5496		0.0260	
18-25	0.2809		0.1470		0.5970	
26-49	0.6245		0.8553		0.6011	
50 or Older	0.0240		0.9203		0.0090	
<b>FEMALE</b>	0.0119		0.4273		0.0206	
18-25	0.1278		0.8304		0.0723	
26-49	0.0075		0.3335		0.0155	
50 or Older	0.5071		*		0.5851	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.15C Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003**

Gender/Age	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	436	397	166	173	395	344
18-25	145	132	53	58	123	116
26-49	309	297	133	132	278	254
50 or Older	235	209	91	95	216	188
<b>MALE</b>	361	328	131	126	341	294
18-25	108	99	34	39	97	91
26-49	261	251	97	90	249	222
50 or Older	208	178	*	80	189	161
<b>FEMALE</b>	221	205	95	121	188	162
18-25	74	77	35	40	62	61
26-49	165	152	83	98	135	120
50 or Older	104	98	32	51	99	83

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.15D Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: Standard Errors of Percentages, 2002 and 2003**

Gender/Age	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.19	0.18	0.90	0.81	0.19	0.17
18-25	0.36	0.34	1.07	1.08	0.37	0.36
26-49	0.30	0.28	1.35	1.17	0.29	0.27
50 or Older	0.29	0.25	2.23	1.92	0.28	0.24
<b>MALE</b>	0.33	0.31	1.88	1.66	0.33	0.30
18-25	0.57	0.55	1.74	1.90	0.59	0.56
26-49	0.50	0.47	2.62	2.30	0.50	0.46
50 or Older	0.56	0.46	*	4.20	0.53	0.45
<b>FEMALE</b>	0.20	0.18	0.84	0.88	0.19	0.17
18-25	0.40	0.41	1.20	1.25	0.42	0.41
26-49	0.32	0.30	1.32	1.28	0.29	0.27
50 or Older	0.24	0.22	1.16	1.68	0.24	0.20

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.15P Alcohol Dependence or Abuse in the Past Year among Persons Aged 18 or Older, by Past Year Serious Mental Illness, Gender, and Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Gender/Age	Total		SERIOUS MENTAL ILLNESS <sup>1</sup>			
			Yes		No	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.3540		0.4024		0.2828	
18-25	0.3726		0.6637		0.3122	
26-49	0.5867		0.6828		0.5057	
50 or Older	0.7062		0.7581		0.6621	
<b>MALE</b>	0.1678		0.2343		0.1789	
18-25	0.1610		0.8556		0.0842	
26-49	0.5533		0.7414		0.5587	
50 or Older	0.3697		*		0.4328	
<b>FEMALE</b>	0.6901		0.9447		0.9358	
18-25	0.7655		0.4406		0.5658	
26-49	0.9230		0.9343		0.6414	
50 or Older	0.3997		0.4728		0.5949	

\*Low precision; no estimate reported.

NOTE: Dependence or abuse is based on definitions found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

<sup>1</sup> Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.16C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Detailed Age Categories: Standard Errors of Numbers in Thousands, 2002 and 2003**

Age Category	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	458	481	294	336	342	316
18	34	42	18	20	31	36
19	34	41	19	22	28	36
20	35	38	16	20	31	31
21	36	36	20	20	28	30
22	33	33	20	19	27	26
23	34	34	23	21	25	28
24	30	35	19	24	23	27
25	31	34	22	22	23	23
26-29	108	130	68	81	84	101
30-34	134	123	99	90	95	89
35-39	162	138	107	97	107	101
40-44	152	146	114	102	99	105
45-49	127	154	103	105	69	109
50-54	173	186	142	149	100	109
55-59	144	132	109	95	97	92
60-64	83	104	36	72	75	75
65 or Older	182	200	72	136	165	144

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year serious mental illness, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.16D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Detailed Age Categories: Standard Errors of Percentages, 2002 and 2003**

Age Category	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.21	0.20	0.93	0.96	0.18	0.17
18	0.72	0.80	3.31	2.91	0.72	0.81
19	0.77	0.95	3.42	3.69	0.71	0.93
20	0.83	0.86	3.49	3.62	0.83	0.80
21	0.80	0.85	3.27	3.45	0.73	0.81
22	0.77	0.78	3.77	3.60	0.74	0.71
23	0.82	0.77	3.71	3.24	0.69	0.74
24	0.78	0.84	3.53	3.70	0.72	0.80
25	0.88	0.84	3.52	3.66	0.76	0.69
26-29	0.68	0.81	3.45	3.82	0.61	0.72
30-34	0.65	0.58	2.74	2.54	0.55	0.50
35-39	0.69	0.60	2.76	2.50	0.55	0.54
40-44	0.64	0.61	2.40	2.47	0.53	0.53
45-49	0.58	0.68	2.50	2.39	0.38	0.60
50-54	0.88	0.90	4.06	3.75	0.62	0.65
55-59	0.94	0.85	3.89	3.65	0.78	0.70
60-64	0.72	0.86	2.96	4.29	0.72	0.73
65 or Older	0.53	0.57	2.30	4.09	0.53	0.45

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year serious mental illness, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.16P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Detailed Age Categories: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Age Category	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL 18 OR OLDER</b>	0.0017		0.0664		0.0120	
18	0.1658		0.4027		0.3140	
19	0.0056		0.1662		0.0165	
20	0.1985		0.1983		0.5087	
21	0.8359		0.5148		0.9497	
22	0.2931		0.9069		0.1194	
23	0.5680		0.6775		0.5262	
24	0.0083		0.0209		0.1304	
25	0.3353		0.0255		0.8312	
26-29	0.0476		0.2036		0.0666	
30-34	0.2562		0.4442		0.3411	
35-39	0.0622		0.6916		0.0528	
40-44	0.7362		0.6414		0.3779	
45-49	0.0097		0.7085		0.0004	
50-54	0.3556		0.4021		0.8069	
55-59	0.6958		0.9086		0.3736	
60-64	0.4444		0.0573		0.4969	
65 or Older	0.1817		0.0058		0.9737	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year serious mental illness, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 6.17C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Demographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Demographic Characteristic	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	458	481	294	336	342	316
<b>AGE</b>						
18-25	106	117	60	63	84	90
26-49	313	308	220	215	209	222
50 or Older	298	320	192	236	232	215
<b>GENDER</b>						
Male	258	282	181	188	191	213
Female	352	374	226	272	267	235
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	435	452	278	319	323	289
White	387	426	248	307	289	263
Black or African American	172	154	111	97	119	112
American Indian or Alaska Native	33	19	*	*	19	15
Native Hawaiian or Other Pacific Islander	14	15	*	*	14	15
Asian	119	85	*	*	81	77
Two or More Races	56	51	*	*	43	26
Hispanic or Latino	149	165	87	90	116	136
<b>EDUCATION</b>						
< High School	229	243	129	136	192	195
High School Graduate	258	257	176	187	191	175
Some College	216	225	160	159	141	145
College Graduate	187	197	130	154	123	138
<b>CURRENT EMPLOYMENT</b>						
Full-Time	284	314	190	208	198	235
Part-Time	150	160	119	124	93	101
Unemployed	107	88	76	55	77	66
Other <sup>3</sup>	288	287	187	205	220	191

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year SMI, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.17D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Demographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Demographic Characteristic	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	0.93	0.96	0.18	0.17
<b>AGE</b>						
18-25	0.30	0.32	1.38	1.26	0.27	0.29
26-49	0.29	0.28	1.18	1.13	0.24	0.26
50 or Older	0.37	0.38	1.83	1.99	0.33	0.30
<b>GENDER</b>						
Male	0.25	0.26	1.75	1.74	0.21	0.22
Female	0.30	0.31	1.09	1.15	0.28	0.25
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.22	0.22	0.95	0.99	0.20	0.18
White	0.25	0.26	1.02	1.06	0.22	0.20
Black or African American	0.67	0.60	4.36	4.16	0.54	0.50
American Indian or Alaska Native	2.60	1.92	*	*	1.89	1.63
Native Hawaiian or Other Pacific Islander	1.97	3.53	*	*	2.03	3.75
Asian	1.39	0.94	*	*	1.05	0.90
Two or More Races	2.53	2.22	*	*	2.37	1.47
Hispanic or Latino	0.58	0.58	3.53	3.77	0.50	0.56
<b>EDUCATION</b>						
< High School	0.56	0.59	2.54	2.75	0.55	0.55
High School Graduate	0.37	0.36	1.75	1.85	0.31	0.29
Some College	0.40	0.40	1.76	1.81	0.31	0.32
College Graduate	0.34	0.34	1.57	1.56	0.27	0.28
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.23	0.25	1.23	1.21	0.19	0.22
Part-Time	0.53	0.53	2.54	2.40	0.40	0.41
Unemployed	1.31	1.05	5.26	4.16	1.12	0.93
Other <sup>3</sup>	0.48	0.46	1.77	1.85	0.44	0.37

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year SMI, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.17P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Demographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Demographic Characteristic	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.0664		0.0120	
<b>AGE</b>						
18-25	0.0723		0.4828		0.2194	
26-49	0.0157		0.8997		0.0015	
50 or Older	0.0631		0.0111		0.8139	
<b>GENDER</b>						
Male	0.0361		0.1782		0.0545	
Female	0.0119		0.2189		0.0595	
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	0.0186		0.0691		0.1512	
White	0.0025		0.0371		0.0631	
Black or African American	0.6960		0.7151		0.7304	
American Indian or Alaska Native	0.2710		*		0.5139	
Native Hawaiian or Other Pacific Islander	0.0821		*		0.0865	
Asian	0.4183		*		0.9094	
Two or More Races	0.7539		*		0.3116	
Hispanic or Latino	0.0056		0.7841		0.0011	
<b>EDUCATION</b>						
< High School	0.0451		0.0141		0.1732	
High School Graduate	0.0708		0.8561		0.0827	
Some College	0.1760		0.7588		0.2021	
College Graduate	0.1676		0.1953		0.4040	
<b>CURRENT EMPLOYMENT</b>						
Full-Time	0.0050		0.8429		0.0026	
Part-Time	0.7102		0.6526		0.3821	
Unemployed	0.5553		0.9917		0.3159	
Other <sup>3</sup>	0.1186		0.0027		0.9838	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year SMI, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>3</sup> Retired person, disabled person, homemaker, student, or other person not in the labor force.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.18C Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2002 and 2003**

Geographic Characteristic	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	458	481	294	336	342	316
<b>GEOGRAPHIC DIVISION</b>						
Northeast	223	210	155	146	142	130
New England	125	87	107	63	53	56
Middle Atlantic	185	191	112	131	132	117
Midwest	194	170	119	135	153	100
East North Central	158	150	97	114	129	95
West North Central	113	80	70	73	82	33
South	279	296	185	200	207	202
South Atlantic	232	221	150	152	175	144
East South Central	105	119	75	85	68	82
West South Central	115	157	79	99	86	116
West	211	265	119	182	176	179
Mountain	79	113	53	94	60	72
Pacific	195	240	107	155	165	164
<b>COUNTY TYPE<sup>3</sup></b>						
Large Metro	337	337	210	234	249	245
Small Metro	259	286	173	198	164	178
250K - 1 Mil. Pop.	208	235	126	154	142	141
< 250K Pop.	157	146	118	114	85	97
Nonmetro	193	197	119	140	158	121
Urbanized	119	155	92	114	67	85
Less Urbanized	145	130	71	91	131	88
Completely Rural	71	52	*	*	62	24

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year serious mental illness, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.18D Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Geographic Characteristics: Standard Errors of Percentages, 2002 and 2003**

Geographic Characteristic	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.21	0.20	0.93	0.96	0.18	0.17
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.51	0.47	2.43	2.13	0.40	0.36
New England	0.98	0.81	5.08	3.14	0.56	0.63
Middle Atlantic	0.59	0.58	2.59	2.78	0.50	0.43
Midwest	0.40	0.35	1.61	1.65	0.37	0.25
East North Central	0.50	0.43	2.01	1.94	0.46	0.34
West North Central	0.68	0.60	2.69	3.18	0.59	0.31
South	0.36	0.36	1.75	1.63	0.30	0.30
South Atlantic	0.57	0.50	2.78	2.28	0.47	0.40
East South Central	0.84	0.82	3.08	4.08	0.65	0.69
West South Central	0.49	0.65	2.69	2.82	0.42	0.57
West	0.41	0.46	1.51	2.44	0.43	0.39
Mountain	0.58	0.74	2.39	3.58	0.54	0.58
Pacific	0.53	0.58	1.88	3.17	0.56	0.49
<b>COUNTY TYPE<sup>3</sup></b>						
Large Metro	0.28	0.27	1.28	1.37	0.25	0.23
Small Metro	0.38	0.36	1.73	1.54	0.29	0.29
250K - 1 Mil. Pop.	0.45	0.45	1.98	1.87	0.37	0.35
< 250K Pop.	0.62	0.55	3.23	2.55	0.40	0.48
Nonmetro	0.51	0.51	2.16	2.47	0.49	0.38
Urbanized	0.71	0.84	3.24	4.05	0.51	0.55
Less Urbanized	0.77	0.70	3.11	3.51	0.80	0.59
Completely Rural	1.54	1.28	*	*	1.56	0.69

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year serious mental illness, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 6.18P Serious Mental Illness in the Past Year among Persons Aged 18 or Older, by Receipt of Past Year Mental Health Treatment/Counseling and Geographic Characteristics: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Geographic Characteristic	Total <sup>1</sup>		MENTAL HEALTH TREATMENT/COUNSELING <sup>2</sup>			
			Received		Not Received	
	2002	2003	2002	2003	2002	2003
<b>TOTAL</b>	0.0017		0.0664		0.0120	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.8462		0.6792		0.8978	
New England	0.6059		0.3150		0.5496	
Middle Atlantic	0.9735		0.8283		0.9052	
Midwest	0.3933		0.4496		0.8187	
East North Central	0.6564		0.5371		0.5560	
West North Central	0.3681		0.0392		0.0787	
South	0.0166		0.2811		0.0753	
South Atlantic	0.3875		0.6857		0.8303	
East South Central	0.5375		0.7258		0.1369	
West South Central	0.0036		0.0408		0.0363	
West	0.0038		0.0299		0.0074	
Mountain	0.0082		0.0588		0.0848	
Pacific	0.0501		0.1994		0.0252	
<b>COUNTY TYPE<sup>3</sup></b>						
Large Metro	0.0123		0.1799		0.0256	
Small Metro	0.0692		0.3671		0.0682	
250K - 1 Mil. Pop.	0.0631		0.0420		0.2729	
< 250K Pop.	0.5859		0.3471		0.0835	
Nonmetro	0.3858		0.4451		0.9191	
Urbanized	0.2456		0.4417		0.2027	
Less Urbanized	0.5679		0.8457		0.9052	
Completely Rural	0.4827		*		0.0499	

\*Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)* and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>1</sup> Estimates in the Total column represent persons aged 18 or older with past year serious mental illness, including those with unknown mental health treatment/counseling information.

<sup>2</sup> Mental Health Treatment/Counseling is defined as having received inpatient care, outpatient care, or using prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with missing treatment/counseling information are excluded.

<sup>3</sup> Estimates for 2002 and 2003 are based on a revised definition of county type and are not comparable with estimates by county type published in prior NSDUH reports.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.