Table 6.33C Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

			AGE GROUP (Years)					
	Total		18-25		26-49		50 or	Older
Location of Treatment/Counseling <sup>1</sup>	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	184	178	34	33	138	132	113	118
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	363	341	50	52	239	267	247	190
Doctor's Office - Not Part of a Clinic	225	209	29	29	140	137	178	150
Outpatient Medical Clinic	116	129	18	19	59	79	97	84
Partial Day Hospital or Day Treatment Program	54	58	14	11	49	50	12	27
School or University Setting/Clinic/Center <sup>2</sup>	30	21	13	10	27	18	*	*
Church/Religious Setting <sup>2</sup>	23	38	7	9	22	34	*	13
Some Other Place	65	65	9	14	42	46	49	45

<sup>\*</sup>Low precision; no estimate reported.

<sup>&</sup>lt;sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Table 6.33D Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2002 and 2003

			AGE GROUP (Years)					
	Total		18-25		26-49		50 or	Older
Location of Treatment/Counseling <sup>1</sup>	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	1.04	1.08	1.40	1.32	1.32	1.38	2.34	2.45
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	1.52	1.50	1.62	1.62	1.71	1.82	3.82	3.27
Doctor's Office - Not Part of a Clinic	1.28	1.16	1.21	1.24	1.41	1.30	3.27	2.89
Outpatient Medical Clinic	0.71	0.81	0.81	0.83	0.64	0.89	2.06	1.86
Partial Day Hospital or Day Treatment Program	0.35	0.38	0.65	0.51	0.54	0.56	0.27	0.63
School or University Setting/Clinic/Center <sup>2</sup>	0.19	0.14	0.62	0.45	0.29	0.21	*	*
Church/Religious Setting <sup>2</sup>	0.15	0.25	0.34	0.41	0.24	0.39	*	0.30
Some Other Place	0.41	0.43	0.43	0.65	0.45	0.52	1.08	1.04

<sup>\*</sup>Low precision; no estimate reported.

<sup>&</sup>lt;sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Table 6.33P Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

			AGE GROUP (Years)					
	Total		18-25		26-49		50 or	Older
Location of Treatment/Counseling <sup>1</sup>	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	0.7657		0.5962		0.2613		0.5113	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.9426		0.8692		0.2813		0.1994	
Doctor's Office - Not Part of a Clinic	0.9170		0.8196		0.9818		0.8969	
Outpatient Medical Clinic	0.6667		0.6710		0.1241		0.5822	
Partial Day Hospital or Day Treatment Program	0.7126		0.1839		0.9742		0.0760	
School or University Setting/Clinic/Center <sup>2</sup>	0.4648		0.1855		0.8416		*	
Church/Religious Setting <sup>2</sup>	0.1804		0.8715		0.3398		*	
Some Other Place	0.7911		0.2236		0.8227		0.7873	

<sup>\*</sup>Low precision; no estimate reported.

<sup>&</sup>lt;sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Table 6.34C Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2002 and 2003

			AGE GROUP (Years)					
	Total		18-25		26-49		50 or Older	
Location of Treatment/Counseling <sup>1</sup>	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	105	129	25	27	86	100	*	71
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	195	199	36	36	140	142	*	*
Doctor's Office - Not Part of a Clinic	126	146	20	21	100	94	*	*
Outpatient Medical Clinic	79	75	11	14	36	53	*	*
Partial Day Hospital or Day Treatment Program	25	42	6	8	21	36	*	20
School or University Setting/Clinic/Center <sup>2</sup>	12	19	7	5	9	18	*	*
Church/Religious Setting <sup>2</sup>	13	19	4	7	12	18	*	*
Some Other Place	33	48	5	10	27	34	*	*

<sup>\*</sup>Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic* and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the Results from the 2003 National Survey on Drug Use and Health: National Findings.

<sup>&</sup>lt;sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Table 6.34D Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2002 and 2003

			AGE GROUP (Years)					
	Total		18-25		26-49		50 or	Older
Location of Treatment/Counseling <sup>1</sup>	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	1.67	1.83	2.12	2.08	2.09	2.35	*	4.49
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	2.35	2.30	2.23	2.29	2.72	2.73	*	*
Doctor's Office - Not Part of a Clinic	1.90	2.04	1.87	1.77	2.37	2.32	*	*
Outpatient Medical Clinic	1.31	1.20	1.07	1.23	1.00	1.47	*	*
Partial Day Hospital or Day Treatment Program	0.43	0.69	0.58	0.78	0.60	0.99	*	1.41
School or University Setting/Clinic/Center <sup>2</sup>	0.21	0.31	0.72	0.51	0.26	0.50	*	*
Church/Religious Setting <sup>2</sup>	0.23	0.31	0.36	0.64	0.35	0.49	*	*
Some Other Place	0.57	0.78	0.48	0.93	0.75	0.92	*	*

<sup>\*</sup>Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic* and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the *Results from the 2003 National Survey on Drug Use and Health: National Findings*.

<sup>&</sup>lt;sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

Table 6.34P Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Mental Illness Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: P-Values from Tests of Differences of Percentages, 2003 Versus 2002

			AGE GROUP (Years)					
	Total		18-25		26-49		50 or	Older
Location of Treatment/Counseling <sup>1</sup>	2002	2003	2002	2003	2002	2003	2002	2003
Outpatient Mental Health Clinic or Center	0.8364		0.7461		0.7629		*	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.6634		0.2427		0.8299		*	
Doctor's Office - Not Part of a Clinic	0.9922		0.9738		0.6385		*	
Outpatient Medical Clinic	0.8452		0.2504		0.2538		*	
Partial Day Hospital or Day Treatment Program	0.1835		0.5155		0.3349		*	
School or University Setting/Clinic/Center <sup>2</sup>	0.9257		0.1610		0.4498		*	
Church/Religious Setting <sup>2</sup>	0.1726		0.3137		0.2476		*	
Some Other Place	0.4804		0.1902		0.7077		*	

<sup>\*</sup>Low precision; no estimate reported.

NOTE: Serious Mental Illness (SMI) is defined as having a diagnosable mental, behavioral, or emotional disorder that met the criteria found in the 4<sup>th</sup> edition of the *Diagnostic* and Statistical Manual of Mental Disorders (DSM-IV) and resulted in functional impairment that substantially interfered with or limited one or more major life activities. See Section B.4 of Appendix B of the Results from the 2003 National Survey on Drug Use and Health: National Findings.

<sup>&</sup>lt;sup>1</sup> Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>2</sup> Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.