

**Table 7.25C Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	196	187	345	321	386	390	559	510
Cigarettes	207	187	321	319	377	384	523	469
Smokeless Tobacco	103	91	124	133	111	131	187	193
Cigars	149	142	192	168	192	176	160	157
Pipe Tobacco	44	55	65	47	94	59	114	90

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.25D Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	1.24	1.16	0.91	0.85	0.58	0.60	0.48	0.43
Cigarettes	1.30	1.16	0.85	0.85	0.57	0.59	0.45	0.40
Smokeless Tobacco	0.65	0.56	0.33	0.35	0.17	0.20	0.16	0.16
Cigars	0.94	0.88	0.51	0.45	0.29	0.27	0.14	0.13
Pipe Tobacco	0.28	0.34	0.17	0.12	0.14	0.09	0.10	0.08

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.25P Tobacco Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	0.9496		0.8157		0.1244		0.5106	
Cigarettes	0.8387		0.7068		0.1932		0.6100	
Smokeless Tobacco	0.4205		0.6743		0.2217		0.2793	
Cigars	0.5773		0.5748		0.7623		0.7399	
Pipe Tobacco	0.5475		0.2545		0.1096		0.9027	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.26C Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	13	13	31	31	27	25	53	50
Cigarettes	15	14	30	29	25	23	48	44
Smokeless Tobacco	11	12	15	16	8	8	18	19
Cigars	15	15	25	23	17	15	26	27
Pipe Tobacco	7	6	8	13	5	4	8	11

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.26D Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	2.02	1.93	1.53	1.54	1.56	1.42	0.26	0.24
Cigarettes	2.37	2.13	1.51	1.47	1.43	1.32	0.23	0.21
Smokeless Tobacco	1.79	1.84	0.75	0.80	0.46	0.44	0.09	0.09
Cigars	2.40	2.27	1.24	1.13	0.97	0.86	0.13	0.13
Pipe Tobacco	1.08	0.98	0.37	0.63	0.31	0.20	0.04	0.05

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.26P Tobacco Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	0.8201		0.8363		0.0885		0.0954	
Cigarettes	0.7166		0.6697		0.2589		0.0204	
Smokeless Tobacco	0.5742		0.9576		0.9108		0.7691	
Cigars	0.7073		0.5105		0.9796		0.7106	
Pipe Tobacco	0.7222		0.1206		0.3140		0.8974	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.27C Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	44	45	73	67	59	58	77	79
Cigarettes	49	50	73	72	59	56	75	77
Smokeless Tobacco	34	35	32	35	16	19	24	20
Cigars	43	43	49	54	31	31	36	37
Pipe Tobacco	12	13	16	14	9	9	11	12

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.27D Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	0.95	0.93	0.91	0.79	0.97	0.92	0.63	0.64
Cigarettes	1.06	1.05	0.90	0.86	0.97	0.90	0.61	0.63
Smokeless Tobacco	0.74	0.74	0.39	0.41	0.26	0.30	0.19	0.16
Cigars	0.93	0.90	0.61	0.64	0.52	0.49	0.30	0.30
Pipe Tobacco	0.26	0.28	0.20	0.17	0.15	0.15	0.09	0.10

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.



**Table 7.27P Tobacco Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	0.6906		0.6985		0.1199		0.5511	
Cigarettes	0.7158		0.4099		0.1550		0.8106	
Smokeless Tobacco	0.3676		0.8110		0.3940		0.0744	
Cigars	0.5937		0.1207		0.7776		0.9505	
Pipe Tobacco	0.7679		0.0220		0.6274		0.9825	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.28C Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	184	182	329	310	374	381	552	491
Cigarettes	194	180	306	306	363	371	516	452
Smokeless Tobacco	98	82	119	122	110	131	186	194
Cigars	142	132	179	163	188	176	153	152
Pipe Tobacco	42	53	62	43	93	58	114	88

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.28D Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2002 and 2003**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	1.74	1.70	1.18	1.14	0.64	0.67	0.67	0.57
Cigarettes	1.83	1.68	1.10	1.12	0.62	0.65	0.62	0.53
Smokeless Tobacco	0.93	0.76	0.43	0.45	0.19	0.23	0.22	0.23
Cigars	1.33	1.23	0.64	0.60	0.32	0.31	0.19	0.18
Pipe Tobacco	0.40	0.49	0.22	0.16	0.16	0.10	0.14	0.10

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.

**Table 7.28P Tobacco Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: P-Values from Tests of Differences of Percentages, 2003 Versus 2002**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2002	2003	2002	2003	2002	2003	2002	2003
<b>ANY TOBACCO<sup>2</sup></b>	0.9974		0.8210		0.1967		0.6220	
Cigarettes	0.8120		0.7729		0.2603		0.7422	
Smokeless Tobacco	0.5080		0.7097		0.2753		0.3380	
Cigars	0.6647		0.8940		0.6796		0.7621	
Pipe Tobacco	0.6209		0.3476		0.1288		0.9215	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Any Tobacco product includes cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002 and 2003.