Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2001: United States, 1999-2001

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates per 100,000 U.S. standard population; see Technical Notes. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years; see Technical Notes. For explanation of asterisks preceding cause-of-death codes, see Technical Notes]

Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	Age												- Age-
	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	adjusted rate ³
All causes													
2001 2000 ⁴ 1999 ⁴	848.5 854.0 857.0	683.4 736.7 736.0	33.3 32.4 34.2	17.3 18.0 18.6	80.7 79.9 79.3	105.2 101.4 102.2	203.6 198.9 198.0	428.9 425.6 418.2	964.6 992.2 1,005.0	2,353.3 2,399.1 2,457.3	5,582.4 5,666.5 5,714.5	15,112.8 15,524.4 15,554.6	854.5 869.0 875.6
Diseases of heart (100-109,I11,I13,I20-I51)	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2	247.8
2000 ⁴	252.6 259.9	13.0 13.8	1.2 1.2	0.7 0.7	2.6 2.8	7.4 7.6	29.2 30.2	94.2 95.7	261.2 269.9	665.6 701.7	1,780.3 1,849.9	5,926.1 6,063.0	257.6 266.5
Malignant neoplasms (C00-C97)	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6	196.0
2000 ⁴	196.5 197.0	2.4 1.8	2.7 2.7	2.5 2.5	4.4 4.5	9.8 10.0	36.6 37.1	127.5 127.6	366.7 374.6	816.3 827.1	1,335.6 1,331.5	1,819.4 1,805.8	199.6 200.8
Cerebrovascular diseases (160-169) 2001	57.4 59.6	2.7 3.3	0.4 0.3	0.2 0.2	0.5 0.5	1.5 1.5	5.5 5.8	15.1 16.0	38.0 41.0	123.4 128.6	443.9 461.3	1,500.2 1.589.2	57.9 60.9
2000 ⁴	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8	61.6
2001 2000 ⁴	43.2 43.4	1.0 0.9	0.3 0.3	0.3 0.3	0.4 0.5	0.7 0.7	2.2 2.1	8.5 8.6	44.1 44.2	167.9 169.4	379.8 386.1	644.7 648.6	43.7 44.2
1999 ⁴ Accidents (unintentional injuries) (V01-X59,Y85-Y86)	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0	45.4
2001 2000 ⁴	35.7 34.8	24.2 23.1	11.2 11.9	6.9 7.3	36.1 36.0	29.9 29.5	35.4 34.1	34.1 32.6	30.3 30.9	42.8 41.9	100.9 95.1	276.4 273.5	35.7 34.9
1999 ⁴ Diabetes mellitus (E10-E14)	35.1 25.1	22.3	12.4	7.6 0.1	35.3 0.4	29.6	33.8	31.8	30.6 37.8	44.6 91.4	100.5	282.4 321.8	35.3
2001 2000 ⁴ 1999 ⁴	24.6 24.5	*	*	0.1 0.1 0.1	0.4 0.4 0.4	1.5 1.6 1.4	4.3 4.3 4.3	13.6 13.1 12.9	37.8 37.8 38.3	90.7 91.8	181.4 179.5 178.0	319.7 317.2	25.3 25.0 25.0
Influenza and pneumonia (J10-J18) 2001	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6	22.0
2000 ⁴ 1999 ⁴	23.2 22.8	7.6 8.4	0.7 0.8	0.2 0.2	0.5 0.5	0.9 0.8	2.4 2.4	4.7 4.6	11.9 11.0	39.1 37.2	160.3 157.0	744.1 751.8	23.7 23.5
Alzheimer's disease (G30) 2001	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3	19.1
2000 ⁴ 1999 ⁴	17.6 16.0	*	*	*	*	*	*	0.2 0.2	2.0 1.9	18.7 17.4	139.6 129.5	667.7 601.3	18.1 16.5
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10.0	2.2	*	*	0.0	0.6	4.7	4.6	10.0	40.0	104.0	007.7	14.0
2001 2000 ⁴ 1999 ⁴	13.9 13.2 12.7	3.3 4.3 4.4	*	0.1 0.1	0.2 0.2 0.2	0.6 0.6 0.6	1.7 1.6 1.6	4.6 4.4 4.0	13.0 12.8 12.0	40.2 38.0 37.1	104.2 100.8 97.6	287.7 277.8 268.9	14.0 13.5 13.0
Septicemia (A40-A41) 2001	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9	11.4
2000 ⁴ 1999 ⁴	11.1 11.0	7.2 7.5	0.6 0.6	0.2 0.2	0.3 0.3	0.7 0.7	1.9 1.8	4.9 4.6	11.9 11.4	31.0 31.2	80.4 79.4	215.7 220.7	11.3 11.3
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) 2001 ⁵	10.8			0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5	10.7
2000 ⁴	10.4 10.5			0.7 0.6	10.2 10.1	12.0 12.7	14.5 14.3	14.4 13.9	12.1 12.2	12.5 13.4	17.6 18.1	19.6 19.3	10.4 10.5
Chronic liver disease and cirrhosis (K70,K73-K74)	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2	9.5
2000 ⁴	9.4 9.4	*	*	*	0.1 0.1	1.0 1.0	7.5 7.3	17.7 17.4	23.8 23.7	29.8 30.6	31.0 31.9	23.1 23.2	9.5 9.6
2001 ⁵ 2000 ⁴	7.1 6.0	8.2 9.2	2.7 2.3	0.8 0.9	13.3 12.6	13.1 10.4	9.5 7.1	6.3 4.7	4.0 3.0	2.9 2.4	2.5 2.4	2.4 2.4	7.1 5.9
1999 ⁴	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4	6.0

See footnotes at end of table.

Table 9. Death rates by age and age-adjusted death rates for the 15 leading causes of death in 2001: United States, 1999-2001 - Con.

[Rates on an annual basis per 100,000 population in specified group; age-adjusted rates per 100,000 U.S. standard population; see Technical Notes. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years; see Technical Notes. For explanation of asterisks preceding cause-of-death codes, see Technical Notes]

	Age												
Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992) and year	All ages ¹	Under 1 year ²	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over	Age- adjusted rate ³
Essential (primary) hypertension and hypertensive renal disease (I10,I12)													
2001	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9	6.8
20004	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9	6.5
1999 ⁴ Pneumonitis due to solids and liquids (J69)	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1	6.2
2001	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4	6.1
2000 ⁴	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6	6.1
19994	5.5	*	*	*	0.1	0.2	0.4	8.0	2.5	9.5	41.1	175.6	5.6

^{*} Figure does not meet standards of reliability or precision, see Technical Notes.
. Category not applicable.
Figures for age not stated included in "All ages" but not distributed among age groups.
Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.
For method of computation, see Technical Notes.
Rates are revised and may differ from rates previously published; see Technical Notes.
Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see Technical Notes and Table D.