Table 97. Nursing home residents 65 years of age and over, according to selected functional status and age, sex, and race: United States, 1985, 1995, and 1999

[Data are based on a sample of nursing home residents]

	Functional status ¹											
	Dependent mobility			Incontinent			Dependent eating			Dependent mobility, eating, and incontinent		
Age, sex, and race	1985	1995	1999	1985	1995	1999	1985	1995	1999	1985	1995	1999
All persons	Percent											
65 years and over,	75.7	79.0	80.3	55.0	63.8	65.7	40.9	44.9	47.3	32.5	36.5	36.9
age adjusted ²	74.8	79.0	80.4	54.5	63.8	65.7	40.5	44.9	47.4	32.1	36.5	37.0
65–74 years	61.2	73.0	73.9	42.9	61.9	58.5	33.5	43.8	43.1	25.7	35.8	31.7
	70.5	76.5	77.8	55.1	62.5	64.2	39.4	45.2	46.6	30.6	35.3	35.4
	83.3	82.4	83.8	58.1	65.3	68.6	43.9	45.0	49.0	35.6	37.5	39.4
Male												
65 years and over,	71.2	76.6	76.6	54.2	63.8	66.6	36.0	42.1	45.2	28.0	34.3	35.0
age adjusted ²	67.8	75.8	75.9	51.9	63.9	66.0	34.9	42.7	45.1	26.9	34.8	35.0
65–74 years	55.8	70.6	70.5	38.8	63.4	59.6	32.8	44.2	45.0	24.1	36.9	34.8
	65.7	76.6	76.9	54.4	64.6	68.9	32.6	44.1	44.7	25.5	35.5	35.2
	79.2	78.2	78.1	58.1	63.4	66.8	39.2	40.2	45.7	30.9	32.7	34.9
Female												
65 years and over,	77.3	79.7	81.5	55.4	63.6	65.0	42.4	45.6	47.8	33.9	36.9	37.2
age adjusted ²	77.1	80.1	81.9	55.4	63.8	65.6	42.4	45.6	48.1	33.8	37.0	37.7
65–74 years	64.5	74.8	76.4	45.4	60.9	57.7	34.0	43.6	41.6	26.7	35.0	29.3
	72.3	76.5	78.2	55.3	61.7	62.2	42.0	45.7	47.4	32.6	35.2	35.6
	84.3	83.3	85.2	58.1	65.7	69.0	45.0	46.0	49.7	36.7	38.6	40.4
White ³												
65 years and over,	75.2	78.5	79.9	54.6	63.2	64.9	40.4	44.2	46.1	32.1	35.7	35.7
age adjusted ²	74.3	78.7	80.2	54.2	63.3	65.1	40.1	44.2	46.2	31.7	35.7	35.8
65–74 years	60.2	71.4	72.6	42.2	60.2	57.1	32.6	41.9	40.7	24.9	33.8	28.8
	69.6	76.4	77.5	54.2	61.8	63.8	38.9	44.9	45.8	30.1	34.7	34.8
	83.1	81.9	83.6	58.2	65.0	67.8	43.5	44.3	47.7	35.5	36.9	38.1
Black or African American ³												
65 years and over,	83.4	83.2	82.1	61.0	69.3	71.9	49.2	52.2	55.9	38.2	44.0	46.8
age adjusted ²	81.1	82.1	81.5	59.9	69.1	70.6	47.9	51.7	54.9	37.7	43.7	45.7
65–74 years	70.9	79.6	78.7	48.6	68.3	64.6	43.1	51.2	53.3	33.8	43.1	42.6
	82.5	77.8	80.1	70.1	68.9	67.5	47.9	49.5	49.7	40.6	42.3	41.0
	87.4	88.0	84.5	57.9	69.8	77.0	51.7	54.3	61.0	37.6	45.5	52.1

¹Nursing home residents who are dependent in mobility and eating require the assistance of a person or special equipment. Nursing home residents who are incontinent have difficulty in controlling bowels and/or bladder or have an ostomy or indwelling catheter.

2Age adjusted by the direct method to the 1995 National Nursing Home Survey population using the following three age groups: 65–74 years, 75–84 years, and 85

NOTES: Age refers to age at time of interview. Excludes residents in personal care or domiciliary care homes. Data for additional years are available (see Appendix III).

SOURCES: Hing E, Sekscenski E, Strahan G. The National Nursing Home Survey: 1985 summary for the United States. National Center for Health Statistics. Vital Health Stat 13(97). 1989; and Centers for Disease Control and Prevention, National Center for Health Statistics, National Nursing Home Survey for other data years.

Health, United States, 2003 287

years and over.

³Beginning in 1999 the instruction for the race item on the Current Resident Questionnaire was changed so that more than one race could be recorded. In previous years only one racial category could be checked. Estimates for racial groups presented in this table are for residents for whom only one race was recorded. Estimates for residents where multiple races were checked are unreliable due to small sample sizes and are not shown.