

FACT SHEET

Office of the Assistant Secretary of Defense (Health Affairs) **Deployment Health Support Directorate**

For more information (703) 578 - 8500 (800) 497 - 6261

Version 10-09-2002

Deseret Test Center

DTC Test 69-12

Shortly after President Kennedy's inauguration in 1961, the Secretary of Defense, Robert McNamara, directed that a total review of the U.S. military be undertaken. The study consisted of 150 separate projects. The chemical and biological warfare review was known as Project 112. As part of the Project 112 review, the Joint Chiefs of Staff convened a working committee that recommended a research, testing, and development program for chemical and biological weapons. To oversee this program, the Deseret Test Center was established at Fort Douglas, Utah, in 1962. Both land-based and ship-based tests were conducted during the period 1962 – 1973. The Deseret Test Center closed in 1973.

In 1967 and 1968, Deseret Test Center Test conducted DTC Test 68-13 (Rapid Tan I, II, III) jointly with the United Kingdom and Canada. Rapid Tan was designed to investigate the extent and duration of hazard following a Tabun, Soman or V nerve agent attack. DTC Test 69-12 was planned as a more sophisticated test than Rapid Tan. DTC Test 69-12 was originally scheduled for conduct near Fort Greely, Alaska; however, the test site was moved to Edgewood Arsenal, Maryland. Only three trials (of 54 scheduled) were completed prior to the imposition of open-air toxic test restrictions and the suspension of the test.

The three completed DTC Test 69-12 trials were conducted at Edgewood Arsenal, Maryland during the spring of 1969.

Test Name	DTC Test 69-12
Testing Organization	US Army Deseret Test Center
Test Dates	Spring 1969
Test Location	Edgewood Arsenal, Maryland
Test Operations	To determine rate of evaporation of Tabun, Sarin, Soman, and VX as a function of contamination density, drop size, and terrain cover under a variety of meteorological conditions in a temperate environment.
Participating Services	Deseret Test Center Personnel
Units and Ships Involved	Not identified
Dissemination Procedures	Not identified
Agents, Simulants, Tracers	Sarin Nerve Agent Soman Nerve Agent Tabun Nerve Agent VX Nerve Agent
Ancillary Testing	Not identified
Decontamination	Not identified
Potential Health Risks Associated with Agents, Simulants, Tracers	Sarin Nerve Agent (GB) Sarin gas is a volatile and lethal nerve agent. It can enter the body by inhalation, ingestion, through the eyes, and to a lesser extent through the skin. After exposure to a sufficient dose, human symptoms may occur within minutes and include runny nose, watery eyes, difficulty breathing, dimness of vision, confusion, drowsiness, coma, and death. Very little information is available regarding long-term health effects following exposures to low levels that do not cause acute symptoms. No information is available regarding potential carcinogenicity. An Institute of

Medicine committee concluded that there was insufficient evidence for or against an association between low-level sarin exposure and long-term health effects.

(Sources: http://www.bt.cdc.gov/Agent/Nerve/Sarin/Sarin.asp [as of February 13, 2002]Institute of Medicine (National Academies), Gulf War and Health (vol.1): Depleted Uranium, Pyridostigmine Bromide, Sarin, Vaccines. National Academy Press, Washington DC, 2000.)

Soman Nerve Agent (GD)

Soman is a colorless liquid, which gives off an odor of rotting fruit when vaporizing. The vapor is colorless. Soman is a persistent agent that can easily remain in a particular area for a day or longer, depending on the atmospheric conditions. Acute health effects associated with exposure to soman include a runny nose, tightness in the chest, constriction of the pupils, difficulty in breathing, coma, and death. There is little information available regarding the long-term human health effects of exposure to soman.

(Source: http://www.sbccom.army.mil/services/edu/soman.htm Zajtchuk R (ed.), Textbook of Military Medicine (part 1, Medical Aspects of Chemical and Biological Warfare, 1997), Office of the Army Surgeon General, Washington DC, 1997.)

Tabun Nerve Agent (GA)

Tabun is an amber, non-persistent liquid, which gives off little odor when vaporizing. The vapor is colorless. When exposed to tabun, the symptoms a victim will experience include a runny nose, tightness in the chest, constriction of the pupils, difficulty breathing,

and nausea. Ultimately the victim will become comatose and will suffocate as a consequence of convulsive spasms. Tabun is mainly absorbed through the skin; however, vapors can also be hazardous. If a person does not receive an immediately lethal dose, death will occur after approximately 20 minutes. Those receiving a less than lethal dose who do not receive immediate medical care may suffer permanent neurological damage. There is little information available regarding the long-term human health effects of exposure to low doses of tabun.

(Source: http://www.sbccom.army.mil/services/edu/tabun.htm Zajtchuk R (ed.), Textbook of Military Medicine (part 1, Medical Aspects of Chemical and Biological Warfare, 1997), Office of the Army Surgeon General, Washington DC, 1997.

<u>VX Nerve Agent</u> – (Synonyms: Phosphonothioic acid, VX)

VX nerve agent is extremely lethal. It is an oily liquid that is clear, odorless, and tasteless. Death usually occurs within 10-15 minutes after absorption of a fatal dosage. VX nerve agent is one of the most toxic substances ever synthesized. Symptoms of overexposure may occur within minutes or hours, depending upon the dose. They include: constriction of pupils, headaches, runny nose, salivation, tightness in the chest, nausea, vomiting, anxiety, difficulty in thinking, muscle twitches, tremors, and weakness. With severe exposure, symptoms progress to convulsions and respiratory failure. There is little information available regarding the long-term human health effects of exposure to low doses of VX.

(Sources: Centers for Disease Control and Prevention http://www.bt.cdc.gov/Agent/Nerve/ VX/ctc0006.asp [as of January 25, 2002] Zajtchuk R (ed.), Textbook of Military Medicine (part 1, Medical Aspects of Chemical and Biological Warfare, 1997), Office of the Army Surgeon General, Washington DC, 1997. SBCCOM Online, Edgewood Chemical Biological Center http://in1.apgea.army.mil:80/RDA/msds/vx.htm [as of April 2, 2002] World Health Organization, Department of Sustainable Development & Environmental Protection, http://209.61.192.180/ phe/factsheet_5.htm [as of April 2, 2002] Department of the Army Pamphlet 40-8: Occupational Health Guidelines for the Evaluation and Control of Occupational Exposure to Nerve Agents GA, GB, GD, and VX http://books.army.mil:80/cgi-bin/bookmgr/BOOKS/ P40_8/CCONTENTS [as of February 5, 2002]).