


Number of Maryland youth projected to die prematurely from their smoking: 85,720

Adult Tobacco Use in Maryland

Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



Youth Tobacco Use

	GRADES 6-8		GRADES 9-12	
	Current Cigarette Smoking	Current Any Tobacco Use	Current Cigarette Smoking	Current Any Tobacco Use
National*	9.2%	12.8%	28.5%	34.8%
MD†	7.3%	11.8%	23.7%	29.9%
Boys†	7.3%	12.5%	23.3%	32.1%
Girls†	7.3%	10.8%	23.8%	27.4%

Current Cigarette Smoking = smoked cigarettes on ≥1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥1 of the 30 days preceding the survey.

*Source: National Youth Tobacco Survey, 1999.

†Source: Maryland Youth Tobacco Survey, 2000.


Health Impact and Costs

AVERAGE ANNUAL DEATHS RELATED TO SMOKING, 1990-1994		AVERAGE ANNUAL YEARS OF POTENTIAL LIFE LOST,* 1990-1994	MEDICAL COSTS RELATED TO SMOKING, 1993	
Overall	7,180	102,119 years or an average of 14.2 years for each death due to smoking. *Calculated to life expectancy	Ambulatory	\$273,890,000
Men	4,480		Hospital	\$303,130,000
Women	2,701		Nursing Home†	\$279,670,000
Death Rate	351/100,000		Drug	\$83,670,000
Rank	26 (No. 1 is lowest death rate)		Other	\$162,780,000
			Total	\$1,103,150,000

Smoking-attributable Medicaid expenditures, Maryland, fiscal year 1993: \$212,304,000

†Preliminary estimates

Lung Cancer Death Rate*



*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.

Maryland





Scheduled 2001 settlement payment to Maryland: \$156,506,355.69

Tobacco Control Funding, 2001

At 71% of the CDC Best Practices recommended lower estimate funding level, Maryland ranks 11th (No. 1 is the highest) for tobacco control funding. Maryland ranks 26th (No. 1 is the lowest death rate) for average annual deaths related to smoking.

<i>FUNDING SOURCE</i>	<i>FY01 AMOUNT</i>	<i>FUNDING CYCLE</i>
State Appropriation—Settlement (Tobacco Only)	\$18,065,486	7/00–6/01
State Appropriation—Excise Tax Revenue	\$1,893,000	7/00–6/01
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$19,958,486	
Federal—CDC Office on Smoking and Health	\$1,370,605	6/00–5/01
Federal—SAMHSA	n/a	
Non-Government Source—American Legacy Foundation	\$99,207	10/00–9/01
Non-Government Source—RWJF/AMA	n/a	
Subtotal: Federal/National Sources	\$1,469,812	
Total Per Capita Funding	\$21,428,298 \$4.05	

Funding as a Percentage of CDC Best Practices Recommendations

	<i>RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST</i>	<i>RECOMMENDED PER CAPITA FUNDING LEVEL</i>	<i>PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS</i>
Lower Estimate	\$30,301,000	\$5.95	71%
Upper Estimate	\$78,601,000	\$15.43	27%

Excise Tax

Cigarette tax per pack	66¢
Rank = 12 (No. 1 is highest tax)	
Number of packages of cigarettes sold and taxed, per capita, 1999	71.4
Rank = 13 (No. 1 is lowest cigarette sales)	
Federal and state taxes as a percentage of retail price	28.6%
Annual gross tax revenue from cigarettes	\$131,338,000
Smokeless tobacco tax15% of wholesale sales price