


Number of South Dakota youth projected to die prematurely from their smoking: 14,626

Adult Tobacco Use in South Dakota

Current Cigarette Smoking Among Adults Aged 18 and Older, 1999



Youth Tobacco Use

	GRADES 6-8		GRADES 9-12	
	Current Cigarette Smoking	Current Any Tobacco Use	Current Cigarette Smoking	Current Any Tobacco Use
National	9.2%*	12.8%*	28.5%*	34.8%*
SD	12.4%†	16.3%†	43.6%**	49.4%**
Boys	14.5%†	20.4%†	41.8%**	51.6%**
Girls	10.0%†	11.7%†	45.5%**	47.4%**

Current Cigarette Smoking = smoked cigarettes on ≥1 of the 30 days preceding the survey.

Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or cigars or kreteks on ≥1 of the 30 days preceding the survey.

*Source: National Youth Tobacco Survey, 1999.

†Source: South Dakota Youth Tobacco Survey, 1999.

**Source: South Dakota Youth Risk Behavior Survey, 1999.


Health Impact and Costs

AVERAGE ANNUAL DEATHS RELATED TO SMOKING, 1990-1994		AVERAGE ANNUAL YEARS OF POTENTIAL LIFE LOST,* 1990-1994	MEDICAL COSTS RELATED TO SMOKING, 1993	
Overall	1,198	14,705 years or an average of 12.3 years for each death due to smoking. *Calculated to life expectancy	Ambulatory	\$24,760,000
Men	860		Hospital	\$40,910,000
Women	338		Nursing Home†	\$47,350,000
Death Rate	309/100,000		Drug	\$6,700,000
Rank (No. 1 is lowest death rate)	9		Other	\$14,840,000
			Total	\$134,550,000

Smoking-attributable Medicaid expenditures, South Dakota, fiscal year 1993: \$20,740,000

†Preliminary estimates

Lung Cancer Death Rate*



*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.

Scheduled 2001 settlement payment to South Dakota: \$24,159,821.39

Tobacco Control Funding, 2001



At 36% of the CDC Best Practices recommended lower estimate funding level, South Dakota ranks 27th (No. 1 is the highest) for tobacco control funding. South Dakota ranks 9th (No. 1 is the lowest death rate) for average annual deaths related to smoking.

<i>FUNDING SOURCE</i>	<i>FY01 AMOUNT</i>	<i>FUNDING CYCLE</i>
State Appropriation—Settlement (Tobacco Only)	\$0	
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$1,700,000	7/00–6/01
Subtotal: State Appropriation	\$1,700,000	
Federal—CDC Office on Smoking and Health	\$886,491	6/00–5/01
Federal—SAMHSA	n/a	
Non-Government Source—American Legacy Foundation	\$500,000	10/00–9/01
Non-Government Source—RWJF/AMA	n/a	
Subtotal: Federal/National Sources	\$1,386,491	
Total	\$3,086,491	
Per Capita Funding	\$4.09	

Funding as a Percentage of CDC Best Practices Recommendations

	<i>RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST</i>	<i>RECOMMENDED PER CAPITA FUNDING LEVEL</i>	<i>PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS</i>
Lower Estimate	\$8,688,000	\$11.77	36%
Upper Estimate	\$18,214,000	\$24.68	17%

Excise Tax

Cigarette tax per pack Rank = 28 (No. 1 is highest tax)	33¢
Number of packages of cigarettes sold and taxed, per capita, 1999 Rank = 24 (No. 1 is lowest cigarette sales)	.83.5
Federal and state taxes as a percentage of retail price	20.9%
Annual gross tax revenue from cigarettes	\$20,327,478
Smokeless tobacco tax	10% of wholesale purchase price