


Number of Wyoming youth projected to die prematurely from their smoking: 10,134**Adult Tobacco Use in Wyoming****Current Cigarette Smoking Among Adults Aged 18 and Older, 1999****Youth Tobacco Use**

	GRADES 6-8		GRADES 9-12	
	Current Cigarette Smoking	Current Any Tobacco Use	Current Cigarette Smoking	Current Any Tobacco Use
National	9.2%*	12.8%*	28.5%*	34.8%*
WY	14.8%†	21.7%†	35.2%**	43.9%**
Boys	15.6%†	25.8%†	34.6%**	49.6%**
Girls	13.4%†	16.9%†	35.9%**	38.0%**

Current Cigarette Smoking = smoked cigarettes on ≥ 1 of the 30 days preceding the survey.Current Any Tobacco Use = current use of cigarettes or smokeless tobacco or pipes or bidis or kreteks on ≥ 1 of the 30 days preceding the survey.

*Source: National Youth Tobacco Survey, 1999.

†Source: Wyoming Youth Tobacco Survey, 2000.


**Source: Wyoming Youth Risk Behavior Survey, 1999.

Health Impact and Costs

AVERAGE ANNUAL DEATHS RELATED TO SMOKING, 1994		AVERAGE ANNUAL YEARS OF POTENTIAL LIFE LOST,* 1994	MEDICAL COSTS RELATED TO SMOKING, 1993	
Overall	712	9,271 years or an average of 13.0 years for each death due to smoking. *Calculated to life expectancy	Ambulatory	\$12,600,000
Men	462		Hospital	\$22,600,000
Women	250		Nursing Home†	\$19,190,000
Death Rate	357/100,000		Drug	\$4,270,000
Rank	30 (No. 1 is lowest death rate)		Other	\$12,560,000
			Total	\$71,210,000

Smoking-attributable Medicaid expenditures, Wyoming, fiscal year 1993: \$11,449,000

†Preliminary estimates

Lung Cancer Death Rate*

*1997 deaths per 100,000 population, adjusted to the 1970 total U.S. population.

Scheduled 2001 settlement payment to Wyoming: \$17,194,554.25

Tobacco Control Funding, 2001

At 25% of the CDC Best Practices recommended lower estimate funding level, Wyoming ranks 32nd (No. 1 is the highest) for tobacco control funding. Wyoming ranks 30th (No. 1 is the lowest death rate) for average annual deaths related to smoking.

<i>FUNDING SOURCE</i>	<i>FY01 AMOUNT</i>	<i>FUNDING CYCLE</i>
State Appropriation—Settlement (Tobacco Only)	\$900,000	7/00–6/01
State Appropriation—Excise Tax Revenue	\$0	
State Appropriation—Other	\$0	
Subtotal: State Appropriation	\$900,000	
Federal—CDC Office on Smoking and Health	\$973,302	6/00–5/01
Federal—SAMHSA	n/a	
Non-Government Source—American Legacy Foundation	\$0	
Non-Government Source—RWJF/AMA	n/a	
Subtotal: Federal/National Sources	\$973,302	
Total Per Capita Funding	\$1,873,302 \$3.79	

In FY2000, \$1,800,000 was appropriated for the biennial budget cycle for tobacco use prevention and control programs.

Funding as a Percentage of CDC Best Practices Recommendations

	<i>RECOMMENDED FOR TOTAL PROGRAM ANNUAL COST</i>	<i>RECOMMENDED PER CAPITA FUNDING LEVEL</i>	<i>PERCENT OF CDC BEST PRACTICES RECOMMENDATIONS</i>
Lower Estimate	\$7,381,000	\$15.39	25%
Upper Estimate	\$14,397,000	\$30.01	13%

Excise Tax

Cigarette tax per pack	12¢
Rank = 46 (No. 1 is highest tax)	
Number of packages of cigarettes sold and taxed, per capita, 1999	104.8
Rank = 40 (No. 1 is lowest cigarette sales)	
Federal and state taxes as a percentage of retail price	13.5%
Annual gross tax revenue from cigarettes	\$6,037,000
Smokeless tobacco tax20% of wholesale purchase price