


EPICHLOROHYDRIN

CAS No. 106-89-8

First Listed in the *Fourth Annual Report on Carcinogens*



CARCINOGENICITY

Epichlorohydrin is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity in experimental animals (IARC 1976, 1982, 1987, 1999). When administered by gavage, epichlorohydrin induced carcinomas and papillomas of the forestomach in male rats. When administered by inhalation, the compound induced carcinomas and papillomas of the nasal cavity in male rats. Subcutaneous injection of epichlorohydrin produced local sarcomas in female mice. Epichlorohydrin also was active as an initiator in a two-stage skin carcinogenesis study of female mice, but it did not induce tumors when applied alone to the skin.

There is inadequate evidence for the carcinogenicity of epichlorohydrin in humans (IARC 1976, 1982, 1987, 1999). One study of factory workers exposed to epichlorohydrin shows a significant excess of respiratory cancer; however, some of the workers may have also been exposed to diisopropyl sulfate.

PROPERTIES

Epichlorohydrin is a colorless liquid with an irritating, chloroform-like odor. It is slightly soluble in water, miscible with alcohol, ether, chloroform, trichloroethylene, and carbon tetrachloride and insoluble in petroleum hydrocarbons (HSDB 2000). Epichlorohydrin hydrolyzes slowly at room temperature and more rapidly in the presence of heat or traces of acid. When heated to decomposition, epichlorohydrin emits toxic fumes of hydrochloric acid and other chlorinated compounds. The commercial product is 98% pure with a maximum of 0.2% water.

USE

Epichlorohydrin has been used in the production of various synthetic materials, including epoxy resins (68% of the epichlorohydrin produced), synthetic glycerin (19%), elastomers (3%), and others (10%) (NCI 1985). It has also been used to cure propylene-base rubbers, as a solvent for cellulose esters and ethers and in resins with high wet-strength for the paper industry (IARC 1999). Epichlorohydrin is also used in the production of Zeospan, a specialty polyether rubber used for automobile parts (Chem. Week 1986). There is widespread use of epichlorohydrin as a stabilizer.

PRODUCTION

Chem Sources (2001) identified 20 U.S. suppliers of epichlorohydrin. The 1997 *Directory of Chemical Producers* listed two domestic producers of the chemical, with a total output of 725 million lb (SRI 1997). The USITC has listed two companies producing undisclosed amounts of epichlorohydrin since 1980 (USITC 1981-1991, 1993-1995). The

Chemical Economics Handbook, however, provided production and import and export data through 1989 (SRI 1991). In 1989, 495 million lb of epichlorohydrin were produced in the U.S. with 18 million lb imported and 60 million lb exported (SRI 1991).

EXPOSURE

The primary routes of potential human exposure to epichlorohydrin are ingestion, inhalation, and dermal contact. Levels of epichlorohydrin in food, food additives, or food packaging are so low that the potential daily intake through ingestion is minimal. According to the FDA, epichlorohydrin is no longer used in the starch industry. Occupational exposure is possible during its production and during the synthesis of its end products. Evidence suggests that epichlorohydrin is readily absorbed when ingested or inhaled and that it is a systemic poison when absorbed through the skin (NCI 1985, NIOSH 1978). The National Occupational Exposure Survey (1981-1983) indicated that 3,306 workers, including 558 women, potentially were exposed to epichlorohydrin (NIOSH 1984). This estimate was derived from observations of the actual use of the compound (80% of total observations) and the use of trade name products known to contain the compound (20%). The National Occupational Hazard Survey, conducted by NIOSH from 1972 to 1974, estimated that 41,380 people in 58 different occupational categories were potentially exposed to epichlorohydrin in the workplace. This estimate was based only on actual observations of the use of the compound and trade name products containing or suspected of containing the compound (NIOSH 1976). Comprehensive industrial surveys conducted for NIOSH included five facilities: two manufacturing epichlorohydrin and three producing epoxy resins and glycerin. The survey results suggested that chemical operators at these plants had the greatest potential for exposure to epichlorohydrin. Time-weighted averages (TWAs) ranged from nondetectable to 2.1 ppm in the two manufacturing plants and from nondetectable to 0.83 ppm in the resin production plants. A limited number of samples also indicated a relatively high potential for epichlorohydrin exposure for shipping workers in epichlorohydrin production plants (0.06 to 0.28 ppm TWA) (NIOSH 1978, NCI 1985).

EPA's Toxic Chemical Release Inventory (TRI) estimated that 353,964 lb of epichlorohydrin were released to the environment from 48 facilities that produced, processed, or used the chemical in the United States in 1996. Of that total, 93.5% was released to air, 5.9% to water, and 0.6% to land. A facility located in Freeport, Texas, reporting under industrial classifications for manufacture of alkalis and chlorine (SIC Code 2812), industrial gases (2813), industrial inorganic chemicals (2819), plastics materials and resins (2821), industrial organic chemicals (2869), and adhesives and sealants (2891), accounted for 41.4% of the total air emissions and 91.6% of the total water release (TRI96 1998). The total amount of epichlorohydrin released on-site in 1999 (TRI99 2001) was 148,532 lb. Total on-site release data from 1988 showed a continued decrease from the 1988 level of 783,298 lb.

REGULATIONS

EPA regulates epichlorohydrin under the Clean Water Act (CWA), Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), Food, Drug, and Cosmetic Act (FD&CA), Resource Conservation and Recovery Act (RCRA), Superfund Amendments and Reauthorization Act (SARA), and Toxic Substances Control Act (TSCA). A reportable quantity (RQ) of 100 lb has been established for epichlorohydrin under CERCLA. Epichlorohydrin is regulated by EPA as a toxic inert ingredient in pesticides under FD&CA. It is listed as a toxic hazardous substance under RCRA and SARA, and it is subject to reporting rules under TSCA.

FDA regulates epichlorohydrin as a food additive.

ACGIH recommends a threshold limit value (TLV) at 0.5 ppm (1.9 mg/m³); the potential for skin absorption was noted. NIOSH has recommended that epichlorohydrin be treated as a human carcinogen, that exposure be limited to the lowest level possible, and that skin exposure be avoided. OSHA has set a permissible exposure limit (PEL) of 5 ppm (19 mg/m³) as an 8-hr TWA. OSHA also regulates epichlorohydrin under the Hazard Communication Standard and as a chemical hazard in laboratories. Regulations are summarized in Volume II, Table 82.

REFERENCES

Chem Sources. Chemical Sources International, Inc. <http://www.chemsources.com>, 2001.

Chemical Week. Vol. 138, No. 8, 1986, p. 21.

HSDB. Hazardous Substances Data Bank. Online database produced by the National Library of Medicine. Epichlorohydrin. Profile last updated June 12, 2000. Last review date, January 29, 2000.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. Cadmium, Nickel, Some Epoxides, Miscellaneous Industrial Chemicals and General Considerations on Volatile Anaesthetics. Vol. 11. 306 pp. Lyon, France: IARC, 1976.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Humans. Chemicals, Industrial Processes and Industries Associated with Cancer in Humans. Supplement 4. 292 pp. Lyon, France: IARC, 1982.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Overall Evaluations of Carcinogenicity. Supplement 7. 440 pp. Lyon, France: IARC, 1987.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Re-evaluation of Some Organic Chemicals, Hydrazine and Hydrogen Peroxide. Vol. 71, part 2. 603 pp. Lyon, France: IARC, 1999.

NCI. National Cancer Institute, Division of Cancer Etiology. Monograph on Human Exposure to Chemicals in the Workplace: Epichlorohydrin. Technical Report No. 84-631. Bethesda, MD: Department of Health and Human Services, 1985.

NIOSH. National Institute for Occupational Safety and Health. National Occupational Hazard Survey (1972-74). Cincinnati, OH: Department of Health, Education, and Welfare, 1976.

NIOSH. National Institute for Occupational Safety and Health. Epichlorohydrin Manufacture and Use -- Industrial Hygiene Survey. DHEW (NIOSH) Publication No. 78-151. Cincinnati, OH: Department of Health, Education, and Welfare, 1978.

NIOSH. National Institute for Occupational Safety and Health. National Occupational Exposure Survey (1980-83). Cincinnati, OH: Department of Health and Human Services, 1984.

SRI. Directory of Chemical Producers, United States, 1997. Stanford Research Institute, Menlo Park, CA: SRI International, 1997.

SRI. Chemical Economics Handbook. Stanford Research Institute, Menlo Park, CA: SRI International, 1991.

TRI96. Toxic Chemicals Release Inventory 1996. Data contained in the Toxic Chemical Release Inventory (TRI). Available from the National Library of Medicine's TOXNET System, 1998.

TRI99. Toxic Chemicals Release Inventory 1999. Data contained in the Toxic Chemical Release Inventory (TRI). Available from the U.S. Environmental Protection Agency Office of Environmental Information, <http://www.epa.gov/triexplorer/reports.htm>, 2001.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1980. USITC Publication No. 1183. Washington, DC: U.S. Government Printing Office, 1981.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1981. USITC Publication No. 1292. Washington, DC: U.S. Government Printing Office, 1982.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1982. USITC Publication No. 1422. Washington, DC: U.S. Government Printing Office, 1983.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1983. USITC Publication No. 1588. Washington, DC: U.S. Government Printing Office, 1984.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1984. USITC Publication No. 1745. Washington, DC: U.S. Government Printing Office, 1985.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1985. USITC Publication No. 1892. Washington, DC: U.S. Government Printing Office, 1986.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1986. USITC Publication No. 2009. Washington, DC: U.S. Government Printing Office, 1987.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1987. USITC Publication No. 2118. Washington, DC: U.S. Government Printing Office, 1988.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1988. USITC Publication No. 2219. Washington, DC: U.S. Government Printing Office, 1989.

Epichlorohydrin (Continued)

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1989. USITC Publication No. 2338. Washington, DC: U.S. Government Printing Office, 1990.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1990. USITC Publication No. 2470. Washington, DC: U.S. Government Printing Office, 1991.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1991. USITC Publication No. 2607. Washington, DC: U.S. Government Printing Office, 1993.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1992. USITC Publication No. 2720. Washington, DC: U.S. Government Printing Office, 1994.

USITC. U.S. International Trade Commission. Synthetic Organic Chemicals, United States Production and Sales, 1994. USITC Publication No. 2933. Washington, DC: U.S. Government Printing Office, 1995.