


## METHYLEUGENOL

CAS No. 93-15-2

First Listed in the *Tenth Report on Carcinogens*



### CARCINOGENICITY

Methyleugenol is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of its carcinogenicity in experimental animals. In animal studies, methyleugenol given orally to rats induced liver and stomach tumors in both sexes and kidney, mammary gland, and skin tumors in males. Methyleugenol given orally to mice induced benign and malignant tumors of the liver. Tumors of the stomach in male mice also were considered related to exposure to methyleugenol (NTP 1998). Earlier studies found that methyleugenol and two similar compounds, the structurally related allylbenzenes, safrole and estragole, induced liver tumors in mice after intraperitoneal injection (IARC 1976, Miller *et al.* 1983). Safrole is classified by IARC as *possibly carcinogenic to humans* (Group 2B) and in the Report on Carcinogens as *reasonably anticipated to be a human carcinogen*.

No adequate human studies of the relationship between exposure to methyleugenol and human cancer were found.

### ADDITIONAL INFORMATION RELEVANT TO CARCINOGENESIS OR POSSIBLE MECHANISMS OF CARCINOGENESIS

Mechanistic data indicate that liver tumors induced by methyleugenol and structurally related allylbenzenes result from metabolism of these compounds to DNA-reactive intermediates. Methyleugenol may be bioactivated by three different pathways: (1) hydroxylation at the 1' position of the allylic side chain to yield 1'-hydroxymethyleugenol, followed by sulfation of this intermediate to form 1'-hydroxymethyleugenol sulfate, (2) oxidation of the 2',3'-double bond of the allylic side chain to form methyleugenol-2,3-oxide, and (3) *O*-demethylation followed by spontaneous rearrangement to form eugenol quinone methide. Formation of protein adducts and DNA adducts in the livers of animals (and in cultured human hepatocytes) exposed to allylbenzenes and induction of liver tumors by these compounds in animals have been attributed to activation via the hydroxylation pathway, because similar effects were produced by the 1'-hydroxy metabolites and because these effects were inhibited by pretreatment with sulfotransferase inhibitors (Miller *et al.* 1983, Boberg *et al.* 1983, Randerath *et al.* 1984, Gardner *et al.* 1996, NTP 1998).

Methyleugenol, safrole, and estragole induce unscheduled DNA synthesis in rat hepatocytes, and their corresponding 1'-hydroxy metabolites are more potent genotoxic agents than are the parent compounds (Howes *et al.* 1990, Chan and Caldwell 1992). Methyleugenol induces morphological transformations in Syrian hamster embryo cells (Kerckaert *et al.* 1996), sister chromatid exchange in Chinese hamster ovary (CHO) cells

(NTP 1998), intrachromosomal recombination in yeast (Schiestl *et al.* 1989), and DNA repair in *Bacillus subtilis* (Sekizawa and Shibamoto 1982). Methyleugenol does not induce mutations in *Salmonella typhimurium* (NTP 1998) or *Escherichia coli* (Sekizawa and Shibamoto 1982), chromosomal aberrations in CHO cells (NTP 1998), or micronucleated erythrocytes in peripheral blood of mice (NTP 1998). A higher frequency of  $\beta$ -*catenin* mutations was observed in liver tumors from mice treated with methyleugenol than in spontaneous liver tumors from control mice (Devereux *et al.* 1999). Methyleugenol's lack of mutagenicity in bacteria may be due to the need for sulfation in the metabolic activation of methyleugenol to its ultimate mutagenic or carcinogenic form.

## PROPERTIES

Methyleugenol is a colorless to pale yellow oily liquid with a boiling point of 254.7°C and a melting point of -4°C. It has a delicate clover-carnation odor and a bitter burning taste. It forms azeotropic mixtures with ethylene glycol, eugenol, and benzoic acid. It slowly darkens and thickens when exposed to air and readily evaporates at room temperature (Lide 1998). It is soluble in water, ether, ethanol, and chloroform and insoluble in glycol and propylene glycol (HSDB 2001).

## USE

Methyleugenol is used as a flavoring agent in jellies, baked goods, nonalcoholic beverages, chewing gum, candy, pudding, relish, and ice cream. Methyleugenol has been used as an anesthetic in rodents (NTP 1998). It also is used as an insect attractant in combination with insecticides (NTP 1998). Methyleugenol has been used as an agent in sunscreens (NTP 2001).

## PRODUCTION

Annual production of methyleugenol in the United States in 1990 was estimated at 25,000 lb (NTP 1998). No current production data were available for methyleugenol.

## EXPOSURE

Methyleugenol is a naturally occurring substance, present in many essential oils (NTP 2001). It is a component of rose, pimento, basil, hyacinth, citronella, anise, nutmeg, mace, cinnamon leaves, pixuri seeds, and laurel fruits and leaves. It also has been found in blackberry essence, bananas, black pepper, and bilberries (NTP 1998). Methyleugenol is used in commercial products as a flavorant at concentrations from 5 to 52 ppm and as a fragrance at concentrations from 0.002% to 0.3%. Methyleugenol has been detected in the wastewater effluent from a paper mill (NTP 1998). Although methyleugenol has been identified in various natural substances, no quantitative studies have assessed environmental (nondietary) exposure to methyleugenol. Methyleugenol exists as a vapor in the ambient atmosphere. Vapor-phase methyleugenol reacts with photochemically produced hydroxyl radicals and degrades with an estimated half-life of five hours (HSDB 2001).

The general population is exposed to methyleugenol through ingestion of foodstuffs or inhalation of fragrances containing the compound (HSDB 2001). In a subset of serum samples from adults participating in the third National Health and Nutrition Examination Survey, methyleugenol was detected in 98% of the 206 samples analyzed. The average methyleugenol concentration was 24 pg/g, and the highest concentration was 390 pg/g (Barr *et al.* 2000).

Daily per capita consumption of methyleugenol in foods was estimated by the World Health Organization to be 0.073 mg (WHO 1981) and, more recently, 0.26 mg/kg body weight (Stroberg and Grundschober 1987, NAS 1989). Occupational exposure to methyleugenol occurs through dermal contact, inhalation, and ingestion. The National Occupational Exposure Survey conducted by NIOSH between 1981 and 1983 estimated that 2,824 workers, including 877 women, potentially were exposed to methyleugenol (NIOSH 1990).

## REGULATIONS

The U.S. Environmental Protection Agency (EPA) regulates methyleugenol under the Federal Insecticide, Fungicide, and Rodenticide Act. EPA allows for exemption from normal tolerances when methyleugenol is used in Oriental fruit fly eradication programs.

The U.S. Food and Drug Administration regulates methyleugenol, allowing it to be used as a synthetic flavoring substance and adjuvant for direct addition to food for human consumption.

The Occupational Safety and Health Administration regulates methyleugenol under the Hazard Communication Standard and as a chemical hazard in laboratories. Regulations are summarized in Volume II, Table 108.

## REFERENCES

Barr, D.B., J.R. Barr, S.L. Bailey, C.R. Lapeza, Jr., M.D. Beeson, S.P. Caudill, V.L. Maggio, A. Schechter, S.A. Masten, G.W. Lucier, L.L. Needham, and E.J. Sampson. Levels of methyleugenol in a subset of adults in the general U.S. population as determined by high resolution mass spectrometry. *Environ. Health Perspect.*, Vol. 108, 2000, pp. 323-328.

Boberg, E.W., E.C. Miller, J.A. Miller, A. Poland, and A. Liem. Strong evidence from studies with brachymorphic mice and pentachlorophenol that 1'-sulfoxysafrole is the major ultimate electrophilic and carcinogenic metabolite of 1'-hydroxysafrole in mouse liver. *Cancer Res.*, Vol. 43, 1983, pp. 5163-5173.

Chan, V.S., and J. Caldwell. Comparative induction of unscheduled DNA synthesis in cultured rat hepatocytes by allylbenzenes and their 1'-hydroxy metabolites. *Food Chem. Toxicol.*, Vol. 30, 1992, pp. 831-836.

Devereux, T.R., C.H. Anna, J.F. Foley, C.M. White, R.C. Sills, and J.C. Barrett. Mutation of beta-catenin is an early event in chemically induced mouse hepatocellular carcinogenesis. *Oncogene*, Vol. 18, 1999, pp. 4726-4733.

Gardner, I., P. Bergin, P. Stening, J.G. Kenna, and J. Caldwell. Immunochemical detection of covalently modified protein adducts in livers of rats treated with methyleugenol. *Chem. Res. Toxicol.*, Vol. 9, 1996, pp. 713-721.

Howes, A.J., V.S. Chan, and J. Caldwell. Structure-specificity of the genotoxicity of some naturally occurring alkenylbenzenes determined by the unscheduled DNA synthesis assay in rat hepatocytes. *Food Chem. Toxicol.*, Vol. 28, 1990, pp. 537-542.

HSDB. Hazardous Substances Data Bank. Online database produced by the National Library of Medicine. Methyleugenol. Profile last updated May 16, 2001. Last review date, May 16, 1996.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Some Naturally Occurring Substances. Vol. 10. Lyon, France: IARC, 1976, 331 pp.

Kerckaert, G.A., R. Brauninger, R.A. LeBoeuf, and R.J. Isfort. Use of the Syrian hamster embryo cell transformation assay for carcinogenicity prediction of chemicals currently being tested by the National Toxicology Program in rodent bioassays. *Environ Health. Perspect.*, Vol. 104, 1996, pp. 1075-1084.

Lide, D.R., ed. *CRC Handbook of Chemistry and Physics*. Washington, DC: CRC Press, 1998.

Miller, E.C., A.B. Swanson, D.H. Phillips, T.L. Fletcher, A. Liem, and J.A. Miller. Structure-activity studies of the carcinogenicities in the mouse and rat of some naturally occurring and synthetic alkenylbenzene derivatives related to safrole and estragole. *Cancer Res.*, Vol. 43, 1983, pp. 1124-1134.

NAS. National Academy of Sciences. Poundage and Technical Effects Update of Substances Added to Food. Committee on Food Additives Survey Data, Food and Nutrition Board, Institute of Medicine. Washington, DC: National Academy of Sciences, 1989.

NIOSH. National Institute for Occupational Safety and Health. National Occupational Exposure Survey (NOES) (1981-1983). Unpublished provisional data as of July 1, 1990, Cincinnati OH.

NTP. National Toxicology Program. Toxicology and Carcinogenesis Studies of Methyleugenol (CAS No. 93-15-12) in F344/N Rats and B6C3F1 Mice (Gavage Studies). Technical Report Series TR-491. Research Triangle Park, NC: NTP, 1998.

NTP. National Toxicology Program. NTP Chemical Repository. Methyleugenol. Last revised August 13, 2001. (<http://ntp-server.niehs.nih.gov> & search 93-15-2).

Randerath, K., R.E. Haglund, D.H. Phillips, and M.V. Reddy. <sup>32</sup>P-post-labelling analysis of DNA adducts formed in the livers of animals treated with safrole, estragole and other naturally-occurring alkenylbenzenes. I. Adult female CD-1 mice. *Carcinogenesis*, Vol. 5, 1984, pp. 1613-1622.

Schiestl, R.H., W.S. Chan, R.D. Gietz, R.D. Mehta, and P.J. Hastings. Safrole, eugenol and methyleugenol induce intrachromosomal recombination in yeast. *Mutat. Res.*, Vol. 224, 1989, pp. 427-436.

Sekizawa, J., and T. Shibamoto. Genotoxicity of safrole-related chemicals in microbial test systems. *Mutat. Res.*, Vol. 101, 1982, pp. 127-140.

Stroberg, J., and F. Grundschober. Consumption ratio and food predominance of flavoring materials. *Perfum Flavor*, Vol. 12, 1987, pp. 27.

WHO. World Health Organization. Evaluation of Certain Food Additives and Contaminants. Twenty-Sixth Report of the Joint FAO/WHO Expert Committee on Food Additives. Technical Report Series 669. Geneva, Switzerland: WHO, 1981, pp. 92-94.