


1,2,3-TRICHLOROPROPANE
CAS No. 96-18-4
First Listed in the *Eighth Report on Carcinogens*



CARCINOGENICITY

1,2,3-Trichloropropane (TCP) is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of malignant tumor formation at multiple sites in multiple species of experimental animals (NTP 1993, Irwin *et al.* 1995).

TCP administered by gavage for 2 years induced multiple tumors in mice and rats. Increased incidences of tumors of the forestomach, liver, and harderian gland were observed in both male and female mice, and increased incidences of tumors of the uterus and oral mucosa were observed in female mice. In male rats, increased incidences of tumors were observed in the preputial gland, kidney, and pancreas. Female rats had increased incidences of tumors in the clitoral gland and mammary gland, and both male and female rats had increased incidences of tumors in the forestomach, oral mucosa, and Zymbal gland (NTP 1993, Irwin *et al.* 1995, IARC 1995).

No adequate data were available to evaluate the carcinogenicity of TCP in humans.

ADDITIONAL INFORMATION RELEVANT TO CARCINOGENESIS OR POSSIBLE MECHANISMS OF CARCINOGENESIS

TCP, when tested *in vitro*, induced gene mutations in bacteria, yeast, and mammalian cells; further, TCP induced sister chromatid exchanges, chromosomal aberrations, micronuclei, and morphological transformation in mammalian cells (Dean and Brooks 1979, Sawin and Hass 1982a,b, IARC 1995, Doherty *et al.* 1996). TCP was active almost exclusively in the presence, but not the absence, of metabolic activation or when tested using metabolically competent cells. In *in vivo* rodent studies, TCP induced DNA damage, including DNA adducts in multiple tissues of rats and mice receiving the chemical by gavage or by intraperitoneal injection (IARC 1995, La *et al.* 1995). TCP also induced cell proliferation in multiple tissues of rats and mice receiving the chemical by gavage or by inhalation (rats only) (Johannsen *et al.* 1988, NTP 1993, Irwin *et al.* 1995). TCP has been reported as negative for the induction of dominant lethal mutations in male rats (IARC 1995). Several metabolites of TCP, including 1,3-dichloroacetone, induce genetic damage in a variety of short-term test systems (IARC 1995). This metabolite is produced by human liver microsomes, although its rate of formation is less than in rats (Weber and Snipes 1992).

No data were available that would suggest that the mechanisms thought to account for tumor induction by TCP in experimental animals would not also operate in humans.

PROPERTIES

TCP is a clear, colorless to straw-colored liquid with a strong acrid odor similar to that of chloroform or trichloroethylene. It has a melting point of -14.7°C and a boiling point of 156.8°C . At normal temperatures, it evaporates at a rate similar to water. TCP is soluble in ethanol, ether, and chloroform, and it is slightly soluble in water. It dissolves several substances, such as oils, waxes, fats, chlorinated rubber, and numerous resins, and reacts with active metals, strong caustics, and oxidizers. It is sensitive to prolonged exposure to light and to heat. TCP is flammable, and when heated to decomposition, it yields highly toxic fumes of hydrogen chloride gas (IARC 1995, HSDB 2000, NTP 2001).

USE

In the past, TCP has been used primarily as a solvent and extractive agent. As a solvent, it has commonly been used as a paint and varnish remover, a cleaning and degreasing agent, and a cleaning and maintenance solvent. No current information is available to indicate that it continues to be used for these purposes. Currently, TCP is used as a chemical intermediate in the production of polysulfone liquid polymers and dichloropropene, synthesis of hexafluoropropylene, and as a crosslinking agent in the synthesis of polysulfides. No data were available to indicate to what extent TCP is currently used for these purposes (ATSDR 1992). TCP has been formulated with dichloropropenes in the manufacture of a soil fumigant D-D (IARC 1995). According to the Farm Chemicals Handbook 1991 (Sine 1991), this soil fumigant is no longer available in the United States.

PRODUCTION

No current production information is available for TCP; however, the estimated production in 1977 ranged from 21 to 110 million lb (ATSDR 1992). In 1985, two manufacturing facilities had a combined annual production greater than 10,000 pounds (NTP 1993). There are currently two U.S. facilities that produce TCP and nine suppliers (ATSDR 1992, HSDB 2000, Chem Sources 2001). No data were available on imports or exports of TCP.

TCP may also be produced in significant quantities as a by-product of the production of other chlorinated compounds such as dichloropropene, propylene chlorohydrin, dichlorohydrin, and glycerol (ATSDR 1992). In addition, TCP is formed as a by-product of epichlorohydrin production (IFIS 1985).

EXPOSURE

The primary routes of potential human exposure to TCP are inhalation of vapors, dermal contact, or ingestion of contaminated water. TCP is not a naturally occurring chemical. Releases to the environment are likely to occur as a result of its manufacture, formulation, and use as a solvent and extractive agent, paint and varnish remover, cleaning and degreasing agent, cleaning and maintenance reagent, and chemical intermediate. TCP has been detected in low concentrations in surface, drinking, and ground waters in various parts of the United States (ATSDR 1992, NTP 1993).

The National Occupational Exposure Survey (1981-1983) indicated that 492 workers, including 9 women, were potentially exposed to TCP (NIOSH 1984). This survey did not contain information on the frequency, concentration, or duration of exposure. The National Occupational Hazard Survey, conducted by NIOSH from 1972 to 1974, estimated that 490 workers were potentially exposed to TCP in the US (NIOSH 1976). TCP is manufactured in closed systems; therefore, occupational exposures are more likely to occur at facilities where the chemical is used rather than at facilities where it is produced. Direct handling of TCP or products containing TCP may occur during purification, formulation of products, sampling and quality control, packaging and storage, leakage of equipment, startup and shutdown procedures, maintenance, cleanup, spills, and other facility emergencies (ATSDR 1992).

In the past, inhalation and dermal exposures likely occurred during the use of consumer products, such as certain paint removers, that contained TCP; however, TCP is no longer used in consumer products. The general population may be exposed to low levels of TCP by ingestion of contaminated well water or by inhalation of contaminated air. Individuals that live near facilities that use or produce the compound or hazardous waste disposal facilities are more likely to be exposed (ATSDR 1992). EPA's Toxic Chemical Release Inventory (TRI99 2001) contains data on environmental releases of TCP from 1995 to 1999. Seven facilities reported releasing 28,860 lb of TCP to the environment in 1999. Total air emissions and releases to land contributed 48% and 28%, respectively, to the total release estimates. Approximately 44% of the total was reported by one waste disposal facility. Historical releases ranged from a low of 2,091 lb in 1996 to a high of 20,895 lb in 1998.

REGULATIONS

TCP is regulated by EPA under the Clean Air Act (CAA), Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA), Resource Conservation and Recovery Act (RCRA), and Safe Drinking Water Act (SDWA). Under CAA, the Synthetic Organic Chemicals Manufacturing Industry is required to monitor equipment leaks of TCP and to develop standards of performance. Under the SDWA, EPA requires that community and non-community water systems monitor for TCP. Under SARA, EPA requires the annual reporting of releases and waste management activities for TCP in amounts exceeding the threshold quantity. TCP is listed as a hazardous constituent of waste under RCRA and is subject to minimum national detection criteria in groundwater from municipal solid waste sites.

The American Conference of Governmental Industrial Hygienists (ACGIH) recommends a threshold limit value (TLV) of 10 ppm (60 mg/m³) for TCP. NIOSH recommends that 1,2,3-trichloropropane be regarded as a potential occupational carcinogen with an exposure level for to 10 ppm as a 10-hr time weighted average (TWA) with the potential for skin absorption noted. OSHA has set a permissible exposure limit (PEL) of 50 ppm (300 mg/m³) as an 8-hour TWA for TCP. OSHA also regulates TCP under the Hazard Communication Standard and as a chemical hazard in laboratories. Regulations are summarized in Volume II, Table 182.

REFERENCES

ATSDR. Agency for Toxic Substances and Disease Registry. Toxicological Profile for 1,2,3-Trichloropropane. (Final Report). Atlanta, GA: ATSDR, Public Health Service, U.S. Department of Health and Human Services. 1992. 117 pp. NTIS Accession No. PB93-110872.

Chem Sources. Chemical Sources International, Inc. <http://www.chemsources.com>, 2001.

Dean, B.J., and T.M. Brooks. In vitro Mutation Studies with 1,2,3-Trichloropropane. [TSCA Section 8(e) submission]. U.S. EPA/OTS Public Files. Record Number: 309026. 1979. Microfiche Number: 0515727.

Doherty, A.T., S. Ellard, E.M. Parry, and J.M. Parry. An Investigation into the Activation and Deactivation of Chlorinated Hydrocarbons to Genotoxins in Metabolically Competent Human Cells. *Mutagenesis* Vol. 11, No. 3, 1996, pp. 247-274.

HSDB. Hazardous Substances Data Bank. Online database produced by the National Library of Medicine. 1,2,3-Trichloropropane. Profile last updated June 12, 2000. Last review date, September 14, 1995.

IARC. International Agency for Research on Cancer. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Dry Cleaning, Some Chlorinated Solvents and Other Industrial Chemicals. Vol. 63. 558 pp. Lyon, France, IARC, 1995.

IFIS. Industry File Index System. EPA Database available on the Chemical Information System. EPA regulations on particular industries and substances. Last update in 1985.

Irwin, R.D., J.K. Haseman, and S.L. Eustis. 1,2,3-Trichloropropane: A Multisite Carcinogen in Rats and Mice. *Fund. Appl. Toxicol.* Vol. 25, 1995, pp. 241-252.

Johannsen, F.R., G.J. Levinskas, G.M. Rusch, J.B. Terrill, and R.E. Schroeder. Evaluation of the Subchronic and Reproductive Effects of a Series of Chlorinated Propanes in the Rat. I. Toxicity of 1,2,3-Trichloropropane. *J. Toxicol. Environ. Health* Vol. 25, 1988, pp. 299-315.

La, D.K., P.D. Lilly, R.J. Anderegg, and J.A. Swenberg. DNA Adduct Formation in B6C3F₁ Mice and Fischer-344 Rats Exposed to 1,2,3-Trichloropropane. *Carcinogenesis* Vol. 16, No. 6, 1995, pp. 1419-1424.

NIOSH. National Institute for Occupational Safety and Health. National Occupational Hazard Survey (1972-74). Cincinnati, OH: Department of Health, Education, and Welfare, 1976.

NIOSH. National Institute for Occupational Safety and Health. National Occupational Exposure Survey (1981-83). Cincinnati, OH: Department of Health and Human Services, 1984.

NTP. National Toxicology Program. Technical Report Series No. 384. Toxicology and Carcinogenesis Studies of 1,2,3-Trichloropropane in F344/N Rats and B6C3F₁ Mice (Gavage Studies). NIH Publication No. 94-2839. National Toxicology Program, Research Triangle Park, NC, 1993.

NTP. National Toxicology Program. NTP Chemical Repository. 1,2,3-Trichloropropane. Last updated August 13, 2001.

Sawin, V.L., and B.S. Hass. Assay of 1,2,3-Trichloropropane for Gene Mutation in Mouse Lymphoma Cells. [TSCA Sections 8(d) and 8(e)(c) submission]. U.S. EPA/OTS Public Files. 1982a. Record Number: 309014. Microfiche Number: 0515721a.

Sawin, V.L., and B.S. Hass. Initial Submission: Assay of 1,2,3-Trichloropropane for Gene Mutation in Mouse Lymphoma Cells (Final Report) with Cover Letter Dated 120391. [TSCA

1,2,3-Trichloropropane (Continued)

Sections 8(d) and 8(e)(p) submission]. U.S. EPA/OTS Public Files. 1982b. Record Number: 420874. Microfiche Number: 0534986b.

Sine, C., Ed. Farm Chemicals Handbook '91. Willoughby, OH: Meister Publishing Co., 1991.

TRI99. Toxic Chemicals Release Inventory 1999. Data contained in the Toxic Chemical Release Inventory (TRI). Available from the U.S. Environmental Protection Agency Office of Environmental Information, <http://www.epa.gov/triexplorer/reports.htm>, 2001.

Weber, G.L., and I.G. Sipes. In Vitro Metabolism and Bioactivation of 1,2,3-Trichloropropane. Toxicol. Appl. Pharmacol. Vol. 113, 1992, pp. 152-158.