

A & S Pharmaceutical
Exhibits

012 3051

Compare to the active
ingredient in Bayer®

**BERKLEY
JENSEN**
Aspirin

**Pain Reliever
Fever Reducer**

Use this aspirin for
cough, cold, flu, sore throat

500 Tablets

DO NOT USE IF THE PAPERED CASE
AROUND THE CAN IS BROKEN OR MISSING.

INDICATIONS: For the temporary relief of headache, pain and fever of acute and chronic infections, rheumatoid fever, rheumatic fever, and other acute and chronic infections.

PRECAUTIONS: Take 2 tablets every 4 hours as needed. Do not exceed 12 tablets in 24 hours unless directed by a doctor. Children under 12 years of age should not use.

ALCOHOL WARNING: If you consume 2 or more alcoholic drinks every day, ask your doctor whether you should take aspirin or other pain relievers. Aspirin may cause stomach bleeding.

WARNING: Do not use with benzoyl peroxide and the product is not for use on the face. Do not use before a dental or medical procedure. Do not use if you are allergic to aspirin or if you are taking any medicine that may interact with aspirin. Do not take this product for more than 10 days or for fever for more than 3 days.

CAUTION: Aspirin should be used with care in patients with a history of asthma, peptic ulcer disease, or if reduced or no stomach acid is present. Aspirin may cause stomach bleeding and the pain of a peptic ulcer. It may also cause stomach bleeding.

WARNING: Do not use if you are pregnant or nursing a child. Aspirin may cause a child to have a fever. It is especially important that you use aspirin during the child's illnesses or fevers. Aspirin may cause stomach bleeding.

WARNING: Do not use if you are taking any medicine that may interact with aspirin. Do not take this product for more than 10 days or for fever for more than 3 days.

WARNING: Do not use if you are pregnant or nursing a child. Aspirin may cause a child to have a fever. It is especially important that you use aspirin during the child's illnesses or fevers. Aspirin may cause stomach bleeding.

WARNING: Do not use if you are taking any medicine that may interact with aspirin. Do not take this product for more than 10 days or for fever for more than 3 days.

WARNING: Do not use if you are pregnant or nursing a child. Aspirin may cause a child to have a fever. It is especially important that you use aspirin during the child's illnesses or fevers. Aspirin may cause stomach bleeding.

WARNING: Do not use if you are taking any medicine that may interact with aspirin. Do not take this product for more than 10 days or for fever for more than 3 days.

WARNING: Do not use if you are pregnant or nursing a child. Aspirin may cause a child to have a fever. It is especially important that you use aspirin during the child's illnesses or fevers. Aspirin may cause stomach bleeding.



00005791000
JANUARY 1988 0530 10000000

Compare to the active
ingredient in Bayer®

**BERKLEY
JENSEN**

Aspirin

*Rain Reliever
Fever Reducer*

With this coating, the
caps melt in your mouth.

500 Tablets

DO NOT USE IF THE PRINTED SEAL
AROUND THE CAP IS BROKEN OR MISSING.



INDICATIONS: For the temporary relief of headache, pain and fever in cases of cold, flu, sore throat, toothache, neuralgia, rheumatism, and other acute and chronic ailments.

CAUTIONS: Aspirin should be taken only as directed. Do not exceed 10 tablets in 24 hours unless directed by a doctor. Avoid or under 12 years. Consult a doctor.

ALCOHOL WARNING: If you consume 3 or more alcoholic drinks every day, you may become addicted to alcohol. Aspirin may cause stomach bleeding.

WARNING: Doctors and patients should tell each other about all medicines being taken. One of the symptoms listed below is associated with aspirin. Do not take this product for pain for more than 10 days or for fever for more than 3 days unless directed by a doctor. It may cause stomach bleeding. If you experience any of the symptoms listed below, stop taking aspirin immediately and consult your doctor.

SYMPTOMS: Stomach pain, heartburn, indigestion, loss of appetite, dizziness, ringing in the ears, blurred vision, tinnitus, and other symptoms.

ADDITIONAL SYMPTOMS: If you experience any of the following symptoms, stop taking aspirin immediately and consult your doctor: Stomach pain, heartburn, indigestion, loss of appetite, dizziness, ringing in the ears, blurred vision, tinnitus, and other symptoms.

CONTRAINDICATIONS: Do not take aspirin if you are allergic to aspirin or any of the ingredients listed on the label. Do not take aspirin if you are taking any other medicine that contains aspirin.

PRECAUTIONS: Do not take aspirin if you are pregnant or breastfeeding. Do not take aspirin if you are taking any other medicine that contains aspirin.

ADDITIONAL PRECAUTIONS: Do not take aspirin if you are taking any other medicine that contains aspirin.

HOW TO TAKE: Take one tablet every 4 to 6 hours as needed. Do not take more than 10 tablets in 24 hours unless directed by a doctor.

HOW TO STORE: Store in a cool, dry place. Do not store in a plastic bag.

DISTRIBUTED BY

BAYER

4 Merritt Road

Whisper, N.J. 07078

1-800-454-1234

LAB 500 AW 22299-0000 Made in U.S.A.



FORMUCARE

Aspirin
TABLETS, 325 mg each

**PAIN RELIEVER/
FEVER REDUCER**

*Ultra thin coating
for easy swallowing*

DO NOT USE IF THE PRINTED SEAL
AROUND THE CAN IS BROKEN OR MISSING.

INDICATIONS: For the temporary relief of headache, neuralgia, fever, and minor aches and pains of various origin.

CAUTION: Adults: 2 tablets every 4 hours as needed. Do not exceed 12 tablets. Adult and child aspirin should not be used in children.

ADVERSE REACTIONS: 1. Some patients may experience dizziness, upset stomach, and heartburn. 2. Some patients may experience allergic reactions.

WARNINGS: 1. Aspirin should not be used in children. 2. Aspirin should not be used in patients with a history of peptic ulcer disease.

CONTRAINDICATIONS: 1. Aspirin should not be used in patients with a history of peptic ulcer disease.

PRECAUTIONS: 1. Aspirin should not be used in patients with a history of peptic ulcer disease.

HOW TO USE: 1. Aspirin should not be used in patients with a history of peptic ulcer disease.

HOW TO STORE: 1. Aspirin should not be used in patients with a history of peptic ulcer disease.

HOW TO OBTAIN: 1. Aspirin should not be used in patients with a history of peptic ulcer disease.

HOW TO ORDER: 1. Aspirin should not be used in patients with a history of peptic ulcer disease.



222448-0100 1-613710-74 T-5570

