References and Included Studies

1. Marsh MN. Gluten, major histocompatibility complex, and the small intestine: A molecular and immunobiologic approach to the spectrum of gluten sensitivity ('celiac sprue'). Gastroenterology 1992;102(1):330-54.

2. McNeish AS, Harms HK, Rey J, Shmerling DH, Visakorpi JK, Walker-Smith JA. The diagnosis of coeliac disease. A commentary on the current practices of members of the European Society for Paediatric Gastroenterology and Nutrition (ESPGAN). Archives of Disease in Childhood 1979;54(10):783-6.

3. Oberhuber G, Granditsch G, Vogelsang H. The histopathology of coeliac disease: time for a standardized report scheme for pathologists. European Journal of Gastroenterology & Hepatology 1999;11(10):1185-94.

4. Walker-Smith JA, Guandalini S, Schmitz J, Shmerling DH, Visakorpi JK. Revised criteria for diagnosis of coeliac disease. Arch Dis Child 1990;65(8):909-11.

5. van de WY, Kooy Y, van Veelen P, Vader W, Koning F, Pena S. Coeliac disease: it takes three to tango! Gut 2000;46(5):734-7.

6. Papadopoulos GK, Wijmenga C, Koning F. Interplay between genetics and the environment in the development of celiac disease: perspectives for a healthy life. Journal of Clinical Investigation 2001;108(9):1261-6.

7. Sollid LM, McAdam SN, Molberg O, Quarsten H, Arentz-Hansen H, Louka AS, et al. Genes and environment in celiac disease. Acta Odontologica Scandinavica 2001;59(3):183-6.

8. Sollid LM, Markussen G, Ek J, Gjerde H, Vartdal F, Thorsby E. Evidence for a primary association of celiac disease to a particular HLA-DQ alpha/beta heterodimer. Journal of Experimental Medicine 1989;169(1):345-50.

9. Ploski R, Ek J, Thorsby E, Sollid LM. On the HLA-DQ(alpha 1*0501, beta 1*0201)-associated susceptibility in celiac disease: a possible gene dosage effect of DQB1*0201. Tissue Antigens 1993;41(4):173-7.

10. Ploski R, Ascher H, Sollid LM. HLA genotypes and the increased incidence of coeliac disease in Sweden. Scand J Gastroenterol 1996;31(11):1092-7.

11. Holopainen P, Mustalahti K, Uimari P, Collin P, Maki M, Partanen J. Candidate gene regions and genetic heterogeneity in gluten sensitivity. Gut 2001;48(5):696-701.

12. Kagnoff MF. Celiac disease pathogenesis: the plot thickens. Gastroenterology 2002;123(3):939-43.

13. Feldman M, Friedman LS, Sleisenger MH. Sleisenger and Fordtran's Gastrointestinal and Liver Disease. 7th edition W.B. Saunders; 2003.

14. Fasano A, Catassi C. Current approaches to diagnosis and treatment of celiac disease: an evolving spectrum. Gastroenterology 2001;120(3):636-51.

15. Fasano A, Berti I, Gerarduzzi T, Not T, Colletti RB, Drago S, et al. Prevalence of celiac disease in at-risk and not-at-risk groups in the United States: a large multicenter study. Archives of Internal Medicine 2003;163(3):286-92.

16. Rostami K, Kerckhaert J, Tiemessen R, von Blomberg BM, Meijer JW, Mulder CJ. Sensitivity of antiendomysium and antigliadin antibodies in untreated celiac disease: disappointing in clinical practice. American Journal of Gastroenterology 1999;94(4):888-94.

17. Rosenthal DS, Roop DR, Huff CA, Weiss JS, Ellis CN, Hamilton T, et al. Changes in photo-aged human skin following topical application of all-trans retinoic acid. Journal of Investigative Dermatology 1990;95(5):510-5.

18. Wahab PJ, Meijer JWR, Goerres MS, Mulder CJJ. Coeliac disease: changing views on gluten-sensitive enteropathy. Scandinavian Journal of Gastroenterology Supplement 2002;(236):60-5.

19. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. BMC Med Res Methodol 2003;3(1):25. Available: PM:14606960.

20. Wells GA, Shea B, O'Connell D, Peterson J,Welch V, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. 3rd Symposium on Systematic Reviews: Beyond the Basics, July 2000 in Oxford.2000.

21. Ophthalmology Study Design Worksheet #3 Noncomparative (nonrandomized, noncontrolled) Interventional Case Series. 2004. Ref Type: Internet Communication

22. Moses LE, Shapiro D, Littenberg B. Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. Stat Med 1993;12(14):1293-316. Available: PM:8210827.

23. Littenberg B, Moses LE. Estimating diagnostic accuracy from multiple conflicting reports: a new meta-analytic method. Med Decis Making 1993;13(4):313-21. Available: PM:8246704.

24. Deeks J. Systematics reviews of evaluations of diagnostic and screening test. Systematic Reviews in Health Care: Meta-analysis in Context. London: BMJ Books, 2001.

25. Deeks J, Altman D, Bradburn J. Statistical methods for examining heterogeneity and combining results from several studies in meta-analysis. Systematic Review in Health Care. Meta-Analysis in Context London: BMJ Books; 2001.

26. Altuntas B, Kansu A, Ensari A, Girgin N. Celiac disease in Turkish short-statured children and the value of antigliadin antibody in diagnosis. Acta Paediatr Jpn 1998;40(5):457-60.

27. Ascher H, Hahn-Zoric M, Hanson LA, Kilander AF, Nilsson LA, Tlaskalova H. Value of serologic markers for clinical diagnosis and population studies of coeliac disease. Scandinavian Journal of Gastroenterology 1996;31(1):61-7.

28. Ascher H, Lanner A, Kristiansson B. A new laboratory kit for anti-gliadin IgA at diagnosis and follow-up of childhood celiac disease. Journal of Pediatric Gastroenterology and Nutrition 1990;10(4):443-50.

29. Bahia M, Rabello A, Brasileiro FG, Penna FJ. Serum antigliadin antibody levels as a screening criterion before jejunal biopsy indication for celiac disease in a developing country. Brazilian Journal of Medical and Biological Research = Revista Brasileira De Pesquisas Medicas E Biologicas / Sociedade Brasileira De Biofisica Et Al 2001;34(11):1415-20.

30. Bardella MT, Trovato C, Cesana BM, Pagliari C, Gebbia C, Peracchi M. Serological markers for coeliac disease: is it time to change? Digestive and Liver Disease - Official Journal of the Italian Society of Gastroenterology and the Italian Association for the Study of the Liver 2001;33(5):426-31.

31. Berger R, Schmidt G. Evaluation of six anti-gliadin antibody assays. Journal of Immunological Methods 1996;191(1):77-86.

32. Biagi F, Pezzimenti D, Campanella J, Vadacca GB, Corazza GR. Endomysial and tissue transglutaminase antibodies in coeliac sera: a comparison not influenced by previous serological testing. Scandinavian Journal of Gastroenterology 2001;36(9):955-8.

33. Bode S, Gudmand-Hoyer E. Evaluation of the gliadin antibody test for diagnosing coeliac disease. Scandinavian Journal of Gastroenterology 1994;29(2):148-52.

34. Bode S, Weile B, Krasilnikoff PA, Gudmand-Hoyer E. The diagnostic value of the gliadin antibody test in celiac disease in children: a prospective study. Journal of Pediatric Gastroenterology and Nutrition 1993;17(3):260-4.

35. Bonamico M, Tiberti C, Picarelli A, Mariani P, Rossi D, Cipolletta E, et al. Radioimmunoassay to detect antitransglutaminase autoantibodies is the most sensitive and specific screening method for celiac disease. American Journal of Gastroenterology 2001;96(5):1536-40.

36. Bottaro G, Volta U, Spina M, Rotolo N, Sciacca A, Musumeci S. Antibody pattern in childhood celiac disease. Journal of Pediatric Gastroenterology and Nutrition 1997;24(5):559-62.

37. Carroccio A, Iacono G, D'Amico D, Cavataio F, Teresi S, Caruso C, et al. Production of anti-endomysial antibodies in cultured duodenal mucosa: usefulness in coeliac disease diagnosis. Scandinavian Journal of Gastroenterology 2002;37(1):32-8.

38. Carroccio A, Iacono G, Montalto G, Cavataio F, Soresi M, Kazmierska I, et al. Immunologic and absorptive tests in celiac disease: can they replace intestinal biopsies? Scandinavian Journal of Gastroenterology 1993;28(8):673-6.

39. Carroccio A, Vitale G, Di Prima L, Chifari N, Napoli S, La Russa C, et al. Comparison of anti-transglutaminase ELISAs and an anti-endomysial antibody assay in the diagnosis of celiac disease:

a prospective study. Clinical Chemistry 2002;48(9):1546-50.

40. Cataldo F, Lio D, Marino V, Picarelli A, Ventura A, Corazza GR. IgG(1) antiendomysium and IgG antitissue transglutaminase (anti-tTG) antibodies in coeliac patients with selective IgA deficiency. Working Groups on Celiac Disease of SIGEP and Club del Tenue. Gut 2000;47(3):366-9.

41. Chan AW, Butzner JD, McKenna R, Fritzler MJ. Tissue transglutaminase enzyme-linked immunosorbent assay as a screening test for celiac disease in pediatric patients. Pediatrics 2001;107(1):E8.

42. Chartrand LJ, Agulnik J, Vanounou T, Russo PA, Baehler P, Seidman EG. Effectiveness of antigliadin antibodies as a screening test for celiac disease in children. Cmaj - Canadian Medical Association Journal = Journal De L'association Medicale Canadienne 1997;157(5):527-33.

43. Chirdo FG, Rumbo M, Carabajal P, Castagnino N, Mavromatopulos E, Cirincione V, et al. Analysis of anti-gliadin antibodies by immunoblot analysis and enzyme-linked immunosorbent assay using gliadin fractions as antigens. Journal of Pediatric Gastroenterology and Nutrition 1999;29(2):171-7.

44. Chirdo FG, Rumbo M, Carabajal P, Mavromatopulos E, Castagnino N, Anon MC, et al. Determination of anti-omega-gliadin antibodies in serologic tests for coeliac disease. Scandinavian Journal of Gastroenterology 2000;35(5):508-16.

45. Dahele A, Aldhous MC, Humphreys K, Ghosh S. Serum IgA tissue transglutaminase antibodies in coeliac disease and other gastrointestinal diseases. Q J Med 2001;94(4):195-205.

46. Di Leo M, Weisz G, Ansaldi BN. Serum and salivary antiendomysium antibodies in the screening of coeliac disease. Panminerva Medica 1999;41(1):68-71.

47. Dickey W, McMillan SA, Hughes DF. Sensitivity of serum tissue transglutaminase antibodies for endomysial antibody positive and negative coeliac disease. Scandinavian Journal of Gastroenterology 2001;36(5):511-4.

48. Falth-Magnusson K, Jansson G, Stenhammar L, Magnusson KE. Serum food antibodies analyzed by enzyme-linked immunosorbent assay (ELISA) and diffusion-in-gel (DIG)-ELISA methods in children with and without celiac disease. Journal of Pediatric Gastroenterology and Nutrition 1994;18(1):56-62.

49. Gillett HR, Freeman HJ. Comparison of IgA endomysium antibody and IgA tissue transglutaminase antibody in celiac disease. Canadian Journal of Gastroenterology = Journal Canadien De Gastroenterologie 2000;14(8):668-71.

50. Gonczi J, Skerritt JH, Mitchell JD. A reliable screening test for coeliac disease: enzyme-linked immunosorbent assay to detect anti-gliadin antibodies in serum. Australian and New Zealand Journal of Medicine 1991;21(5):723-31.

51. Hallstrom O. Comparison of IgA-class reticulin and endomysium antibodies in coeliac disease and dermatitis herpetiformis. Gut 1989;30(9):1225-32.

52. Hansson T, Dahlbom I, Hall J, Holtz A, Elfman L, Dannaeus A, et al. Antibody reactivity against human and guinea pig tissue transglutaminase in children with celiac disease. Journal of Pediatric Gastroenterology and Nutrition 2000;30(4):379-84.

53. Iltanen S, Rantala I, Laippala P, Holm K, Partanen J, Maki M. Expression of HSP-65 in jejunal epithelial cells in patients clinically suspected of coeliac disease. Autoimmunity 1999;31(2):125-32.

54. Kaukinen K, Turjanmaa K, Maki M, Partanen J, Venalainen R, Reunala T, et al. Intolerance to cereals is not specific for coeliac disease. Scandinavian Journal of Gastroenterology 2000;35(9):942-6.

55. Kolho KL, Savilahti E. IgA endomysium antibodies on human umbilical cord: an excellent diagnostic tool for celiac disease in childhood. Journal of Pediatric Gastroenterology and Nutrition 1997;24(5):563-7.

56. Kumar V, Lerner A, Valeski JE, Beutner EH, Chorzelski TP, Rossi T. Endomysial antibodies in the diagnosis of celiac disease and the effect of gluten on antibody titers. Immunological Investigations 1989;18(1-4):533-44.

57. Ladinser B, Rossipal E, Pittschieler K. Endomysium antibodies in coeliac disease: an improved method. Gut 1994;35(6):776-8.

58. Lerner A, Kumar V, Iancu TC. Immunological diagnosis of childhood coeliac disease: comparison between antigliadin, antireticulin and antiendomysial antibodies. Clinical and Experimental Immunology 1994;95(1):78-82.

59. Lindberg T, Nilsson LA, Borulf S, Cavell B, Fallstrom SP, Jansson U, et al. Serum IgA and IgG gliadin antibodies and small intestinal mucosal damage in children. Journal of Pediatric Gastroenterology and Nutrition 1985;4(6):917-22.

60. Lindquist BL, Rogozinski T, Moi H, Danielsson D, Olcen P. Endomysium and gliadin IgA antibodies in children with coeliac disease. Scandinavian Journal of Gastroenterology 1994;29(5):452-6.

61. Lock RJ, Pitcher MC, Unsworth DJ. IgA anti-tissue transglutaminase as a diagnostic marker of gluten sensitive enteropathy. Journal of Clinical Pathology 1999;52(4):274-7.

62. Maki M, Holm K, Lipsanen V, Hallstrom O, Viander M, Collin P, et al. Serological markers and HLA genes among healthy first-degree relatives of patients with coeliac disease. Lancet 1991;338(8779):1350-3.

63. McMillan SA, Haughton DJ, Biggart JD, Edgar JD, Porter KG, McNeill TA. Predictive value for coeliac disease of antibodies to gliadin, endomysium, and jejunum in patients attending for jejunal biopsy. Bmj (Clinical Research Ed) 1991;303(6811):1163-5.

64. Meini A, Pillan NM, Villanacci V, Monafo V, Ugazio AG, Plebani A. Prevalence and diagnosis of celiac disease in IgA-deficient children. Annals of Allergy, Asthma & Immunology - Official Publication of the American College of Allergy, Asthma, & Immunology 1996;77(4):333-6.

65. Pacht A, Sinai N, Hornstein L, Kumar V, Ish-Shalom N, Lerner A. The diagnostic reliability of anti-endomysial antibody in celiac disease: the north Israel experience. Israel Journal of Medical Sciences 1995;31(4):218-20.

66. Picarelli A, Sabbatella L, Di Tola M, Gabrielli F, Greco R, Di Cello T, et al. Celiac disease diagnosis in misdiagnosed children. Pediatric Research 2000;48(5):590-2.

67. Poddar U, Thapa BR, Nain CK, Prasad A, Singh K. Celiac disease in India: are they true cases of celiac disease? Journal of Pediatric Gastroenterology and Nutrition 2002;35(4):508-12.

68. Rich EJ, Christie DL. Anti-gliadin antibody panel and xylose absorption test in screening for celiac disease. Journal of Pediatric Gastroenterology and Nutrition 1990;10(2):174-8.

69. Russo PA, Chartrand LJ, Seidman E. Comparative analysis of serologic screening tests for the initial diagnosis of celiac disease. Pediatrics 1999;104(1 Pt 1):75-8.

70. Salmaso C, Ocmant A, Pesce G, Altrinetti V, Montagna P, Descalzi D, et al. Comparison of ELISA for tissue transglutaminase autoantibodies with antiendomysium antibodies in pediatric and adult patients with celiac disease. Allergy 2001;56(6):544-7.

71. Sategna-Guidetti C, Grosso S, Bruno M, Grosso SB. Comparison of serum anti-gliadin, anti-endomysium, and anti-jejunum antibodies in adult celiac sprue. Journal of Clinical Gastroenterology 1995;20(1):17-21.

72. Sblattero D, Berti I, Trevisiol C, Marzari R, Tommasini A, Bradbury A, et al. Human recombinant tissue transglutaminase ELISA: an innovative diagnostic assay for celiac disease. American Journal of Gastroenterology 2000;95(5):1253-7.

73. Sulkanen S, Collin P, Laurila K, Maki M. IgA- and IgG-class antihuman umbilical cord antibody tests in adult coeliac disease. Scandinavian Journal of Gastroenterology 1998;33(3):251-4.

74. Sulkanen S, Halttunen T, Laurila K, Kolho KL, Korponay-Szabo IR, Sarnesto A, et al. Tissue transglutaminase autoantibody enzyme-linked immunosorbent assay in detecting celiac disease. Gastroenterology 1998;115(6):1322-8.

75. Tesei N, Sugai E, Vazquez H, Smecuol E, Niveloni S, Mazure R, et al. Antibodies to human recombinant tissue transglutaminase may detect coeliac disease patients undiagnosed by endomysial antibodies. Alimentary Pharmacology & Therapeutics 2003;17(11):1415-23.

76. Troncone R, Maurano F, Rossi M, Micillo M, Greco L, Auricchio R, et al. IgA antibodies to tissue transglutaminase: An effective diagnostic test for celiac disease. Journal of Pediatrics 1999;134(2):166-71.

77. Valdimarsson T, Franzen L, Grodzinsky E, Skogh T, Strom M. Is small bowel biopsy necessary in adults with suspected celiac disease and IgA anti-endomysium antibodies? 100% positive predictive value for celiac disease in adults. Digestive Diseases and Sciences 1996;41(1):83-7.

78. Valentini RA, Andreani ML, Corazza GR, Gasbarrini G. IgA endomysium antibody: a valuable tool in the screening of coeliac disease

but not its follow-up. Italian Journal of Gastroenterology 1994;26(6):279-82.

79. Vitoria JC, Arrieta A, Ortiz L, Ayesta A. Antibodies to human tissue transglutaminase for the diagnosis of celiac disease. Journal of Pediatric Gastroenterology and Nutrition 2001;33(3):349-50.

80. Vogelsang H, Genser D, Wyatt J, Lochs H, Ferenci P, Granditsch G, et al. Screening for celiac disease: a prospective study on the value of noninvasive tests. American Journal of Gastroenterology 1995;90(3):394-8.

81. Volta U, Molinaro N, De Franceschi L, Fratangelo D, Bianchi FB. IgA anti-endomysial antibodies on human umbilical cord tissue for celiac disease screening. Save both money and monkeys. Digestive Diseases and Sciences 1995;40(9):1902-5.

82. Whelan A, Willoughby R, Weir D. Human umbilical vein endothelial cells: a new easily available source of endomysial antigens. European Journal of Gastroenterology & Hepatology 1996;8(10):961-6.

83. Wolters V, Vooijs-Moulaert A, Burger H, Brooimans R, De Schryver J, Rijkers G, et al. Human tissue transglutaminase enzyme linked immunosorbent assay outperforms both the guinea pig based tissue transglutaminase assay and anti-endomysium antibodies when screening for coeliac disease. European Journal of Pediatrics 2002;161(5):284-7.

84. Di Leo M, Weisz G, Ansaldi BN. Serum and salivary antiendomysium antibodies in the screening of coeliac disease. Panminerva Medica 1999;41(1):68-71.

85. Artan R. Antigliadin antibody measurement as a screening test for childhood coeliac disease. Int Med J 1998;5(3):209-12.

86. Dahele A, Kingstone K, Bode J, Anderson D, Ghosh S. Anti-endomysial antibody negative celiac disease: does additional serological testing help? Digestive Diseases and Sciences 2001;46(1):214-21.

87. Brandborg LL, Goldberg SB, Breidenbach WC. Human coccidiosis--a possible cause of malabsorption. New England Journal of Medicine 1970;283(24):1306-13.

88. Hazama H, Omagari K, Masuda J, Ohba K, Kinoshita H, Matsuo I, et al. Automated enzymatic mitochondrial antibody assay for the diagnosis of primary biliary cirrhosis: applications of a routine diagnostic tool for the detection of antimitochondrial antibodies. Journal of Gastroenterology and Hepatology 2002;17(3):316-23.

89. Cheli R, Giacosa A. Inflammatory cell count and identification in specific duodenitis. (Celiac disease, Whipple's disease and Crohn's disease). Comparison with jejunal findings. Endoscopy 1977;9(3):147-51.

90. Talbot HS. A report on sexual function in paraplegics. J Urol 1949;61:265-70.

91. Fernandez-Arquero M, Polanco I, Escobar H, Figueredo MA, de la Concha EG, Clerici-Larradet N, et al. HLA-DQ alleles and susceptibility to celiac disease in Spanish children. Tissue Antigens 1995;45(2):145-7.

92. Palavecino EA, Mota AH, Awad J, DeRosa S, Herrera M, Chertkoff L, et al. HLA and celiac disease in Argentina: involvement of the DQ subregion. Disease Markers 1990;8(1):5-10.

93. Spurkland A, Ingvarsson G, Falk ES, Knutsen I, Sollid LM, Thorsby E. Dermatitis herpetiformis and celiac disease are both primarily associated with the HLA-DQ (alpha 1*0501, beta 1*02) or the HLA-DQ (alpha 1*03, beta 1*0302) heterodimers. Tissue Antigens 1997;49(1):29-34.

94. Hall MA, Lanchbury JS, Lee JS, Welsh K, I, Ciclitira PJ. HLA-DQ2 second-domain polymorphisms may explain increased trans-associated risk in celiac disease and dermatitis herpetiformis. Human Immunology 1993;38(4):284-92.

95. Spurkland A, Sollid LM, Ronningen KS, Bosnes V, Ek J, Vartdal F, et al. Susceptibility to develop celiac disease is primarily associated with HLA-DQ alleles. Human Immunology 1990;29(3):157-65.

96. Bao F, Yu L, Babu S, Wang T, Hoffenberg EJ, Rewers M, et al. One third of HLA DQ2 homozygous patients with type 1 diabetes express celiac disease-associated transglutaminase autoantibodies. Journal of Autoimmunity 1999;13(1):143-8.

97. Martin-Villa JM, Lopez-Suarez JC, Perez-Blas M, Martinez-Laso J, Ferre-Lopez S, Garcia-Torre C, et al. Coeliac- and enteropathy-associated autoantibodies in Spanish insulin-dependent diabetes mellitus patients and their relation to HLA antigens. Journal of Diabetes and Its Complications 2001;15(1):38-43.

98. Kaukinen K, Partanen J, Maki M, Collin P. HLA-DQ typing in the diagnosis of celiac disease. American Journal of Gastroenterology 2002;97(3):695-9.

99. Saukkonen T, Ilonen J, Akerblom HK, Savilahti E. Prevalence of coeliac disease in siblings of patients with Type I diabetes is related to the prevalence of DQB1*02 allele. Diabetologia 2001;44(8):1051-3.

100. Congia M, Frau F, Lampis R, Frau R, Mele R, Cucca F, et al. A high frequency of the A30, B18, DR3, DRw52, DQw2 extended haplotype in Sardinian celiac disease patients: further evidence that disease susceptibility is conferred by DQ A1*0501, B1*0201. Tissue Antigens 1992;39(2):78-83.

101. Herrera M, Theiler G, Augustovski F, Chertkoff L, Fainboim L, DeRosa S, et al. Molecular characterization of HLA class II genes in celiac disease patients of Latin American Caucasian origin. Tissue Antigens 1994;43(2):83-7.

102. Petronzelli F, Ferrante P, Triglione P, Bonamico M, Mazzilli MC. Oligotyping of celiac multiplex families with the 11th International Histocompatibility Workshop reagents. Tissue Antigens 1991;38(5):238-9.

103. Mantovani V, Corazza GR, Angelini G, Delfino L, Frisoni M, Mirri P, et al. Molecular analysis of HLA-DQ A alleles in coeliac disease lack of a unique disease-associated sequence. Clinical and Experimental Immunology 1991;83(1):74-8.

104. Agardh D, Nilsson A, Carlsson A, Kockum I, Lernmark A, Ivarsson SA. Tissue transglutaminase autoantibodies and human leucocyte antigen in Down Syndrome patients with coeliac disease. Acta Paediatrica (Oslo, Norway - 1992) 2002;91(1):34-8.

105. Bonamico M, Mazzilli MC, Morellini M, Vania A, Carpino F, Nicotra MR, et al. Expression of class II MHC antigens in the intestinal epithelium of pediatric celiac disease. Journal of Pediatric Gastroenterology and Nutrition 1989;9(3):269-75.

106. Bonamico M, Morellini M, Mariani P, Triglione P, Trabace S, Lulli P, et al. HLA antigens and antigliadin antibodies in coeliac disease. Dis Markers 1991;9(6):313-7.

107. Brett PM, Yiannakou JY, Morris MA, Vaughan R, Curtis D, Ciclitira PJ. Common HLA alleles, rather than rare mutants, confer susceptibility to coeliac disease. Annals of Human Genetics 1999;63(Pt 3):217-25.

108. Collin P, Syrjanen J, Partanen J, Pasternack A, Kaukinen K, Mustonen J. Celiac disease and HLA DQ in patients with IgA nephropathy. American Journal of Gastroenterology 2002;97(10):2572-6.

109. Csizmadia CG, Mearin ML, Oren A, Kromhout A, Crusius JB, von Blomberg BM, et al. Accuracy and cost-effectiveness of a new strategy to screen for celiac disease in children with Down syndrome. Journal of Pediatrics 2000;137(6):756-61.

110. Falchuk ZM, Katz AJ, Shwachman H, Rogentine GN, Strober W. Gluten-sensitive enteropathy: genetic analysis and organ culture study in 35 families. Scandinavian Journal of Gastroenterology 1978;13(7):839-43.

111. Fedrick JA, Pandey JP, Verkasalo M, Teppo AM, Fudenberg HH. Immunoglobulin allotypes and the immune response to wheat gliadin in a Finnish population with celiac disease. Experimental and Clinical Immunogenetics 1985;2(4):185-90.

112. Hansson T, Anneren G, Sjoberg O, Klareskog L, Dannaeus A. Celiac disease in relation to immunologic serum markers, trace elements, and HLA-DR and DQ antigens in Swedish children with Down syndrome. J Pediatr Gastroenterol Nutr 1999;29(3):286-92.

113. Lampasona V, Bonfanti R, Bazzigaluppi E, Venerando A, Chiumello G, Bosi E, et al. Antibodies to tissue transglutaminase C in type I diabetes. Diabetologia 1999;42(10):1195-8.

114. Lorini R, Scotta MS, Cortona L, Avanzini MA, Vitali L, De Giacomo C, et al. Celiac disease and type I (insulin-dependent) diabetes mellitus in childhood: follow-up study. Journal of Diabetes and Its Complications 1996;10(3):154-9.

115. Lundin KE, Sollid LM, Anthonsen D, Noren O, Molberg O, Thorsby E, et al. Heterogeneous reactivity patterns of HLA-DQ-restricted, small intestinal T-cell clones from patients with celiac disease. Gastroenterology 1997;112(3):752-9.

116. Mearin ML, Koninckx CR, Biemond I, Polanco I, Pena AS. Influence of genetic factors on the serum levels of antigliadin antibodies in celiac disease. Journal of Pediatric Gastroenterology and Nutrition 1984;3(3):373-7.

117. Silva EM, Fernandes M, I, Galvao LC, Sawamura R, Donadi EA. Human leukocyte antigen class II alleles in white Brazilian patients with celiac disease. Journal of Pediatric

Gastroenterology and Nutrition 2000;31(4):391-4.

118. Stern M. Comparative evaluation of serologic tests for celiac disease: a European initiative toward standardization. Journal of Pediatric Gastroenterology and Nutrition 2000;31(5):513-9.

119. Bevan S, Popat S, Braegger CP, Busch A, O'Donoghue D, Falth-Magnusson K, et al. Contribution of the MHC region to the familial risk of coeliac disease. Journal of Medical Genetics 1999;36(9):687-90.

120. Catassi C, Doloretta MM, Ratsch IM, De Virgiliis S, Cucca F. The distribution of DQ genes in the Saharawi population provides only a partial explanation for the high celiac disease prevalence. Tissue Antigens 2001;58(6):402-6.

121. Clot F, Gianfrani C, Babron MC, Bouguerra F, Southwood S, Kagnoff MF, et al. HLA-DR53 molecules are associated with susceptibility to celiac disease and selectively bind gliadin-derived peptides. Immunogenetics 1999;49(9):800-7.

122. Holm K, Savilahti E, Koskimies S, Lipsanen V, Maki M. Immunohistochemical changes in the jejunum in first degree relatives of patients with coeliac disease and the coeliac disease marker DQ genes. HLA class II antigen expression, interleukin-2 receptor positive cells and dividing crypt cells. Gut 1994;35(1):55-60.

123. Karell K, Holopainen P, Mustalahti K, Collin P, Maki M, Partanen J. Not all HLA DR3 DQ2 haplotypes confer equal susceptibility to coeliac disease: transmission analysis in families. Scand J Gastroenterol 2002;37(1):56-61.

124. Meddeb-Garnaoui A, Zeliszewski D, Mougenot JF, Djilali-Saiah I, Caillat-Zucman S, Dormoy A, et al. Reevaluation of the relative risk for susceptibility to celiac disease of HLA-DRB1, -DQA1, -DQB1, -DPB1, and -TAP2 alleles in a French population. Human Immunology 1995;43(3):190-9.

125. Ruiz MY, Olivares JL. Three-loci HLA haplotypes in Spanish celiac children and healthy subjects: estimation of linkage disequilibrium and haplotype frequencies. Am J Gastroenterol 2001;96(5):1455-9.

126. Volta U, Bellentani S, Bianchi FB, Brandi G, De Franceschi L, Miglioli L, et al. High prevalence of celiac disease in Italian general population. Dig Dis Sci 2001;46(7):1500-5.

127. O'Driscoll BR, Stevens FM, O'Gorman TA, Finnegan P, McWeeney JJ, Little MP, et al. HLA type of patients with coeliac disease and malignancy in the west of Ireland. Gut 1982;23(8):662-5.

128. Hoffenberg EJ, MacKenzie T, Barriga KJ, Eisenbarth GS, Bao F, Haas JE, et al. A prospective study of the incidence of childhood celiac disease. J Pediatr 2003;143(3):308-14.

129. Holm KH. Correlation of HLA-DR alleles to jejunal mucosal morphology in healthy first-degree relatives of coeliac disease patients. Eur J Gastroenterol Hepatol 1993;5(1):35-9.

130. Louka AS, Moodie SJ, Karell K, Bolognesi E, Ascher H, Greco L, et al. A collaborative European search for non-DQA1 *05-DQB1 *02 Celiac disease loci on HLA-Dr3 haplotypes: Analysis of transmission from homozygous parents. Hum Immunol 2003;64(3):350-8.

131. Agrawal S, Gupta A, Yachha SK, Muller-Myhsok B, Mehrotra P, Agarwal SS. Association of human leucocyte-DR and DQ antigens in coeliac disease: a family study. Journal of Gastroenterology and Hepatology 2000;15(7):771-4.

132. Louka AS, Lie BA, Talseth B, Ascher H, Ek J, Gudjonsdottir AH, et al. Coeliac disease patients carry conserved HLA-DR3-DQ2 haplotypes revealed by association of TNF alleles. Immunogenetics 2003;55(5):339-43.

133. Van Belzen MJ, Meijer Jos WR, Sandkuijl LA, Bardoel Alfons FJ, Mulder Chris JJ, Pearson PL, et al. A major non-HLA locus in celiac disease maps to chromosome 19. Gastroenterology 2003;125(4):1032-41.

134. Castro M, Crino A, Papadatou B, Purpura M, Giannotti A, Ferretti F, et al. Down Syndrome and celiac disease: the prevalence of high IgA-antigliadin antibodies and HLA-DR and DQ antigens in trisomy 21. Journal of Pediatric Gastroenterology and Nutrition 1993;16(3):265-8.

135. Fine KD, Do K, Schulte K, Ogunji F, Guerra R, Osowski L, et al. High prevalence of celiac sprue-like HLA-DQ genes and enteropathy in patients with the microscopic colitis syndrome. American Journal of Gastroenterology 2000;95(8):1974-82.

136. Iltanen S, Holm K, Partanen J, Laippala P, Maki M. Increased density of jejunal gammadelta+ T cells in patients having normal mucosa--marker

of operative autoimmune mechanisms? Autoimmunity 1999;29(3):179-87.

137. Perez-Bravo F, Araya M, Mondragon A, Rios G, Alarcon T, Roessler JL, et al. Genetic differences in HLA-DQA1* and DQB1* allelic distributions between celiac and control children in Santiago, Chile. Human Immunology 1999;60(3):262-7.

138. Arranz E, Telleria JJ, Sanz A, Martin JF, Alonso M, Calvo C, et al. HLA-DQA1*0501 and DQB1*02 homozygosity and disease susceptibility in Spanish coeliac patients. Experimental and Clinical Immunogenetics 1997;14(4):286-90.

139. Boy MF, La Nasa G, Balestrieri A, Cherchi M, V, Usai P. Distribution of HLA-DPB1, -DQB1 -DQA1 alleles among Sardinian celiac patients. Disease Markers 1995;12(3):199-204.

140. Djilali-Saiah I, Caillat-Zucman S, Schmitz J, Chaves-Vieira ML, Bach JF. Polymorphism of antigen processing (TAP, LMP) and HLA class II genes in celiac disease. Human Immunology 1994;40(1):8-16.

141. Djilali-Saiah I, Schmitz J, Harfouch-Hammoud E, Mougenot JF, Bach JF, Caillat-Zucman S. CTLA-4 gene polymorphism is associated with predisposition to coeliac disease. Gut 1998;43(2):187-9.

142. Erkan T, Kutlu T, Yilmaz E, Cullu F, Tumay GT. Human leukocyte antigens in Turkish pediatric celiac patients. Turkish Journal of Pediatrics 1999;41(2):181-8.

143. Fernandez-Arquero M, Figueredo MA, Maluenda C, de la Concha EG. HLA-linked genes acting as additive susceptibility factors in celiac disease. Human Immunology 1995;42(4):295-300.

144. Ferrante P, Petronzelli F, Mariani P, Bonamico M, Mazzilli MC. Oligotyping of Italian celiac patients with the 11th International Histocompatibility Workshop reagents. Tissue Antigens 1992;39(1):38-9.

145. Mazzilli MC, Ferrante P, Mariani P, Martone E, Petronzelli F, Triglione P, et al. A study of Italian pediatric celiac disease patients confirms that the primary HLA association is to the DQ(alpha 1*0501, beta 1*0201) heterodimer. Human Immunology 1992;33(2):133-9.

146. Michalski JP, McCombs CC, Arai T, Elston RC, Cao T, McCarthy CF, et al. HLA-DR, DQ genotypes of celiac disease patients and healthy subjects from the West of Ireland. Tissue Antigens 1996;47(2):127-33.

147. Ruiz del Prado MY, Olivares Lopez JL, Lazaro AA, Lasierra Diaz MP. HLA system. Phenotypic and gene frequencies in celiac and healthy subjects from the same geographical area. Revista Espanola De Enfermedades Digestivas - Organo Oficial De La Sociedad Espanola De Patologia Digestiva 2001;93(2):106-13.

148. Tighe MR, Hall MA, Ashkenazi A, Siegler E, Lanchbury JS, Ciclitira PJ. Celiac disease among Ashkenazi Jews from Israel. A study of the HLA class II alleles and their associations with disease susceptibility. Human Immunology 1993;38(4):270-6.

149. Tighe MR, Hall MA, Barbado M, Cardi E, Welsh K, I, Ciclitira PJ. HLA class II alleles associated with celiac disease susceptibility in a southern European population. Tissue Antigens 1992;40(2):90-7.

150. Tuysuz B, Dursun A, Kutlu T, Sokucu S, Cine N, Suoglu O, et al. HLA-DQ alleles in patients with celiac disease in Turkey. Tissue Antigens 2001;57(6):540-2.

151. Howell WM, Leung ST, Jones DB, Nakshabendi I, Hall MA, Lanchbury JS, et al. HLA-DRB, -DQA, and -DQB polymorphism in celiac disease and enteropathy-associated T-cell lymphoma. Common features and additional risk factors for malignancy. Hum Immunol 1995;43(1):29-37.

152. Sacchetti L, Calcagno G, Ferrajolo A, Sarrantonio C, Troncone R, Micillo M, et al. Discrimination between celiac and other gastrointestinal disorders in childhood by rapid human lymphocyte antigen typing. Clinical Chemistry 1998;44(8 Pt 1):1755-7.

153. Pettersson A, Sjoberg K, Lernmark A, Eriksson S. HLA genotypes in coeliac disease and healthy individuals carrying gliadin antibodies. Eur J Gastroenterol Hepatol 1993;5(6):445-50.

154. Arnason A, Skaftadottir I, Sigmundsson J, Mooney E, Bjornsson J, Cariglia N, et al. The association between coeliac disease, dermatitis herpetiformis and certain HLA-antigens in Icelanders. European Journal of Immunogenetics - Official Journal of the British Society for Histocompatibility and Immunogenetics 1994;21(6):457-60.

155. Balas A, Vicario JL, Zambrano A, Acuna D, Garcia-Novo D. Absolute linkage of celiac disease and dermatitis herpetiformis to HLA-DQ. Tissue Antigens 1997;50(1):52-6.

156. Colonna M, Mantovani W, Corazza GR, Barboni P, Gasbarrini G, Ferrara GB, et al. Reassessment

of HLA association with celiac disease in special reference to the DP association. Human Immunology 1990;29(4):263-74.

157. Congia M, Cucca F, Frau F, Lampis R, Melis L, Clemente MG, et al. A gene dosage effect of the DQA1*0501/DQB1*0201 allelic combination influences the clinical heterogeneity of celiac disease. Human Immunology 1994;40(2):138-42.

158. Lio D, Bonanno CT, D'Anna C, De Luca S, Gervasi F, Cavataio F, et al. Gluten stimulation induces an in vitro expansion of peripheral blood T gamma delta cells from HLA-DQ2-positive subjects of families of patients with celiac disease. Experimental and Clinical Immunogenetics 1998;15(1):46-55.

159. Tumer L, Altuntas B, Hasanoglu A, Soylemezoglu O, Arinsoy T. Pattern of human leukocyte antigens in Turkish children with celiac disease. Pediatrics International - Official Journal of the Japan Pediatric Society 2000;42(6):678-81.

160. Book L, Hart A, Black J, Feolo M, Zone JJ, Neuhausen SL. Prevalence and clinical characteristics of celiac disease in Downs syndrome in a US study. American Journal of Medical Genetics 2001;98(1):70-4.

161. Farre C, Humbert P, Vilar P, Varea V, Aldeguer X, Carnicer J, et al. Serological markers and HLA-DQ2 haplotype among first-degree relatives of celiac patients. Catalonian Coeliac Disease Study Group. Digestive Diseases and Sciences 1999;44(11):2344-9.

162. Iltanen S, Collin P, Korpela M, Holm K, Partanen J, Polvi A, et al. Celiac disease and markers of celiac disease latency in patients with primary Sjogren's syndrome. American Journal of Gastroenterology 1999;94(4):1042-6.

163. Kaur G, Sarkar N, Bhatnagar S, Kumar S, Rapthap CC, Bhan MK, et al. Pediatric celiac disease in India is associated with multiple DR3-DQ2 haplotypes. Human Immunology 2002;63(8):677-82.

164. Neuhausen SL, Weizman Z, Camp NJ, Elbedour K, Sheffield VC, Zone JJ, et al. HLA DQA1-DQB1 genotypes in Bedouin families with celiac disease. Human Immunology 2002;63(6):502-7.

165. Sumnik Z, Kolouskova S, Cinek O, Kotalova R, Vavrinec J, Snajderova M. HLA-DQA1*05-DQB1*0201 positivity predisposes to coeliac disease in Czech diabetic children. Acta Paediatrica (Oslo, Norway - 1992) 2000;89(12):1426-30.

166. Liu J, Juo S, Holopainen P, Terwilliger J, Tong X, Grunn A, et al. Genomewide linkage analysis of celiac disease in Finnish families. American Journal of Human Genetics 2002;70(1):51-9.

167. Polvi A, Eland C, Koskimies S, Maki M, Partanen J. HLA DQ and DP in Finnish families with celiac disease. Eur J Immunogenet 1996;23(3):221-34.

168. Larizza D, Calcaterra V, Luinetti O, Villani L, De Silvestri A, Autelli M, et al. Evidence for immunogenetic predisposition in children with celiac disease and autoimmune thyroid disease. Int J Med Biol Environ 2001;29(2):143-8.

169. Book L, Zone JJ, Neuhausen SL. Prevalence of celiac disease among relatives of sib pairs with celiac disease in U.S. families. American Journal of Gastroenterology 2003;98(2):377-81.

170. Failla P, Ruberto C, Pagano MC, Lombardo M, Bottaro G, Perichon B, et al. Celiac disease in Down Syndrome with HLA serological and molecular studies. J Pediatr Gastroenterol Nutr 1996;23(3):303-6.

171. Maki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, et al. Prevalence of Celiac disease among children in Finland. New England Journal of Medicine 2003;348(25):2517-24.

172. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, et al. Coeliac disease among healthy members of multiple case coeliac disease families. Scandinavian Journal of Gastroenterology 2002;37(2):161-5.

173. Zubillaga P, Vidales MC, Zubillaga I, Ormaechea V, Garcia-Urkia N, Vitoria JC. HLA-DQA1 and HLA-DQB1 genetic markers and clinical presentation in celiac disease. Journal of Pediatric Gastroenterology and Nutrition 2002;34(5):548-54.

174. Bouguerra F, Babron MC, Eliaou JF, Debbabi A, Clot J, Khaldi F, et al. Synergistic effect of two HLA heterodimers in the susceptibility to celiac disease in Tunisia. Genetic Epidemiology 1997;14(4):413-22.

175. Popat S, Hearle N, Wixey J, Hogberg L, Bevan S, Lim W, et al. Analysis of the CTLA4 gene in Swedish coeliac disease patients. Scand J Gastroenterol 2002;37(1):28-31.

176. Karell K, Louka AS, Moodie SJ, Ascher H, Clot F, Greco L, et al. HLA types in celiac disease patients not carrying the DQA1 *05-DQB1 *02 (DQ2) heterodimer: Results from the European

genetics cluster on celiac disease. Hum Immunol 2003;64(4):469-77.

177. Lewis C, Book L, Black J, Sawitzke A, Cannon-Albright L, Zone J, et al. Celiac disease and human leukocyte antigen genotype: accuracy of diagnosis in self-diagnosed individuals, dosage effect, and sibling risk. J Pediatr Gastroenterol Nutr 2000;31(1):22-7.

178. Howard FM, Carter CO, Candy DC, Harries JT. A family study of protracted diarrhoea in infancy. Journal of Medical Genetics 1981;18(2):81-6.

179. Volta U, Molinaro N, De Franceschi L, Bianchi FB. Human umbilical cord as substrate for IgA antiendomysial antibodies allows large scale screening for celiac sprue. J Clin Gastroenterol 1996;23(1):18-20.

180. Trevisiol C, Ventura A, Baldas V, Tommasini A, Santon D, Martelossi S, et al. A reliable screening procedure for coeliac disease in clinical practice. Scand J Gastroenterol 2002;37(6):679-84.

181. Bonamico M, Mariani P, Mazzilli MC, Triglione P, Lionetti P, Ferrante P, et al. Frequency and clinical pattern of celiac disease among siblings of celiac children. J Pediatr Gastroenterol Nutr 1996;23(2):159-63.

182. Catassi C, Ratsch IM, Fabiani E, Ricci S, Bordicchia F, Pierdomenico R, et al. High prevalence of undiagnosed coeliac disease in 5280 Italian students screened by antigliadin antibodies. Acta Paediatr 1995;84(6):672-6.

183. Catassi C, Ratsch IM, Fabiani E, Rossini M, Bordicchia F, Candela F, et al. Coeliac disease in the year 2000: exploring the iceberg. Lancet 1994;343(8891):200-3.

184. Grodzinsky E, Hed J, Lieden G, Sjogren F, Strom M. Presence of IgA and IgG antigliadin antibodies in healthy adults as measured by micro-ELISA. Effect of various cutoff levels on specificity and sensitivity when diagnosing coeliac disease. Int Arch Allergy Appl Immunol 1990;92(2):119-23.

185. Grodzinsky E, Franzen L, Hed J, Strom M. High prevalence of celiac disease in healthy adults revealed by antigliadin antibodies. Ann Allergy 1992;69(1):66-70.

186. Lagerqvist C, Ivarsson A, Juto P, Persson LA, Hernell O. Screening for adult coeliac disease - which serological marker(s) to use? J Intern Med 2001;250(3):241-8.

187. Weile B, Grodzinsky E, Skogh T, Jordal R, Cavell B, Krasilnikoff PA. Screening Danish blood donors for antigliadin and antiendomysium antibodies. Acta Paediatr 1996;412(Suppl):46.

188. McMillan SA, Watson RP, McCrum EE, Evans AE. Factors associated with serum antibodies to reticulin, endomysium, and gliadin in an adult population. Gut 1996;39(1):43-7.

189. Stenhammar L, Brandt A, Wagermark J. A family study of coeliac disease. Acta Paediatr Scand 1982;71(4):625-8.

190. Maki M, Holm K, Lipsanen V, Hallstrom O, Viander M, Collin P, et al. Serological markers and HLA genes among healthy first-degree relatives of patients with coeliac disease. Lancet 1991;338(8779):1350-3.

191. Sjoberg K, Wassmuth R, Reichstetter S, Eriksson KF, Ericsson UB, Eriksson S. Gliadin antibodies in adult insulin-dependent diabetes--autoimmune and immunogenetic correlates. Autoimmunity 2000;32(4):217-28.

192. Carlsson AK, Axelsson IE, Borulf SK, Bredberg AC, Lindberg BA, Sjoberg KG, et al. Prevalence of IgA-antiendomysium and IgA-antigliadin autoantibodies at diagnosis of insulin-dependent diabetes mellitus in Swedish children and adolescents. Pediatrics 1999;103(6 Pt 1):1248-52.

193. Ivarsson A, Persson LA, Nystrom L, Ascher H, Cavell B, Danielsson L, et al. Epidemic of coeliac disease in Swedish children. Acta Paediatr 2000;89(2):165-71.

194. Maki M, Holm K. Incidence and prevalence of coeliac disease in Tampere. Coeliac disease is not disappearing. Acta Paediatrica Scandinavica 1990;79(10):980-2.

195. Ivarsson A, Persson LA, Nystrom L, Hernell O. The Swedish coeliac disease epidemic with a prevailing twofold higher risk in girls compared to boys may reflect gender specific risk factors. Eur J Epidemiol 2003;18(7):677-84.

196. Weile B, Krasilnikoff PA. Extremely low incidence rates of celiac disease in the Danish population of children. J Clin Epidemiol 1993;46(7):661-4.

197. Bode S, Gudmand-Hoyer E. Incidence and prevalence of adult coeliac disease within a defined geographic area in Denmark. Scand J Gastroenterol 1996;31(7):694-9.

198. Maki M, Kallonen K, Lahdeaho ML, Visakorpi JK. Changing pattern of childhood coeliac disease in Finland. Acta Paediatrica Scandinavica 1988;77(3):408-12.

199. Collin P, Reunala T, Rasmussen M, Kyronpalo S, Pehkonen E, Laippala P, et al. High incidence and prevalence of adult coeliac disease. Augmented diagnostic approach. Scand J Gastroenterol 1997;32(11):1129-33.

200. Jansen Th TLA, Mulder CJJ, Karssen PHZ, Wagenaar CGJ. Epidemiological survey of the Dutch Coeliac Disease Society: An update 1992. Eur J Gastroenterol Hepatol 1993;5(2):73-8.

201. Hawkes ND, Swift GL, Smith PM, Jenkins HR. Incidence and presentation of coeliac disease in South Glamorgan. Eur J Gastroenterol Hepatol 2000;12(3):345-9.

202. Corrao G, Usai P, Galatola G, Ansaldi N, Meini A, Pelli MA, et al. Estimating the incidence of coeliac disease with capture-recapture methods within four geographic areas in Italy. J Epidemiol Community Health 1996;50(3):299-305.

203. Magazzu G, Bottaro G, Cataldo F, Iacono G, Di Donato F, Patane R, et al. Increasing incidence of childhood celiac disease in Sicily: results of a multicenter study. Acta Paediatr 1994;83(10):1065-9.

204. Lopez-Rodriguez MJ, Canal Macias ML, Lavado Garcia JM, Sanchez BM, Robledo AP, Pedrera Zamorano JD. Epidemiological changes in diagnosed coeliac disease in a population of Spanish children. Acta Paediatr 2003;92(2):165-9.

205. Talley NJ, Valdovinos M, Petterson TM, Carpenter HA, Melton LJ. Epidemiology of celiac sprue: a community-based study. Am J Gastroenterol 1994;89(6):843-6.

206. Fasano A, Berti I, Gerarduzzi T, Not T, Colletti RB, Drago S, et al. Prevalence of celiac disease in at-risk and not-at-risk groups in the United States: a large multicenter study. Arch Intern Med 2003;163(3):286-92.

207. Green PH, Shane E, Rotterdam H, Forde KA, Grossbard L. Significance of unsuspected celiac disease detected at endoscopy. Gastroenterol Int 2000;51(1):60-5.

208. Not T, Horvath K, Hill ID, Partanen J, Hammed A, Magazzu G, et al. Celiac disease risk in the USA: high prevalence of antiendomysium antibodies in healthy blood donors. Scand J Gastroenterol 1998;33(5):494-8.

209. Carlsson AK, Axelsson IE, Borulf SK, Bredberg AC, Ivarsson SA. Serological screening for celiac disease in healthy 2.5-year-old children in Sweden. Pediatrics 2001;107(1):42-5.

210. Collin P, Rasmussen M, Kyronpalo S, Laippala P, Kaukinen K. The hunt for coeliac disease in primary care. Q J Med 2002;95(2):75-7.

211. Grodzinsky E. Screening for coeliac disease in apparently healthy blood donors. Acta Paediatr 1996;412(Suppl):36-8.

212. Hovdenak N, Hovlid E, Aksnes L, Fluge G, Erichsen MM, Eide J. High prevalence of asymptomatic coeliac disease in Norway: a study of blood donors. Eur J Gastroenterol Hepatol 1999;11(2):185-7.

213. Ivarsson A, Persson LA, Juto P, Peltonen M, Suhr O, Hernell O. High prevalence of undiagnosed coeliac disease in adults: a Swedish population-based study. J Intern Med 1999;245(1):63-8.

214. Kolho KL, Farkkila MA, Savilahti E. Undiagnosed coeliac disease is common in Finnish adults. Scand J Gastroenterol 1998;33(12):1280-3.

215. Maki M, Mustalahti K, Kokkonen J, Kulmala P, Haapalahti M, Karttunen T, et al. Prevalence of Celiac disease among children in Finland. New Engl J Med 2003;348(25):2517-24.

216. Sjoberg K, Alm R, Ivarsson SA, Lindstrom C, Eriksson S. Prevalence and clinical significance of gliadin antibodies in healthy children and adults. Scand J Gastroenterol 1994;29(3):248-54.

217. Sjoberg K, Eriksson S. Regional differences in coeliac disease prevalence in Scandinavia? Scand J Gastroenterol 1999;34(1):41-5.

218. Weile I, Grodzinsky E, Skogh T, Jordal R, Cavell B, Krasilnikoff PA. High prevalence rates of adult silent coeliac disease, as seen in Sweden, must be expected in Denmark. APMIS 2001;109(11):745-50.

219. Borch K, Grodzinsky E, Petersson F, Jonsson K-A, Mardh S, Valdimarsson T. Prevalence of coeliac disease and relations to Helicobacter pylori infection and duodenitis in a Swedish adult population sample: A histomorphological and serological survey. Inflammopharmacology 2000;8(4):341-50.

220. Catassi C, Fabiani E, Ratsch IM, Coppa G, V, Giorgi PL, Pierdomenico R, et al. The coeliac

iceberg in Italy. A multicentre antigliadin antibodies screening for coeliac disease in school-age subjects. Acta Paediatr 1996;412(Suppl):29-35.

221. Catassi C, Fanciulli G, D'Appello AR, El Asmar R, Rondina C, Fabiani E, et al. Antiendomysium versus antigliadin antibodies in screening the general population for coeliac disease. Scand J Gastroenterol 2000;35(7):732-6.

222. Corazza GR, Andreani ML, Biagi F, Corrao G, Pretolani S, Giulianelli G, et al. The smaller size of the 'coeliac iceberg' in adults. Scand J Gastroenterol 1997;32(9):917-9.

223. Mazzetti dP, Giorgetti GM, Gregori M, De Simone M, Leonardi C, Barletta PA, et al. Subclinical coeliac disease. Ital J Gastroenterol 1992;24(6):352-4.

224. Pittschieler K, Ladinser B. Coeliac disease: screened by a new strategy. Acta Paediatr 1996;412(Suppl):42-5.

225. Trevisiol C, Not T, Berti I, Buratti E, Citta A, Neri E, et al. Screening for coeliac disease in healthy blood donors at two immuno-transfusion centres in north-east Italy. Ital J Gastroenterol Hepatol 1999;31(7):584-6.

226. Johnston SD, Watson RG, McMillan SA, Sloan J, Love AH. Coeliac disease detected by screening is not silent--simply unrecognized. Q J Med 1998;91(12):853-60.

227. Sanders DS, Patel D, Stephenson TJ, Ward AM, McCloskey EV, Hadjivassiliou M, et al. A primary care cross-sectional study of undiagnosed adult coeliac disease. Eur J Gastroenterol Hepatol 2003;15(4):407-13.

228. West J, Logan RFA, Hill PG, Lloyd A, Lewis S, Hubbard R, et al. Seroprevalence, correlates, and characteristics of undetected coeliac disease in England. Gut 2003;52(7):960-5.

229. Dickey W, McMillan SA, Bharucha C, Porter KG. Antigliadin antibodies in blood donors in Northern Ireland. Eur J Gastroenterol Hepatol 1992;4(9):739-41.

230. Riestra S, Fernandez E, Rodrigo L, Garcia S, Ocio G. Prevalence of Coeliac disease in the general population of northern Spain. Strategies of serologic screening. Scand J Gastroenterol 2000;35(4):398-402.

231. Rostami K, Mulder CJ, Werre JM, van Beukelen FR, Kerchhaert J, Crusius JB, et al. High prevalence of celiac disease in apparently healthy blood donors suggests a high prevalence of undiagnosed celiac disease in the Dutch population. Scand J Gastroenterol 1999;34(3):276-9.

232. Csizmadia CGDS, Mearin ML, Von Blomberg BME, Brand R, Verloove-Vanhorick SP. An iceberg of childhood coeliac disease in the Netherlands. Lancet 1999;353(9155):813-4.

233. Rutz R, Ritzler E, Fierz W, Herzog D. Prevalence of asymptomatic celiac disease in adolescents of eastern Switzerland. Swiss Med Wkly 2002;132(3-4):43-7.

234. Jaeger C, Hatziagelaki E, Petzoldt R, Bretzel RG. Comparative analysis of organ-specific autoantibodies and celiac disease--associated antibodies in type 1 diabetic patients, their first-degree relatives, and healthy control subjects. Diabetes Care 2001;24(1):27-32.

235. Book L, Zone JJ, Neuhausen SL. Prevalence of celiac disease among relatives of sib pairs with celiac disease in U.S. families. Am J Gastroenterol 2003;98(2):377-81.

236. Corazza G, Valentini RA, Frisoni M, Volta U, Corrao G, Bianchi FB, et al. Gliadin immune reactivity is associated with overt and latent enteropathy in relatives of celiac patients. Gastroenterology 1992;103(5):1517-22.

237. Farre C, Humbert P, Vilar P, Varea V, Aldeguer X, Carnicer J, et al. Serological markers and HLA-DQ2 haplotype among first-degree relatives of celiac patients. Catalonian Coeliac Disease Study Group. Dig Dis Sci 1999;44(11):2344-9.

238. Hill I, Fasano A, Schwartz R, Counts D, Glock M, Horvath K. The prevalence of celiac disease in at-risk groups of children in the United States. J Pediatr 2000;136(1):86-90.

239. Korponay-Szabo I, Kovacs J, Lorincz M, Torok E, Goracz G. Families with multiple cases of gluten-sensitive enteropathy. Z Gastroenterol 1998;36(7):553-8.

240. Kotze LM, Utiyama SR, Nisihara RM, Zeni MP, de Sena MG, Amarante HM. Antiendomysium antibodies in Brazilian patients with celiac disease and their first-degree relatives. Arq Gastroenterol 2001;38(2):94-103.

241. Mustalahti K, Sulkanen S, Holopainen P, Laurila K, Collin P, Partanen J, et al. Coeliac disease among healthy members of multiple case coeliac disease families. Scand J Gastroenterol 2002;37(2):161-5.

242. Robinson DC, Watson AJ, Wyatt EH, Marks JM, Roberts DF. Incidence of small-intestinal mucosal abnormalities and of clinical coeliac disease in the relatives of children with coeliac disease. Gut 1971;12(10):789-93.

243. Rolles CJ, Myint TO, Sin WK, Anderson M. Proceedings: Family study of coeliac disease. Gut 1974;15(10):827.

244. Rostami K, Mulder CJ, van Overbeek FM, Kerckhaert J, Meijer JW, von Blomberg MB, et al. Should relatives of coeliacs with mild clinical complaints undergo a small-bowel biopsy despite negative serology? Eur J Gastroenterol Hepatol 2000;12(1):51-5.

245. Stokes PL, Ferguson R, Holmes GK, Cooke WT. Familial aspects of coeliac disease. Quarterly Journal of Medicine 1976;45(180):567-82.

246. Vitoria JC, Arrieta A, Astigarraga I, Garcia-Masdevall D, Rodriguez-Soriano J. Use of serological markers as a screening test in family members of patients with celiac disease. J Pediatr Gastroenterol Nutr 1994;19(3):304-9.

247. Hogberg L, Falth-Magnusson K, Grodzinsky E, Stenhammar L. Familial prevalence of coeliac disease: A twenty-year follow-up study. Scand J Gastroenterol 2003;38(1):61-5.

248. Pittschieler K, Gentili L, Niederhofer H. Onset of coeliac disease: A prospective longitudinal study. Acta Paediatr Int J Paediatr 2003;92(10):1149-52.

249. Tursi A, Brandimarte G, Giorgetti GM, Inchingolo CD. Effectiveness of the sorbitol HSUB2 breath test in detecting histological damage among relatives of coeliacs. Scand J Gastroenterol 2003;38(7):727-31.

250. Acerini CL, Ahmed ML, Ross KM, Sullivan PB, Bird G, Dunger DB. Coeliac disease in children and adolescents with IDDM: clinical characteristics and response to gluten-free diet. Diabet Med 1998;15(1):38-44.

251. Aktay AN, Lee PC, Kumar V, Parton E, Wyatt DT, Werlin SL. The prevalence and clinical characteristics of celiac disease in juvenile diabetes in Wisconsin. J Pediatr Gastroenterol Nutr 2001;33(4):462-5.

252. Arato A, Korner A, Veres G, Dezsofi A, Ujpal I, Madacsy L. Frequency of coeliac disease in Hungarian children with type 1 diabetes mellitus. Eur J Pediatr 2002;162(1):1-5.

253. Bao F, Yu L, Babu S, Wang T, Hoffenberg EJ, Rewers M, et al. One third of HLA DQ2 homozygous patients with type 1 diabetes express celiac disease-associated transglutaminase autoantibodies. J Autoimmun 1999;13(1):143-8.

254. Barera G, Bianchi C, Calisti L, Cerutti F, Dammacco F, Frezza E, et al. Screening of diabetic children for coeliac disease with antigliadin antibodies and HLA typing. Arch Dis Child 1991;66(4):491-4.

255. Barera G, Bonfanti R, Viscardi M, Bazzigaluppi E, Calori G, Meschi F, et al. Occurrence of celiac disease after onset of type 1 diabetes: a 6-year prospective longitudinal study. Pediatrics 2002;109(5):833-8.

256. Calero P, Ribes-Koninckx C, Albiach V, Carles C, Ferrer J. IgA antigliadin antibodies as a screening method for nonovert celiac disease in children with insulin-dependent diabetes mellitus. J Pediatr Gastroenterol Nutr 1996;23(1):29-33.

257. Cronin CC, Feighery A, Ferriss JB, Liddy C, Shanahan F, Feighery C. High prevalence of celiac disease among patients with insulin-dependent (type I) diabetes mellitus. Am J Gastroenterol 1997;92(12):2210-2.

258. De Block CE, De L, I, Vertommen JJ, Rooman RP, Du CM, V, Van Campenhout CM, et al. Beta-cell, thyroid, gastric, adrenal and coeliac autoimmunity and HLA-DQ types in type 1 diabetes. Clin Exp Immunol 2001;126(2):236-41.

259. De V, I, Ghirlanda G, Gasbarrini G. Prevalence of coeliac disease in type I diabetes: a multicentre study. Acta Paediatr 1996;412(Suppl):56-7.

260. Fraser-Reynolds KA, Butzner JD, Stephure DK, Trussell RA, Scott RB. Use of immunoglobulin A-antiendomysial antibody to screen for celiac disease in North American children with type 1 diabetes. Diabetes Care 1998;21(11):1985-9.

261. Gillett PM, Gillett HR, Israel DM, Metzger DL, Stewart L, Chanoine JP, et al. High prevalence of celiac disease in patients with type 1 diabetes detected by antibodies to endomysium and tissue transglutaminase. Can J Gastroenterol 2001;15(5):297-301.

262. Hansen D, Bennedbaek FN, Hansen LK, Hoier-Madsen M, Hegedu LS, Jacobsen BB, et al. High prevalence of coeliac disease in Danish children with type I diabetes mellitus. Acta Paediatr 2001;90(11):1238-43.

263. Kaukinen K, Collin P, Mykkanen AH, Partanen J, Maki M, Salmi J. Celiac disease and

autoimmune endocrinologic disorders. Dig Dis Sci 1999;44(7):1428-33.

264. Kordonouri O, Dieterich W, Schuppan D, Webert G, Muller C, Sarioglu N, et al. Autoantibodies to tissue transglutaminase are sensitive serological parameters for detecting silent coeliac disease in patients with Type 1 diabetes mellitus. Diabet Med 2000;17(6):441-4.

265. Lampasona V, Bonfanti R, Bazzigaluppi E, Venerando A, Chiumello G, Bosi E, et al. Antibodies to tissue transglutaminase C in type I diabetes. Diabetologia 1999;42(10):1195-8.

266. Li Voon Chong JSW, Leong KS, Wallymahmed M, Sturgess R, MacFarlane IA. Is coeliac disease more prevalent in young adults with coexisting Type 1 diabetes mellitus and autoimmune thyroid disease compared with those with Type 1 diabetes mellitus alone? Diabet Med 2002;19(4):334-7.

267. Lorini R, Scotta MS, Cortona L, Avanzini MA, Vitali L, De Giacomo C, et al. Celiac disease and type I (insulin-dependent) diabetes mellitus in childhood: follow-up study. J Diabetes Complications 1996;10(3):154-9.

268. Not T, Tommasini A, Tonini G, Buratti E, Pocecco M, Tortul C, et al. Undiagnosed coeliac disease and risk of autoimmune disorders in subjects with Type I diabetes mellitus. Diabetologia 2001;44(2):151-5.

269. Page SR, Lloyd CA, Hill PG, Peacock I, Holmes GK. The prevalence of coeliac disease in adult diabetes mellitus. Q J Med 1994;87(10):631-7.

270. Rensch MJ, Merenich JA, Lieberman M, Long BD, Davis DR, McNally PR. Gluten-sensitive enteropathy in patients with insulin-dependent diabetes mellitus. Ann Intern Med 1996;124(6):564-7.

271. Roldan MB, Barrio R, Roy G, Parra C, Alonso M, Yturriaga R, et al. Diagnostic value of serological markers for celiac disease in diabetic children and adolescents. J Pediatr Endocrinol Metab 1998;11(6):751-6.

272. Rossi TM, Albini CH, Kumar V. Incidence of celiac disease identified by the presence of serum endomysial antibodies in children with chronic diarrhea, short stature, or insulin-dependent diabetes mellitus. J Pediatr 1993;123(2):262-4.

273. Sategna-Guidetti C, Grosso S, Pulitano R, Benaduce E, Dani F, Carta Q. Celiac disease and insulin-dependent diabetes mellitus. Screening in an adult population. Dig Dis Sci 1994;39(8):1633-7.

274. Saukkonen T, Savilahti E, Reijonen H, Ilonen J, Tuomilehto-Wolf E, Akerblom HK. Coeliac disease: frequent occurrence after clinical onset of insulin-dependent diabetes mellitus. Childhood Diabetes in Finland Study Group. Diabet Med 1996;13(5):464-70.

275. Schober E, Bittmann B, Granditsch G, Huber WD, Huppe A, Jager A, et al. Screening by anti-endomysium antibody for celiac disease in diabetic children and adolescents in Austria. J Pediatr Gastroenterol Nutr 2000;30(4):391-6.

276. Sigurs N, Johansson C, Elfstrand PO, Viander M, Lanner A. Prevalence of coeliac disease in diabetic children and adolescents in Sweden. Acta Paediatr 1993;82(9):748-51.

277. Sjoberg K, Eriksson KF, Bredberg A, Wassmuth R, Eriksson S. Screening for coeliac disease in adult insulin-dependent diabetes mellitus. J Intern Med 1998;243(2):133-40.

278. Spiekerkoetter U, Seissler J, Wendel U. General screening for celiac disease is advisable in children with type 1 diabetes. Horm Metab Res 2002;34(4):192-5.

279. Talal AH, Murray JA, Goeken JA, Sivitz W, I. Celiac disease in an adult population with insulin-dependent diabetes mellitus: use of endomysial antibody testing. Am J Gastroenterol 1997;92(8):1280-4.

280. Valerio G, Maiuri L, Troncone R, Buono P, Lombardi F, Palmieri R, et al. Severe clinical onset of diabetes and increased prevalence of other autoimmune diseases in children with coeliac disease diagnosed before diabetes mellitus. Diabetologia 2002;45(12):1719-22.

281. Vitoria JC, Castano L, Rica I, Bilbao JR, Arrieta A, Garcia-Masdevall MD. Association of insulin-dependent diabetes mellitus and celiac disease: a study based on serologic markers. J Pediatr Gastroenterol Nutr 1998;27(1):47-52.

282. Agardh D, Nilsson A, Tuomi T, Lindberg B, Carlsson AK, Lernmark A, et al. Prediction of silent celiac disease at diagnosis of childhood type 1 diabetes by tissue transglutaminase autoantibodies and HLA. Pediatr Diabetes 2001;2(2):58-65.

283. Ackerman Z, Eliakim R, Stalnikowicz R, Rachmilewitz D. Role of small bowel biopsy in the endoscopic evaluation of adults with iron

deficiency anemia. Am J Gastroenterol 1996;91(10):2099-102.

284. Annibale B, Capurso G, Chistolini A, D'Ambra G, DiGiulio E, Monarca B, et al. Gastrointestinal causes of refractory iron deficiency anemia in patients without gastrointestinal symptoms. Am J Med 2001;111(6):439-45.

285. Corazza GR, Valentini RA, Andreani ML, D'Anchino M, Leva MT, Ginaldi L, et al. Subclinical coeliac disease is a frequent cause of iron-deficiency anaemia. Scand J Gastroenterol 1995;30(2):153-6.

286. Dickey W, Kenny BD, McMillan SA, Porter KG, McConnell JB. Gastric as well as duodenal biopsies may be useful in the investigation of iron deficiency anaemia. Scand J Gastroenterol 1997;32(5):469-72.

287. Howard MR, Turnbull AJ, Morley P, Hollier P, Webb R, Clarke A. A prospective study of the prevalence of undiagnosed coeliac disease in laboratory defined iron and folate deficiency. J Clinl Path 2002;55(10):754-7.

288. Kepczyk T, Kadakia SC. Prospective evaluation of gastrointestinal tract in patients with iron-deficiency anemia. Dig Dis Sci 1995;40(6):1283-9.

289. McIntyre AS, Long RG. Prospective survey of investigations in outpatients referred with iron deficiency anaemia. Gut 1993;34(8):1102-7.

290. Oxentenko AS, Grisolano SW, Murray JA, Burgart LJ, Dierkhising RA, Alexander JA. The insensitivity of endoscopic markers in celiac disease. Am J Gastroenterol 2002;97(4):933-8.

291. Ransford Rupert AJ, Hayes M, Palmer M, Hall MJ. A controlled, prospective screening study of celiac disease presenting as iron deficiency anemia. J Clin Gastroenterol 2002;35(3):228-33.

292. Unsworth DJ, Lock RJ, Harvey RF. Improving the diagnosis of coeliac disease in anaemic women. Br J Haematol 2000;111(3):898-901.

293. Annibale B, Lahner E, Chistolini A, Gallucci C, Di Giulio E, Capurso G, et al. Endoscopic evaluation of the upper gastrointestinal tract is worthwhile in premenopausal women with iron-deficiency anaemia irrespective of menstrual flow. Scand J Gastroenterol 2003;38(3):239-45.

294. Van Mook WNKA, Bourass-Bremer IHDN, Bos LP, Verhoeven HMJM, Engels LGJB. The outcome of esophagogastroduodenoscopy (EGD) in asymptomatic outpatients with iron deficiency anemia after a negative colonoscopy. Eur J Intern Med 2001;12(2):122-6.

295. Gonzalez D, Sugai E, Gomez JC, Oliveri MB, Gomez AC, Vega E, et al. Is it necessary to screen for celiac disease in postmenopausal osteoporotic women? Calcif Tissue Int 2002;71(2):141-4.

296. Lindh E, Ljunghall S, Larsson K, Lavo B. Screening for antibodies against gliadin in patients with osteoporosis. J Intern Med 1992;231(4):403-6.

297. Mather KJ, Meddings JB, Beck PL, Scott RB, Hanley DA. Prevalence of IgA-antiendomysial antibody in asymptomatic low bone mineral density. Am J Gastroenterol 2001;96(1):120-5.

298. Nuti R, Martini G, Valenti R, Giovani S, Salvadori S, Avanzati A. Prevalence of undiagnosed coeliac syndrome in osteoporotic women. J Intern Med 2001;250(4):361-6.

299. Hin H, Bird G, Fisher P, Mahy N, Jewell D. Coeliac disease in primary care: case finding study. BMJ 1999;318(7177):164-7.

300. Bardella MT, Molteni N, Cesana B, Baldassarri AR, Binanchi PA. IgA antigliadin antibodies, cellobiose/mannitol sugar test, and carotenemia in the diagnosis of and screening for celiac disease. Am J Gastroenterol 1991;86(3):309-11.

301. Bardella MT, Trovato C, Cesana BM, Pagliari C, Gebbia C, Peracchi M. Serological markers for coeliac disease: is it time to change? Dig Liver Dis 2001;33(5):426-31.

302. Bode S, Weile B, Krasilnikoff PA, Gudmand-Hoyer E. The diagnostic value of the gliadin antibody test in celiac disease in children: a prospective study. J Pediatr Gastroenterol Nutr 1993;17(3):260-4.

303. Carroccio A, Vitale G, Di Prima L, Chifari N, Napoli S, La Russa C, et al. Comparison of anti-transglutaminase ELISAs and an anti-endomysial antibody assay in the diagnosis of celiac disease: a prospective study. Clin Chem 2002;48(9):1546-50.

304. Chan AW, Butzner JD, McKenna R, Fritzler MJ. Tissue transglutaminase enzyme-linked immunosorbent assay as a screening test for celiac disease in pediatric patients. Pediatrics 2001;107(1):E8.

305. Chartrand LJ, Agulnik J, Vanounou T, Russo PA, Baehler P, Seidman EG. Effectiveness of antigliadin antibodies as a screening test for celiac disease in children. CMAJ 1997;157(5):527-33.

306. Day AS, Cook HB, Whitehead M, Abbott GD. Anti-endomysial and anti-gliadin antibodies in screening for coeliac disease in children at greater risk of developing coeliac disease. N Z Med J 2000;113(1119):412-3.

307. Fitzpatrick KP, Sherman PM, Ipp M, Saunders N, Macarthur C. Screening for celiac disease in children with recurrent abdominal pain. J Pediatr Gastroenterol Nutr 2001;33(3):250-2.

308. Thomas AG, Phillips AD, Walker-Smith JA. The value of proximal small intestinal biopsy in the differential diagnosis of chronic diarrhoea. Arch Dis Child 1992;67(6):741-3.

309. Ventura A, Facchini S, Amantidu C, Andreotti MF, Andrighetto A, Baggiani A, et al. Searching for celiac disease in pediatric general practice. Clin Pediatr 2001;40(10):575-7.

310. Ascher H, Krantz I, Kristiansson B. Increasing incidence of coeliac disease in Sweden. Arch Dis Child 1991;66(5):608-11.

311. Cavell B, Stenhammar L, Ascher H, Danielsson L, Dannaeus A, Lindberg T, et al. Increasing incidence of childhood coeliac disease in Sweden. Results of a national study. Acta Paediatr 1992;81(8):589-92.

312. Stenhammar L, Johansson CG. The incidence of coeliac disease in children in south-east Sweden. Acta Paediatrica Scandinavica 1981;70(3):379-81.

313. Weile B, Cavell B, Nivenius K, Krasilnikoff PA. Striking differences in the incidence of childhood celiac disease between Denmark and Sweden: a plausible explanation. J Pediatr Gastroenterol Nutr 1995;21(1):64-8.

314. Weile B, Krasilnikoff PA. Low incidence rates by birth of symptomatic coeliac disease in a Danish population of children. Acta Paediatr 1992;81(5):394-8.

315. Stevens FM, Egan-Mitchell B, Cryan E, McCarthy CF, McNicholl B. Decreasing incidence of coeliac disease. Arch Dis Child 1987;62(5):465-8.

316. Logan RF, Rifkind EA, Busuttil A, Gilmour HM, Ferguson A. Prevalence and "incidence" of celiac disease in Edinburgh and the Lothian region of Scotland. Gastroenterology 1986;90(2):334-42.

317. Hoffenberg EJ, Bao F, Eisenbarth GS, Uhlhorn C, Haas JE, Sokol RJ, et al. Transglutaminase antibodies in children with a genetic risk for celiac disease. Journal of Pediatrics 2000;137(3):356-60.

318. Corazza GR, Frisoni M, Treggiari EA, Valentini RA, Filipponi C, Volta U, et al. Subclinical celiac sprue. Increasing occurrence and clues to its diagnosis. J Clin Gastroenterol 1993;16(1):16-21.

319. Sanders DS, Hurlstone DP, Stokes RO, Rashid F, Milford-Ward A, Hadjivassiliou M, et al. Changing face of adult coeliac disease: experience of a single university hospital in South Yorkshire. Postgraduate Medical Journal 2002;78(915):31-3.

320. Sategna-Guidetti C, Grosso S. Changing pattern in adult coeliac disease: A 24-year survey. Eur J Gastroenterol Hepatol 1994;6(1):15-9.

321. Holmes GK. Non-malignant complications of coeliac disease. Acta Paediatrica Supplement 1996;412:68-75.

322. Suharjono. Intestinal biopsy and coeliac disease. Paediatrica Indonesiana 1971;11(3):116-34.

323. Mannell A, van Heerden JA, Weiland LH, Ilstrup DM. Factors influencing survival after resection for ductal adenocarcinoma of the pancreas. Annals of Surgery 1986;203(4):403-7.

324. Atherton DJ. Diagnosis and management of skin disorders caused by food allergy. Annals of Allergy 1984;53(6 Pt 2):623-8.

325. Di Stefano M, Jorizzo RA, Veneto G, Cecchetti L, Gasbarrini G, Corazza GR. Bone mass and metabolism in dermatitis herpetiformis. Dig Dis Sci 1999;44(10):2139-43.

326. Ots M, Uibo O, Metskula K, Uibo R, Salupere V. IgA-antigliadin antibodies in patients with IgA nephropathy: the secondary phenomenon? American Journal of Nephrology 1999;19(4):453-8.

327. Collin P, Reunala T, Pukkala E, Laippala P, Keyrilainen O, Pasternack A. Coeliac disease--associated disorders and survival. Gut 1994;35(9):1215-8.

328. Cooper BT, Holmes GK, Cooke WT. Lymphoma risk in coeliac disease of later life. Digestion 1982;23(2):89-92.

329. Harris OD, Cooke WT, Thompson H, Waterhouse JA. Malignancy in adult coeliac disease and idiopathic steatorrhoea. Am J Med 1967;42(6):899-912.

330. Holmes GK, Stokes PL, Sorahan TM, Prior P, Waterhouse JA, Cooke WT. Coeliac disease, gluten-free diet, and malignancy. Gut 1976;17(8):612-9.

331. Peters U, Askling J, Gridley G, Ekbom A, Linet M. Causes of death in patients with celiac disease in a population-based Swedish cohort. Archives of Internal Medicine 2003;163(13):1566-72.

332. Cooper BT, Holmes GKT, Ferguson R. Celiac disease and malignancy. Medicine 1980;59(4):249-61.

333. Holmes GK, Prior P, Lane MR, Pope D, Allan RN. Malignancy in coeliac disease--effect of a gluten free diet. Gut 1989;30(3):333-8.

334. Sorensen HT, Fonager K. Risk estimation of disorders associated with coeliac disease. A 16-year Danish nationwide follow-up study based on hospital discharge data. Implications for screening. Int J Risk Saf Med 1996;8(2):137-40.

335. Cottone M, Termini A, Oliva L, Magliocco A, Marrone C, Orlando A, et al. Mortality and causes of death in celiac disease in a Mediterranean area. Dig Dis Sci 1999;44(12):2538-41.

336. Logan RF, Rifkind EA, Turner ID, Ferguson A. Mortality in celiac disease. Gastroenterology 1989;97(2):265-71.

337. Askling J, Linet M, Gridley G, Halstensen TS, Ekstrom K, Ekbom A. Cancer incidence in a population-based cohort of individuals hospitalized with celiac disease or dermatitis herpetiformis. Gastroenterology 2002;123(5):1428-35.

338. Collin P, Pukkala E, Reunala T. Malignancy and survival in dermatitis herpetiformis: a comparison with coeliac disease. Gut 1996;38(4):528-30.

339. Corrao G, Corazza GR, Bagnardi V, Brusco G, Ciacci C, Cottone M, et al. Mortality in patients with coeliac disease and their relatives: a cohort study. Lancet 2001;358(9279):356-61.

340. Green PHR, Fleischauer AT, Bhagat G, Goyal R, Jabri B, Neugut AI. Risk of malignancy in patients with celiac disease. Am J Med 2003;115(3):191-5.

341. Selby WS, Gallagher ND. Malignancy in a 19-year experience of adult celiac disease. Dig Dis Sci 1979;24(9):684-8.

342. Delco F, El Serag HB, Sonnenberg A. Celiac sprue among US military veterans: associated disorders and clinical manifestations. Dig Dis Sci 1999;44(5):966-72.

343. Nielsen OH, Jacobsen O, Pedersen ER, Rasmussen SN, Petri M, Laulund S, et al. Non-tropical sprue. Malignant diseases and mortality rate. Scand J Gastroenterol 1985;20(1):13-8.

344. Shainoff JR, Valenzuela R, Urbanic DA, DiBello PM, Lucas F, V, Graor R. Fibrinogen A alpha and gamma-chain dimers as potential differential indicators of atherosclerotic and thrombotic vascular disease. Blood Coagulation & Fibrinolysis - an International Journal in Haemostasis and Thrombosis 1990;1(4-5):499-503.

345. Ribes-Koninckx C, Alfonso P, Ortigosa L, Escobar H, Suarez L, Arranz E, et al. A beta-turn rich oats peptide as an antigen in an ELISA method for the screening of coeliac disease in a paediatric population. European Journal of Clinical Investigation 2000;30(8):702-8.

346. Kemppainen TA, Kosma VM, Janatuinen EK, Julkunen RJ, Pikkarainen PH, Uusitupa M, I. Nutritional status of newly diagnosed celiac disease patients before and after the institution of a celiac disease diet--association with the grade of mucosal villous atrophy. Am J Clin Nutr 1998;67(3):482-7.

347. Amin R, Murphy N, Edge J, Ahmed ML, Acerini CL, Dunger DB. A longitudinal study of the effects of a gluten-free diet on glycemic control and weight gain in subjects with type 1 diabetes and celiac disease. Diabetes Care 2002;25(7):1117-22.

348. Bardella MT, Fredella C, Prampolini L, Molteni N, Giunta AM, Bianchi PA. Body composition and dietary intakes in adult celiac disease patients consuming a strict gluten-free diet. Am J Clin Nutr 2000;72(4):937-9.

349. Barera G, Mora S, Brambilla P, Ricotti A, Menni L, Beccio S, et al. Body composition in children with celiac disease and the effects of a gluten-free diet: a prospective case-control study. Am J Clin Nutr 2000;72(1):71-5.

350. Boersma B, Houwen RHJ, Blum WF, van Doorn J, Wit JM. Catch-up growth and endocrine changes in childhood celiac disease. Endocrine

changes during catch-up growth. Horm Res 2002;58(Suppl 1):57-65.

351. Ciacci C, Cirillo M, Auriemma G, Di Dato G, Sabbatini F, Mazzacca G. Celiac disease and pregnancy outcome. Am J Gastroenterol 1996;91(4):718-22.

352. Rea F, Polito C, Marotta A, Di Toro A, Iovene A, Collini R, et al. Restoration of body composition in celiac children after one year of gluten-free diet. J Pediatr Gastroenterol Nutr 1996;23(4):408-12.

353. Sategna-Guidetti C, Grosso SB, Grosso S, Mengozzi G, Aimo G, Zaccaria T, et al. The effects of 1-year gluten withdrawal on bone mass, bone metabolism and nutritional status in newly-diagnosed adult coeliac disease patients. Aliment Pharmacol Ther 2000;14(1):35-43.

354. Sategna-Guidetti C, Volta U, Ciacci C, Usai P, Carlino A, De Franceschi L, et al. Prevalence of thyroid disorders in untreated adult celiac disease patients and effect of gluten withdrawal: an Italian multicenter study. Am J Gastroenterol 2001;96(3):751-7.

355. Valdimarsson T, Lofman O, Toss G, Strom M. Reversal of osteopenia with diet in adult coeliac disease. Gut 1996;38(3):322-7.

356. Valdimarsson T, Toss G, Lofman O, Strom M. Three years' follow-up of bone density in adult coeliac disease: significance of secondary hyperparathyroidism. Scand J Gastroenterol 2000;35(3):274-80.

357. Westman E, Ambler GR, Royle M, Peat J, Chan A. Children with coeliac disease and insulin dependent diabetes mellitus--growth, diabetes control and dietary intake. J Pediatr Endocrinol Metab 1999;12(3):433-42.

358. Annibale B, Severi C, Chistolini A, Antonelli G, Lahner E, Marcheggiano A, et al. Efficacy of gluten-free diet alone on recovery from iron deficiency anemia in adult celiac patients. Am J Gastroenterol 2001;96(1):132-7.

359. Arato A, Korner A, Veres G, Dezsofi A, Ujpal I, Madacsy L. Frequency of coeliac disease in Hungarian children with type 1 diabetes mellitus. Eur J Pediatr 2002;162(1):1-5.

360. Atkinson K, Tokmakajian S, Watson W, Gregor J. Evaluation of the endomysial antibody for celiac disease: operating properties and associated cost implications in clinical practice. Can J Gastroenterol 1997;11(8):673-7.

361. Bardella MT, Molteni N, Quatrini M, Velio P, Ranzi T, Bianchi PA. Clinical, biochemical and histological abnormalities in adult celiac patients on gluten-free diet. Gastroenterol Clin Biol 1985;9(11):787-9.

362. Corrao G, Corazza GR, Bagnardi V, Brusco G, Ciacci C, Cottone M, et al. Mortality in patients with coeliac disease and their relatives: a cohort study. Lancet 2001;358(9279):356-61.

363. Fabiani E, Catassi C, Villari A, Gismondi P, Pierdomenico R, Ratsch IM, et al. Dietary compliance in screening-detected coeliac disease adolescents. Acta Paediatr 1996;412(Suppl):65-7.

364. Fabiani E, Taccari LM, Ratsch IM, Di Giuseppe S, Coppa G, V, Catassi C. Compliance with gluten-free diet in adolescents with screening-detected celiac disease: a 5-year follow-up study. J Pediatr 2000;136(6):841-3.

365. Greco L, Mayer M, Ciccarelli G, Troncone R, Auricchio S. Compliance to a gluten-free diet in adolescents, or "what do 300 coeliac adolescents eat every day?". Ital J Gastroenterol Hepatol 1997;29(4):305-10.

366. Harewood GC, Murray JA. Diagnostic approach to a patient with suspected celiac disease: a cost analysis. Dig Dis Sci 2001;46(11):2510-4.

367. Holmes GK, Stokes PL, Sorahan TM, Prior P, Waterhouse JA, Cooke WT. Coeliac disease, gluten-free diet, and malignancy. Gut 1976;17(8):612-9.

368. Johnston SD, Watson RG, McMillan SA, Sloan J, Love AH. Coeliac disease detected by screening is not silent--simply unrecognized. Qjm 1998;91(12):853-60.

369. Poddar U, Thapa BR, Nain CK, rasad A, ingh K. Celiac disease in India: are they true cases of celiac disease? J Pediatr Gastroenterol Nutr 2002;35(4):508-12.

370. Saukkonen T, Vaisanen S, Akerblom HK, Savilahti E. Coeliac disease in children and adolescents with type 1 diabetes: a study of growth, glycaemic control, and experiences of families. Acta Paediatr 2002;91(3):297-302.

371. Smecuol E, Gonzalez D, Mautalen C, Siccardi A, Cataldi M, Niveloni S, et al. Longitudinal study on the effect of treatment on body composition and anthropometry of celiac disease patients. Am J Gastroenterol 1997;92(4):639-43.

372. Bowron A, Moorghen M, Morgan JE, Osborne JR, Stansbie D, Stone JE. Cost-effective strategy for the serological investigation of coeliac disease. Ann Clin Biochem 2000;37(4):467-70.

373. Thomason K, West J, Logan RFA, Coupland C, Holmes GKT. Fracture experience of patients with coeliac disease: a population based survey. Gut 2003;52(4):518-22.

374. Addolorato G, Capristo E, Ghittoni G, Valeri C, Masciana R, Ancona C, et al. Anxiety but not depression decreases in coeliac patients after one-year gluten-free diet: a longitudinal study. Scand J Gastroenterol 2001;36(5):502-6.

375. Bai JC, Gonzalez D, Mautalen C, Mazure R, Pedreira S, Vazquez H, et al. Long-term effect of gluten restriction on bone mineral density of patients with coeliac disease. Aliment Pharmacol Ther 1997;11(1):157-64.

376. Kemppainen T, Kroger H, Janatuinen E, Arnala I, Lamberg-Allardt C, Karkkainen M, et al. Bone recovery after a gluten-free diet: a 5-year follow-up study. Bone 1999;25(3):355-60.

377. Mora S, Barera G, Beccio S, Menni L, Proverbio MC, Bianchi C, et al. A prospective, longitudinal study of the long-term effect of treatment on bone density in children with celiac disease. J Pediatr 2001;139(4):516-21.

378. Mustalahti K, Collin P, Sievanen H, Salmi J, Maki M. Osteopenia in patients with clinically silent coeliac disease warrants screening. Lancet 1999;354(9180):744-5.

379. Zaccari G, Mazzetti dP, Paone FM, Guidiceandrea P. A proposal for coeliac disease screening of all infants at the age of fifteen months. Gastroenterol Int 1996;9(1):11-5.

380. Bowron A, Moorghen M, Morgan JE, Osborne JR, Stansbie D, Stone JE. Cost-effective strategy for the serological investigation of coeliac disease. Ann Clin Biochem 2000;37(4):467-70.

381. Bernstein CN, Leslie WD, Leboff MS. AGA technical review on osteoporosis in gastrointestinal diseases. Gastroenterology 2003;124(3):795-841.

382. Gomez JC, Selvaggio G, Pizarro B, Viola MJ, La Motta G, Smecuol E, et al. Value of a screening algorithm for celiac disease using tissue transglutaminase antibodies as first level in a population-based study. American Journal of Gastroenterology 2002;97(11):2785-90.

383. Kemppainen T, Kroger H, Janatuinen E, Arnala I, Kosma VM, Pikkarainen P, et al. Osteoporosis in adult patients with celiac disease. Bone 1999;24(3):249-55.

384. Mora S, Barera G, Beccio S, Proverbio MC, Weber G, Bianchi C, et al. Bone density and bone metabolism are normal after long-term gluten-free diet in young celiac patients. Am J Gastroenterol 1999;94(2):398-403.

385. West J, Logan Richard FA, Card TR, Smith C, Hubbard R. Fracture risk in people with celiac disease: a population-based cohort study. Gastroenterology 2003;125(2):429-36.

386. Ciacci C, Maurelli L, Klain M, Savino G, Salvatore M, Mazzacca G, et al. Effects of dietary treatment on bone mineral density in adults with celiac disease: factors predicting response. Am J Gastroenterol 1997;92(6):992-6.

387. McFarlane XA, Bhalla AK, Robertson DA. Effect of a gluten free diet on osteopenia in adults with newly diagnosed coeliac disease. Gut 1996;39(2):180-4.

388. Vestergaard P, Mosekilde L. Fracture risk in patients with celiac Disease, Crohn's disease, and ulcerative colitis: a nationwide follow-up study of 16,416 patients in Denmark. Am J Epidemiol 2002;156(1):1-10. Available: PM:12076883.

389. Vasquez H, Mazure R, Gonzalez D, Flores D, Pedreira S, Niveloni S, et al. Risk of fractures in celiac disease patients: a cross-sectional, case-control study. Am J Gastroenterol 2000;95(1):183-9. Available: PM:10638580.

390. Fickling WE, McFarlane XA, Bhalla AK, Robertson DA. The clinical impact of metabolic bone disease in coeliac disease. Postgrad Med J 2001;77(903):33-6. Available: PM:11123392.

391. Bianchi ML, Bardella MT. Bone and celiac disease. Calcif Tissue Int 2002;71(6):465-71. Available: PM:12232681.

392. Moreno ML, Vazquez H, Mazure R, Smecuol E, Niveloni S, Pedreira S, et al. Stratification of bone fracture risk in patients with celiac disease. Clin Gastroenterol Hepatol 2004;2(2):127-34. Available: PM:15017617.

393. Norgard B, Fonager K, Sorensen HT, Olsen J. Birth outcomes of women with celiac disease: A nationwide historical cohort study. Am J Gastroenterol 1999;94(9):2435-40.

394. Thomason K, West J, Logan RFA, Coupland C, Holmes GKT. Fracture experience of patients with coeliac disease: a population based survey. Gut 2003;52(4):518-22.

395. Cook HB, Burt MJ, Collett JA, Whitehead MR, Frampton CM, Chapman BA. Adult coeliac disease: prevalence and clinical significance. J Gastroenterol Hepatol 2000;15(9):1032-6.

396. Bartholomeusz RC, Labrooy JT, Davidson GP, Hetzel P, Johnson RB, Shearman DJ. Polymeric IgA antibody to gliadin in the serum of patients with coeliac disease. J Gastroenterol Hepatol 1990;5(6):675-81.

397. Fotoulaki M, Nousia-Arvanitakis S, Augoustidou-Savvopoulou P, Kanakoudi-Tsakalides F, Zaramboukas T, Vlachonikolis J. Clinical application of immunological markers as monitoring tests in celiac disease. Dig Dis Sci 1999;44(10):2133-8.

398. Kaukinen K, Sulkanen S, Maki M, Collin P. IgA-class transglutaminase antibodies in evaluating the efficacy of gluten-free diet in coeliac disease. Eur J Gastroenterol Hepatol 2002;14(3):311-5.

399. Bardella MT, Trovato C, Cesana BM, Pagliari C, Gebbia C, Peracchi M. Serological markers for coeliac disease: Is it time to change? Dig Liver Dis 2001;33(5):426-31.

400. Burgin-Wolff A, Gaze H, Hadziselimovic F, Huber H, Lentze MJ, Nussle D, et al. Antigliadin and antiendomysium antibody determination for coeliac disease. Arch Dis Child 1991;66(8):941-7.

401. Scalici C, Manzoni D, Licastro G, Varia F, Di Prima L, Vitali R. Reliability of EMA assay in the evaluation of gluten-free diet compliance in celiac patients during follow-up. Acta Med Mediterr 2003;19(1):67-9.

402. Vahedi K, Mascart-Lemone F, Mary JY, Laberenne JE, Bouhnik Y, Morin MC, et al. Are Anti-Endomysial (AEM) and Anti-Transglutaminase (TTG) Antibodies Reliable Markers of Strict Diet Compliance in Adult Celiacs on a Gluten Free Diet (GFD)? Gastroenterology 2000;118(4 Suppl. 2 Part 1):AGA.

403. Martini S, Mengozzi G, Aimo G, Giorda L, Pagni R, Guidetti CS. Comparative evaluation of serologic tests for celiac disease diagnosis and follow-up. Clin Chem 2002;48(6 Pt 1):960-3.

404. Valletta EA, Trevisiol D, Mastella G. IgA anti-gliadin antibodies in the monitoring of gluten challenge in celiac disease. J Pediatr Gastroenterol Nutr 1990;10(2):169-73.

405. Wahab PJ, Meijer Jos WR, Mulder Chris JJ. Histologic follow-up of people with celiac disease on a gluten-free diet: slow and incomplete recovery. American Journal of Clinical Pathology 2002;118(3):459-63.

406. McNicholl B, Egan-Mitchell B, Stevens F, Keane R, Baker S, McCarthy CF, et al. Mucosal recovery in treated childhood celiac disease (gluten-sensitive enteropathy). J Pediatr 1976;89(3):418-24.

407. Valentini RA, Andreani ML, Corazza GR, Gasbarrini G. IgA endomysium antibody: a valuable tool in the screening of coeliac disease but not its follow-up. Ital J Gastroenterol 1994;26(6):279-82.

408. Fabiani E, Catassi C. The serum IgA class anti-tissue transglutaminase antibodies in the diagnosis and follow up of coeliac disease. Results of an international multi-centre study. International Working Group on Eu-tTG. Eur J Gastroenterol Hepatol 2001;13(6):659-65.

409. Dickey W, Hughes DF, McMillan SA. Disappearance of endomysial antibodies in treated celiac disease does not indicate histological recovery. Am J Gastroenterol 2000;95(3):712-4.

410. Fabiani E, Catassi C, Villari A, Gismondi P, Pierdomenico R, Ratsch IM, et al. Dietary compliance in screening-detected coeliac disease adolescents. Acta Paediatr 1996;412:65-7.

411. Lee SK, Lo W, Memeo L, Rotterdam H, Green Peter HR. Duodenal histology in patients with celiac disease after treatment with a gluten-free diet. Gastrointest Endosc 2003;57(2):187-91.

412. Pacht A, Sinai N, Hornstein L, Kumar V, Ish-Shalom N, Lerner A. The diagnostic reliability of anti-endomysial antibody in celiac disease: the north Israel experience. Isr J Med Sci 1995;31(4):218-20.

413. Sategna-Guidetti C, Grosso S, Bruno M, Grosso SB. Reliability of immunologic markers of celiac sprue in the assessment of mucosal recovery after gluten withdrawal. J Clin Gastroenterol 1996;23(2):101-4.

414. Selby WS, Painter D, Collins A, Faulkner-Hogg KB, Loblay RH. Persistent mucosal abnormalities in coeliac disease are not related to the ingestion of trace amounts of gluten. Scand J Gastroenterol 1999;34(9):909-14.

415. Troncone R, Mayer M, Spagnuolo F, Maiuri L, Greco L. Endomysial antibodies as unreliable markers for slight dietary transgressions in adolescents with celiac disease. J Pediatr Gastroenterol Nutr 1995;21(1):69-72.

416. Anson O, Weizman Z, Zeevi N. Celiac disease: parental knowledge and attitudes of dietary compliance. Pediatrics 1990;85(1):98-103.

417. Fabiani E, Taccari LM, Ratsch IM, Di Giuseppe S, Coppa G, V, Catassi C. Compliance with gluten-free diet in adolescents with screening-detected celiac disease: a 5-year follow-up study. J Pediatr 2000;136(6):841-3.

418. Jackson PT, Glasgow JF, Thom R. Parents' understanding of coeliac disease and diet. Arch Dis Child 1985;60(7):672-4.

419. Lamontagne P, West GE, Galibois I. Quebecers with celiac disease: analysis of dietary problems. Canadian Journal of Dietetic Practice and Research - a Publication of Dietitians of Canada = Revue Canadienne De La Pratique Et De La Recherche En Dietetique - Une Publication Des Dietetistes Du C 2001;62(4):175-81.

420. Ljungman G, Myrdal U. Compliance in teenagers with coeliac disease--a Swedish follow-up study. Acta Paediatr 1993;82(3):235-8.

421. Hogberg L, Grodzinsky E, Stenhammar L. Better dietary compliance in patients with coeliac disease diagnosed in early childhood. Scand J Gastroenterol 2003;38(7):751-4.

422. Patin NM, Johnson F, Kirks BA. Promoting dietary compliance in gluten-intolerant children. J Nutr Educ 1989;21(2):100D.

423. Lijmer JG, Mol BW, Heisterkamp S, Bonsel GJ, Prins MH, van der Meulen JH, et al. Empirical evidence of design-related bias in studies of diagnostic tests. JAMA 1999;282(11):1061-6. Available: PM:10493205.

424. Tursi A, Brandimarte G, Giorgetti GM. Prevalence of antitissue transglutaminase antibodies in different degrees of intestinal damage in celiac disease. Journal of Clinical Gastroenterology 2003;36(3):219-21.

425. Tursi A, Brandimarte G, Giorgetti G, Gigliobianco A, Lombardi D, Gasbarrini G. Low prevalence of antigliadin and anti-endomysium antibodies in subclinical/silent celiac disease. American Journal of Gastroenterology 2001;96(5):1507-10.

426. Tursi A, Brandimarte G, Giorgetti GM. Sorbitol H2-breath test versus anti-endomysium antibodies for the diagnosis of subclinical/silent coeliac disease. Scand J Gastroenterol 2001;36(11):1170-2.

427. Demir H, Yuce A, Kocak N, Ozen H, Gurakan F. Celiac disease in Turkish children: presentation of 104 cases. Pediatrics International - Official Journal of the Japan Pediatric Society 2000;42(5):483-7.

428. Kotze Lorete Maria da Silva, Utiyama Shirley Ramos da Rosa, Nisihara RM, de C, V, Ioshii SO. IgA class anti-endomysial and anti-tissue transglutaminase antibodies in relation to duodenal mucosa changes in coeliac disease. Pathology 2003;35(1):56-60.

429. Sategna-Guidetti C, Bruno M, Pulitano R, Ferfoglia G. Disease specificity of IgA class anti-endomysium antibodies (IgA-EmA) in adult coeliac disease. Eur J Gastroenterol Hepatol 1991;3(3):251-4.

430. Ciacci C, Cavallaro R, Della VN, D'argenio G. The Use of Serum Ttg-Ab Assay in Patients on Gluten-Free Diet as a Measure of Dietetic Compliance. Gastroenterology 2002;122(2):588.

431. Garcia VE, De Lourdes de Abreu Ferrari, Alves BE, Affonso Barbosa AJ, Brasileiro FG, Hartung TN, et al. Agreement between pathologists concerning assessment of intestinal biopsies from adult celiac disease patients. Gastroenterol Int 2002;15(1-2):1-8.

432. Weile B, Hansen BF, Hagerstrand I, Hansen JP, Krasilnikoff PA. Interobserver variation in diagnosing coeliac disease. A joint study by Danish and Swedish pathologists. Apmis - Acta Pathologica, Microbiologica, Et Immunologica Scandinavica 2000;108(5):380-4.

433. Landis JR, Koch GG. The measurement of observer agreement for categorical data. Biometrics 1977;33(1):159-74. Available: PM:843571.

434. Risdon RA, Keeling JW. Quantitation of the histological changes found in small intestinal biopsy specimens from children with suspected coeliac disease. Gut 1974;15(1):9-18.

435. Glasgow JF, Corkey CW, Molla A. Critical assessment of small bowel biopsy in children. Archives of Disease in Childhood 1979;54(8):604-8.

436. Jarvinen TT, Kaukinen K, Laurila K, Kyronpalo S, Rasmussen M, Maki M, et al. Intraepithelial

lymphocytes in celiac disease. American Journal of Gastroenterology 2003;98(6):1332-7.

437. Corazza GR, Bonvicini F, Frazzoni M, Gatto M, Gasbarrini G. Observer variation in assessment of jejunal biopsy specimens. A comparison between subjective criteria and morphometric measurement. Gastroenterology 1982;83(6):1217-22. Available: PM:7129029.

438. Stenhammar L, Brandt A, Wagermark J. A family study of coeliac disease. Acta Paediatr Scand 1982;71(4):625-8. Available: PM:7136679.

439. Troncone R, Catassi C, Lambertini A, Zaniboni MG, Lazzari R, Bottaro G, et al. Latent coeliac disease in Italy. Acta Paediatr Int J Paediatr 1995;84(11):1252-7.

440. Maki M, Lahdeaho ML, Hallstrom O, Viander M, Visakorpi JK. Postpubertal gluten challenge in coeliac disease. Arch Dis Child 1989;64(11):1604-7. Available: PM:2604420.

441. Schmitz J, Arnaud-Battandier F JJ, Rey J. Long term follow-up of childhood coeliac disease (CD): Is there a "natural recovery"? Pediatr Res 1984;18;1052.

442. Ferguson A, McClure JP, Townley RR. Intraepithelial lymphocyte counts in small intestinal biopsies from children with diarrhoea. Acta Paediatrica Scandinavica 1976;65(5):541-6.

443. Kutlu T, Brousse N, Rambaud C, Le Deist F, Schmitz J, Cerf-Bensussan N. Numbers of T cell receptor (TCR) alpha beta+ but not of TcR gamma delta+ intraepithelial lymphocytes correlate with the grade of villous atrophy in coeliac patients on a long term normal diet. Gut 1993;34(2):208-14.

444. O'Farrelly C, Graeme-Cook F, Hourihane DO, Feighery C, Weir DG. Histological changes associated with wheat protein antibodies in the absence of villous atrophy. Journal of Clinical Pathology 1987;40(10):1228-30.

445. Saputo V, Losi S, Mancosu M, Della Morte MA, Moschettini GF. Intraepithelial lymphocytes in jejunal mucosa: diagnostic significance of changes in their number during chronic intestinal disease, with particular reference to coeliac disease. Schweizerische Rundschau Fur Medizin Praxis = Revue Suisse De Medecine Praxis 1981;70(30):1342-8.

446. Mino M, Lauwers GY. Role of lymphocytic immunophenotyping in the diagnosis of gluten-sensitive enteropathy with preserved villous architecture. Am J Surg Pathol 2003;27(9):1237-42.

447. Goldstein NS, Underhill J. Morphologic features suggestive of gluten sensitivity in architecturally normal duodenal biopsy specimens. American Journal of Clinical Pathology 2001;116(1):63-71.

448. Kuitunen P, Kosnai I, Savilahti E. Morphometric study of the jejunal mucosa in various childhood enteropathies with special reference to intraepithelial lymphocytes. Journal of Pediatric Gastroenterology and Nutrition 1982;1(4):525-31.

449. Kaukinen K, Collin P, Holm K, Karvonen AL, Pikkarainen P, Maki M. Small-bowel mucosal inflammation in reticulin or gliadin antibody-positive patients without villous atrophy. Scandinavian Journal of Gastroenterology 1998;33(9):944-9.

450. Wahab PJ, Crusius JB, Meijer JW, Mulder CJ. Gluten challenge in borderline gluten-sensitive enteropathy. American Journal of Gastroenterology 2001;96(5):1464-9.

451. Wahab PJ, Meijer Jos WR, Dumitra D, Goerres MS, Mulder Chris JJ. Coeliac disease: more than villous atrophy. Romanian Journal of Gastroenterology 2002;11(2):121-7.

452. Mahadeva S, Wyatt J, I, Howdle PD. Is a raised intraepithelial lymphocyte count with normal duodenal villous architecture clinically relevant? Journal of Clinical Pathology 2002;55(6):424-8.

453. Kaukinen K, Maki M, Partanen J, Sievanen H, Collin P. Celiac disease without villous atrophy: revision of criteria called for. Dig Dis Sci 2001;46(4):879-87.

454. Catassi C, Fabiani E, Corrao G, Barbato M, De Renzo A, Carella AM, et al. Risk of non-Hodgkin lymphoma in celiac disease. JAMA 2002;287(11):1413-9.

455. Johnston SD, Watson RGP. Small bowel lymphoma in unrecognized coeliac disease: A cause for concern? Eur J Gastroenterol Hepatol 2000;12(6):645-8.

456. Green PHR, Stavropoulos SN, Panagi SG, Goldstein SL, Mcmahon DJ, Absan H, et al. Characteristics of adult celiac disease in the USA: results of a national survey. Am J Gastroenterol 2001;96(1):126-31.

457. Wahab PJ, Meijer Jos WR, Mulder CJJ. Histologic follow-up of people with celiac

disease on a gluten-free diet: slow and incomplete recovery. Am J Clin Pathol 2002;118(3):459-63.

458. Goerres MS, Meijer JWR, Wahab PJ, Kerckhaert JAM, Groenen PJTA, Van Krieken JHJM, et al. Azathioprine and prednisone combination therapy in refractory coeliac disease. Aliment Pharmacol Ther 2003;18(5):487-94.

459. Cellier C, Delabesse E, Helmer C, Patey N, Matuchansky C, Jabri B, et al. Refractory sprue, coeliac disease, and enteropathy-associated T-cell lymphoma. French Coeliac Disease Study Group. Lancet 2000;356(9225):203-8.

460. Ciacci C, Cirillo M, Cavallaro R, Mazzacca G. Long-term follow-up of celiac adults on gluten-free diet: prevalence and correlates of intestinal damage. Digestion 2002;66(3):178-85.

461. Kluge F, Koch HK, Grosse-Wilde H, Lesch R, Gerok W. Follow-up of treated adult celiac disease: clinical and morphological studies. Hepatogastroenterology 1982;29(1):17-23.