References

Acker DB, Sachs BP, Friedman EA. Risk factors for shoulder dystocia. Obstet Gynecol 1985;66(6):762-8.

ACOG. Routine ultrasound in low-risk pregnancy. 1997.

ACOG. Fetal macrosomia. ACOG Practice Bulletin No. 22. 2000.

Alfirevic Z, Howarth G, Gaussmann A. Oral misoprostol for induction of labor (Cochrane Review). In: The Cochrane Library, Issue 2, 2000. Oxford: Update Software.

Alfirevic Z, Luckas M, Walkinshaw SA, et al. A randomised comparison between amniotic fluid index and maximum pool depth in the monitoring of post-term pregnancy. Br J Obstet Gynaecol 1997;104(2):207-11.

Alfirevic Z, Walkinshaw SA. A randomised controlled trial of simple compared with complex antenatal fetal monitoring after 42 weeks of gestation. Br J Obstet Gynaecol 1995;102(8):638-43.

Allott HA, Palmer CR. Sweeping the membranes: a valid procedure in stimulating the onset of labour? Br J Obstet Gynaecol 1993:100(10):898-903.

Anonymous. ACOG practice patterns. Management of postterm pregnancy. Number 6, October 1997. American College of Obstetricians and Gynecologists. Int J Gynaecol Obstet 1997;60(1):86-91.

Arabin B, Snyjders R, Mohnhaupt A, et al. Evaluation of the fetal assessment score in pregnancies at risk for intrauterine hypoxia. Am J Obstet Gynecol 1993;169(3):549-54.

Arias F. Predictability of complications associated with prolongation of pregnancy. Obstet Gynecol 1987;70(1):101-6.

Atad J, Hallak M, Auslender R, et al. A randomized comparison of prostaglandin E2, oxytocin, and the double-balloon device in inducing labor. Obstet Gynecol 1996;87(2):223-7.

Augensen K, Bergsjø P, Eikeland T, et al. Randomised comparison of early versus late induction of labour in post-term pregnancy. BMJ 1987;294(6581):1192-5.

Ayres-de-Campos D, Bernardes J, Costa-Pereira A, et al. Inconsistencies in classification by experts of cardiotocograms and subsequent clinical decision. Br J Obstet Gynaecol 1999;106(12):1307-10.

Bastian H, Keirse MJ, Lancaster PA. Perinatal death associated with planned home birth in Australia: population based study. BMJ 1998;317(7155):384-8.

Battaglia C, Larocca E, Lanzani A, et al. Doppler velocimetry in prolonged pregnancy. Obstet Gynecol 1991;77(2):213-6.

Bell RJ, Permezel M, MacLennan A, et al. A randomized, double-blind, placebo-controlled trial of the safety of vaginal recombinant human relaxin for cervical ripening. Obstet Gynecol 1993;82(3):328-33.

Bennett BB. Uterine rupture during induction of labor at term with intravaginal misoprostol. Obstet Gynecol 1997;89(5 Pt 2):832-3.

Berghella V, Rogers RA, Lescale K. Stripping of membranes as a safe method to reduce prolonged pregnancies. Obstet Gynecol 1996;87(6):927-31.

Bergsjø P, Huang GD, Yu SQ, et al. Comparison of induced versus non-induced labor in post-term pregnancy. A randomized prospective study. Acta Obstet Gynecol Scand 1989;68(8):683-7.

Bernardes J, Costa-Pereira A, Ayres-de-Campos D, et al. Evaluation of interobserver agreement of cardiotocograms. Int J Gynaecol Obstet 1997;57(1):33-7.

Bishop EH. Pelvic scoring for elective induction. Obstet Gynecol 1964;24:266-8.

Blanchette HA, Nayak S, Erasmus S. Comparison of the safety and efficacy of intravaginal misoprostol (prostaglandin E1) with those of dinoprostone (prostaglandin E2) for cervical ripening and induction of labor in a community hospital. Am J Obstet Gynecol 1999;180(6 Pt 1):1551-9.

Bochner CJ, Medearis AL, Ross MG, et al. The role of antepartum testing in the management of postterm pregnancies with heavy meconium in early labor. Obstet Gynecol 1987;69(6):903-7.

Bochner CJ, Williams J 3d, Castro L, et al. The efficacy of starting postterm antenatal testing at 41 weeks as compared with 42 weeks of gestational age. Am J Obstet Gynecol 1988;159(3):550-4.

Boulvain M, Fraser WD, Marcoux S, et al. Does sweeping of the membranes reduce the need for formal induction of labour? A randomised controlled trial. Br J Obstet Gynaecol 1998;105(1):34-40.

Boulvain M, Irion O. Stripping/sweeping the membranes for inducing labour or preventing post-term pregnancy (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Brar HS, Horenstein J, Medearis AL, et al. Cerebral, umbilical, and uterine resistance using Doppler velocimetry in postterm pregnancy. J Ultrasound Med 1989;8(4):187-91.

Brennand JE, Calder AA, Leitch CR, et al. Recombinant human relaxin as a cervical ripening agent. Br J Obstet Gynaecol 1997;104(7):775-80.

Brown VA, Sawers RS, Parsons RJ, et al. The value of antenatal cardiotocography in the management of high-risk pregnancy: a randomized controlled trial. Br J Obstet Gynaecol 1982;89(9):716-22.

Buser D, Mora G, Arias F. A randomized comparison between misoprostol and dinoprostone for cervical ripening and labor induction in patients with unfavorable cervices. Obstet Gynecol 1997;89(4):581-5.

Buttino LT, Garite TJ. Intracervical prostaglandin in postdate pregnancy. A randomized trial. J Reprod Med 1990;35(2):155-8.

Cammu H, Haitsma V. Sweeping of the membranes at 39 weeks in nulliparous women: a randomised controlled trial. Br J Obstet Gynaecol 1998;105(1):41-4.

Campbell MK, Ostbye T, Irgens LM. Post-term birth: Risk factors and outcomes in a 10-year cohort of Norwegian births. Obstet Gynecol 1997;89(4):543-8.

Cardozo L, Fysh J, Pearce JM. Prolonged pregnancy: the management debate. BMJ 1986;293(6554):1059-63.

Chang CH, Chang FM. Randomized comparison of misoprostol and dinoprostone for preinduction cervical ripening and labor induction. J Formos Med Assoc 1997;96(5):366-9.

Chatterjee MS, Ramchandran K, Ferlita J, et al. Prostaglandin E2 (PGE2) vaginal gel for cervical ripening. Eur J Obstet Gynecol Reprod Biol 1991;38(3):197-202.

Chauhan SP, Sullivan CA, Lutton TC, et al. Parous patients' estimate of birth weight in postterm pregnancy. J Perinatol 1995;15(3):192-4.

Chauhan SP, Sullivan CA, Magann EF, et al. Estimate of birthweight among post-term pregnancy: Clinical versus sonographic. J Matern Fetal Med 1994;3(5):208-11.

Chayen B, Tejani N, Verma U. Induction of labor with an electric breast pump. J Reprod Med 1986;31(2):116-8.

Chervenak JL, Divon MY, Hirsch J, et al. Macrosomia in the postdate pregnancy: is routine ultrasonographic screening indicated? Am J Obstet Gynecol 1989;161(3):753-6.

Chuck FJ, Huffaker BJ. Labor induction with intravaginal misoprostol versus intracervical prostaglandin E2 gel (Prepidil gel): randomized comparison. Am J Obstet Gynecol 1995;173(4):1137-42.

Cotzias CS, Paterson-Brown S, Fisk NM. Prospective risk of unexplained stillbirth in singleton pregnancies at term: population based analysis. BMJ 1999;319(7205):287-8.

Crane J, Bennett K, Young D, et al. The effectiveness of sweeping membranes at term: a randomized trial. Obstet Gynecol 1997;89(4):586-90.

Crowley P. Interventions for preventing or improving the outcome of delivery at or beyond term (Cochrane Review). In: The Cochrane Library, Issue 2, 2000. Oxford: Update Software.

Crowley P, O'Herlihy C, Boylan P. The value of ultrasound measurement of amniotic fluid volume in the management of prolonged pregnancies. Br J Obstet Gynaecol 1984;91(5):444-8.

Curtis PD, Matthews TG, Clarke TA, et al. Neonatal seizures: the Dublin Collaborative Study. Arch Dis Child 1988;63(9):1065-8.

Devoe LD, Sholl JS. Postdates pregnancy. Assessment of fetal risk and obstetric management. J

Reprod Med 1983;28(9):576-80.

Divon MY, Haglund B, Nisell H, et al. Fetal and neonatal mortality in the postterm pregnancy: the impact of gestational age and fetal growth restriction. Am J Obstet Gynecol 1998;178(4):726-31.

Doany W, McCarty J. Outpatient management of the uncomplicated postdate pregnancy with intravaginal prostaglandin E2 gel and membrane stripping. J Matern Fetal Med 1997;6(2):71-8.

Donker DK, van Geijn HP, Hasman A. Interobserver variation in the assessment of fetal heart rate recordings. Eur J Obstet Gynecol Reprod Biol 1993;52(1):21-8.

Dyson DC, Miller PD, Armstrong MA. Management of prolonged pregnancy: induction of labor versus antepartum fetal testing. Am J Obstet Gynecol 1987;156(4):928-34.

Eden RD, Gergely RZ, Schifrin BS, et al. Comparison of antepartum testing schemes for the management of the postdate pregnancy. Am J Obstet Gynecol 1982;144(6):683-92.

Eden RD, Seifert LS, Winegar A, et al. Perinatal characteristics of uncomplicated postdate pregnancies. Obstet Gynecol 1987;69(3 Pt 1):296-9.

Egarter C, Kofler E, Fitz R, et al. Is induction of labor indicated in prolonged pregnancy? Results of a prospective randomised trial. Gynecol Obstet Invest 1989;27(1):6-9.

El-Torkey M, Grant JM. Sweeping of the membranes is an effective method of induction of labour in prolonged pregnancy: a report of a randomized trial. Br J Obstet Gynaecol 1992;99(6):455-8.

Elliott CL, Brennand JE, Calder AA. The effects of mifepristone on cervical ripening and labor induction in primigravidae. Obstet Gynecol 1998;92(5):804-9.

Elliott JP, Flaherty JF. The use of breast stimulation to prevent postdate pregnancy. Am J Obstet Gynecol 1984;149(6):628-32.

Escudero F, Contreras H. A comparative trial of labor induction with misoprostol versus oxytocin. Int J Gynaecol Obstet 1997;57(2):139-43.

Evans MI, Dougan MB, Moawad AH, et al. Ripening of the human cervix with porcine ovarian relaxin. Am J Obstet Gynecol 1983;147(4):410-4.

Farah LA, Sanchez-Ramos L, Rosa C, et al. Randomized trial of two doses of the prostaglandin E1 analog misoprostol for labor induction. Am J Obstet Gynecol 1997;177(2):364-9.

Farmakides G, Schulman H, Winter D, et al. Prenatal surveillance using nonstress testing and Doppler velocimetry. Obstet Gynecol 1988;71(2):184-7.

Fleischer A, Schulman H, Farmakides G, et al. Antepartum nonstress test and the postmature pregnancy. Obstet Gynecol 1985;66(1):80-3.

Fletcher H, Mitchell S, Frederick J, et al. Intravaginal misoprostol versus dinoprostone as cervical ripening and labor-inducing agents. Obstet Gynecol 1994;83(2):244-7.

Fletcher HM, Mitchell S, Simeon D, et al. Intravaginal misoprostol as a cervical ripening agent. Br J Obstet Gynaecol 1993;100(7):641-4.

Flynn AM, Kelly J, Mansfield H, et al. A randomized controlled trial of non-stress antepartum cardiotocography. Br J Obstet Gynaecol 1982;89(6):427-33.

Froen JF, Arnestad M, Frey K, et al. Risk factors for sudden intrauterine unexplained death: epidemiologic characteristics of singleton cases in Oslo, Norway, 1986-1995. American Journal of Obstetrics & Gynecology 2001;184(4):694-702.

Frydman R, Lelaidier C, Baton-Saint-Mleux C, et al. Labor induction in women at term with mifepristone (RU 486): a double-blind, randomized, placebo-controlled study. Obstet Gynecol 1992;80(6):972-5.

Garry D, Figueroa R, Guillaume J, et al. Use of castor oil in pregnancies at term. Altern Ther Health Med 2000;6(1):77-9.

Giacalone PL, Targosz V, Laffargue F, et al. Cervical ripening with mifepristone before labor induction: A randomized study. Obstet Gynecol 1998;92(4 I):487-92.

Gilby JR, Williams MC, Spellacy WN. Fetal abdominal circumference measurements of 35 and 38 cm as predictors of macrosomia. J Reprod Med 2000;45:936-8.

Gilson GJ, O'Brien ME, Vera RW, et al. Prolonged pregnancy and the biophysical profile. A birthing center perspective. J Nurse Midwifery 1988;33(4):171-7.

Gjessing HK, Skjaerven R, Wilcox AJ. Errors in gestational age: evidence of bleeding early in pregnancy. Am J Public Health 1999;89(2):213-8.

Goeree R, Hannah M, Hewson S. Cost-effectiveness of induction of labour versus serial antenatal monitoring in the Canadian Multicentre Postterm Pregnancy Trial. CMAJ 1995;152(9):1445-50.

Goldberg J, Newman RB, Rust PF. Interobserver reliability of digital and endovaginal ultrasonographic cervical length measurements. Am J Obstet Gynecol 1997;177(4):853-8.

Gonen O, Rosen DJ, Dolfin Z, et al. Induction of labor versus expectant management in macrosomia: a randomized study. Obstet Gynecol 1997;89(6):913-7.

Gottschall DS, Borgida AF, Mihalek JJ, et al. A randomized clinical trial comparing misoprostol with prostaglandin E2 gel for preinduction cervical ripening. Am J Obstet Gynecol 1997;177(5):1067-70.

Graves GR, Baskett TF, Gray JH, et al. The effect of vaginal administration of various doses of prostaglandin E2 gel on cervical ripening and induction of labor. Am J Obstet Gynecol 1985;151(2):178-81.

Grünberger W, Spona J. The effect of pericervical PGEinf 2 instillation on levels of maternal serum 13,14 dihydro-15-keto-PGF(2alpha) and progesterone. Arch Gynecol 1986;239(2):93-9.

Gupta R, Vasishta K, Sawhney H, et al. Safety and efficacy of stripping of membranes at term. Int J Gynaecol Obstet 1998;60(2):115-21.

Hann L, McArdle C, Sachs B. Sonographic biophysical profile in the postdate pregnancy. J Ultrasound Med 1987;6(4):191-5.

Hannah ME, Hannah WJ, Hellmann J, et al. Induction of labor as compared with serial antenatal monitoring in post-term pregnancy. A randomized controlled trial. The Canadian Multicenter Post-term Pregnancy Trial Group [published erratum appears in N Engl J Med 1992 Jul 30;327(5):368]. N Engl J Med 1992;326(24):1587-92.

Hannah ME, Huh C, Hewson SA, et al. Postterm pregnancy: putting the merits of a policy of induction of labor into perspective. Birth 1996;23(1):13-9.

Harris BA, Huddleston JF, Sutliff G, et al. The unfavorable cervix in prolonged pregnancy. Obstet Gynecol 1983;62(2):171-4.

Hedén L, Ingemarsson I, Ahlström H, et al. Induction of labor versus conservative management in prolonged pregnancy: controlled study. Int J Feto-Matern Med 1991;4(4):231-6.

Henriksen TB, Wilcox AJ, Hedegaard M, et al. Bias in studies of preterm and postterm delivery due to ultrasound assessment of gestational age. Epidemiology 1995;6(5):533-7.

Herabutya Y, Prasertsawat P, Pokpirom J. A comparison of intravaginal misoprostol and intracervical prostaglandin E2 gel for ripening of unfavorable cervix and labor induction. J Obstet Gynaecol Res 1997;23(4):369-74.

Herabutya Y, Prasertsawat PO, Tongyai T, et al. Prolonged pregnancy: the management dilemma. Int J Gynaecol Obstet 1992;37(4):253-8.

Hilder L, Costeloe K, Thilaganathan B. Prolonged pregnancy: evaluating gestation-specific risks of fetal and infant mortality. Br J Obstet Gynaecol 1998;105(2):169-73.

Hofmeyr GJ, Gulmezoglu AM. Vaginal misoprostol for cervical ripening and labour induction in late pregnancy (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Holcomb WL, Smeltzer JS. Cervical effacement: variation in belief among clinicians. Obstet Gynecol 1991;78(1):43-5.

Howarth GR, Funk M, Steytler P, et al. A randomised controlled trial comparing vaginally administered misoprostol to vaginal dinoprostone gel in labour induction. J Obstet Gynaecol 1996;16:474-8.

Idrisa A, Obisesan KA, Adeleye JA. Fetal membrane sweeping for stimulation of labour in prolonged pregnancy: a controlled study. J Obstet Gynaecol 1993;13(4):235-7.

Imai M, Tani A, Saito M, et al. Significance of fetal fibronectin and cytokine measurement in the cervicovaginal secretions of women at term in predicting term labor and post-term pregnancy. Eur J Obstet Gynecol Reprod Biol 2001;97:53-8.

Jazayeri A, Heffron JA, Phillips R, et al. Macrosomia prediction using ultrasound fetal abdominal circumference of 35 centimeters or more. Obstet Gynecol 1999;93(4):523-6.

Johanson RB, Menon BKV. Vacuum extraction vs forcepts for assisted vaginal delivery (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Jüni P, Altman DG, Egger M. Assessing the quality of controlled clinical trials. BMJ 2001;323:42-6.

Jüni P, Witschi A, Bloch R, et al. The hazards of scoring the quality of clinical trials for meta-analysis. JAMA 1999;282(11):1054-60.

Kadanali S, Küçüközkan T, Zor N, et al. Comparison of labor induction with misoprostol vs. oxytocin/prostaglandin E2 in term pregnancy. Int J Gynaecol Obstet 1996;55(2):99-104.

Kadar N, Tapp A, Wong A. The influence of nipple stimulation at term on the duration of pregnancy. J Perinatol 1990;10(2):164-6.

Katz Z, Yemini M, Lancet M, et al. Non-aggressive management of post-date pregnancies. Eur J Obstet Gynecol Reprod Biol 1983;15(2):71-9.

Kelly AJ, Kavanagh J, Thomas J. Castor oil, bath and/or enema for cervical priming and induction of labour (Cochrane Review). In: The Cochrane Library, Issue 2, 2001. Oxford: Update Software.

Kemp B, Winkler M, Rath W. Induction of labor by prostaglandin E2 in relation to the Bishop score. Int J Gynecol Obstet 2000;71:13-7.

Keng MTY, Eng BTS. Oral contraceptives postdatism as a complication. Med J Malaysia 1982;37(4):338-43.

Kidd LC, Patel NB, Smith R. Non-stress antenatal cardiotocography—a prospective randomized clinical trial. Br J Obstet Gynaecol 1985;92(11):1156-9.

Knox GE, Huddleston JF, Flowers CE. Management of prolonged pregnancy: results of a prospective randomized trial. Am J Obstet Gynecol 1979;134(4):376-84.

Kramer RL, Gilson GJ, Morrison DS, et al. A randomized trial of misoprostol and oxytocin for induction of labor: safety and efficacy. Obstet Gynecol 1997;89(3):387-91.

Lee HY. A randomised double-blind study of vaginal misoprostol vs dinoprostone for cervical ripening and labour induction in prolonged pregnancy. Singapore Med J 1997;38(7):292-4.

Leveno KJ, Quirk JG, Cunningham FG, et al. Prolonged pregnancy. I. Observations concerning the causes of fetal distress. Am J Obstet Gynecol 1984;150(5 Pt 1):465-73.

Lidegaard O, Bottcher LM, Weber T. Description, evaluation and clinical decision making according to various fetal heart rate patterns. Inter-observer and regional variability. Acta Obstet Gynecol Scand 1992;71(1):48-53.

Lien JM, Morgan MA, Garite TJ, et al. Antepartum cervical ripening: applying prostaglandin E2 gel in conjunction with scheduled nonstress tests in postdate pregnancies. Am J Obstet Gynecol 1998;179(2):453-8.

Lumley J, Lester A, Anderson I, et al. A randomized trial of weekly cardiotocography in high-risk obstetric patients. Br J Obstet Gynaecol 1983;90(11):1018-26.

Lydon-Rochelle M, Holt VL, Easterling TR, et al. Risk of uterine rupture during labor among women with a prior cesarean delivery. N Engl J Med 2001;345(1):3-8.

MacKenzie IZ, Burns E. Randomised trial of one versus two doses of prostaglandin E2 for induction of labour: 1. Clinical outcome. Br J Obstet Gynaecol 1997;104(9):1062-7.

Magann EF, Chauhan SP, Nevils BG, et al. Management of pregnancies beyond forty-one weeks' gestation with an unfavorable cervix. Am J Obstet Gynecol 1998;178(6):1279-87.

Magann EF, McNamara MF, Whitworth NS, et al. Can we decrease postdatism in women with an unfavorable cervix and a negative fetal fibronectin test result at term by serial membrane sweeping? Am J Obstet Gynecol 1998;179(4):890-4.

Mahmood TA. A prospective comparative study on the use of prostaglandin E2 gel (2 mg) and prostaglandin E2 tablet (3 mg) for the induction of labour in primigravid women with unfavorable cervices. Eur J Obstet Gynecol Reprod Biol 1989;33(2):169-75.

Mannino F. Neonatal complications of postterm gestation. J Reprod Med Obstet Gynecol 1988;33(3):271-6.

Martin JN, Sessums JK, Howard P, et al. Alternative approaches to the management of gravidas with

prolonged-postterm-postdate pregnancies. J Miss State Med Assoc 1989;30(4):105-11.

McColgin SW, Hampton HL, McCaul JF, et al. Stripping membranes at term: can it safely reduce the incidence of post-term pregnancies? Obstet Gynecol 1990:76(4):678-80.

McColgin SW, Patrissi GA, Morrison JC. Stripping the fetal membranes at term. Is the procedure safe and efficacious? Journal of Reproductive Medicine 1990;35(8):811-4.

Mehl-Madrona L, Madrona MM. Physician- and midwife-attended home births. Effects of breech, twin, and post-dates outcome data on mortality rates. J Nurse-Midwifery 1997;42(2):91-8.

Misra M, Vavre S. Labour induction with intracervical prostaglandin Einf 2 gel and intravenous oxytocin in women with a very unfavourable cervix. Aust N Z J Obstet Gynaecol 1994;34(5):511-5.

Moher D, Jadad AR, Tugwell P. Assessing the quality of randomized controlled trials. Current issues and future directions. Internat J Tech Assess Health Care 1996;12(2):195-208.

Monaghan J, O'Herlihy C, Boylan P. Ultrasound placental grading and amniotic fluid quantitation in prolonged pregnancy. Obstet Gynecol 1987;70(3 Pt 1):349-52.

Montan S, Malcus P. Amniotic fluid index in prolonged pregnancy: A cohort study. J Matern Fetal Invest 1995;5(1):4-7.

Mouw RJ, Egberts J, Kragt H, et al. Cervicovaginal fetal fibronectin concentrations: predictive value of impending birth in postterm pregnancies. Eur J Obstet Gynecol Reprod Biol 1998;80(1):67-70.

Mundle WR, Young DC. Vaginal misoprostol for induction of labor: a randomized controlled trial. Obstet Gynecol 1996;88(4 Pt 1):521-5.

National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. A clinical trial of induction of labor versus expectant management in postterm pregnancy. The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. Am J Obstet Gynecol 1994;170(3):716-23.

Nationwide Inpatient Sample [NIS]. [electronic database]. Release 6; 1997 data. Agency for Healthcare Research and Quality: Rockville, MD. Available from: http://www.ntis.gov/fcpc/cpn8834.htm. 1997.

Neilson JP. Mifepristone for induction of labour (Cochrane Review). In: The Cochrane Library, Issue 3, 2001. Oxford: Update Software.

Nocon JJ, McKenzie DK, Thomas LJ, et al. Shoulder dystocia: an analysis of risks and obstetric maneuvers. Am J Obstet Gynecol 1993;168(6 Pt 1):1732-7.

O'Brien JM, Mercer BM, Cleary NT, et al. Efficacy of outpatient induction with low-dose intravaginal prostaglandin E2: a randomized, double-blind, placebo-controlled trial. Am J Obstet Gynecol 1995;173(6):1855-9.

O'Reilly-Green CP, Divon MY. Predictive value of amniotic fluid index for oligohydramnios in patients with prolonged pregnancies. J Matern Fetal Med 1996;5(4):218-26.

O'Reilly-Green CP, Divon MY. Receiver operating characteristic curves of sonographic estimated fetal weight for prediction of macrosomia in prolonged pregnancies. Ultrasound Obstet Gynecol 1997;9(6):403-8.

Ohel G, Rahav D, Rothbart H, et al. Randomised trial of outpatient induction of labor with vaginal PGE2 at 40-41 weeks of gestation versus expectant management. Arch Gynecol Obstet 1996;258(3):109-12.

Papageorgiou I, Tsionou C, Minaretzis D, et al. Labor characteristics of uncomplicated prolonged pregnancies after induction with intracervical prostaglandin E2 gel versus intravenous oxytocin. Gynecol Obstet Invest 1992;34(2):92-6.

Pattison N, McCowan L. Cardiotocography for antepartum fetal assessment (Cochrane Review). In: The Cochrane Library, Issue 1, 2001. Oxford: Update Software.

Pearce JM, Cardozo C. Prolonged pregnancy: the management debate. BMJ 1988;297:715-7.

Phelan JP, Platt LD, Yeh SY, et al. The role of ultrasound assessment of amniotic fluid volume in the management of the postdate pregnancy. Am J Obstet Gynecol 1985;151(3):304-8.

Phelan JP, Platt LD, Yeh SY, et al. Continuing role

of the nonstress test in the management of postdates pregnancy. Obstet Gynecol 1984;64(5):624-8.

Phelps JY, Higby K, Smyth MH, et al. Accuracy and intraobserver variability of simulated cervical dilatation measurements. Am J Obstet Gynecol 1995;173(3 Pt 1):942-5.

Plaut MM, Schwartz ML, Lubarsky SL. Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. Am J Obstet Gynecol 1999;180(6 Pt 1):1535-42.

Pollack RN, Hauer-Pollack G, Divon MY. Macrosomia in postdates pregnancies: the accuracy of routine ultrasonographic screening. Am J Obstet Gynecol 1992;167(1):7-11.

Ramrekersingh-White P, Farkas AG, Chard T, et al. Self-selected expectant management of post dates pregnancy including the use of Doppler ultrasound. J Obstet Gynaecol 1993;13(1):16-9.

Rayburn W, Gosen R, Ramadei C, et al. Outpatient cervical ripening with prostaglandin E2 gel in uncomplicated postdate pregnancies. Am J Obstet Gynecol 1988;158(6 Pt 1):1417-23.

Rayburn WF, Motley ME, Stempel LE, et al. Antepartum prediction of the postmature infant. Obstet Gynecol 1982;60(2):148-53.

Roberts LJ, Young KR. The management of prolonged pregnancy—an analysis of women's attitudes before and after term. Br J Obstet Gynaecol 1991;98(11):1102-6.

Rouse DJ, Owen J, Goldenberg RL, et al. The effectiveness and costs of elective cesarean delivery for fetal macrosomia diagnosed by ultrasound. JAMA 1996;276(18):1480-6.

Royal College of Obstetricians and Gynaecologists. Induction of labour: evidence-based clinical guideline number 9. London: Royal College of Obstetricians and Gynaecologists, 2001.

Salamalekis E, Vitoratos N, Kassanos D, et al. Sweeping of the membranes versus uterine stimulation by oxytocin in nulliparous women. A randomized controlled trial. Gynecol Obstet Invest 2000;49(4):240-3.

Sanchez-Ramos L, Kaunitz AM, Del Valle GO, et al. Labor induction with the prostaglandin E1 methyl analogue misoprostol versus oxytocin: a randomized trial. Obstet Gynecol 1993;81(3):332-6.

Sarkar PK, Duthie SJ. The clinical significance of reduced amniotic fluid index in post-term pregnancy: a retrospective study. J Obstet Gynaecol 1997;17(3):274-5.

Sarno AP, Hinderstein WN, Staiano RA. Fetal macrosomia in a military hospital: Incidence, risk factors, and outcome. Mil Med 1991;156(2):55-8.

Satin AJ, Hankins GD, Yeomans ER. A prospective study of two dosing regimens of oxytocin for the induction of labor in patients with unfavorable cervices. Am J Obstet Gynecol 1991;165(4 Pt 1):980-4.

Sawai SK, O'Brien WF, Mastrogiannis DS, et al. Patient-administered outpatient intravaginal prostaglandin E2 suppositories in post-date pregnancies: a double-blind, randomized, placebo-controlled study. Obstet Gynecol 1994;84(5):807-10.

Sawai SK, Williams MC, O'Brien WF, et al. Sequential outpatient application of intravaginal prostaglandin E2 gel in the management of postdates pregnancies. Obstet Gynecol 1991;78(1):19-23.

Schreyer P, Bar-Natan N, Sherman DJ, et al. Fetal breathing movements before oxytocin induction in prolonged prenancies. Am J Obstet Gynecol 1991;165(3):577-81.

Sciscione AC, Nguyen L, Manley J, et al. A randomized comparison of transcervical Foley catheter to intravaginal misoprostol for preinduction cervical ripening. Obstet Gynecol 2001;97:603-7.

Sherer DM, Onyeije CI, Binder D, et al. Uncomplicated baseline fetal tachycardia or bradycardia in postterm pregnancies and perinatal outcome. Am J Perinatol 1998;15(5):335-8.

Shime J, Librach CL, Gare DJ, et al. The influence of prolonged pregnancy on infant development at one and two years of age: a prospective controlled study. Am J Obstet Gynecol 1986;154(2):341-5.

Small ML, Phelan JP, Smith CV, et al. An active management approach to the postdate fetus with a reactive nonstress test and fetal heart rate decelerations. Obstet Gynecol 1987;70(4):636-40.

Smith GCS. Life-table analysis of the risk of perinatal death at term and post term in singleton pregnancies. Am J Obstet Gynecol 2001;184:489-96.

Srisomboon J, Piyamongkol W, Aiewsakul P. Comparison of intracervical and intravaginal misoprostol for cervical ripening and labour induction in patients with an unfavourable cervix. J Med Assoc Thai 1997;80(3):189-94.

Stenlund PM, Ekman G, Aedo A-R, et al. Induction of labor with mifepristone: a randomized, double-blind study versus placebo. Acta Obstet Gynecol Scand 1999;78(9):793-8.

Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. JAMA 2000;283(15):2008-12.

Surbek DV, Boesiger H, Hoesli I, et al. A double-blind comparison of the safety and efficacy of intravaginal misoprostol and prostaglandin E2 to induce labor. Am J Obstet Gynecol 1997;177(5):1018-23.

Tai-Seale M, Rodwin M, Wedig G. Drive-through delivery: where are the "savings"? Med Care Res Rev 1999;56(1):30-46.

Tam W-H, Tai SMB, Rogers MS. Prediction of cervical response to prostaglandin Einf 2 using fetal fibronectin. Acta Obstet Gynecol Scand 1999;78(10):861-5.

Tongsong T, Srisomboon J. Amniotic fluid volume as a predictor of fetal distress in postterm pregnancy. Int J Gynaecol Obstet 1993;40(3):213-7.

Toppozada MK, Anwar MY, Hassan HA, et al. Oral or vaginal misoprostol for induction of labor. Int J Gynaecol Obstet 1997;56(2):135-9.

Tuffnell DJ, Bryce F, Johnson N, et al. Simulation of cervical changes in labour: reproducibility of expert assessment. Lancet 1989;2(8671):1089-90.

Varaklis K, Gumina R, Stubblefield PG. Randomized controlled trial of vaginal misoprostol and intracervical prostaglandin E2 gel for induction of labor at term. Obstet Gynecol 1995;86(4 Pt 1):541-4.

Ventura SJ, Martin JA, Curtin SC, et al. Births: Final data for 1998. National Vital Statistics Report, vol 48, no 3. Hyattsville, MD. National Center for Health Statistics, 2000.

Voss DH, Cumminsky KC, Cook VD, et al. Effects of three concentrations of intracervical prostaglandin Einf 2 gel for cervical ripening. J Matern Fetal Med 1996;5(4):186-93.

Weeks JW, Pitman T, Spinnato JA 2nd. Fetal macrosomia: does antenatal prediction affect delivery route and birth outcome? Am J Obstet Gynecol 1995;173(4):1215-9.

Weiner Z, Farmakides G, Schulman H, et al. Computerized analysis of fetal heart rate variation in postterm pregnancy: prediction of intrapartum fetal distress and fetal acidosis. Am J Obstet Gynecol 1994;171(4):1132-8.

Weiner Z, Reichler A, Zlozover M, et al. The value of Doppler ultrasonography in prolonged pregnancies. Eur J Obstet Gynecol Reprod Biol 1993;48(2):93-7.

Wing DA, Fassett MJ, Mishell DR. Mifepristone for preinduction cervical ripening beyond 41 weeks' gestation: a randomized controlled trial. Obstet Gynecol 2000;96(4):543-8.

Wing DA, Jones MM, Rahall A, et al. A comparison of misoprostol and prostaglandin E2 gel for preinduction cervical ripening and labor induction. Am J Obstet Gynecol 1995;172(6):1804-10.

Wing DA, Ortiz-Omphroy G, Paul RH. A comparison of intermittent vaginal administration of misoprostol with continuous dinoprostone for cervical ripening and labor induction. Am J Obstet Gynecol 1997;177(3):612-8.

Wing DA, Paul RH. A comparison of differing dosing regimens of vaginally administered misoprostol for preinduction cervical ripening and labor induction [published erratum appears in Am J Obstet Gynecol 1997 Jun;176(6):1423]. Am J Obstet Gynecol 1996;175(1):158-64.

Wing DA, Rahall A, Jones MM, et al. Misoprostol: an effective agent for cervical ripening and labor induction. Am J Obstet Gynecol 1995;172(6):1811-6.

Wiriyasirivaj B, Vutyavanich T, Ruangsri RA. A randomized controlled trial of membrane stripping at term to promote labor. Obstet Gynecol 1996;87(5 Pt 1):767-70.

Witter FR, Weitz CM. A randomized trial of induction at 42 weeks gestation versus expectant management for postdates pregnancies. Am J Perinatol 1987;4(3):206-11.

Witter FR, Weitz CM. Cervical examination prior to induction in postdate pregnancies. Surg Gynecol Obstet 1989;168(3):214-6.

Yudkin PL, Wood L, Redman CWG. Risk of unexplained stillbirth at different gestational ages. Lancet 1987;1:1192-4.