Bibliography

Efficacy of felbamate in childhood epileptic encephalopathy (Lennox-Gastaut syndrome). The Felbamate Study Group in Lennox-Gastaut Syndrome. N Engl J Med 1993 Jan;328(1):29-33.

Gabapentin in partial epilepsy. UK Gabapentin Study Group. Lancet 1990 May;335(8698):1114-7.

ILAE Commission Report. The epidemiology of the epilepsies: future directions. International League Against Epilepsy. Epilepsia 1997 May;38(5):614-8.

National Institutes of Health Consensus Development Conference Statement: surgery for epilepsy, March 19-21, 1990. Epilepsia 1990;31(6):806-12.

Optimizing epilepsy management: seizure control, medication tolerability, and co-morbidities. Neurology 2002 Apr;58(8 Suppl 5):S1-39.

Physician's desk reference 2001. 55th ed. Montvale (NJ): Medical Economics/Thomson Healthcare; 2001. 3506 p.

Practice parameter: management issues for women with epilepsy (summary statement). Report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology 1998 Oct;51(4):944-8.

Proposal for revised classification of epilepsies and epileptic syndromes. Commission on Classification and Terminology of the International League Against Epilepsy. Epilepsia 1989;30(4):389-99.

Proposal for revised clinical and electroencephalographic classification of epileptic seizures. From the Commission on Classification and Terminology of the International League Against Epilepsy. Epilepsia 1981 Aug;22(4):489-501.

Quality of life in epilepsy QOLIE-31 (Version 1.0). Patient inventory. [internet]. Santa Monica (CA): RAND Health Communications; 1993 [cited 2002 Jan 24]. [9 p]. Available: http://www.rand.org/health/surveys/QOLIE31.pdf.

Quality of life in epilepsy QOLIE-89 (Version 1.0). Patient inventory. [internet]. Santa Monica (CA): RAND Health Communications; [cited 2002 Jan 24]. [15 p]. Available: http://www.rand.org/health/surveys/QOLIE89.survey.pdf.

A randomized controlled trial of chronic vagus nerve stimulation for treatment of medically intractable seizures. The Vagus Nerve Stimulation Study Group. Neurology 1995 Feb;45(2):224-30.

Scoring manual for the quality of life in epilepsy inventory-89 (QOLIE-89). [internet]. Santa Monica (CA): RAND Health Communications; [cited 2002 Jan 24]. [14 p]. Available: http://www.rand.org/health/surveys/QOLIE89.scoring.pdf.

Mortality statistics: general. Review of the Registrar General on deaths in England and Wales, 1999. London (UK): The Stationery Office; 2001. Table 4: Death rates per 1,000 population within sex and age-groups, and infant mortality rates per 1,000 live births 1841-1999. (Series DH1; no. 32).p. 5-10. Also available: http://www.statistics.gov.uk/statbase.

Mortality statistics: general. Review of the Registrar General on deaths in England and Wales, 1999. London (UK): The Stationery Office; 2001. Table 7: Death rates for 1,000,000 population: age-group and sex, infant mortality rates and standardized mortality ratios for selected underlying causes, 1999. (Series DH1; no. 32).p. 12-37.

Topiramate in medically intractable partial epilepsies: double-blind placebo-controlled randomized parallel group trial. Korean Topiramate Study Group. Epilepsia 1999 Dec;40(12):1767-74.

Statistical abstract of the United States. Washington (DC): U.S. Census Bureau; 2000. Vital statistics. p. various. Also available: http://www.census.gov/prod/www/statistical-abstract.html.

Aarli JA. Epilepsy and the immune system. Arch Neurol 2000 Dec;57(12):1689-92.

Acciarri N, Giulioni M, Padovani R, et al. Surgical management of cerebral cavernous angiomas causing epilepsy. J Neurosurg Sci 1995 Mar;39(1):13-20.

Acharya JN, Dinner DS. Use of the intracarotid amobarbital procedure in the evaluation of memory. J Clin Neurophysiol 1997 Jul;14(4):311-25.

Adam C, Clemenceau S, Semah F, et al. Variability of presentation in medial temporal lobe epilepsy: a study of 30 operated cases. Acta Neurol Scand 1996 Jul;94(1):1-11.

Adler J, Erba G, Winston KR, et al. Results of surgery for extratemporal partial epilepsy that began in childhood. Arch Neurol 1991 Feb;48(2):133-40.

Agency for Healthcare Research and Quality (AHRQ). Management of newly diagnosed patients with epilepsy: a systematic review of the literature: summary. Rockville (MD): AHRQ; 2001 Feb 1. 158 p. (Evidence report/technology assessment; no. 39). Also available: http://www.ahrq.gov.

Aiken SP, Brown WM. Treatment of epilepsy: existing therapies and future developments. Front Biosci 2000 Nov;5:E124-52.

Aldenkamp AP. Cognitive and behavioural assessment in clinical trials: when should they be done? Epilepsy Res 2001 May;45(1-3):155-7; discussion 159-61.

Aldenkamp AP, Van de Veerdonk SH, Majoie HJ, et al. Effects of 6 months of treatment with vagus nerve stimulation on behavior in children with Lennox-Gastaut syndrome in an open clinical nonrandomized study. Epilepsy Behav 2001 Aug;2(4):343-50.

Alsaadi T, Morris GL, Mueller WM. Prognostic value of the consistency of lateralizing ictal features and ictal EEG in patients undergoing temporal lobectomy for refractory complex partial seizures. Clin Electroencephalogr 2001 Apr;32(2):87-91.

Altshuler L, Rausch R, Delrahim S, et al. Temporal lobe epilepsy, temporal lobectomy, and major depression. J Neuropsychiatry Clin Neurosci 1999 Fall;11(4):436-43.

Alving J. Serum prolactin levels are elevated also after pseudo-epileptic seizures. Seizure 1998 Apr;7(2):85-9.

Andersen B, Rogvi-Hansen B, Kruse-Larsen C, et al. Corpus callosotomy: seizure and psychosocial outcome. A 39-month follow-up of 20 patients. Epilepsy Res 1996 Feb;23(1):77-85.

Andrews DJ, Schonfeld WH. Predictive factors for controlling seizures using a behavioural approach. Seizure 1992 Jun;1(2):111-6.

Andriola MR, Vitale SA. Vagus nerve stimulation in the developmentally disabled. Epilepsy Behav 2001 Apr;2(2):129-34.

Anhoury S, Brown RJ, Krishnamoorthy ES, et al. Psychiatric outcome after temporal lobectomy: a predictive study. Epilepsia 2000 Dec;41(12):1608-15.

Anhut H, Ashman P, Feuerstein TJ, et al. Gabapentin (Neurontin) as add-on therapy in patients with partial seizures: a double-blind, placebo-controlled study. The International Gabapentin Study Group. Epilepsia 1994;35(4):795-801.

Annegers JF, Coan SP. SUDEP: overview of definitions and review of incidence data. Seizure 1999 Sep;8(6):347-52.

Annegers JF, Coan SP, Hauser WA, et al. Epilepsy, vagal nerve stimulation by the NCP system, all-cause mortality, and sudden, unexpected, unexplained death. Epilepsia 2000 May;41(5):549-53.

Annegers JF, Coan SP, Hauser WA, et al. Epilepsy, vagal nerve stimulation by the NCP system, mortality, and sudden, unexpected, unexplained death. Epilepsia 1998 Feb;39(2):206-12.

Anzola GP. Predictivity of plasma prolactin levels in differentiating epilepsy from pseudoseizures: a prospective study. Epilepsia 1993;34(6):1044-8.

Appleton R, Fichtner K, LaMoreaux L, et al. Gabapentin as add-on therapy in children with refractory partial seizures: a 12-week, multicentre, double-blind, placebo-controlled study. Gabapentin Paediatric Study Group. Epilepsia 1999 Aug;40(8):1147-54.

Armour DJ, Veitch GB. Is valproate monotherapy a practical possibility in chronically uncontrolled epilepsy? J Clin Pharm Ther 1988 Feb;13(1):53-64.

Arnold LM, Privitera MD. Psychopathology and trauma in epileptic and psychogenic seizure patients. Psychosomatics 1996;37(5):438-43.

Arroyo S, Brodie MJ, Avanzini G, et al. Is refractory epilepsy preventable? Epilepsia 2002 Apr;43(4):437-44.

Assaf BA, Ebersole JS. Visual and quantitative ictal EEG predictors of outcome after temporal lobectomy. Epilepsia 1999 Jan;40(1):52-61.

Atherton Skaff PJ, Sloan JA. Design and analysis of equivalence clinical trials via the SAS system. In: Proceedings of the 23rd annual SAS users group international. March 22-25, 1998; Nashville (TN). Also available: http://www2.sas.com/proceedings/sugi23/Stats/p218.pdf.

Barry JJ, Atzman O, Morrell MJ. Discriminating between epileptic and nonepileptic events: the utility of hypnotic seizure induction. Epilepsia 2000 Jan;41(1):81-4.

Battaglia D, Di Rocco C, Iuvone L, et al. Neuro-cognitive development and epilepsy outcome in children with surgically treated hemimegalencephaly. Neuropediatrics 1999 Dec;30(6):307-13.

Bauer J, Burr W. Course of chronic focal epilepsy resistant to anticonvulsant treatment. Seizure 2001 Jun;10(4):239-46.

Baxendale S. Amnesia in temporal lobectomy patients: historical perspective and review. Seizure 1998 Feb;7(1):15-24.

Bazil CW, Pedley TA. Advances in the medical treatment of epilepsy. Annu Rev Med 1998;49:135-62.

Becu M, Becu N, Manzur G, et al. Self-help epilepsy groups: an evaluation of effect on depression and schizophrenia. Epilepsia 1993;34(5):841-5.

Begg CB, Greenes RA. Assessment of diagnostic tests when disease verification is subject to selection bias. Biometrics 1983 Mar;39(1):207-15.

Ben-Menachem E, Falter U. Efficacy and tolerability of levetiracetam 3000 mg/d in patients with refractory partial seizures: a multicenter, double-blind, responder-selected study evaluating monotherapy. Epilepsia 2000;41(10):1276-83.

Ben-Menachem E, Hellstrom K, Waldton C, et al. Evaluation of refractory epilepsy treated with vagus nerve stimulation for up to 5 years. Neurology 1999 Apr;52(6):1265-7.

Ben-Menachem E, Henriksen O, Dam M, et al. Double-blind, placebo-controlled trial of topiramate as add-on therapy in patients with refractory partial seizures. Epilepsia 1996 Jun;37(6):539-43.

Benbadis SR. Provocative techniques should be used for the diagnosis of psychogenic nonepileptic seizures. Arch Neurol 2001 Dec;58(12):2063-5.

Benbadis SR. What can EEG-video monitoring do for you and your patients? J Fla Med Assoc 1997 Jun-1997 Jul;84(5):320-2.

Benbadis SR, Tatum WO 4th. Advances in the treatment of epilepsy. Am Fam Physician 2001 Jul;64(1):91-8.

Benbadis SR, Wolgamuth BR, Goren H, et al. Value of tongue biting in the diagnosis of seizures. Arch Intern Med 1995 Nov;155(21):2346-9.

Berg AT, Shinnar S. The contributions of epidemiology to the understanding of childhood seizures and epilepsy. J Child Neurol 1994 Oct;9(Suppl 2):19-26.

Berg AT, Shinnar S. Do seizures beget seizures? An assessment of the clinical evidence in humans. J Clin Neurophysiol 1997 Mar;14(2):102-10.

Bergey GK, Morris HH, Rosenfeld W, et al. Gabapentin monotherapy: I. An 8-day, double-blind, dose-controlled, multicenter study in hospitalized patients with refractory complex partial or secondarily generalized seizures. The U.S. Gabapentin Study Group 88/89. Neurology 1997 Sep;49(3):739-45.

Berkovic SF, Andermann F, Olivier A, et al. Hippocampal sclerosis in temporal lobe epilepsy demonstrated by magnetic resonance imaging. Ann Neurol 1991 Feb;29(2):175-82.

Berkovic SF, McIntosh AM, Kalnins RM, et al. Preoperative MRI predicts outcome of temporal lobectomy: an actuarial analysis. Neurology 1995 Jul;45(7):1358-63.

Berlin JA, Santanna J, Schmid CH, et al. Individual patient- versus group-level data meta-regressions for the investigation of treatment effect modifiers: ecological bias rears its ugly head. Stat Med 2002 Feb;21(3):371-87.

Betts T, Waegemans T, Crawford P. A multicentre, double-blind, randomized, parallel group study to evaluate the tolerability and efficacy of two oral doses of levetiracetam, 2000 mg daily and 4000 mg daily, without titration in patients with refractory epilepsy. Seizure 2000 Mar;9(2):80-7.

Beydoun A. Monotherapy trials with gabapentin for partial epilepsy. Epilepsia 1999;40(Suppl 6):S13-6; discussion S73-4.

Beydoun A, Fischer J, Labar DR, et al. Gabapentin monotherapy: II. A 26-week, double-blind, dose-controlled, multicenter study of conversion from polytherapy in outpatients with refractory complex partial or secondarily generalized seizures. The U.S. Gabapentin Study Group 82/83. Neurology 1997b Sep;49(3):746-52.

Beydoun A, Sachdeo RC, Rosenfeld WE, et al. Oxcarbazepine monotherapy for partial-onset seizures: a multicenter, double-blind, clinical trial. Neurology 2000 Jun;54(12):2245-51.

Beydoun A, Sackellares JC, Shu V. Safety and efficacy of divalproex sodium monotherapy in partial epilepsy: a double-blind, concentration-response design clinical trial. Depakote Monotherapy for Partial Seizures Study Group. Neurology 1997a Jan;48(1):182-8.

Bidzinski J. Temporal lobectomy. Acta Neurochir Suppl (Wien) 1990;50:119-21.

Binnie CD. Vagus nerve stimulation for epilepsy: a review. Seizure 2000 Apr;9(3):161-9.

Birnbach CD, Wilensky AJ, Dodrill CB. Predictors of early mortality and sudden death in epilepsy: a multidisciplinary approach. J Epilepsy 1991;4(1):11-7.

Biton V, Montouris GD, Ritter F, et al. A randomized, placebo-controlled study of topiramate in primary generalized tonic-clonic seizures. Topiramate YTC Study Group. Neurology 1999 Apr;52(7):1330-7.

Bittencourt PR, Antoniuk SA, Bigarella MM, et al. Carbamazepine and phenytoin in epilepsies refractory to barbiturates: efficacy, toxicity and mental function. Epilepsy Res 1993 Oct;16(2):147-55.

Bizzi JW, Bruce DA, North R, et al. Surgical treatment of focal epilepsy in children: results in 37 patients. Pediatr Neurosurg 1997 Feb;26(2):83-92.

Blackwelder WC. 'Proving the null hypothesis' in clinical trials. Control Clin Trials 1982 Dec;3(4):345-53.

Bladin PF. Post-temporal lobectomy seizures. Clin Exp Neurol 1987;24:77-83.

Bladin PF. Psychosocial difficulties and outcome after temporal lobectomy. Epilepsia 1992;33(5):898-907.

Blumcke I, Beck H, Lie AA, et al. Molecular neuropathology of human mesial temporal lobe epilepsy. Epilepsy Res 1999 Sep;36(2-3):205-23.

Blume WT. Current trends in electroencephalography. Curr Opin Neurol 2001 Apr;14(2):193-7.

Blume WT. Temporal lobe epilepsy surgery in childhood: rationale for greater use. Can J Neurol Sci 1997 May;24(2):95-8.

Blume WT, Desai HB, Girvin JP, et al. Effectiveness of temporal lobectomy measured by yearly follow-up and multivariate analysis. J Epilepsy 1994;7:203-14.

Blume WT, Girvin JP, McLachlan RS, et al. Effective temporal lobectomy in childhood without invasive EEG. Epilepsia 1997 Feb;38(2):164-7.

Blume WT, Whiting SE, Girvin JP. Epilepsy surgery in the posterior cortex. Ann Neurol 1991 Jun;29(6):638-45.

Blumer D, Wakhlu S, Davies K, et al. Psychiatric outcome of temporal lobectomy for epilepsy: incidence and treatment of psychiatric complications. Epilepsia 1998 May;39(5):478-86.

Boling W, Andermann F, Reutens D, et al. Surgery for temporal lobe epilepsy in older patients. J Neurosurg 2001;95(2):242-8.

Boon P, De Reuck J, Drieghe C, et al. Long-term video-EEG monitoring revisited. The value of interictal and ictal video-EEG recording, a follow-up study. Eur Neurol 1994;34(Suppl 1):33-9.

Boon P, Vonck K, D'Have M, et al. Cost-benefit of vagus nerve stimulation for refractory epilepsy. Acta Neurol Belg 1999 Dec;99(4):275-80.

Boon PA, Williamson PD. The diagnosis of pseudoseizures. Clin Neurol Neurosurg 1993 Mar;95(1):1-8.

Bouilleret V, Dupont S, Spelle L, et al. Insular cortex involvement in mesiotemporal lobe epilepsy: a positron emission tomography study. Ann Neurol 2002 Feb;51(2):202-8.

Bourgeois B, Leppik IE, Sackellares JC, et al. Felbamate: a double-blind controlled trial in patients undergoing presurgical evaluation of partial seizures. Neurology 1993 Apr;43(4):693-6.

Brown MC, Levin BE, Ramsay RE, et al. Characteristics of patients with nonepileptic seizures. J Epilepsy 1991;4(4):225-9.

Brown WM, Aiken SP. Felbamate: clinical and molecular aspects of a unique antiepileptic drug. Crit Rev Neurobiol 1998;12(3):205-22.

Browne TR, Holmes GL. Primary care: epilepsy. N Engl J Med 2001 Apr;344(15):1145-51.

Bruni J. Efficacy of topiramate. Can J Neurol Sci 1998 Aug;25(3):S6-7.

Bruni J. Outcome evaluation of gabapentin as add-on therapy for partial seizures. 'NEON' Study Investigators Group. Neurontin Evaluation of Outcomes in Neurological Practice. Can J Neurol Sci 1998 May;25(2):134-40.

Bulteau C, Jambaque I, Viguier D, et al. Epileptic syndromes, cognitive assessment and school placement: a study of 251 children. Dev Med Child Neurol 2000 May;42(5):319-27.

Bus AG, van IJzendoorn MH. Phonological awareness and early reading: a meta-analysis of experimental training studies. J Educ Psychol 1999;91(3):403-14.

Bye A, Lamont P, Healy L. Commencement of a paediatric EEG-video telemetry service. Clin Exp Neurol 1990;27:83-8.

Callaghan N, O'Dwyer R, Keating J. Unnecessary polypharmacy in patients with frequent seizures. Acta Neurol Scand 1984 Jan;69(1):15-9.

Canizares S, Torres X, Boget T, et al. Does neuroticism influence cognitive self-assessment after epilepsy surgery? Epilepsia 2000 Oct;41(10):1303-9.

Cappabianca P, Alfieri A, Maiuri F, et al. Supratentorial cavernous malformations and epilepsy: seizure outcome after lesionectomy on a series of 35 patients. Clin Neurol Neurosurg 1997 Aug;99(3):179-83.

Carey P, O'Moore B, Sheahan K, et al. Experience of temporal lobectomy as a treatment modality for epilepsy, using inter-ictal EEG data alone to localize the epileptogenic focus. Ir Med J 1985;78(3):74-7.

Carmant L, Holmes GL, Lombroso CT. Outcome following corpus callosotomy. J Epilepsy 1998;11(4):224-8.

Carreno M, Wyllie E, Bingaman W, et al. Seizure outcome after functional hemispherectomy for malformations of cortical development. Neurology 2001 Jul;57(2):331-3.

Carson BS Sr. Indications and outcomes for lobectomy, corpus callosotomy, and hemispherectomy in pediatric neurosurgical patients. Clin Neurosurg 2000;47:385-99.

Casazza M, Avanzini G, Ciceri E, et al. Lesionectomy in epileptogenic temporal lobe lesions: preoperative seizure course and postoperative outcome. Acta Neurochir Suppl (Wien) 1997;68:64-9.

Castillo S, Schmidt DB, White S. Oxcarbazepine add-on for drug-resistant partial epilepsy (Cochrane Review). In: The Cochrane Library. Issue 4. Oxford: Update Software; 2001.

Cereghino JJ, Biton V, Abou-Khalil B, et al. Levetiracetam for partial seizures: results of a double-blind, randomized clinical trial. Neurology 2000 Jul;55(2):236-42.

Chadwick D. Epilepsy. J Neurol Neurosurg Psychiatry 1994 Mar;57(3):264-77.

Chadwick D. Monotherapy clinical trials of new antiepileptic drugs: design, indications, and controversies. Epilepsia 1997;38(Suppl 9):S16-20.

Chadwick D, Leiderman DB, Sauermann W, et al. Gabapentin in generalized seizures. Epilepsy Res 1996 Nov;25(3):191-7.

Chassoux F, Devaux B, Landre E, et al. Postoperative motor deficits and recovery after cortical resections. Adv Neurol 1999;81:189-99.

Chayasirisobhon S, Griggs L, Westmoreland S, et al. The usefulness of one to two hour video EEG monitoring in patients with refractory seizures. Clin Electroencephalogr 1993 Apr;24(2):78-84.

Chayasirisobhon S, Koulouris S, Parker E, et al. Vagus nerve stimulation for refractory epilepsy. Perm J 2001 Fall;5(4):21-6.

Chelune GJ, Naugle RI, Lueders H, et al. Individual change after epilepsy surgery: Practice effects and base-rate information. Neuropsychology 1993 Jan;7(1):41-52.

Chen LS, Mitchell WG, Horton EJ, et al. Clinical utility of video-EEG monitoring. Pediatr Neurol 1995 Apr;12(3):220-4.

Chene G, Thompson SG. Methods for summarizing the risk associations of quantitative variables in epidemiologic studies in a consistent form. Am J Epidemiol 1996 Sep;144(6):610-21.

Chesson AL, Kasarskis EJ, Small VW. Postictal elevation of serum creatine kinase level. Arch Neurol 1983;40(5):315-7.

Chugani HT, Chugani DC. Basic mechanisms of childhood epilepsies: studies with positron emission tomography. Adv Neurol 1999;79:883-91.

Claverie B, Rougier A. Life comfort and psychosocial adjustment linked to age at the time of anterior callosotomy. J Epilepsy 1995;8(4):321-31.

Clemmons DC, Fraser RT, Trejo W. The General Aptitude Test Battery: implications for vocational counseling and employment in epilepsy rehabilitation. J Appl Rehabil Couns 1987 Fall;18(3):33-8.

Cohen MJ, Holmes GL, Campbell R, et al. Cognitive functioning following anterior two-thirds corpus callosotomy in children and adolescents: a one-year prospective report. J Epilepsy 1991;4(2):63-5.

Collins WC, Lanigan O, Callaghan N. Plasma prolactin concentrations following epileptic and pseudoseizures. J Neurol Neurosurg Psychiatry 1983 Jun;46(6):505-8.

Cook TD, Campbell DT. Quasi-experimentation: design and analysis issues for field settings. Boston (MA): Houghton Mifflin Company; 1979. 405 p.

Cornaggia C, Canevini MP, Giuccioli D, et al. Monotherapy and polytherapy for intractable epilepsies. Ital J Neurol Sci 1985 Jun;6(2):201-5.

Corradetti R, Pugliese AM. Electrophysiological effects of felbamate. Life Sci 1998;63(13):1075-88.

Cramer JA, Arrigo C, Van Hammee G, et al. Effect of levetiracetam on epilepsy-related quality of life. N132 Study Group. Epilepsia 2000 Jul;41(7):868-74.

Culy CR, Goa KL. Lamotrigine. A review of its use in childhood epilepsy. Paediatr Drugs 2000;2(4):299-330.

Cutfield RG, Wrightson P. Surgery for temporal lobe epilepsy. N Z Med J 1987 Mar;100(820):163-6.

Dahl J, Melin L, Lund L. Effects of a contingent relaxation treatment program on adults with refractory epileptic seizures. Epilepsia 1987;28(2):125-32.

Dasheiff RM. Sudden unexpected death in epilepsy: a series from an epilepsy surgery program and speculation on the relationship to sudden cardiac death. J Clin Neurophysiol 1991 Apr;8(2):216-22.

Davies KG, Weeks RD. Results of cortical resection for intractable epilepsy using intra-operative corticography without chronic intracranial recording. Br J Neurosurg 1995;9(1):7-12.

Davis R. Cerebellar stimulation for cerebral palsy spasticity, function, and seizures. Arch Med Res 2000;31(3):290-9.

Deepak KK, Manchanda SK, Maheshwari MC. Meditation improves clinicoelectroencephalographic measures in drug-resistant epileptics. Biofeedback Self Regul 1994 Mar;19(1):25-40.

DeGiorgio CM, Schachter SC, Handforth A, et al. Prospective long-term study of vagus nerve stimulation for the treatment of refractory seizures. Epilepsia 2000 Sep;41(9):1195-200.

Delgado-Escueta AV, Walsh GO. Type I complex partial seizures of hippocampal origin: excellent results of anterior temporal lobectomy. Neurology 1985 Feb;35(2):143-54.

Denio LS, Drake ME Jr, Pakalnis A. The effect of exercise on seizure frequency. J Med 1989;20(2):171-6.

Derby LE, Tennis P, Jick H. Sudden unexplained death among subjects with refractory epilepsy. Epilepsia 1996 Oct;37(10):931-5.

Derry PA, McLachlan RS. The MMPI-2 as an adjunct to the diagnosis of pseudoseizures. Seizure 1996;5:35-40.

Derry PA, Rose KJ, McLachlan RS. Moderators of the effect of preoperative emotional adjustment on postoperative depression after surgery for temporal lobe epilepsy. Epilepsia 2000 Feb;41(2):177-85.

DerSimonian R, Laird N. Meta-analysis in clinical trials. Control Clin Trials 1986 Sep;7(3):177-88.

DeToledo JC, Lowe MR. Seizures, lateral decubitus, aspiration, and shoulder dislocation: time to change the guidelines? Neurology 2001 Feb;56(3):290-1.

DeToledo JC, Lowe MR, Ramsay RE. Restraining patients and shoulder dislocations during seizures. J Shoulder Elbow Surg 1999;8(4):300-2.

Devinsky O. Patients with refractory seizures. N Engl J Med 1999 May;340(20):1565-70.

Devinsky O, Faught RE, Wilder BJ, et al. Efficacy of felbamate monotherapy in patients undergoing presurgical evaluation of partial seizures. Epilepsy Res 1995 Mar;20(3):241-6.

Devinsky O, Sanchez-Villasenor F, Vazquez B, et al. Clinical profile of patients with epileptic and nonepileptic seizures. Neurology 1996 Jun;46(6):1530-3.

Devinsky O, Thacker K. Nonepileptic seizures. Neurol Clin 1995 May;13(2):299-319.

Di Rocco C, Iannelli A. Hemimegalencephaly and intractable epilepsy: complications of hemispherectomy and their correlations with the surgical technique. A report on 15 cases. Pediatr Neurosurg 2000 Oct;33(4):198-207.

Dirik E, Sen A, Anal O, et al. Serum cortisol and prolactin levels in childhood paroxysmal disorders. Acta Paediatr Jpn 1996 Apr;38(2):118-20.

Dlugos DJ. The early identification of candidates for epilepsy surgery. Arch Neurol 2001 Oct;58(10):1543-6.

Dodrill CB, Arnett JL, Hayes AG, et al. Cognitive abilities and adjustment with gabapentin: results of a multisite study. Epilepsy Res 1999 Jun;35(2):109-21.

Dodrill CB, Arnett JL, Shu V, et al. Effects of tiagabine monotherapy on abilities, adjustment, and mood. Epilepsia 1998 Jan;39(1):33-42.

Dodrill CB, Arnett JL, Sommerville KW, et al. Cognitive and quality of life effects of differing dosages of tiagabine in epilepsy. Neurology 1997 Apr;48(4):1025-31.

Dodrill CB, Morris GL. Effects of vagal nerve stimulation on cognition and quality of life in epilepsy. Epilepsy Behav 2001 Feb;2(1):46-53.

Dodrill CB, Van Belle G, Wilkus RJ. Stability of predictors of outcome of surgical treatment for epilepsy. J Epilepsy 1990;3(1):29-35.

Dooley M, Plosker GL. Levetiracetam. A review of its adjunctive use in the management of partial onset seizures. Drugs 2000 Oct;60(4):871-93.

Drake J, Hoffman HJ, Kobayashi J, et al. Surgical management of children with temporal lobe epilepsy and mass lesions. Neurosurgery 1987 Dec;21(6):792-7.

Drake ME Jr, Huber SJ, Pakalnis A, et al. Neuropsychological and event-related potential correlates of nonepileptic seizures. J Neuropsychiatry Clin Neurosci 1993 Winter;5(1):102-4.

Drury I, Selwa LM, Schuh LA, et al. Value of inpatient diagnostic CCTV-EEG monitoring in the elderly. Epilepsia 1999 Aug;40(8):1100-2.

Duchowny M, Jayakar P, Koh S. Selection criteria and preoperative investigation of patients with focal epilepsy who lack a localized structural lesion. Epileptic Disord 2000 Dec;2(4):219-26.

Duchowny M, Pellock JM, Graf WD, et al. A placebo-controlled trial of lamotrigine add-on therapy for partial seizures in children. Lamictal Pediatric Partial Seizure Study Group. Neurology 1999 Nov;53(8):1724-31.

Duchowny MS, Resnick TJ, Deray MJ, et al. Video EEG diagnosis of repetitive behavior in early childhood and its relationship to seizures. Pediatr Neurol 1988;4(3):162-4.

Duncan JS. Imaging and epilepsy. Brain 1997 Feb;120(Pt 2):339-77.

Duncan JS, Shorvon SD, Trimble MR. Discontinuation of phenytoin, carbamazepine, and valproate in patients with active epilepsy. Epilepsia 1990;31(3):324-33.

Duncan JS, Shorvon SD, Trimble MR. Effects of removal of phenytoin, carbamazepine, and valproate on cognitive function. Epilepsia 1990;31(5):584-91.

Duncan JS, Shorvon SD, Trimble MR. Withdrawal symptoms from phenytoin, carbamazepine and sodium valproate. J Neurol Neurosurg Psychiatry 1988 Jul;51(7):924-8.

Dunlap WP, Cortina JM, Vaslow JB, et al. Meta-analysis of experiments with matched groups or repeated measures designs. Psychol Methods 1996 Jun;1(2):170-7.

Dupont S, Semah F, Clemenceau S, et al. Accurate prediction of postoperative outcome in mesial temporal lobe epilepsy: a study using positron emission tomography with 18fluorodeoxyglucose. Arch Neurol 2000 Sep;57(9):1331-6.

Duse M, Notarangelo LD, Tiberti S, et al. Intravenous immune globulin in the treatment of intractable childhood epilepsy. Clin Exp Immunol 1996 May;104(Suppl 1):71-6.

Eberhardt KE, Stefan H, Buchfelder M, et al. The significance of bilateral CSI changes for the postoperative outcome in temporal lobe epilepsy. J Comput Assist Tomogr 2000;24(6):919-26.

Edelstein SF, Chisholm M. Management of intractable childhood seizures using the non-MCT oil ketogenic diet in 20 patients. J Am Diet Assoc 1996 Nov;96(11):1181-2.

Egli M, O'Kane M, Mothersill I, et al. Monitoring at the Swiss Epilepsy Center. Electroencephalogr Clin Neurophysiol Suppl 1985;37:371-84.

El Desoky E, Kandil MR, Afifi AH, et al. Spotlight on the continual applicability of routine plasma monitoring antiepileptic drugs in the treatment of epilepsy. Pharmacol Res 1999 Apr;39(4):269-74.

Eller DP, Patterson CA, Webb GW. Maternal and fetal implications of anticonvulsive therapy during pregnancy. Obstet Gynecol Clin North Am 1997 Sep;24(3):523-34.

Elterman RD, Glauser TA, Wyllie E, et al. A double-blind, randomized trial of topiramate as adjunctive therapy for partial-onset seizures in children. Topiramate YP Study Group. Neurology 1999 Apr;52(7):1338-44.

Elwes RD, Dunn G, Binnie CD, et al. Outcome following resective surgery for temporal lobe epilepsy: a prospective follow up study of 102 consecutive cases. J Neurol Neurosurg Psychiatry 1991 Nov;54(11):949-52.

Emilien G, Maloteaux JM. Pharmacological management of epilepsy. Mechanism of action, pharmacokinetic drug interactions, and new drug discovery possibilities. Int J Clin Pharmacol Ther 1998 Apr;36(4):181-94.

Engel J Jr, editor. Surgical treatment of the epilepsies. New York (NY): Raven Press; 1986. Outcome with respect to epileptic seizures. p. 553-71.

Engel J Jr. Intractable epilepsy: definition and neurobiology. Epilepsia 2001;42(Suppl 6):3.

Engel J Jr. The timing of surgical intervention for mesial temporal lobe epilepsy: a plan for a randomized clinical trial. Arch Neurol 1999 Nov;56(11):1338-41.

Ergene E, Behr PK, Shih JJ. Quality-of-life assessment in patients treated with vagus nerve stimulation. Epilepsy Behav 2001 Jun;2(3):284-7.

Eriksen HR, Ellertsen B, Gronningsaeter H, et al. Physical exercise in women with intractable epilepsy. Epilepsia 1994;35(6):1256-64.

Eriksson S, Malmgren K, Rydenhag B, et al. Surgical treatment of epilepsy–clinical, radiological and histopathological findings in 139 children and adults. Acta Neurol Scand 1999 Jan;99(1):8-15.

Erlichman M. Electroencephalographic (EEG) video monitoring. Health Technol Assess Rep 1990(4):1-14.

Escobedo LG, Zack MM. Comparison of sudden and nonsudden coronary deaths in the United States. Circulation 1996 Jun;93(11):2033-6.

Estes ML, Morris HH 3rd, Luders H, et al. Surgery for intractable epilepsy. Clinicopathologic correlates in 60 cases. Cleve Clin J Med 1988;55(5):441-7.

Ettinger AB, Coyle PK, Jandorf L, et al. Postictal SPECT in epileptic versus nonepileptic seizures. J Epilepsy 1998;11:67-73.

Fandino-Franky J, Torres M, Narino D, et al. Corpus callosotomy in Colombia and some reflections on care and research among the poor in developing countries. Epilepsia 2000;41(Suppl 4):S22-7.

Faught E, Ayala R, Montouris GG, et al. Randomized controlled trial of zonisamide for the treatment of refractory partial-onset seizures. Neurology 2001 Nov;57(10):1774-9.

Faught E, Sachdeo RC, Remler MP, et al. Felbamate monotherapy for partial-onset seizures: an active-control trial. Neurology 1993 Apr;43(4):688-92.

Faught E, Wilder BJ, Ramsay RE, et al. Topiramate placebo-controlled dose-ranging trial in refractory partial epilepsy using 200-, 400-, and 600-mg daily dosages. Topiramate YD Study Group. Neurology 1996 Jun;46(6):1684-90.

Ferrier CH, Alarcon G, Engelsman J, et al. Relevance of residual histologic and electrocorticographic abnormalities for surgical outcome in frontal lobe epilepsy. Epilepsia 2001 Mar;42(3):363-71.

Ferrier CH, Engelsman J, Alarcon G, et al. Prognostic factors in presurgical assessment of frontal lobe epilepsy. J Neurol Neurosurg Psychiatry 1999 Mar;66(3):350-6.

Ficker DM, So EL, Shen WK, et al. Population-based study of the incidence of sudden unexplained death in epilepsy. Neurology 1998 Nov;51(5):1270-4.

Fisher RS, Chan DW, Bare M, et al. Capillary prolactin measurement for diagnosis of seizures. Ann Neurol 1991 Feb;29(2):187-90.

Fisher RS, Uematsu S, Krauss GL, et al. Placebo-controlled pilot study of centromedian thalamic stimulation in treatment of intractable seizures. Epilepsia 1992;33(5):841-51.

Fisher RS, Vickrey BG, Gibson P, et al. The impact of epilepsy from the patient's perspective I. Descriptions and subjective perceptions. Epilepsy Res 2000 Aug;41(1):39-51.

Fletcher RH, Fletcher SW, Wagner EH. Clinical epidemiology: the essentials. 2nd ed. Baltimore (MD): Williams & Wilkins; 1988. 246 p.

Foldvary N, Nashold B, Mascha E, et al. Seizure outcome after temporal lobectomy for temporal lobe epilepsy: a Kaplan-Meier survival analysis. Neurology 2000 Feb;54(3):630-4.

Foley CM, Legido A, Miles DK, et al. Long-term computer-assisted outpatient electroencephalogram monitoring in children and adolescents. J Child Neurol 2000 Jan;15(1):49-55.

Foley CM, Legido A, Miles DK, et al. Diagnostic value of pediatric outpatient video-EEG. Pediatr Neurol 1995 Feb;12(2):120-4.

Fraser RT, Clemmons DC, Dodrill CB, et al. The difficult-to-employ in epilepsy rehabilitation: predictions of response to an intensive intervention. Epilepsia 1986;27(3):220-4.

Freeman JM, Vining EP, Pillas DJ, et al. The efficacy of the ketogenic diet-1998: a prospective evaluation of intervention in 150 children. Pediatrics 1998 Dec;102(6):1358-63.

French J, Smith M, Faught E, et al. Practice advisory: the use of felbamate in the treatment of patients with intractable epilepsy: report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Epilepsy Society. Neurology 1999 May;52(8):1540-5.

Fried R. Breathing training for the self-regulation of alveolar CO2 in the behavioural control of idiopathic epileptic seizures. In: Mostofsky DI, Loyning Y, editors. The neurobehavioural treatment of epilepsy. Hillsdale (NJ): Lawrence Erlbaum Associates; 1993. p. 19-66.

Fried R, Fox MC, Carlton RM. Effect of diaphragmatic respiration with end-tidal COinf 2 biofeedback on respiration, EEG, and seizure frequency in idiopathic epilepsy(a,b). Ann N Y Acad Sci 1990;602:67-96.

Friis ML. Valproate in the treatment of epilepsy in people with intellectual disability. J Intellect Disabil Res 1998 Dec;42(Suppl 1):32-5.

Fryback DG, Thornbury JR. The efficacy of diagnostic imaging. Med Decis Making 1991 Apr-1991 Jun;11(2):88-94.

Fuiks KS, Wyler AR, Hermann BP, et al. Seizure outcome from anterior and complete corpus callosotomy. J Neurosurg 1991 Apr;74(4):573-8.

Gann PH. Prostate-specific antigen screening for prostate cancer: issues involving test validity. Endocr Related Cancer 1996 Fall;3(3):179-89.

Garcia-Flores E. Corpus callosum section for patients with intractable epilepsy. Appl Neurophysiol 1987;50(1-6):390-7.

Garcia Sola R, Miravet J. Surgical treatment for epilepsy. Results after a minimum follow-up of five years. Acta Neurochir Suppl (Wien) 1991;52:157-60.

Garg SK, Gupta MC, Handu SS, et al. Therapeutic drug monitoring of antiepileptic drugs- a preliminary experience. Indian J Pharmacol 2000;32(1):28-30.

Gates JR, Rosenfeld WE, Maxwell RE, et al. Response of multiple seizure types to corpus callosum section. Epilepsia 1987;28(1):28-34.

Gates JR, Wada JA, Reeves AG, et al. Reevaluation of corpus callosotomy. In: Surgical treatment of the epilepsies. 2nd ed. New York (NY): Raven Press; 1993. p. 637-48.

Gavaghan DJ, Moore RA, McQuay HJ. An evaluation of homogeneity tests in meta-analyses in pain using simulations of individual patient data. Pain 2000 Apr;85(3):415-24.

Genton P, Guerrini R, Perucca E. Tiagabine in clinical practice. Epilepsia 2001;42(Suppl 3):42-5.

Giesler RB, Ashton CM, Brody B, et al. Assessing the performance of utility techniques in the absence of a gold standard. Med Care 1999 Jun;37(6):580-8.

Gillham RA. Refractory epilepsy: an evaluation of psychological methods in outpatient management. Epilepsia 1990;31(4):427-32.

Gilliam F, Vazquez B, Sackellares JC, et al. An active-control trial of lamotrigine monotherapy for partial seizures. Neurology 1998 Oct;51(4):1018-25.

Gilman JT, Duchowny M, Jayakar P, et al. Medical intractability in children evaluated for epilepsy surgery. Neurology 1994 Jul;44(7):1341-3.

Glauser TA. Expanding first-line therapy options for children with partial seizures. Neurology 2000;55(11 Suppl 3):S30-7.

Glauser TA, Nigro M, Sachdeo R, et al. Adjunctive therapy with oxcarbazepine in children with partial seizures. The Oxcarbazepine Pediatric Study Group. Neurology 2000 Jun;54(12):2237-44.

Goldstein R, Harvey AS, Duchowny M, et al. Preoperative clinical, EEG, and imaging findings do not predict seizure outcome following temporal lobectomy in childhood. J Child Neurol 1996 Nov;11(6):445-50.

Goodin DS, Aminoff MJ. Does the interictal EEG have a role in the diagnosis of epilepsy? Lancet 1984 Apr;1(8381):837-9.

Grubb BP, Gerard G, Roush K, et al. Differentiation of convulsive syncope and epilepsy with head-up tilt testing. Ann Intern Med 1991 Dec;115(11):871-6.

Guldvog B, Loyning Y, Hauglie-Hanssen E, et al. Surgical treatment for partial epilepsy among Norwegian adults. Epilepsia 1994b;35(3):540-53.

Guldvog B, Loyning Y, Hauglie-Hanssen E, et al. Surgical treatment for partial epilepsy among Norwegian children and adolescents. Epilepsia 1994a;35(3):554-65.

Guldvog B, Loyning Y, Hauglie-Hanssen E, et al. Surgical versus medical treatment for epilepsy. I. Outcome related to survival, seizures, and neurologic deficit. Epilepsia 1991;32(3):375-88.

Haafiz A, Kissoon N. Status epilepticus: current concepts. Pediatr Emerg Care 1999 Apr;15(2):119-29.

Handforth A, DeGiorgio CM, Schachter SC, et al. Vagus nerve stimulation therapy for partial-onset seizures: a randomized active-control trial. Neurology 1998 Jul;51(1):48-55.

Hanna NJ, Black M, Sander JW, et al. The national sentinal clinical audit of epilepsy-related death: epilepsy--death in the shadows. The Stationary Office; 2002 May. 134 p.

Harbord MG, Manson JI. Temporal lobe epilepsy in childhood: reappraisal of etiology and outcome. Pediatr Neurol 1987;3(5):263-8.

Hasselblad V, Hedges LV. Meta-analysis of screening and diagnostic tests. Psychol Bull 1995 Jan;117(1):167-78.

Helmstaedter C, Gleibner U, Zentner J, et al. Neuropsychological consequences of epilepsy surgery in frontal lobe epilepsy. Neuropsychologia 1998 Apr;36(4):333-41.

Hennessy MJ, Elwes RD, Honavar M, et al. Predictors of outcome and pathological considerations in the surgical treatment of intractable epilepsy associated with temporal lobe lesions. J Neurol Neurosurg Psychiatry 2001a Apr;70(4):450-8.

Hennessy MJ, Elwes RD, Rabe-Hesketh S, et al. Prognostic factors in the surgical treatment of medically intractable epilepsy associated with mesial temporal sclerosis. Acta Neurol Scand 2001b Jun;103(6):344-50.

Hennessy MJ, Langan Y, Elwes RD, et al. A study of mortality after temporal lobe epilepsy surgery. Neurology 1999 Oct;53(6):1276-83.

Henry TR. Functional neuroimaging with positron emission tomography. Epilepsia 1996 Dec;37(12):1141-54.

Henry TR, Drury I. Non-epileptic seizures in temporal lobectomy candidates with medically refractory seizures. Neurology 1997 May;48(5):1374-82.

Henry TR, Ross DA, Schuh LA, et al. Indications and outcome of ictal recording with intracerebral and subdural electrodes in refractory complex partial seizures. J Clin Neurophysiol 1999 Sep;16(5):426-38.

Hermann BP, Wyler AR, VanderZwagg R, et al. Predictors of neuropsychological change following anterior temporal lobectomy: role of regression toward the mean. J Epilepsy 1991;4(3):139-48.

Hermanns G, Noachtar S, Tuxhorn I, et al. Systematic testing of medical intractability for carbamazepine, phenytoin, and phenobarbital or primidone in monotherapy for patients considered for epilepsy surgery. Epilepsia 1996 Jul;37(7):675-9.

Ho SS, Kuzniecky RI. Algorithm on the clinical evaluation of epilepsy. J Neuroimaging 1997 Oct;7(4):236-41.

Ho SS, Newton MR, McIntosh AM, et al. Perfusion patterns during temporal lobe seizures: relationship to surgical outcome. Brain 1997 Nov;120(Pt 1):1921-8.

Hodaie M, Musharbash A, Otsubo H, et al. Image-guided, frameless stereotactic sectioning of the corpus callosum in children with intractable epilepsy. Pediatr Neurosurg 2001 Jun;34(6):286-94.

Holmes GL. Intractable epilepsy in children. Epilepsia 1996;37(Suppl 3):14-27.

Holmes GL, Sackellares JC, McKiernan J, et al. Evaluation of childhood pseudoseizures using EEG telemetry and video tape monitoring. J Pediatr 1980 Oct;97(4):554-8.

Holmes MD, Born DE, Kutsy RL, et al. Outcome after surgery in patients with refractory temporal lobe epilepsy and normal MRI. Seizure 2000 Sep;9(6):407-11.

Holmes MD, Dodrill CB. What is the significance of subjective events recorded during long-term EEG video monitoring? Epilepsia 1998 Aug;39(8):857-62.

Holmes MD, Dodrill CB, Ojemann LM, et al. Five-year outcome after epilepsy surgery in nonmonitored and monitored surgical candidates. Epilepsia 1996 Aug;37(8):748-52.

Holmes MD, Dodrill CB, Wilkus RJ, et al. Is partial epilepsy progressive? Ten-year follow-up of EEG and neuropsychological changes in adults with partial seizures. Epilepsia 1998 Nov;39(11):1189-93.

Holmes MD, Wilensky AJ, Ojemann GA, et al. Hippocampal or neocortical lesions on magnetic resonance imaging do not necessarily indicate site of ictal onsets in partial epilepsy. Ann Neurol 1999 Apr;45(4):461-5.

Hong SC, Kang KS, Seo DW, et al. Surgical treatment of intractable epilepsy accompanying cortical dysplasia. J Neurosurg 2000 Nov;93(5):766-73.

Hopkins IJ, Klug GL. Temporal lobectomy for the treatment of intractable complex partial seizures of temporal lobe origin in early childhood. Dev Med Child Neurol 1991 Jan;33(1):26-31.

Hoppe C, Helmstaedter C, Scherrmann J, et al. No evidence for cognitive side effects after 6 months of vagus nerve stimulation in epilepsy patients. Epilepsy Behav 2001 Aug;2(4):351-6.

Hosain S, Nikalov B, Harden C, et al. Vagus nerve stimulation treatment for Lennox-Gastaut syndrome. J Child Neurol 2000 Aug;15(8):509-12.

Hufnagel A, Zentner J, Fernandez G, et al. Multiple subpial transection for control of epileptic seizures: effectiveness and safety. Epilepsia 1997;38(6):678-88.

Huttenlocher PR, Hapke RJ. A follow-up study of intractable seizures in childhood. Ann Neurol 1990 Nov;28(5):699-705.

Iannelli A, Guzzetta F, Battaglia D, et al. Surgical treatment of temporal tumors associated with epilepsy in children. Pediatr Neurosurg 2000 May;32(5):248-54.

Irwig L, Tosteson AN, Gatsonis C, et al. Guidelines for meta-analyses evaluating diagnostic tests. Ann Intern Med 1994 Apr;120(8):667-76.

Ivnik RJ, Sharbrough FW, Laws ER Jr. Anterior temporal lobectomy for the control of partial complex seizures: information for counseling patients. Mayo Clin Proc 1988 Aug;63(8):783-93.

Jallon P. The problem of intractability: the continuing need for new medical therapies in epilepsy. Epilepsia 1997;38(Suppl 9):S37-42.

Jan MM, Sadler M, Rahey SR. Lateralized postictal EEG delta predicts the side of seizure surgery in temporal lobe epilepsy. Epilepsia 2001 Mar;42(3):402-5.

Jawad S, Richens A, Goodwin G, et al. Controlled trial of lamotrigine (Lamictal) for refractory partial seizures. Epilepsia 1989;30(3):356-63.

Jedrzejczak J, Owczarek K. Patients with coexistent psychogenic pseudoepileptic and epileptic seizures: a psychological profile. Seizure;10:566-9.

Jick SS, Cole TB, Mesher RA, et al. Sudden unexplained death in young persons with primary epilepsy. Pharmacoepidemiol Drug Safe 1992;1:59-64.

Jokeit H, Ebner A. Long term effects of refractory temporal lobe epilepsy on cognitive abilities: a cross sectional study. J Neurol Neurosurg Psychiatry 1999 Jul;67(1):44-50.

Jooma R, Yeh HS, Privitera MD, et al. Lesionectomy versus electrophysiologically guided resection for temporal lobe tumors manifesting with complex partial seizures. J Neurosurg 1995a Aug;83(2):231-6.

Jooma R, Yeh HS, Privitera MD, et al. Seizure control and extent of mesial temporal resection. Acta Neurochir (Wien) 1995b;133(1-2):44-9.

Joy Andrews D, Reiter JM, Schonfeld W, et al. A neurobehavioral treatment for unilateral complex partial seizure disorders: a comparison of right- and left-hemisphere patients. Seizure 2000 Apr;9(3):189-97.

Jozwiak S, Terczynski A. Open study evaluating lamotrigine efficacy and safety in add-on treatment and consecutive monotherapy in patients with carbamazepine- or valproate-resistant epilepsy. Seizure 2000 Oct;9(7):486-92.

Juhasz C, Chugani DC, Muzik O, et al. Relationship between EEG and positron emission tomography abnormalities in clinical epilepsy. J Clin Neurophysiol 2000 Jan;17(1):29-42.

Kalviainen R. Tiagabine: a new therapeutic option for people with intellectual disability and partial epilepsy. J Intellect Disabil Res 1998 Dec;42(Suppl 1):63-7.

Kalviainen R, Salmenpera T, Partanen K, et al. Recurrent seizures may cause hippocampal damage in temporal lobe epilepsy. Neurology 1998 May;50(5):1377-82.

Kanemoto K, Kim Y, Miyamoto T, et al. Presurgical postictal and acute interictal psychoses are differentially associated with postoperative mood and psychotic disorders. J Neuropsychiatry Clin Neurosci 2001 Spring;13(2):243-7.

Kanner AM PJFM. The 'forgotten' cross-tolerance between phenobarbital and primidone: it can prevent acute primidone-related toxicity. Kanner AM, Parra J, Frey M. The 'forgotten' cross-tolerance between phenobarbital and primidone: it can prevent acute primidone-related toxicity. Epilepsia 2000 Oct;41(10):1310-4.: EPC5cites_everything, Helen_transfer , EPC0005 . Epilepsia. 41. 2000:1310-4.

Karande SC, Joshi MV, Kshirsagar NA, et al. 'An analysis of epileptic patients nonresponsive to drugs'. J Assoc Physicians India 1992 Jul;40(7):445-7.

Kaufman DW, Kelly JP, Anderson T, et al. Evaluation of case reports of aplastic anemia among patients treated with felbamate. Epilepsia 1997 Dec;38(12):1265-9.

Keene DL, Higgins MJ, Ventureyra EC. Outcome and life prospects after surgical management of medically intractable epilepsy in patients under 18 years of age. Childs Nerv Syst 1997 Oct;13(10):530-5.

Keene DL, Whiting S, Ventureyra EC. Electrocorticography. Epileptic Disord 2000 Mar;2(1):57-63.

Kelsey JL, Whittemore AS, Evans AS, et al. Methods in observational epidemiology. 2nd ed. New York (NY): Oxford University Press; 1996. 432 p. (Monographs in epidemiology and biostatistics; vol. 26).

Kendrick AM, Duncan JS, Trimble MR. Effects of discontinuation of individual antiepileptic drugs on mood. Hum Psychopharmacol 1993;8(4):263-270.

Kennedy PA, Schallert G. Practical issues and concepts in vagus nerve stimulation: a nursing review. J Neurosci Nurs 2001 Apr;33(2):105-11.

Ketter TA, Malow BA, Flamini R, et al. Felbamate monotherapy has stimulant-like effects in patients with epilepsy. Epilepsy Res 1996 Mar;23(2):129-37.

Kilpatrick C, Cook M, Kaye A, et al. Non-invasive investigations successfully select patients for temporal lobe surgery. J Neurol Neurosurg Psychiatry 1997 Sep;63(3):327-33.

King DW, Flanigin HF, Gallagher BB, et al. Temporal lobectomy for partial complex seizures: evaluation, results, and 1-year follow-up. Neurology 1986 Mar;36(3):334-9.

Klenerman P, Sander JW, Shorvon SD. Mortality in patients with epilepsy: a study of patients in long term residential care. J Neurol Neurosurg Psychiatry 1993 Feb;56(2):149-52.

Kloster R, Engelskjon T. Sudden unexpected death in epilepsy (SUDEP): a clinical perspective and a search for risk factors. J Neurol Neurosurg Psychiatry 1999 Oct;67(4):439-44.

Kloster R, Larsson PG, Lossius R, et al. The effect of acupuncture in chronic intractable epilepsy. Seizure 1999 May;8(3):170-4.

Kohler CG, Carran MA, Bilker W, et al. Association of fear auras with mood and anxiety disorders after temporal lobectomy. Epilepsia 2001 May;42(5):674-81.

Kotchoubey B, Busch S, Strehl U, et al. Changes in EEG power spectra during biofeedback of slow cortical potentials in epilepsy. Appl Psychophysiol Biofeedback 1999 Dec;24(4):213-33.

Kotchoubey B, Schneider D, Schleichert H, et al. Self-regulation of slow cortical potentials in epilepsy: a retrial with analysis of influencing factors. Epilepsy Res 1996 Nov;25(3):269-76.

Kotchoubey B, Strehl U, Holzapfel S, et al. Negative potential shifts and the prediction of the outcome of neurofeedback therapy in epilepsy. Clin Neurophysiol 1999 Apr;110(4):683-6.

Kral T, Kuczaty S, Blumcke I, et al. Postsurgical outcome of children and adolescents with medically refractory frontal lobe epilepsies. Childs Nerv Syst 2001 Oct;17(10):595-601.

Kumlien E, Doss RC, Gates JR. Treatment outcome in patients with mesial temporal sclerosis. Seizure 2001;0:1-5.

Kuyk J, Spinhoven P, van Dyck R. Hypnotic recall: a positive criterion in the differential diagnosis between epileptic and pseudoepileptic seizures. Epilepsia 1999 Apr;40(4):485-91.

Kwan SY, Wong TT, Chang KP, et al. Seizure outcomes after anterior callosotomy in patients with posterior-dominant and with anterior-dominant epileptiform discharges. Childs Nerv Syst 2001 Jan;17(1-2):71-5.

Labar D, Murphy J, Tecoma E. Vagus nerve stimulation for medication-resistant generalized epilepsy. E04 VNS Study Group. Neurology 1999 Apr;52(7):1510-2.

Labar D, Nikolov B, Tarver B, et al. Vagus nerve stimulation for symptomatic generalized epilepsy: a pilot study. Epilepsia 1998 Feb;39(2):201-5.

Lambert PC, Sutton AJ, Abrams KR, et al. A comparison of summary patient-level covariates in meta-regression with individual patient data meta-analysis. J Clin Epidemiol 2002 Jan;55(1):86-94.

Lancman ME, Asconape JJ, Craven WJ, et al. Predictive value of induction of psychogenic seizures by suggestion. Ann Neurol 1994 Mar;35(3):359-61.

Langtry HD, Gillis JC, Davis R. Topiramate. A review of its pharmacodynamic and pharmacokinetic properties and clinical efficacy in the management of epilepsy. Drugs 1997 Nov;54(5):752-73.

Lantz DL, Sterman MB. Neuropsychological assessment of s ubjects with uncontrolled epilepsy: effects of EEG feedback training. Epilepsia 1988;29(2):163-71.

Laxer KD, Mullooly JP, Howell B. Prolactin changes after seizures classified by EEG monitoring. Neurology 1985 Jan;35(1):31-5.

Lee BI, Markand ON, Stewart WA, et al. Evaluation of patients with intractable seizures for epilepsy surgery at Indiana University Medical Center. Indiana Med 1986 May;79(5):420-4.

Leestma JE, Annegers JF, Brodie MJ, et al. Sudden unexplained death in epilepsy: observations from a large clinical development program. Epilepsia 1997 Jan;38(1):47-55.

Leppik IE. Monotherapy and polypharmacy. Neurology 2000;55(11 Suppl 3):S25-9.

Leppik IE. Rational monotherapy for epilepsy. Baillieres Clin Neurol 1996 Dec;5(4):749-55.

Leppik IE. Tiagabine: the safety landscape. Epilepsia 1995;36(Suppl 6):S10-S13.

Leppik IE, Dreifuss FE, Pledger GW, et al. Felbamate for partial seizures: results of a controlled clinical trial. Neurology 1991 Nov;41(11):1785-9.

Lesser RP, Arroyo S, Crone N, et al. Motor and sensory mapping of the frontal and occipital lobes. Epilepsia 1998;39(Suppl 4):S69-80.

Lesser RP, Fisher RS, Uematsu S. Assessment of surgical outcome. Epilepsy Res 1992(Suppl 5):217-29.

Lesser RP, Pippenger CE, Luders H, et al. High-dose monotherapy in treatment of intractable seizures. Neurology 1984;34:707-11.

Leung GK, Fan YW, Fong KY. Temporal lobe resection for intractable epilepsy: review of 11 cases. Hong Kong Med J 1999 Dec;5(4):329-336.

Levisohn PM. Epilepsy surgery in children with developmental disabilities. Semin Pediatr Neurol 2000 Sep;7(3):194-203.

Lewis DV. Febrile convulsions and mesial temporal sclerosis. Curr Opin Neurol 1999 Apr;12(2):197-201.

Lhatoo SD, Johnson AL, Goodridge DM, et al. Mortality in epilepsy in the first 11 to 14 years after diagnosis: multivariate analysis of a long-term, prospective, population-based cohort. Ann Neurol 2001 Mar;49(3):336-44.

Libman MD, Potvin L, Coupal L, et al. Seizure vs. syncope: measuring serum creatine kinase in the emergency department. J Gen Intern Med 1991;6(5):408-12.

Lieb JP, Engel J Jr, Babb TL. Interhemispheric propagation time of human hippocampal seizures. I. Relationship to surgical outcome. Epilepsia 1986;27(3):286-93.

Lieb JP, Walsh GO, Babb TL, et al. A comparison of EEG seizure patterns recorded with surface and depth electrodes in patients with temporal lobe epilepsy. Epilepsia 1976;17:137-60.

Lijmer JG, Mol BW, Heisterkamp S, et al. Empirical evidence of design-related bias in studies of diagnostic tests. JAMA 1999 Sep;282(11):1061-6.

Lindsay J, Ounsted C, Richards P. Hemispherectomy for childhood epilepsy: a 36-year study. Dev Med Child Neurol 1987 Oct;29(5):592-600.

Liporace J, Hucko D, Morrow R, et al. Vagal nerve stimulation: adjustments to reduce painful side effec ts. Neurology 2001 Sep;57(5):885-6.

Littenberg B, Moses LE. Estimating diagnostic accuracy from multiple conflicting reports: a new meta-analytic method. Med Decis Making 1993 Oct-1993 Dec;13(4):313-21.

Liu MY, Yeh HS, Blisard K, et al. Surgical management of epilepsy associated with developmental anomalies of the brain. Surg Neurol 1995 Feb;43(2):182-8; discussion 188-90.

Loiseau P. Review of controlled trials of gabitril (tiagabine): a clinician's viewpoint. Epilepsia 1999;40(Suppl 9):S14-9.

Loring DW, Meador KJ. Cognitive and behavioral effects of epilepsy treatment. Epilepsia 2001;42(Suppl 8):24-32.

Lundgren J, Amark P, Blennow G, et al. Vagus nerve stimulation in 16 children with refractory epilepsy. Epilepsia 1998 Aug;39(8):809-13.

Lusic I, Pintaric I, Hozo I, et al. Serum prolactin levels after seizure and syncopal attacks. Seizure 1999 Jun;8(4):218-22.

Mackenzie RA, Matheson JM, Smith JS, et al. Surgery for refractory epilepsy. Med J Aust 1990 Jul;153(2):69-72, 76.

Maehara T, Shimizu H. Surgical outcome of corpus callosotomy in patients with drop attacks. Epilepsia 2001 Jan;42(1):67-71.

Maher J, McLachlan RS. Antiepileptic drug treatment following temporal lobectomy. Neurology 1998 Jul;51(1):305-7.

Mak SC, Chi CS, Wan CJ. Clinical experience of ketogenic diet on children with refractory epilepsy. Acta Paediatr Taiwan 1999;40(2):97-100.

Makuch R, Simon R. Sample size requirements for evaluating a conservative therapy. Cancer Treat Rep 1978 Jul;62(7):1037-40.

Manchanda R, Schaefer B, McLachlan RS, et al. Psychiatric disorders in candidates for surgery for epilepsy. J Neurol Neurosurg Psychiatry 1996 Jul;61(1):82-9.

Marino R Jr, Radvany J, Huck FR, et al. Selective electroencephalograph-guided microsurgical callosotomy for refractory generalized epilepsy. Surg Neurol 1990 Oct;34(4):219-28.

Markand ON, Salanova V, Whelihan E, et al. Health-related quality of life outcome in medically refractory epilepsy treated with anterior temporal lobectomy. Epilepsia 2000 Jun;41(6):749-59.

Mathern GW, Giza CC, Yudovin S, et al. Postoperative seizure control and antiepileptic drug use in pediatric epilepsy surgery patients: the UCLA experience, 1986-1997. Epilepsia 1999 Dec;40(12):1740-9.

Matsuo F, Bergen D, Faught E, et al. Placebo-controlled study of the efficacy and safety of lamotrigine in patients with partial seizures. U.S. Lamotrigine Protocol 0.5 Clinical Trial Group. Neurology 1993 Nov;43(11):2284-91.

Matsuzaka T, Ono K, Baba H, et al. Quantitative EEG analyses and surgical outcome after corpus callosotomy. Epilepsia 1999 Sep;40(9):1269-78.

May TW, Bulmahn A, Wohlhuter M, et al. Effects of withdrawal of phenytoin on cognitive and psychomotor functions in hospitalized epileptic patients on polytherapy. Acta Neurol Scand 1992 Aug;86(2):165-70.

McCabe PH, McNew CD, Michel NC. Effect of divalproex-lamotrigine combination therapy in frontal lobe seizures. Arch Neurol 2001 Aug;58(8):1264-8.

McInerney J, Siegel AM, Nordgren RE, et al. Long-term seizure outcome following corpus callosotomy in children. Stereotactic Funct Neurosurg 1999;73(1-4):79-83.

McKee JR, Bodfish JW. Sudden unexpected death in epilepsy in adults with mental retardation. Am J Ment Retard 2000 Jul;105(4):229-35.

McLachlan RS, Maher J. Management of antiepileptic drugs following epilepsy surgery: a review. Can J Neurol Sci 2000 May;27(Suppl 1):S106-10; discussion S121.

McLachlan RS, Rose KJ, Derry PA, et al. Health-related quality of life and seizure control in temporal lobe epilepsy. Ann Neurol 1997 Apr;41(4):482-9.

McLean MJ, Morrell MJ, Willmore LJ, et al. Safety and tolerability of gabapentin as adjunctive therapy in a large, multicenter study. Epilepsia 1999 Jul;40(7):965-72.

McLean MJ, Ramsay RE, Leppik LE, et al. Gabapentin as add-on therapy in refractory partial epilepsy: a double-blind, placebo-controlled, parallel-group study. Neurology 1993;43(11):2292-8.

Meador KJ, Baker GA. Behavioral and cognitive effects of lamotrigine. J Child Neurol 1997 Nov;12(Suppl 1):S44-7.

Messenheimer J, Ramsay RE, Willmore LJ, et al. Lamotrigine therapy for partial seizures: a multicenter, placebo-controlled, double-blind, cross-over trial. Epilepsia 1994;35(1):113-21.

Meyer FB, Marsh WR, Laws ER Jr, et al. Temporal lobectomy in children with epilepsy. J Neurosurg 1986 Mar;64(3):371-6.

Mikati MA, Holmes GL. Lamotrigine in absence and primary generalized epilepsies. J Child Neurol 1997 Nov;12(Suppl 1):S29-37.

Milano Collaborative Group for Studies on Epilepsy. Long-term intensive monitoring in the difficult patient. Preliminary results of 16 months of observations - usefulness and limitations. In: Gardner-Thorpe C, Janz D, Meinardi H, et al, editors. Antiepileptic drug monitring. Kent: Pitman Medical; 1977. p. 197-213.

Miranda C, Smith ML. Predictors of intelligence after temporal lobectomy in children with epilepsy. Epilepsy Behav 2001 Feb;2(1):13-9.

Mirza WU, Credeur LJ, Penry JK. Results of antiepileptic drug reduction in patients with multiple handicaps and epilepsy. Drug Invest 1993;5(6):320-6.

Mishra V, Gahlaut DS, Kumar S. Value of serum prolactin in differentiating epilepsy from pseudoseizures. J Assoc Physicians India 1990;38:846-7.

Mitchell LA, Jackson GD, Kalnins RM, et al. Anterior temporal abnormality in temporal lobe epilepsy: a quantitative MRI and histopathologic study. Neurology 1999 Jan;52(2):327-36.

Mizrahi EM, Kellaway P, Grossman RG, et al. Anterior temporal lobectomy and medically refractory temporal lobe epilepsy of childhood. Epilepsia 1990;31(3):302-12.

Mohan KK, Chugani DC, Chugani HT. Positron emission tomography in pediatric neurology. Semin Pediatr Neurol 1999 Jun;6(2):111-9.

Mohan KK, Markand ON, Salanova V. Diagnostic utility of video EEG monitoring in paroxysmal events. Acta Neurol Scand 1996 Nov;94(5):320-5.

Moran NF, Lemieux L, Maudgil D, et al. Analysis of temporal lobe resections in MR images. Epilepsia 1999 Aug;40(8):1077-84.

Morrell F, Kanner AM, de Toledo-Morrell L, et al. Multiple subpial transection. Adv Neurol 1999;81:259-70.

Morrell F, Whisler WW, Bleck TP. Multiple subpial transection: a new approach to the surgical treatment of focal epilepsy. J Neurosurg 1989 Feb;70(2):231-9.

Morrell F, Whisler WW, Smith MC, et al. Landau-Kleffner syndrome. Treatment with subpial intracortical transection. Brain 1995 Dec;118(Pt 6):1529-46.

Morrell MJ, McLean MJ, Willmore LJ, et al. Efficacy of gabapentin as adjunctive therapy in a large, multicenter study. The Steps Study Group. Seizure 2000 Jun;9(4):241-8.

Morris GL 3rd. Efficacy and tolerability of gabapentin in clinical practice. Clin Ther 1995;17(5):891-900.

Moses LE, Shapiro D, Littenberg B. Combining independent studies of a diagnostic test into a summary ROC curve: data-analytic approaches and some additional considerations. Stat Med 1993 Jul;12(14):1293-316.

Mosewich RK, So EL, O'Brien TJ, et al. Factors predictive of the outcome of frontal lobe epilepsy surgery. Epilepsia 2000 Jul;41(7):843-9.

Mosquera DA, Goldman MD. Surgical audit without autopsy: tales of the unexpected. Ann R Coll Surg Engl 1993 Mar;75(2):115-7.

Mulligan LP, Spencer DD, Spencer SS. Multiple subpial transections: the Yale experience. Epilepsia 2001 Feb;42(2):226-9.

Murro AM, Flanigin HF, Gallagher BB, et al. Corpus callosotomy for the treatment of intractable epilepsy. Epilepsy Res 1988;2(1):44-50.

Murthy JM, Yangala R. Etiological spectrum of symptomatic localization related epilepsies: a study from South India. J Neurol Sci 1998 Jun;158(1):65-70.

Myers JE, Thompson ML. Meta-analysis and occupational epidemiology. Occup Med (Lond) 1998 Feb;48(2):99-101.

Nagakubo S, Niwa S, Kumagai N, et al. Effects of TJ-960 on Sternberg's paradigm results in epileptic patients. Jpn J Psychiatry Neurol 1993 Sep;47(3):609-20.

Nakken KO, Bjorholt PG, Johannessen SI, et al. Effect of physical training on aerobic capacity, seizure occurrence, and serum level of antiepileptic drugs in adults with epilepsy. Epilepsia 1990;31(1):88-94.

Nashef L, Fish DR, Sander JW, et al. Incidence of sudden unexpected death in an adult outpatient cohort with epilepsy at a tertiary referral centre. J Neurol Neurosurg Psychiatry 1995 Apr;58(4):462-4.

Nashef L, Sander JW. Sudden unexpected deaths in epilepsy–where are we now? Seizure 1996 Sep;5(3):235-8.

Naylor AS, Rogvi-Hansen B, Kessing L, et al. Psychiatric morbidity after surgery for epilepsy: short term follow up of patients undergoing amygdalohippocampectomy. J Neurol Neurosurg Psychiatry 1994;57(11):1375-81.

Nees H, Moriarty J, Kitchen ND, et al. Psychosocial and neurobehavioral factors related to surgical treatment for partial epilepsy: a multivariate analysis. Epilepsy Behav 2001 Apr;2(2):135-9.

Neufeld MY, Korczyn AD. New drugs and vagal stimulation for treatment of epilepsy--the Israeli experience. Neurol Neurochir Pol 1996;30 Suppl 2:113-20.

Neufeld MY, Treves TA, Chistik V, et al. Sequential serum creatine kinase determination differentiates vaso-vagal syncope from generalized tonic-clonic seizures. Acta Neurol Scand 1997 Mar;95(3):137-9.

Nilsson L, Farahmand BY, Persson PG, et al. Risk factors for sudden unexpected death in epilepsy: a case-control study. Lancet 1999 Mar;353(9156):888-93.

Nordgren RE, Reeves AG, Viguera AC, et al. Corpus callosotomy for intractable seizures in the pediatric age group. Arch Neurol 1991 Apr;48(4):364-72.

Nordli DR. Epilepsy surgery in children, with special attention to focal cortical resections. Semin Pediatr Neurol 2000 Sep;7(3):204-15.

Nordli DR Jr, Kelley KR. Selection and evaluation of children for epilepsy surgery. Pediatr Neurosurg 2001 Jan;34(1):1-12.

Noseworthy JH, Ebers GC, Vandervoort MK, et al. The impact of blinding on the results of a randomized, placebo-controlled multiple sclerosis clinical trial. Neurology 1994 Jan;44(1):16-20.

Nousiainen U, Suomalainen T, Mervaala E. Clinical benefits of scalp EEG studies in intractable seizure disorders. Acta Neurol Scand 1992 Mar;85(3):181-6.

Novotny E, Ashwal S, Shevell M. Proton magnetic resonance spectroscopy: an emerging technology in pediatric neurology research. Pediatr Res 1998 Jul;44(1):1-10.

Office for National Statistics, General Register Office for Scotland, Northern Ireland Statistics and Research Agency. 7.3 Age-standardised mortality rates: by cause, gender and NHS regional office area 1999. [internet]. United Kingdom: National Health Service; 1999 [cited 2001 Oct 11]. [1 p]. Available: http://www.statistics.gov.uk/statbase/.

Office for National Statistics. Mortality statistics: general. Review of the Registrar General on deaths in England and Wales, 1999. London (UK): The Stationery Office; 2001. (Series DH1; no. 32). Also available: http://www.statistics.gov.uk/statbase.

Oguni H, Olivier A, Andermann F, et al. Anterior callosotomy in the treatment of medically intractable epilepsies: a study of 43 patients with a mean follow-up of 39 months. Ann Neurol 1991 Sep;30(3):357-64.

Ogunmekan AO, Hwang PA. A randomized, double-blind, placebo-controlled, clinical trial of D-alpha-tocopheryl acetate (vitamin E), as add-on therapy, for epilepsy in children. Epilepsia 1989;30(1):84-9.

Ohman R, Walinder J, Balldin J, et al. Prolactin response to electroconvulsive therapy. Lancet 1976 Oct;2(7992):936-7.

Ohtsuka Y, Ogino T, Asano T, et al. Long-term follow-up of vitamin B(6)-responsive West syndrome. Pediatr Neurol 2000 Sep;23(3):202-6.

Ojemann GA. Treatment of temporal lobe epilepsy. Annu Rev Med 1997;48:317-28.

Ojemann GA, Dodrill CB. Verbal memory deficits after left temporal lobectomy for epilepsy. Mechanism and intraoperative prediction. J Neurosurg 1985 Jan;62(1):101-7.

Olivier A. Surgical techniques in temporal lobe epilepsy. Clin Neurosurg 1997;44:211-41.

Olson DM. Evaluation of children for epilepsy surgery. Pediatr Neurosurg 2001 Mar;34(3):159-65.

Orbach D, Romanelli P, Devinsky O, et al. Late seizure recurrence after multiple subpial transections. Epilepsia 2001;49(9):1130-33.

Ozuna J. Ethical dilemmas in people with epilepsy who drive. Clin Nurs Pract Epilepsy 1998 Mar;5(1):7-10.

Pacia SV, Devinsky O, Perrine K, et al. Multiple subpial transection for intractable partial seizures: seizure outcome. J Epilepsy 1997;10(2):86-91.

Palmini A. Medical and surgical strategies for epilepsy care in developing countries. Epilepsia 2000;41(Suppl 4):S10-7.

Palmini A, Andermann F, Olivier A, et al. Focal neuronal migration disorders and intractable partial epilepsy: results of surgical treatment. Ann Neurol 1991 Dec;30(6):750-7.

Panjwani U, Selvamurthy W, Singh SH, et al. Effect of Sahaja yoga practice on seizure control & EEG changes in patients of epilepsy. Indian J Med Res 1996 Mar;103:165-72.

Parker AP, Polkey CE, Binnie CD, et al. Vagal nerve stimulation in epileptic encephalopathies. Pediatrics 1999 Apr;103(4 Pt 1):778-82.

Parrent AG, Blume WT. Stereotactic amygdalohippocampotomy for the treatment of medial temporal lobe epilepsy. Epilepsia 1999 Oct;40(10):1408-16.

Patil AA, Andrews RV, Torkelson R. Surgical treatment of intractable seizures with multilobar or bihemispheric seizure foci (MLBHSF). Surg Neurol 1997 Jan;47(1):72-7; discussion 77-8.

Patrick DL, Deyo RA. Generic and disease-specific measures in assessing health status and quality of life. Med Care 1989 Mar;27(3 Suppl):S217-32.

Peacock WJ, Wehby-Grant MC, Shields WD, et al. Hemispherectomy for intractable seizures in children: a report of 58 cases. Childs Nerv Syst 1996 Jul;12(7):376-84.

Pellock JM. Treatment of seizures and epilepsy in children and adolescents. Neurology 1998 Nov;51(5 Suppl 4):S8-14.

Pellock JM, Watemberg N. New antiepileptic drugs in children: present and future. Semin Pediatr Neurol 1997 Mar;4(1):9-18.

Perry TR, Gumnit RJ, Gates JR, et al. Routine EEG vs. intensive monitoring in the evaluation of intractable epilepsy. Public Health Rep 1983;98(4):384-9.

Perucca E. The management of refractory idiopathic epilepsies. Epilepsia 2001;42(Suppl 3):31-5.

Perucca E. The new generation of antiepileptic drugs: advantages and disadvantages. Br J Clin Pharmacol 1996 Nov;42(5):531-43.

Perucca E. Trial duration and follow-up. Epilepsy Res 2001 May;45(1-3):123-7; discussion 129-31.

Petitti DB. Approaches to heterogeneity in meta-analysis. Stat Med 2001 Dec;20(23):3625-33.

Phelps CE, Hutson A. Estimating diagnostic test accuracy using a 'fuzzy gold standard'. Med Decis Making 1995;15(1):44-57.

Pierelli F, Chatrian GE, Erdly WW, et al. Long-term EEG-video-audio monitoring: detection of partial epileptic seizures and psychogenic episodes by 24-hour EEG record review. Epilepsia 1989;30(5):513-23.

Pinard JM, Delalande O, Chiron C, et al. Callosotomy for epilepsy after West syndrome. Epilepsia 1999 Dec;40(12):1727-34.

Powell GE, Polkey CE, McMillan T. The new Maudsley series of temporal lobectomy. I: Short-term cognitive effects. Br J Clin Psychol 1985 May;24 (Pt 2):109-24.

Pritchard PB 3rd, Wannamaker BB, Sagel J, et al. Serum prolactin and cortisol levels in evaluation of pseudoepileptic seizures. Ann Neurol 1985 Jul;18(1):87-9.

Privitera M, Fincham R, Penry J, et al. Topiramate placebo-controlled dose-ranging trial in refractory partial epilepsy using 600-, 800-, and 1,000-mg daily dosages. Topiramate YE Study Group. Neurology 1996 Jun;46(6):1678-83.

Provinciali L, Del Pesce M, Censori B, et al. Evolution of neuropsychological changes after partial callosotomy in intractable epilepsy. Epilepsy Res 1990 Jul;6(2):155-65.

Provini F, Plazzi G, Lugaresi E. From nocturnal paroxysmal dystonia to nocturnal frontal lobe epilepsy. Clin Neurophysiol 2000 Sep;111(Suppl 2):S2-8.

Pulsifer MB, Gordon JM, Brandt J, et al. Effects of ketogenic diet on development and behavior: preliminary report of a prospective study. Dev Med Child Neurol 2001 May;43(5):301-6.

Purves SJ, Wada JA, Woodhurst WB, et al. Results of anterior corpus callosum section in 24 patients with medically intractable seizures. Neurology 1988 Aug;38(8):1194-201.

Puskarich CA, Whitman S, Dell J, et al. Controlled examination of effects of progressive relaxation training on seizure reduction. Epilepsia 1992;33(4):675-80.

Racoosin JA, Feeney J, Burkhart G, et al. Mortality in antiepileptic drug development programs. Neurology 2001 Feb;56(4):514-9.

Radhakrishnan K, So EL, Silbert PL, et al. Predictors of outcome of anterior temporal lobectomy for intractable epilepsy: a multivariate study. Neurology 1998 Aug;51(2):465-71.

Ransohoff DF, Feinstein AR. Problems of spectrum and bias in evaluating the efficacy of diagnostic tests. N Engl J Med 1978 Oct;299(17):926-30.

Rao MB, Radhakrishnan K. Is epilepsy surgery possible in countries with limited resources? Epilepsia 2000;41 Suppl 4:S31-4.

Rao ML, Stefan H, Bauer J. Epileptic but not psychogenic seizures are accompanied by simultaneous elevation of serum pituitary hormones and cortisol levels. Neuroendocrinology 1989 Jan;49(1):33-9.

Rasmussen T. Tailoring of cortical excisions for frontal lobe epilepsy. Can J Neurol Sci 1991 Nov;18(4 Suppl):606-10.

Rasmussen T, Feindel W. Temporal lobectomy: review of 100 cases with major hippocampectomy. Can J Neurol Sci 1991 Nov;18(4 Suppl):601-2.

Reeves AL, So EL, Evans RW, et al. Factors associated with work outcome after anterior temporal lobectomy for intractable epilepsy. Epilepsia 1997 Jun;38(6):689-95.

Regis Y, Roberts DW. Gamma Knife radiosurgery relative to microsurgery: epilepsy. Stereotactic Funct Neurosurg 1999;72(Suppl 1):11-21.

Reinikainen KJ, Keranen T, Halonen T, et al. Comparison of oxcarbazepine and carbamazepine: a double-blind study. Epilepsy Res 1987 Sep;1(5):284-9.

Reiss WG, Oles KS. Acetazolamide in the treatment of seizures. Ann Pharmacother 1996 May;30(5):514-9.

Reiter JM, Andrews DJ. A neurobehavioral approach for treatment of complex partial epilepsy: efficacy. Seizure 2000 Apr;9(3):198-203.

Renowden SA, Matkovic Z, Adams CB, et al. Selective amygdalohippocampectomy for hippocampal sclerosis: postoperative MR appearance. AJNR Am J Neuroradiol 1995 Oct;16(9):1855-61.

Reutens DC, Bye AM, Hopkins IJ, et al. Corpus callosotomy for intractable epilepsy: seizure outcome and prognostic factors. Epilepsia 1993;34(5):904-9.

Rey E, Treluyer JM, Pons G. Pharmacokinetic optimization of benzodiazepine therapy for acute seizures. Focus on delivery routes. Clin Pharmacokinet 1999 Jun;36(6):409-24.

Rho JM, Sankar R. The pharmacologic basis of antiepileptic drug action. Epilepsia 1999 Nov;40(11):1471-83.

Ring HA, Moriarty J, Trimble MR. A prospective study of the early postsurgical psychiatric associations of epilepsy surgery. J Neurol Neurosurg Psychiatry 1998 May;64(5):601-4.

Roberts DW, Darcey TM. The evaluation and image-guided surgical treatment of the patient with a medically intractable seizure disorder. Neurosurg Clin N Am 1996 Apr;7(2):215-27.

Roberts R, Fitch P. Monitoring at the National Hospital, Queen Square, London. Electroencephalogr Clin Neurophysiol Suppl 1985;37:423-36.

Robinson S, Park TS, Blackburn LB, et al. Transparahippocampal selective amygdalohippocampectomy in children and adolescents: efficacy of the procedure and cognitive morbidity in patients. J Neurosurg 2000 Sep;93(3):402-9.

Rockstroh B, Elbert T, Birbaumer N, et al. Cortical self-regulation in patients with epilepsies. Epilepsy Res 1993 Jan;14(1):63-72.

Rodary C, Com-Nougue C, Tournade MF. How to establish equivalence between treatments: a one-sided clinical trial in paediatric oncology. Stat Med 1989 May;8(5):593-8.

Rohmel J. Therapeutic equivalence investigations: statistical considerations. Stat Med 1998 Aug;17(15-16):1703-14.

Rose KJ, Derry PA, Wiebe S, et al. Determinants of health-related quality of life after temporal lobe epilepsy surgery. Qual Life Res 1996 Jun;5(3):395-402.

Rossi GF, Colicchio G, Marchese E, et al. Callosotomy for severe epilepsies with generalized seizures: outcome and prognostic factors. Acta Neurochir (Wien) 1996;138(2):221-7.

Rossi GF, Pompucci A, Colicchio G, et al. Factors of surgical outcome in tumoural epilepsy. Acta Neurochir (Wien) 1999;141(8):819-24.

Rowan AJ, Siegel M, Rosenbaum DH. Daytime intensive monitoring: comparison with prolonged intensive and ambulatory monitoring. Neurology 1987 Mar;37(3):481-4.

Ryan SW, Forsythe I, Hartley R, et al. Slow release carbamazepine in treatment of poorly controlled seizures. Arch Dis Child 1990 Sep;65(9):930-5.

Sachdeo R, Beydoun A, Schachter S, et al. Oxcarbazepine (Trileptal) as monotherapy in patients with partial seizures. Neurology 2001 Sep;57(5):864-71.

Sachdeo RC, Glauser TA, Ritter F, et al. A double-blind, randomized trial of topiramate in Lennox-Gastaut syndrome. Topiramate YL Study Group. Neurology 1999 Jun;52(9):1882-7.

Sachdeo RC, Leroy RF, Krauss GL, et al. Tiagabine therapy for complex partial seizures. A dose-frequency study. The Tiagabine Study Group. Arch Neurol 1997b May;54(5):595-601.

Sachdeo RC, Reife RA, Lim P, et al. Topiramate monotherapy for partial onset seizures. Epilepsia 1997a Mar;38(3):294-300.

Sachdev P. Schizophrenia-like psychosis and epilepsy: the status of the association. Am J Psychiatry 1998 Mar;155(3):325-36.

Sakas DE, Phillips J. Anterior callosotomy in the management of intractable epileptic seizures: significance of the extent of resection. Acta Neurochir (Wien) 1996;138(6):700-7.

Salanova V, Andermann F, Olivier A, et al. Occipital lobe epilepsy: Electroclinical manifestations, electrocorticography, cortical stimulation and outcome in 42 patients treated between 1930 and 1991. Surgery of occipital lobe epilepsy. Brain 1992;115(6):1655-80.

Salanova V, Andermann F, Rasmussen T, et al. Parietal lobe epilepsy. Clinical manifestations and outcome in 82 patients treated surgically between 1929 and 1988. Brain 1995 Jun;118(Pt 3):607-27.

Salanova V, Andermann F, Rasmussen T, et al. Tumoural parietal lobe epilepsy. Clinical manifestations and outcome in 34 patients treated between 1934 and 1988. Brain 1995;118(5):1289-304.

Salanova V, Markand O, Worth R. Longitudinal follow-up in 145 patients with medically refractory temporal lobe epilepsy treated surgically between 1984 and 1995. Epilepsia 1999 Oct;40(10):1417-23.

Salinsky MC, Uthman BM, Ristanovic RK, et al. Vagus nerve stimulation for the treatment of medically intractable seizures. Results of a 1-year open-extension trial. Vagus Nerve Stimulation Study Group. Arch Neurol 1996 Nov;53(11):1176-80.

Sander JW, Shorvon SD. Incidence and prevalence studies in epilepsy and their methodological problems: a review. J Neurol Neurosurg Psychiatry 1987 Jul;50(7):829-39.

Sass KJ, Novelly RA, Spencer DD, et al. Postcallosotomy language impairments in patients with crossed cerebral dominance. J Neurosurg 1990;72(1):85-90.

Sawhney IM, Robertson IJ, Polkey CE, et al. Multiple subpial transection: a review of 21 cases. J Neurol Neurosurg Psychiatry 1995 Mar;58(3):344-9.

Saygi S, Katz A, Marks DA, et al. Frontal lobe partial seizures and psychogenic seizures: comparison of clinical and ictal characteristics. Neurology 1992 Jul;42(7):1274-7.

Schachter SC. Tiagabine monotherapy in the treatment of partial epilepsy. Epilepsia 1995;36(Suppl 6):S2-S6.

Schachter SC, Vazquez B, Fisher RS, et al. Oxcarbazepine: double-blind, randomized, placebo-control, monotherapy trial for partial seizures. Neurology 1999 Mar;52(4):732-7.

Scharfman HE. Epileptogenesis in the parahippocampal region. Parallels with the dentate gyrus. Ann N Y Acad Sci 2000 Jun;911:305-27.

Schmid-Schoenbein C. Improvement of seizure control by psychological methods in patients with intractable epilepsies. Seizure 1998 Aug;7(4):261-70.

Schmidt D. Reduction of two-drug therapy in intractable epilepsy. Epilepsia 1983b Jun;24(3):368-76.

Schmidt D. Single drug therapy for intractable epilepsy. J Neurol 1983a;229(4):221-6.

Schmidt D, Gram L, Brodie M, et al. Tiagabine in the treatment of epilepsy – a clinical review with a guide for the prescribing physician. Epilepsy Res 2000 Oct;41(3):245-51.

Schmidt D, Jacob R, Loiseau P, et al. Zonisamide for add-on treatment of refractory partial epilepsy: a European double-blind trial. Epilepsy Res 1993 May;15(1):67-73.

Schramm J, Behrens E, Entzian W. Hemispherical deafferentation: an alternative to functional hemispherectomy. Neurosurgery 1995 Mar;36(3):509-15; discussion 515-6.

Schramm J, Kral T, Clusmann H, et al. Transsylvian keyhole functional hemispherectomy. Neurosurgery 2001;49(4):891-901.

Schramm J, Kral T, Grunwald T, et al. Surgical treatment for neocortical temporal lobe epilepsy: clinical and surgical aspects and seizure outcome. J Neurosurg 2001 Jan;94(1):33-42.

Schulz KF, Grimes DA. Blinding in randomised trials: hiding who got what. Lancet 2002 Feb;359:696-700.

Schulz KF, Grimes DA. Case-control studies: research in reverse. Lancet 2002 Feb;359(9304):431-34.

Schwartz RH, Eaton J, Bower BD, et al. Ketogenic diets in the treatment of epilepsy: short-term clinical effects. Dev Med Child Neurol 1989 Apr;31(2):145-51.

Schwartz TH, Bazil CW, Walczak TS, et al. The predictive value of intraoperative electrocorticography in resections for limbic epilepsy associated with mesial temporal sclerosis. Neurosurgery 1997 Feb;40(2):302-9; discussion 309-11.

Seidenberg M. Academic achievement and school performance of children with epilepsy. In: Hermann BP, Seidenberg M, editors. Childhood epilepsies: neuropsychological, psychosocial and intervention aspects. New York (NY): John Wiley & Sons, Ltd.; 1989. p. 105-18.

Semah F, Gimenez F, Longer E, et al. Carbamazepine and its epoxide: an open study of efficacy and side effects after carbamazepine dose increment in refractory partial epilepsy. Ther Drug Monit 1994 Dec;16(6):537-40.

Shah AK, Shein N, Fuerst D, et al. Peripheral WBC count and serum prolactin level in various seizure types and nonepileptic events. Epilepsia 2001;42(11):1472-5.

Sharief M, Viteri C, Ben-Menachem E, et al. Double-blind, placebo-controlled study of topiramate in patients with refractory partial epilepsy. Epilepsy Res 1996 Nov;25(3):217-24.

Sheiner LB. Bioequivalence revisited. Stat Med 1992 Sep;11(13):1777-88.

Shen WK, Edwards WD, Hammill SC, et al. Sudden unexpected nontraumatic death in 54 young adults: a 30-year population-based study. Am J Cardiol 1995 Jul;76(3):148-52.

Shi ZY, Gong BT, Jia YW, et al. The efficacy of electro-acupuncture on 98 cases of epilepsy. J Tradit Chin Med 1987 Mar;7(1):21-2.

Shihabuddin B, Abou-Khalil B, Fakhoury T. The value of combined ambulatory cassette-EEG and video monitoring in the differential diagnosis of intractable seizures. Clin Neurophysiol 1999 Aug;110(8):1452-7.

Shimizu H, Maehara T. Modification of peri-insular hemispherotomy and surgical results. Neurosurgery 2000 Aug;47(2):367-72; discussion 372-3.

Shimizu H, Maehara T. Neuronal disconnection for the surgical treatment of pediatric epilepsy. Epilepsia 2000;41(Suppl 9):28-30.

Shimizu H, Suzuki I, Ishijima B, et al. Multiple subpial transection (MST) for the control of seizures that originated in unresectable cortical foci. Jpn J Psychiatry Neurol 1991 Jun;45(2):354-6.

Shorvon SD. The epidemiology and treatment of chronic and refractory epilepsy. Epilepsia 1996;37(Suppl 2):S1-S3.

Shorvon SD, Reynolds EH. Unnecessary polypharmacy for epilepsy. Br Med J 1977 Jun;1(6077):1635-7.

Sidorenko VN. Effects of the medical resonance therapy music in the complex treatment of epileptic patients. Integr Physiol Behav Sci 2000;35(3):212-7.

Siegel AM, Jobst BC, Thadani VM, et al. Medically intractable, localization-related epilepsy with normal mri: presurgical evaluation and surgical outcome in 43 patients. Epilepsia 2001 Jul;42(7):883-8.

Sigler M, Strassburg HM, Boenigk HE. Effective and safe but forgotten: methsuximide in intractable epilepsies in childhood. Seizure 2001 Mar;10(2):120-4.

Silander HC, Blom S, Malmgren K, et al. Surgical treatment for epilepsy: a retrospective Swedish multicenter study. Acta Neurol Scand 1997;95(6):321-30.

Sillanpaa M. Long-term outcome of epilepsy. Epileptic Disord 2000 Jun;2(2):79-88.

Sillanpaa M, Jalava M, Kaleva O, et al. Long-term prognosis of seizures with onset in childhood. N Engl J Med 1998 Jun;338(24):1715-22.

Sills MA, Forsythe WI, Haidukewych D, et al. The medium chain triglyceride diet and intractable epilepsy. Arch Dis Child 1986 Dec;61(12):1168-72.

Singh UK, Jana UK. Serum prolactin and cortisol in children with some paroxysmal disorders. Indian J Pediatr 1994;61(1):57-61.

Sirven J, Whedon B, Caplan D, et al. The ketogenic diet for intractable epilepsy in adults: preliminary results. Epilepsia 1999 Dec;40(12):1721-6.

Sirven JI, Sperling MR, French JA, et al. Significance of simple partial seizures in temporal lobe epilepsy. Epilepsia 1996 May;37(5):450-4.

Sisodiya SM. Surgery for malformations of cortical development causing epilepsy. Brain 2000 Jun;123(Pt 6):1075-91.

Sisodiya SM, Moran N, Free SL, et al. Correlation of widespread preoperative magnetic resonance imaging changes with unsuccessful surgery for hippocampal sclerosis. Ann Neurol 1997 Apr;41(4):490-6.

Sivenius J, Kalviainen R, Ylinen A, et al. Double-blind study of Gabapentin in the treatment of partial seizures. Epilepsia 1991;32(4):539-42.

Sivenius J, Keranen T, Reinikainen K, et al. Diagnostic evaluation of ambulatory, cassette EEG monitoring in 100 epileptic patients. Acta Neurol Scand 1984;69(Suppl 98):99-100.

Slater JD, Brown MC, Jacobs W, et al. Induction of pseudoseizures with intravenous saline placebo. Epilepsia 1995 Jun;36(6):580-5.

Smith JR, Lee MR, King DW, et al. Results of lesional vs. nonlesional frontal lobe epilepsy surgery. Stereotactic Funct Neurosurg 1997;69(1-4 Pt 2):202-9.

Smith MC. Multiple subpial transection in patients with extratemporal epilepsy. Epilepsia 1998;39(Suppl 4):S81-9.

Snodgrass SR. Ambulatory management of children with epilepsy. Pediatr Ann 1999 Apr;28(4):246-53.

So N, Olivier A, Andermann F, et al. Results of surgical treatment in patients with bitemporal epileptiform abnormalities. Ann Neurol 1989 May;25(5):432-9.

Son YJ, Chung CK, Lee SK, et al. Comparison of localizing values of various diagnostic tests in non-lesional medial temporal lobe epilepsy. Seizure 1999 Dec;8(8):465-70.

Sorenson JM, Wheless JW, Baumgartner JE, et al. Corpus callosotomy for medically intractable seizures. Pediatr Neurosurg 1997 Nov;27(5):260-7.

Sotero de Menezes MA, Connolly M, Bolanos A, et al. Temporal lobectomy in early childhood: the need for long-term follow-up. J Child Neurol 2001 Aug;16(8):585-90.

Specht U, Boenigk HE, Wolf P. Discontinuation of clonazepam after long-term treatment. Epilepsia 1989;30(4):458-63.

Spencer SS. Corpus callosum section and other disconnection procedures for medically intractable epilepsy. Epilepsia 1988;29(Suppl 2):S85-99.

Spencer SS, Spencer DD, Williamson PD, et al. Corpus callosotomy for epilepsy. I. Seizure effects. Neurology 1988 Jan;38(1):19-24.

Sperling MR, Feldman H, Kinman J, et al. Seizure control and mortality in epilepsy. Ann Neurol 1999 Jul;46(1):45-50.

Sperling MR, Lieb JP, Engel J Jr, et al. Prognostic significance of independent auras in temporal lobe seizures. Epilepsia 1989;30(3):322-31.

Sperling MR, O'Connor MJ. Auras and subclinical seizures: characteristics and prognostic significance. Ann Neurol 1990 Sep;28(3):320-8.

Sperling MR, Saykin AJ, Roberts FD, et al. Occupational outcome after temporal lobectomy for refractory epilepsy. Neurology 1995 May;45(5):970-7.

Spilker B, Segreti A. Validation of the phenomenon of regression of seizure frequency in epilepsy. Epilepsia 1984 Aug;25(4):443-9.

Sramka M, Chkhenkeli SA. Clinical experience in intraoperational determination of brain inhibitory structures and application of implanted neurostimulators in epilepsy. Stereotactic Funct Neurosurg 1990;54-55:56-9.

Srikumar G, Bhatia M, Jain S, et al. Usefulness of short term video-EEG monitoring in children with frequent intractable episodes. Neurol India 2000 Mar;48(1):29-32.

Stafstrom CE, Lynch M, Sutula TP. Consequences of epilepsy in the developing brain: implications for surgical management. Semin Pediatr Neurol 2000 Sep;7(3):147-57.

Storzbach D, Binder LM, Salinsky MC, et al. Improved prediction of nonepileptic seizures with combined MMPI and EEG measures. Epilepsia 2000 Mar;41(3):332-7.

Sturniolo MG, Galletti F. Idiopathic epilepsy and school achievement. Arch Dis Child 1994 May;70(5):424-8.

Sutula TP, Sackellares JC, Miller JQ, et al. Intensive monitoring in refractory epilepsy. Neurology 1981 Mar;31(3):243-7.

Swartz BE, Delgado-Escueta AV, Walsh GO, et al. Surgical outcomes in pure frontal lobe epilepsy and foci that mimic them. Epilepsy Res 1998;29(2):97-108.

Szabo CA, Wyllie E, Stanford LD, et al. Neuropsychological effect of temporal lobe resection in preadolescent children with epilepsy. Epilepsia 1998 Aug;39(8):814-9.

Takkouche B, Cadarso-Suarez C, Spiegelman D. Evaluation of old and new tests of heterogeneity in epidemiologic meta-analysis. Am J Epidemiol 1999 Jul;150(2):206-15.

Tallian KB, Nahata MC, Tsao CY. Role of the ketogenic diet in children with intractable seizures. Ann Pharmacother 1998 Mar;32(3):349-61.

Tamer SK. Cognitive and behavioural concerns in epileptic children. Indian J Pediatr 1999;66(6):877-86.

Tanenbaum L, Drayer BP, Anderson RE, et al. Epilepsy. American College of Radiology. ACR Appropriateness Criteria. Radiology 2000 Jun;215 Suppl:459-70.

Tassinari CA, Michelucci R, Chauvel P, et al. Double-blind, placebo-controlled trial of topiramate (600 mg daily) for the treatment of refractory partial epilepsy. Epilepsia 1996 Aug;37(8):763-8.

Tatum WO 4th, Benbadis SR, Vale FL. The neurosurgical treatment of epilepsy. Arch Fam Med 2000;9(10):1142-7.

Tennis P, Cole TB, Annegers JF, et al. Cohort study of incidence of sudden unexplained death in persons with seizure disorder treated with antiepileptic drugs in Saskatchewan, Canada. Epilepsia 1995 Jan;36(1):29-36.

Thadani VM, Williamson PD, Berger R, et al. Successful epilepsy surgery without intracranial EEG recording: criteria for patient selection. Epilepsia 1995 Jan;36(1):7-15.

Theodore WH, Albert P, Stertz B, et al. Felbamate monotherapy: implications for antiepileptic drug development. Epilepsia 1995 Nov;36(11):1105-10.

Theodore WH, Raubertas RF, Porter RJ, et al. Felbamate: a clinical trial for complex partial seizures. Epilepsia 1991;32(3):392-7.

Thompson PJ, Huppert FA. Problems in the development of measures to test cognitive performance in adult epileptic patients. In: Kulig B, Meinardi H, Stores G, editors. Epilepsy and behavior '79. Lisse: Swets & Zeitlinger; 1980. p. 37-42.

Thompson PJ, Trimble MR. Anticonvulsant drugs and cognitive functions. Epilepsia 1982 Oct;23(5):531-44.

Thompson PJ, Trimble MR. Sodium valproate and cognitive functioning in normal volunteers. Br J Clin Pharmacol 1981;12:819-24.

Timmings PL. Sudden unexpected death in epilepsy: a local audit. Seizure 1993 Dec;2(4):287-90.

Tinuper P, Andermann F, Villemure JG, et al. Functional hemispherectomy for treatment of epilepsy associated with hemiplegia: rationale, indications, results, and comparison with callosotomy. Ann Neurol 1988 Jul;24(1):27-34.

Trauner DA. Medium-chain triglyceride (MCT) diet in intractable seizure disorders. Neurology 1985 Feb;35(2):237-8.

Trimble MR. Anticonvulsant-induced psychiatric disorders. The role of forced normalisation. Drug Saf 1996 Sep;15(3):159-66.

Trimble MR. Serum prolactin in epilepsy and hysteria. Br Med J 1978 Dec;2(6153):1682.

Tudehope DI, Harris A, Hawes D, et al. Clinical spectrum and outcome of neonatal convulsions. Aust Paediatr J 1988 Aug;24(4):249-53.

Uhlmann C, Froscher W. Biofeedback treatment in patients with refractory epilepsy: changes in depression and control orientation. Seizure 2001 Jan;10(1):34-8.

U.S. Preventive Services Task Force (USPSTF). Cost-effectiveness analyses of colorectal cancer screening: a systematic review for the US Preventive Services Task Force. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2002. 37 p.

Usiskin SC. The role of counselling in an out-patient epilepsy clinic: a three year study. Seizure 1993 Jun;2(2):111-4.

Uthman BM. Vagus nerve stimulation for seizures. Arch Med Res 2000;31(3):300-3.

Uthman BM, Rowan AJ, Ahmann PA, et al. Tiagabine for complex partial seizures: a randomized, add-on, dose-response trial. Arch Neurol 1998 Jan;55(1):56-62.

Vanderzant CW, Giordani B, Berent S, et al. Personality of patients with pseudoseizures. Neurology 1986;36:664-8.

Velasco F, Velasco M, Jimenez F, et al. Predictors in the treatment of difficult-to-control seizures by electrical stimulation of the centromedian thalamic nucleus. Neurosurgery 2000 Aug;47(2):295-304; discussion 304-5.

Velasco M, Velasco F, Velasco AL, et al. Subacute electrical stimulation of the hippocampus blocks intractable temporal lobe seizures and paroxysmal EEG activities. Epilepsia 2000 Feb;41(2):158-69.

Verma A, Lewis D, VanLandingham KE, et al. Lateralized seizure termination: relationship to outcome following anterior temporal lobectomy. Epilepsy Res 2001 Nov;47(1-2):9-15.

Vickrey B. Mortality in a consecutive cohort of 248 adolescents and adults who underwent diagnostic evaluation for epilepsy surgery. Epilepsia 1997;38(Suppl 11):S67-9.

Vickrey BG, Hays RD, Rausch R, et al. Outcomes in 248 patients who had diagnostic evaluations for epilepsy surgery. Lancet 1995 Dec;346(8988):1445-9.

Victor M, Ropper AH. Adams and Victor's principles of neurology. 7th ed. New York (NY): McGraw-Hill Companies, Inc.; 2001. Epilepsy and other seizure disorders. p. 331-65.

Villemure JG, de Tribolet N. Epilepsy in patients with central nervous system tumors. Curr Opin Neurol 1996 Dec;9(6):424-8.

Vining EP, Freeman JM, Ballaban-Gil K, et al. A multicenter study of the efficacy of the ketogenic diet. Arch Neurol 1998 Nov;55(11):1433-7.

Vining EP, Freeman JM, Pillas DJ, et al. Why would you remove half a brain? The outcome of 58 children after hemispherectomy-the Johns Hopkins experience: 1968 to 1996. Pediatrics 1997 Aug;100(2 Pt 1):163-71.

Vinters HV, Park SH, Johnson MW, et al. Cortical dysplasia, genetic abnormalities and neurocutaneous syndromes. Dev Neurosci 1999 Nov;21(3-5):248-59.

Virudhagirinathan BS, Biswakumar B, Sultana MS. Psychological management of temporal lobe epilepsy. Indian J Clin Psychol 1986 Sep;13(2):119-24.

Visudhiphan P, Bunyaratavej S, Visudtibhan A, et al. Temporal lobectomy for intractable complex partial seizures in pediatric patients. J Med Assoc Thai 1999 Aug;82(8):778-83.

Vossler DG, Wilkus RJ, Ojemann GA. Preoperative EEG correlates of seizure outcome from epilepsy surgery in children. J Epilepsy 1995;8(3):236-45.

Wada JA. Differential diagnosis of epilepsy. Electroencephalogr Clin Neurophysiol Suppl 1985;37:285-311.

Walczak TS, Leppik IE, D'Amelio M, et al. Incidence and risk factors in sudden unexpected death in epilepsy: a prospective cohort study. Neurology 2001 Feb;56(4):519-25.

Walczak TS, Radtke RA, McNamara JO, et al. Anterior temporal lobectomy for complex partial seizures: evaluation, results, and long-term follow-up in 100 cases. Neurology 1990 Mar;40(3 Pt 1):413-8.

Walczak TS, Williams DT, Berten W. Utility and reliability of placebo infusion in the evaluation of patients with seizures. Neurology 1994 Mar;44(3 Pt 1):394-9.

Walker IA, Slovis CM. Lidocaine in the treatment of status epilepticus. Acad Emerg Med 1997 Sep;4(9):918-22.

Wang T. Effects of Chinese medicine zhenxianling in 239 cases of epilepsy. J Tradit Chin Med 1996 Jun;16(2):94-7.

Weller SC, Mann NC. Assessing rater performance without a 'Gold Standard' using consensus theory. Med Decis Making 1997;17(1):71-9.

Welty TE, Privitera M, Shukla R. Increased seizure frequency associated with felbamate withdrawal in adults. Arch Neurol 1998 May;55(5):641-5.

Wennberg R, Quesney LF, Lozano A, et al. Role of electrocorticography at surgery for lesion-related frontal lobe epilepsy. Can J Neurol Sci 1999 Feb;26(1):33-9.

Westerveld M, Sass KJ, Chelune GJ, et al. Temporal lobectomy in children: cognitive outcome. J Neurosurg 2000 Jan;92(1):24-30.

White HS, Wolf HH, Woodhead JH, et al. The National Institutes of Health Anticonvulsant Drug Development Program: screening for efficacy. Adv Neurol 1998;76:29-39.

Whitman S, Dell J, Legion V, et al. Progressive relaxation for seizure reduction. J Epilepsy 1990;3(1):17-22.

Wiebe S, Blume WT, Girvin JP, et al. A randomized, controlled trial of surgery for temporal-lobe epilepsy. N Engl J Med 2001 Aug;345(5):311-8.

Wieser HG. Selective amygdalohippocampectomy: indications and follow-up. Can J Neurol Sci 1991 Nov;18(4 Suppl):617-27.

Wieser HG, Blume WT, Fish D, et al. ILAE Commission Report. Proposal for a new classification of outcome with respect to epileptic seizures following epilepsy surgery. Epilepsia 2001 Feb;42(2):282-6.

Wijdicks EF, Scott JP. Causes and outcome of mechanical ventilation in patients with hemispheric ischemic stroke. Mayo Clin Proc 1997;72(3):210-3.

Williamson PD, Spencer DD, Spencer SS, et al. Complex partial seizures of frontal lobe origin. Ann Neurol 1985 Oct;18(4):497-504.

Willmer JP, Brunet DG. The value of prolonged electroencephalographic and video monitoring in diagnosis of seizure disorders. Can J Neurol Sci 1986 Nov;13(4):327-30.

Willmore LJ, Shu V, Wallin B. Efficacy and safety of add-on divalproex sodium in the treatment of complex partial seizures. The M88-194 Study Group. Neurology 1996 Jan;46(1):49-53.

Wilson SJ, Bladin PF, Saling MM, et al. The longitudinal course of adjustment after seizure surgery. Seizure 2001 Apr;10(3):165-72.

Wisniewski KE. Pheno/genotypic correlations of neuronal ceroid lipofuscinoses. Neurology 2001 Aug;57(4):576-81.

Wolf P. Treatment of epilepsy following rejection of epilepsy surgery. Seizure 1998 Feb;7(1):25-9.

Wong IC, Chadwick DW, Fenwick PB, et al. The long-term use of gabapentin, lamotrigine, and vigabatrin in patients with chronic epilepsy. Epilepsia 1999;40(10):1439-45.

Wong IC, Mawer GE, Sander JW. Adverse event monitoring in lamotrigine patients: a pharmacoepidemiologic study in the United Kingdom. Epilepsia 2001 Feb;42(2):237-44.

Wroe SJ, Henley R, John R, et al. The clinical value of serum prolactin measurement in the differential diagnosis of complex partial seizures. Epilepsy Res 1989;3(3):248-52.

Wurm G, Wies W, Schnizer M, et al. Advanced surgical approach for selective amygdalohippocampectomy through neuronavigation. Neurosurgery 2000 Jun;46(6):1377-82; discussion 1382-3.

Wyler AR. Recent advances in epilepsy surgery: temporal lobectomy and multiple subpial transections. Neurosurgery 1997 Dec;41(6):1294-301; discussion 1301.

Wyler AR, Hermann BP, Somes G. Extent of medial temporal resection on outcome from anterior temporal lobectomy: a randomized prospective study. Neurosurgery 1995 Nov;37(5):982-90; discussion 990-1.

Wyllie E. Surgery for catastrophic localization-related epilepsy in infants. Epilepsia 1996;37(Suppl 1):S22-5.

Wyllie E, Comair YG, Kotagal P, et al. Seizure outcome after epilepsy surgery in children and adolescents. Ann Neurol 1998 Nov;44(5):740-8.

Wyllie E, Luders H, MacMillan JP, et al. Serum prolactin levels after epileptic seizures. Neurology 1984 Dec;34(12):1601-4.

Yagi K. Epilepsy: comprehensive care, quality of life, and factors preventing people with epilepsy from being employed. Clin Ther 1998;20(Suppl A):A19-29.

Yamamoto LG, Yim GK. The role of intravenous valproic acid in status epilepticus. Pediatr Emerg Care 2000 Aug;16(4):296-8.

Yeh HS, Kashiwagi S, Tew JM Jr, et al. Surgical management of epilepsy associated with cerebral arteriovenous malformations. J Neurosurg 1990 Feb;72(2):216-23.

Yerby MS, van Belle G, Friel PN, et al. Serum prolactins in the diagnosis of epilepsy: sensitivity, specificity, and predictive value. Neurology 1987 Jul;37(7):1224-6.

Zaidi A, Clough P, Cooper P, et al. Misdiagnosis of epilepsy: many seizure-like attacks have a cardiovascular cause. J Am Coll Cardiol 2000 Jul;36(1):181-4.

Zelnik N, Kahana L, Rafael A, et al. Prolactin and cortisol levels in various paroxysmal disorders in childhood. Pediatrics 1991 Sep;88(3):486-9.

Zupanc ML. Neuroimaging in the evaluation of children and adolescents with intractable epilepsy: I. Magnetic resonance imaging and the substrates of epilepsy. Pediatr Neurol 1997 Jul;17(1):19-26.