References

1. MacMahon S, Peto R Cutler J, Collins R, Sorlie P, Neaton J, Abbott R, Godwin J, Dyer A, and Stamler J. Blood pressure, stroke, and coronary heart disease. Part 1, Prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. Lancet 1990;335(8692):765-74.

2. Stamler J, Stamler R, and Neaton JD. Blood pressure, systolic and diastolic, and cardiovascular risks. US population data. Arch Intern Med 1993;153(5):598-615.

3. Burt VL, Whelton P, Roccella EJ, Brown C, Cutler JA, Higgins M, Horan MJ, and Labarthe D. Prevalence of hypertension in the US adult population. Results from the Third National Health and Nutrition Examination Survey, 1988-1991. Hypertension 1995;25(3):305-13.

4. Collins R, Peto R, MacMahon S, Hebert P, Fiebach NH, Eberlein KA, Godwin J, Qizilbash N, Taylor JO, and Hennekens CH. Blood pressure, stroke, and coronary heart disease. Part 2, Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. Lancet 1990;335(8693):827-38.

5. Kostis JB, Davis BR, Cutler J, Grimm RH Jr, Berge KG, Cohen JD, Lacy CR, Perry HM Jr, Blaufox MD, Wassertheil-Smoller S, Black HR, Schron E, Berkson DM, Curb JD, Smith WM, McDonald R, and Applegate WB. Prevention of heart failure by antihypertensive drug treatment in older persons with isolated systolic hypertension. SHEP Cooperative Research Group. JAMA 1997; 278(3):212-6.

6. Appel LJ. The role of diet in the prevention and treatment of hypertension. Curr Atheroscler Rep 2000;2(6):521-8.

7. The sixth report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. Arch Intern Med 1997;157(21):2413-46.

8. Sennett C. Implementing the new HEDIS hypertension performance measure. Manag Care 2000;9(4 Suppl):2-17; quiz 18-21.

9. Appel LJ and Stason WB. Ambulatory blood pressure monitoring and blood pressure self-measurement in the diagnosis and management of hypertension. Ann Intern Med 1993;118(11): 867-82.

10. Pickering TG, Coats A, Mallion JM, Mancia G, and Verdecchia P. Blood Pressure Monitoring. Task force V: White-coat hypertension. Blood Press Monit 1999 ;4(6):333-41.

11. Beevers G, Lip GY, and O'Brien E. Blood pressure measurement. Part ii-conventional sphygmomanometry: technique of auscultatory blood pressure measurement. BMJ 2001;322(7293):1043-1047.

12. Jones DW, Frohlich ED, Grim CM, Grim CE, and Taubert KA. Mercury Sphygmomanometers Should Not be Abandoned: An Advisory Statement From the Council for High Blood Pressure Research, American Heart Association. Hypertension 2001;37(2):185-186.

13. O'Brien E, Waeber B, Parati G, Staessen J, and Myers MG. Blood pressure measuring devices: recommendations of the European Society of Hypertension. BMJ 2001;322(7285):531-536.

14. O'Brien E, Beevers G, and Lip GY. Blood pressure measurement. Part iv-automated sphygmomanometry: self blood pressure measurement. BMJ 2001;322(7295):1167-70 .

15. Verdecchia P, Schillaci G, and Porcellati C. Dippers versus non-dippers. J Hypertens Suppl 1991;9(8):S42-4.

16. Carr AA, Bottini PB and Prisant LM. Ambulatory blood pressure monitoring for evaluation and management of hypertensives: effect on outcome and cost effectiveness. J Clin Pharmacol 1992 Jul;32 (7):610-3.

17. Dickersin K , Manheimer E, Wieland L, Robinson K, Lefebvre C, and McDonald S. Development of a centralized register of controlled clinical trials: The Cochrane Collaboration's CENTRAL. Evaluation and the Health Professions Supplement Issue: The Cochrane Collaboration 2002;25(1): 38-64.

18. Antivalle M, Lattuada S, Paravicini M, Rindi M, and Libretti A. Twenty-four hour non-invasive ambulatory blood pressure monitoring in the assessment of early hypertension. J Hypertens 1986;4(Suppl 5):S322-S324.

19. Melina D, Colivicchi F, Melina G, and Pristipino C. Left ventricular hypertrophy and diastolic dysfunction in alcohol-associated hypertension. Minerva Cardioangiol 1993;41(7-8):293-6.

20. Pickering TG, Mann SJ, and James GD. Clinic and ambulatory blood pressure measurements for the evaluation of borderline hypertension in smokers and non-smokers. Arch Mal Coeur Vaiss 1991;84 Spec No 3:17-9.

21. Abe H, Yokouchi M, Saitoh F, Deguchi F, Kimura G, Kojima S, Yoshimi H, Ito K, Kuramochi M, Ikeda M and others. Hypertensive complications and home blood pressure: comparison with blood pressure measured in the doctor's office. J Clin Hypertens 1987;3(4):661-9.

22. Jula A, Puukka P, and Karanko H. Multiple clinic and home blood pressure measurements versus ambulatory blood pressure monitoring. Hypertension 1999 ;34(2):261-6.

23. Mengden T, Battig B, and Vetter W. Self-measurement of blood pressure improves the accuracy and reduces the number of subjects in clinical trials. J Hypertens Suppl 1991;9(6):S336-7.

24. Nielsen PE, Myschetzky P, Andersen AR, and Andersen GS. Home readings of blood pressure in assessment of hypertensive subjects. Acta Med Scand Suppl 1986;714:147-51.

25. Stergiou GS , Skeva II, Zourbaki AS, and Mountokalakis TD. Self-monitoring of blood pressure at home: how many measurements are needed? J Hypertens 1998b;16(6):725-31.

26. Weisser B, Grune S, Burger R, Blickenstorfer H, Iseli J, Michelsen SH, Opravil R, Rageth S, Sturzenegger ER, Walker P and others. The Dubendorf Study: a population-based investigation on normal values of blood pressure self-measurement. J Hum Hypertens 1994;8(4):227-31.

27. Ironson GH, Gellman MD, Spitzer SB, Llabre MM, De Carlo Pasin R, Weidler DJ, and Schneiderman N. Predicting home and work blood pressure measurements from resting baselines and laboratory reactivity in black and white Americans. Psychophysiology 1989;26(2):174-84.

28. Khoury S, Yarows SA, O'Brien TK, and Sowers JR. Ambulatory blood pressure monitoring in a nonacademic setting. Effects of age and sex. Am J Hypertens 1992;5(9):616-23.

29. Modesti PA, Pieri F, Cecioni I, Valenti R, Mininni S, Toccafondi S, Vocioni F, Salvati G, Gensini GF, and Neri Serneri GG. Comparison of ambulatory blood pressure monitoring and conventional office measurement in the workers of a chemical company. Int J Cardiol 1994;46(2):151-7.

30. Myers MG, Oh PI, Reeves RA, and Joyner CD. Prevalence of white coat effect in treated hypertensive patients in the community. Am J Hypertens 1995b;8(6):591-7.

31. Narkiewicz K, Piccolo D, Borella P, Businaro R, Zonzin P, and Palatini P. Response to orthostatic stress predicts office-daytime blood pressure difference, but not nocturnal blood pressure fall in mild essential hypertensives: results of the harvest trial. Clin Exp Pharmacol Physiol 1995;22(10):743-7.

32. Staessen JA, Thijs L, Fagard R, O'Brien ET, Clement D, de Leeuw PW, Mancia G, Nachev C, Palatini P, Parati G, Tuomilehto J, and Webste, J. Predicting cardiovascular risk using conventional vs ambulatory blood pressure in older patients with systolic hypertension. Systolic Hypertension in Europe Trial Investigators. JAMA 1999;282(6):539-46.

33. Thijs L, Celis H, Clement D, Gil-Extremera B, Kawecka-Jaszcz K, Mancia G, Parati G, Salvetti A, Sarti C, van den Meiracker AH, O'Brien E, Staessen JA, and Fagard R. Conventional and ambulatory blood pressure measurement in older patients with isolated systolic hypertension:second progress report on the ambulatory blood pressure monitoring project in the Syst-Eur trial. Blood Press Monit 1996;1(2):95-103.

34. Zachariah PK, Sheps SG, Ilstrup DM, Long CR, Bailey KR, Wiltgen CM, and Carlson CA. Blood pressure load--a better determinant of hypertension. Mayo Clin Proc 1988;63(11):1085-91.

35. Zachariah PK, Sheps SG, Bailey KR, Wiltgen CM, and Moore AG. Age-related characteristics of ambulatory blood pressure load and mean blood pressure in normotensive subjects. JAMA 1991;265(11):1414-7.

36. Zawadzka A, Bird R, Casadei B, and Conway J. Audit of ambulatory blood pressure monitoring in the diagnosis and management of hypertension in practice. J Hum Hypertens 1998;12(4):249-52.

37. Sega G, Bravi C, Cesana G, Valagussa F, Mancia G, and Zanchetti A. Ambulatory and home blood pressure normality: the Pamela Study. J Cardiovasc Pharmacol 1994;23 Suppl 5:S12-5.

38. Stergiou GS, Skeva II, Baibas NM, Kalkana CB, Roussias LG, and Mountokalakis TD. Diagnosis of hypertension using home or ambulatory blood pressure monitoring: comparison with the conventional strategy based on repeated clinic blood pressure measurements. J Hypertens 2000;18(12):1745-51.

39. Manning G, Rushton L, and Millar-Craig MW. Clinical implications of white coat hypertension: an ambulatory blood pressure monitoring study. J Hum Hypertens 1999;13(12):817-22.

40. MacDonald MB, Laing GP, Wilson MP, and Wilson TW. Prevalence and predictors of white-coat response in patients with treated hypertension. CMAJ 1999;161(3):265-9.

41. Owens P, Atkins N, and O'Brien E. Diagnosis of white coat hypertension by ambulatory blood pressure monitoring. Hypertension 1999;34(2):267-72.

42. Tochikubo O, Miyajima E, Shigemasa T, and Ishii M. Relation between body fat-corrected ECG voltage and ambulatory blood pressure in patients with essential hypertension. Hypertension 1999;33(5):1159-63.

43. Martinez MA, Garcia-Puig J, Martin JC, Guallar-Castillon P, Aguirre de Carcer A, Torre A, Armada E, Nevado A, and Madero RS. Frequency and determinants of white coat hypertension in mild to moderate hypertension: a primary care-based study. Monitorizacion Ambulatoria de la Presion Arterial (MAPA)-Area 5 Working Group. Am J Hypertens 1999;12(3):251-9.

44. Inden Y, Tsuda M, Hayashi H, Takezawa H, Iino S, Kondo T, Yoshida Y, Akahoshi,M, Terasawa M, Itoh, T Saito, H, and Hirai M. Relationship between Joint National Committee-VI classification of hypertension and ambulatory blood pressure in patients with hypertension diagnosed by casual blood

pressure. Clin Cardiol 1998;21(11):801-6.

45. Stergiou GS , Zourbaki AS, Skeva II, and Mountokalakis TD. White coat effect detected using self-monitoring of blood pressure at home: comparison with ambulatory blood pressure. Am J Hypertens 1998a;11(7):820-7.

46. Myers MG and Reeves RA. White coat effect in treated hypertensive patients: sex differences. J Hum Hypertens 1995a;9(9):729-33.

47. Verdecchia P, Schillaci G, Borgioni C, Ciucci A, Zampi I, Gattobigio R, Sacchi N, and Porcellati C. White coat hypertension and white coat effect. Similarities and differences. Am J Hypertens 1995;8(8):790-8.

48. Pierdomenico SD, Mezzetti A, Lapenna D, Guglielmi MD, Mancini M, Salvatore L, Antidormi T, Costantini F, and Cuccurullo F. 'White-coat' hypertension in patients with newly diagnosed hypertension: evaluation of prevalence by ambulatory monitoring and impact on cost of health care. Eur Heart J 1995;16(5):692-7.

49. Verdecchia P, Schillaci G, Boldrini F, Guerrieri, M, and Porcellati, C. Sex, cardiac hypertrophy and diurnal blood pressure variations in essential hypertension. J Hypertens 1992;10(7):683-92.

50. Martinez MA, Moreno A, Aguirre de Carcer A, Cabrera R, Rocha R, Torre A, Nevado A, Ramos T, Neri J, Anton G, Miranda I, Fernande, P, Rodriguez E, Miquel A, Martinez JL, Rodriguez M, Eisman C, and Puig JG. Frequency and determinants of microalbuminuria in mild hypertension: a primary-care-based study. MAPA--Madrid Working Group. J Hypertens 2001;19(2):319-26.

51. Helmers KF, Baker B, O'Kelly B, and Tobe S. Anger expression, gender, and ambulatory blood pressure in mild, unmedicated adults with hypertension. Ann Behav Med 2000;22(1):60-4.

52. Aylett M, Marples G, and Jones K. Home blood pressure monitoring: its effect on the management of hypertension in general practice. Br J Gen Pract 1999;49(446):725-8.

53. Hoegholm A, Kristensen KS, Bang LE, and Gustavsen PH. White coat hypertension and blood pressure variability. Am J Hypertens 1999;12(1O Pt 1):966-72.

54. Verdecchia P, Schillaci G, Boldrini F, Zampi I, and Porcellati C. Variability between current definitions of 'normal' ambulatory blood pressure. Implications in the assessment of white coat hypertension. Hypertension 1992;20(4):555-62.

55. Palatini P, Dorigatti F, Roman E, Giovinazzo P, Piccolo D, De Venuto G, Mattarei M, Cozzutti E, Gregori S, Mormino P, and Pessina AC. White-coat hypertension: a selection bias? Harvest Study Investigators. Hypertension and Ambulatory Recording Venetia Study. J Hypertens 1998;16(7):977-84.

56. Verdecchia P, Schillaci G, Borgioni C, Ciucci A, Gattobigio R, Sacchi N, Guerrieri M, Comparato E, and Porcellati C. Identification of subjects with white-coat hypertension and persistently normal ambulatory blood pressure. Blood Press Monit 1996;1(3):217-22.

57. Gosse P, Bougaleb M, and Clementy J. Long term reproducibility of ambulatory blood pressure monitoring. Therapie 1996;51(1):5-9.

58. Sahn DJ, DeMaria A, Kisslo J, and Weyman A. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. Circulation 1978 Dec;58(6):1072-83.

59. Devereux RB, Alonso DR, Lutas EM, Gottlieb GJ, Campo E, Sachs I, and Reichek N. Echocardiographic assessment of left

ventricular hypertrophy: comparison to necropsy findings. Am J Cardiol 1986 Feb 15;57(6):450-8.

60. Ohkubo T, Imai Y, Tsuji I, Nagai K, Kato J, Kikuch, N, Nishiyama A, Aihara A, Sekino M, Kikuya M, Ito S, Satoh H, and Hisamichi S. Home blood pressure measurement has a stronger predictive power for mortality than does screening blood pressure measurement: a population-based observation in Ohasama, Japan. J Hypertens 1998;16(7):971-5.

61. Sakuma M, Imai Y, Tsuji I, Naga K, Ohkubo T, Watanabe N, Sakum, H, Satoh H, and Hisamichi S. Predictive value of home blood pressure measurement in relation to stroke morbidity: a population-based pilot study in Ohasama, Japan. Hypertens Res 1997;20(3):167-74.

62. Bailey B, Carney SL, Gillies AA, and Smith AJ. Antihypertensive drug treatment: a comparison of usual care with self blood pressure measurement. J Hum Hypertens 1999;13(2):147-50.

63. Binstock M L and Franklin KL. A comparison of compliance techniques on the control of high blood pressure. Am J Hypertens 1988;1(3 Pt 3):192S-194S.

64. Carnahan JE and Nugent CA. The effects of self-monitoring by patients on the control of hypertension. Am J Med Sci 1975;269(1):69-73.

65. Earp JA, Ory MG, and Strogatz DS. The effects of family involvement and practitioner home visits on the control of hypertension. Am J Public Health 1982;72(10):1146-54.

66. Friedman RH, Kazis LE, Jette A, Smith MB, Stollerman J, Torgerson J, and Carey K. A telecommunications system for monitoring and counseling patients with hypertension. Impact on medication adherence and blood pressure control . Am J Hypertens 1996;9(4 Pt 1):285-92.

67. Johnson AL, Taylor DW, Sackett DL, Dunnett CW, and Shimizu AG. Self-recording of blood pressure in the management of hypertension. Can Med Assoc J 1978;119(9):1034-9.

68. Lehner, H, Kaluza K, Vetter H, Losse H, and Dorst K. Long-term effects of a complex behavioral treatment of essential hypertension. Psychosom Med 1987;49(4):422-30.

69. Midanik LT, Resnick B, Hurley LB, Smith EJ, and McCarthy M. Home blood pressure monitoring for mild hypertensives. Public Health Rep 1991;106(1):85-9.

70. Rogers MA, Small D, Buchan DA, Butch CA, Stewart CM, Krenzer BE, and HusovskyHL. Home monitoring service improves mean arterial pressure in patients with essential hypertension. A randomized, controlled trial. Ann Intern Med 2001;134(11):1024-32.

71. Soghikian K, Casper SM, Fireman BH, Hunkeler EM, Hurley LB, Tekawa IS, and Vog TM. Home blood pressure monitoring. Effect on use of medical services and medical care costs. Med Care 1992;30 (9):855-65.

72. Stahl SM, Kelley CR, Neill PJ, Grim CE, and Mamlin J. Effects of home blood pressure measurement on long-term BP control. Am J Public Health 1984;74(7):704-9.

73. Vetter W, Hess,L, and Brignoli R. Influence of self-measurement of blood pressure on the responder rate in hypertensive patients treated with losartan: results of the SVATCH Study. Standard vs Automatic Treatment Control of COSAAR in Hypertension. J Hum Hypertens 2000;14(4):235-41.

74. Cuspidi C, Lonati L, Sampieri L, Macca G, Michev I, Salerno M, Fusi V, Leonetti G, and Zanchetti A. Impact of blood pressure control on prevalence of left ventricular hypertrophy in treated hypertensive patients.

Cardiology 2000;93(3):149-54.

75. Zakopoulos NA, Toumanidis ST, Barlas GJ, Nanas SN, Lekakis JP, Stamatelopoulos SF, and Moulopoulos SD. A pressure-time index' for assessing the severity of essential hypertension. J Hypertens 1999;17(10):1387-93.

76. Bauduceau B, Genes N, Chamontin B, Vaur L, Renault M, Etienne S, and Marre M. Ambulatory blood pressure and urinary albumin excretion in diabetic (non-insulin-dependent and insulin-dependent) hypertensive patients: relationships at baseline and after treatment by the angiotensin converting enzyme inhibitor trandolapril. Am J Hypertens 1998;11(9):1065-73.

77. Palatini P, Mormino P, Santonastaso M, Mos L, Dal Foll M, Zanata G, and Pessina AC. Target-organ damage in stage I hypertensive subjects with white coat and sustained hypertension: results from the HARVEST study. Hypertension 1998;31(1):57-63.

78. Ferrara LA, Guida L, Pasanisi F, Celentano A, Palmieri V, Iannuzzi R, Gaeta I, Leccia G, and Crivaro M. Isolated office hypertension and end-organ damage. J Hypertens 1997;15(9):979-85.

79. Gosse, P, Ansoborlo P, Lemetayer P, and Clementy J. Left ventricular mass is better correlated with arising blood pressure than with office or occasional blood pressure. Am J Hypertens 1997;10 (5 Pt 1):505-10.

80. Pose-Reino A, Gonzalez-Juanatey JR, Pastor C, Mendez I, Estevez JC, Alvarez D, Valdes L, and Cabezas-Cerrato J. Clinical implications of white coat hypertension. Blood Press 1996;5(5):264-73.

81. Redon, J Baldo E, Lurbe E, Bertolin V, Lozano JV, Miralles A, and Pascual JM. Microalbuminuria, left ventricular mass and ambulatory blood pressure in essential hypertension. Kidney Int Suppl 1996;55:S81-4.

82. Pierdomenico SD, Lapenna D, Guglielmi MD, Antidormi T, Schiavone C, Cuccurullo F, and Mezzetti A. Target organ status and serum lipids in patients with white coat hypertension. Hypertension 1995;26(5):801-7.

83. Lemne C, Lindvall K, Georgiades A, Fredrikson M, and de Faire U. Structural cardiac changes in relation to 24-h ambulatory blood pressure levels in borderline hypertension. J Intern Med 1995;238(1):49-57.

84. Chen CH, Ting CT, Lin SJ, Hsu TL, Chou P, Kuo HS, Wang SP, Yin FC, and Chang MS. Relation between diurnal variation of blood pressure and left ventricular mass in a Chinese population. Am J Cardiol 1995;75(17):1239-43.

85. Weber MA, Neutel JM, Smith DH, and Graettinger WF. Diagnosis of mild hypertension by ambulatory blood pressure monitoring. Circulation 1994;90(5):2291-8.

86. Redon J, Liao Y, Lozano JV, Miralles A, Pascual JM, and Cooper RS. Ambulatory blood pressure and microalbuminuria in essential hypertension: role of circadian variability. J Hypertens 1994;12 (8):947-53.

87. Gosse P, Promax H, Durande, P, and Clementy J. 'White coat' hypertension. No harm for the heart. Hypertension 1993;22(5):766-70.

88. Hansen KW, Christensen CK, Andersen PH, Pedersen MM, Christiansen JS, and Mogensen CE. Ambulatory blood pressure in microalbuminuric type 1 diabetic patients. Kidney Int 1992;41(4):847-54.

89. Verdecchia P, Schillaci G, Guerrieri M, Gatteschi C, Benemio G, Boldrini F, and Porcellati C. Circadian blood pressure changes and left ventricular hypertrophy in essential hypertension. Circulation 1990;81(2):528-36.

90. Baguet JP, De Gaudemaris R, Antoniadis A, Tremel F, Siche JP, and Mallion JM. Use of

ambulatory blood pressure monitoring data to predict left ventricular mass in hypertension. Blood Press Monit 2001;6(2):73-80.

91. Devereux RB , Pickering TG, Harshfield GA, Kleinert HD, Denby L, Clark L, Pregibon D, Jason M, Kleiner B, Borer JS, and Laragh JH. Left ventricular hypertrophy in patients with hypertension: importance of blood pressure response to regularly recurring stress. Circulation 1983;68(3):470-6.

92. Hoegholm A, Bang LE, Kristensen KS, Nielsen JW , and Holm J. Microalbuminuria in 411 untreated individuals with established hypertension, white coat hypertension, and normotension. Hypertension 1994;24(1):101-5.

93. Schulte KL, Liederwald K, Meyer-Sabellek W, van Gemmeren D, Lenz T, and Gotzen R. Relationships between ambulatory blood pressure, forearm vascular resistance, and left ventricular mass in hypertensive and normotensive subjects. Am J Hypertens 1993;6(9):786-93.

94. Amar J, Vernier I, Rossignol E, Bongard V, Arnaud C, Conte JJ, Salvador M, and Chamontin B. Nocturnal blood pressure and 24-hour pulse pressure are potent indicators of mortality in hemodialysis patients. Kidney Int 2000;57(6):2485-91.

95. Fagard RH, Staessen JA, Thijs L, Gasowsk J, Bulpitt CJ, Clement D, de Leeuw PW, Dobovisek J, Jaaskivi M, Leonetti G, O'Brien E, Palatini P, Parati G, Rodicio JL, Vanhanen H, and Webster J. Response to antihypertensive therapy in older patients with sustained and nonsustained systolic hypertension. Systolic Hypertension in Europe (Syst-Eur) Trial Investigators. Circulation 2000;102(10):1139-44.

96. Gosse P, Gasparoux P, Ansoborlo P, Lemetayer P, and Clementy J. Prognostic value of ambulatory measurement of the timing of Korotkoff sounds in elderly hypertensives: a pilot study. Am J Hypertens 1997;10(5 Pt 1):552-7.

97. Nakano S, Ogihara M, Tamura C, Kitazawa M, Nishizawa M, Kigoshi T, and Uchida K. Reversed circadian blood pressure rhythm independently predicts endstage renal failure in non-insulin-dependent diabetes mellitus subjects. J Diabetes Complications 1999;13(4):224-31.

98. Ohkubo T, Imai Y, Tsuji I, Nagai K, Watanabe N, Minami N, Kato J, Kikuchi N, Nishiyama A, Aihara A, Sekino M, Satoh H, and Hisamichi S. Relation between nocturnal decline in blood pressure and mortality. The Ohasama Study. Am J Hypertens 1997a;10(11):1201-7.

99. Ohkubo T, Imai Y, Tsuji I, Nagai K, Watanabe N, Minami N, Itoh O, Bando T, Sakuma M, Fukao A, Satoh H, Hisamichi S, and Abe K. Prediction of mortality by ambulatory blood pressure monitoring versus screening blood pressure measurements: a pilot study in Ohasama. J Hypertens 1997b;15(4):357-64.

100. Ohkubo T, Hozawa A, Nagai K, Kikuya M, Tsuji I, Ito S, Satoh H, Hisamichi S, and Imai Y. Prediction of stroke by ambulatory blood pressure monitoring versus screening blood pressure measurements in a general population: the Ohasama study. J Hypertens 2000;18(7):847-54.

101. Perloff D, Sokolow M, Cowan RM , and Juste RP. Prognostic value of ambulatory blood pressure measurements: further analyses. J Hypertens Suppl 1989;7(3):S3-10.

102. Redon J, Campos C, Narciso ML, Rodicio JL, Pascual JM, and Ruilope LM. Prognostic value of ambulatory blood pressure monitoring in refractory hypertension: a prospective study. Hypertension 1998;31(2):712-8.

103. Suzuki Y, Kuwajima I, Aono T, Kanemaru A, Nishinaga M, Shibata H, and Ozawa T. Prognostic value of nighttime blood pressure

in the elderly: a prospective study of 24-hour blood pressure. Hypertens Res 2000;23(4):323-30.

104. Verdecchia P, Porcellati C, Schillaci G, Borgioni C, Ciucci A, Battistelli M, Guerrieri M , Gatteschi C, Zampi I, Santucci A and others. Ambulatory blood pressure. An independent predictor of prognosis in essential hypertension. Hypertension 1994;24(6):793-801.

105. Verdecchia P, Schillaci G, Borgioni C, Ciucci A, Pede S, and Porcellati C. Ambulatory pulse pressure: a potent predictor of total cardiovascular risk in hypertension. Hypertension 1998;32(6):983-8.

106. Zweiker R, Eber B, Schumache, M, Toplak H, and Klein W. "Non-dipping" related to cardiovascular events in essential hypertensive patients. Acta Med Austriaca 1994;21(3):86-9.

107. Staessen JA, Byttebier G, Buntinx F, Celis H, O'Brien ET, and Fagard R. Antihypertensive treatment based on conventional or ambulatory blood pressure measurement. A randomized controlled trial. Ambulatory Blood Pressure Monitoring and Treatment of Hypertension Investigators. JAMA 1997;278(13):1065-72.

108. Schrader J, Luders S, Zuchner C, Herbold M, and Schrandt G. Practice vs ambulatory blood pressure measurement under treatment with ramipril (PLUR Study): a randomised, prospective long-term study to evaluate the benefits of ABPM in patients on antihypertensive treatment. J Hum Hypertens 2000;14(7):435-40.