Bibliography

Abitbol MM, Castillo I, Taylor UB, et al. Vaginal birth after cesarean section: The patient's point of view. Am Fam Physician 1993;47(1):129-34.

Abitbol MM, Taylor UB, Castillo I, et al. The cephalopelvic disproportion index. Combined fetal sonography and x-ray pelvimetry for early detection of cephalopelvic disproportion. J Reprod Med 1991;36(5):369-73.

Abma JC, Chandra A, Mosher WD, et al. Fertility, family planning, and women's health: new data from the 1995 National Survey of Family Growth. Vital & Health Statistics - Series 23, Data From the National Survey of Family Growth 1997(19):1-114.

Abraham R, Sadovsky E. Delay in the diagnosis of rupture of the uterus due to epidural anesthesia in labor. Gynecol Obstet Invest 1992;33(4):239-40.

Abramowitz L, Sobhani I, Ganansia R, et al. Are sphincter defects the cause of anal incontinence after vaginal delivery? Results of a prospective study. Dis Colon Rectum 2000;43(5):590-6; discussion 596-8.

Abu-Ghazzeh YM, Barqawi R. An appraisal of computed tomography pelvimetry in patients with previous caesarean section. East Medit Health J 2000;6(2-3):260-4.

ACOG Committee on Obstetric Practice. Vaginal delivery after a previous cesarean birth. ACOG Committee opinion. Number 143-October 1994. Int J Gynaecol Obster 1995;48(1):127-9.

ACOG Committee on Obstetric Practice. ACOG committee opinion. Rate of vaginal births after cesarean delivery. Number 179, November 1996. Int J Gynaecol Obster 1997;57(1):89-90.

ACOG Committee on Obstetric Practice. ACOG committee opinion. Monitoring during induction of labor with dinoprostone. Number 209, October 1998. Int J Gynaecol Obster 1999;64(2):200.

Adair CD, Sanchez-Ramos L, Gaudier FL, et al. Labor induction in patients with previous cesarean section. Am J Perinatol 1995;12(6):450-4.

Adair CD, Sanchez-Ramos L, Kaunitz AM, et al. A trial of labor complicated by uterine rupture following amnioinfusion. South Med J 1995;88(8):847-8.

Adair CD, Sanchez-Ramos L, Whitaker D, et al. Trial of labor in patients with a previous lower uterine vertical cesarean section. Am J Obstet Gynecol 1996;174(3):966-970.

Adami J, Glimelius B, Cnattingius S, et al. Maternal and perinatal factors associated with non-Hodgkin's lymphoma among children. Int J Cancer 1996;65(6):774-7.

Adams DM, Druzin ML, Cederqvist LL. Intrapartum uterine rupture. Obstet Gynecol 1989;73(3 Pt 2):471-3.

Adams EK, Mauldin PD, Mauldin JG, et al. Determining cost savings from attempted cephalic version in an inner city delivering population. Health Care Management Science 2000;3(3):185-92.

Adolph HP, Adolph BJ. Surgery on the edge of the desert. Bull Am Coll Surg 1997;82(6):8-19.

Afriat CI. Vaginal birth after cesarean section: A review of the literature. J Perinat Neonat Nurs 1990;3(3):1-13.

Ahdoot D, Van Nostrand KM, Nguyen NJ, et al. The effect of route of delivery on regression of abnormal cervical cytologic findings in the postpartum period. Am J Obstet Gynecol 1998;178(6):1116-20.

Akar ME, Yilmaz Z, Gokmen O. Fetal survival despite unrecognized uterine rupture resulting from previous unknown corporeal scar. Arch Gynecol Obstet 2001;265(2):89-90.

Akyol D, Mungan T, Unsal A, et al. Prelabour rupture of the membranes at term - No advantage of delaying induction for 24 hours. Aust N Z J Obstet Gynaecol 1999;39(3):291-295.

Al Nuaim LA, Mustafa MS. Vaginal breech delivery or a third cesarean section for intrauterine fetal demise. Int J Gynaecol Obster 1996;52(2):183-4.

Al Sakka M, Dauleh W, Al Hassani S. Case series of uterine rupture and subsequent pregnancy outcome. Int J Fertil Womens Med 1999;44(6):297-300.

Al Sakka M, Hamsho A, Khan L. Rupture of the pregnant uterus--a 21-year review. Int J Gynaecol Obster 1998;63(2):105-8.

Al-Sibai MH, Rahman J, Rahman MS, et al. Emergency hysterectomy in obstetrics--a review of 117 cases. Aust N Z J Obstet Gynaecol 1987;27(3):180-4.

Aladjem S. Twice a cesarean, always a cesarean? Obstet Gynecol 1989;74(1):137-8.

Albers LL, Savitz DA. Hospital setting for birth and use of medical procedures in low-risk women. J Nurse Midwifery 1991;36(6):327-33.

Albrechtsen S, Rasmussen S, Reigstad H, et al. Evaluation of a protocol for selecting fetuses in breech presentation for vaginal delivery or cesarean section. Am J Obstet Gynecol 1997;177(3):586-92.

Ales KL, Druzin ML, Santini DL. Impact of advanced maternal age on the outcome of pregnancy. Surg Gynecol Obstet 1990;171(3):209-16.

Alexander JM, Ramus R, Cox SM, et al. Outcome of twin gestations with a single anomalous fetus. Am J Obstet Gynecol 1997;177(4):849-52.

Alexander JM, Sharma SK, McIntire DD, et al. Intensity of labor pain and cesarean delivery. Anesth Analg 2001;92(6):1524-8.

Alfirevic Z. Oral misoprostol for induction of labour. Cochrane Database of Systematic Reviews 2000(4):CD001338.

Alfirevic Z, Howarth G, Gaussmann A. Oral misoprostol for induction of labour with a viable fetus. Cochrane Database of Systematic Reviews 2000(2):CD001338.

Allenby K, Rand RJ. Pregnancies in a woman with a vesico-uterine fistula following lower segment caesarean section. Br J Obstet Gynaecol 1996;103(1):87-9.

Amer-Wahlin I, Hellsten C, Noren H, et al. Cardiotocography only versus cardiotocography plus ST analysis of fetal electrocardiogram for intrapartum fetal monitoring: A Swedish randomised controlled trial. Lancet 2001;358(9281):534-8.

American Academy of Family Physicians. FACTS about Family Practice. Table 67. (www.aafp.org). 2001.

American College of Obstetricians and Gynecologists. News release: Court-Ordered Treatment Rarely Justified for Obstetrical Care. 1987(August).

American College of Obstetricians and Gynecologists. Vaginal birth after previous cesarean delivery: Clinical Management guidelines for obstetrician-gynecologists. Washington: The College 1999;ACOG Practice Bulletin No. 5.

Amin P, Wiener JJ. Vaginal delivery after abdominal wall replacement by Marlex mesh. J Obstet Gynaecol 1999;19(5):543.

Amini SB, Dierker LJ, Catalano PM, et al. Trends in an obstetric patient population: An eighteen-year study. Am J Obstet Gynecol 1994;171(4):1014-21.

Amir W, Peter J, Etan Z. Trial of labor without oxytocin in patients with a previous cesarean section. Am J Perinatol 1987;4(2):140-3.

Amon E, Sibai BM, Anderson GD. How perinatologists manage the problem of the presenting breech. Am J Perinatol 1988;5(3):247-50.

Ananth CV, Smulian JC, Vintzileos AM. The association of placenta previa with history of cesarean delivery and abortion: A metaanalysis. Am J Obstet Gynecol 1997;177(5):1071-8.

Anderson GM, Lomas J. Explaining variations in cesarean section rates: Patients, facilities or policies? Can Med Assoc J 1985;132(3):253-6.

Anderson R, Greener D. A descriptive analysis of home births attended by CNMs in two nurse-

midwifery services. J Nurse Midwifery 1991;36(2):95-103.

Anderson RE, Anderson DA. The cost-effectiveness of home birth. J Nurse Midwifery 1999;44(1):30-5.

Andreasson B, Bock JE, Larsen J. Induction of labor. A double-blind randomized controlled study of prostaglandin E2 vaginal suppositories compared with intranasal oxytocin and with sequential treatment. Acta Obstet Gynecol Scand 1985;64(2):157-61.

Andres RL, Piacquadio KM, Resnik R. A reappraisal of the need for autologous blood donation in the obstetric patient. Am J Obstet Gynecol 1990;163(5 Pt 1):1551-3.

Annas GJ. Protecting the liberty of pregnant patients. N Engl J Med 1987;316(19):1213-4.

Annibale DJ, Hulsey TC, Wagner CL, et al. Comparative neonatal morbidity of abdominal and vaginal deliveries after uncomplicated pregnancies. Arch Pediatr Adolesc Med 1995;149(8):862-7.

Anonymous. Caesarean childbirth. Summary of an NIH consensus statement. Br Med J (Clin Res Ed) 1981;282(6276):1600-4.

Anonymous. Cesarean childbirth: an NIH consensus development conference. Clin Pediatr 1981;20(9):555-60.

Anonymous. Cesarean Childbirth: Report of a Consensus Development Conference Sponsored by the NICHHD in Conjunction with the National Center for Health Care Technology and Assisted by the Office for Medical Applications of Research. NIH Publication 1981;82-2067.

Anonymous. The National Institutes of Health Consensus Development statement on cesarean childbirth. A summary. J Reprod Med 1981;26(3):103-12.

Anonymous. NIH Consensus Development Conference summary--Cesarean childbirth. J Tenn Med Assoc 1981;74(10):734-40.

Anonymous. NIH Consensus Development Task force statement on cesarean childbirth. Am J Obstet Gynecol 1981;139(8):902-9.

Anonymous. Panel discussion: cesarean section. March of Dimes Symposium - perinatal care in the 1980's. November 17, 1980. J Perinat Med 1981;9(4):195-9.

Anonymous. Breech: Vaginal delivery or caesarean section? Br Med J (Clin Res Ed) 1982;285(6350):1275-6.

Anonymous. Kielland or Caesar? Br Med J (Clin Res Ed) 1983;287(6392):609-11.

Anonymous. Forum on malpractice issues in childbirth. Public Health Rep 1985;100(6):629-33.

Anonymous. Vaginal delivery after cesarean section. Am Fam Physician 1985;32(3):90.

Anonymous. Indications for cesarean section: Final statement of the Panel of the National Consensus Conference on Aspects of Cesarean Birth. Can Med Assoc J 1986;134(12):1348-52.

Anonymous. Cesarean births and trial of labor rates. JAMA 1987;257(20):2757-9.

Anonymous. Court-ordered obstetrical interventions. N Engl J Med 1987;317(19):1223-5.

Anonymous. Delivery after caesarean section. Br Med J (Clin Res Ed) 1987;295(6594):388.

Anonymous. Obstetrics '87. A report of the Canadian Medical Association on obstetrical care in Canada. Can Med Assoc J 1987;Suppl:1-26.

Anonymous. Births by cesarean: cost changes, 1982-83 to 1986. Stat Bull Metrop Insur Co 1988;1988(69):18-25.

Anonymous. Costs of normal births: Regional variations, 1986. Stat Bull Metrop Insur Co 1988;69(4):24-32.

Anonymous. The costs of maternity care and childbirth in the United States. Health Insurance Association of America. Research Bulletin 1989;1 (R1589).

Anonymous. Practice parameters: Cesarean birth. Minnesota Clinical Comparison and Assessment Project. QA Rev 1991;3(5):6.

Anonymous. Charges for a cesarean section: United States, 1990. Stat Bull Metrop Insur Co 1992;73(1):12-8.

Anonymous. Induction and augmentation of labor: ACOG Technical bulletin number 157. International Journal of Gynecology & Obstetrics 1992;39(2):139-142.

Anonymous. Improving the timeliness of emergency C-sections at Southwestern Vermont Medical Center leads to improved patient care and increased physician satisfaction. Qual Lett Healthc Lead 1993;5(1):6-8.

Anonymous. Mothers who use nurse midwives have shorter hospital stays and fewer cesarean sections and episiotomies. J Nurs Adm 1993;23(7-8):11, 40.

Anonymous. Rates of cesarean delivery-United States, 1991. MMWR Morb Mortal Wkly Rep 1993;42:285-289.

Anonymous. Health care guideline: Vaginal birth after cesarean. In: Institute for Clinical Systems Integration 1993/1994 health care guidelines 1994;Bloomington(MN):ICSI.

Anonymous. AAFP Task Force on Clinical Policies for Patient Care. Trial of labor vs. elective repeat cesarean section for the woman with a previous cesarean section. American Academy of Family Physicians 1995.

Anonymous. Australia's mothers and babies. Aust Health Rev 1995;18(3):137-8.

Anonymous. Obstetrical stays--IMS, AMA policy. Iowa Med 1995;85(9):358.

Anonymous. Trial of labor vs. elective repeat cesarean section. AAFP Task Force on Clinical Policies for Patient Care. Am Fam Physician 1995;52(6):1763-5.

Anonymous. Vaginal delivery after previous cesarean birth. ACOG Prac Pat 1995;1:1-8.

Anonymous. Average postpartum length of stay for uncomplicated deliveries--New Jersey, 1995. MMWR Morb Mortal Wkly Rep 1996;45(32):700-4.

Anonymous. Health care guideline: Vaginal birth after cesarean. In: Institute for Clinical Systems Integration 1996;1996:577-90.

Anonymous. Multimillion dollar verdict in missed uterine rupture case. Obstet Gynecol Malpract Prev 1996;3(6):41-43.

Anonymous. Vaginal birth after previous caesarean birth. In: The Society of Obstetricians and Gynaecologists of Canada: policy statement, committee opinion, clinical practice guidelines. Ottawa, ON : SOGC 1996;4.

Anonymous. Vaginal delivery after previous cesarean birth. Number 1--August 1995. Committee on Practice Patterns. American College of Obstetricians and Gynecologists. Int J Gynaecol Obster 1996;52(1):90-8.

Anonymous. VBAC (Vaginal Birth After Cesareans): Are cost concerns outweighing possible safety risks? Hosp Case Manag 1996;4(11):161-4.

Anonymous. 1998 ACOG Socioeconomic Survey of Fellows (unpublished.). ACOG NET 1997;www.acog.com.

Anonymous. ACOG practice patterns. External cephalic version. Number 4, July 1997. American College of Obstetricians and Gynecologists. Int J Gynaecol Obster 1997;59(1):73-80.

Anonymous. Electronic fetal heart rate monitoring: Research guidelines for interpretation. National Institute of Child Health and Human Development Research Planning Workshop. American Journal of Obstetrics & Gynecology. 1997;177(6):1385-90.

Anonymous. Approach VBAC deliveries cautiously, experts say. Hosp Peer Rev 1998;23(10):189.

Anonymous. C-sections give way to VBACs and better quality. Hosp Peer Rev 1998;23(10):184, 189.

Anonymous. Case records of the Massachusetts General Hospital. Weekly clinicopathological exercises. Case 9-1998. Cardiovascular collapse after vaginal delivery in a patient with a history of cesarean section. N Engl J Med 1998;338(12):821-6.

Anonymous. Charges and LOS for deliveries

vary widely by geographic region, analysis concludes. Data Strat Benchmarks 1998;2(10):154-7.

Anonymous. Outreach program slashes Florida health network's rate of cesarean sections. Health Care Cost Reeng Rep 1998;3(9):137-8.

Anonymous. ACOG practice bulletin. Vaginal birth after previous cesarean delivery. Number 2, October 1998. Clinical management guidelines for obstetrician-gynecologists. American College of Obstetricians and Gynecologists. Int J Gynaecol Obster 1999;64(2):201-8.

Anonymous. Misoprostol for cervical ripening: Is it a safe, cost-effective alternative to dinoprostone? Formulary 1999;34(1):77-78.

Anonymous. When the patient changed her mind. Schreiber vs. Physicians Ins. Co. Hosp Law Newsl 1999;16(5):1-3.

Anonymous. From the Centers for Disease Control and Prevention. Use of hospital discharge data to monitor uterine rupture--Massachusetts, 1990-1997. JAMA 2000;283(16):2098-100.

Anonymous. ACOG committee opinion: number 265, December 2001. Mode of term single breech delivery. Obstet Gynecol 2001;98(6):1189-90.

Anonymous. Stats & facts. Vaginal births after cesarean sections: A need to revisit. Manag Care Interface 2001;14(11):34-5.

Appleby J. Data briefing. Caesarean births. Health Serv J 2001;111(5756):31.

Appleton B, Targett C, Rasmussen M, et al. Knowledge and attitudes about vaginal birth after Caesarean section in Australian hospitals. VBAC Study Group. Vaginal Birth After Caesarean. Aust N Z J Obstet Gynaecol 2000;40(2):195-9.

Appleton B, Targett C, Rasmussen M, et al. Vaginal birth after caesarean section: an Australian multicentre study. VBAC Study Group. Aust N Z J Obstet Gynaecol 2000;40(1):87-91.

Apuzzio JJ, Pelosi M, Frisoli G, et al. Bacterial colonization of amniotic fluid in patients with intact membranes during labor. South Med J 1984;77(9):1133-5.

Arkutu AA. Spontaneous uterine rupture with placenta percreta: a case report. Int J Gynaecol Obster 1980;17(5):485-7.

Arnold M. Midwifery care on trial? Midwife Health Visit Community Nurse 1986;22(6):207-8.

Artinian BM. Collaborative planning of patient care in the prenatal, labor and delivery, and neonatal settings. JOGN Nurs 1984;13(2):105-10.

Arulkumaran S, Chua S, Ratnam SS. Symptoms and signs with scar rupture--value of uterine activity measurements. Aust N Z J Obstet Gynaecol 1992;32(3):208-12.

Arulkumaran S, Gibb DM, Ingemarsson I, et al. Uterine activity during spontaneous labour after previous lower-segment caesarean section. Br J Obstet Gynaecol 1989;96(8):933-8.

Arulkumaran S, Ingemarsson I, Ratnam SS. Oxytocin augmentation in dysfunctional labour after previous caesarean section. Br J Obstet Gynaecol 1989;96(8):939-941.

Asaad K, Alaily B. Oxytocin use and delivery outcome in women with one previous caesarean section and pre-labour rupture of the membranes at term. J Obstet Gynaecol 1994;14(6):420-422.

Asakura H, Myers SA. More than one previous cesarean delivery: a 5-year experience with 435 patients. Obstet Gynecol 1995;85(6):924-9.

Asakura H, Nakai A, Ishikawa G, et al. Prediction of uterine dehiscence by measuring lower uterine segment thickness prior to the onset of labor: Evaluation by transvaginal ultrasonography. Nippon Ika Daigaku Zasshi - Journal of the Nippon Medical School 2000;67(5):352-6.

Atalla RK, Thompson JR, Oppenheimer CA, et al. Reactive thrombocytosis after caesarean section and vaginal delivery: Implications for maternal thromboembolism and its prevention. BJOG 2000;107(3):411-4.

Atiba EO, Adeghe AJ, Murphy PJ, et al. Patients' expectation and caesarean section rate. Lancet

1993;341(8839):246.

Austin SE. Childbirth classes for couples desiring VBAC (Vaginal Birth After Cesarean). MCN Am J Matern Child Nurs 1986;11(4):250-5.

Awad IT, Shorten GD. Amniotic fluid embolism and isolated coagulopathy: Atypical presentation of amniotic fluid embolism. Eur J Anaesthesiol 2001;18(6):410-3.

Ayhan A, Yuce K, Kisnisci HA. Analysis of 20,291 deliveries in a Turkish institution. Int J Gynaecol Obster 1989;29(2):131-4.

Ayres AW, Johnson TR, Hayashi R. Characteristics of fetal heart rate tracings prior to uterine rupture. Int J Gynaecol Obster 2001;74(3):235-40.

Azem F, Jaffa A, Lessing JB, et al. Uterine rupture with the use of a low-dose vaginal PGE2 tablet. Acta Obstet Gynecol Scand 1993;72(4):316-7.

Baird AG, Jewell D, Walker JJ. Management of labour in an isolated rural maternity hospital. BMJ 1996;312(7025):223-6.

Bais JM, van der Borden DM, Pel M, et al. Vaginal birth after caesarean section in a population with a low overall caesarean section rate. Eur J Obstet Gynecol Reprod Biol 2001;96(2):158-62.

Baker K. Vaginal delivery after lower uterine cesarean section. Surg Gynecol Obstet 1955;100:690.

Bakshi S, Meyer BA. Indications for and outcomes of emergency peripartum hysterectomy. A five-year review. J Reprod Med 2000;45(9):733-7.

Balaban D, McCall N, Jones-Bauer E. Quality of Medicaid Managed Care: An evaluation of the Arizona Health Care Cost Containment System (AHCCCS). San Francisco: Laguna Research Associates; 1994.

Baldwin LM, Hart LG, Lloyd M, et al. Defensive medicine and obstetrics. JAMA 1995;274(20):1606-10.

Barclay AM, Knapp DP, Kallail KJ. The provision of labor and delivery services by graduates of four Kansas family practice residencies. Kans Med 1996;97(1):19-23.

Barclay L, Andre CA, Glover PA. Women's business: the Challenge of childbirth. Midwifery 1989;5(3):122-33.

Barnhart RJ. Cesarean section: Changing indications, 1968--1977. J Kans Med Soc 1980;81(2):57-62, 78.

Barnsley JM, Vayda E, Lomas J, et al. Cesarean section in Ontario: Practice patterns and responses to hypothetical cases. Can J Surg 1990;33(2):128-32.

Barratt K, von Briesen JD. The continued vitality of a patient's informed consent, or, when the patient says 'no'. Wis Med J 1999;98(2):60-1.

Barros FC, Vaughan JP, Victora CG, et al. Epidemic of caesarean sections in Brazil. Lancet 1991;338(8760):167-9.

Barton A. Caesarean section and litigation. Lancet 2001;357(9255):556.

Baskett TF, McMillen RM. Cesarean section: Trends and morbidity. Can Med Assoc J 1981;125(7):723-6.

Bassett KL, Iyer N, Kazanjian A. Defensive medicine during hospital obstetrical care: a Byproduct of the technological age. Soc Sci Med 2000;51(4):523-37.

Baumann P, Blackwell SC, Schild C, et al. Mathematic modeling to predict abruptio placentae. Am J Obstet Gynecol 2000;183(4):815-22.

Beall M, Eglinton GS, Clark SL, et al. Vaginal delivery after cesarean section in women with unknown types of uterine scar. J Reprod Med 1984;29(1):31-5.

Beckett VA, Regan L. Vaginal birth after cesarean: The European experience. Clin Obstet Gynecol 2001;44(3):594-603.

Beckley S, Gee H, Newton JR. Scar rupture in labour after previous lower uterine segment caesarean section: The role of uterine activity measurement. Br J Obstet Gynaecol 1991;98(3):265-9.

Bedoya C, Bartha JL, Rodriguez I, et al. A trial of labor after cesarean section in patients with or without a prior vaginal delivery. Int J Gynaecol Obster 1992;39(4):285-9.

Beebe LA, Rayburn WF, Beaty CM, et al. Indications for labor induction. Differences between university and community hospitals. J Reprod Med 2000;45(6):469-75.

Benedetti TJ. Birth injury and method of delivery. N Engl J Med 1999;341(23):1758-9.

Benedetti TJ, Platt L, Druzin M. Vaginal delivery after previous cesarean section for a nonrecurrent cause. Am J Obstet Gynecol 1982;142(3):358-9.

Bennett BB. Uterine rupture during induction of labor at term with intravaginal misoprostol. Obstet Gynecol 1997;89(5 Pt 2):832-3.

Bennett KA, Butt K, Crane JMG, et al. A masked randomized comparison of oral and vaginal administration of misoprostol for labor induction. Obstet Gynecol 1998;92(4 I):481-486.

Bennett MJ, Laurence DT. Induction of labour with prostaglandin E2 vaginal pessaries in high risk pregnancies. J Obstet Gynaecol 1981;2(2):71-73.

Bennetts AB, Lubic RW. The free-standing birth centre. Lancet 1982;1(8268):378-80.

Benson MD. Therapeutic equivalents in clinical practice. Int J Fertil Womens Med 2001;46(2):89-94.

Berard J, Dufour P, Vinatier D, et al. Fetal macrosomia: risk factors and outcome. A study of the outcome concerning 100 cases >4500 g. Eur J Obstet Gynecol Reprod Biol 1998;77(1):51-9.

Berchuck A, Sokol RJ. Previous cesarean section, placenta increta, and uterine rupture in second-trimester abortion. Am J Obstet Gynecol 1983;145(6):766-7.

Berg A, Yuval D, Ivancovsky M, et al. Patient perception of involvement in medical care during labor and delivery. Isr Med Assoc J 2001;3(5):352-6.

Berghella V, Rogers RA, Lescale K. Stripping of membranes as a safe method to reduce prolonged pregnancies. Obstet Gynecol 1996;87(6):927-931.

Bergman A, Henderson M, Cline JL. Technology and outpatient review: a Preliminary evaluation. Qrb. Quality Review Bulletin 1990;16(6):234-9.

Bergsjo P, Denman DW, 3rd, Hoffman HJ, et al. Duration of human singleton pregnancy. A population-based study. Acta Obstet Gynecol Scand 1990;69(3):197-207.

Bergsjo P, Schmidt E, Pusch D. Differences in the reported frequencies of some obstetrical interventions in Europe. Br J Obstet Gynaecol 1983;90(7):628-32.

Berkman H. A discussion of medical malpractice and cesarean section. Spec Law Dig Health Care Law 1993(170):9-30.

Berkowitz GS, Fiarman GS, Mojica MA, et al. Effect of physician characteristics on the cesarean birth rate. Am J Obstet Gynecol 1989;161(1):146-9.

Berkowitz GS, Skovron ML, Lapinski RH, et al. Delayed childbearing and the outcome of pregnancy. N Engl J Med 1990;322(10):659-64.

Berkus MD, Ramamurthy RS, O'Connor PS, et al. Cohort study of Silastic obstetric vacuum cup deliveries: II. Unsuccessful vacuum extraction. Obstet Gynecol 1986;68(5):662-6.

Bernstein P, Leyland N, Gurland P, et al. Cervical ripening and labor induction with prostaglandin E2 gel: a Placebo-controlled study. Am J Obstet Gynecol 1987;156(2):336-40.

Bertollini R, DiLallo D, Spadea T, et al. Cesarean section rates in Italy by hospital payment mode: An analysis based on birth certificates. Am J Public Health 1992;82(2):257-61.

Bex P, Gunasekera PC, Phipps JH. Difficulties with controlled release prostaglandin E2 pessaries (II). Lancet 1990;336(8707):119.

Bhal PS, Sharma A, Asaad K. Vaginal delivery after caesarean section: factors influencing success rates. Aust N Z J Obstet Gynaecol 1996;36(4):497-8.

Bianco A, Stone J, Lynch L, et al. Pregnancy outcome at age 40 and older. Obstet Gynecol 1996;87(6):917-22.

Bick DE, MacArthur C. Attendance, content and relevance of the six week postnatal examination. Midwifery 1995;11(2):69-73.

Bickell NA, Zdeb MS, Applegate MS, et al. Effect of external peer review on cesarean delivery rates: a statewide program. Obstet Gynecol 1996;87(5 Pt 1):664-7.

Bique C, Bugalho A, Bergstrom S. Labor induction by vaginal misoprostol in grand multiparous women. Acta Obstet Gynecol Scand 1999;78(3):198-201.

Biran G, Mazor M, Shoham I, et al. Premature delivery of small versus appropriate-for-gestational-age neonates. A comparative study of maternal characteristics. J Reprod Med 1994;39(1):39-44.

Blackwell SC, Hassan SS, Wolfe HM, et al. Vaginal birth after cesarean in the diabetic gravida. J Reprod Med 2000;45(12):987-90.

Blakemore KJ, Qin NG, Petrie RH, et al. A prospective comparison of hourly and quarter-hourly oxytocin dose increase intervals for the induction of labor at term. Obstet Gynecol 1990;75(5):757-61.

Blanchette H, Blanchette M, McCabe J, et al. Is vaginal birth after cesarean safe? Experience at a community hospital. Am J Obstet Gynecol 2001;184(7):1478-84; discussion 1484-7.

Blanchette HA, Nayak S, Erasmus S, et al. Comparison of the safety and efficacy of intravaginal misoprostol (prostaglandin e1) with those of dinoprostone (prostaglandin E2) for cervical ripening and induction of labor in a community hospital. Am J Obstet Gynecol 1999;180(6 Pt I):1551-1559.

Blanco JD, Collins M, Willis D, et al. Prostaglandin E2 gel induction of patients with a prior low transverse cesarean section. Am J Perinatol 1992;9(2):80-3.

Blegen MA, Reiter RC, Goode CJ, et al. Outcomes of hospital-based managed care: a Multivariate analysis of cost and quality. Obstet Gynecol 1995;86(5):809-14.

Blickstein I, Lancet M, Kessler I. Re-evaluation of the obstetrical risk for the older primipara. Int J Gynaecol Obster 1987;25(2):107-12.

Blumenthal PD, Gaffikin L, Atad J, et al. Nonpharmaceutical ripening of the unfavorable cervix and induction of labor by a novel double balloon device [8]. Obstet Gynecol 1991;77(6):967-968.

Boivin J. Baby and me--out 1,2,3. Nursing Spectrum (D.C./Baltimore Metro Edition) 1995;5(17):3.

Bolaji, II, Meehan FP. Caesarean section survey in Galway--1973 through 1987. Eur J Obstet Gynecol Reprod Biol 1993;48(1):1-8.

Bonham DG. Caesarean birth. N Z Med J 1983;96(728):205-6.

Bonnar J, Norris LA, Greene R. Low molecular weight heparin for thromboprophylaxis during caesarean section. Thromb Res 1999;96(4):317-22.

Bors-Koefoed R, Zylstra S, Resseguie LJ, et al. Statistical models of outcome in malpractice lawsuits involving death or neurologically impaired infants. J Matern Fetal Med 1998;7(3):124-31.

Bortolus R. Determinants of response to intracervical prostaglandin E2 for cervical ripening. Eur J Obstet Gynecol Reprod Biol 1999;87(2):137-141.

Bossert R, Rayburn WF, Stanley JR, et al. Early postpartum discharge at a university hospital. Outcome analysis. J Reprod Med 2001;46(1):39-43.

Bottoms SF, Hirsch VJ, Sokol RJ. Medical management of arrest disorders of labor: a Current overview. Am J Obstet Gynecol 1987;156(4):935-9.

Boucher M, Tahilramaney MP, Eglinton GS, et al. Maternal morbidity as related to trial of labor after previous cesarean delivery. A quantitative analysis. J Reprod Med 1984;29(1):12-6.

Boulot P, Hoffet M, Bachelard B, et al. Late vaginal induced abortion after a previous cesarean birth: Potential for uterine rupture. Gynecol Obstet Invest 1993;36(2):87-90.

Boulvain M, Fraser WD, Brisson-Carroll G, et al. Trial of labour after caesarean section in sub-Saharan Africa: a Meta-analysis. Br J Obstet Gynaecol 1997;104(12):1385-90.

Boulvain M, Irion O. Stripping/sweeping the membranes for inducing labour or preventing post-term pregnancy. Cochrane Database of Systematic Reviews 2000(2):CD000451.

Boulvain M, Irion O, Marcoux S, et al. Sweeping of the membranes to prevent post-term pregnancy and to induce labour: a Systematic review. Br J Obstet Gynaecol 1999;106(5):481-5.

Boulvain M, Stan C, Irion O. Elective delivery in diabetic pregnant women. Cochrane Database of Systematic Reviews 2000(2):CD001997.

Bowers SK, MacDonald HM, Shapiro ED. Prevention of iatrogenic neonatal respiratory distress syndrome: Elective repeat cesarean section and spontaneous labor. Am J Obstet Gynecol 1982;143(2):186-9.

Boyce PM, Todd AL. Increased risk of postnatal depression after emergency caesarean section. Med J Aust 1992;157(3):172-4.

Boyd ME, Usher RH, McLean FH, et al. Obstetric consequences of postmaturity. Am J Obstet Gynecol 1988;158(2):334-8.

Boyers SP, Gilbert WM. Elective repeat caesarean section versus trial of labour: the neonatologist's view. Lancet 1998; 351(9097):155.

Brady K, Read JA. Vaginal delivery of twins after previous cesarean section. N Engl J Med 1988;319(2):118-9.

Brahams D. Caesarean sections by court order. Lancet 1996;348(9030):770.

Bramadat IJ, Driedger M. Satisfaction with childbirth: Theories and methods of measurement. Birth 1993;20(1):22-9.

Brandsma C, Calhoun BC, Vannatta JE. Uncomplicated pregnancy: Clinical pathway genesis based on the nursing process. Mil Med 2000;165(11):839-43.

Braveman P, Egerter S, Edmonston F, et al. Racial/ethnic differences in the likelihood of cesarean delivery, California. Am J Public Health 1995;85(5):625-30.

Braveman PA. Short hospital stays for mothers and newborns. J Fam Pract 1996;42(5):523-5.

Bredfeldt RC, Thomas JM, Massie M. Pregnancy care in family practice: Medical student perspectives on specialty and residency selection. Fam Med 1994;26(3):145-8.

Bremme K, Bygdeman M. A comparative study of uterine activity and fetal heart rate pattern in labor induced with oral prostaglandin E2 or oxytocin. Acta Obstet Gynecol Scand Suppl 1980;92:23-9.

Bremme K, Bygdeman M. Induction of labor by oxytocin or prostaglandin E2. Acta Obstet Gynecol Scand Suppl 1980;92:11-21.

Brengman SL, Burns MK. Vaginal delivery after C-section. Am J Nurs 1983;83(11):1544-7.

Bretelle F, Cravello L, Shojai R, et al. Vaginal birth following two previous cesarean sections. Eur J Obstet Gynecol Reprod Biol 2001; 94(1):23-6.

Bricker L, Luckas M. Amniotomy alone for induction of labour. Cochrane Database of Systematic Reviews 2000(4):CD002862.

Brink S. C-sections rise but may not be the kindest cut. US News & World Report 2000;129(9):63.

Brisson-Carroll G, Fraser W, Breart G, et al. The effect of routine early amniotomy on spontaneous labor: a Meta-analysis. Obstet Gynecol 1996;87(5 Pt 2):891-6.

Britton JR. Postpartum early hospital discharge and follow-up practices in Canada and the United States. Birth 1998;25(3):161-8.

Brody CZ, Kosasa TS, Nakayama RT, et al. Vaginal birth after cesarean section in Hawaii. Experience at Kapiolani Medical Center for Women and Children. Hawaii Med J 1993;52(2):38-42.

Broe S, Khoo SK. How safe is caesarean section in current practice? A survey of mortality and serious morbidity. Aust N Z J Obstet Gynaecol

1989;29(2):93-8.

Bromham DR, Anderson RS. Uterine scar rupture in labour induced with vaginal prostaglandin E2. Lancet 1980;2(8192):485-486.

Brooten D, Roncoli M, Finkler S, et al. A randomized trial of early hospital discharge and home follow-up of women having cesarean birth. Obstet Gynecol 1994;84(5):832-8.

Brow CS, Garcia JI, Pellegrini JE. Comparison of the length of stage II labor and incidence of forceps and cesarean delivery after epidural infusion of 0.125% bupivacaine with 2 mu/mL fentanyl as compared with 0.0625% bupivacaine with 2 mu/mL fentanyl. CRNA 2000;11(2):51-6.

Brown CE, Stettler RW, Twickler D, et al. Puerperal septic pelvic thrombophlebitis: incidence and response to heparin therapy. Am J Obstet Gynecol 1999;181(1):143-8.

Brown S. Mothers who use nurse midwives have fewer C-sections, episiotomies and shorter hospital stays. Penn Nurse 1993;48(4):9.

Brown S. Women's views of care in labour and birth: Is anyone listening? Proceedings of the 2nd Annual Conference of the Perinatal Society of Autralia and New Zealand 1998.

Brown S, Lumley J. Satisfaction with care in labor and birth: a Survey of 790 Australian women. Birth 1994;21(1):4-13.

Brown S, Lumley J. Changing childbirth: Lessons from an Australian survey of 1336 women. Br J Obstet Gynaecol 1998;105(2):143-55.

Brown S, Lumley J. Maternal health after childbirth: Results of an Australian population based survey. Br J Obstet Gynaecol 1998;105(2):156-61.

Bryan H, Hawrylyshyn P, Hogg-Johnson S, et al. Perinatal factors associated with the respiratory distress syndrome. Am J Obstet Gynecol 1990;162(2):476-81.

Bryant DR, Leonardi MR, Landwehr JB, et al. Limited usefulness of fetal weight in predicting neonatal brachial plexus injury. Am J Obstet Gynecol 1998;179(3 Pt 1):686-9.

Buechner JS. Rate of vaginal births among women with previous cesarean deliveries. Med Health Rhode Island 1997;80(2):63-4.

Buist R, Brown J, McNamara T. For whom is the caesarean section rate high? N Z Med J 1999;112(1101):469-71.

Bujold E, Gauthier RJ. Neonatal morbidity associated with uterine rupture: What are the risk factors? Am J Obstet Gynecol 2002;186(2):311-4.

Burke MS, Porreco RP. Obstetric factors causing early periventricular-intraventricular hemorrhage. Obstet Gynecol 1985;65(6):858-9.

Burns LR, Chilingerian JA, Wholey DR. The effect of physician practice organization on efficient utilization of hospital resources. Health Serv Res 1994;29(5):583-603.

Burns LR, Geller SE, Wholey DR. The effect of physician factors on the cesarean section decision. Med Care 1995;33(4):365-82.

Burns LR, Wholey DR, Abeln MO. Hospital utilization and mortality levels for patients in the Arizona Health Care Cost Containment System. Inquiry 1993;30(2):142-56.

Burrows J. The parturient woman: Can there be room for more than 'one person with full and equal rights inside a single human skin'? J Adv Nurs 2001;33(5):689-95.

Burt RD, Vaughan TL, Daling JR. Evaluating the risks of cesarean section: Low Apgar score in repeat C-section and vaginal deliveries. AmJ Public Health 1988;78(10):1312-4.

Bussinger EK. Vaginal birth after cesarean section in a rural private setting. Nebr Med J 1993;78(11):358-60.

Butler J, Abrams B, Parker J, et al. Supportive nurse-midwife care is associated with a reduced incidence of cesarean section. Am J Obstet Gynecol 1993;168(5):1407-13.

Byrne JP, Crowther CA, Moss JR. A randomised controlled trial comparing birthing centre care with delivery suite care in Adelaide, Australia. Aust N Z J Obstet Gynaecol 2000;40(3):268-74.

Cadet JJ. Occult uterine rupture: Role of

ultrasonography. J Natl Med Assoc 1998;90(6):374-6.

Cain M. Choice of plan in Michigan Medicaid and cesarean section delivery. AHSR FSHR Annu Meet Abstr Book;13:2a-4a.

Callahan C, Chescheir N, Steiner BD. Safety and efficacy of attempted vaginal birth after cesarean beyond the estimated date of delivery. J Reprod Med 1999;44(7):606-10.

Carey TS, Weis K, Homer C. Prepaid versus traditional Medicaid plans: Lack of effect on pregnancy outcomes and prenatal care. Health Serv Res 1991;26(2):165-81.

Carlan SJ, Bouldin S, O'Brien WF. Extemporaneous preparation of misoprostol gel for cervical ripening: a Randomized trial. Obstet Gynecol 1997;90(6):911-5.

Carley ME, Turner RJ, Scott DE, et al. Obstetric history in women with surgically corrected adult urinary incontinence or pelvic organ prolapse. J Am Assoc Gynecol Laparosc 1999;6(1):85-9.

Carlomagno G, Candussi G, Conforti BR, et al. Vaginal birth after caesarean section: Further contribution to counteract caesarean section epidemic. Ann Ig 1992;4(4):199-202.

Carlsson C, Nybell-Lindahl G, Ingemarsson I. Extradural block in patients who have previously undergone caesarean section. Br J Anaesth 1980;52(8):827-30.

Caron A, Neuhauser D. The effect of public accountability on hospital performance: Trends in rates for cesarean sections and vaginal births after cesarean section in Cleveland, Ohio. Qual Manag Health Care 1999;7(2):1-10.

Carpenter MW, Soule D, Yates WT, et al. Practice environment is associated with obstetric decision making regarding abnormal labor. Obstet Gynecol 1987;70(4):657-62.

Carr CJ, Patton R. A comparison of varying doses of vaginal prostaglandin E2 for induction of labour. Ir Med J 1984;77(3):70-2.

Carr CJ, Winckworth T, Finneran E. Trial of vaginal prostaglandin pessary for induction of labour. Ir J Med Sci 1987;156(9):265-7.

Carroll SG, Turner MJ, Stronge JM, et al. Management of antepartum spontaneous membrane rupture after one previous caesarean section. Eur J Obstet Gynecol Reprod Biol 1990;35(2-3):173-8.

Carter LV, Dooley DJ, Elliott CE, et al. Obstetrics in small hospitals. Med J Aust 1980;1(8):359-62.

Carty EM, Tier T. Birth planning. A reality-based script for building confidence. J Nurse Midwifery 1989;34(3):111-4.

Caspi B, Lancet M, Kessler I. Sinusoidal pattern of uterine contractions in abruptio placentae. Int J Gynaecol Obster 1980;17(6):615-6.

Castledine G, Oliver J, Turton P, et al. Fact-finding mission. Nurs Times 1985;81(21):23.

Catanese ML. Vaginal birth after cesarean: Recommendations, risks, realities, and the client's right to know. Holist Nurs Pract 1987;2(1):35-43.

Catanzarite VA. Risk of vaginal delivery after a prior cesarean section. Am J Obstet Gynecol 1987;157(1):213.

Catanzarite VA, Foster E, Robinette P, et al. Maternal death due to rupture of a low transverse cesarean section incision during labor at home. West J Med 1992;157(4):454-5.

Catanzarite VA, Mehalek KE, Wachtel T, et al. Sonographic diagnosis of traumatic and later recurrent uterine rupture. Am J Perinatol 1996;13(3):177-80.

Catlin AJ. When pregnant women and their physicians disagree on the need for cesarean section: No simple solution. Adv Pract Nurs Q 1998;4(2):23-9.

Caughey AB, Shipp TD, Repke JT, et al. Trial of labor after cesarean delivery: The effect of previous vaginal delivery. Am J Obstet Gynecol 1998;179(4):938-41.

Caughey AB, Shipp TD, Repke JT, et al. Rate of uterine rupture during a trial of labor in women with one or two prior cesarean deliveries. Am J Obstet Gynecol 1999;181(4):872-6.

Cavero CM, Fullerton JT, Bartlome JA.

Assessment of the process and outcomes of the first 1,000 births of a nurse-midwifery service. J Nurse Midwifery 1991;36(2):104-10.

Cepicky P, Stembera Z, Zeman J, et al. When is it possible to meet the wish of a woman to terminate her labour by caesarean section? Eur J Obstet Gynecol Reprod Biol 1991;38(2):109-12.

Chalmers B, Meyer D. What women say about their birth experiences: a Cross-cultural study. Journal of Psychosomatic Obstetrics & Gynecology 1994;15(4):211-8.

Chambliss LR, Daly C, Medearis AL, et al. The role of selection bias in comparing cesarean birth rates between physician and midwifery management. Obstet Gynecol 1992;80(2):161-5.

Chan BC, Lao TT. Influence of parity on the obstetric performance of mothers aged 40 years and above. Hum Reprod 1999;14(3):833-7.

Chan CC, To WW. Antepartum hemorrhage of unknown origin--What is its clinical significance? Acta Obstet Gynecol Scand 1999;78(3):186-90.

Chang C. Trial of labor with prior vertical cesarean incision. J Fam Pract 1997;45(5):380-1.

Chang MY, Tsieh TT, Chu KK, et al. An analysis of 121 trial labors following a previous cesarean section. Changgeng Yi Xue Za Zhi 1987;10(1):28-36.

Chanrachakul B, Chaturachinda K, Phuapradit W, et al. Cesarean and postpartum hysterectomy. Int J Gynaecol Obster 1996;54(2):109-13.

Chanrachakul B, Herabutya Y, Punyavachira P. Potential efficacy of nitric oxide for cervical ripening in pregnancy at term. Int J Gynaecol Obster 2000;71(3):217-9.

Chanrachakul B, Herabutya Y, Udomsubpayakul U. Epidemic of cesarean section at the general, private and university hospitals in Thailand. Journal of Obstetrics & Gynaecology Research 2000;26(5):357-61.

Chapman DJ, Perez-Escamilla R. Identification of risk factors for delayed onset of lactation. J Am Diet Assoc 1999;99(4):450-4; quiz 455-6.

Chapman K, Meire H, Chapman R. The value of serial ultrasounds in the management of recurrent uterine scar rupture. Br J Obstet Gynaecol 1994;101(6):549-51.

Chapman RH, Stone PW, Sandberg EA, et al. A comprehensive league table of cost-utility ratios and a sub-table of "panel-worthy" studies. Med Decis Making 2000;20(4):451-67.

Chapman SJ, Owen J, Hauth JC. One- versus two-layer closure of a low transverse cesarean: the Next pregnancy. Obstet Gynecol 1997;89(1):16-8.

Chaska BW, Mellstrom MS, Grambsch PM, et al. Influence of site of obstetric care and delivery on pregnancy management and outcome. J Am Board Fam Pract 1988;1(3):152-63.

Chattopadhyay K, Sengupta BS, Edress YB, et al. Vaginal birth after cesarean section: Management debate. Int J Gynaecol Obster 1988;26(2):189-96.

Chattopadhyay SK, Sherbeeni MM, Anokute CC. Planned vaginal delivery after two previous caesarean sections. Br J Obstet Gynaecol 1994;101(6):498-500.

Chauhan SP, Magann EF, Carroll CS, et al. Mode of delivery for the morbidly obese with prior cesarean delivery: Vaginal versus repeat cesarean section. Am J Obstet Gynecol 2001;185(2):349-54.

Chauhan SP, Troyer LR, Hendrix NW, et al. Neonatal acidemia with trial of labor among parturients with prior cesarean delivery: a Case-control study. J Matern Fetal Med 2000;9(5):278-81.

Chazotte C, Cohen WR. Catastrophic complications of previous cesarean section. Am J Obstet Gynecol 1990;163(3):738-42.

Chazotte C, Madden R, Cohen WR. Labor patterns in women with previous cesareans. Obstet Gynecol 1990;75(3 Pt 1):350-5.

Chelmow D, Kilpatrick SJ, Laros RK. Maternal and neonatal outcomes after prolonged latent phase. Obstet Gynecol 1993;81(4):486-91.

Chelmow D, Laros RK. Maternal and neonatal outcomes after oxytocin augmentation in patients undergoing a trial of labor after prior cesarean

delivery. Obstet Gynecol 1992;80(6):966-71.

Chen CH, Wang SY, Chang MY. Women's perceptions of helpful and unhelpful nursing behaviors during labor: a Study in Taiwan. Birth 2001;28(3):180-5.

Chen DC, Nommsen-Rivers L, Dewey KG, et al. Stress during labor and delivery and early lactation performance. Am J Clin Nutr 1998;68(2):335-44.

Chen HY, Chen SJ, Hsieh FJ. Observation of cesarean section scar by transvaginal ultrasonography. Ultrasound Med Biol 1990;16(5):443-7.

Chen KC, Hsieh TT. Rupture of gravid uterus: a Eight-year clinical analysis and review of the literature. Changgeng Yi Xue Za Zhi 1992;15(1):15-22.

Chen LH, Tan KH, Yeo GS. A ten-year review of uterine rupture in modern obstetric practice. Ann Acad Med Singapore 1995;24(6):830-5.

Cheng M, Hannah M. Breech delivery at term: a Critical review of the literature. Obstet Gynecol 1993;82(4 Pt 1):605-18.

Chervenak FA, Herslinger R, Freedman R, et al. Current perspectives on iatrogenic neonatal respiratory distress syndrome. J Reprod Med 1986;31(1):53-7.

Chervenak FA, McCullough LB. An ethically justified algorithm for offering, recommending, and performing cesarean delivery and its application in managed care practice. Obstet Gynecol 1996;87(2):302-5.

Chetty VK. Stochastic technology, production organization and costs. J Health Econ 1998;17(2):187-210.

Cheung TH, Leung A, Chang A. Macrosomic babies. Aust N Z J Obstet Gynaecol 1990;30(4):319-22.

Chew S, Biswas A. Caesarean and postpartum hysterectomy. Singapore Med J 1998;39(1):9-13.

Chew SY. Uterine rupture in labour. A 10-year review. Singapore Med J 1984;25(1):24-9.

Chez RA. Vaginal birth after cesarean section. Am J Obstet Gynecol 1989;160(5 Pt 1):1251.

Chez RA. Cervical ripening and labor induction after previous cesarean delivery. Clin Obstet Gynecol 1995;38(2):287-92.

Chez RA, Droegemueller W, Gant NF, Jr., et al. Clinical experience reported by candidates for the American Board of Obstetrics and Gynecology 1995 and 1997 oral examinations. Am J Obstet Gynecol 2001;185(6):1429-32.

Choy-Hee L, Raynor BD. Misoprostol induction of labor among women with a history of cesarean delivery. Am J Obstet Gynecol 2001;184(6):1115-7.

Chua S, Arulku maran S, Kurup A, et al. Oxytocin titration for induction of labour: a Prospective randomized study of 15 versus 30 minute dose increment schedules. Aust N Z J Obstet Gynaecol 1991;31(2):134-7.

Chua S, Arulkumaran S, Singh P, et al. Trial of labour after previous caesarean section: Obstetric outcome. Aust N Z J Obstet Gynaecol 1989;29(1):12-7.

Chua S, Kurup A, Arulkumaran S, et al. Augmentation of labor: Does internal tocography result in better obstetric outcome than external tocography? Obstet Gynecol 1990;76(2):164-7.

Chuang JH, Jenders RA. Trial of labor versus elective repeat cesarean section for the women with a previous cesarean section: a Decision analysis. Proceedings / AMIA Annual Symposium 1999:226-30.

Chuck FJ, Huffaker BJ. Labor induction with intravaginal misoprostol versus intracervical prostaglandin E2 gel (Prepidil gel): Randomized comparison. Am J Obstet Gynecol 1995;173(4):1137-42.

Chung A, Macario A, El-Sayed YY, et al. Cost-effectiveness of a trial of labor after previous cesarean. Obstet Gynecol 2001;97(6):932-41.

Clark JF, Westney LS, Lawyer CJ. Adolescent pregnancy: a 25-year review. J Natl Med Assoc 1987;79(4):377-80.

Clark L, Mugford M, Paterson C. How does the mode of delivery affect the cost of maternity care? Br J Obstet Gynaecol 1991;98(6):519-23.

Clark SL. Rupture of the scarred uterus. Obstetrics & Gynecology Clinics of North America 1988;15(4):737-44.

Clark SL, Eglinton GS, Beall M, et al. Effect of indication for previous cesarean section on subsequent delivery outcome in patients undergoing a trial of labor. J Reprod Med 1984;29(1):22-5.

Clark SL, Hankins GD, Dudley DA, et al. Amniotic fluid embolism: Analysis of the national registry. Am J Obstet Gynecol 1995;172(4 Pt 1):1158-67; discussion 1167-9.

Clark SL, Scott JR, Porter TF, et al. Is vaginal birth after cesarean less expensive than repeat cesarean delivery? Am J Obstet Gynecol 2000;182(3):599-602.

Clark SL, Yeh SY, Phelan JP, et al. Emergency hysterectomy for obstetric hemorrhage. Obstet Gynecol 1984;64(3):376-80.

Clarke SC, Taffel S. Changes in cesarean delivery in the United States, 1988 and 1993. Birth 1995;22(2):63-7.

Clarke SC, Taffel SM. Rates of cesarean and VBAC delivery, United States, 1994. Birth 1996;23(3):166-8.

Clarke SC, Taffel SM. State variation in rates of cesarean and VBAC delivery: 1989 and 1993. Stat Bull Metrop Insur Co 1996;77(1):28-36.

Cnattingius S, Forman MR, Berendes HW, et al. Delayed childbearing and risk of adverse perinatal outcome. A population-based study. JAMA 1992;268(7):886-90.

Coco AS, Gates TJ, Gallagher ME, et al. Association of attending physician specialty with the cesarean delivery rate in the same patient population. Fam Med 2000;32(9):639-44.

Cohen M, Carson BS. Respiratory morbidity benefit of awaiting onset of labor after elective cesarean section. Obstet Gynecol 1985;65(6):818-24.

Cohen NW, LaLeike S. What factors cause unnecessary cesareans? Midwifery Today Childbirth Educ 1992(23):16-7.

Cohen RL. Factors influencing maternal choice of childbirth alternatives. J Am Acad Child Psychiatry 1981;20(1):1-15.

Cohen S, Mitchell JL, Marx GF. Severe systemic reactions following administration of different uterotonic drugs. N Y State J Med 1983;83(8-10):1060-1061.

Coleman TL, Randall H, Graves W, et al. Vaginal birth after cesarean among women with gestational diabetes. Am J Obstet Gynecol 2001;184(6):1104-7.

Coltart TM, Davies JA, Katesmark M. Outcome of a second pregnancy after a previous elective caesarean section. Br J Obstet Gynaecol 1990;97(12):1140-3.

Combs CA, Murphy EL, Laros RK. Cost-benefit analysis of autologous blood donation in obstetrics. Obstet Gynecol 1992;80(4):621-5.

Combs CA, Singh NB, Khoury JC. Elective induction versus spontaneous labor after sonographic diagnosis of fetal macrosomia. Obstet Gynecol 1993;81(4):492-6.

Comried LA. Cost analysis: Initiation of HBMC and first CareMap. Nurs Econ 1996;14(1):34-9.

Connolly GA, Razak ARA, Conroy R, et al. A five year review of scar dehiscence in the Rotunda Hospital Dublin. Ir Med J 2001;94(6):176-178.

Coody D, Yetman RJ, Montgomery D, et al. Early hospital discharge and the timing of newborn metabolic screening. Clin Pediatr 1993;32(8):463-6.

Coonrod DV, Bay RC, Kishi GY. The epidemiology of labor induction: Arizona, 1997. Am J Obstet Gynecol 2000;182(6):1355-62.

Cotzias CS, Paterson-Brown S, Fisk NM. Obstetricians say yes to maternal request for elective caesarean section: a Survey of current opinion. Eur J Obstet Gynecol Reprod Biol 2001;97(1):15-6.

Coulter CH, Lehrfeld R. When push comes to shove: Implementing VBAC practice guidelines. Physician Exec 1995;21(6):30-5.

Cowan RK, Kinch RA, Ellis B, et al. Trial of labor following cesarean delivery. Obstet

Gynecol 1994;83(6):933-6.

Cragin EB. Conservatism in obstetrics. N Y Med J 1916;104(1):1-3.

Crane S, Chun B, Acker D. Treatment of obstetrical hemorrhagic emergencies. Curr Opin Obstet Gynecol 1993;5(5):675-82.

Cranley MS, Hedahl KJ, Pegg SH. Women's perceptions of vaginal and cesarean deliveries. Nurs Res 1983;32(10-15).

Creedy DK, Shochet IM, Horsfall J. Childbirth and the development of acute trauma symptoms: Incidence and contributing factors. Birth 2000;27(2):104-11.

Cronk M. Home birth after LSCS. Midwives Chronicle 1991;104(1239):110.

Crowley P. Interventions for preventing or improving the outcome of delivery at or beyond term. Cochrane Database of Systematic Reviews 2000(2):CD000170.

Cunha M, Bugalho A, Bique C, et al. Induction of labor by vaginal misoprostol in patients with previous cesarean delivery. Acta Obstet Gynecol Scand 1999;78(7):653-4.

Cunningham JD. Experiences of Australian mothers who gave birth either at home, at a birth centre, or in hospital labour wards. Soc Sci Med 1993;36(4):475-83.

Curet LB, Zachman RD, Rao AV, et al. Effect of mode of delivery on incidence of respiratory distress syndrome. Int J Gynaecol Obster 1988;27(2):165-70.

Curtin SC. Rates of cesarean birth and vaginal birth after previous cesarean, 1991-1995. Natl Vital Stat Rep 1997;45(11):Suppl 3.

Curtin SC. Recent changes in birth attendant, place of birth, and the use of obstetric interventions, United States, 1989-1997. J Nurse Midwifery 1999;44(4):349-54.

Curtin SC, Kozak LJ. Cesarean delivery rates in 1995 continue to decline in the United States. Birth 1997;24(3):194-6.

Curtin SC, Martin JA. Births: Preliminary data for 1999. Natl Vital Stat Rep 2000;48(14):1-20.

Curtin SC, Mathews TJ. U.S. obstetric procedures, 1998. Birth 2000;27(2):136-8.

Curtin SC, Park MM. Trends in the attendant, place, and timing of births, and in the use of obstetric interventions: United States, 1989-97. Natl Vital Stat Rep 1999;47(27):1-12.

Cyr RM, Usher RH, McLean FH. Changing patterns of birth asphyxia and trauma over 20 years. Am J Obstet Gynecol 1984;148(5):490-8.

d'Orsi E, Chor D, Giffin K, et al. Factors associated with vaginal birth after cesarean in a maternity hospital of Rio de Janeiro. Eur J Obstet Gynecol Reprod Biol 2001;97(2):152-7.

Dagher AP, Fishman EK. Uterine and bladder rupture during vaginal delivery in a patient with a prior cesarean section: Case report. Urol Radiol 1992;14(3):200-1.

Dainer M, Hall CD, Choe J, et al. Pregnancy following incontinence surgery. Int Urogynecol J Pelvic Floor Dysfunct 1998;9(6):385-90.

Dani C, Reali MF, Bertini G, et al. Risk factors for the development of respiratory distress syndrome and transient tachypnoea in newborn infants. Italian Group of Neonatal Pneumology. Eur Respir J 1999;14(1):155-9.

Dannenbring D, Stevens MJ, House AE. Predictors of childbirth pain and maternal satisfaction. J Behav Med 1997;20(2):127-42.

Darby M. Reimbursement has small impact on c-section rates. In: Report on Medical Guidelines and Outcome Research. Alexandria, VA: Capitol Publications; 1992. p. 8-10.

Darling MR. Elective or selective caesarean delivery of the small baby? Lancet 1997;349(9053):666.

Daus KM. Spontaneous rupture of low transverse cesarean scar. South Med J 1995;88(5):600.

Davies GA, Hahn PM, McGrath MM. Vaginal birth after cesarean. Physicians' perceptions and practice. J Reprod Med 1996;41(7):515-20.

Davis LG, Riedmann GL, Sapiro M, et al. Cesarean section rates in low-risk private patients managed by certified nurse-midwives

and obstetricians. J Nurse Midwifery 1994;39(2):91-7.

Davis-Floyd R, Davis E. Intuition as authoritative knowledge in midwifery and homebirth. Med Anthropol Q 1996;10(2):237-69.

Daviss BA. Vaginal delivery after caesarean section. Study's focus on induction v spontaneous labour neglects spontaneous deliver. BMJ 2001;323(7324):1307.

Dawson NV, Acheson LS. Trade-offs in clinical policy making. Arch Fam Med 1997;6(2):127-8.

Day A, MacLennan A, Green R. A comparison of intravaginal PGF2 alpha and intravenous oxytocin to stimulate labour after membrane rupture. Aust N Z J Obstet Gynaecol 1985;25(4):252-5.

de Jong P. Trial of labor following cesarean section--a Study of 212 patients. Int J Gynaecol Obster 1987;25(5):405-11.

de Meeus JB, Ellia F, Magnin G. External cephalic version after previous cesarean section: a Series of 38 cases. Eur J Obstet Gynecol Reprod Biol 1998;81(1):65-8.

de Mello e Souza C. C-sections as ideal births: the Cultural constructions of beneficence and patients' rights in Brazil. Camb Q Healthc Ethics 1994;3(3):358-66.

De Muylder X. Vaginal delivery after caesarean section: Is it safe in a developing country? Aust N Z J Obstet Gynaecol 1988;28(2):99-102.

de Vries LS, Regev R, Dubowitz LM, et al. Perinatal risk factors for the development of extensive cystic leukomalacia. Am J Dis Child 1988;142(7):732-5.

DeJoy S, O'Grady JP, Burkman RT. The risks of lowering the cesarean-delivery rate. N Engl J Med 1999;341(1):53-4; discussion 54-5.

Del Valle GO, Adair CD, Sanchez-Ramos L, et al. Cervical ripening in women with previous cesarean deliveries. International Journal of Gynecology & Obstetrics 1994;47(1):17-21.

Delarue T, Pele P. Prevention of rupture of the uterus before or during birth in women with previous caesarian sections: Report on 14 cases. J Gynecol Obstet Biol Reprod 1981;10(3):259-67.

Delarue T, Pele P, Pelletier P, et al. Prognosis for delivery in women with previous cesarean section. Apropos of 209 cases. J Gynecol Obstet Biol Reprod 1983;12(2):193-206.

Dellinger EH, Boehm FH, Crane MM. Electronic fetal heart rate monitoring: early neonatal outcomes associated with normal rate, fetal stress, and fetal distress. Am J Obstet Gynecol 2000;182(1 Pt 1):214-20.

Delpapa EH, Mueller-Heubach E. Pregnancy outcome following ultrasound diagnosis of macrosomia. Obstet Gynecol 1991;78(3 Pt 1):340-3.

DeMaio H, Edwards RK, Euliano TY, et al. Vaginal birth after cesarean delivery: An historic cohort cost analysis. Am J Obstet Gynecol 2002;186:890-2.

Demianczuk NN, Hunter DJ, Taylor DW. Trial of labor after previous cesarean section: Prognostic indicators of outcome. Am J Obstet Gynecol 1982;142(6 Pt 1):640-2.

DeMott RK, Sandmire HF. The Green Bay cesarean section study. I. The physician factor as a determinant of cesarean birth rates. Am J Obstet Gynecol 1990;162(6):1593-9; discussion 1599-602.

Deutchman ME, Sills D, Connor PD. Perinatal outcomes: a Comparison between family physicians and obstetricians. J Am Board Fam Pract 1995;8(6):440-7.

Devlin R. Childbirth on trial. Nurs Times 1986;82(9):19-20.

Devoe LD, Croom CS, Youssef AA, et al. The prediction of "controlled" uterine rupture by the use of intrauterine pressure catheters. Obstet Gynecol 1992;80(4):626-9.

Devoe LD, Samuel S, Prescott P, et al. Predicting the duration of the first stage of spontaneous labor using a neural network. J Matern Fetal Med 1996;5(5):256-61.

Dhall K, Mittal SC, Grover V, et al. Childbirth following primary cesarean section--Evaluation

of a scoring system. Int J Gynaecol Obster 1987;25(3):199-205.

Diani F, Venanzi S, Zanconato G, et al. Fetal macrosomia and management of delivery. Clin Exp Obstet Gynecol 1997;24(4):212-4.

Dickson MJ. Vaginal delivery after previous caesarean section for failure of second stage of labour. Br J Obstet Gynaecol 1999;106(7):749.

Dildy GA, Jackson GM, Fowers GK, et al. Very advanced maternal age: Pregnancy after age 45. Am J Obstet Gynecol 1996;175(3 Pt 1):668-74.

Dilks FM, Beal JA. Role of self-efficacy in birth choice. J Perinat Neonat Nurs 1997;11(1):1-9.

DiMatteo MR, Morton SC, Lepper HS, et al. Cesarean childbirth and psychosocial outcomes: a Meta-analysis. Health Psychol 1996;15(4):303-14.

Dishman RK, Buckworth J. Increasing physical activity: a Quantitative synthesis. Med Sci Sports Exerc 1996;28(6):706-19.

Donnelly JP, Franzoni KT. Uterine rupture. A 30-year survey. Obstet Gynecol 1964;23:774.

Donnelly V, Fynes M, Campbell D, et al. Obstetric events leading to anal sphincter damage. Obstet Gynecol 1998;92(6):955-61.

Drummond M, O'Brien B, Stoddart G, et al. Methods for the Economic Evaluation of Health Care Programmes, 2nd Ed. New York: Oxford University Press; 1997.

Dubay L, Kaestner R, Waidmann T. The impact of malpractice fears on cesarean section rates. J Health Econ 1999;18(4):491-522.

Duff C, Sinclair M. Exploring the risks associated with induction of labour: a Retrospective study using the NIMATS database. Northern Ireland Maternity System. J Adv Nurs 2000;31(2):410-7.

Duff P, Southmayd K, Read JA. Outcome of trial of labor in patients with a single previous low transverse cesarean section for dystocia. Obstet Gynecol 1988;71(3 Pt 1):380-4.

Dulitzki M, Soriano D, Schiff E, et al. Effect of very advanced maternal age on pregnancy outcome and rate of cesarean delivery. Obstet Gynecol 1998;92(6):935-9.

Durik AM, Hyde JS, Clark R. Sequelae of cesarean and vaginal deliveries: psychosocial outcomes for mothers and infants. Dev Psychol 2000;36(2):251-60.

Dyack C, Hughes PF, Simbakalia JB. Vaginal birth after cesarean section in the grand multipara with a previous lower segment scar. Int J Gynaecol Obster 1996;55(2):167-8.

Dyack C, Hughes PF, Simbakalia JB. Vaginal delivery in the grand multipara following previous lower segment cesarian section. Journal of Obstetrics & Gynaecology Research 1997;23(2):219-22.

Eakes M. Economic considerations for epidural anesthesia in childbirth. Nurs Econ 1990;8(5):329-32.

Eakins PS, O'Reilly WB, May LJ, et al. Obstetric outcomes at the birth place in Menlo Park: the First seven years. Birth 1989;16(3):123-9.

Eddy DM. Clinical decision making: from Theory to practice. Guidelines for policy statements: the Explicit approach. JAMA 1990;263(16):2239-40.

Edelin KC, Oellerich D, Larrieux JR. The effect of induced abortion on the risks and outcome of a trial of labor after a previous cesarean birth. J Natl Med Assoc 1988;80(9):993-5.

Eden RD, Parker RT, Gall SA. Rupture of the pregnant uterus: a 53-year review. Obstet Gynecol 1986;68(5):671-4.

Edwards DR, Porter SA, Stein GS. A pilot study of postnatal depression following caesarean section using two retrospective self-rating instruments. J Psychosom Res 1994;38(2):111-7.

Egarter C, Kofler E, Fitz R, et al. Is induction of labor indicated in prolonged pregnancy? Results of a prospective randomised trial. Gynecol Obstet Invest 1989;27(1):6-9.

Eglinton GS, Phelan JP, Yeh S, et al. Outcome of a trial of labor after prior cesarean delivery. J Reprod Med 1984;29(1):3-8.

Eidelman AI, Schimmel MS, Bromiker R, et al.

Pediatric coverage of the delivery room: an Analysis of manpower utilization. J Perinatol 1998;18(2):131-4.

Eisenberg JM. Sociological influences on the decision-making by clinicians. Ann Intern Med 1979;90:957.

Eisenberg JM. Doctors' Decisions and the Cost of Medical Care. Ann Arbor, Mich: Health Administration Press; 1986.

Ekblad U, Erkkola R. Intracervical prostaglandin E2 gel for cervical ripening. Ann Chir Gynaecol Suppl 1987;202:23-5.

Ekblad U, Vilpa T. Pregnancy in women over forty. Ann Chir Gynaecol Suppl 1994;208:68-71.

Ekeocha CE, Jackson P. The 'birth plan' experience. Br J Obstet Gynaecol 1985;92(1):97-101.

Ekman G, Granstrom L, Ulmsten U. Induction of labor with intravenous oxytocin or vaginal PGE2 suppositories. A randomized study. Acta Obstet Gynecol Scand 1986;65(8):857-9.

el Gammal NA, Jallad KB, O'Deh H M. Breech vaginal delivery after one cesarean section: a Retrospective study. Int J Gynaecol Obster 1990;33(2):99-102.

Elferink-Stinkens PM, Brand R, le Cessie S, et al. Large differences in obstetrical intervention rates among Dutch hospitals, even after adjustment for population differences. Eur J Obstet Gynecol Reprod Biol 1996;68(1-2):97-103.

Elferink-Stinkens PM, Brand R, Van Hemel OJ. Trends in caesarean section rates among high- and medium-risk pregnancies in The Netherlands 1983-1992. Eur J Obstet Gynecol Reprod Biol 1995;59(2):159-67.

Elferink-Stinkens PM, Van Hemel OJ, Brand R, et al. The Perinatal Database of the Netherlands. Eur J Obstet Gynecol Reprod Biol 2001;94(1):125-38.

Elkady AA, Bayomy HM, Bekhiet MT, et al. A review of 126 cases of ruptured gravid uterus. Int Surg 1993;78(3):231-5.

Elkins T, Onwuka E, Stovall T, et al. Uterine rupture in Nigeria. J Reprod Med 1985;30(3):195-9.

Elliott JP, Flaherty JF. The use of breast stimulation to ripen the cervix in term pregnancies. Am J Obstet Gynecol 1983;145(5):553-6.

Elliott JP, Flaherty JF. The use of breast stimulation to prevent postdate pregnancy. Am J Obstet Gynecol 1984;149(6):628-32.

Elliott JP, Russell MM, Dickason LA. The labor-adjusted cesarean section rate--a more informative method than the cesarean section "rate" for assessing a practitioner's labor and delivery skills. Am J Obstet Gynecol 1997;177(1):139-43.

Emerson MV, Pieramici DJ, Stoessel KM, et al. Incidence and rate of disappearance of retinal hemorrhage in newborns. Ophthalmology 2001;108(1):36-9.

Engelsen IB, Albrechtsen S, Iversen OE. Peripartum hysterectomy - incidence and maternal morbidity. Acta Obstet Gynecol Scand 2001;80(5):409-12.

Enkin M. Labour and Delivery follwing previous Caesarean Section. In: Chalmers I, Enkin M, Kierse MJNC, editors. Effective Care in Pregnancy and Childbirth. Oxford: Oxford University Press; 1989. p. 1196-1215.

Enthoven A, Kronick R. A consumer-choice health plan for the 1990s. Universal health insurance in a system designed to promote quality and economy (2). N Engl J Med 1989;320(2):94-101.

Erb L, Hill G, Houston D. A survey of parent's attitudes toward their cesaran births in Manitoba hospitals. Birth 1983;10(2):85-92.

Eriksen NL, Buttino L. Vaginal birth after cesarean: a Comparison of maternal and neonatal morbidity to elective repeat cesarean section. Am J Perinatol 1989;6(4):375-9.

Esposito MA, Menihan CA, Malee MP. Association of interpregnancy interval with uterine scar failure in labor: a Case-control study. Am J Obstet Gynecol 2000;183(5):1180-3.

Evans MI, Dougan MB, Moawad AH, et al. Ripening of the human cervix with porcine ovarian relaxin. Am J Obstet Gynecol 1983;147(4):410-4.

Evans MI, Richardson DA, Sholl JS, et al. Cesarean section. Assessment of the convenience factor. J Reprod Med 1984;29(9):670-6.

Ewen SP, Notley RG, Coats PM. Bladder laceration associated with uterine scar rupture during vaginal delivery. Br J Urol 1994;73(6):712-3.

Fanaroff AA, Wright LL, Stevenson DK, et al. Very-low-birth-weight outcomes of the National Institute of Child Health and Human Development Neonatal Research Network, May 1991 through December 1992. Am J Obstet Gynecol 1995;173(5):1423-31.

Farmakides G, Duvivier R, Schulman H, et al. Vaginal birth after two or more previous cesarean sections. Am J Obstet Gynecol 1987;156(3):565-6.

Farmer KC, Schwartz WJ, Rayburn WF, et al. A cost-minimization analysis of intracervical prostaglandin E2 for cervical ripening in an outpatient versus inpatient setting. Clin Ther 1996;18(4):747-56; discussion 702.

Farmer RM, Kirschbaum T, Potter D, et al. Uterine rupture during trial of labor after previous cesarean section. Am J Obstet Gynecol 1991;165(4 Pt 1):996-1001.

Fawcett J, Henklein JC. Antenatal education for cesarean birth: Extension of a field test. J Obstet Gynecol Neonat Nurs 1987;16(1):61-5.

Fawcett J, Knauth D. The factor structure of the perception of birth scale. Nurs Res 1996;45(2):83-6.

Fawcett J, Pollio N, Tully A. Women's perceptions of cesarean and vaginal delivery: Another look. Res Nurs Health 1992;15(6):439-46.

Fawcett J, Tulman L, Spedden J. Responses to vaginal birth after cesarean section. J Obstet Gynecol Neonat Nurs 1994;23(3):253-9.

Faxelius G, Hagnevik K, Lagercrantz H, et al. Catecholamine surge and lung function after delivery. Arch Dis Child 1983;58(4):262-6.

Fedorkow DM, Nimrod CA, Taylor PJ. Ruptured uterus in pregnancy: a Canadian hospital's experience. Can Med Assoc J 1987;137(1):27-9.

Feldman GB, Freiman JA. Prophylactic cesarean section at term? N Engl J Med 1985;312(19):1264-7.

Ferguson JE, 2nd, Newberry YG, DeAngelis GA, et al. The fetal-pelvic index has minimal utility in predicting fetal-pelvic disproportion. Am J Obstet Gynecol 1998;179(5):1186-92.

Field MA. Controlling the woman to protect the fetus. Law Med Health Care 1989;17(2):114-29.

Finkelstein BS, Harper DL, Rosenthal GE. Assessing variation, validity, and reproducibility in patient satisfaction with hospital obstetrical care. AHSR FSHR Annu Meet Abstr Book;13.

Finkelstein BS, Harper DL, Rosenthal GE. Does length of hospital stay during labor and delivery influence patient satisfaction? Results from a regional study. Am J Manag Care 1998;4(12):1701-8.

Finkelstein BS, Harper DL, Rosenthal GE. Patient assessments of hospital maternity care: a Useful tool for consumers? Health Serv Res 1999;34(2):623-40.

Finkelstein BS, Singh J, Silvers JB, et al. Patient and hospital characteristics associated with patient assessments of hospital obstetrical care. Med Care 1998;36(8 Suppl):AS68-78.

Finkler MD, Wirtschafter DD. Cost-effectiveness and obstetric services. Med Care 1991;29(10):951-63.

Finkler MD, Wirtschafter DD. Why pay extra for cesarean-section deliveries? Inquiry 1993;30(2):208-15.

Finkler MD, Wirtschafter DD. One health maintenance organization's experience: Obstetric costs depend more on staffing patterns than on mode of delivery. J Perinatol 1997;17(2):148-55.

Finkler SA. The distinction between cost and charges. Ann Intern Med 1982;96(1):102-9.

Finley BE, Gibbs CE. Emergent cesarean

delivery in patients undergoing a trial of labor with a transverse lower-segment scar. Am J Obstet Gynecol 1986;155(5):936-9.

Firth PA, Chu J, Bell MA, et al. Changing practice habits of King County obstetricians. Obstet Gynecol 1988;72(3 Pt 1):419-22.

Flamm B, Mac Donald D, Shearer E, et al. Should the electronic fetal monitor always be used for women in labor who are having a vaginal birth after a previous cesarean section? Birth 1992;19(1):31-5.

Flamm BL. Vaginal birth after cesarean section: Controversies old and new. Clin Obstet Gynecol 1985;28(4):735-44.

Flamm BL. The patient who demands cesarean delivery. In: EJ Q, Flamm B, editors. Cesarean section: Guidelines for appropriate utilization. New York: Springer-Verlag; 1995. p. 207-21.

Flamm BL. Once a cesarean, always a controversy. Obstet Gynecol 1997;90(2):312-5.

Flamm BL, Anton D, Goings JR, et al. Prostaglandin E2 for cervical ripening: a Multicenter study of patients with prior cesarean delivery. Am J Perinatol 1997;14(3):157-60.

Flamm BL, Dunnett C, Fischermann E, et al. Vaginal delivery following cesarean section: Use of oxytocin augmentation and epidural anesthesia with internal tocodynamic and internal fetal monitoring. Am J Obstet Gynecol 1984;148(6):759-763.

Flamm BL, Fried MW, Lonky NM, et al. External cephalic version after previous cesarean section. Am J Obstet Gynecol 1991;165(2):370-2.

Flamm BL, Geiger AM. Vaginal birth after cesarean delivery: an Admission scoring system. Obstet Gynecol 1997;90(6):907-10.

Flamm BL, Goings JR. Vaginal birth after cesarean section: is Suspected fetal macrosomia a contraindication? [see comments]. Obstet Gynecol 1989;74(5):694-7.

Flamm BL, Goings JR, Fuelberth NJ, et al. Oxytocin during labor after previous cesarean section: Results of a multicenter study. Obstet Gynecol 1987;70(5):709-12.

Flamm BL, Goings JR, Liu Y, et al. Elective repeat cesarean delivery versus trial of labor: a Prospective multicenter study. Obstet Gynecol 1994;83(6):927-32.

Flamm BL, Lim OW, Jones C, et al. Vaginal birth after cesarean section: Results of a multicenter study. Am J Obstet Gynecol 1988;158(5):1079-84.

Flamm BL, Newman LA, Thomas SJ, et al. Vaginal birth after cesarean delivery: Results of a 5-year multicenter collaborative study. Obstet Gynecol 1990;76(5 Pt 1):750-4.

Flanagan TA, Mulchahey KM, Korenbrot CC, et al. Management of term breech presentation. Am J Obstet Gynecol 1987;156(6):1492-502.

Fleissig A. Are women given enough information by staff during labour and delivery? Midwifery 1993;9(2):70-5.

Fleissig A. Prevalence of procedures in childbirth. BMJ 1993;306(6876):494-5.

Fletcher H, McCaw-Binns A. Rupture of the uterus with misoprostol (prostaglandin E1) used for induction of labour. J Obstet Gynaecol 1998;18(2):184-187.

Fogarty JP. Twin vaginal delivery after a previous cesarean delivery for twins. J Am Board Fam Pract 1993;6(6):600-3.

Foote AJ, Giles WB. Review of obstetric operative intervention rates. Asia Oceania J Obstet Gynaecol 1992;18(3):195-8.

Forsnes EV, Browning JE, Gherman RB. Bladder rupture associated with uterine rupture. A report of two cases occurring during vaginal birth after cesarean. J Reprod Med 2000;45(3):240-2.

Francis KT. Status of the year 2000 health goals for physical activity and fitness. Phys Ther 1999;79(4):405-14.

Franco S. VBAC: the Lesson of labor. Midwifery Today 2000(53):41.

Fraser W, Maunsell E, Hodnett E, et al. Randomized controlled trial of a prenatal vaginal birth after cesarean section education and support program. Childbirth Alternatives Post-

Cesarean Study Group. Am J Obstet Gynecol 1997;176(2):419-25.

Fraser W, Usher RH, McLean FH, et al. Temporal variation in rates of cesarean section for dystocia: Does "convenience" play a role? Am J Obstet Gynecol 1987;156(2):300-4.

French L. Trial of labor after cesarean section. J Fam Pract 1996;43(6):538-9.

Fribourg S. Previous cesarean delivery and trial of labor. Am J Obstet Gynecol 1987;157(4 Pt 1):1014-5.

Friedman EA. Correlation between maximum cervical dilatation at cesarean delivery and subsequent vaginal birth after cesarean delivery. Obstet Gynecol 1997;90(2):316-7.

Frigoletto FD, Jr., Lieberman E, Lang JM, et al. A clinical trial of active management of labor. [erratum appears in N Engl J Med 1995 Oct 26;333(17):1163.]. N Engl J Med 1995;333(12):745-50.

Fruchter O. Trial of labor compared with an elective second cesarean section. N Engl J Med 1997;336(9):658; discussion 659.

Fujii T, Kozuma S, Unno N, et al. Successful pregnancy following antenatal closure of uterine wall defect. Int J Gynaecol Obster 2000;68(3):261-2.

Gaffney G, Flavell V, Johnson A, et al. Cerebral palsy and neonatal encephalopathy. Arch Dis Child Fetal Neonatal Ed 1994;70(3):F195-200.

Gafni A, Goeree R, Myhr TL, et al. Induction of labour versus expectant management for prelabour rupture of the membranes at term: an Economic evaluation. TERMPROM Study Group. Term Prelabour Rupture of the Membranes. Can Med Assoc J 1997;157(11):1519-25.

Galbraith RS. Incidence of neonatal sixth nerve palsy in relation to mode of delivery. Am J Obstet Gynecol 1994;170(4):1158-9.

Gamble JA, Creedy DK. Women's request for a cesarean section: a Critique of the literature. Birth 2000;27(4):256-63.

Gamble JA, Creedy DK. Women's preference for a cesarean section: Incidence and associated factors. Birth 2001;28(2):101-10.

Garcia FA, Miller HB, Huggins GR, et al. Effect of academic affiliation and obstetric volume on clinical outcome and cost of childbirth. Obstet Gynecol 2001;97(4):567-76.

Gardeil F, Daly S, Turner MJ. Uterine rupture in pregnancy reviewed. Eur J Obstet Gynecol Reprod Biol 1994;56(2):107-10.

Garel M, Lelong N, Marchand A, et al. Psychosocial consequences of caesarean childbirth: a Four-year follow-up study. Early Hum Dev 1990;21(2):105-14.

Garite TJ, Freeman RK, Nageotte MP. Fetal maturity cascade: a Rapid and cost-effective method for fetal lung maturity testing. Obstet Gynecol 1986;67(5):619-22.

Gates PE. Think globally, act locally: an Approach to implementation of clinical practice guidelines. Jt Comm J Qual Improv 1995;21(2):71-84.

Geary M, Fanagan M, Boylan P. Maternal satisfaction with management in labour and preference for mode of delivery. J Perinat Med 1997;25(5):433-9.

Gee H. Delivery following previous Caesarean section. Curr Obstet Gynaecol 2000;10(2):86-90.

Gellman E, Goldstein MS, Kaplan S, et al. Vaginal delivery after cesarean section. Experience in private practice. JAMA 1983;249(21):2935-7.

Gemer O, Kopmar A, Sassoon E, et al. Neglected transverse lie with uterine rupture. Arch Gynecol Obstet 1993;252(3):159-60.

Gemer O, Segal S, Sassoon E. Detection of scar dehiscence at delivery in women with prior cesarean section. Acta Obstet Gynecol Scand 1992;71(7):540-2.

Gerbaud L, Biolay S, Venet M, et al. [Comparative evaluation of maternity hospitals in Auvergne: from planning to contracting]. Cah Sociol Demogr Med 1998;38(4):297-323.

Gherman RB, Heath T. Trial of labor after cesarean delivery: a Pilot study of oral

misoprostol for preinduction cervical ripening. Obstet Gynecol 2001;97(4 Suppl. 1):S68.

Gherman RB, McBrayer S, Browning J. Uterine rupture associated with vaginal birth after cesarean section: a Complication of intravaginal misoprostol? Gynecol Obstet Invest 2000;50(3):212-3.

Gibbs CE. Planned vaginal delivery following cesarean section. Clin Obstet Gynecol 1980;23(2):507-15.

Gibson DH. Vaginal delivery after caesarean section in primigravidae. Ir J Med Sci 1988;157(9):290-2.

Gifford DS, Keeler E, Kahn KL. Reductions in cost and cesarean rate by routine use of external cephalic version: a Decision analysis. Obstet Gynecol 1995;85(6):930-6.

Gifford DS, Morton SC, Fiske M, et al. Lack of progress in labor as a reason for cesarean. Obstet Gynecol 2000;95(4):589-95.

Gifford DS, Pearson M, Flamm B, et al. Factors which influence the decision to perform a cesarean delivery. Abstr Book Assoc Health Serv Res;14:271-2.

Gilbert WM, Nesbitt TS, Danielsen B. Childbearing beyond age 40: Pregnancy outcome in 24,032 cases. Obstet Gynecol 1999;93(1):9-14.

Gillette RD. Cost of vaginal delivery vs. repeated cesarean section. Am Fam Physician 1996;53(7):2284, 2287.

Gilson GJ, Izquierdo LA, Chatterjee MS, et al. Prevention of cesarean section - Does intracervical dinoprostone work? West J Med 1993;159(2):149-152.

Gilstrap LC, 3rd, Hauth JC, Schiano S, et al. Neonatal acidosis and method of delivery. Obstet Gynecol 1984;63(5):681-5.

Gilstrap LC, 3rd, Leveno KJ, Burris J, et al. Diagnosis of birth asphyxia on the basis of fetal pH, Apgar score, and newborn cerebral dysfunction. Am J Obstet Gynecol 1989;161(3):825-30.

Gimovsky ML, Bayer-Zwirello LA, Plevyak M. Fetal heart rate monitoring casebook. Amnioinfusion with uterine dehiscence and fetal distress. J Perinatol 1997; 17(1):83-6.

Girard M. Episiotomy and faecal incontinence. Lancet 1999;354(9196):2169.

Glasser M. Strategies to avoid unnecessary cesarean sections. J Fam Pract 1988;27(5):514-8.

Glasser M. Cesarean section: Science or ritual surgery? Med Hypotheses 1991;34(1):73-80.

Gleicher N. Cesarean section rates in the United States. The short-term failure of the National Consensus Development Conference in 1980. JAMA 1984;252(23):3273-6.

Gleicher N. The cesarean-section epidemic. Mt Sinai J Med 1986;53(7):563-5.

Gleicher N. Mandatory trial of labor after cesarean delivery: an Alternative viewpoint. Obstet Gynecol 1991;78(4):727-8.

Gobis LJ. Recent developments in health law relevant to health care providers. Nurse Pract 1992;17(3):77-80.

Goer H. Misoprostol and uterine rupture. Birth 2000;27(3):224-5.

Goeree R, Hannah M, Hewson S. Cost-effectiveness of induction of labour versus serial antenatal monitoring in the Canadian Multicentre Postterm Pregnancy Trial. Can Med Assoc J 1995;152(9):1445-50.

Goetzl L, Shipp TD, Cohen A, et al. Oxytocin dose and the risk of uterine rupture in trial of labor after cesarean. Obstet Gynecol 2001;97(3):381-4.

Goh JT. Genital tract fistula repair on 116 women. Aust N Z J Obstet Gynaecol 1998;38(2):158-61.

Gold M, Siegel J, Russell L, et al. Cost-Effectiveness in Health and Medicine. New York: Oxford University Press; 1996.

Gold RB, Kenney AM, Singh S. Paying for maternity care in the United States. Fam Plann Perspect 1987;19(5):190-206.

Goldberger SB, Rosen DJ, Michaeli G, et al. The

use of PGE2 for induction of labor in parturients with a previous cesarean section scar. Acta Obstet Gynecol Scand 1989;68(6):523-6.

Goldfarb MG. Who recieves cesareans: Patient and hospital characteristics. Hospital Cost and Utilization Project-Research Note No. 4. National Center for Health Services Research 1984.

Goldfarb MG, Coffey RM. Case-mix differences between teaching and nonteaching hospitals. Inquiry 1987;24(1):68-84.

Goldfarb NI, Hillman AL, Eisenberg JM, et al. Impact of a mandatory Medicaid case management program on prenatal care and birth outcomes. A retrospective analysis. Med Care 1991;29(1):64-71.

Goldman G, Pineault R, Bilodeau H, et al. Effects of patient, physician and hospital characteristics on the likelihood of vaginal birth after previous cesarean section in Quebec. Can Med Assoc J 1990;143(10):1017-24.

Goldman G, Pineault R, Potvin L, et al. Factors influencing the practice of vaginal birth after cesarean section. Am J Public Health 1993;83(8):1104-8.

Goldman GA, Kaplan B, Neri A, et al. The grand multipara. Eur J Obstet Gynecol Reprod Biol 1995;61(2):105-9.

Goldman GA, Kaplan B, Rabinerson D, et al. Vaginal delivery following caesarean section - the use of oxytocin and prostaglandins. J Obstet Gynaecol 1998;18(4):328-330.

Gonen O, Rosen DJ, Dolfin Z, et al. Induction of labor versus expectant management in macrosomia: a Randomized study. Obstet Gynecol 1997;89(6):913-7.

Gonzalves PE, Hardin JJ. Coordinated care early discharge of postpartum patients at Irwin Army Community Hospital. Mil Med 1993;158(12):820-2.

Goodlin RC. Low birth weight infants. Obstet Gynecol 1984;64(3):446-7.

Goodlin RC. Uterine activity during spontaneous labour after previous lower-segment caesarean section. Br J Obstet Gynaecol 1990;97(3):277-8.

Goodlin RC. Correlation between maximum cervical dilatation at cesarean delivery and subsequent vaginal birth after cesarean delivery. Obstet Gynecol 1997;90(2):316.

Gordon D, Milberg J, Daling J, et al. Advanced maternal age as a risk factor for cesarean delivery. Obstet Gynecol 1991;77(4):493-7.

Gordon H. Uterine rupture and its association with oxytocic drugs: The Northwick Park Hospital experience. J Obstet Gynaecol 1988;8(SUPPL. 1):S16-S17.

Gordon NP, Walton D, McAdam E, et al. Effects of providing hospital-based doulas in health maintenance organization hospitals. Obstet Gynecol 1999;93(3):422-6.

Gotoh H, Masuzaki H, Yoshida A, et al. Predicting incomplete uterine rupture with vaginal sonography during the late second trimester in women with prior cesarean. Obstet Gynecol 2000;95(4):596-600.

Gough GW, Randall NJ, Genevier ES, et al. Head-to-cervix forces and their relationship to the outcome of labor. Obstet Gynecol 1990;75(4):613-8.

Gould JB, Davey B, Stafford RS. Socioeconomic differences in rates of cesarean section. [see comments.]. N Engl J Med 1989;321(4):233-9.

Gower RH, Toraya J, Miller JM. Laminaria for preinduction cervical ripening. Obstet Gynecol 1982;60(5):617-9.

Goyert GL, Bottoms SF, Treadwell MC, et al. The physician factor in cesarean birth rates. N Engl J Med 1989;320(11):706-9.

Graham AR. Trial labor following previous cesarean section. Am J Obstet Gynecol 1984;149(1):35-45.

Graham EM, Forouzan I, Morgan MA. A retrospective analysis of Erb's palsy cases and their relation to birth weight and trauma at delivery. J Matern Fetal Med 1997;6(1):1-5.

Graham WJ, Hundley V, McCheyne AL, et al. An investigation of women's involvement in the decision to deliver by caesarean section. Br J Obstet Gynaecol 1999;106(3):213-20.

Granovsky-Grisaru S, Shaya M, Diamant YZ. The management of labor in women with more than one uterine scar: is a repeat cesarean section really the only "safe" option? J Perinat Med 1994;22(1):13-7.

Grazier KL, McGuire TG. Payment systems and hospital resource use: a comparative analysis of psychiatric, medical and obstetric services. Adv Health Econ Health Serv Res Suppl 1987;8:75-95.

Green JM. Expectations, experiences and psychological outcomes of childbirth: a prospective study of 825 women. Birth 1990;17(1):15-24.

Green JM, Coupland VA, Kitzinger JV. Great expectations: A prospective study of women's expectations and experiences of childbirth. Cambridge: University of Cambridge, Childcare and Develment Group; 1988.

Green L. VBAC clinical policy: an AAFP guideline. Am Fam Physician 1995;52(6):1655.

Green-Thompson RW. Antepartum haemorrhage. Clin Obstet Gynaecol 1982;9(3):479-515.

Greene MF. Vaginal delivery after cesarean section - Is the risk acceptable? N Engl J Med 2001;345(1):54-55.

Greene MF. Vaginal delivery after cesarean section--is the risk acceptable? [letter; comment]. N Engl J Med 2001;345(1):54-5.

Greenhalgh R, Slade P, Spiby H. Fathers' coping style, antenatal preparation, and experiences of labor and the postpartum. Birth 2000;27(3):177-84.

Gregory K, Henry O, Anderson E. Vaginal birth after cesarean: what should we be monitoring. AHSR FSHR Annu Meet Abstr Book;11:92.

Gregory KD, Henry OA, Gellens AJ, et al. Repeat cesareans: how many are elective? Obstet Gynecol 1994;84(4):574-8.

Gregory KD, Henry OA, Ramicone E, et al. Maternal and infant complications in high and normal weight infants by method of delivery. Obstet Gynecol 1998;92(4 Pt 1):507-13.

Gregory KD, Korst LM, Cane P, et al. Vaginal birth after cesarean and uterine rupture rates in California. Obstet Gynecol 1999;94(6):985-9.

Gregory KD, Ramicone E, Chan L, et al. Cesarean deliveries for Medicaid patients in public and private hospitals. Abstr Book Assoc Health Serv Res;14:125-6.

Gregory KD, Ramicone E, Chan L, et al. Cesarean deliveries for medicaid patients: a comparison in public and private hospitals in Los Angeles county. Am J Obstet Gynecol 1999;180(5):1177-84.

Greis JB, Bieniarz J, Scommegna A. Comparison of maternal and fetal effects of vacuum extraction with forceps or cesarean deliveries. Obstet Gynecol 1981;57(5):571-7.

Greulich B, Paine LL, McClain C, et al. Twelve years and more than 30,000 nurse-midwife-attended births: the Los Angeles County + University of Southern California women's hospital birth center experience. J Nurse Midwifery 1994;39(4):185-96.

Gries DM, Phyall G, Barfield WD. Evaluation of an early discharge program for infants after childbirth in a military population. Mil Med 2000;165(8):616-21.

Griffith P. Midwives' attitudes: promoting normal birth. Aust J Adv Nur 1988;5(3):33-9.

Griffiths ML. The scarred uterus and subsequent pregnancy and delivery. S Afr Med J 1982;62(1):5.

Grobman WA, Peaceman AM, Socol ML. Cost-effectiveness of elective cesarean delivery after one prior low transverse cesarean. Obstet Gynecol 2000;95(5):745-51.

Gross PA, Greenfield S, Cretin S, et al. Optimal methods for guideline implementation: conclusions from Leeds Castle meeting. Med Care 2001;39(8 Suppl 2):II85-92.

Groutz A, Gordon D, Wolman I, et al. Persistent postpartum urinary retention in contemporary obstetric practice. Definition, prevalence and clinical implications. J Reprod Med 2001;46(1):44-8.

Grubb DK, Kjos SL, Paul RH. Latent labor with

an unknown uterine scar. Obstet Gynecol 1996;88(3):351-5.

Grullon KE, Grimes DA. The safety of early postpartum discharge: a review and critique. Obstet Gynecol 1997;90(5):860-5.

Grzybowski SC, Cadesky AS, Hogg WE. Rural obstetrics: a 5-year prospective study of the outcomes of all pregnancies in a remote northern community. Can Med Assoc J 1991;144(8):987-94.

Gudgeon CW. Uterine rupture and scar dehiscence. Anaesth Intensive Care 1997;25(4):434.

Guerdan BR, McKenna JP, Wright JC. Vaginal birth after cesarean section in a community hospital: a family practice residency experience. J Am Board Fam Pract 1989;2(3):169-71.

Guerresi E, Gori G, Beccari A, et al. Influence of spasmolytic treatment and amniotomy on delivery times: A factorial clinical trial. Clin Ther 1981;3(5):382-388.

Guillemette J, Fraser WD. Differences between obstetricians in caesarean section rates and the management of labour. Br J Obstet Gynaecol 1992;99(2):105-8.

Guldholt I, Espersen T. Maternal febrile morbidity after cesarean section. Acta Obstet Gynecol Scand 1987;66(8):675-9.

Gupta U, Ganesh K. Emergency hysterectomy in obstetrics: review of 15 years. Asia Oceania J Obstet Gynaecol 1994;20(1):1-5.

Guyer B, Strobino DM, Ventura SJ, et al. Annual summary of vital statistics-1994. Pediatrics 1995;96(6):1029-39.

Haas JS, Udvarhelyi S, Epstein AM. The effect of health coverage for uninsured pregnant women on maternal health and the use of cesarean section. JAMA 1993;270(1):61-4.

Hadley CB, Mennuti MT, Gabbe SG. An evaluation of the relative risks of a trial of labor versus elective repeat cesarean section. Am J Perinatol 1986;3(2):107-14.

Hadley J, Hoffman J, Feder J. Relationships between health insurance coverage and selected health and hospital use characteristics of newborns and pregnant women. Washington: Center for Health Policy Studies, George Washington University; 1989.

Hage ML, Helms MJ, Dudley A, et al. Acute childbirth morbidity: its measurement using hospital charges. Am J Obstet Gynecol 1992;166(6 Pt 1):1853-9; discussion 1859-62.

Haines CJ, Rogers MS, Leung DH. Neonatal outcome and its relationship with maternal age. Aust N Z J Obstet Gynaecol 1991;31(3):209-12.

Haire DB, Elsberry CC. Maternity care and outcomes in a high-risk service: the North Central Bronx Hospital experience. Birth 1991;18(1):33-7.

Hale RW. Trial of labor compared with an elective second cesarean section. N Engl J Med 1997;336(9):659.

Hales KA, Morgan MA, Thurnau GR. Influence of labor and route of delivery on the frequency of respiratory morbidity in term neonates. Int J Gynaecol Obster 1993;43(1):35-40.

Hall MH. Mortality associated with elective caesarean section. BMJ 1994;308(6943):1572.

Halliday HL. Elective delivery at "term": implications for the newborn. Acta Paediatr 1999;88(11):1180-1.

Halperin ME, Moore DC, Hannah WJ. Classical versus low-segment transverse incision for preterm caesarean section: maternal complications and outcome of subsequent pregnancies. Br J Obstet Gynaecol 1988;95(10):990-6.

Hamilton EF, Bujold E, McNamara H, et al. Dystocia among women with symptomatic uterine rupture. Am J Obstet Gynecol 2001;184(4):620-4.

Hamrick-Turner JE, Cranston PE, Lantrip BS. Gravid uterine dehiscence: MR findings. Abdom Imaging 1995;20(5):486-8.

Handa VL, Harris TA, Ostergard DR. Protecting the pelvic floor: obstetric management to prevent incontinence and pelvic organ prolapse. Obstet Gynecol 1996;88(3):470-8.

Haney EI, Reiter JA, MacGregor SN, et al. Optional vaginal delivery rate. An informative indicator of intrapartum care. J Reprod Med 1999;44(10):842-8.

Hangsleben KL, Taylor MA, Lynn NM. VBAC program in a nurse-midwifery service. Five years of experience. J Nurse Midwifery 1989;34(4):179-84.

Hankins GD, Hammond TL, Snyder RR, et al. Transverse lie. Am J Perinatol 1990;7(1):66-70.

Hanley ML, Smulian JC, Lake MF, et al. Analysis of repeat cesarean delivery indications: implications of heterogeneity. Am J Obstet Gynecol 1996;175(4 Pt 1):883-8.

Hannah ME, Hannah WJ, Hodnett ED, et al. Outcomes at 3 months after planned cesarean vs planned vaginal delivery for breech presentation at term: the international randomized Term Breech Trial. JAMA 2002;287(14):1822-31.

Hannah ME, Ohlsson A, Farine D, et al. Induction of labor compared with expectant management for prelabor rupture of the membranes at term. N Engl J Med 1996;334(16):1005-1010.

Hansell RS, McMurray KB, Huey GR. Vaginal birth after two or more cesarean sections: a five-year experience. Birth 1990;17(3):146-50; discussion 150-1.

Hansen JP. Older maternal age and pregnancy outcome: a review of the literature. Obstetrical & Gynecological Survey 1986;41(11):726-42.

Harlass FE, Duff P. The duration of labor in primiparas undergoing vaginal birth after cesarean delivery. Obstet Gynecol 1990;75(1):45-7.

Harrington LC, Miller DA, McClain CJ, et al. Vaginal birth after cesarean in a hospital-based birth center staffed by certified nurse-midwives. J Nurse Midwifery 1997;42(4):304-7.

Harris LH. Counselling women about choice. Best Prac Res Clin Obstet Gynaecol 2001;15(1):93-107.

Harris RP, Helfand M, Woolf SH, et al. Methods of the third U.S. Preventive Services Task Force. Am J Prev Med 2001;20(3S):21-35.

Hart GL, Dobie SA, Baldwin LM, et al. Rural and urban differences in physician resource use for low-risk obstetrics. Seattle: WAMI RHRC 1995;42.

Hart LG, Dobie SA, Baldwin LM, et al. Rural and urban differences in physician resource use for low-risk obstetrics. Health Serv Res 1996;31(4):429-52.

Hart MA, Foster SN. Couples' attitudes toward childbirth participation: relationship to evaluation of labor and delivery. J Perinat Neonat Nurs 1997;11(1):10-20.

Harwood MI. Are there adverse maternal and neonatal outcomes associated with induction of labor when there is no well-accepted indication? J Fam Pract 2001;50(2):106.

Haupt BJ. Deliveries in short-stay hospitals: United States, 1980. Adv Data 1982(83):1-11.

Hawe JA, Olah KS. Posterior uterine rupture in a patient with a lower segment caesarean section scar complicating prostaglandin induction of labour. Br J Obstet Gynaecol 1997;104(7):857-8.

Haynes de Regt R, Minkoff HL, Feldman J. Relations of private or clinic care to the cesarean birth rate. N Engl J Med 1986;315:619-24.

Heddleston LN, Watson WJ. Vaginal birth after cesarean section in a small hospital. Mil Med 1991;156(5):239-40.

Heidrick WP. Rupture of the gravid uterus--lessons and questions. Nebr Med J 1983;68(7):197-8.

Heilbrunn JZ, Park E. Variations in the Use of Cesarean Section: Literature Review. Santa Monica, Calif.: RAND; 1992.

Helfand M, Marton K, Ueland K. Factors involved in the interpretation of fetal monitor tracings. Am J Obstet Gynecol 1985;151(6):737-44.

Hemminki E. Obstetric practice in Finland, 1950-1980. Changes in technology and its relation to health. Med Care 1983;21(12):1131-43.

Hemminki E. Pregnancy and birth after cesarean section: a survey based on the Swedish birth

register. Birth 1987;14(1):12-7.

Hemminki E. Long term maternal health effects of caesarean section. J Epidemiol Community Health 1991;45(1):24-8.

Hemminki E, Kojo-Austin H, Malin M, et al. Variation in obstetric interventions by midwife. Scand J Caring Sci 1992;6(2):81-6.

Henderson J, McCandlish R, Kumiega L, et al. Systematic review of economic aspects of alternative modes of delivery. BJOG 2001;108(2):149-57.

Henry OA, Gregory KD, Hobel CJ, et al. Using ICD-9 codes to identify indications for primary and repeat cesarean sections: agreement with clinical records. Am J Public Health 1995;85(8 Pt 1):1143-6.

Heres MH, Pel M, Elferink-Stinkens PM, et al. The Dutch obstetric intervention study--variations in practice patterns. Int J Gynaecol Obster 1995;50(2):145-50.

Heriot MJ. Fetal rights versus the female body: contested domains. Med Anthropol Q 1996;10(2):176-94.

Heritage C, Cunningham M. Association of elective repeat cesarean delivery and persistent pulmonary hypertension of the newborn. Am J Obstet Gynecol 1983;152:627.

Hibbard JH, Weeks EC. Does the dissemination of comparative data on physician fees affect consumer use of services? Med Care 1989;27(12):1167-74.

Hibbard JU, Ismail MA, Wang Y, et al. Failed vaginal birth after a cesarean section: how risky is it? I. Maternal morbidity. Am J Obstet Gynecol 2001;184(7):1365-71; discussion 1371-3.

Hickson GB, Altemeier WA, Perrin JM. Physician reimbursement by salary or fee-for-service: effect on physician practice behavior in a randomized prospective study. Pediatrics 1986;80:344-50.

Hill DA, Chez RA, Quinlan J, et al. Uterine rupture and dehiscence associated with intravaginal misoprostol cervical ripening. J Reprod Med 2000;45(10):823-6.

Hill DJ, Beischer NA. Hysterectomy in obstetric practice. Aust N Z J Obstet Gynaecol 1980;20(3):151-3.

Hillan EM. Postoperative morbidity following Caesarean delivery. J Adv Nurs 1995;22(6):1035-42.

Hillman AL. Health maintenance organizations, financial incentives, and physicians' judgments. Ann Intern Med 1990;112(12):891-3.

Hoffmann RAM, Anthony J, Fawcus S. Oral misoprostol vs. placebo in the management of prelabor rupture of membranes at term. Int J Gynecol Obstet 2001;72(3):215-221.

Hofmeyr GJ, Alfirevic Z, Matonhodze B, et al. Titrated oral misoprostol solution for induction of labour: A multi-centre, randomised trial. Br J Obstet Gynaecol 2001;108(9):952-959.

Hofmeyr GJ, Kulier R. Operative versus conservative management for 'fetal distress' in labour. Cochrane Database of Systematic Reviews 2000(2):CD001065.

Holden D, Vere M, Manyonda I. Vesico-uterine fistula occurring in a woman with a previous caesarean section and two subsequent normal vaginal deliveries. Br J Obstet Gynaecol 1994;101(4):354-6.

Holland JG, Dupre AR, Blake PG, et al. Trial of labor after cesarean delivery: experience in the non-university level II regional hospital setting. Obstet Gynecol 1992;79(6):936-9.

Holmes P, Oppenheimer LW, Wen SW. The relationship between cervical dilatation at initial presentation in labour and subsequent intervention. BJOG 2001;108(11):1120-4.

Holt VL, Mueller BA. Attempt and success rates for vaginal birth after caesarean section in relation to complications of the previous pregnancy. Paediatr Perinat Epidemiol 1997;11(Suppl 1):63-72.

Hook B, Kiwi R, Amini SB, et al. Neonatal morbidity after elective repeat cesarean section and trial of labor. Pediatrics 1997;100(3 Pt 1):348-53.

Horenstein JM, Eglinton GS, Tahilramaney MP, et al. Oxytocin use during a trial of labor in

patients with previous cesarean section. J Reprod Med Obstet Gynecol 1984;29(1):26-30.

Horenstein JM, Phelan JP. Previous cesarean section: the risks and benefits of oxytocin usage in a trial of labor. Am J Obstet Gynecol 1985;151(5):564-9.

Hornbrook MC, Goldfarb MG. Patterns of obstetrical care in hospitals. Med Care 1981;19(1):55-67.

Horowitz BJ, Edelstein SW, Lippman L. Once a cesarean, always a cesarean. Obstet Gynecol Survey 1981;36:592.

Hoskins IA, Gomez JL. Correlation between maximum cervical dilatation at cesarean delivery and subsequent vaginal birth after cesarean delivery. Obstet Gynecol 1997;89(4):591-3.

Hourvitz A, Alcalay M, Korach J, et al. A prospective study of high- versus low-dose oxytocin for induction of labor. Acta Obstet Gynecol Scand 1996;75(7):636-41.

Hsiao WC, Braun P, Dunn D, et al. Results and policy implications of the resource-based relative-value study. N Engl J Med 1988;319(13):881-8.

Hsu CD, Chen S, Feng TI, et al. Rupture of uterine scar with extensive maternal bladder laceration after cocaine abuse. Am J Obstet Gynecol 1992;167(1):129-30.

Hueston WJ. Obstetric referral in family practice. Factors Affecting Cesarean Section (FACS) Study Group. J Fam Pract 1994;38(4):368-72.

Hueston WJ, Applegate JA, Mansfield CJ, et al. Practice variations between family physicians and obstetricians in the management of low-risk pregnancies. J Fam Pract 1995;40(4):345-51.

Hueston WJ, Rudy M. A comparison of labor and delivery management between nurse midwives and family physicians. J Fam Pract 1993;37(5):449-54.

Hueston WJ, Rudy M. Factors predicting elective repeat cesarean delivery. Obstet Gynecol 1994;83(5 Pt 1):741-4.

Hueston WJ, Rudy M. Differences in labor and delivery experience in family physician- and obstetrician-supervised teaching services. Fam Med 1995;27(3):182-7.

Hughes EG, Kelly AJ, Kavanagh J. Dinoprostone vaginal insert for cervical ripening and labor induction: a meta-analysis. Obstet Gynecol 2001;97(5 Pt 2):847-55.

Hundley VA, Milne JM, Glazener CM, et al. Satisfaction and the three C's: continuity, choice and control. Women's views from a randomised controlled trial of midwife-led care. Br J Obstet Gynaecol 1997;104(11):1273-80.

Hunter IW, Cato E, Ritchie JW. Induction of labor using high-dose or low-dose prostaglandin vaginal pessaries. Obstet Gynecol 1984;63(3):418-20.

Hurst M, Summey PS. Childbirth and social class: the case of cesarean delivery. Soc Sci Med 1984;18(8):621-31.

Hutchins V, Kessel SS, Placek PJ. Trends in maternal and infant health factors associated with low infant birth weight, United States, 1972 and 1980. Public Health Rep 1984;99(2):162-72.

Iglesias S, Burn R, Saunders LD. Reducing the cesarean section rate in a rural community hospital. Can Med Assoc J 1991;145(11):1459-64.

Iloabachie GC, Njoku O. Vesico-uterine fistula. Br J Urol 1985;57(4):438-9.

Impey L, O'Herlihy C. First delivery after cesarean delivery for strictly defined cephalopelvic disproportion. Obstet Gynecol 1998;92(5):799-803.

Ingemarsson E, Ingemarsson I, Svenningsen NW. Impact of routine fetal monitoring during labor on fetal outcome with long-term follow-up. Am J Obstet Gynecol 1981;141(1):29-38.

Innes G, Rosen P. An unusual cause of abdominal pain and shock in pregnancy: case report and review of the literature. J Emerg Med 1985;2(5):361-6.

Irion O, Hirsbrunner Almagbaly P, Morabia A. Planned vaginal delivery versus elective caesarean section: a study of 705 singleton term breech presentations. Br J Obstet Gynaecol

1998;105(7):710-7.

Ito M, Nawa T, Mikamo H, et al. Lower segment uterine rupture related to early pregnancy by in vitro fertilization and embryo transfer after a previous cesarean delivery. J Med 1998;29(1-2):85-91.

Jackson DJ, Lang J, Swartz W, et al. Outcomes from the San Diego Birth Center Study: Examining an alternative model of perinatal service delivery. Abstr Book Assoc Health Serv Res;15:95.

Jackson DJ, Lang JM, Ganiats TG. Epidemiological issues in perinatal outcomes research. Paediatr Perinat Epidemiol 1999;13(4):392-404.

Jackson R. L.A. County learns costly lesson on mandated labor guidelines. Qual Lett Healthc Lead 1998;10(4):18-9.

Jacoby A. Women's preferences for and satisfaction with current procedures in childbirth--findings from a national study. Midwifery 1987;3(3):117-24.

Jacoby I, Meyer GS, Haffner W, et al. Modeling the future workforce of obstetrics and gynecology. Obstet Gynecol 1998;92(3):450-6.

Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? Control Clin Trials 1996;17(1):1-12.

Jagani N, Schulman H, Chandra P, et al. The predictability of labor outcome from a comparison of birth weight and x-ray pelvimetry. Am J Obstet Gynecol 1981;139(5):507-11.

Jain M, Pandey S, Pandey LK, et al. Obstetric prospects after caesarean section. J Indian Med Assoc 1987;85(11):324-6.

Jakobi P, Weissman A, Paldi E. Uniform presentation of data concerning cesarean sections. Acta Obstet Gynecol Scand 1987;66(7):657-8.

Jakobi P, Weissman A, Peretz BA, et al. Evaluation of prognostic factors for vaginal delivery after cesarean section. J Reprod Med 1993;38(9):729-33.

Jakobi P, Zimmer EZ, Weissman A. Oxytocin administration and uterine exploration in the management of vaginal birth after cesarean--not routine procedures. Am J Obstet Gynecol 1989;160(6):1538-40.

Jakobi P, Zimmer EZ, Weissman A, et al. Oxytocin administration and uterine exploration in the management of vaginal birth after cesarean - Not routine procedures. Am J Obstet Gynecol 1989;160(6):1538-1540.

Jamelle RN. Outcome of unplanned vaginal deliveries after two previous caesarean sections. J Obstet Gynaecol Res 1996;22(5):431-6.

Janke JR. Breastfeeding duration following cesarean and vaginal births. J Nurse Midwifery 1988;33(4):159-64.

Janowitz B, Higgins JE, Clopton DC, et al. Access to postpartum sterilization in southeast Brazil. Med Care 1982;20(5):526-34.

Janowitz B, Wallace S, Araujo G, et al. Method of payment and the cesarean birth rate in a hospital in northeast Brazil. J Health Polit Policy Law 1984;9(3):515-26.

Janssen PA, Klein MC, Soolsma JH. Differences in institutional cesarean delivery rates-the role of pain management. J Fam Pract 2001;50(3):217-23.

Jarrell MA, Ashmead GG, Mann LI. Vaginal delivery after cesarean section: a five-year study. Obstet Gynecol 1985;65(5):628-32.

Jarvelin MR, Hartikainen-Sorri AL, Rantakallio P. Labour induction policy in hospitals of different levels of specialisation. Br J Obstet Gynaecol 1993;100(4):310-5.

Johanson R, Burr R, Leighton N, et al. Informed choice? Evidence of the persuasive power of professionals. J Public Health Med 2000;22(3):439-40.

Johanson RB, Heycock E, Carter J, et al. Maternal and child health after assisted vaginal delivery: five-year follow up of a randomised controlled study comparing forceps and ventouse. Br J Obstet Gynaecol 1999;106(6):544-9.

Johnson C, Oriol N. The role of epidural

anesthesia in trial of labor. Reg Anesth 1990;15(6):304-8.

Johnson C, Oriol N, Flood K. Trial of labor: a study of 110 patients. J Clin Anesth 1991;3(3):216-8; discussion 214-5.

Johnson KC, Gaskin IM. Vaginal delivery after caesarean section. Safety of single-layer suturing in caesarean sections must be proved. BMJ 2001;323(7324):1307-8.

Johnson N, Ansell D. Variation in caesarean and instrumental delivery rates in New Zealand hospitals. Aust N Z J Obstet Gynaecol 1995;35(1):6-11.

Johnson SR, Elkins TE, Strong C, et al. Obstetric decision-making: responses to patients who request cesarean delivery. Obstet Gynecol 1986;67(6):847-50.

Jolly J, Walker J, Bhabra K. Subsequent obstetric performance related to primary mode of delivery. Br J Obstet Gynaecol 1999;106(3):227-32.

Jones L, LoGerfo J, Shy K, et al. StORQS: Washington's Statewide Obstetrical Review and Quality System: overview and provider evaluation. Qrb. Quality Review Bulletin 1993;19(4):110-8.

Jones RO, Nagashima AW, Hartnett-Goodman MM, et al. Rupture of low transverse cesarean scars during trial of labor. Obstet Gynecol 1991;77(6):815-7.

Jongen VH, Halfwerk MG, Brouwer WK. Vaginal delivery after previous caesarean section for failure of second stage of labour. Br J Obstet Gynaecol 1998;105(10):1079-81.

Joseph GF, Stedman CM, Robichaux AG. Vaginal birth after cesarean section: the impact of patient resistance to a trial of labor. Am J Obstet Gynecol 1991;164(6 Pt 1):1441-4; discussion 1444-7.

Kafkas S, Taner CE. Ruptured uterus. Int J Gynaecol Obster 1991;34(1):41-4.

Kahn K, Fiske M, DiMatteo R, et al. Mother's thoughts about cesarean as compared with vaginal deliveries and the effect of those thoughts on method of delivery. AHSR FSHR Annu Meet Abstr Book;11:23.

Kaplan B, Rabinerson D, Harel L, et al. Multiple repeat cesarean sections. Int J Gynaecol Obster 1996;55(2):173-4.

Kaplan B, Royburt M, Peled Y, et al. Routine revision of uterine scar after prior cesarean section. Acta Obstet Gynecol Scand 1994;73(6):473-475.

Kattan SA. Maternal urological injuries associated with vaginal deliveries: change of pattern. Int Urol Nephrol 1997;29(2):155-61.

Kazandjian VA, Thomson RG, Law WR, et al. Do performance indicators make a difference? Jt Comm J Qual Improv 1996;22(7):482-91.

Keeler EB, Brodie M. Economic incentives in the choice between vaginal delivery and cesarean section. Milbank Q 1993;71(3):365-404.

Keeler EB, Fok T. Equalizing physician fees had little effect on cesarean rates. Med Care Res Rev 1996;53(4):465-71.

Keirse M, De Koning Gans HJ. Randomized comparison of the effects of endocervical and vaginal prostaglandin E2 gel in women with various degrees of cervical ripeness. Am J Obstet Gynecol 1995;173(6):1859-1864.

Keirse MJ. Amniotomy or oxytocin for induction of labor. Re-analysis of a randomized controlled trial. Acta Obstet Gynecol Scand 1988;67(8):731-5.

Keirse MJ. Prostaglandins in preinduction cervical ripening. Meta-analysis of worldwide clinical experience. J Reprod Med 1993;38(1 Suppl):89-100.

Kennedy CM, Peleg D, Syrop C, et al. Antenatal diagnosis of vesicouterine fistula. Obstet Gynecol 1999;94(5 Pt 2):808-9.

Kennedy EK. VBAC trend put in reverse. Hosp Peer Rev 1997;22(9):129-31.

Kennedy JH, Stewart P, Barlow DH, et al. Induction of labour: a comparison of a single prostaglandin E2 vaginal tablet with amniotomy and intravenous oxytocin. Br J Obstet Gynaecol 1982;89(9):704-7.

Kennell J, Klaus M, McGrath S, et al. Continuous emotional support during labor in a US hospital. A randomized controlled trial. JAMA 1991;265(17):2197-201.

Khan KS, Rizvi A. The partograph in the management of labor following cesarean section. Int J Gynaecol Obster 1995;50(2):151-7.

Khan KS, Rizvi A, Rizvi JH. Risk of uterine rupture after the partographic 'alert' line is crossed--an additional dimension in the quest towards safe motherhood in labour following caesarean section. JPMA J Pak Med Assoc 1996;46(6):120-2.

Khanna S. Inversion of uterus through an upper segment scar. Br J Obstet Gynaecol 1999;106(11):1225-6.

Kieback DG, Zahradnik HP, Quaas L, et al. Clinical evaluation of endocervical prostaglandin E2-triacetin-gel for preinduction cervical softening in pregnant women at term. Prostaglandins 1986;32(1):81-5.

Kildea S. Trial of scar--team midwifery makes a difference. J Aust Coll Midwives 1996;9(3):21-2.

Kilpatrick SJ, Safford KL. Repeat external cephalic version. Is it worth the effort? J Reprod Med 1995;40(11):775-8.

Kimball FA, Ruppel PL, Noah ML, et al. The effect of endocervical PGE2-gel (Prepidil) gel on plasma levels of 13,14-dihydro-15-keto-PGE2 (PGEM) in women at term. Prostaglandins 1986;32(4):527-37.

Kindig M, Cardwell M, Lee T. Delayed postpartum uterine dehiscence. A case report. J Reprod Med 1998;43(7):591-2.

King DE, Lahiri K. Socioeconomic factors and the odds of vaginal birth after cesarean delivery. JAMA 1994;272(7):524-9.

Kirk EP, Doyle KA, Leigh J, et al. Vaginal birth after cesarean or repeat cesarean section: medical risks or social realities? Am J Obstet Gynecol 1990;162(6):1398-403; discussion 403-5.

Kishor T, Singh C, Barman SD, et al. Study of vaginal delivery in patients with one previous lower segment caesarean section. Aust N Z J Obstet Gynaecol 1986;26(4):245-8.

Kitzinger S. Sheila Kitzinger's letter from Europe: awake, aware --and action! Birth 2001;28(3):210-2.

Kizer KW, Ellis A. C-section rate related to payment source. Am J Public Health 1988;78(1):96-7.

Klein M, Rosen A, Beck A. Diagnostic potential of cardiotocography (CTG) for silent uterine rupture. Acta Obstet Gynecol Scand 1989;68(7):653-6.

Klein MC, Gauthier RJ, Robbins JM, et al. Relationship of episiotomy to perineal trauma and morbidity, sexual dysfunction, and pelvic floor relaxation. Am J Obstet Gynecol 1994;171(3):591-8.

Kline J, Arias F. Analysis of factors determining the selection of repeated cesarean section or trial of labor in patients with histories of prior cesarean delivery. J Reprod Med 1993;38(4):289-92.

Knight R, Fynes M, Permezel M. Faecal incontinence after vaginal delivery: a review. Aust N Z J Obstet Gynaecol 2001;41(2):161-6.

Kolder VE, Gallagher J, Parsons MT. Court-ordered obstetrical interventions. N Engl J Med 1987;316(19):1192-6.

Koska MT. Survey: lower cesarean rate reveal cost-quality tradeoffs. Trustee 1989;42(4):16.

Kotagal UR, Atherton HD, Eshett R, et al. Safety of early discharge for Medicaid newborns. JAMA 1999;282(12):1150-6.

Kozak LJ. Surgical and nonsurgical procedures associated with hospital delivery in the United States: 1980-1987. Birth 1989;16(4):209-13.

Kramer RL, Gilson GJ, Morrison DS, et al. A randomized trial of misoprostol and oxytocin for induction of labor: safety and efficacy. Obstet Gynecol 1997;89(3):387-91.

Krantz ME, Wennergren M, Bengtson LG, et al. Epidemiological analysis of the increased risk of disturbed neonatal respiratory adaptation after caesarean section. Acta Paediatr Scand 1986;75(5):832-9.

Kraus JF, Bulterys M, Greenland S. A nested case-control study of oxytocin and sudden infant death syndrome. Am J Obstet Gynecol 1990;162(2):604-605.

Krieger N. Analyzing socioeconomic and racial/ethnic patterns in health and health care. Am J Public Health 1993;83(8):1086-7.

Krikke EH, Bell NR. Relation of family physician or specialist care to obstetric interventions and outcomes in patients at low risk: a western Canadian cohort study. Can Med Assoc J 1989;140(6):637-43.

Krishnamurthy S, Fairlie F, Cameron AD, et al. The role of postnatal x-ray pelvimetry after caesarean section in the management of subsequent delivery. Br J Obstet Gynaecol 1991;98(7):716-8.

Kumar S, Maouris P. Induction of labour for trial of vaginal birth after caesarean section in a remote district hospital. Aust N Z J Obstet Gynaecol 1996;36(4):417-20.

Kwon JS, Davies GAL, Mackenzie VP. A comparison of oral and vaginal misoprostol for induction of labour at term: A randomised trial. Br J Obstet Gynaecol 2001;108(1):23-26.

Lagrew DC, Jr., Adashek JA. Lowering the cesarean section rate in a private hospital: comparison of individual physicians' rates, risk factors, and outcomes. Am J Obstet Gynecol 1998;178(6):1207-14.

Lagrew DC, Morgan MA. Decreasing the cesarean section rate in a private hospital: success without mandated clinical changes. Am J Obstet Gynecol 1996;174(1 Pt 1):184-91.

Lai SF, Sidek S. Delivery after a lower segment caesarean section. Singapore Med J 1993;34(1):62-6.

Laken MP, Ager J. Use of the postpartum survey for program planing and ongoing evaluation. Abstr Book Assoc Health Serv Res;14:347-8.

Land R, Parry E, Rane A, et al. Personal preferences of obstetricians towards childbirth. Aust N Z J Obstet Gynaecol 2001;41(3):249-52.

Lange AP, Secher NJ, Nielsen FH, et al. Stimulation of labor in cases of premature rupture of the membranes at or near term. A consecutive randomized study of prostaglandin E2-tablets and intravenous oxytocin. Acta Obstet Gynecol Scand 1981;60(2):207-10.

Lantinen AJ. Labor following previous cesarean section. Am J Obstet Gynecol 1980;137(4):517.

Lao TT, Chin RK, Leung BF. Is X-ray pelvimetry useful in a trial of labour after caesarean section? Eur J Obstet Gynecol Reprod Biol 1987;24(4):277-83.

Lao TT, Leung BF. Rupture of the gravid uterus. Eur J Obstet Gynecol Reprod Biol 1987;25(3):175-80.

Lao TT, Leung BFH. Labor induction for planned vaginal delivery in patients with previous cesarean section. Acta Obstet Gynecol Scand 1987;66(5):413-416.

Lau TK, Chan F. Unknown uterine scars, unknown risks. Aust N Z J Obstet Gynaecol 1994;34(2):216-7.

Lau TK, MRCOG, Wong SH, et al. A study of patient's acceptance towards vaginal birth after cesarean section. Aust N Z J Obstet Gynaecol 1996;36(2):155-158.

Laube DW, Zlatnik FJ, Pitkin RM. Preinduction cervical ripening with prostaglandin E2 intracervical gel. Obstet Gynecol 1986;68(1):54-7.

Laufer A, Hodenius V, Friedman L, et al. Vaginal birth after cesarean section. Nurse-midwifery management. J Nurse Midwifery 1987;32(1):41-7.

Lavender T, Wallymahmed AH, Walkinshaw SA. Managing labor using partograms with different action lines: a prospective study of women's views. Birth 1999;26(2):89-96.

Lavin JP, Stephens RJ, Miodovnik M, et al. Vaginal delivery in patients with a prior cesarean section. Obstet Gynecol 1982;59(2):135-48.

Law YY, Lam KY. A randomized controlled trial comparing midwife-managed care and obstetrician-managed care for women assessed to be at low risk in the initial intrapartum period. J Obstet Gynaecol Res 1999;25(2):107-12.

Lawler PE, Bulfin MJ, Lawler FC, et al. A review of vainal delivery follwoing cesarean section, from private practice. Am J Obstet Gynecol 1956;72:252.

Le Coutour X, Infante-Rivard C, Danzon A. [Regionalization of health care and obstetric practice]. Rev Epidemiol Sante Publique 1990;38(3):211-20.

Leaphart WL, Meyer MC, Capeless EL. Labor induction with a prenatal diagnosis of fetal macrosomia. J Matern Fetal Med 1997;6(2):99-102.

Learman LA, Evertson LR, Shiboski S. Predictors of repeat cesarean delivery after trial of labor: do any exist? J Am Coll Surg 1996;182(3):257-62.

Lee AH, Xiao J, Vemuri SR, et al. A discordancy test approach to identify outliers of length of hospital stay. Stat Med 1998;17(19):2199-206.

Lee JY, Cass AS. Spontaneous bladder and uterine rupture with attempted vaginal delivery after cesarean section. J Urol 1992;147(3):691-2.

Lehmann DK, Chism J. Pregnancy outcome in medically complicated and uncomplicated patients aged 40 years or older. Am J Obstet Gynecol 1987;157(3):738-42.

Lelaidier C, Baton C, Benifla JL, et al. Mifepristone for labour induction after previous caesarean section. Br J Obstet Gynaecol 1994;101(6):501-3.

Lenke R, Osterkamp T. A prolapsed umbilical cord into the abdominal cavity in a woman with a previous cesarean section. Am J Obstet Gynecol 1980;138(8):1224-5.

Lescale KB, Inglis SR, Eddleman KA, et al. Conflicts between physicians and patients in non-elective cesarean delivery: incidence and the adequacy of informed consent. Am J Perinatol 1996;13(3):171-6.

Leung AS, Farmer RM, Leung EK, et al. Risk factors associated with uterine rupture during trial of labor after cesarean delivery: A case-control study. Am J Obstet Gynecol 1993;168(5):1358-1363.

Leung AS, Leung EK, Paul RH. Uterine rupture after previous cesarean delivery: maternal and fetal consequences. Am J Obstet Gynecol 1993;169(4):945-50.

Leung KM, Elashoff RM, Rees KS, et al. Hospital- and patient-related characteristics determining maternity length of stay: a hierarchical linear model approach. Am J Public Health 1998;88(3):377-81.

Leung TY, Lau TK, Lo KW, et al. A survey of pregnant women's attitude towards breech delivery and external cephalic version. Aust N Z J Obstet Gynaecol 2000;40(3):253-9.

Leuthard JL, Davison M, Lorenz D. Obstetrical practice survey report. Healthy Futures Program, May 2, 1991. J Oklahoma State Med Assoc 1991;84(12):607-11.

Leveno KJ, Cunningham FG, Nelson S, et al. A prospective comparison of selective and universal electronic fetal monitoring in 34,995 pregnancies. N Engl J Med 1986;315(10):615-9.

Leveno KJ, Cunningham FG, Pritchard JA. Cesarean section: the House of Horne revisited. Am J Obstet Gynecol 1989;160(1):78-9.

Leyland A. Socioeconomic and racial differences in obstetric procedures. [letter; comment]. Am J Public Health 1993;83(8):1178-9.

Li WJ, Li ZL, Ha KW. Effect of hyaluronidase on cervical ripening. Chin Med J 1994;107(7):552-3.

Lieberman E. Risk factors for uterine rupture dining a trial of labor after cesarean. Clin Obstet Gynecol 2001;44(3):609-621.

Lieberman E. Risk factors for uterine rupture during a trial of labor after cesarean. Clin Obstet Gynecol 2001;44(3):609-621.

Lieberman E, Lang JM, Heffner LJ, et al. Assessing the role of case mix in cesarean delivery rates. Obstet Gynecol 1998;92(1):1-7.

Lilford RJ, van Coeverden de Groot HA, Moore PJ, et al. The relative risks of caesarean section (intrapartum and elective) and vaginal delivery: a detailed analysis to exclude the effects of medical disorders and other acute pre-existing physiological disturbances. Br J Obstet Gynaecol

1990;97(10):883-92.

Lim R. A window to Bali. Midwifery Today Childbirth Educ 1993(28):22-3.

Lind B, Hoel TM. Alleviation of labor pain in Norway. An interview investigation in 1969 and 1986. Acta Obstet Gynecol Scand 1989;68(2):125-9.

Lipscomb KR, Gregory K, Shaw K. The outcome of macrosomic infants weighing at least 4500 grams: Los Angeles County + University of Southern California experience. Obstet Gynecol 1995;85(4):558-64.

Lipson JG. Repeat cesarean births. Social and psychological issues. JOGN Nurs 1984;13(3):157-62.

Lledo R, Rodriguez T, Trilla A, et al. Perceived quality of care in pregnancy assessment before and after delivery. Eur J Obstet Gynecol Reprod Biol 2000;88(1):35-42.

Localio AR, Lawthers AG, Bengtson JM, et al. Relationship between malpractice claims and cesarean delivery. JAMA 1993;269(3):366-73.

Lomas J, Anderson G, Enkin M, et al. The role of evidence in the consensus process. Results from a Canadian consensus exercise. JAMA 1988;259(20):3001-5.

Lomas J, Anderson GM, Domnick-Pierre K, et al. Do practice guidelines guide practice? The effect of a consensus statement on the practice of physicians. N Engl J Med 1989;321(19):1306-11.

Lomas J, Enkin M. Variations in operative delivery rates. Effictive care in pregnancy and childbirth, In: Chalmers I, Enkin M, Keirse MJNC, editors. 1989;2 (Childbirth):1182-95.

Lomas J, Enkin M, Anderson GM, et al. Opinion leaders vs audit and feedback to implement practice guidelines. Delivery after previous cesarean section. JAMA 1991;265(17):2202-7.

Lonky NM, Worthen N, Ross MG. Prediction of cesarean section scars with ultrasound imaging during pregnancy. J Ultrasound Med 1989;8(1):15-9.

Lopez-Zeno JA, Peaceman AM, Adashek JA, et al. A controlled trial of a program for the active management of labor. N Engl J Med 1992;326(7):450-4.

Lorenz RP, Botti JJ, Chez RA, et al. Variations of biologic activity of low-dose prostaglandin E2 on cervical ripening. Obstet Gynecol 1984;64(1):123-7.

Lovell R. Vaginal delivery after Caesarean section: factors influencing success rates. Aust N Z J Obstet Gynaecol 1996;36(1):4-8.

Lucas VA. Birth: nursing's role in today's choices. RN 1993;56(6):38-44.

Luckas M, Bricker L. Intravenous prostaglandin for induction of labour. Cochrane Database of Systematic Reviews 2000(4):CD002864.

Lumley J. Assessing satisfaction with childbirth. Birth 1985;12:141-145.

Lurie S, Hagay Z, Goldschmit R, et al. Routine previous cesarean scar exploration following successful vaginal delivery. Is it necessary? Eur J Obstet Gynecol Reprod Biol 1992;45(3):185-6.

Lydon-Rochelle M, Holt VL, Easterling TR, et al. Cesarean delivery and postpartum mortality among primiparas in Washington State, 1987-1996(1). Obstet Gynecol 2001;97(2):169-74.

Lydon-Rochelle M, Holt VL, Easterling TR, et al. Risk of uterine rupture during labor among women with a prior cesarean delivery. N Engl J Med 2001;345(1):3-8.

Lydon-Rochelle M, Holt VL, Martin DP, et al. Association between method of delivery and maternal rehospitalization. JAMA 2000;283(18):2411-6.

Lydon-Rochelle MT, Holt VL, Martin D. The association between method of delivery and maternal rehospitalization among primiparous women: Washington State 1987-1996. Abstr Book Assoc Health Serv Res;16:116.

Lydon-Rochelle MT, Holt VL, Martin DP. Delivery method and self-reported postpartum general health status among primiparous women. Paediatr Perinat Epidemiol 2001;15(3):232-40.

Lynch JC, Pardy JP. Uterine rupture and scar dehiscence. A five-year survey. Anaesth

Intensive Care 1996;24(6):699-704.

MacDonald D. Previous obstetrical or gynaecological surgery. Clin Obstet Gynaecol 1982;9(1):147-69.

Macfarlane A. At last--maternity statistics for England. BMJ 1998;316(7131):566-7.

Macfarlane A, Chamberlain G. What is happening to caesarean section rates? Lancet 1993;342(8878):1005-6.

MacKenzie IZ, Bradley S, Embrey MP. Vaginal prostaglandins and labour induction for patients previously delivered by caesarean section. Br J Obstet Gynaecol 1984;91(1):7-10.

MacLennan AH, Chan FY, Eckert K. The safety of vaginal prostaglandin F2alpha for the stimulation of labour. Aust N Z J Obstet Gynaecol 1994;34(2):154-158.

MacLennan AH, Green RC. A double blind dose trial of intravaginal prostaglandin F2 alpha for cervical ripening and the induction of labour. Aust N Z J Obstet Gynaecol 1980;20(2):80-3.

MacLennan AH, Green RC, Bryant-Greenwood GD, et al. Ripening of the human cervix and induction of labour with purified porcine relaxin. Lancet 1980;1(8162):220-3.

MacLennan AH, Green RC, Bryant-Greenwood GD, et al. Cervical ripening with combinations of vaginal prostaglandin F2-alpha estradiol, and relaxin. Obstet Gynecol 1981;58(5):601-4.

MacLennan AH, Green RC, Grant P, et al. Ripening of the human cervix and induction of labor with intracervical purified porcine relaxin. Obstet Gynecol 1986;68(5):598-601.

Macones GA. The utility of clinical tests of eligibility for a trial of labour following a caesarean section: a decision analysis. Br J Obstet Gynaecol 1999;106(7):642-6.

Macones GA, Hausman N, Edelstein R, et al. Predicting outcomes of trials of labor in women attempting vaginal birth after cesarean delivery: a comparison of multivariate methods with neural networks. Am J Obstet Gynecol 2001;184(3):409-13.

Magann EF, Nolan TE. Pregnancy outcome in an active-duty population. Obstet Gynecol 1991;78(3 Pt 1):391-3.

Maher CF, Cave DG, Haran MV. Caesarean section rate reduced. Aust N Z J Obstet Gynaecol 1994;34(4):389-92.

Mahmood T. The role of radiological pelvimetry in the management of patients who have had a previous caesarean section. J Obstet Gynaecol 1987;8(8):24-28.

Mahmood TA. Maternal height, birthweight, obstetric conjugate and their influence on the management of parturients with a previous cesarean scar. Acta Obstet Gynecol Scand 1989;68(7):595-8.

Makinen JI, Meltomaa SS, Ekblad UU. Hospital stay due to various hysterectomies, caesarean section and normal delivery in Turku University Central Hospital area from 1983 to 1992. Ann Chir Gynaecol Suppl 1994;208:54-7.

Maouris P. Successful vaginal delivery after caesarean scar rupture: a case report. Clin Exp Obstet Gynecol 1989;16(1):1-2.

Marchesoni D, Franco F, Mozzanega B, et al. Vaginal delivery after previous cesarean section. Clin Exp Obstet Gynecol 1984;11(4):150-1.

Marcus S, Cheng E, Goff B. Extrauterine pregnancy resulting from early uterine rupture. Obstet Gynecol 1999;94(5 Pt 2):804-5.

Mardon RE, Shafer KD, Hinton SA. Cesarean deliveries in Florida: current trends and related factors. J Fla Med Assoc 1997;84(5):310-5.

Marieskind HI. Cesarean section in the United States: has it changed since 1979? Birth 1989;16(4):196-202.

Marta MR. Current topics in obstetrical risk management--Part I. J Healthc Qual 1994;16(5):6-9, quiz 9, 19.

Marta MR. Current topics in obstetrical risk management--Part II. J Healthc Qual 1994;16(6):6-9; quiz 16, 48.

Martin JA, Hamilton BE, Ventura SJ, et al. Births: final data for 2000. Natl Vital Stat Rep 2002;50(5):1-101.

Martin JA, Smith BL, Mathews TJ, et al. Births and deaths: preliminary data for 1998. Natl Vital Stat Rep 1999;47(25):1-45.

Martin JN, Harris BA, Huddleston JF, et al. Vaginal delivery following previous cesarean birth. Am J Obstet Gynecol 1983;146(3):255-63.

Martin JN, Jr., Morrison JC, Wiser WL. Vaginal birth after cesarean section: the demise of routine repeat abdominal delivery. Obstet Gynecol Clin North Am 1988;15(4):719-36.

Maslow AS, Sweeny AL. Elective induction of labor as a risk factor for cesarean delivery among low-risk women at term. Obstet Gynecol 2000;95(6 Pt 1):917-22.

Mathews J. Placenta praevia percreta with bladder penetration. Med J Aust 1983;1(4):173-4.

Mathews TJ. Trends in stimulation and induction of labor 1989-1995. Stat Bull Metrop Insur Co 1997;78(4):20-6.

Mauldin JG, Mauldin PD, Feng TI, et al. Determining the clinical efficacy and cost savings of successful external cephalic version. Am J Obstet Gynecol 1996;175(6):1639-44.

Maymon R, Haimovich L, Shulman A, et al. Third-trimester uterine rupture after prostaglandin E2 use for labor induction. J Reprod Med 1992;37(5):449-52.

Mc Gurgan P, Coulter-Smith S, PJ OD. A national confidential survey of obstetrician's personal preferences regarding mode of delivery. Eur J Obstet Gynecol Reprod Biol 2001;97(1):17-9.

McClain CS. Why women choose trial of labor or repeat cesarean section. J Fam Pract 1985;21(3):210-6.

McClain CS. Patient decision making: the case of delivery method after a previous cesarean section. Cult Med Psychiatry 1987;11(4):495-508.

McClain CS. The making of a medical tradition: vaginal birth after cesarean. Soc Sci Med 1990;31(2):203-10.

McCloskey L, Kosecoff J, Brook RJ. Conformity to recommendations on childbirth by cesarean delivery. In: Kanouse DE, Winkler JD, Kosecoff J, editors. Changing medical practice through technology assessment: an evaluation of the National Institutes of Health Consensus Development Program. Ann Arbor, MI: Health Administration Press; 1989. p. 28.

McCloskey L, Petitti DB, Hobel CJ. Variations in the use of cesarean delivery for dystocia: lessons about the source of care. Med Care 1992;30(2):126-35.

McGurgan P, Coulter-Smith S, PJ OD. A national confidential survey of obstetrician's personal preferences regarding mode of delivery. Eur J Obstet Gynecol Reprod Biol 2001;97(1):17-9.

McIntosh DG, Rayburn WF. Patient-controlled analgesia in obstetrics and gynecology. Obstet Gynecol 1991;78(6):1129-35.

McIntosh ID. Hospital effects of maternity early discharge. Med Care 1984;22(7):611-9.

McKenna JP, Guerdan BR, Wright JC. Vaginal birth after cesarean section. A safe option in carefully selected patients. Postgrad Med 1988;84(6):211-5.

McKenzie L, Stephenson PA. Variation in cesarean section rates among hospitals in Washington State. Am J Public Health 1993;83(8):1109-12.

McMahon MJ. Vaginal birth after cesarean. Clin Obstet Gynecol 1998;41(2):369-81.

McMahon MJ, Luther ER, Bowes WA, et al. Comparison of a trial of labor with an elective second cesarean section. N Engl J Med 1996;335(10):689-95.

McNally OM, Turner MJ. Induction of labour after 1 previous Caesarean section. Aust N Z J Obstet Gynaecol 1999;39(4):425-9.

McNellis D, Medearis AL, Fowler S, et al. A clinical trial of induction of labor versus expectant management in postterm pregnancy: The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. Am J Obstet Gynecol 1994;170(3):716-723.

Meehan FP. Delivery following prior cesarean section: an obstetrician's dilemma? Obstet Gynecol Survey 1988;43(10):582-9.

Meehan FP. Trial of scar with induction/oxytocin in delivery following prior section. Clin Exp Obstet Gynecol 1988;15(4):117-123.

Meehan FP, Burke G. Trial of labour following prior section; a 5 year prospective study (1982-1987). Eur J Obstet Gynecol Reprod Biol 1989;31(2):109-17.

Meehan FP, Burke G, Casey C, et al. Delivery following cesarean section and perinatal mortality. Am J Perinatol 1989;6(1):90-4.

Meehan FP, Burke G, Kehoe JT. Update on delivery following prior cesarean section: a 15-year review 1972-1987. Int J Gynaecol Obster 1989;30(3):205-12.

Meehan FP, Burke G, Kehoe JT, et al. True rupture/scar dehiscence in delivery following prior section. Int J Gynaecol Obster 1990;31(3):249-55.

Meehan FP, Magani IM. True rupture of the caesarean section scar (a 15 year review, 1972-1987). Eur J Obstet Gynecol Reprod Biol 1989;30(2):129-35.

Meehan FP, Moolgaoker AS, Stallworthy J. Vaginal delivery under caudal analgesia after caesarean section and other major uterine surgery. BMJ 1972;2(816):740-2.

Megafu U. Factors influencing maternal survival in ruptured uterus. Int J Gynaecol Obster 1985;23(6):475-80.

Meier PR, Porreco RP. Trial of labor following cesarean section: a two-year experience. Am J Obstet Gynecol 1982;144(6):671-8.

Melchior J, Bernard N. Incidence and pattern of fetal heart rate alterations during labor. In: Kunzel W, editor. Heart rate monitoring: clinical practice and physiology. Berlin: Springer-Verlag; 1985.

Melnikow J, Romano P, Gilbert WM, et al. Vaginal birth after cesarean in California. Obstet Gynecol 2001;98(3):421-6.

Menacker F, Curtin SC. Trends in Cesarean Birth and Vaginal Birth After Previous Cesarean, 1991-99. Natl Vital Stat Rep 2001;49(13):1-16.

Menihan CA. Uterine rupture in women attempting a vaginal birth following prior cesarean birth. J Perinatol 1998;18(6 Pt 1):440-3.

Menihan CA. The effect of uterine rupture on fetal heart rate patterns. J Nurse Midwifery 1999;44(1):40-6.

Menticoglou SM, Manning FA, Morrison I, et al. Must macrosomic fetuses be delivered by a caesarean section? A review of outcome for 786 babies greater than or equal to 4,500 g. Aust N Z J Obstet Gynaecol 1992;32(2):100-3.

Merrill DC, Zlatnik FJ. Randomized, double-masked comparison of oxytocin dosage in induction and augmentation of labor. Obstet Gynecol 1999;94(3):455-63.

Mesleh RA, Kurdi AM, Algwiser AA. Intrapartum rupture of the uterus - 19 Years' experience. Int J Gynecol Obstet 1999; 64(3):311-312.

Metcalfe J. Midwives' Journal. An image restored. Nurs Times 1986;82(22):66.

Michaels WH, Thompson HO, Boutt A, et al. Ultrasound diagnosis of defects in the scarred lower uterine segment during pregnancy. Obstet Gynecol 1988;71(1):112-20.

Miklos JR, Sze E, Parobeck D, et al. Vesicouterine fistula: a rare complication of vaginal birth after cesarean. Obstet Gynecol 1995;86(4 Pt 2):638-9.

Miksovsky P, Watson WJ. Obstetric vacuum extraction: state of the art in the new millennium. Obstet Gynecol Survey 2001;56(11):736-51.

Miles AM, Monga M, Waller K, et al. Risk factors for symptomatic uterine rupture during a trial of labor: The 1990s. Am J Perinatol 2000;17(7):385-389.

Miller CF, Sutter CS. Vaginal birth after cesarean. J Obstet Gynecol Neonat Nurs 1985;14(5):383-9.

Miller DA, Diaz FG, Paul RH. Vaginal birth after cesarean: a 10-year experience. Obstet

Gynecol 1994;84(2):255-8.

Miller DA, Goodwin TM, Gherman RB, et al. Intrapartum rupture of the unscarred uterus. Obstet Gynecol 1997;89(5 Pt 1):671-3.

Miller DA, Mullin P, Hou D, et al. Vaginal birth after cesarean section in twin gestation. Am J Obstet Gynecol 1996;175(1):194-8.

Miller ES, Partezana J, Montgomery RL. Vaginal birth after cesarean: a 5-year experience in a family practice residency program. J Am Board Fam Pract 1995;8(5):357-60.

Miller J. The role of the legal system in the cesarean childbirth controversy. Nurs Law Ethics 1980;1(10):1-2, 6, 8.

Miller JM, Jr. Maternal and neonatal morbidity and mortality in cesarean section. Obstetrics & Gynecology Clinics of North America 1988;15(4):629-38.

Miller M, Leader LR. Vaginal delivery after caesarean section. Aust N Z J Obstet Gynaecol 1992;32(3):213-6.

Minor AF. The cost of maternity care and childbirth in the United States, 1989. Washington: Health Insurance Association of America, Research Bulletin 1989;December.

Misra M, Vavre S. Labour induction with intracervical prostaglandin E2 gel and intravenous oxytocin in women with a very unfavourable cervix. Aust N Z J Obstet Gynaecol 1994;34(5):511-515.

Mocanu EV, Greene RA, Byrne BM, et al. Obstetric and neonatal outcome of babies weighing more than 4.5 kg: an analysis by parity. Eur J Obstet Gynecol Reprod Biol 2000;92(2):229-33.

Moller M, Thomsen AC, Sorensen J, et al. Oxytocin- or low-dose prostaglandin F2 alpha-infusion for stimulation of labor after primary rupture of membranes. A prospective, randomized trial. Acta Obstet Gynecol Scand 1987;66(2):103-6.

Molloy BG, Sheil O, Duignan NM. Delivery after caesarean section: review of 2176 consecutive cases. Br Med J (Clin Res Ed) 1987;294(6588):1645-7.

Moore TR, Origel W, Key TC, et al. The perinatal and economic impact of prenatal care in a low-socioeconomic population. Am J Obstet Gynecol 1986;154(1):29-33.

Mootabar H, Dwyer JF, Surur F, et al. Vaginal delivery following previous cesarean section in 1983. Int J Gynaecol Obstet 1984;22(2):155-60.

Mor-Yosef S, Zeevi D, Samueloff A, et al. Vaginal delivery following one previous cesarean birth: nation wide survey. Asia Oceania J Obstet Gynaecol 1990;16(1):33-7.

Moran C, Carranza-Lira S, Ochoa R, et al. Gastrin levels in mothers and neonates at delivery in various perinatal conditions. Acta Obstet Gynecol Scand 1996;75(7):608-11.

Morewood GA, O'Sullivan MJ, McConney J. Vaginal delivery after cesarean section. Obstet Gynecol 1973;42(4):589-95.

Morgan MA, Thurnau GR. Efficacy of the fetal-pelvic index for delivery of neonates weighing 4000 grams or greater: a preliminary report. Am J Obstet Gynecol 1988;158(5):1133-7.

Morgan MA, Thurnau GR, Fishburne JI. The fetal-pelvic index as an indicator of fetal-pelvic disproportion: a preliminary report. Am J Obstet Gynecol 1986;155(3):608-13.

Moriya M, Kusaka H, Shimizu K, et al. Spontaneous rupture of the uterus caused by placenta percreta at 28 weeks of gestation: a case report. J Obstet Gynaecol Res 1998;24(3):211-4.

Morrison JJ, Rennie JM, Milton PJ. Neonatal respiratory morbidity and mode of delivery at term: influence of timing of elective caesarean section. Br J Obstet Gynaecol 1995;102(2):101-6.

Moskovitz H, O'Grady JP, Gimovsky M. Fetal heart rate monitoring casebook. Uterine rupture and sinusoidal heart rate. J Perinatol 1994;14(2):154-8.

Mould TA, Chong S, Spencer JA, et al. Women's involvement with the decision preceding their caesarean section and their degree of satisfaction. Br J Obstet Gynaecol 1996;103(11):1074-7.

Mousa HA, Mahmood TA. Do practice guidelines guide practice? A prospective audit of

induction of labor three years experience. Acta Obstet Gynecol Scand 2000; 79(12):1086-92.

Mozurkewich EL, Hutton EK. Elective repeat cesarean delivery versus trial of labor: a meta-analysis of the literature from 1989 to 1999. Am J Obstet Gynecol 2000;183(5):1187-97.

Mullings AM, Hall JS. Rupture of uterus and bladder in vaginal delivery following previous caesarean section. West Indian Med J 1987;36(1):51-3.

Mundle WR, Young DC. Vaginal misoprostol for induction of labor: a randomized controlled trial. Obstet Gynecol 1996;88(4 Pt 1):521-5.

Murphy MC, Harvey SM. Choice of a childbirth method after cesarean. Women Health 1989;15(2):67-85.

Murray TM. Case 9-1998: uterine rupture. N Engl J Med 1998;339(4):268; discussion 269.

Mushinski M. Average charges for uncomplicated cesarean and vaginal deliveries, United States, 1993. Stat Bull Metrop Insur Co 1994;75(4):27-36.

Mushinski M. Average charges for uncomplicated vaginal, cesarean and VBAC deliveries: regional variations, United States, 1996. Stat Bull Metrop Insur Co 1998;79(3):17-28.

Myers SA, Gleicher N. Breech delivery: why the dilemma? Am J Obstet Gynecol 1986;156(1):6-10.

Myers SA, Gleicher N. A successful program to lower cesarean-section rates. N Engl J Med 1988;319(23):1511-6.

Myers SA, Gleicher N. A program to lower cesarean section rates [letter]. N Engl J Med 1989;320:1693.

Myers SA, Gleicher N. 1988 U.S. cesarean-section rate: good news or bad? N Engl J Med 1990;323(3):200.

Myers SA, Gleicher N. Implementation of a cesarean section program: the quality assurance process. QRB. Qual Rev Bull 1991;17(5):162-166.

Myers SA, Gleicher N. The Mount Sinai cesarean section reduction program: an update after 6 years. Soc Sci Med 1993;37(10):1219-22.

Myerscough P. Systematic reviews: insufficient evidence on which to base medicine. Br J Obstet Gynaecol 1998;105(11):1235; discussion 1236.

Myles T. Vaginal birth of twins after a previous Cesarean section. J Matern Fetal Med 2001;10(3):171-4.

Nachman KL, Oleske DM, Linn ES, et al. The effect of obstetrical quality process initiatives on the likelihood of newborn birth trauma in Medicaid managed care, Medicaid fee-for-service, and private managed care. Abstr Book Assoc Health Serv Res;15:198.

Naef RW, Ray MA, Chauhan SP, et al. Trial of labor after cesarean delivery with a lower-segment, vertical uterine incision: is it safe? Am J Obstet Gynecol 1995;172(6):1666-73; discussion 1673-4.

Naeye RL. Maternal age, obstetric complications, and the outcome of pregnancy. Obstet Gynecol 1983;61(2):210-6.

Nagarkatti RS, Ambiye VR, Vaidya PR. Ruptured uterus: changing trends in etiology and management. J Postgrad Med 1991;37:136-139.

Naiden J, Deshpande P. Using active management of labor and vaginal birth after previous cesarean delivery to lower cesarean delivery rates: aA 10-year experience. Am J Obstet Gynecol 2001;184(7):1535-41; discussion 1541-3.

National Institute of Child Health & Human Development Research Planning Workshop. Electronic fetal heart rate monitoring: research guidelines for interpretation. Am J Obstet Gynecol 1997;177(6):1385-90.

Neilson DR, Prins RP, Bolton RN, et al. A comparison of prostaglandin E2 gel and prostaglandin F2 alpha gel for preinduction cervical ripening. Am J Obstet Gynecol 1983;146(5):526-32.

Nelson LJ. Legal dimensions of maternal-fetal conflict. Clin Obstet Gynecol 1992;35(4):738-48.

Nelson LJ, Milliken N. Compelled medical treatment of pregnant women. Life, liberty, and law in conflict. JAMA 1988;259(7):1060-6.

Nesbitt TS, Tanji JL, Scherger JE, et al. Obstetric care, Medicaid, and family physicians. How policy changes affect physicians' attitudes. West J Med 1991;155(6):653-7.

Newton ER, Higgins CS. Factors associated with hospital-specific cesarean birth rates. J Reprod Med 1989;34(6):407-11.

Ngu A, Quinn MA. Vaginal delivery following caesarean section. Aust N Z J Obstet Gynaecol 1985;25(1):41-3.

Nguyen TV, Dinh TV, Suresh MS, et al. Vaginal birth after cesarean section at the University of Texas. J Reprod Med 1992;37(10):880-2.

Nielsen TF, Hokegard KH. Postoperative cesarean section morbidity: a prospective study. Am J Obstet Gynecol 1983;146(8):911-5.

Nielsen TF, Hokegard KH. The course of subsequent pregnancies after previous cesarean section. Acta Obstet Gynecol Scand 1984;63(1):13-6.

Nielsen TF, Hokegard KH. The incidence of acute neonatal respiratory disorders in relation to mode of delivery. Acta Obstet Gynecol Scand 1984;63(2):109-14.

Nielsen TF, Hokegard KH, Moldin PG. X-ray pelvimetry and trial of labor after previous cesarean section. A prospective study. Acta Obstet Gynecol Scand 1985;64(6):485-90.

Nielsen TF, Ljungblad U, Hagberg H. Rupture and dehiscence of cesarean section scar during pregnancy and delivery. Am J Obstet Gynecol 1989;160(3):569-73.

Nimrod C, Currie J, Yee J, et al. Cervical ripening and labor induction with intracervical triacetin base prostaglandin E2 gel: a placebo-controlled study. Obstet Gynecol 1984;64(4):476-9.

Nissen E, Gustavsson P, Widstrom AM, et al. Oxytocin, prolactin, milk production and their relationship with personality traits in women after vaginal delivery or Cesarean section. J Psychosomat Obstet Gynecol 1998;19(1):49-58.

Noah ML, DeCoster JM, Fraser TJ, et al. Preinduction cervical softening with endocervical PGE2 gel. A multi-center trial. Acta Obstet Gynecol Scand 1987;66(1):3-7.

Nordin AJ, Richardson JA. Lower segment uterine scar rupture during induction of labour with vaginal prostaglandin E2 [1]. Postgrad Med J 1993;69(813):592.

Norman M, Ekman G. Preinductive cervical ripening with prostaglandin E2 in women with one previous cesarean section. Acta Obstet Gynecol Scand 1992;71(5):351-355.

Norman P. Vaginal birth after caesarean section. Lancet 1995;345(8943):142.

Norman P, Kostovcik S, Lanning A. Elective repeat cesarean sections: how many could be vaginal births? Can Med Assoc J 1993;149(4):431-5.

Notzon FC. International differences in the use of obstetric interventions. JAMA 1990;263(24):3286-91.

Notzon FC, Cnattingius S, Bergsjo P, et al. Cesarean section delivery in the 1980s: international comparison by indication. Am J Obstet Gynecol 1994;170(2):495-504.

Novas J, Myers SA, Gleicher N. Obstetric outcome of patients with more than one previous cesarean section. Am J Obstet Gynecol 1989;160(2):364-7.

O'Boyle AL, Woodman PJ, O'Boyle JD, et al. Pelvic organ support in nulliparous pregnant and nonpregnant women: a case control study. Am J Obstet Gynecol 2002;187(1):99-102.

O'Brien JM, Barton JR, Donaldson ES. The management of placenta percreta: conservative and operative strategies. Am J Obstet Gynecol 1996;175(6):1632-8.

O'Bryant C. QQ: VBAC waterbirth. Midwifery Today 2000(54):34.

O'Connor KM. How safe is induction of labour following a previous caesarean section? J Obstet Gynaecol 1983;4(2):86-87.

O'Connor RA, Gaughan B. Pregnancy following simple repair of the ruptured gravid uterus. Br J

Obstet Gynaecol 1989;96(8):942-4.

O'Driscoll K, Foley M, MacDonald D, et al. Cesarean section and perinatal outcome: response from the House of Horne. Am J Obstet Gynecol 1988;158(3 Pt 1):449-52.

O'Flaherty L. Oral misoprostol and intracervical dinoprostone for cervical ripening and labor induction: a randomized comparison. J Midwifery Womens Health 2001;46(3):203-4.

O'Meara C. An evaluation of consumer perspectives of childbirth and parenting education. Midwifery 1993;9(4):210-9.

O'Neil JD, Gilbert P, Kusugak N, et al. Obstetric policy for the Keewatin Region, N.W.T.: results of the childbirth experience survey. Arctic Med Res 1991;Suppl:572-6.

O'Sullivan MJ, Fumia F, Holsinger K, et al. Vaginal delivery after cesarean section. Clin Perinatol 1981;8(1):131-43.

Obara H, Minakami H, Koike T, et al. Vaginal birth after cesarean delivery: results in 310 pregnancies. Journal of Obstetrics & Gynaecology Research 1998;24(2):129-34.

Oberer D, Auckerman L. Best practice: clinical pathways for uncomplicated births. Best Pract Benchmarking Healthc 1996;1(1):43-50.

Oberman LH. Obstetrics: reducing C-sections without financial ruin. Trustee 1989;42(4):20.

Obst TE, Nauenberg E, Buck GM. Maternal health insurance coverage as a determinant of obstetrical anesthesia care. Journal of Health Care for the Poor & Underserved 2001;12(2):177-91.

Odeh M, Tarazova L, Wolfson M, et al. Evidence that women with a history of cesarean section can deliver twins safely. Acta Obstet Gynecol Scand 1997;76(7):663-6.

Odibo AO, O'Coigligh J, Selinger M. Planned vaginal delivery after two previous caesarean sections. Br J Obstet Gynaecol 1995;102(3):262-3.

Ohel G, Haddad S, Linder N. Fetal urine production after induction of labor with prostaglandin E2. Am J Obstet Gynecol 1994;171(5):1242-1243.

Ojo VA, Okwerekwu FO. Vaginal delivery in first pregnancy following a primary caesarean section. Asia Oceania J Obstet Gynaecol 1989;15(2):121-6.

Oleske DM, Glandon GL, Giacomelli GJ, et al. The cesarean birth rate: influence of hospital teaching status. Health Serv Res 1991;26(3):325-37.

Oleske DM, Glandon GL, Tancredi DJ, et al. Information dissemination and the cesarean birth rate. The Illinois experience. Int J Technol Assess Health Care 1992;8(4):708-18.

Oleske DM, Linn ES, Nachman KL, et al. Effect of Medicaid managed care on pregnancy complications. Obstet Gynecol 2000;95(1):6-13.

Oleske DM, Linn ES, Nachman KL, et al. Cesarean and VBAC delivery rates in Medicaid managed care, Medicaid fee-for-service, and private managed care. Birth 1998;25(2):125-7.

Ollendorff DA, Goldberg JM, Minogue JP, et al. Vaginal birth after cesarean section for arrest of labor: is success determined by maximum cervical dilatation during the prior labor? Am J Obstet Gynecol 1988;159(3):636-9.

Onuora VC, al Ariyan R, Koko AH, et al. Major injuries to the urinary tract in association with childbirth. East Afr Med J 1997;74(8):523-6.

Ophir E, Oettinger M, Yagoda A, et al. Breech presentation after cesarean section: always a section? Am J Obstet Gynecol 1989;161(1):25-8.

Ophir E, Yagoda A, Rojansky N, et al. Trial of labor following cesarean section: dilemma. Obstet Gynecol Survey 1989;44(1):19-24.

Osmers R, Ulbrich R, Schauer A, et al. Sonographic detection of an asymptomatic rupture of the uterus due to necrosis during the third trimester. Int J Gynaecol Obster 1988;26(2):279-84.

Ottinger WS, Menard MK, Brost BC, et al. A randomized clinical trial of prostaglandin E2 intracervical gel and a slow release vaginal pessary for preinduction cervical ripening. Am J Obstet Gynecol 1998;179(2):349-353.

Oumachigui A, Rajagopalan G, Reddy R, et al. Placenta accreta and percreta: a review of 5 cases. Int J Gynaecol Obster 1981;19(4):337-40.

Ouzounian JG, Miller DA, Paul RH. Amnioinfusion in women with previous cesarean births: a preliminary report. Am J Obstet Gynecol 1996;174(2):783-6.

Oyelese Y, Landy HJ, Collea JV. Cervical laceration associated with misoprostol induction. Int J Gynaecol Obster 2001;73(2):161-2.

Ozumba BC, Nwabue R. Spontaneous uterine rupture following low upper segment transverse incision. Trop Doct 1997;27(2):111-2.

Paech MJ. The King Edward Memorial Hospital 1,000 mother survey of methods of pain relief in labour. Anaesth Intensive Care 1991;19(3):393-9.

Paraskevaides E, Stuart B, Gardeil F. Secondary postpartum haemorrhage from nondehisced lower caesarean section scar: a case for hysteroscopy. Aust N Z J Obstet Gynaecol 1993;33(4):427.

Parrish KM, Holt VL, Connell FA, et al. Variations in the accuracy of obstetric procedures and diagnoses on birth records in Washington State, 1989. Am J Epidemiol 1993;138(2):119-27.

Parrish KM, Holt VL, Easterling TR, et al. Effect of changes in maternal age, parity, and birth weight distribution on primary cesarean delivery rates. JAMA 1994;271(6):443-7.

Paterson CM, Saunders NJ. Mode of delivery after one caesarean section: audit of current practice in a health region. BMJ 1991;303(6806):818-21.

Paterson CM, Saunders NS. The fetal maturity and mode of delivery of a woman's first baby influences the characteristics of her next labour. Eur J Obstet Gynecol Reprod Biol 1991;42(3):187-93.

Pauerstein CJ. Labor after cesarean section: from precept to practice. J Reprod Med 1981;26(8):409-12.

Pauerstein CJ, Karp L, Muher S. Trial of labor after low segment cesarean section. South Med J 1969;62(8):925-8.

Paul RH. Cesarean section: at term on request [editorial]. Perinatol Neonatol 1988;12(4):7.

Paul RH. Toward fewer cesarean sections--the role of a trial of labor. N Engl J Med 1996;335(10):735-6.

Paul RH, Phelan JP, Yeh SY. Trial of labor in the patient with a prior cesarean birth. Am J Obstet Gynecol 1985;151(3):297-304.

Pauly MV, Redisch M. The not-for-profit hospital as a physicians' cooperative. Am Econ Rev 1973;63(1):87-99.

Peaceman AM, Lopez-Zeno JA, Minogue JP, et al. Factors that influence route of delivery--active versus traditional labor management. Am J Obstet Gynecol 1993;169(4):940-4.

Pearl ML, Escamilla G, Karpel BM, et al. Conservative management of placenta percreta with involvement of the ileum. Arch Gynecol Obstet 1996;258(3):147-50.

Pel M, Heres MH, Hart AA, et al. Provider-associated factors in obstetric interventions. Eur J Obstet Gynecol Reprod Biol 1995;61(2):129-34.

Peleg D, Hannah ME, Hodnett ED, et al. Predictors of cesarean delivery after prelabor rupture of membranes at term. Obstet Gynecol 1999;93(6):1031-5.

Pello LC, Rosevear SK, Dawes GS, et al. Computerized fetal heart rate analysis in labor. Obstet Gynecol 1991;78(4):602-10.

Pelosi MA. Vesicouterine fistula: a rare complication of vaginal birth after cesarean. Obstet Gynecol 1996;87(1):160-2.

Penso C. Vaginal birth after cesarean section: An update on physician trends and patient perceptions. Curr Opin Obstet Gynecol 1994;6(5):417-425.

Perveen F, Shah Q. Obstetric outcome after one previous caesarean section. J Obstet Gynaecol Res 1997;23(4):341-6.

Peters C, Cowley M, Standiford L. The process of outcomes management in an acute care

facility. Nurs Adm Q 1999;24(1):75-89.

Petitti D. Maternal mortality and morbidity in cesarean section. Clin Obstet Gynecol 1985;28(4):763-69.

Petitti DB, Cefalo RC, Shapiro S, et al. In-hospital maternal mortality in the United States: time trends and relation to method of delivery. Obstet Gynecol 1982;59(1):6-12.

Petrikovsky BM. Endoscopic assessment of the integrity of the postcesarean uterine wall before a trial of labor. Transcervical Endoscopy Registry. J Reprod Med 1994;39(6):464-6.

Petrikovsky BM. Rupture of the scarred uterus: prediction and diagnosis. Lancet 1996;347(9004):838-9.

Petrou S, Henderson J, Glazener C. Economic aspects of caesarean section and alternative modes of delivery. Best Prac Res Clin Obstet Gynaecol 2001;15(1):145-63.

Phelan JP. Delivery following cesarean section and perinatal mortality. Am J Perinatol 1989;6(1):95-6.

Phelan JP. VBAC: Time to reconsider? Obstet Gynecol Manag 1996;11:62-68.

Phelan JP, Ahn MO, Diaz F, et al. Twice a cesarean, always a cesarean? Obstet Gynecol 1989;73(2):161-5.

Phelan JP, Ahn MO, Korst L, et al. Intrapartum fetal asphyxial brain injury with absent multiorgan system dysfunction. J Matern Fetal Med 1998;7(1):19-22.

Phelan JP, Boucher M, Mueller E, et al. The nonlaboring transverse lie. A management dilemma. J Reprod Med 1986;31(3):184-6.

Phelan JP, Clark SL, Diaz F, et al. Vaginal birth after cesarean. Am J Obstet Gynecol 1987;157(6):1510-5.

Phelan JP, Eglinton GS, Horenstein JM, et al. Previous cesarean birth. Trial of labor in women with macrosomic infants. J Reprod Med 1984;29(1):36-40.

Phelan JP, Korst LM, Settles DK. Uterine activity patterns in uterine rupture: a case-control study. Obstet Gynecol 1998;92(3):394-7.

Phibbs CS, Mark DH, Luft HS, et al. Choice of hospital for delivery: a comparison of high-risk and low-risk women. Health Serv Res 1993;28(2):201-22.

Phillips RN, Thornton J, Gleicher N. Physician bias in cesarean sections. JAMA 1982;248(9):1082-4.

Pickhardt MG, Martin JN, Meydrech EF, et al. Vaginal birth after cesarean delivery: are there useful and valid predictors of success or failure? Am J Obstet Gynecol 1992;166(6 Pt 1):1811-5; discussion 1815-9.

Placek PJ, Taffel SM. Recent patterns in cesarean delivery in the United States. Obstet Gynecol Clin North Am 1988;15(4):607-27.

Placek PJ, Taffel SM. Vaginal birth after cesarean (VBAC) in the 1980s. Am J Public Health 1988;78(5):512-5.

Placek PJ, Taffel SM, Moien M. 1986 C-sections rise; VBACs inch upward. Am J Public Health 1988;78(5):562-3.

Plauche WC, Von Almen W, Muller R. Catastrophic uterine rupture. Obstet Gynecol 1984;64(6):792-7.

Plaut MM, Schwartz ML, Lubarsky SL, et al. Uterine rupture associated with the use of misoprostol in the gravid patient with a previous cesarean section. Am J Obstet Gynecol 1999;180(6 Pt I):1535-1542.

Poma PA. Effects of obstetrician characteristics on cesarean delivery rates. A community hospital experience. Am J Obstet Gynecol 1999;180(6 Pt 1):1364-72.

Poma PA. Rupture of a cesarean-scarred uterus: a community hospital experience. J Natl Med Assoc 2000;92(6):295-300.

Porreco RP. High cesarean section rate: a new perspective. Obstet Gynecol 1985;65(3):307-11.

Porreco RP, Klaus MH, Shearer E, et al. Commentaries: the cesarean section rate is 25 percent and rising. Why? What can be done about it? Birth 1989;16(3):118-22.

Porreco RP, Meier PR. Trial of labor in patients with multiple previous cesarean sections. J Reprod Med 1983;28(11):770-2.

Prabulos AM, Philipson EH. Umbilical cord prolapse. Is the time from diagnosis to delivery critical? J Reprod Med 1998;43(2):129-32.

Prasad RN, Ratnam SS. Uterine rupture after induction of labour for intrauterine death using the prostaglandin E2 analogue sulprostone. Aust N Z J Obstet Gynaecol 1992;32(3):282-3.

Prendergast MS, Anderle LJ. Experience with vaginal delivery after prior cesarean birth in a community hospital. J Am Osteopath Assoc 1985;85(4):238-40.

Press F, Katz M, Leiberman JR, et al. Obstetric performance in Ethiopian immigrants compared with Israeli parturients. Isr J Med Sci 1993;29(6-7):403-7.

Pridjian G. Labor after prior cesarean section. Clin Obstet Gynecol 1992;35(3):445-456.

Pridjian G, Hibbard JU, Moawad AH. Cesarean: changing the trends. Obstet Gynecol 1991;77(2):195-200.

Proctor S. What determines quality in maternity care? Comparing the perceptions of childbearing women and midwives. Birth 1998;25(2):85-93.

Pruett KM, Kirshon B, Cotton DB. Unknown uterine scar and trial of labor. Am J Obstet Gynecol 1988;159(4):807-10.

Pruett KM, Kirshon B, Cotton DB, et al. Is vaginal birth after two or more cesarean sections safe? Obstet Gynecol 1988;72(2):163-5.

Prysak M, Castronova FC. Elective induction versus spontaneous labor: a case-control analysis of safety and efficacy. Obstet Gynecol 1998;92(1):47-52.

Purdie FR, Nieto JM, Summerson DJ, et al. Rupture of the uterus with DIC. Ann Emerg Med 1983;12(3):174-6.

Puza S, Roth N, Macones GA, et al. Does cesarean section decrease the incidence of major birth trauma? J Perinatol 1998;18(1):9-12.

Quilligan EJ. Vaginal birth after Cesarean section: 270 degrees. Journal of Obstetrics & Gynaecology Research 2001;27(4):169-73.

Quinlivan JA, Petersen RW, Nichols CN. Patient preference the leading indication for elective Caesarean section in public patients--results of a 2-year prospective audit in a teaching hospital. Aust N Z J Obstet Gynaecol 1999;39(2):207-14.

Quirk JG, Jr., Fortney J, Collins HB, 2nd, et al. Outcomes of newborns with gastroschisis: the effects of mode of delivery, site of delivery, and interval from birth to surgery. Am J Obstet Gynecol 1996;174(4):1134-8; discussion 1138-40.

Rabinerson D, Rafael ZB, Dekel A. Vaginal birth after cesarean delivery? Am J Obstet Gynecol 2001;184(4):780.

Rachagan SP, Raman S, Balasundram G, et al. Rupture of the pregnant uterus--a 21-year review. Aust N Z J Obstet Gynaecol 1991;31(1):37-40.

Rageth JC, Juzi C, Grossenbacher H. Delivery after previous cesarean: a risk evaluation. Swiss Working Group of Obstetric and Gynecologic Institutions. Obstet Gynecol 1999;93(3):332-7.

Rajaram P, Oumachigui A. Trial of labor after cesarean section. Am J Obstet Gynecol 1984;149(4):471-2.

Ramsey PS, Johnston BW, Welter VE, et al. Artifactual fetal electrocardiographic detection using internal monitoring following intrapartum fetal demise during VBAC trial. J Matern Fetal Med 2000;9(6):360-1.

Ranzinger M, Fuentes A, Smyk LV. Spontaneous rupture of a low transverse cesarean scar. South Med J 1994;87(10):1001-2.

Raskin KS, Dachauer JD, Doeden AL, et al. Uterine rupture after use of a prostaglandin E2 vaginal insert during vaginal birth after cesarean: A report of two cases. J Reprod Med Obstet Gynecol 1999;44(6):571-574.

Ratten GJ. Medicolegal matters involving a major obstetric and gynaecological teaching hospital. Aust N Z J Obstet Gynaecol 1997;37(2):192-4.

Raube K, Merrell K, Gomberg MC. Maternal

minimum-stay legislation: cost and policy implications. Abstr Book Assoc Health Serv Res;14:302.

Ravasia DJ, Brain PH, Pollard JK. Incidence of uterine rupture among women with mullerian duct anomalies who attempt vaginal birth after cesarean delivery. Am J Obstet Gynecol 1999;181(4):877-81.

Ravasia DJ, Wood SL, Pollard JK. Uterine rupture during induced trial of labor among women with previous cesarean delivery. Am J Obstet Gynecol 2000;183(5):1176-9.

Rayburn W, Gosen R, Ramadei C, et al. Outpatient cervical ripening with prostaglandin E2 gel in uncomplicated postdate pregnancies. Am J Obstet Gynecol 1988;158(6 Pt 1):1417-23.

Rayburn WF. Prostaglandin E2 gel for cervical ripening and induction of labor: a critical analysis. Am J Obstet Gynecol 1989;160(3):529-34.

Rayburn WF. Clinical experience with a controlled-release, prostaglandin E2 intravaginal insert in the USA. Br J Obstet Gynaecol 1997;104(Suppl 15):8-12; discussion 20-5.

Rayburn WF, Gittens LN, Lucas MJ, et al. Weekly administration of prostaglandin E2 gel compared with expectant management in women with previous cesareans. Obstet Gynecol 1999;94(2):250-254.

Raynor BD. The experience with vaginal birth after cesarean delivery in a small rural community practice. Am J Obstet Gynecol 1993;168(1 Pt 1):60-2.

Read AW, Waddell VP, Prendiville WJ, et al. Trends in caesarean section in Western Australia, 1980-1987. Med J Aust 1990;153(6):318-23.

Read JA. The scheduling of repeat cesarean section operations: prospective management protocol experience. Am J Obstet Gynecol 1985;151(5):557-63.

Reddy UM, DiVito MM, Armstrong JC, et al. Population adjustment of the definition of the vaginal birth after cesarean rate. Am J Obstet Gynecol 2000;183(5):1166-9.

Redwine DB. Vaginal delivery in patients with a prior cesarean section. Obstet Gynecol 1983;61(3):399-400.

Reid AJ, Carroll JC, Ruderman J, et al. Differences in intrapartum obstetric care provided to women at low risk by family physicians and obstetricians. Can Med Assoc J 1989;140(6):625-33.

Reid MC. Use of methodological standards in diagnostic testing research. JAMA 1995;274:645-57.

Resick LK, Erlen JA. Vaginal birth after cesarean: issues and implications. J Am Acad Nurs Pract 1990;2(3):100-6.

Resnick MB, Ariet M, Carter RL, et al. Prospective pricing system by diagnosis-related groups: comparison of federal diagnosis-related groups with high-risk obstetric care groups. Am J Obstet Gynecol 1987;156(3):567-73.

Reynolds JL. Post-traumatic stress disorder after childbirth: the phenomenon of traumatic birth. Can Med Assoc J 1997;156(6):831-5.

Rhodes MK. Early discharge of mothers and infants following vaginal childbirth at the United States Air Force Academy: A three-year study. Mil Med 1994;159(3):227-30.

Rice DP, Kleinman JC. National health data for policy and planning. Health Policy Educ 1980;1(2):129-41.

Rieger N, Schloithe A, Saccone G, et al. The effect of a normal vaginal delivery on anal function. Acta Obstet Gynecol Scand 1997;76(8):769-72.

Riva HL, Teich JC. Vaginal delivery after cesarean section. Am J Obstet Gynecol 1961;81(3):501-10.

Roberts LJ, Beardsworth SA, Trew G. Labour following caesarean section: current practice in the United Kingdom. Br J Obstet Gynaecol 1994;101(2):153-5.

Roberts M, Nordin AJ. Posterior uterine rupture in a patient with a lower segment caesarean section scar complicating prostaglandin induction of labour. Br J Obstet Gynaecol 1997;104(12):1421.

Roberts RG, Bell HS, Wall EM, et al. Trial of labor or repeated cesarean section. The woman's choice. Arch Fam Med 1997;6(2):120-5.

Rochat RW, Koonin LM, Atrash HK, et al. Maternal mortality in the United States: report from the Maternal Mortality Collaborative. Obstet Gynecol 1988;72(1):91-7.

Rock SM. Malpractice premiums and primary cesarean section rates in New York and Illinois. Public Health Rep 1988;103(5):459-63.

Rockner G, Jonasson A, Olund A. The effect of mediolateral episiotomy at delivery on pelvic floor muscle strength evaluated with vaginal cones. Acta Obstet Gynecol Scand 1991;70(1):51-4.

Rodriguez MH, Masaki DI, Phelan JP, et al. Uterine rupture: are intrauterine pressure catheters useful in the diagnosis? Am J Obstet Gynecol 1989;161(3):666-9.

Rodriguez MH, Wang R, Clark SL, et al. Previous cesarean birth: management considerations in the patient with acute puerperal uterine inversion. Am J Obstet Gynecol 1984;150(4):433-5.

Rogers RG, Gardner MO, Tool KJ, et al. Active management of labor: a cost analysis of a randomized controlled trial. West J Med 2000;172(4):240-3.

Roland EH, Poskitt K, Rodriguez E, et al. Perinatal hypoxic-ischemic thalamic injury: clinical features and neuroimaging. Ann Neurol 1998;44(2):161-6.

Rooks JP, Weatherby NL, Ernst EK. The National Birth Center Study. Part III--Intrapartum and immediate postpartum and neonatal complications and transfers, postpartum and neonatal care, outcomes, and client satisfaction. J Nurse Midwifery 1992;37(6):361-97.

Rooks JP, Weatherby NL, Ernst EK, et al. Outcomes of care in birth centers. The National Birth Center Study. N Engl J Med 1989;321(26):1804-11.

Rose V. ACOG urges a cautious approach to vaginal birth after cesarean delivery. Am Fam Physician 1999;59(2):474.

Rose VL. ACOG urges a cautious approach to vaginal birth after cesarean delivery. Am Fam Physician 1999;60(4):1245-6.

Rosen MG, Bilenker RM, Thompson K. Assessment of developmental time periods and risks of brain damage in the fetus and neonate. J Reprod Med 1986;31(5):297-303.

Rosen MG, Dickinson JC. Vaginal birth after cesarean: a meta-analysis of indicators for success. Obstet Gynecol 1990;76(5 Pt 1):865-9.

Rosen MG, Dickinson JC, Westhoff CL. Vaginal birth after cesarean: a meta-analysis of morbidity and mortality. Obstet Gynecol 1991;77(3):465-70.

Rothstein EP, Faccenda DR, Rubenstein SD. Neonatal morbidity of abdominal and vaginal deliveries after uncomplicated pregnancies. Arch Pediatr Adolesc Med 1996;150(6):653-4.

Roumen FJ, Janssen AA, Vrouenraets FP. The course of delivery after previous cesarean section. Eur J Obstet Gynecol Reprod Biol 1990;34(1-2):15-20.

Rourke JT. Trends in small hospital obstetric services in Ontario. Can Fam Physician 1998;44:2117-24.

Rouse DJ, Owen J, Goldenberg RL, et al. The effectiveness and costs of elective cesarean delivery for fetal macrosomia diagnosed by ultrasound. JAMA 1996;276(18):1480-6.

Rowbottom SJ, Critchley LA, Gin T. Uterine rupture and epidural analgesia during trial of labour. Anaesthesia 1997;52(5):486-8.

Rowlands S, Bell R, Donath S, et al. Misoprostol versus dinoprostone for cervical priming prior to induction of labour in term pregnancy: A randomised controlled trial. Aust N Z J Obstet Gynaecol 2001;41(2):145-152.

Rozenberg P, Goffinet F, Philippe HJ, et al. Thickness of the lower uterine segment: its influence in the management of patients with previous cesarean sections. Eur J Obstet Gynecol Reprod Biol 1999;87(1):39-45.

Rozenberg P, Goffinet F, Phillippe HJ, et al. Ultrasonographic measurement of lower uterine

segment to assess risk of defects of scarred uterus. Lancet 1996;347(8997):281-4.

Rubin GL, Peterson HB, Rochat RW, et al. Maternal death after cesarean section in Georgia. Am J Obstet Gynecol 1981;139(6):681-5.

Ruderman J, Carroll JC, Reid AJ, et al. Are physicians changing the way they practise obstetrics? Can Med Assoc J 1993;148(3):409-15.

Rudick V, Niv D, Hetman-Peri M, et al. Epidural analgesia for planned vaginal delivery following previous cesarean section. Obstet Gynecol 1984;64(5):621-3.

Rust OA, Place JC, Melendez D, et al. Lowering the cesarean rate at a small USAF hospital. Mil Med 1993;158(1):22-6.

Ryding EL. Psychosocial indications for cesarean section. A retrospective study of 43 cases. Acta Obstet Gynecol Scand 1991;70(1):47-9.

Saad AH. Trial of labor compared with an elective second cesarean section. N Engl J Med 1997;336(9):658-9; discussion 659.

Sachs BP. A program to lower cesarean-section rates [letter]. N Engl J Med 1989;320:1694.

Sachs BP, Kobelin C, Castro MA, et al. The risks of lowering the cesarean-delivery rate. N Engl J Med 1999;340(1):54-7.

Sack RA. The effect of utilization on health care costs. Am J Obstet Gynecol 1980;137(2):270-5.

Saglamtas M, Vicdan K, Yalcin H, et al. Rupture of the uterus. Int J Gynaecol Obster 1995;49(1):9-15.

Saisto T, Salmela-Aro K, Nurmi JE, et al. Psychosocial characteristics of women and their partners fearing vaginal childbirth. BJOG 2001;108(5):492-8.

Saisto T, Salmela-Aro K, Nurmi JE, et al. Psychosocial predictors of disappointment with delivery and puerperal depression. A longitudinal study. Acta Obstet Gynecol Scand 2001;80(1):39-45.

Saisto T, Salmela-Aro K, Nurmi JE, et al. A randomized controlled trial of intervention in fear of childbirth. Obstet Gynecol 2001;98(5 Pt 1):820-6.

Saisto T, Ylikorkala O, Halmesmaki E. Factors associated with fear of delivery in second pregnancies. Obstet Gynecol 1999;94(5 Pt 1):679-82.

Sakala C. Midwifery care and out-of-hospital birth settings: how do they reduce unnecessary cesarean section births? Soc Sci Med 1993;37(10):1233-50.

Sakala EP, Erhard LN, White JJ. Elective cesarean section improves outcomes of neonates with gastroschisis. Am J Obstet Gynecol 1993;169(4):1050-3.

Sakala EP, Kaye S, Murray RD, et al. Epidural analgesia. Effect on the likelihood of a successful trial of labor after cesarean section. J Reprod Med 1990;35(9):886-90.

Sakala EP, Kaye S, Murray RD, et al. Oxytocin use after previous cesarean: Why a higher rate of failed labor trial? Obstet Gynecol 1990;75(3 I):356-359.

Saldana LR, Schulman H, Reuss L. Management of pregnancy after cesarean section. Am J Obstet Gynecol 1979;135(5):555-61.

Samueloff A, Mor-Yosef S, Seidman DS, et al. The 1984 national perinatal census: design, organization and uses for assessing obstetric services in Israel. Isr J Med Sci 1989;25(11):629-34.

Sanchez-Ramos L. Dinoprostone vaginal insert for cervical ripening and labor induction: a meta-analysis. Obstet Gynecol 2001;98(3):519-20.

Sanchez-Ramos L, Farah LA, Kaunitz AM, et al. Preinduction cervical ripening with commercially available prostaglandin E2 gel: a randomized, double-blind comparison with a hospital-compounded preparation. Am J Obstet Gynecol 1995;173(4):1079-84.

Sanchez-Ramos L, Gaudier FL, Kaunitz AM. Cervical ripening and labor induction after previous cesarean delivery. Clin Obstet Gynecol 2000;43(3):513-23.

Sanchez-Ramos L, Kaunitz AM. Misoprostol for

cervical ripening and labor induction: a systematic review of the literature. Clin Obstet Gynecol 2000;43(3):475-88.

Sanchez-Ramos L, Kaunitz AM. Uterine rupture associated with the use of prostaglandin E1 in patients with previous cesarean delivery. Am J Obstet Gynecol 2000;182(4):990-1.

Sanchez-Ramos L, Kaunitz AM, Connor PM. Hygroscopic cervical dilators and prostaglandin E2 gel for preinduction cervical ripening. A randomized, prospective comparison. J Reprod Med 1992;37(4):355-9.

Sanchez-Ramos L, Kaunitz AM, Delke I, et al. Cervical ripening and labor induction with a controlled-release dinoprostone vaginal insert: a meta-analysis. Obstet Gynecol 1999;94(5 Pt 2):878-83.

Sanchez-Ramos L, Kaunitz AM, Peterson HB, et al. Reducing cesarean sections at a teaching hospital. Am J Obstet Gynecol 1990;163(3):1081-7; discussion 1087-8.

Sanchez-Ramos L, Kaunitz AM, Wears RL, et al. Misoprostol for cervical ripening and labor induction: a meta-analysis. Obstet Gynecol 1997;89(4):633-42.

Sandmire HF. Whither ultrasonic prediction of fetal macrosomia? Obstet Gynecol 1993;82(5):860-2.

Sandmire HF. The labor-adjusted cesarean section: a more informative method than the cesarean section "rate" for assessing a practitioner's labor and delivery skills. Am J Obstet Gynecol 1998;178(1 Pt 1):196-7.

Sandmire HF, DeMott RK. The Green Bay cesarean section study. III. Falling cesarean birth rates without a formal curtailment program. Am J Obstet Gynecol 1994;170(6):1790-9; discussion 1799-802.

Sandmire HF, DeMott RK. The Green Bay cesarean section study. IV. The physician factor as a determinant of cesarean birth rates for the large fetus. Am J Obstet Gynecol 1996;174(5):1557-64.

Santerre RE. The effect of the ACOG guideline on vaginal births after cesarean. Med Care Res Rev 1996;53(3):315-29.

Sarah R. Power, certainty, and the fear of death. Women Health 1987;13(1-2):59-71.

Sarno AP, Jr., Phelan JP, Ahn MO, et al. Vaginal birth after cesarean delivery. Trial of labor in women with breech presentation. J Reprod Med 1989;34(10):831-3.

Satin AJ, Hankins GD, Yeomans ER. A prospective study of two dosing regimens of oxytocin for the induction of labor in patients with unfavorable cervices. Am J Obstet Gynecol 1991;165(4 Pt 1):980-4.

Satin AJ, Leveno KJ, Sherman ML, et al. High-dose oxytocin: 20- versus 40-minute dosage interval. Obstet Gynecol 1994;83(2):234-8.

Sato K, Takahashi K, Shioda K. Uterine rupture during a trial of labor in a case with a unicornuate uterus and a previous cesarean section. Gynecol Obstet Invest 1993;36(2):124-6.

Sato T, Konishi F, Minakami H, et al. Pelvic floor disturbance after childbirth: vaginal delivery damages the upper levels of sphincter innervation. Dis Colon Rectum 2001;44(8):1155-61.

Savage W. Midwives' Journal. Changing attitudes to intervention. Nurs Times 1986;82(22):63-4.

Schachter M, Kogan S, Blickstein I. External cephalic version after previous cesarean section--a clinical dilemma. Int J Gynaecol Obster 1994;45(1):17-20.

Schauberger CW, Rooney BL, Beguin EA, et al. Evaluating the thirty minute interval in emergency cesarean sections. J Am Coll Surg 1994;179(2):151-5.

Schimmel LM, Hogan P, Boehler B, et al. The Yolo County Midwifery Service. A descriptive study of 496 singleton birth outcomes, 1990. J Nurse Midwifery 1992;37(6):398-403.

Schimmel LM, Schimmel LD, DeJoseph J. Toward lower cesarean birth rates and effective care: five years' outcomes of joint private obstetric practice. Birth 1997;24(3):181-7.

Schiotz HA. Rupture of the uterus in labour. An

unusual case followed with sonography. Arch Gynecol Obstet 1991;249(1):43-5.

Schneider A. Delivery after caesarean section. Br Med J (Clin Res Ed) 1987;295(6597):554-5.

Schneider G. Palpation of the uterine scar after vaginal delivery. Can Med Assoc J 1989;141(2):98.

Schneider J, Gallego D, Benito R. Trial of labour in patients with a prior caesarean section and an intervening vaginal delivery. Aust N Z J Obstet Gynaecol 1987;27(3):178-9.

Schneider J, Gallego D, Benito R. Trial of labor after an earlier cesarean section. A conservative approach. J Reprod Med 1988;33(5):453-6.

Schneider KTM, Luftner D, Rath W. Efficacy and safety of a 2-tier prostaglandin labor induction scheduIe. J Perinat Med 1994;22(5):399-407.

Schnitker KA. Uterine rupture during trial of labor: risk management recommendations. J Healthc Risk Manag 1999;19(4):12-6.

Schuitemaker N, van Roosmalen J, Dekker G, et al. Maternal mortality after cesarean section in The Netherlands. Acta Obstet Gynecol Scand 1997;76(4):332-4.

Schussman LC, Lutz LJ. Hazards and uses of prenatal diagnostic X-radiation. J Fam Pract 1982;14(3):473-80.

Sciscione AC, Nguyen L, Manley J, et al. A randomized comparison of transcervical Foley catheter to intravaginal misoprostol for preinduction cervical ripening. Obstet Gynecol 2001;97(4):603-7.

Sciscione AC, Nguyen L, Manley JS, et al. Uterine rupture during preinduction cervical ripening with misoprostol in a patient with a previous Caesarean delivery. Aust N Z J Obstet Gynaecol 1998;38(1):96-7.

Scott JR. Mandatory trial of labor after cesarean delivery: an alternative viewpoint. Obstet Gynecol 1991;77(6):811-4.

Scott JR. Avoiding labor problems during vaginal birth after cesarean delivery. Clin Obstet Gynecol 1997;40(3):533-41.

Sebitloane MH, Moodley J. Emergency peripartum hysterectomy. East Afr Med J 2001;78(2):70-4.

Secher NJ, Lange AP, Nielsen FH, et al. Induction of labor with and without primary amniotomy. A randomized study of prostaglandin E2 tablets and intravenous oxytocin. Acta Obstet Gynecol Scand 1981;60(3):237-41.

Seffah JD, Amaniampong K, Wilson JB. The use of the partograph in monitoring labor in a prior cesarean section. Int J Gynaecol Obster 1994;45(3):281.

Segal S, Gemer O, Zohav E, et al. Evaluation of breast stimulation for induction of labor in women with a prior cesarean section and in grandmultiparas. Acta Obstet Gynecol Scand 1995;74(1):40-1.

Seguin L, Therrien R, Champagne F, et al. The components of women's satisfaction with maternity care. Birth 1989;16(3):109-13.

Seidman DS, Laor A, Gale R, et al. Long-term effects of vacuum and forceps deliveries. Lancet 1991;337(8757):1583-5.

Seitchik J. Cesarean delivery in nulliparous women for failed oxytocin-augmented labor: route of delivery in subsequent pregnancy. Am J Obstet Gynecol 1982;143(4):393-7.

Setubal A, Clode N, Bruno-Paiva JL, et al. Vesicouterine fistula after manual removal of placenta in a woman with previous cesarean section. Eur J Obstet Gynecol Reprod Biol 1999;84(1):75-6.

Sewell JE. Cesarean Section--A Brief History. Washington, DC: Historian Librarian, The American College of Obstetricians and Gynecologists; 1993.

Shalev E, Battino S, Giladi Y, et al. External cephalic version at term--using tocolysis. Acta Obstet Gynecol Scand 1993;72(6):455-7.

Shaver DC, Bada HS, Korones SB, et al. Early and late intraventricular hemorrhage: the role of obstetric factors. Obstet Gynecol 1992;80(5):831-7.

Shearer EL. Cesarean section: medical benefits

and costs. Soc Sci Med 1993;37(10):1223-31.

Shearer MH. Commentary:Pondering the Study of Women's Psychosocial Outcomes. Birth 1990;17(1):24.

Sherman SJ, Greenspoon JS, Nelson JM, et al. Obstetric hemorrhage and blood utilization. J Reprod Med 1993;38(12):929-34.

Shimonovitz S, Botosneano A, Hochner-Celnikier D. Successful first vaginal birth after cesarean section: a predictor of reduced risk for uterine rupture in subsequent deliveries. Isr Med Assoc J 2000;2(7):526-8.

Shin YK. Shoulder pain in a trial of labor after cesarean delivery. South Med J 1989;82(10):1320.

Shiono PH, Fielden JG, McNellis D, et al. Recent trends in cesarean birth and trial of labor rates in the United States. JAMA 1987;257(4):494-7.

Shiono PH, McNellis D, Rhoads GG. Reasons for the rising cesarean delivery rates: 1978-1984. Obstet Gynecol 1987;69(5):696-700.

Shipp TD, Zelop CM, Repke JT, et al. Intrapartum uterine rupture and dehiscence in patients with prior lower uterine segment vertical and transverse incisions. Obstet Gynecol 1999;94(5 Pt 1):735-40.

Shipp TD, Zelop CM, Repke JT, et al. Labor after previous cesarean: influence of prior indication and parity. Obstet Gynecol 2000;95(6 Pt 1):913-6.

Shipp TD, Zelop CM, Repke JT, et al. Interdelivery Interval and risk of asymptomatic uterine rupture. Obstet Gynecol 2001;97(2):175-7.

Shipp TD, Zelop CM, Repke JT, et al. Interdelivery Interval and Risk of Symptomatic Uterine Rupture. Obstet Gynecol 2001;97(2):175-7.

Shorten A, Lewis DE, Shorten B. Trial of labour versus elective repeat caesarean section: a cost-effectiveness analysis. Aust Health Rev 1998;21(1):8-28.

Shy KK, Frost F, Ullom J. Out-of-hospital delivery in Washington State, 1975 to 1977. Am J Obstet Gynecol 1980;137(5):547-52.

Shy KK, LoGerfo JP, Karp LE. Evaluation of elective repeat cesarean section as a standard of care: an application of decision analysis. Am J Obstet Gynecol 1981;139(2):123-9.

Siddiqui D, Stiller RJ, Collins J, et al. Pregnancy outcome after successful external cephalic version. Am J Obstet Gynecol 1999;181(5 Pt 1):1092-5.

Sieck CC. Vaginal birth after cesarean section: a comparison of rural and metropolitan rates in Oklahoma. J Oklahoma State Med Assoc 1997;90(8):444-9.

Silberstein T, Wiznitzer A, Katz M, et al. Routine revision of uterine scar after cesarean section: has it ever been necessary? Eur J Obstet Gynecol Reprod Biol 1998;78(1):29-32.

Silver RK, Gibbs RS. Predictors of vaginal delivery in patients with a previous cesarean section, who require oxytocin. Am J Obstet Gynecol 1987;156(1):57-60.

Silver RK, Minogue J. When does a statistical fact become an ethical imperative? [see comments]. Am J Obstet Gynecol 1987;157(2):229-33.

Sims EJ, Newman RB, Hulsey TC. Vaginal birth after cesarean: to induce or not to induce. Am J Obstet Gynecol 2001;184(6):1122-4.

Sims PD, Cabral D, Daley W, et al. The incentive plan: an approach for modification of physician behavior. Am J Public Health 1984;74(2):150-2.

Sinusas K. Deliveries by family physicians in Connecticut: results of a practice-based research network. Arch Fam Med 2000;9(5):434-8.

Sjogren B. Reasons for anxiety about childbirth in 100 pregnant women. J Psychosomat Obstet Gynecol 1997;18(4):266-72.

Skelton AG. The relationship among cost, quality, and competition: an analysis of obstetrics services in Missouri hospitals. J Health Care Finance 1997;24(1):30-44.

Skinner G. Action: promoting normal birth. Pract

Midwife 2002;5(3):4.

Skupski DW, Chervenak FA, McCullough LB, et al. Cesarean delivery for intrapartum fetal heart rate abnormalities: incorporating survey data into clinical judgment. Obstet Gynecol 1996;88(1):60-4.

Slavin R. Best-evidence synthesis: An alternative to meta-analytic and traditional reviews. Educ Res 1986;15:5-11.

Small R, Lumley J, Donohue L, et al. Randomised controlled trial of midwife led debriefing to reduce maternal depression after operative childbirth. BMJ 2000;321(7268):1043-7.

Smith DW, Murphy NJ. Alaska's obstetrical delivery systems: a descriptive epidemiologic study. Alaska Med 2000;42(3):78-84.

Smith GCS, Pell JP, Cameron AD, et al. Risk of perinatal death associated with labor after previous cesarean delivery in uncomplicated term pregnancis. JAMA 2002;287(20):2684-90.

Smith LF. Development of a multidimensional labour satisfaction questionnaire: dimensions, validity, and internal reliability. Quality in Health Care 2001;10(1):17-22.

Smith LP, DeLeon A, Funnell WR, et al. A research-oriented system for McGill obstetrical and noenatal data (Mond). Acta Obstet Gynecol Scand 1982;109(Supplement):49-50.

Smith LP, Nagourney BA, McLean FH, et al. Hazards and benefits of elective induction of labor. Am J Obstet Gynecol 1984;148(5):579-85.

Socol M L, Garcia PM, Peaceman AM, et al. Reducing cesarean births at a primarily private university hospital. Am J Obstet Gynecol 1993;168(6 Pt 1):1748-54; discussion 1754-8.

Socol ML, Peaceman AM. Vaginal birth after cesarean: an appraisal of fetal risk. Obstet Gynecol 1999;93(5 Pt 1):674-9.

Soliman SR, Burrows RF. Cesarean section: analysis of the experience before and after the National Consensus Conference on Aspects of Cesarean Birth. Can Med Assoc J 1993;148(8):1315-20.

Soltan MH, Khashoggi T, Adelusi B. Pregnancy following rupture of the pregnant uterus. Int J Gynaecol Obster 1996;52(1):37-42.

Somell C. Induction of labor and cervical ripening with oral PGE2 in risk pregnancies. A placebo-controlled study. Acta Obstet Gynecol Scand 1987;66(7):633-7.

Somell C, Larsson B. Priming and induction of labor with oral PGE2 in patients with low Bishop score. Acta Obstet Gynecol Scand 1983;62(4):315-20.

Sorensen SS, Brocks V, Lenstrup C. Induction of labor and cervical ripening by intracervical prostaglandin E2. Obstet Gynecol 1985;65(1):110-4.

Sox HC, Jr. Evaluation of diagnostic testing. Ann Rev Med 1996;47:463-71.

Spaulding LB. Delivery through the maternal bladder during trial of labor. Obstet Gynecol 1992;80(3 Pt 2):512-4.

Spellacy WN. Vaginal birth after cesarean: a reward/penalty system for national implementation. Obstet Gynecol 1991;78(2):316-7.

Spellacy WN, Miller SJ, Winegar A. Pregnancy after 40 years of age. Obstet Gynecol 1986;68(4):452-4.

Spiegelhalter D, Thomas A, Best N. WinBUGS Version 1.2 User Manual. In. 1.2 ed. Cambridge: MRC Biostatistics Unit; 1999.

Spong FW. Deliveries, readmission rates and practice guidelines to reduce variations in obstetrical practice. Health Care Innov 1997;7(1):18-23.

Springen K. The right to choose. Cesarean sections are on the rise again. Public-health officials want to limit them, but many patients and doctors are resisting. Newsweek 2000;136(23):73-4.

Stafford RS. Alternative strategies for controlling rising cesarean section rates. JAMA 1990;263(5):683-7.

Stafford RS. Cesarean section use and source of payment: an analysis of California hospital

discharge abstracts. Am J Public Health 1990;80(3):313-5.

Stafford RS. The impact of nonclinical factors on repeat cesarean section. JAMA 1991;265(1):59-63.

Stafford RS, Sullivan SD, Gardner LB. Trends in cesarean section use in California, 1983 to 1990. Am J Obstet Gynecol 1993;168(4):1297-302.

Stalnaker BL, Maher JE, Kleinman GE, et al. Characteristics of successful claims for payment by the Florida Neurologic Injury Compensation Association Fund. Am J Obstet Gynecol 1997;177(2):268-71; discussion 271-3.

Stanco LM, Schrimmer DB, Paul RH, et al. Emergency peripartum hysterectomy and associated risk factors. Am J Obstet Gynecol 1993;168(3 Pt 1):879-83.

Stark CM, Smith RS, Lagrandeur RM, et al. Need for urgent delivery after third-trimester amniocentesis. Obstet Gynecol 2000;95(1):48-50.

Steer PJ, Carter MC, Choong K, et al. A multicentre prospective randomized controlled trial of induction of labour with an automatic closed-loop feedback controlled oxytocin infusion system. Br J Obstet Gynaecol 1985;92(11):1127-33.

Steer PJ, Kennedy JH, Calder AA, et al. Induction of labour: A comparison of a single prostaglandin E2 vaginal tablet with amniotomy and intravenous oxytocin. Br J Obstet Gynaecol 1983;90(2):186-187.

Stein ZA. A woman's age: childbearing and child rearing. Am J Epidemiol 1985;121(3):327-42.

Stone C, Halliday J, Lumley J, et al. Vaginal births after Caesarean (VBAC): a population study. Paediatr Perinat Epidemiol 2000;14(4):340-8.

Stone J, Lockwood CJ, Berkowitz GS, et al. Morbidity of failed labor in patients with prior cesarean section. Am J Obstet Gynecol 1992;167(6):1513-7.

Stone JL, Lockwood CJ, Berkowitz G, et al. Use of cervical prostaglandin E2 gel in patients with previous cesarean section. Am J Perinatol 1994;11(4):309-12.

Stone SE, Brown MP, Westcott JP. Nurse-midwifery service in a rural setting. J Nurse Midwifery 1996;41(5):377-82.

Stovall TG, Shaver DC, Solomon SK, et al. Trial of labor in previous cesarean section patients, excluding classical cesarean sections. Obstet Gynecol 1987;70(5):713-717.

Strong TH, Jr., Brown WL, Jr., Brown WL, et al. Experience with early postcesarean hospital dismissal. Am J Obstet Gynecol 1993;169(1):116-9.

Strong TH, Phelan JP, Ahn MO, et al. Vaginal birth after cesarean delivery in the twin gestation. Am J Obstet Gynecol 1989;161(1):29-32.

Strong TH, Vega JS, O'Shaughnessy MJ, et al. Amnioinfusion among women attempting vaginal birth after cesarean delivery. Obstet Gynecol 1992;79(5 (Pt 1)):673-4.

Stronge J. Factors Affecting Mode of Delivery In Labour Following A Single Previous Birth by Caesarean Section. J Obstet Gynaecol 1996;16(5):353-7.

Stuart KA, Krakauer H, Schone E, et al. Labor epidurals improve outcomes for babies of mothers at high risk for unscheduled cesarean section. J Perinatol 2001;21(3):178-85.

Studnicki J, Remmel R, Campbell R, et al. The impact of legislatively imposed practice guidelines on cesarean section rates: the Florida experience. Am J Med Qual 1997;12(1):62-8.

Sturdee DW, Rushton DI. Caesarean and post-partum hysterectomy 1968-1983. Br J Obstet Gynaecol 1986;93(3):270-4.

Suk Wai N, Wing Kee T, Lao T, et al. Cervical priming with oral misoprostol in pre-labor rupture of membranes at term. Obstet Gynecol 1996;87(6):923-926.

Sultan AH, Kamm MA. Faecal incontinence after childbirth. Br J Obstet Gynaecol 1997;104(9):979-82.

Sultan AH, Kamm MA, Hudson CN. Pudendal nerve damage during labour: prospective study

before and after childbirth. Br J Obstet Gynaecol 1994;101(1):22-8.

Sultan AH, Kamm MA, Hudson CN, et al. Anal-sphincter disruption during vaginal delivery. N Engl J Med 1993;329(26):1905-11.

Sultan AH, Stanton SL. Preserving the pelvic floor and perineum during childbirth--elective caesarean section? Br J Obstet Gynaecol 1996;103(8):731-4.

Sun MS, Hseu SS, Chang DS, et al. Anesthetic management in parturients with uterine rupture preoperatively--report of two cases. Acta Anaesthesiol Sin 1997;35(3):167-70.

Suner S, Jagminas L, Peipert JF, et al. Fatal spontaneous rupture of a gravid uterus: case report and literature review of uterine rupture. J Emerg Med 1996;14(2):181-5.

Suonio S, Saarikoski S, Kaariainen J, et al. Intrapartum rupture of uterus diagnosed by ultrasound: a case report. Int J Gynaecol Obster 1984;22(5):411-3.

Sutcliffe E. Factors predicting elective repeat cesarean delivery. Obstet Gynecol 1994;84(4):641-2.

Suzuki S, Sawa R, Yoneyama Y, et al. Preoperative diagnosis of dehiscence of the lower uterine segment in patients with a single previous Caesarean section. Aust N Z J Obstet Gynaecol 2000;40(4):402-4.

Swaim LS, Holste CS, Waller DK. Umbilical cord blood pH after prior cesarean delivery. Obstet Gynecol 1998;92(3):390-3.

Sweeten KM, Graves WK, Athanas siou A. Spontaneous rupture of the unscarred uterus. Am J Obstet Gynecol 1995;172(6):1851-5; discussion 1855-6.

Sykes GS, Molloy PM, Johnson P, et al. Fetal distress and the condition of newborn infants. Br Med J (Clin Res Ed) 1983;287(6397):943-5.

Taffe l SM. Cesarean section in America: dramatic trends, 1970 to 1987. Stat Bull Metrop Insur Co 1989;70(4):2-11.

Taffel SM, Placek PJ. Complications in cesarean and non-cesarean deliveries: United States, 1980. Am J Public Health 1983;73(8):856-60.

Taffel SM, Placek PJ, Liss T. Trends in the United States cesarean section rate and reasons for the 1980-85 rise. Am J Public Health 1987;77(8):955-9.

Taffel SM, Placek PJ, Moien M, et al. 1989 U.S. cesarean section rate steadies--VBAC rate rises to nearly one in five. Birth 1991;18(2):73-7.

Tahilramaney MP, Boucher M, Eglinton GS, et al. Previous cesarean section and trial of labor. Factors related to uterine dehiscence. J Reprod Med 1984;29(1):17-21.

Tamale-Sali EG, Iskandar MN. Is thre a risk of lower segment scar rupture in pregnancy after multiple caesarean sections? J Obstet Gynaecol 1992;12:19-21.

Tammelleo AD. "Asleep on duty" record impacts delay in delivery case. Case in point: St. Paul Medical Center v. Cecil (842 S.W.2d 808--TX [1992]). Regan Rep Nurs Law 1993;33(10):2.

Tammelleo AD. Failure to "fully communicate": catastrophic results. Case in point: Baptist Medical Center v. Wilson 618 So. 2d 1335--AL (1993). Regan Rep Nurs Law 1993;34(3):2.

Tammelleo AD. Nurse opines on non-natural childbirth: freedom of speech. Regan Rep Nurs Law 1993;34(2):1.

Tanik A, Ustun C, Cil E, et al. Sonographic evaluation of the wall thickness of the lower uterine segment in patients with previous cesarean section. J Clin Ultrasound 1996;24(7):355-7.

Targett C. Caesarean section and trial of scar. Aust N Z J Obstet Gynaecol 1988;28(4):249-62.

Tauer CA. Lives at stake. How to respond to a woman's refusal of cesarean surgery when she risks losing her child or her life. Health Prog 1992;73(7):18-27.

Tauer CA. When pregnant patients refuse interventions. AWHONNS 1993;4(4):596-605.

Taylor AVG, Sellers S, Ah-Moye M, et al. A prospective random allocation trial to compare vaginal prostaglandin E2 with intravenous oxytocin for labour induction in women

previously delivered by caesarean section. J Obstet Gynaecol 1993;13(5):333-336.

Taylor HA, Hayes SM, McGowan LM. The maternity care practice of Navy family practice residency graduates while on active duty. Mil Med 1997;162(9):620-2.

Tetzschner T, Sorensen M, Jonsson L, et al. Delivery and pudendal nerve function. Acta Obstet Gynecol Scand 1997;76(4):324-31.

Tews G, Ebner T, Yaman C, et al. Maternal ascites into the abdomen in a patient with status post adnexectomy and uterine rupture. Acta Obstet Gynecol Scand 2001;80(5):474-5.

Thacker SB, Peterson HB, Stroup DF. Metaanalysis for the obstetrician-gynecologist. Am J Obstet Gynecol 1996;174(5):1403-7.

Thacker SB, Stroup DF. Continuous electronic heart rate monitoring for fetal assessment during labor. Cochrane Database of Systematic Reviews 2000(2):CD000063.

Thiery M, Decoster JM, Parewijck W, et al. Endocervical prostaglandin E2 gel for preinduction cervical softening. Prostaglandins 1984;27(3):429-39.

Thomas IL, Chenoweth JN, Tronc GN, et al. Preparation for induction of labour of the unfavourable cervix with Foley catheter compared with vaginal prostaglandin. Aust N Z J Obstet Gynaecol 1986;26(1):30-5.

Thompson MS, Cohen AB. Should we measure personal valuations of perinatal outcomes? Mead Johnson Symp Perinat Dev Med 1982(20):47-52.

Thorp JM, Bowes WA. Vaginal birth after abdominal delivery. Am J Obstet Gynecol 1990;163(3):1101-2.

Thubisi M, Ebrahim A, Moodley J, et al. Vaginal delivery after previous caesarean section: is X-ray pelvimetry necessary? [see comments]. Br J Obstet Gynaecol 1993;100(5):421-4.

Thurnau GR, Scates DH, Morgan MA. The fetal-pelvic index: a method of identifying fetal-pelvic disproportion in women attempting vaginal birth after previous cesarean delivery. Am J Obstet Gynecol 1991;165(2):353-8.

Tomlinson AJ, Archer PA, Hobson S. Induction of labour: A comparison of two methods with particular concern to patient acceptability. J Obstet Gynaecol 2001;21(3):239-241.

Torres A, Reich MR. The shift from home to institutional childbirth: a comparative study of the United Kingdom and The Netherlands. Int J Health Serv 1989;19(3):405-14.

Traynor JD, Peaceman AM. Maternal hospital charges associated with trial of labor versus elective repeat cesarean section. Birth 1998;25(2):81-4.

Trofatter KF, Bowers D, Gall SA, et al. Preinduction cervical ripening with prostaglandin E2 (Prepidil) gel. Am J Obstet Gynecol 1985;153(3):268-71.

Troyer LR, Parisi VM. Obstetric parameters affecting success in a trial of labor: designation of a scoring system. Am J Obstet Gynecol 1992;167(4 Pt 1):1099-104.

Tucker JM, Hauth JC, Hodgkins P, et al. Trial of labor after a one- or two-layer closure of a low transverse uterine incision. Am J Obstet Gynecol 1993;168(2):545-6.

Tulman L, Fawcett J, Groblewski L, et al. Changes in functional status after childbirth. Nurs Res 1990;39:70-75.

Tuncer R, Erkaya S, Sipahi T, et al. Emergency postpartum hysterectomy. Journal of Gynecologic Surgery 1995;11(4):209-13.

Turan C, Kutlay B. Cesarean section rates and perinatal outcomes in resident and midwife attended low risk deliveries. Eur J Obstet Gynecol Reprod Biol 1995;62(1):3-5.

Turnbull D, Holmes A, Shields N, et al. Randomised, controlled trial of efficacy of midwife-managed care. Lancet 1996;348(9022):213-8.

Turnbull DA, Wilkinson C, Yaser A, et al. Women's role and satisfaction in the decision to have a caesarean section. Med J Aust 1999;170(12):580-3.

Turner MJ. Delivery after one previous cesarean section. Am J Obstet Gynecol 1997;176(4):741-4.

Turner MJ, McNally O, Gardeil F. Trial of labor compared with an elective second cesarean section. N Engl J Med 1997;336(9):658; discussion 659.

Turnquest MA, James T, Marcell C, et al. Vaginal birth after cesarean section in a university setting. J Ky Med Assoc 1994;92(6):216-21.

Tussing AD, Wojtowycz MA. The cesarean decision in New York State, 1986. Economic and noneconomic aspects. Med Care 1992;30(6):529-40.

Tyson H. Outcomes of 1001 midwife-attended home births in Toronto, 1983-1988. Birth 1991;18(1):14-9.

U.S. Department of Health and Human Services. Healthy People 2010. In: Office of Disease Prevention and Health Promotion; 2000.

Udo-Inyang A, Lee C, Evans E. Gravid uterine rupture following cesarean section and intervening vaginal delivery. Henry Ford Hosp Med J 1986;34(3):215-7.

Udom NU, Betley CL. Effects of maternity-stay legislation on 'drive-through deliveries'. Health Aff 1998;17(5):208-15.

Udvarhelyi IS, Colditz GA, Rai A, et al. Cost-effectiveness and cost-benefit analyses in the medical literature. Are the methods being used correctly? Ann Intern Med. 1992;116(3):238-44.

Uppington J. Epidural analgesia and previous caesarean section. Anaesthesia 1983;38(4):336-41.

van Alten D, Eskes M, Treffers PE. Midwifery in The Netherlands. The Wormerveer study; selection, mode of delivery, perinatal mortality and infant morbidity. Br J Obstet Gynaecol 1989;96(6):656-62.

van Amerongen D. Vaginal birth after cesarean section in an HMO. HMO Pract 1989;3(3):104-7.

van Amerongen D. Vaginal birth after cesarean section. Experience in a community-based practice. J Reprod Med 1989;34(8):531-4.

van der Walt WA, Cronje HS, Bam RH. Vaginal delivery after one cesarean section. Int J Gynaecol Obster 1994;46(3):271-7.

van Gelderen CJ, England MJ, Naylor GA, et al. Labour in patients with a caesarean section scar. The place of oxytocin augmentation. S Afr Med J 1986;70(9):529-32.

Van Tuinen I, Wolfe SM. Unnecessary Cesarean Sections: halting a National Epidemic. Washington: Public Citizens Health Research Group; 1992.

Vandevusse L. Decision making in analyses of women's birth stories. Birth 1999;26(1):43-50.

Varma A, Gunn J, Gardiner A, et al. Obstetric anal sphincter injury: prospective evaluation of incidence. Dis Colon Rectum 1999;42(12):1537-43.

Varma TR, Norman J. A comparison of three dosages of prostaglandin E2 pessaries for ripening the unfavourable cervix prior to induction of labor. Acta Obstet Gynecol Scand 1984;63(1):17-21.

Vause S, Macintosh M. Evidence based case report: use of prostaglandins to induce labour in women with a caesarean section scar. BMJ 1999;318(7190):1056-8.

Vedat A, Hasan B, Ismail A. Rupture of the uterus in labor: a review of 150 cases. Isr J Med Sci 1993;29(10):639-43.

Veridiano NP, Thorner NS, Ducey J. Vaginal delivery after cesarean section. Int J Gynaecol Obster 1989;29(4):307-11.

Vestermark V, Hogdall CK, Birch M, et al. Influence of the mode of delivery on initiation of breast-feeding. Eur J Obstet Gynecol Reprod Biol 1991;38(1):33-8.

Videla FL, Satin AJ, Barth WH, et al. Trial of labor: a disciplined approach to labor management resulting in a high rate of vaginal delivery. Am J Perinatol 1995;12(3):181-4.

Vimercati A, Greco P, Kardashi A, et al. Choice of cesarean section and perception of legal pressure. J Perinat Med 2000;28(2):111-7.

Vintzileos AM, Gaffney SE, Salinger LM, et al. The relationship between fetal biophysical

profile and cord pH in patients undergoing cesarean section before the onset of labor. Obstet Gynecol 1987;70(2):196-201.

Vintzileos AM, Nochimson DJ, Guzman ER, et al. Intrapartum electronic fetal heart rate monitoring versus intermittent auscultation: a meta-analysis. Obstet Gynecol 1995;85(1):149-55.

Vinueza CA, Chauhan SP, Barker L, et al. Predicting the success of a trial of labor with a simple scoring system. J Reprod Med 2000;45(4):332-6.

Vroman S, Sian AYL, Thiery M, et al. Elective induction of labor conducted under lumbar epidural block I. Labor induction by anniotomy and intravenous oxytocin. Eur J Obstet Gynecol Reprod Biol 1977;7:159-80.

Wadhawan S, Narone JN. Outcome of labor following previous cesarean section. Int J Gynaecol Obster 1983;21(1):7-10.

Wagner CL, Metts AK. Rates of successful vaginal delivery after cesarean for patients with private versus public insurance. J Perinatol 1999;19(1):14-8.

Waldenstrom U. Experience of labor and birth in 1111 women. J Psychosom Res 1999;47(5):471-82.

Wall EM. "But doctor, I'd prefer to have a cesarean section": when public policy conflicts with patient preference. J Am Board Fam Pract 1995;8(5):414-6.

Walmsley K, Hobbs L. Vaginal birth after lower segment caesarean section. Modern Midwife 1994;4(4):20-1.

Walton DL, Ludlow D, Willis DC. Vaginal birth after cesarean section. Acceptance and outcome at a rural hospital. J Reprod Med 1993;38(9):716-8.

Wang PH, Yuan CC, Chao HT, et al. Posterior uterine wall rupture during labour. Hum Reprod 2000;15(5):1198-9.

Ware JE, Snow KK, Kosinski MA, et al. SF-36 health Survey Manual and Interpretation Guide. Boston: The Health Institute, New England Medical Center; 1997.

Watson KM. Induction of labour by prostaglandin tablets in a general practitioner maternity unit. Practitioner 1981;225(1354):545-549.

Wax JR, Philput C, Mather J, et al. Twin vaginal birth after cesarean. Conn Med 2000;64(4):205-8.

Wax JR, Sutula K, Lerer T, et al. Labor and delivery following successful external cephalic version. Am J Perinatol 2000;17(4):183-6.

Wear A. Problems in labor. Nursing - Oxford 1986;3(2):46-51.

Wearing MP. Comparison of vaginal delivery and cesarean section in breech presentation. Am J Obstet Gynecol 1980;136(8):1083.

Webb JC, Gilson G, Gordon L. Late second stage rupture of the uterus and bladder with vaginal birth after cesarean section: a case report and review of the literature. J Matern Fetal Med 2000;9(6):362-5.

Weber DO. At Saddleback Memorial, care maps detail birth process interventions, cutting C-section rate and raising VBACs. Strategies for Healthcare Excellence 1997;10(5):7-9.

Weeks JW, Pitman T, Spinnato JA. Fetal macrosomia: does antenatal prediction affect delivery route and birth outcome? Am J Obstet Gynecol 1995;173(4):1215-9.

Weerasekera DS, Grant JM. Sweeping of the membranes is an effective method of induction of labour in prolonged pregnancy: A report of a randomised trial [2]. Br J Obstet Gynaecol 1993;100(2):193-194.

Weinstein D, Benshushan A, Ezra Y, et al. Vaginal birth after cesarean section: current opinion. Int J Gynaecol Obster 1996;53(1):1-10.

Weinstein D, Benshushan A, Tanos V, et al. Predictive score for vaginal birth after cesarean section. Am J Obstet Gynecol 1996;174(1 Pt 1):192-8.

Weiss J, Nannini A, Fogerty S, et al. Use of hospital discharge data to monitor uterine rupture--Massachusetts, 1990-1997; US Department of Health & Human Services. MMWR Morb Mortal Wkly Rep

2000;49(12):245-8.

Weiss RR. Case 9-1998: uterine rupture. N Engl J Med 1998;339(4):268; discussion 269.

Wen SW, Liu S, Marcoux S, et al. Trends and variations in length of hospital stay for childbirth in Canada. Can Med Assoc J 1998;158(7):875-80.

Wennberg JE, Barnes BA, Zubkoff M. Professional uncertainty and the problem of supplier-induced demand. Soc Sci Med 1982;16(7):811-24.

Westgate J, Williams JA. Evaluation of a controlled release vaginal prostaglandin E2 pessary with a retrieval system for the induction of labour. J Obstet Gynaecol 1994;14(3):146-150.

Westgren M, Ingemarsson E, Ingemarsson I, et al. Intrapartum electronic fetal monitoring in low-risk pregnancies. Obstet Gynecol 1980;56(3):301-4.

Whiteside DC, Mahan CS, Cook JC. Factors associated with successful vaginal delivery after cesarean section. J Reprod Med 1983;28(11):785-8.

Whitford HM, Hillan EM. Women's perceptions of birth plans. Midwifery 1998;14(4):248-53.

Whitsel AI, Capeless EC, Abel DE, et al. Adjustment for case mix in comparisons of cesarean delivery rates: university versus community hospitals in Vermont. Am J Obstet Gynecol 2000;183(5):1170-5.

Wijma K, Soderquist J, Wijma B. Posttraumatic stress disorder after childbirth: a cross sectional study. J Anxiety Disord 1997;11(6):587-97.

Wilf RT, Franklin JB. Six years' experience with vaginal births after cesareans at Booth Maternity Center in Philadelphia. Birth 1984;11(1):5-9.

Wilkinson C, McIlwaine G, Patel N. Vaginal birth after caesarean section. Scottish Caesarean Section Audit Group. Lancet 1995;345(8951):725.

Williams JK, Lewis ML, Cohen GR, et al. The sequential use of estradiol and prostaglandin E2 topical gels for cervical ripening in high-risk term pregnancies requiring induction of labor. Am J Obstet Gynecol 1988;158(1):55-8.

Williams JK, Wilkerson WG, O'Brien WF, et al. Use of prostaglandin E2 topical cervical gel in high-risk patients: a critical analysis. Obstet Gynecol 1985;66(6):769-73.

Williams LS. Relations between Alberta midwives, MDs appear to be thawing despite high-profile trial. Can Med Assoc J 1991;145(5):497-50.

Williams MA, Luthy DA, Zingheim RW, et al. Preinduction prostaglandin E2 gel prior to induction of labor in women with a previous cesarean section. Gynecol Obstet Invest 1995;40(2):89-93.

Williams MC, Knuppel RA, O'Brien WF, et al. A randomized comparison of assisted vaginal delivery by obstetric forceps and polyethylene vacuum cup. Obstet Gynecol 1991;78(5 Pt 1):789-94.

Williams RL, Chen PM. Identifying the sources of the recent decline in perinatal mortality rates in California. N Engl J Med 1982;306(4):207-14.

Williams RL, Chen PM. Controlling the rise in cesarean section rates by the dissemination of information from vital records. Am J Public Health 1983;73(8):863-7.

Wilner S, Schoenbaum SC, Monson RR, et al. A comparison of the quality of maternity care between a health-maintenance organization and fee-for-service practices. N Engl J Med 1981;304(13):784-7.

Wilson PD, Herbison RM, Herbison GP. Obstetric practice and the prevalence of urinary incontinence three months after delivery. Br J Obstet Gynaecol 1996;103(2):154-61.

Wing DA, Lovett K, Paul RH. Disruption of prior uterine incision following misoprostol for labor induction in women with previous cesarean delivery. Obstet Gynecol 1998;91(5 Pt 2):828-30.

Wing DA, Paul RH. Induction of labor with misoprostol for premature rupture of membranes beyond thirty-six weeks' gestation. Am J Obstet Gynecol 1998;179(1):94-9.

Wing DA, Paul RH. Vaginal birth after cesarean section: selection and management. Clin Obstet Gynecol 1999;42(4):836-48.

Winslow R. Patient data may reshape health care. Efforst to cut costs, pinpoint treatment gain. Wall Street Journal 1989(April 17).

Wiqvist I, Norstrom A, Wiqvist N. Induction of labor by intra-cervical PGE2 in viscous gel. Mechanism of action and clinical treatment routines. Acta Obstet Gynecol Scand 1986;65(5):485-92.

Witlin AG, Mattar F, Sibai BM. Postpartum stroke: a twenty-year experience. Am J Obstet Gynecol 2000;183(1):83-8.

Witter FR, Rocco LE, Johnson TR. A randomized trial of prostaglandin E2 in a controlled-release vaginal pessary for cervical ripening at term. Am J Obstet Gynecol 1992;166(3):830-4.

Wittich AC. Uterine scar separation in patients undergoing trial of labor (TOL) in one army hospital. Mil Med 2000;165(10):730-2.

Woo GM, Twickler DM, Stettler RW, et al. The pelvis after cesarean section and vaginal delivery: normal MR findings. AJR Am J Roentgenol 1993;161(6):1249-52.

Woods JR, Saywell RM, Jr., Benson JT. Comparative costs of a cooperative care program versus inpatient hospital care for obstetric patients. Med Care 1988;26(6):596-606.

Woolbright LA. Why is the cesarean delivery rate so high in Alabama? An examination of risk factors, 1991-1993. Birth 1996;23(1):20-5.

Woolery W. Informed consent issues throughout the birthing process. J Leg Med 2000;21(2):241-55.

Wright CH, Gardin TH, Wright CL. Obstetric care in a health maintenance organization and a private fee-for-service practice: a comparative analysis. Am J Obstet Gynecol 1984;149(8):848-56.

Wright JB, Wright AL, Simpson NA, et al. A survey of trainee obstetricians preferences for childbirth. Eur J Obstet Gynecol Reprod Biol 2001;97(1):23-5.

Wright M. The value of X-ray pelvimetry in previous caesarean section pregnancies. S Afr Med J 1985;68(6):409-11.

Wynn P. Hospitals, health plans and employers unite to reduce cesarean deliveries. Manag Care 1996;5(12):57-8.

Xenakis E-J, Langer O, Piper JM, et al. Low-dose versus high-dose oxytocin augmentation of labor - A randomized trial. Am J Obstet Gynecol 1995;173(6):1874-1878.

Yamani TY, Rouzi AA. Induction of labor with vaginal prostaglandin-E2 in grand multiparous women. International Journal of Gynecology & Obstetrics 1998;62(3):255-259.

Yamani TY, Rouzi AA. Induction of labor with vaginal prostaglandin-E2 in grand multiparous women with one previous cesarean section. Int J Gynaecol Obster 1999;65(3):251-3.

Yan JS, Chang YK, Yin CS. Elective cesarean section for macrosomia? Chin Med J 1994;53(3):141-5.

Yang JY, Fang LJ, Tsou Yau KI. Labor pain before elective cesarean section reduces neonatal respiratory distress. Chung-Hua Min Kuo Hsiao Erh Ko i Hsueh Hui Tsa Chih 1997;38(1):38-43.

Yanover MJ, Jones D, Miller MD. Perinatal care of low-risk mothers and infants. Early discharge with home care. N Engl J Med 1976;294(13):702-5.

Yap OW, Kim ES, Laros RK, Jr. Maternal and neonatal outcomes after uterine rupture in labor. Am J Obstet Gynecol 2001;184(7):1576-81.

Yasumizu T, Nozawa A, Kinoshita T, et al. Trial of Labor in the Patient with a Prior Cesarean Section. Yamanashi Med J 1993;8(3):113-119.

Yasumizu T, Nozawa A, Kinoshita T, et al. Trial of vaginal birth following cesarean section for arrest disorders of labor: analysis of patients with well-documented medical records. Asia Oceania J Obstet Gynaecol 1994;20(4):407-13.

Yeast JD, Jones A, Poskin M. Induction of labor and the relationship to cesarean delivery: A review of 7001 consecutive inductions. [see comments.]. Am J Obstet Gynecol 1999;180(3 Pt 1):628-33.

Yeh S, Huang X, Phelan JP. Postterm pregnancy after previous cesarean section. J Reprod Med 1984;29(1):41-4.

Yetman TJ, Nolan TE. Vaginal birth after cesarean section: a reappraisal of risk. Am J Obstet Gynecol 1989;161(5):1119-23.

Yip SK, Fung HY, Wong WS, et al. Vesico-uterine fistula--a rare complication of vacuum extraction in a patient with previous caesarean section. Br J Urol 1997;80(3):502-3.

Yla-Outinen A, Heinonen PK, Tuimala R. Predisposing and risk factors of umbilical cord prolapse. Acta Obstet Gynecol Scand 1985;64(7):567-70.

Young D. A new push to reduce cesareans in the United States. Birth 1997;24(1):1-3.

Young D. Confrontation in Kansas City: elective cesareans and maternal choice. Birth 2000;27(3):153-5.

Yudkin P. Identifying the sources of the recent decline in perinatal mortality rates in California: two commentaries and a reply. Birth 1983;10(1):39-46.

Zahniser SC, Kendrick JS, Franks AL, et al. Trends in obstetric operative procedures, 1980 to 1987. Am J Public Health 1992;82(10):1340-4.

Zakut H, Blankstein J, Kramer J, et al. The outcome of pregnancy following abdominal surgery during pregnancy. Clin Exp Obstet Gynecol 1981;8(2):45-9.

Zelop CM, Shipp TD, Cohen A, et al. Trial of labor after 40 weeks' gestation in women with prior cesarean. Obstet Gynecol 2001;97(3):391-3.

Zelop CM, Shipp TD, Repke JT, et al. Uterine rupture during induced or augmented labor in gravid women with one prior cesarean delivery. Am J Obstet Gynecol 1999;181(4):882-6.

Zelop CM, Shipp TD, Repke JT, et al. Effect of previous vaginal delivery on the risk of uterine rupture during a subsequent trial of labor. Am J Obstet Gynecol 2000;183(5):1184-6.

Zelop CM, Shipp TD, Repke JT, et al. Outcomes of trial of labor following previous cesarean delivery among women with fetuses weighing >4000 g. Am J Obstet Gynecol 2001;185(4):903-5.

Zhang J, Klebanoff MA, DerSimonian R. Epidural analgesia in association with duration of labor and mode of delivery: a quantitative review. Am J Obstet Gynecol 1999;180(4):970-7.

Zhou SW, Chi IC. Immediate postpartum IUD insertions in a Chinese hospital--a two year follow-up. Int J Gynaecol Obster 1991;35(2):157-64.

Ziadeh SM, El-Jallad MF, Sunn'a EI. Obstetric uterine rupture: a four-year clinical analysis. Gynecol Obstet Invest 1999;48(3):176-8.

Zinberg S. Vaginal delivery after previous cesarean delivery: A continuing controversy. Clin Obstet Gynecol 2001;44(3):561-570.

Zisow DL. Uterine rupture as a cause of shoulder dystocia. Obstet Gynecol 1996;87(5 Pt 2):818-9.

Zorlu CG, Danisman N, Caglar T, et al. Vaginal birth following unmonitored labor in patients with prior cesarean section. Gynecol Obstet Invest 1996;42(4):222-6.

Zorlu CG, Turan C, Isik AZ, et al. Emergency hysterectomy in modern obstetric practice. Changing clinical perspective in time. Acta Obstet Gynecol Scand 1998;77(2):186-90.

Zweig S, Kruse J, LeFevre M. Patient satisfaction with obstretric care. J Fam Pract 1986;23:131-136.