

RESOURCE NOTES

NO. 34

DATE 08/01/00

Effects of Contraception and Removal Treatments on Pryor Mountain Wild Horse Population Demographics and Genetics

*By: John E. Gross
CSIRO Davies Laboratory
Queensland, Australia*

*The twelfth in a series of 13,
Session 4*

Background

Most free-ranging horse populations in the United States inhabit Bureau of Land Management (BLM) lands and the BLM is mandated to manage these herds. Wild horse populations have the ability to rapidly increase in size, and land managers are thus faced with the need to control the size of a rapidly increasing population.

Because the management objective for many horse populations is small (<150 horses), there are also concerns about the long-term genetic viability of these small populations. Managers are thus faced with conflicting needs to minimize population

size to control habitat damage or forage use, and to maximize population size to preserve genetic variation. To evaluate consequences of population control strategies that used contraception, periodic removals, or a combination of the two, I developed a population viability model that simulated population dynamics and changes in genetic variation and applied the model to horses on the Pryor Mountain Wild Horse Range. Simulated management actions were applied to horses that were young, old, or a random selection from the population.

Discussion

Computer simulations were used to evaluate management alternatives that included changes in population objective (AML), contraceptive treatment, and removals. Management actions (contraception and/or removal) focused on treatment/removal of young horses, old horses, or a random selection of individuals. Population responses included average population size, variation in population size, population age structure, growth rate, changes in genetic variation (heterozygosity), and loss of alleles.

Model results showed that simulated control strategies had striking differences in terms of population structure and persistence of genetic diversity. When comparing contraceptives to removal, it was found that use of contraceptives can greatly reduce the variation in yearly population size, and average population sizes remained much closer to the objective. However, population growth rate is relatively insensitive to low levels of infertility, and there are sharp contrasts between the dynamics of populations controlled by harvest or contraception. With contraception, populations can increase rapidly if a high level of infertility is not achieved, but population size declines slowly because natural mortality is typically low. In contrast, removals permit managers to rapidly reduce population size, modify sex ratio, or adjust the age structure of the population. For most wild horse populations, about 70% of all reproductively active females will need to be maintained in an infertile state to achieve a stable population size. For the conditions simulated, the age of animal treated had a huge impact on the rate of loss of genetic variation. Control

WILD HORSE AND
BURRO PROGRAM



strategies that delayed the age of first reproduction, such as contraception or removal of young animals, greatly reduced loss of genetic variation. In contrast, strategies that emphasized reproduction by young animals (contraception or removal of old horses) increased the rate of loss of genetic diversity. To retain genetic diversity, a particularly poor suggestion is to permit a horse to reproduce and then rendering it infertile. These effects result from changes in generation time – rate of loss of genetic variation is directly proportional to generation time, and strategies that increase generation time (e.g., delaying reproduction) reduce the rate of loss of genetic resources. For the sizes of populations simulated

(AML of 90 to 180 horses), the age of treatment had a greater effect than an overall change in AML (total population size).

Conclusion

Regardless of control strategy, genetic variation is lost much more slowly if young animals are treated (e.g., removed or rendered temporarily infertile). The most practical control program will likely involve both contraceptives and periodic removals. Contraceptives can reduce growth rate and are likely to be cost-effective, while removals permit managers to rapidly adjust sex ratio, age structure, or overall population size. If single-year contraceptives are used to maintain infertility, a very intensive management

program will be necessary. In small populations, changes in the age of horses treated will likely have more impact on loss of genetic variation than changes in the number of horses in the population.

Contact

John E. Gross, PhD.
CSIRO Davies Laboratory
PMB Aitkenvale
Queensland, 4814 Australia
phone 61 7 4753 8509
fax 61 7 4753 8600
email:
John.Gross@tag.csiro.au

RESOURCE NOTES are intended to be early announcements of technical and informational topics for Bureau of Land Management personnel and some of their customers. Information in this RESOURCE NOTE is based on the opinion and experience of the author and has not been peer-reviewed. Conclusions and opinions expressed herein do not necessarily represent those of BLM. Use of trade names does not imply U.S. Government endorsement of commercial products.

If you have received a copy of or found out about RESOURCE NOTES in an indirect way and would like to be included in future mailings, please send the following:

NAME, TITLE, MAILING ADDRESS and a list of the two or three subject areas that you are most interested in or that most directly relate to your job. Send this information to Phil Dittberner, BLM, RS-140, P.O. Box 25047, Denver, CO. 80225-0047 or phil_dittberner@blm.gov or FAX 303-236-3508.

If you would like to prepare a RESOURCE NOTE for distribution, or you have an idea and author in mind for a good RESOURCE NOTE, please contact Phil Dittberner at 303-236-1833, FAX 303-236-3508 or phil_dittberner@blm.gov with the topic and the name of writer, including an address, phone number, and e-mail address.

Thank you for your interest in RESOURCE NOTES.



National Science &
Technology Center 
Bureau of Land Management